


9608  
C187

DL596  
S7E6



**Cornell University Library**  
Ithaca, New York

FROM THE  
**UNITED STATES GOVERNMENT**  
THROUGH THE  
SUPERINTENDENT OF DOCUMENTS

§ 506 Rev. Stat. prohibits the withdrawal of this book for home use.



# DATE DUE

DEC 21 1974			
MAY 19 1975			
AUG 7 1975			
DEC 11 1975			
DEC 11 1975			
MAY 13 1976			
AUG 20 1976			
DEC 16 1976			
MAY 13 1977			
MAY 14 1985			
SEP 6 1985			
AUG 22 1990			

RETURN TO

ENTOMOLOGY LIBRARY

Cornell University  
Ithaca, N. Y.

Cornell University Library

QL 596.S7E6

Genera et species staphylinorum insector



3 1924 018 330 732

ent





*Very Sun*

GENERA ET SPECIES

STAPHYLINORUM

INSECTORUM COLEOPTERORUM

FAMILIAE.

GENERA ET SPECIES

STAPHYLINORUM

AUCTORE

**GUIL. F. ERICHSON.**

---

1847

STAPHYLINORUM  
GENERA ET SPECIES

LECTOR

GUIL. F. ERICHSON.



# GENERA ET SPECIES STAPHYLINORUM

## INSECTORUM COLEOPTERORUM FAMILIAE

AUCTORE

**GUIL. F. ERICHSON**

MEDICINAE ET PHILOSOPHIAE DOCTORE HISTOR. NAT. IN UNIVERS. LITTER. REG. FRIDERICA  
GUILIELMA BEROLIN. PRIV. DOCENTE ACAD. SCIENC. REG. TAURIN. SOCIO ACAD. NAT. CURIOS.  
CAESAR. LEOPOLDINAE SOCIET. IMPERIAL. NAT. SCRUT. MOSQUENSIS ET SOC. ENTOMOL.  
LONDINENSIS SODALI SOCIET. NAT. SCRUTAT. BEROLINENSIS SOCIO HONORARIO

---

ACCEDUNT TABULAE AENEAE QUINQUE

---

**BEROLINI**

**F. H. MORIN**

MDCCCL

CORNELL  
UNIVERSITY  
LIBRARY

GENERA ET SPECIES

Countdown  
4/11/72

DL 596

STAPHYLINUM

INSECTORUM COLEOPTERORUM

FAMILIAE

9608

C 187

GUIL. F. ERICHSON

ORDINIS ET PHILOSOPHIAE DOCTOR HISTOR. NAT. TO. CARLUS. FILIUS. REG. BRUNSV. /  
GUILIAMS BEROLIN. PRIN. DOCTOR. MAG. SCIENT. REG. LITHUEN. SOCIO. ACAD. NAT. LONDIN. /  
LITVON. GEOPHYSIC. SOCIET. IMPERIAL. NAT. SCIENT. MOSCOWENSIS ET SOC. ENTOMOL. /  
LONDINENSIS SOCIET. NAT. SCIENT. BRITANNICAE SOCIO. HONORARIO.

C. 570

ACCEDUNT TABULAE AERAE QUINQUE

BEROLINI

F. M. O. M. A. M.

LIBRARY

UNIVERSITY  
LIBRARY

Staphylini

20/72



## L. S.

**A**mplam hanc innumerae Insectorum classis familiam, a Gravenhorstio olim ingeniose illustratam, iam, specierum copia prodigiosum in modum gliscente, omnibus forte Coleopterorum saltem reliquis familiis minus excultam, labore peculiari tractatam in medium me proferre, non est quod praefando excusam: de rationibus autem, quas hoc in opere conformando secutus sum, pauca praemoneam, cum non tam speciebus quam plurimis enumeratis quam rite examinatis, distinctis, dispositis historiae naturalis scientia promoveri videatur.

Nitidissimus huius Coleopterorum familiae cultor, cel. Gravenhorst nuper \*) professus est, dispositiones nostras systematicas omnino arbitrarías neque ullo modo constitutas esse a natura, quae formas alteras extinguere, alteras novas procreare continuaret, ita ut sectiones omnes confluerent et, simulatque characteribus circumscriberentur, artificiales evaderent: quam quidem opinionem non magis atque illa de speciebus hybridis olim adducta sententia, observationibus accuratioribus nullo modo confirmata et ab omnibus neglecta, arridere naturae scrutatores puto, et cui tanto minus accedere possum, quanto magis ex historiae naturalis penitiori studio persuasum me habeo, non generum solum, sed familiarum etiam tribuumque et reliquarum divisionum limites adeo esse firmos, ut non alio ac specierum modo a natura ipsa positi videantur. Quo autem dicto methodos systematicas omnes omnium auctorum nolo approbare, quorum recentiores plurimi, historiae naturalis studium vix periclitati, differentias sibi observatas quantum valeant perpendere inscii, temerarias venditant dispositiones. Interest, ut characteres naturales rite cognoscantur: naturalibus enim na-

\*) Germar Zeitsch. f. d. Entomologie, II. p. 210.



turales, artificialibus artificiales sectiones definiuntur: characteres autem quinam naturales et quinam artificiales sint, non subitaria observatione sed intimo studio perspiciuntur, praesertim cum saepe accadat, ut notas easdem, quibus in aliis divisionibus determinandis utimur certis et stabilibus, in aliis fallaces omnino et variabiles reperiamus, et ne proximae quidem in quibusdam generibus species earum inter se conveniant partium structura, quae in aliis constantes praebent differentias. Qui iisdem sectionibus naturalibus omnium partium structuram eandem esse volunt, non rigidas figuras geometricas nos tractae obliviscuntur, sed corpora animalium, quae alma natura ad vitam procreavit et conformavit.

Sectiones huius familiae principales a Latreillio conditas conservare non potui, quippe quas non solum magis artificiales quam naturales, sed vacillantibus etiam notis circumscriptas reperi. Nova dispositione usus tribus et subtribus proposui, quarum plurimae generibus Gravenhorstianis respondent, et quas generum primariorum nominibus designavi. Staphylini genuini certe huius familiae principes sunt, et principium tenuissent, nisi ab Aleocharinis adscendendo, et ad Omalinos et Proteininos descendendo tribuum ordo naturalis minus interruptus esse videatur. Non omnes enim natura divisiones alteram alteri subiunxit, sed plurimas iuxta posuit, omnes autem vario intricavit affinitatum nexu: unde cum non possimus, quin in corporibus naturalibus enumerandis et describendis seriem simplicem evolva- mus, ordo omnino naturalis nostris in libris constitui nequit.

Genera nova pauca addidi, multa a variis auctoribus proposita suppressi, reliqua recepi, omnia scrupulose examinavi, et speciebus mihi cognitis cunctis respectis descripsi, ne specie singula pro generis typo usus characteres adhibeam, qui reliquis non congruant. Saepius antennarum formam, imprimis articulorum proportionem, nonnunquam palporum etiam et pedum structuram variam in variis eiusdem generis speciebus reperi, et potius inter speciei quam inter generis notas recensui. Oris structuram licet saepenumero non sine difficultate perspiciendam quam plurimis in speciebus inquirendam nunquam neglexi, et partes eius interiores omnium fere generum delineavi, quo facilius differentiae, quae plerumque sat insignes et semper magni momenti sunt, recognoscantur. Observationibus his instituendis microscopii compositi mei Schiekiani lenticulis magnitudinem naturalem centies augmentibus usus sum.

Describendis speciebus id praecipue agendum esse arbitror, ut notae differentiaeque speciales auctori perspectae sint, ut a lectore percipiantur: novi enim descriptiones, quae verbis uberrimae omnia



continere videntur nisi notas distinctivas, quibus neglectis species descriptae non lectori solum sed auctori etiam ipsi dubiae remanent, et specierum cognitio saepe magis perturbatur quam illustratur. Verborum copiam descriptiones nequaquam distinctiores reddere, Linnaei, Fabriciique et Illigeri exemplo liquet, brevitate vero neminem praecellere video Sayum Americanum, qui descriptiones adeo edidit concisas, ut diagnoseos volumen vix superent, nihilominus adeo lucidas ut vix speciem quandam ab illo exhibitam unquam invenies dubiam. Sunt autem auctores quoque plurimi, simili describendi modo usi, illis tamen viris ingenio impares, qui descriptionibus abbreviatis species nonnisi dubias et obscuras provulgant. Tanto magis igitur evitandum mihi putavi, quod mancae quam quod verbosae evadant descriptiones, quanto difficiliore et aegrius distinguendae plurimae Staphylinorum species apparent.

Quas enumeravi species omnes ipse examinavi, exceptis nonnullis mihi invisis, ubi de loco, quo inserendae sint, minus haesi, et quas auctorum descriptionibus adhibitis addidi: reliquas ab auctoribus nimis mance descriptas et nimis dubias omnino omisi. Gravenhorstianos typos plurimos in Museo Regio asservatos, nonnullos ab auctore ipso transmissos lustravi: specierum Gyllenhalianarum plerarumque specimina ab auctore et a cel. Schönherro missa, a Schüppelio et Germaro communicata, vidi, unde gravissimorum horum auctorum synonymia vix unquam dubia remansit. Staphylinos quoque a reliquis auctoribus descriptos plerosque cognoscere mihi succesit. Species autem Anglicae numerosae a Stephensio editae mihi haud extricatae sunt: libro illius de Entomologia Brittanica petenti mihi a cel. D. von Heyden, Consule Francofurthensi, benigne communicato et usui longinquo commisso, ut species pro novis descriptas plurimas cognoscerem summum quidem studium adhibui, quo saepenumero iterato, semper autem pari modo frustrato, tandem non sine taedio, licet invitatus ab hoc opere decessi, nihil persecutus, nisi quod genera, quibus Angli utuntur, recenserem, et specierum icones ab elegantissimo Westwoodio pictas allegarem. Videant perinde, qui cognoscendis typis Stephensianis prosperantur, ut speciebus, quascunque ab illo prius descriptas reperiant, nomina priora scientiae e legibus restituant: vocabula nonnulla a me proposita peritura fore parum refert: observatis scientiam auxisse iuvat.

Studio meo summa Dom. Prof. Klugii benevolentia commissae erant opes Musei Regii Berolinensis, collectione Hellwig-Hoffmanns-eggiana olim locupletissima et celeberrima conditi, collectione Knöchiana etc. aucti, plurimorum peregrinatorum collectis stipati, cui Dom.

Schüppel generose totam suam collectionem, speciebus Europaeis certe ditissimam, scientiae in usum tradidit, et cui ipse omnia addidi, quaecunque olim iuvenili fervore collegi. D. Prof. Germar, D. Dr. Aubé, D. Chevrolat collectionum suarum Staphylinos cunctos mihi transmiserunt, D. Prof. Gené Staphylinos in Sardinia sibi lectos cunctos communicavit; D. Prof. Germar magis e scientiae quam collectionis suae studio omnes huius familiae, quae singulae etiam ei erant, species liberaliter donavit, ut typi quam plurimi in Museo Berolinensi conserventur. D. D. Prof. Gené et Dr. Aubé omnia quae ultra dupla eiusdem speciei individua in eorum collectionibus invenerim, humaniter concederunt, Dom. Westermann omnes, quas duplas, multas etiam quas singulas habuit species, comiter dedit; multi praeterea multa contulerunt, unde successit, ut non solum uberrimam speciminum copiam examinandi et insignem specierum numerum describendi facultatem assecutus sim, sed quo etiam factum est, ut typi quam plurimi in Museo Regio Berolinensi collecti sint. Species omnes, quibus describendis speciminibus hoc in Museo conservatis usus sum, asterisco notavi.

Opere iam finito et integro edendo nihil restat, nisi quod gratias agam publicas, Vobis, D. D. Klug et Schüppel, quorum consilia mihi nunquam defuerunt, et Vobis omnibus, qui ex amicitia et scientiae studio laboribus meis subvenire benevole voluistis.

Berol. d. III. ante Cal. Aug. MDCCCXL.



## LIBRI CITATI.

---

- Act. Acad. Caes. Leop. Carol. Nat. Cur.*: Nova Acta Physico-Medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosorum.
- Act. Acad. nat. Scienc. Philadelph.*: Journal of the Academy of natural Sciences of Philadelphía.
- Act. Holm.*: Kongl. Svenska Vetenskaps Academiens Handlingar.
- Act. nov. Halens.*: Neue Schriften der naturforschenden Gesellschaft zu Halle.
- Act. Nidr.*: Det Trondhjemske og Norske Videnskabers Selskabs Skrifter. Kiöbenh.
- Act. Reg. Scienc. Acad. Taurin.*: Memorialia Regiae Scienciarum Academiae Taurinensis.
- Ahrens. Faun. Ins. Europ.*: A. Ahrensii Fauna Insectorum Europae. Hal. 1812. etc. (Cont. Germar.)
- Annal. Scienc. nat.*: Annales des Sciences naturelles. Paris.
- Beck Beitr.*: Beck, Beiträge zur bairischen Insectenfaune, oder Beschreibung und Abbildung neuentdeckter Käfer, etc. Augsburg 1817.
- Block Ins. vall. Plauens.* cont. in: Becker, der Plauensche Grund bei Dresden. Nürnberg. 1799.
- Boisd. Faun. Ent. Ocean.*: Boisduval, Faune Entomologique de l'Océanie. Par. 1835.
- Boisd. et Lacord. Faun. Ent. Paris.*: Boisduval et Lacordaire, Faune Entomologique des environs de Paris. Par. 1835.
- Bouché Hist. nat. Ins.*: Bouché, Naturgeschichte der Insecten, besonders in Hinsicht ihrer ersten Zustände als Larven und Puppen. Berl. 1834.
- Brullé Hist. nat. Ins.*: Audouin et Brullé, Histoire naturelle des Insectes. Paris 1834. etc.
- Bull. d. l. Soc. Imp. d. Moskou*: Bulletin de la Société Imperiale des Naturalistes de Moskou.
- Cederh. Faun. Ingr.*: Cederhjelm, Faunae Ingriae Prodrömus, exhibens methodicam descriptionem Insectorum Agri Petropolensis. Lips. 1798.
- Charp. Hor. Ent.*: Charpentier, Horae Entomologicae. Wratislav. 1825.
- Creutz. Ent. Vers.*: Creutzer Entomologische Versuche. Wien 1799.
- Curtis Britt. Ent.*: Curtis, Brittish Entomology, being illustrations and descriptions of the genera of Insects found in Great Britain and Ireland. Lond. 1824. etc.
- Dalm. Anal. Ent.*: Dalman, Analecta Entomologica. Holm. 1823.
- Degeer Ins.*: Degeer, Memoires pour servir a l'histoire des insectes. Stockholm 1752. etc.
- Dej. Cat.*: Catalogue des Coléoptères de la collection de M. le Comte Dejean. Paris 1837.
- Don. Britt. Ins.*: Donovan, the Natural History of Brittish Insects. Lond. 1792—1816.
- Encycl. Meth.*: Encyclopedie Methodique, Histoire naturelle des Insectes. Paris 1782—1824.

- Ent. Mag.*: The Entomological Magazine. Lond. 1833 — 38.  
*Er. Col. March.*: Erichson, die Käfer der Mark Brandenburg. Berl. 1837 — 39.  
*Fab. Syst. Ent.*: Fabricii Systema Entomologiae. Flensb. et Lips. 1775.  
*Fab. Gen. Ins.*: Fabricii Genera Insectorum. Chilon. 1776.  
*Fab. It. Norvag.*: Fabricius, Reise nach Norwegen. Hamb. 1779.  
*Fab. Spec. Ins.*: Fabricii Species Insectorum. Hamb. et Kilon. 1781.  
*Fab. Mant. Ins.*: Fabricii Mantissa Insectorum. Hafn. 1787.  
*Fab. Ent. Syst.*: Fabricii Entomologia Systematica emendata et aucta. Hafn. 1792 — 98.  
*Fab. Syst. El.*: Fabricii Systema Eleutheratorum. Kil. 1801.  
*Fourcr. Ent. Par.*: Fourcroy, Entomologia Parisiensis s. Catalogus Ins., quae in agro Parisiensi reperiuntur, secundum methodum Geoffroyanam etc. Par. 1785.  
*Frisch Ins. Germ.*: Frisch, Beschreibung von allerlei Insecten in Deutschland. Berlin 1766.  
*Fuessl. Verz. Schweiz. Ins.*: Fuessly, Verzeichniss der ihm bekannten Schweizerischen Insecten. Zürich u. Winterth. 1775.  
*Geoffr. Hist. Ins.*: Geoffroy, Histoire abrégée des Insectes. Paris 1764.  
*Germ. Faun. Ins. Europ.*: v. Abrens.  
*Germ. Mag.*: Germar, Magazin der Entomologie. Halle 1813 — 21.  
*Germ. Spec. Ins.*: Germar, Insectorum Species novae aut minus cognitae. Hal. 1824.  
*Goeze Ent. Beitr.*: Goeze, Entomologische Beiträge zu des Ritters Linné zwölften Ausgabe des Natursystems. Leipz. 1777.  
*Grav. Micr.*: Gravenhorst, Microptera Brunsvicensia etc. Brunsv. 1802.  
*Grav. Mon.*: Gravenhorst, Monographia Micropterorum. Gotting. 1806.  
*Gray An. Kingd.*: The Animal Kingdom, arrang. by Cuvier, with supplementary additions to each order, by Griffith, Vol. XIV., The Insecta, by Gray. London 1832.  
*Guér. Icon. Regn. an.*: Guérin, Iconographie du Règne Animal de Mr. le B. Cuvier. Paris.  
*Guér. Mag. Zool.*: Guér. Magasin de Zoologie.  
*Guér. Voy. d. l. Coquille*: Voyage autour du monde, exécuté p. o. d. r. sur la Corvette d. s. M. la Coquille p. l. a. 1822—25. par Duperrey. Zoologie Atl. 1826, Texte: Crust. Arachn. et Ins. par. Guérin Meneville. 1838.  
*Gyll. Ins. Suec.*: Gyllenhal, Insecta Suecica. Scar. 1808 — 28.  
*Harrer Besch. d. Ins.*: Harrer, Beschreibung derjenigen Insecten, welche Herr D. J. Schäffer in 280 ausgemalten Kupfertafeln herausgegeben hat. Regenb. 1784.  
*Heer Obs. Ent.*: Heer Observationes Entomologicae. Turic. 1836.  
*Herbst Arch.*: Archiv der Insectengeschichte, herausgegeben von J. C. Fuessly. Zürich u. Winterth. 1781.  
*Herbst Col.*: Natursystem aller bekannten in- und ausländischen Insecten von C. G. Jablonsky, fortgesetzt von J. F. W. Herbst. Berl. 1789 — 1801.  
*Illig. Käf. Pr.*: Illiger, Verzeichniss der Käfer Preussens. Halle 1798.  
*Illig. Mag.*: Illiger, Magazin für Insectenkunde. Braunschw. 1801 — 7.  
*Isis*: Isis, Encyclopädische Zeitschrift, vorzüglich für Naturgesch., vergl. Anatomie und Physiologie, von Oken. Leipz.  
*Kirby Faun. Boreali-Amer.*: Richardson Fauna Boreali-Americana, or the Zoology of the Northern parts of America, etc. — Part IV. The Insects, by W. Kirby. Norwich 1837.  
*Kirby et Spence Introd.*: Kirby and Spence, Introduction to Entomology. London 1818 — 1827.  
*Klug Col. Madag.*: Klug Bericht über eine auf Madagascar veranstaltete Sammlung von Insecten aus der Ordnung Coleoptera. Berl. 1834. (Physicalische Abhandl. der Königl. Acad. der Wissensch. zu Berlin, aus d. J. 1832.)  
*Klug Jahrb.*: Klug Jahrbücher der Insectenkunde. Berl. 1834.  
*Laporte Etud. Ent.*: Laporte Etudes Entomologiques. Paris 1835.  
*Latreille Précis d. caract.*: Latreille Précis des caractères génériques des Insectes. Briv. 1796.  
*Latreille Hist. nat. Crust. et Ins.*: Latreille Histoire naturelle générale et particulière des Crustacés et des Insectes. Paris 1802 — 5.  
*Latreille Gen. Crust. et Ins.*: Latreille Genera Crustaceorum et Insectorum. Paris et Argentorat. 1806 — 7.

- Latreille Regn. An.*: Cuvier Regne Animal, distribué d'après son organisation. Nouv. Ed.: Crustac. Arachn. et Ins. par Latreille. Paris 1829.
- Ledebour It.*: Ledebour Reise durch das Altai-Gebirge und die Songarische Kirgisensteppe. Berl. 1829.
- Linn. Faun. Suec.*: Linné Fauna Suecica, Editio altera, auctior. Stockholmiae 1761.
- Linn. Iter Gotl.*: Linné Oeländska och Gothländska Resa förättad år 1741. Stockh. och Upsal. 1745.
- Linn. Syst. Nat.*: Linné Systema Naturae per Regna tria naturae, Ed. 12. Holm 1767.
- Linn. Transact.*: Transactions of the Linnean Society of London.
- Mag. Zool. and Bot.*: Magazine of Zoology and Botany, cond. by Jardine, Selby and Johnston. Lond. Edinb. and Dublin 1836 etc.
- Mannerh. Brachelytr.*: Mannerh. Précis d'un nouvel arrangement de la famille des Brachélytres. St. Pétersb. 1830. (Mém. présentés à l'Acad. Imp. des scienc. de St. Pétersbourg par divers savans, T. I.)
- Marsh. Ent. Britt.*: Marsham Entomologia Britannica. London 1802.
- Mart. Engl. Ent.*: Martyn, the English Entomologist, exhibiting all the Coleopterous Insects found in England. London 1792.
- Ménétr. Catal.*: Ménétries, Catalogue raisonnée des Objects de Zoologie recueillis dans un voyage au Caucase etc. St. Pétersbourg 1832.
- Meyen It.*: Meyen, Reise um die Erde. Dritter Theil. Zoologischer Bericht. Breslau u. Bonn 1834.
- Müll. Faun. Frid.*: Müller, Fauna Insectorum Fridrichsdalina. Hafn. et Lips. 1764.
- Müll. Prodr.*: Müller, Zoologiae Danicae Prodomus. Hafn. 1776.
- Nicol. Col. Hal.*: Nicolai, Dissertatio sistens Coleopterorum species agri Halensis. Hal. 1822.
- Nordm. Symb.*: Nordmann, Symbolae ad Monographiam Staphylinorum. Petrop. 1837. (Mém. présentés à l'Acad. Imp. des sciences de St. Pétersbourg. T. IV.)
- Nov. Ann. Mus. Hist. nat.*: Nouvelles Annales du Muséum d'histoire naturelle. Par. 1832 etc.
- Nouv. Mém. d. l. Soc. Imp. des nat. de Mosk.*: Nouveaux mémoires de la Société Imperiale des Naturalistes de Moskou.
- Oliv. Ent.*: Olivier, Entomologie ou Histoire naturelle des Insectes. Par. 1789—1808.
- D'Orbign. It.*: D'Orbigny Voyage dans l'Amerique méridionale (le Brésil, la républ. orient. de l'Uruguay, la Patagonie, la républ. Argentine, du Chili, du Pérou, de Bolivia) exécuté dans le cours des années 1826—33. Par.
- Pall. It.*: Pallas, Reise durch verschiedene Provinzen des Russischen Reichs. St. Petersburg 1772—76.
- Panz. Ent. Germ.*: Panzer, Entomologia Germanica, exhibens Insecta per Germaniam Indigena. Norimb. 1795.
- Panz. Faun. Germ.*: Panzer, Faunae Insectorum German. Initia. Norimb. 1789—1810.
- Payk. Faun. Suec.*: Paykull, Fauna Suecica. Upsal. 1800.
- Payk. Mon. Staph.*: Paykull, Monographia Staphylinorum Sueciae Upsal. 1789.
- Payk. Mon. Car.*: Paykull, Monographia Caraborum Sueciae. Upsal. 1790.
- Payk. Mon. Curc.*: Paykull, Monographia Curculionum Sueciae, Upsal. 1792.
- Payk. Mon. Hist.*: Paykull Monographia Histeroidum. Upsal. 1811.
- Percheron Gén. Ins.*: Guérin et Percheron Genera des Insectes. 1835.
- Perty Delect. An. art.*: Delectus Animalium articulorum quae in itinere per Brasiliam annis 1817—20 collegerunt Spix et Martius, dig. descr. ping. cur. Perty, ed. Martius. Monach. 1830.
- Petiver Op.*: Petiveri Opera, historiam naturalem spectantia. Lond. 1767.
- Preysl. Böhm. Ins.*: Preyssler, Verzeichniss Böhmischer Insecten. Prag 1790.
- Ratzeburg Forstins.*: Ratzeburg, die Forstinsecten, oder Abbildung und Beschreibung der in den Wäldern Preussens und der Nachbarstaaten als schädlich oder nützlich bekannt gewordenen Insecten. Berlin 1837.
- Rossi Faun. Etr.*: Fauna Etrusca, sistens Insecta, quae in Provinciis Florentina et Pisana praesertim collegit P. Rossius. Liburn. 1790. Eiusd. Mantissa Insect., exhibens species nuper in Etruria collectas. Pis. 1792—94.
- Ed. Hellw.*: Rossii Fauna Etrusca, iterum edita et annotationibus perpetuis aucta a Hellwig. Helmst. 1795.
- Royle Himalaja*: Forbes Royle, Illustrations of the Botany and other Branches of the natural history of the Himalajan Mountains and of the Flora of Cashmere. Lond. 1835.



- Runde Brachelytr. Hal.*: Runde Brachelytrorum species agri Halensis, Hal. 1835.  
*Sahlb. Ins. Fenn.*: Sahlberg, Insecta Fennica, dissertationibus academicis ed. Hel-singf. 1817 etc.  
*Schäff. Abhandl.*: Schäffers Abhandlungen von Insecten. Regenb. 1764 — 79.  
*Schäff. Elem. Ent.*: Schäfferi Elementa Entomologica. Ratisbon. 1766.  
*Schäff. Icon.*: Schäfferi Icones Insectorum circa Ratisbonam indigenorum. Re-gensb. 1769.  
*Schneid. Mag.*: Schneider, Neuestes Magazin für Liebhaber der Entomologie. Stral-sund 1791—94.  
*Schrank Beitr. z Naturgesch.*: Schrank Beiträge zur Naturgeschichte. Leipz. 1776.  
*Schrank Enum. Ins. Austr.*: Schrank Enumeratio Insectorum Austriae. Aug. Vin-del. 1781.  
*Schrank Faun. Boic.*: Schrank Fauna Boica, durchgedachte Geschichte der in Bajern einheimischen und zahmen Thiere. Nürnberg. 1798.  
*Scop. Ent. Carn.*: Scopoli Entomologia Carniolica. Vindabon, 1763.  
*Scop. Ann. hist. nat.*: Scopoli Annus historico-naturalis I—V. Lips. 1769 — 72.  
*Shaw Gener. Zool.*: Shaw General Zoology or Systematic Natural History. Lond. 1806.  
*Steph. Illustr.*: Stephens Illustrations of British Entomology, Mandibulata. Lond. 1827 etc.  
*Sturm Catal.*: Sturm, Catalog meiner Insectensammlung. I. Theil. Käfer. Nürnberg. 1826.  
*Sulz. Kennz.*: Sulzer, die Kennzeichen der Insecten. Zürich 1761.  
*Sulz. Hist. Ins.*: Sulzer, abgekürzte Geschichte der Insecten. Winterth. 1776.  
*Transact. Amer. Philosoph. Soc.*: Transactions of the American Philosophical Society, held at Philadelphia.  
*Transact. Ent. Soc.*: The Transactions of the Entomological Society of London. Lond. 1834 etc.  
*Transact. Hist. Nat. Soc. Hartf.*: Transactions of the Natural History Society of Hartford. Hartf. 1836.  
*De Vill. Ent.*: C. Linnaei Entomologia, cur. et aug. De Villers. Lugd. 1789.  
*Walk. Faun. Paris.*: Walkenaer Faune Parisienne, Insectes, ou Histoire abrégée des Insects des environs de Paris. Par. 1802.  
*Waltl It. Andal.*: Waltl, Reise durch Tyrol, Oberitalien und Piemont nach dem südlichen Spanien. Passau 1835.  
*Weber et Mohr Archiv.*: Weber und Mohr, Archiv für systematische Naturgeschichte. Leipzig 1804.  
*Weber Beitr.*: Weber, Beiträge zur Naturkunde. Kiel 1805 — 10.  
*Wiedem. Zool. Mag.*: Wiedemann, zoologisches Magazin. Kiel u. Altona 1817 — 23.  
*Zetterst. Faun. Lappon.*: Zetterstedt Fauna Insectorum Lapponiae. Hamm 1828.  
*Zetterst. Ins. Lappon.*: Zetterstedt Insecta Lapponica. Lips. 1838.  
*Zool. Journ.*: The Zoological Journal. Edited by Vigors. Lond. 1825 etc.

## **S t a p h y l i n i .**

---

### **D e c h a r a c t e r e   n a t u r a l i .**

Magna haec familia, notis licet ab antennarum insertione, figura et articulorum numero, ab oris partibus, a tarsorum etiam articulorum numero derivatis omnino instabilibus, tamen, quod in omnibus familiis naturalibus fieri solet, a ceteris distinguitur constante caractere: abdomine segmentis sex vel septem distinctis, liberis, omnibus corneis, aut toto nudo, aut basi ab elytris obtecto, his abbreviatis, sutura commissis, alas complicatas omnino obtegentibus. Priores plerique auctores Staphylinorum familiam coleopteris abbreviatis, quorum etiam causa a Latreillio Brachelytrorum, a Gravenhorstio Micropteriorum nomine signata est, distinxerunt, elytra abbreviata autem in plerisque aliis Coleopterorum familiis obveniunt, quae vero abdominis segmenta, si posteriora sola nuda, anteriora autem ab elytris obtecta sunt, illa sola cornea, haec membranea habent (ut Silphae, Histeri, Nitidulae etc.), cum in Staphylinis segmenta dorsalia anteriora, etiamsi nonnunquam elytrorum postrema pars iis imposita est, pariter cornea atque in illis reperiamus, qui elytris pectus non superantibus, abdomine toto nudo instructi sunt. Alia nonnulla coleoptera, quorum elytra adeo abbreviata sunt, ut abdomen iis sit omnino nudum et segmentis dorsalibus omnibus corneis instructum, a Staphylinis aut alis sub elytra non recipiendis (*Molorchus*, *Atractocerus*), aut, si alis carent, elytris dehiscentibus (*Meloe*) divergunt: Staphylinis enim omnibus, apteris etiam, coleoptera sutura commissa sunt. Sola coleopterorum familia elytris abbreviatis, pectus haud excedentibus, alis sub elytris complicandis, abdomine toto nudo et segmentis omnibus dorsalibus corneis cum Staphylinis congruens, *Pselaphi*, praeter plures alios caracteres, abdomine segmentis distinctis tantum quinque compositis differt, in Staphylinis enim segmenta sex semper distincta, septimum etiam saepius conspicuum. Quaedam huius familiae genera (*Anthophagus*, *Lestea*) Caraborum quorundam habitum prae se ferunt, praeter alias autem graviore notas abdominis etiam structura distinguuntur, cum *Carabis* segmenta tria anteriora connata, illis omnia sint libera.

Staphylini a Latreillio Coleopterorum sectioni Pentamerorum adnumerantur, et a Gravenhorstio etiam praeter „elytra dimidiata” tarsis 5-articulatis a reliquis Coleopteris facillime distinguendi laudantur. At studii huius familiae cultor excellens, Com. Mannerheim (Brachelytr. p. 5.) Oxytelorum tarsos triarticulatos esse observavit. Hanc Latreille quidem observationem, quam difficillimam habet, non satis exactam fuisse suspicatus est, (Nouv. Annal. du Mus. d. hist. nat. I. p. 79.) et Oxytelorum tarsos non infra 4 articulis compositos esse contendit, (Regn. anim. ed. II. IV. p. 438.), attamen ex meis observationibus non solum Oxyteli, sed quaedam etiam alia genera revera sunt trimera, alia tetramera, alia quodammodo heteromera, ut tarsorum articulorum numerus, in aliis plurimis familiis constantissimus, in hac eo minus gravioris sit momenti, ac genera simillima et proxima saepius articulorum tarsalium numero differe videamus. Plurimi Staphylini revera tarsis omnibus 5-articulatis instructi sunt. Tarsis posterioribus 5-, anticis 4-articulatis plura Aleocharinorum genera distinguuntur: Myrmedonia, Antalia, Falagria, Bolitochara, Tachyusa, Homalota, Phytosus, Gyrophæna, Peliusa, Placusa, Euryusa, Dinarda, Lomechusa, Silusa, Pronomæa, Myllaena, quae hoc modo Heteromera quasi inversa vocanda sunt. Eandem, quam in his Aleocharinis, articulorum dispositionem observamus in Malachiis quibusdam, antennarum, articulis nonnullis in maribus difformiter dilatatis, e quibus species nonnullae a Fabricio Pausso adiectae, et ad quos genera Laius Guérin et Megadenterus Westwood referenda sunt. Apud hos autem maribus solis tarsi antici 4-articulati, feminae 5-articulati sunt, sicuti apud Mycetophagos tarsos anticos 3-articulatos in maribus, in feminis cum posticoribus 4 articulatos esse monuit Müller (Illig. Mag. IV. p. 206.); in tota Staphylinorum familia vero tarsi in utroque sexu eodem articulorum numero gaudent. Tarsi revera heteromeri in hac familia nondum innotuerunt. — Tarsi 4-articulati obveniunt inter Aleocharinos in Hygronoma, Oligota, Diglossa, inter Tachyporinos in Hypocypto et Tangynatho, inter Steninos in Euæstheto. Tarsis 3-articulatis sunt Oxytelini genuini et inter Proteininos Glyptoma et Micropeplus instructi. Systemati Latreilliano hanc familiam omnino adversariam esse patet.

#### De affinitatibus.

Affinitates Staphylinorum cum aliis familiis naturalibus variae sunt atque insequentibus accuratius illustrandae. Caraborum quorundam (Dromiorum) habitum offerunt genera Anthophagus et Lestæa, ut non solum species quaedam a Linnaeo Staphylini caraboidis nomine signata sit, sed plures etiam auctores, Fabricius et Illiger ipse, Caraborum generi nonnullas huius generis species adscripserint. Quamvis omnino victu carnivoro cum Carabis Staphylini conveniunt, pluribus notis tamen ab iis distinguuntur, et praeter oris fabricam, qua Carabi ab omnibus Coleopteris terrestribus satis distant, genera illa abdominis structura supra memorata, nec non fronte ocellis duobus instructa plurimisque levioribus notis habitualibus differunt.

Silphis Staphylini summa affinitate iunguntur, ut facile pro Silphis



elytris fortius abbreviatis haberi possint, non modo antennis saepius apicem versus leviter incrassatis, sed abdomine etiam in utraque familia segmentis sex composito, et evenit, ut a Fabricio, Illigero pluribusque aliis auctoribus plures Omalinorum species, quae corpore latiusculo et abdomine partim ab elytris oblecto magis earum habitum prae se ferunt, pro meris Silphis descriptae sint. Silphis autem abdominis segmenta dorsalia anteriora membranacea sunt, et abdominis segmentum primum minus distinctum. Praeterea Silphis genuinis Staphylini genuini, Catopibus Tachyporini propiores mihi videntur.

Pselaphi elytris pectus haud excedentibus et abdomine nudo, segmentis omnibus corneis composito Staphylinis sunt proximi, neque victu multum ab iis discedunt, ut Euplectos, qui corpore elongato, depressiusculo Staphylinorum habitum praebent, sub nomine Stenostheti in primo collectionis suae Catalogo cel. Com. Dejean inter Staphylinos enumeraverit. Sunt etiam, qui Pselaphos omnino Staphylinis adiungunt. (Laporte Etud. Ent. p. 137.) Abdomen illis autem segmentis tantummodo 5 compositum, et praeter multas alias notas habituales praecipue maxillarum mala exteriora falciformi a Staphylinis differunt, licet iis affines et Aleocharinis imprimis propinqui videantur.

Scaphidia elytris abbreviatis et abdomine prominulo quoque Staphylinis similia atque abdomine acuminato, segmento primo maiore, reliqua recipiente et elytris longioribus Tachyporinis quibusdam habitu certe affinia sunt, et Hypocryptum longicornem etiam Paykull et Gyllenhal inter Scaphidia posuerunt. Discedunt Scaphidia a Staphylinis abdomine distinctius 5-annulato, segmentis superioribus, quae ab elytris conduntur, haud corneis, coxis posticis distantibus, etc.

Nitidulae multae elytris magis minusve abbreviatis et abdominis pluribus segmentis nudis Staphylinorum habitum saepius prae se ferunt, ut non solum Nitidulas quasdam inter Stenos (St. spissicornis, conicus, Ficus) collocaverit Fabricius, sed recentioribus etiam temporibus Com. Laporte Nitidularum quandam formam, statura magis elongata Ipum, elytris autem fortiter abbreviatis instructam, nomine Cillaei pro Omalinorum genere descripserit. (Etud. Ent. p. 133.) E Staphylinis praecipue Omalini corporis habitu et victu pluribus Nitidulis affines videntur, propiores autem iis sunt coxarum anticarum insertionem, ocellis nullis et antennis distinctius apice incrassatis Proteinini, ut Micropeplus a Latreille ipso et principio Proteinus Nitidulis adiectae sint. Proteinus certe ad Nitidularum, imprimis ad Cercorum affinitatem distincte se vergit, abdomine licet segmentis sex composito, et coxis posterioribus approximatis ab iis discedit. Micropeplus autem, quamquam plerisque ab auctoribus Nitidulis adiungitur, tamen antennae in scrobiculo in inferiore thoracis facie sito recipiendis, et coxis intermediis fortiter distantibus potius Histeris appropinquatur, et sculptura etiam corporis quandam Onthophilorum similitudinem ei praebet, abdomine segmentis sex composito autem, et coxis posticis approximatis pluribusque aliis notis Staphylinis adiungitur, licet antennae capitulatis iis quodammodo alienus videtur. A Staphylinis Nitidulae differunt praeterea maxillarum mala singula, Histeri abdominis structura, de qua conf. quae Klug Jahrb. d. Insectenk. I. p. 84 et Col. March. I. p. 649 monui.

Xylophagis Latr. nonnulla Staphylinorum genera victu non solum, sed corporis etiam habitu affinia sunt. Glyptomati Monotomae habitus est, et

Piestini omnino Cucuios elytris abbreviatis simulant, ut Piesti et Leptochiri speciem quandam Fabricius ipse inter Cucuios descripserit. Magis vero, praecipue thoracis etiam forma *Isomalus* Cucuiis minoribus similis videtur. Dom. Laporte inter Omalinos novum genus induxit (Etud. Ent. p. 135.) Ino vocatum, quod licet elytris abbreviatis instructum sit, tamen revera ad Cucuios pertinet, et a Hemipeplo Latr. haud diversum videtur. Abdomine segmentis 5 composito Cucuii a Staphylinis, et elytris pectus non excedentibus Piestini a Cucuiis etiam brachypteris distinguuntur.

### De corporis structura exteriori.

Corpus plurimis elongatum, lineare, plerisque subdepressum (*Staphylinus*), rarius complanatum, (*Hygromoma*, *Holisis*, *Achenium*, *Isomalus*), nonnunquam cylindricum (*Procirrus*, *Stenus*, *Osorius*), aliis convexum, aut utrinque aut saltem apicem versus attenuatum (*Tachyporini*), aliis breviusculum, interdum subquadratum seu oblongo-subquadratum (*Omaliini* *Proteinini*que plures), nonnunquam totum tomentosum (*Staphylini* quidam), aut totum pube subtili brevi depressa sericante vestitum (*Conurus*), rarius totum glaberrimum (*Oxyporus*), plerumque tenuiter pubescens, capite thoraceque saepius pube rariore subtiliore vestitis, nonnunquam glaberrimis, et nonnisi versus latera parce pilosellis (*Philonthus*). Abdomen quoque praeter pubescentiam subtiliorem plerumque parce pilosum, praecipue latera versus.

Caput thoraci aut insertum, basi saepius magis minusve in colli formam coarctatum, rarius nullo modo constrictum (*Tachyporini*), aut exsertum, thoracis apici attenuato affixum, (*Autalia*, *Falagria*, *Stilicus*, *Echiaster*, *Ophites*), rarius deflexum (*Tachyporini*), vel inflexum (*Aleochara*), plerumque porrectum, fronte superiore rarius prona aut retusa, antrorsum spectante (*Staphylinus versicolor*). Caput inferius planum, superius saepius magis minusve convexum, fronte integra, clypeo distincto nullo, spatio eius autem nonnunquam impresso et linea impressa circumdato (*Oxytelus*), nonnunquam cornuta (*Piestus*, *Ziroporus*, *Bledius*, *Platystethus*), aut margine dilatato ampliata (*Phloeobium*).

Oculi laterales, rotundati ovate, saepius minuti et minus prominuli, interdum magni, prominentes, (*Stenus*, *Megalops*), ocellis plerumque minutis et parum convexis, rarius maioribus et fere hemisphaericis (*Oxytelini*), semiautem hexagonis compositi.

Ocelli frontales duo omnibus Omalinis, singulum *Phloeobio* clypeato, reliquis *Staphylinis* nulli. Errore quodam *Euaesthetum* ocellis instructum esse affert Kirby (Introduct. III. p. 506.)

Labrum sub frontis margine anteriore insertum, porrectum, plerumque transversum, rarius latitudine longius (*Boletobius atricapillus* etc.), aut adeo amplum, ut reliquas oris partes obtegat (*Stilicus*), saepius brevissimum (*Pinophilus*), aliis integrum, (*Aleocharini*, *Tachyporini*), aliis bilobum, lobis rotundatis (*Staphylinus*, *Philonthus*, *Lathrobium*) vel divergentibus (*Oxyporus*, *Achenium*), aliis apice denticulatum (*Scytalinus*, *Lithocharis*, *Stilicus*, *Ophites*, *Sunius*, *Echiaster*), his totum corneum (*Aleocharini*, *Tachyporini* plerique etc.), illis corneum, limbo aut laterali et anteriore (*Staphylinus*, *Philonthus*, etc.), aut laterali solo membranaceo (*Xantholinus*, *Stilicus*, etc.),



pluribus simplex, ut reliquis coleopteris, multis etiam laciniiis membraneis auctum, aut in angulo utraque anteriore affixis, acuminatis, paraglossis similes (Oxytelini, Piestini), aut limbi membranei saepius bilobi ad instar sub labii margine anteriore prominulis (Bledius, Omalini, Proteinini). Limbus ille membranaceus, quem apud plurimos Staphylininos observamus, substantia cornea versus margines evanescente exoritur, a labro ipso nullo modo distinctus, laciniae haec membranae, quibus praecipue Oxytelini et Piestini insignes sunt, appendices peculiare videntur, e membrana, qua interior s. inferior labri facies vestitur, nascentes. Limbus membranaceus saepius supra fortiter pilosus setosusque, laciniae membranae margine ciliatae. Labrum totum membranaceum videtur in Tanygnatho. Insolitam formam labrum inducit in Megalope: spinis duabus acutis, sub frontis margine prominentibus, porrectis referri videtur.

Mandibulae corneae, acutae, plerumque falcatae, interdum rectae (Piestus), aliis muticae, aut breviores, haud prominentes (Aleocharini, Tachyporini, Omalini plerique etc.), aut elongatae (Oxyopus fam. III, Belonuchus, Oxyporus etc.), aliis plerisque medio vel ante apicem dentatis, dentibus vel simplicibus, duplicibus triplicibusve, vel bi- aut tricuspidibus, saepius inaequalibus, mandibula sinistra plerumque fortius dentata, iam breviores et validiores (Staphylinus, Philonthus), iam elongatae, tenues (Paederini, Pinophilini, Stenini), aliis interne fortiter dentatae, (Leptochirus), aliis externe ante apicem denticulo prominulo instructae (Piestus), ore clauso plerumque complicatae et haud prominentes, nonnunquam prominentes, decussatae (Oxyporus, Belonuchus), aut porrectae, rectae (Leptochirus, Piestus). Margo interior lacinia membranea, interne subtiliter ciliata auctus, plerumque mandibulae margini interiori connata, ita ut mandibulis breviusculis apex solus liber esse soleat, in tota autem Staphylininorum tribu libera, basi sola mandibulae radici affixa.

Maxillae cardine brevi stipiteque corneis, hoc in Piestinis nonnullis infra palporum basin dilatato. Mala utraque distincta. Mala interior stipitis margini interiori connata, plerumque membranea membranaceave, mutica, interne apicem versus barbata, basin versus ciliata, apice nonnunquam unco corneo armata (Dinarda, Proteinini et omnes Omalini), saepius simplici, rarius duplici (Lomechusa, Proteinus), quam plurimum rhombea; mala exteriore brevior, rarius elongata, mala exteriori aequali, cornea, interne apicem versus denticulata (Aleocharini nonnulli et Trichophyus), rarissime mala exteriore longior (Silusa). — Mala exterior stipitis apici inserta, plerumque segmentis duobus composita, inferiore aut corneo aut coriaceo, iam breviori, iam longiore, apicali aut membraneo aut membranaceo, hoc aut toto aut apice plerumque introrsum vergente barbato, illo semper glabro, malam interiorem plerumque superans, interdum linearis, saepius apicem versus leviter dilatata, margine exteriore nonnunquam ampliato, in Hypocrypto sinuato, interdum cum mala interiore elongata, tota cornea, semper mutica.

Palpi maxillares maxillae stipiti exterius ad apicem prope malam exteriorem inserti, 4-articulati, articulo primo quam plurimum minuto, brevi, Stenini solis elongato, secundo tertioque plerumque sat elongatis, hoc nonnunquam paulo breviori, interdum brevi (Megarthrus, Micropeplus), ubi autem articulus quartus subulatus est, maiore, magis minusve, praecipue apicem versus, incrassato, articulo quarto aut reliquis subaequali, apice leviter

acuminato, rarius truncato (Scariphaeus), vel securiformi (Astrapaeus, Taenodema, Palaminus, Oedichirus), vel transversim introrsum producto (Pinophilus), saepius minuto, subuliformi, (Aleocharini, Hypocyptus, Conurus, Tachyporus, Mycetoporus, Leptacinus, Holisus, Diochus, Paederini, Oxytelini genuini), vel minutissimo, non nisi oculo fortiter armato conspiciendo, quo palpi triarticulati videntur (Stilicus, Echiaster, Sunius et Stenini), nonnunquam pariter minimo, at obtuso (Doliceon, Paederus).

Labium segmentis 4, ut in reliquis insectis, compositum, nondum satis illustratis, in hac familia plerumque omnibus conspicuis et distinctis.

Segmentum primum est mentum, semper corneum, plerumque breve, transversum, apicem versus sensim angustatum, apice truncatum vel leviter lateque emarginatum, raro maius et reliquas labii partes omnino aut maxima e parte obtegens (Pronomaea, Micropeplus).

Segmentum secundum, innominatum, membranaceum coriaceumve, in reliquis coleopteris plerisque sub mento reconditum, in hac familia autem quam plurimum magis minusve exsertum, in nonnullis, ut videtur, retractile, basi menti apici latitudine subaequali, apicem versus sensim leniter angustatum, apice truncatum, in Oxyporo bidentatum. In hac familia haec pars a Fabricio pro ligula recensita est.

Segmentum tertium, in aliis etiam plurimis coleopteris distinctum et conspicuum, segmento secundo saepius paulo latius, vel apicem versus leviter dilatatum, coriaceum corneumve, palporum labialium stipites cylindricos, corneos, saepe sat distinctos continens. Eius apici ligula et paraglossae insertae sunt.

Segmentum quartum e ligula cum paraglossis constat. Ligula semper distincta, angusta, linearis, plerumque bifida, saepius elongata in Aleocharinis, minuta, brevis in Staphylininis, in utraque tribu reliquis labii partibus multo angustior, paraglossis distinctioribus, ligulam longe superantibus in Staphylininis, subaequalibus vel ea brevioribus vel plerumque omnino nullis in Aleocharinis, paraglossis et ligula membraneis, illis nonnunquam membranaceis vel coriaceis in Staphylininis nonnullis. In reliqua hac familia ligula latior, segmento labii tertio haud angustior, apice rarius truncata, plerumque medio magis minusve emarginata, vel incisione profundiore biloba, his lobis nonnunquam divergentibus (Leptochirus), nonnunquam spatio latiore distantibus, (Lithocharis, Stilicus, Echiaster, Sunius, Paederus,) paraglossis iam liberis, interdum ligulam excedentibus, nonnunquam ei aequalibus eique fere incumbenibus, iam ei connatis, ut sutura earum nonnisi ciliarum subtilium serie cognoscatur, iam omnino nullis (Boletobius, Mycetoporus, Platyprosopus, Oxyporus et Omalini omnes); ligula plerumque cum paraglossis membranea, rarius membranacea, nonnunquam cornea, paraglossis autem membraneis (Scopaeus, Megalops, Osorius, Holotrochus).

Palpi labiales labii segmento tertio inserti, plerisque ut reliquis coleopteris triarticulati, articulo tertio nonnunquam securiformi (Scytalinus, Staphylini et Ocypodes nonnulli, Astrapaeus, Euryporus), vel hemisphaerico (Staphyl. canaliculatus), vel lunato (Oxyporus), saepius minuto, subulato (Hypocyptus, Mycetoporus, Heterothops, Leptacinus, Holisus, Diochus, Paederini, Stenini), in plerisque filiformes, in Aleocharis etiam, in Conuro, Tachyporo, Phoeocharide et in Oxytelinis genuinis, qui palpis maxillaribus subuliformibus instructi sunt. Quadri-articulatos eos observavi in Aleochara,



biarticulatos in nonnullis Aleocharinorum generibus, neque in Trichophyo ultra duos articulos vidi. Palpis labialibus exarticulatis, setaceis nonnulla Aleocharinorum genera insignia sunt.

Antennae vel plerumque infra capitis marginem lateralem inter oculos et mandibulas, vel superne iam in frontis margine anteriore, intra mandibularum basin (Staphylinini), iam ad oculorum marginem interiorem (Aleocharini), iam in media fronte (Stenini) insertae, plerumque articulis 11, nonnunquam 10 (Oligota, Hypocyptus), vel 9 (Micropeplus) compositae, quam plurimum filiformes, vel apicem versus subincrassatae, nonnunquam articulis ultimis tribus praecedentibus paulo crassioribus, (Bledius, Acrognathus), interdum clavatae (Stenini, Megalops, Glyptoma), vel capitulatae (Micropeplus), paucis capillares, verticillae-pilosellae (Habrocerus, Trichophyus), plerisque rectae, aliis geniculatae, flagello antrorsum directo (Xantholinini genuini, Acylophorus, Cryptobium, Ophites), aut refractae, flagello retrorsum vergente (Oxytelini plurimi). Articulus primus magis minusve validior, et ubi antennae geniculatae sunt, saepius sat elongatus, secundus tertiusque sequentibus quam plurimum paulo maiores, praecipue longiores, hic plerumque illo magis elongatus, primo gracilior. Antennae interdum caput et thoracem longitudine excedunt, raro corporis fere longitudine (Piestus), saepius capite thoraceque breviores, nonnunquam capite vix longiores sunt.

Thorax sensu stricto seu prothorax forma magnopere varians, plerisque marginatus, aliis immarginatus (Lathrobium), vel cylindricus (Stenus), nonnunquam fortiter elongatus (Ophites), apice interdum, ubi caput omnino exsertum est, attenuatus, (Autalia, Falagria, Ophites, Stillicus, Echiaster, Sumius), nonnunquam utrinque oblique truncatus (Xantholinus, Lithocharis), plerumque truncatus, in multis etiam emarginatus, angulis anterioribus prominulis, caput amplectentibus (Tachyporini), basi nonnunquam coleopteris arcte applicatus vel ea amplectens, (Tachyporini), interdum etiam ab iis remotus, (Leptochirus), aliis rotundatus, aliis truncatus; infra prosterno triangulari, coxarum anticarum insertionem haud excedente, epimeris et episternis nullis, in plerisque spatio pone coxas anticas absque integumento corneo, nonnisi membrana clauso, stigmate prothoracico utrinque pone coxam anticam sito aut peritremate corneo distincto, nudo in Aleocharinis, Tachyporinis et Staphylinis Xantholininis et genuinis, vel processu membranaceo corneove triangulari oblecto in Staphylininis Oxyporinis, iam stigmate peritremate corneo nullo, sub integumento corneo pone coxas anticas introrsum producto occulto, in Paederinis, Oxytelinis, Phloeocharinis, Omalinis; in nonnullis integumento corneo integro, spatium pone coxas anticas etiam omnino claudente (Pinophilini, Stenini, Piestini, Proteinini).

Scutellum plerumque distinctum, triangulare aut rotundatum, rarius inconspicuum (Stenini, Oxytelini genuini plerique).

Elytra abbreviata, plerumque pectoris longitudine vel nonnunquam eodem paulo breviora (Aleochara), rarius pectore longiora, (Hypocyptus, Tachyporus, Conurus, Habrocerus, Tachinus, Omalini, Proteinini) et in his saepius abdominis summam basin, rarius maiorem eius partem (Lathrimaeum, Anthobium, Proteinus) obtegentia, apice truncata, saepius coniuncta emarginata, rarius angulo interiore producto (Anthobia quaedam), angulo exteriori iam rotundato (Omalini), iam saepius recto, iam in multis profundius incisio (Oxygota, Tanygnathus), lateribus in omnibus inflexis, saepe marginatis,

sutura plerumque ut in reliquis plurimis coleopteris integerrima, in Xantholinis genuinis imbricata.

Alae in plerisque Staphylinis obviae, exsertae abdominis apicem attingentes, longitudinaliter complicandae, plicaturis transversis duabus, altera infra, altera ultra medium, pone plicaturam priorem stigmate oblongo coriaceo instructae. Basi nervi longitudinales tres oriuntur, anterior in stigma, reliqui ad marginem posteriorem, intermedium ad plicaturam secundam, posterior ad plicaturam primam excurrunt. Nervi praeterea duo longitudinales pone plicaturam primam, anterior ex stigmate, posterior ante nervum e alae basi nascentem intermedium orientes longitudinaliter ad alae apicem vergunt, et anterior eorum ante, posterior pone alae apicem sensim tenuiores excurrunt.

Pectus mesothoracis breve, metathoracis amplius, mesosterno minuto, triangulari, plerumque simplice, interdum carina longitudinali subtili, (Conurus) rarius fortiore et magis prominente (Tachinus\*) instructum, metasterno sat amplo, pectoris maximam partem occupante, simplex, episternis et epimeris mesothoracis rhomboideis, methathoracis ad metasterni latera sitis, maxima parte ab elytrorum margine inflexo obtectis, angustis, coniunctis parallelipedis, sutura diagonali divisis, episternis posterioribus, epimeris antrorsum angustatis, his nonnunquam metasternum excedentibus, posterius prominulis (Aleocharini).

Abdomen quam plurimum totum aut maxima e parte nudum, plerumque lineare, saepius apicem versus angustatum, in nonnullis Tachyusis etiam basin versus attenuatum, in Omalinis latiusculum, saepius marginatum, rarius immarginatum (Conurus, Palaminus, Oedichirus, Procirrus, Steni permulti, Euaesthetus quidam, Osorius, Holotrochus, Leptochirus, Lissinus, Glyptoma), marginatum plerumque dorso saltem depressiusculum, immarginatum plerumque cylindricum, vivis mobile, praecipue elevandum et deflectendum, nonnullis apice deorsum curvatum (Boletobius, Mycetoporus), aliis sursum revolutum (Aleocharini permulti, praecipue Myrmedonia, Autalia, Falagria, Lomechusa), pluribus etiam, quando cursum accelerant, elevatum; segmentis omnibus liberis, anterioribus plerumque subaequalibus, rarius inaequalibus, (Tachysae quaedam, Hypocyptus), quinto nonnunquam reliquis paulo longiore. Segmenta sex semper valde distincta, dorso inter metathoracem et segmentum abdominis primum segmentum peculiare observandum, licet brevissimum, stigmate vero instructum, ita ut revera primum sit, a reliquis autem interstitio membraneo distinctum, metathoraci arctius adhaerens, inferne autem vestigio nullo detegendum, in descriptionibus mihi neglectum, ne in enumerandis segmentis segmenta dorsalia et ventralia numero incongrua lectorem irriterent.

Abdominis segmentum septimum organa sexualia exteriora continens et includens, quadrivalve, valvula dorsali integra, ventrali in maribus integra, in feminis longitudinaliter oblique fissa, ceterum valvulae dorsali subaequali, nonnunquam utrinque in processum styloformem elongata (Gymnusa), valvulis lateralibus saepe cylindricis, fortius pilosis setosisque, a reliquis magis remotis, stylos anales referentibus (Tachinus, Tachyporus, Staphylinini plerique et permulti Paederini Pinophilique etc.), nonnunquam basi compressis et valvuliformibus, apice styloformibus, vel uncinatis acuminatisque (Achenium), aut valvulis intermediis longioribus (Lathrobium quadratum), aut iisdem aequalibus (Staphylinus), aut iisdem brevioribus (Gymnusa) —



in reliquis totis valvuliformibus, valvulis intermediis arcte applicatis. In multis segmentum septimum totum reconditum (Aleocharini multi, Piestini, Omalini, Proteinii), ut nisi nonnunquam in maribus, ubi segmentum sextum excisum sive emarginatum est, conspici nequeat, in plerisque autem exserendum, ut quanto magis abdomen potensum sit, tanto magis promineat, in aliis etiam totum exsertum et haud retractile, licet segmentis reliquis minus (Phytosus, Lomechusa, et Oxytelini omnes).

Staphylinini plurimi, prehensi aut alio modo excitati prope anum utrinque papillam cutaneam exserunt, quibus forte odorem emittunt, quarum indoles autem nondum satis constat.

Pedes plerumque breviores, inter se longitudine subaequales, anteriores saepius paulo breviores et validiores, posteriores plerumque paulo longiores, raro reliquis debiliores (Xantholinus), antici et postici semper insertione approximati, intermedii saepe distantes, interstitio plerumque angusto, nonnunquam etiam latissimo, ut pedes intermedii pectoris lateribus insertae videantur (Oxyporus).

Coxae anticae maiusculae, conicae, exsertae, retrorsum vergentes, thoracis spatium integumento corneo destitutum obtegentes in plerisque, minutae, parum prominulae in Steninis, parvae, globosae, haud exsertae in Piestinis, subcylindricae, oblique positae, ex acetabulis vix prominentes in Proteinis. Coxae intermediae oblique positae, conicae aut subcylindricae, ex acetabulis haud prominentes. Coxae posticae subconicae, minores, quodammodo mobiles in Staphylininis, Paederinis, Pinophilinis, Steninis, transversae, subcylindricae, pariter paululum secundum axem transversalem mobiles in Oxytelinis, Piestinis, Phloeocharinis, Omalinis et Proteinis, transversim parallelopipedae, fixae in Aleocharinis et Tachyporinis, in his nonnunquam amplioribus, scutiformibus (Boletobius, Mycetoporus).

Trochanteres anteriores simplices, postici aut simplices (Staphylinini, Paederini, Stenini, Oxytelini, Piestini), aut fulcrantes (Aleocharini, Tachyporini, Phloeocharini, Omalini, Proteinini).

Femora plerumque simplicia et mutica, nonnunquam antica incrassata (Lathrobium), aut antica et postica inferne spinularum serie duplici instructa.

Tibiae nonnunquam spinulosae, et apice spinis apicalibus validioribus instructae, hae autem saepe minutissimae, anticae in aliis minus (Staphilinus, Philonthus etc.) in aliis magis spinulosae (Bledius), nonnunquam externe incrassatae et spinulis frequentibus muricatae (Xantholinus), interdum compressae et spinularum serie externe armatae (Oxytelus), aut denticulatae (Leptochirus, Piestus), plerumque integrae, nonnunquam ante apicem excisae (Oxytelini nonnulli); posteriores nonnunquam apice dilatatae, aut oblique truncatae (Lathrobium), aut emarginatae vel excisae (Pinophilini).

Tarsi articularum numero inconstante supra iam accuratus illustrati, antici aliis simplices, aliis articulis 4 primis magis minusve dilatatis, subtus plerumque tomento denso vestitis, nonnunquam lobis membranceis instructis (Pinophilus); posteriores simplices articulis primis aut aequalibus, brevibus, aut inaequalibus, primo magis minusve elongato, sequentibus sensim brevioribus, penultimo nonnunquam bilobo (Paederus, Steni permulti), aut minuto, subtus lobo membraneo aucto (Sunius, Pinophilini), ultimo s. unguiculari plerumque magis elongato, leviter clavato, unguiculis binis, aequalibus, simplicibus.



Stigmata 10 in utroque corporis latere, primum (prothoracicum) pone coxas anticas in inferiore corporis pagina situm, magnum, in Aleocharinis, Tachyporinis et Staphylininis peritremate corneo instructo, secundo (mesothoracico) pone elytrorum insertionem inter mesothoracis dorsum et episternum situm, occultum, tertium (metathoracicum) in dorsi margine laterali sub elytris inter metathoracis scutum dorsale et eius epimera situm, permagnum, reliquis abdominalibus, primo in segmento dorsali basali, reliquis in segmentis abdominis sex sequentibus dorsalibus prope marginem lateralem sitis, parvis, omnibus oblongis, fimbria laciniata subtiliter ciliata clausis.

### De corporis structura interiore.

Tracheae simplices, stipitibus maioribus in thorace, ubi stigmata sunt grandiora, fortioribus.

Systema nervorum cerebro bilobo, ganglio capitis, (cerebello) sub cerebro sito, ganglio prothoracico in anteriore thoracis parte sito, mesothoracico et metathoracico coalitis, ganglium singulum, reliquis multo maius, globosum, in mesothorace ante pedum intermediorum insertionem situm, formantibus; gangliis abdominalibus 7, tribus primis in medio metathorace sitis, approximatis, primo maiore, tertio in summa abdominis (segmenti primi) basi, quinto in segmento secundo, reliquis duobus in segmento tertio sitis, approximatis, septimo (ganglio sexuali) oblongo, in mare inter testiculos, in femina inter ovaria sito. Nervorum intestinalium systematis imparis ganglion frontale sub anteriore frontis margine situm, systematis paris ganglia bina pone et fere infra cerebrum oesophagi lateribus apposita, approximata.

Tubus intestinalis corpore vix sesqui longior. Oesophagus tenuis, sensim in ventriculum transiens. Ventriculus in mesothorace situs, parum amplus, subovalis, interne costis corneis longitudinalibus dense villosis, per paria alternatim fortioribus et fortius villosis, paribus earum fortioribus et subtilioribus quaternis. Duodenum amplissimum, a metathoracis basi ad abdominis apicem usque extensum, rectum, undique papillis intestinuliformibus dense quasi villosum. Intestinum ieium flexuosum, minus elongatum, colon simplex, absque coeco, rectum haud distinctum. Vasa biliaria bina, ansata, orificiis quatuor intestini duodeni apici inserta, libera (int. coloni haud affixa).

Organa sexualia mascula: Testiculi duo, integra, oblongi, vase deferente tenuissimo breviori ductui eiaculatorio inserti. Vasorum secermentum paria duo, in ductus eiaculatorii apice sita, interiora saepius longiora, cylindrica, intestinuliformia, exteriora sacciformia. Ductus eiaculatorius sat longus, flexuosus, apicem versus spinam longam tenuem corneam continens. Testiculorum structuram inquirens Leon Dufour vasis spermatici nullum vestigium, sed pulpam aequalem in iis invenit. Ipse in Staphylinino caesareo testiculum vasis cuiusdam spermatici ramosi finibus plurimis, intestinuliformibus, apicibus incrassatis glandulae conglobatae fere speciem praebentibus, accuratius autem oculo fortiter armato examinatis, convolutis, apice summo puncto obscuro, forte acino instructo. Tracheae numerosae in testiculo pertextae sunt. Penis corneus, subcylindricus, seta cornea laterali fultus.

**Organa sexualia feminea:** Ovaria fusiformia, funiculis ova continentibus paucis (tres in *Ocypod.* olente observavit Leon Dufour, ipse in *Staph. maxilloso* quatuor). Calices elongati. Oviductus communis bipartitus. Portio anterior calices excipiens dilatata, subglobosa vel fere transversa, uteriformis, portio posterior longior, cylindrica; bursa copulatrix eiusdem apici inserta, pyriformis, apice vasculo filiformi instructa. Appendices aliae nullae.

**Structura corporis interna** Staphylini minus *Carabis*, qui vasis analibus, testiculo e vase singulo convoluto formato et vasis biliaris ternis discedunt, quam *Silphis* et *Histeris* propinqui sunt, cum illis testiculorum structura, cum his oviductu basi simplice dilatato conveniunt. Intestino (duodeno) papilloso maximo et amplo praecipue excellunt. Conf. de tubi intestinalis structura: Leon Dufour *Annal. des Sciences naturelles* III. p. 219. t. 10. f. 6. 7. (*Staph. erythropterus*) f. 8. (*Staph. punctatissimus* (?)) f. 9. (*Paederus riparius*). — Ramdohr *Verdauungswerkzeuge der Insecten* p. 87. t. 3. f. 4. 5. 6. (*Staph. politus*) — de organis genitalibus masculis: Leon Dufour l. c. VI. p. 161. t. 5. f. 5 — 8. (*Staph. olens*, *maxillosus*, *erythropterus*, *punctatissimus*, *Paederus riparius*), de organis genitalibus femineis: Leon Dufour *ibid.* VI. p. 435. t. 17. f. 7. (*Staph. olens*). Ipse Staphylino maxilloso et caesareo dissectis observationes anatomicas supplavi.

### De metamorphosi.

Ova Staphylinorum oblonga, sat magna, at pauca.

Larvae adultae corpus elongatum, apicem versus attenuatum.

Caput magnum, orbiculatum, basi constrictum, collo quasi seu isthmo brevissimo thoraci insertum, subtus planum, supra leviter convexum, fronte apice denticulata, clypeo haud distincto.

Ocelli 4, quadratim positi, approximati, in utroque capitis latere pone mandibularum basin siti.

Os clausum.

Labrum brevissimum, sub frontis margine occultum, aegre observandum.

Mandibulae magnae, porrectae, falcatae, acutae, simplices, interne ante apicem apertura instructae.

Maxillae infra mandibulas insertae, a labio distantes, corneae, stipite elongato cylindrico, cardine brevi, mala singula, simplice.

Palpi maxillares stipiti crassiusculo, brevi, articulum basalem simulanti insertae, 4-articulati, articulis primis duobus longioribus, subaequalibus, tertibus, tertio minore, tenuiore, apicem versus sensim attenuato, quarto minimo, subulato.

Mentum membraneum, transversum.

Labium subquadratum, basi corneum, apice membranaceum.

Ligula minuta, angusta, porrecta, cornea.

Palpi labiales triarticulati, articulo primo longiore, tertio minimo, subulato.

Antennae 5-articulae, articulo primo brevi, secundo elongato, tertio quartoque decrescentibus, quinto minimo, subulato.

Segmenta thoracica supra scutis corneis instructa, infra membra-

nea, prothoracico maiore, infra antice sterno triangulari scutulisque duobus adiacentibus instructo, meso- et metathoracico fortiter transversis.

Pedes breves, ad latera segmentorum thoracicorum inserti; coxis cylindricis, oblique introrsum directis, trochanteribus brevioribus, femoribus teretibus, inferne spinulosis, tibiis teretibus, undique spinulosis, tarsis minutis, articulo simplice et ungue leviter falcato, acuminato, singulo compositis.

Segmenta abdominalia 9, latitudine decrescentia, penultimis paulo longioribus, supra et infra areis binis coriaceis instructa, segmenta prima supra scuto corneo oblecto, ultimo toto coriaceo, cylindrico, apice caudis duabus articulatis instructo.

Caudae anales biarticulati, articulo primo longissimo, secundo tenuiore, apice seta longa rigida terminato.

Anus fortiter prominens, tubulosus, descendens.

Stigmata 18, vel 9 paria, pari primo prothoracico, reliquis pari singulo in abdominis segmentis 1 — 8 singulis siti.

Stigmata prothoracis maiora, infra in membrana interstitiali inter pro- et mesothoracem posita. Mesothorax et metathorax stigmatibus carent.

Stigmata abdominalia in segmenti cuiusque facie superiore, in areae coriaceae dorsalis margine laterali, paulo ante medium posita, minuta, orificio elevato.

Segmenta thoracica ante quam penultima cuticula deponitur, membranea, abdominis segmentis aequalia, testibus Frisch (Ins. Germ. V. p. 49.) et Heer (Obs. ent. p. 20).

Staphylinorum larvae inter se satis convenire et nonnisi notis levioribus discedere videntur; ab aliorum Coleopterorum larvis facile distinguendae sunt. Caraborum larvae discedunt ocellis senis, ore aperto, mandibulis dentatis, maxillis prope labium insertis, mala biarticulata, palpiformi, tarsis bi-ungiculatis, abdominis segmento ultimo utrinque corniculato, caudis analibus nullis, stigmatibus abdominalibus lateralibus, inter segmenta dorsalia et ventralia sitis. Dytiscorum larvae corporis habitu, ore clauso, mandibulis simplicibus perforatis, maxillis distantibus, mala simplici instructis, palpisque gracilioribus eximie cum Staphylinorum larvis congruunt, differunt autem ocellis senis, maxillarum mala unciformi, ligula lata, transversa, tarsis bi-ungiculatis, ano haud exserto, caudis analibus elongatis, exarticulatis, tubulos respiratorios referentibus, stigmatibus, licet conspicuis, clausis.

Magis propincae sunt Silpharum et Histerorum larvae. Illae (Silphae cuiusdam et Catopis fuscii larvis comparatis) differunt capite minore, deflexo, ore aperto, labro exserto, mandibulis haud prominulis, maxillis mala interiore sat forti, interne denticulata aut spinulis armata, exteriori nonnunquam distinguenda, licet parva et obsoleta, labio lato, emarginato, ligula nulla, palpis minus elongatis, maxillaribus 3-, labialibus 2-articulatis, ocellis utrinque sex, 4 approximatis lateribus supra antennarum basin ut in Staphylinis, duobus ab his remotis in genis infra antenas versus mandibularum basin sitis, stigmatibus in inferiore corporis facie obviis — haec (secundum Hist. merdarii larvam, cuius specimen a cel. Müllero sibi missum Dom. Schüppel mecum communicavit, a Paykullo Mon. Hist. t. I. f. I. perperam delineatam) multo magis affines, praecipue capite porrecto, labro occulto, palpis elongatis, gracilibus, mandibulis autem dentatis, maxillis mala



destitutis, ligula nulla, ocellis omnino nullis, corporis segmento primo (prothoracico) solo corneo, pedibus omnibus brevissimis, ano tubuloso haud exserto.

Nympha membrana tenui obducta, uti reliquae coleopterorum nymphae, margine prothoracis anteriore setis erectis instructo insignis.

Primus Staphylinorum metamorphosin observavit noster Frisch, qui (Ins. Germ. V. p. 49. t. XXV.) speciei cuiusdam mediae magnitudinis, quam hucusque pro Staph. erythroptero caesareove habuerunt auctores, quae potius autem ad Tachinum quendam referenda videtur, historiam accuratius descripsit. Alterius cuiusdam speciei minutae, forte Oxyteli cuiusdam, (ibid. p. 50. t. XXVI.) cuius larvam in fimo siccior auctor invenit, historia non indubia videtur, cum larva pedibus ventralibus instructa exhibeatur, quam ob rem commutatione quadam Lepidopteri cuiusdam pusillam erucam pro Staphilini larva h. l. suppositam esse suspicor. Cel. Gravenhorst deinde (Micr. p. LXI.), similitudine quadam exteriore deceptus, Carabi genuini larvam pro Staphylini cuiusdam maioris larvam perspicue descripsit. Recentioribus temporibus plures et magis exacte Staphylinorum larvae observatae sunt. Dom. Bouché Philonthi aenei (Hist. nat. I. 179. 1. t. VII. f. 29 — 35.), Quedii fulgidi (variabilis) (ibid. 180. 2. t. VIII. f. 1 — 8.), Xantholini punctulati (ibid. 181. 3. t. VIII. f. 9 — 13.) et Platystheti morsitantis (ibid. 182. 4. t. VIII. f. 14 — 21.) metamorphoses illustravit, Dom. Heer Ocypodis olentis historiam et larvae structuram accuratius exhibuit (Obs. Ent. 16. 5. t. 3. A.), eiusdem speciei larvam et nympham Dom. Blanchard (Guerin Mag. Zool. VI. t. 165.) mance, Dom. Ratzeburg (Forstinsect. I. p. 30. t. 1. f. 14.) eleganter delineaverunt, Dom. Waterhouse denique (Transact. Soc. Ent. Lond. I. p. 32. t. 3. f. 2.) Quedii tristis larvam proposuit. Descriptiones auctorum modo laudatorum secundum singulum larvae cuiusdam (nescio num Staphylini an Ocypodis, anne potius Philonthi cuiusdam maioris, sine dubio autem Staphylinini cuiusdam genuini) specimen, plures per annos mihi conservatum, supplevi, et observationes quasdam maioris momenti, quae auctores reliquos fugerunt, de eius structura addidi. Plurimorum Staphylinorum metamorphoses illustrandas posteris relinquimus.

### De victu.

Victu Staphilini plerique non solum putredine delectari, sed carnavori esse videntur, ut plurimi in fimo, in vegabilibus putridis et in boletis obvii, aliis etiam insectis, praecipue Dipterorum larvis, quibus materies illae putrescentes abundant, ibidem insidiari videantur. Multi etiam animalium cadavera copiosi aggrediuntur et cum Silphis carne putrescente vescuntur. Stercus praecipue habitant Tachini, Staphylini nonnulli, Philonthi, Oxyteli, vegetabilia putrida Aleocharini plerique, Conuri, Tachypori, Quedii, Paederini plurimi, boletos Aleocharini permulti (Autalia, Bolitochara, Gyrophæna etc.), inter Tachyporinos Boletobii et Mycetopori, inter Staphylininos imprimis Oxypori. Caraborum more Staphylini maiores (Staphylini genuini, Ocypodes), interdum in terra, sub musco, foliis deciduis et lapidibus latitant, noctu circumvagantes insecta venantur, alii interdum in arena riparia humida summa agilitate cursitant (Steni, Tachysae). Permulti etiam sub arborum cortice latent: inter Aleocharinos Phloeopora nonnullae, Ho-

malotae et secunda Oxypodarum sectio, inter Staphylininos Quedii nonnulli (*Q. laevigatus*, *scitus*), dein Piestini omnes et Phloeocharini, Glyptoma. Osorii sub cortice putrescente cuniculos sibi parant. Bledii quoque, alii in arena riparia vel littorali, alii in terra pingua cuniculos, nonnunquam sat profundos effodiunt. Simili ac nonnulli littorales Bledii vitae ratione Aleocharinorum species, *Diglossa mersa*, insignis, quae in arena littorali aestu inundata degens, alternis vicibus fluctu obtegitur. Floribus delectantur Nitidularum more Omalinorum plurimi et Proteinus.

In formicetis etiam quidam Staphilini quaerendi, praecipue Aleocharini: *Lomechusa*, *Dinarda*, *Euryusa*, *Myrmedoniae*, e reliquis etiam generibus species singulae, e. g. *Falagria thoracica*, *Homalota anceps*, *Aleochara angulata*, e reliquis Staphylinis pariter species singulae, e. c. *Stenus aterrimus*.

Permuli Staphylini in ripis occurrunt, praecipue Paederi et Steni, multi Aleocharini et complures Oxytelini. Nonnulli etiam nonnisi in maris littoribus obvii sunt, aut sub fucis putrescentibus (*Philonthus xantholoma*, *fucicola*), aut in humo fluctu salso humectata (*Bledius fureatus*, *unicornis*); hi pariter in lacuum salsorum ripis obveniunt, qui similem sale praegnantem glebam iis praebent.

Larvis eadem vitae ratio ac imaginibus est. Plurimae pariter vagae sunt. Ocypodis olentis larva, larvarum Cicindelarum more, in cuniculo latet, quem, mandibulis terram arripiens et pedibus anterioribus proiciens, dein ex profundo mandibulis terram tollendam auferens, dimidium vel totum pedem profundum, oblique descendantem parat. Plerumque larva in cuniculi orificio cubans, mandibulis apertis insectis praetereuntibus insidiatur, quae, simulatque eaprehendere valet, minora in cuniculi fundum secum detrahit, maiora, pedibus anterioribus praedam arripiens, posticis adnixa, in cuniculi orificio depascit.

O. olentis larvam erucam, quam ei praebuerat, devoravisse, alterius erucae, brevi tempore post ei oblatae tantummodo sanguinem bibisse tradit Dom. Heer, at patet ex oris omnino clausi, mandibulis solis perviis praediti structura, eam nonnisi sugendo ex praeda sua nutrimentum capere. Haec res accuratissimum Frischium non omnino fugit. Helices etiam larvam illam avide arripuisse, at ore muco illarum obducto relinquisse, helicem dissectam autem paulatim devoravisse Dom. Heer observavit. Nocte semel larvam extra cuniculum circumvagantem vidit. Ad cuticulam deponendam in cuniculi inam recedit partem, orificium terra clausum reliquens. In nymphae statum transiens speluncam ovatam, intus laevigatam et politam sibi parat. Quinque prope menses hoc statu peragit.

Larvae praecipue versus autumnum, per hiemem et primo vere obviae sunt. Imagines etiam plurimi in terra, in fimo, sub foliis putridis, musco, lapidibus et sub arborum cortice hibernant. Vere et autumnum tempore meridiano, aestate vesperi coelo sereno plurimi volitant.

### De distributione geographica.

Cum omnino insectorum, quae non solum e remotis orbis nostri terrarum regionibus adportantur, sed in nostris etiam terris quotidie nova reperiuntur, numero in immensum quantum continuo gliscente, tot tantisque terrarum plagis ne a fugitivo quidem naturae strutatoris calce tactis, plerisque

regionibus magis ex eo, quod forte in peregrinatorum manus incidit, cognitis, quam rite perlustratis, insectis ipsis denique, imprimis minutis, quae undique obveniunt, nondum satis examinatis et illustratis, ut, quae e diversis regionibus allata specie divergant convenientve satis sit recensitum, iam omnino in eo esse nobis videri non possumus, ut nisi quod periculi speciem prae se ferat aliquid de distributione insectorum proferre valeamus, tum praecipue de Staphylinorum, qui ob studiū diffultatem et corporis habitum ingratum et quam plurimum exiguum ab observationibus et collectoribus insectorum plerisque maxime neglecti sunt, rationibus geographicis minus nondum instructos nos esse elucet. Etiam si observationes meae hac de re paucae et imperfectae sunt, tamen eo minus eas promere dubito, quanto magis studio huius familiae speciali intersunt.

Numerus Staphylinorum relativus certe non adeo exiguus est, quam e faunis plerisque et e catalogis collectionum patere videtur. In collectione Dejeaniana, quam in omnibus coleopterorum familiis aequaliter fere locupletem esse constat, Staphylini vix ultra tricesimam partem occupant, et Carabis numero plus triplo inferiora sunt. In Stephensii contra enumeratione Insectorum Angliae Staphylini quartam paene Coleopterorum partem amplectuntur, et Carabis duplo prope sunt numerosiora, et idem fere in Fauna mea Marchica reperio, ubi Staphylini Carabos duplo numero superant. Hanc eandem proportionem etiam omnino valere mihi persuasum, nam specierum exoticarum copia eam ob causam exigua in collectionibus reperitur, quod peregrinatores Staphylinis plerisque parvulis et victu absconditis minus quam aliis Coleopteris colligendis operam navaverunt, et quamquam auctores quidem (Perty, Lacordaire) Staphylinos in tropicis terris multo minus copiosos esse quam apud nos affirmarunt, tamen hanc rem non solum ob causam modo laudatam revera sic sese habere acclamare nequeo, sed eo etiam de facto, quod insectorum quidem collector sedulus gnarusque, Dom. Moritz, ex America aequinoctiali tantum ex hac familia specierum numerum apportavit, ut non modo ne minutarum quidem specierum copia tropicis regionibus terrae nostrae praestare videantur, sed ex eadem etiam proportionem, quam apud nos valere memoravi, duplum fere eorum quam Caraborum numerum collegerit. Cave autem, cum in quavis regione Staphylinorum copiam maiorem esse Caraborum numero constare videatur, ne ex eadem ratione omnino Staphylinos numero duplo superare Carabos concludas, cum Staphylinorum species singulae multo magis dissipatae sint, ut in regionibus aliquantum distantibus Caraborum fauna maxima ex parte alia, Staphylinorum eadem sit. Specierum Europaearum maiorum pars per totam hanc orbis terrarum partem obvia est, ut nostrarum plurimae pariter Lapponiam atque Siciliam habitent, et plurimae quae in tota Asia septentrionali (Sibiria) reperiantur species cum Europaeis congruant, multae etiam per totam Orientem (Asiam minorem, Syrian, Aegyptum) inveniantur, aliae pariter in America septentrionali indigenae sint, nonnullae remotioribus etiam locis reperiantur. Sequentia sunt quae hac de re hucusque observavi facta:

Totam fere Europam et Orientem habitant:

*Homalota gregaria.*

*Aleochara tristis.*

— *hisignata.*

— *nitida.*



*Leptacinus batychnous.*

*Philonthus intermedius.*

— *cephalotes.*

— *ebenus.*

— *bipustulatus.*

— *quisquiliarius.*

*Stenus bipunctatus.*

*Platysthetus cornutus.*

*Oxytelus piceus.*

Per omnem Europam et in China meridionali occurrit:

*Conurus pubescens.*

Per omnem praeter maxime borealem Europam, in Mauritania et in Tene-  
riffa invenitur:

*Ocypus olens.*

In omni Europa et in Africae fine meridionali obvenit:

*Oxytelus sculpturatus.*

Plures sunt species Americae septentrionali cum Europa communes, partim  
ab auctoribus secundum patriam alteram altero nomine impositae:

*Homalota analis.*

— *lividipennis* (*Oxypoda livida* Dej.)

*Aleochara fuscipes* (*Aleoch. lata* Grav.)

*Tachinus silphoides* (*Tachyporus marginalis* Grav.)

*Staph. fossor* (*Staph. fodiens* Grav. forte):

*Ocypus ater* (*Staph. nigripes* Dej.)

*Philonthus cyanipennis* (*Staph. coeruleipennis* Mannerh.)

— *aeneus.*

— *umbratilis* (*Staph. subfuscus* Gyll.)

— *ventralis* (*Staph. anthrax* Grav.)

— *fulvipes.*

— *aterrimus.*

*Quedius fulgidus* (*Staph. Groenlandicus* Zett.)

— *molochinus* (*Staph. laticollis* Grav.)

*Stenus Juno.*

e quibus *Philonthus ventralis* etiam in Americae meridionalis (Indiae  
occidentalis) insulis occurrit.

Nonnullae etiam Europaeae species pariter in Orientem atque in Americam  
septentrionalem patriam extendunt:

*Philonthus debilis.*

*Oxytelus sculptus.*

— *nitidulus.*

In locis magis distantibus et remotis observantur:

*Philonthus varians* in Europa, in Oriente, ad Promontorium Bo-  
nae Spei atque in Americae meridionalis insulis.

*Aleochara morion.*

*Philonthus discoideus.*

*Lithocharis ochracea* in Europa et in variis Americae meridio-  
nalis locis.

Species Americanae magis quam Europaeae certis limitibus circumscri-  
ptae videntur, ita ut Staphylini Columbiae plerique ab iis Brasiliae, Ame-

ricae meridionalis ab iis Americae septentrionalis distinctae sint. Speciem nonnisi singulam novi, quae totam Americam habitare videtur: *Oxytelus insignitus* (O. Americanus Dej.), et paucae sunt ex iis, quae Americam septentrionalem incolunt, quae ultra eius fines, sicut *Staph. villosus* in Mexico quoque et in Cuba obvius est, vel ultra mare Antillarum, sicut *Belonuchus formosus* (*Staph. rufipennis* Fab.) Cayennam et Columbiam habitat, indigenae observantur.

Ut e reliquis insectis formae maiores secundum geographicas rationes distributae, minores et praecipue minimae per omnes plagas dissipatae esse videntur, sic apud Staphylinos etiam genera pleraque e speciebus parvulis constituta per totum terrarum orbem occurrere videntur, speciebus tropicis et septentrionalibus simillimis. Steni in Europa numerosi, multi in America vel septentrionali vel meridionali, nonnulli in Madagascar occurrunt et in India orientali etiam posteriore eosdem sibi observatos esse literis certiores nos fecit Dr. Helfer. Falagriae, Homalotae, Aleocharae, Conuri, Bledii, *Oxyteli* eodem modo ubique obvenire videntur. Nonnullae etiam genera, paucas tantum species continentia, pariter apud nos et in America tropica reperiuntur, ut *Tanygnathus*, *Heterothops*. Alia autem sunt genera numerosiora nonnulla, quae regiones temperatas non excedere videntur, ut *Tachypori* ultra Europam et Americam septentrionalem haud observati sunt, et *Tachinorum* sectio altera praecipue tropicis plagis, altera Europae et Americae septentrionali minus calidae peculiaris videtur. Sunt etiam quae nonnisi tropos habitant, ut *Piesti*, *Leptochiri*. Et quamvis pleraque *Staphylinorum* genera, licet iis, quae singulam vel paucas species continent, exceptis, Europae cum remotioribus etiam regionibus communia esse solent, tamen non omnino desunt, quae in variis Asiae, Africae, et Americae utriusque regionibus obveniunt, quibus Europa omnino caret, ut *Pinophilus*, *Osorius*.

Fauna Europaea *Omalinorum* copia insignis videtur, et in ipsa Europa haec tribus in regionibus septentrionalibus et alpestribus frequentius occurrere videtur. Species boreales omnino facile in regionum temperatarum montibus reperiuntur, in quibus sat elevatis boreali climate foventur. Sic *Tachinus* elongatus non modo in Suecia et in Unalaschka, sed in Helvetiae etiam alibus obviis. Alpibus aut montibus elevationibus praeterea complures propriae videntur species, praecipue e genere *Ocypode* (O. macrocephalus, *Italicus*, alpestris, megacephalus, alpicola). Species alpinae saepenumero apterae: sic *Paederus littoralis* etiam, qui regiones elevationes habitat, alis destitutus est. Species salinae minime arctis limitibus terminantur: *Philonthus xantholoma* in omni Europae littore reperitur. *Bledius furcatus* et *unicornis* Europae meridionalis et occidentalis littora nec non Germaniae interioris lacus salsos habitant. *Bledius bicornis* pariter ad eosdem Germaniae atque Iberiae lacus salsos observatur.

Asiae totae septentrionalis fauna in hac familia ab Europaea parum distincta videtur, ut et tota Asia occidentalis, praecipue Asia minor et Syria nec non Mesopotamia easdem plerasque species atque Europa proferunt. Indiae *Staphylini* parum innotuerunt, nec peculiaris quidquam habere videntur. Genera *Myrmedonia*, *Tachinus*\*, *Xantholinus*, *Staphylinus*, *Philonthus*, *Quedius*, *Paederus*, *Pinophilus*, *Osorius*, *Bledius*, *Leptochirus* hucusque ex India nobis allata sunt, et formas Europaeas plerasque inveniri, licet speciminibus raris, neque copiosis ut apud nos, a

Helfero literis nobis relatum. Palaestrini forte genus Indiae orientali proprium.

Africani Staphylini non magis quam Indiae orientalis species innotuerunt. Madagascarienses plures legit Goudot, e quibus alii Europaeis simillimi et proximi (*Tachinus pictus*, *Paederus Madagascariensis*), alii magis Staphylinorum Americae meridionalis habitum prae se ferunt (*Steni*, *Holotrochus*, *Lispinus*, *Isomalus*). Staphylini Capenses pauci adhuc lecti, nonnulli ab Europaeis haud distincti (v. s.), alii speciebus Aegyptiis et Indicis approximati (*Myrmedonia*, *Pinophilus*). Africa aequinoctialis in Staphylinorum familia eximiam suam faunam nondam aperuit; quas cognovimus species paucae ad Aegyptum usque fines suos extendunt. Aegyptia fauna, ut in reliquis insectis, valde cum Europaea convenit, speciebus plurimis omnino iisdem, immixtis autem multis formis partim Africae aequinoctiali, partim Indiae orientali propriis.

Americae septentrionalis Staphylini Europaeis omnino simillimi sunt et saepe omnino iidem. Genera nonnulla his ambobus orbis terrarum partibus propria videntur, ut *Oxypori* in America septentrionali, *Anthophagi* in Europa frequentiores occurrunt. America meridionalis magis peculiarem characterem prae se fert, putredine autem in calidis eius regionibus tantopere accelerata, observantibus Lacordaire et Perty, individua ibidem rariora. Specierum autem numerus, tot specierum minutarum copia a Beskio in Brasilia, a Moritzio in Columbia et in pluribus insulis lecta, nequaquam imminutus videtur, quamvis hac contentione Lacordaire et Perty consentiunt. Tropicae Americae cum Madagascariensi et Indiae orientalis insulis communes sunt *Leptochiri* et *Osorii*, peculiare videntur *Piesti*. Americae omni propria videntur quaedam genera, quae et in America septentrionali et meridionali occurrunt, e. g. *Palaminus*, *Megalops*, quae autem succedentibus temporibus in Africa etiam et in India orientali forte reperiuntur. In Americae meridionalis (s. Indiae occidentalis) insulis et praecipue in Mexico Americae meridionalis et septentrionalis faunae conveniunt eo modo, ut plurimis speciebus peculiaribus aliae immixtae sint A. septentrionali (e. g. *Staphylinus villosus*), aliae Americae meridionali (e. g. *Staph. versicolor*) propriae.

Novae Hollandiae et reliquae Australiae Staphylinorum fauna adhuc parum cognita, ad quantum eam novimus, licet speciebus sat distinctis, formas ab indigenis differentes non profert.

#### De dispositione.

Totam Staphylinorum nostram familiam genus Linneanum *Staphylinus* amplectitur, in Fauna Suecica 21, in Systemate Naturae (Ed. X.) 26 species continens, e quibus *St. sanguineus* ad *Pselaphos*, *St. cantarellus* ad *Malthini* genus pertinentes excludendae sunt. Sectionibus Linnaeus specierum numerum tam exiguum nullis distinxit, sequentes autem nonnulli auctores aut capite thoraci aequali eodemve angustiore (*Paykul* Mon. Staph.), aut thoracis in aliis paralleli, in aliis apicem, in aliis basin versus angustati, in aliis cylindrici forma (*Müller*), divisiones instituerunt. Fabricius primus genera altera duo addidit, *Oxyporum* et *Paederum*, illum palpis posterioribus securiformibus, hunc palpis anterioribus clavatis a Staphylinis palpis omnibus filiformibus distinctum. Ceterum Staphy-



lino plurimas licet diversissimas species continenti, *St. erythroptero* pro typo proposito, labium membranaceum, sub palpis elongatum, trifidum, lamina intermedia latiore, apice rotundata, subemarginata, lateralibus paulo longioribus, distantibus, acutis, ligula paraglossisque computatis, *Oxyporo* labium „elongatum, membranaceum, cylindricum, emarginatum cum mucrone” et *Paedero* „porrectum, membranaceum, cylindricum, apice truncatum, obtusum, integrum,” in utroque ligula et paraglossis omnino neglectis, tribuit, *Oxyporo* plurimos quos cognovit *Tachyporinos*, quamvis distincte palpos eorum filiformes esse monuit acutissimusque quisque auctor, pertinaciter adiciens. In *Systemate Eleutheratorum* denique quartum genus, *Stenus*, a *Latreillio* conditum, a *Paykullo* receptum, palpis inaequalibus, anterioribus longissimis et antennis clava solida distinctum, addidit, generis autem illustrationem ex oris fabrica erroribus pluribus omnino perversam perperam a *Paykullo*, nisi *Paykull* ipse hanc misellam oris descriptionem a *Fabricio* acceperit, mutuavit, neque minus improspere *Nitidulas* quasdam elytris abbreviatis et *Omalium* quoddam generi huic quam maxime naturali adscripsit.

Maxime de hac familia meritis est cel. *Gravenhorst*, qui duabus *Monographiis* eandem (*Microptera* vocatam) illustravit et primus genera naturalia invenit et singulari habilitate distinxit. Notis imprimis habitibus genera 14 cel. auctor in *Micropt. Brunsv.* hoc modo disposuit:

Palpis anterioribus	3-articulatis. Antennarum articulo ultimo	{	ovato. Articulo ultimo palporum posteriorum cylindrico	{	acuminato . . . . . ovato . . . . .	3. <i>Paederus</i> . 12. <i>Stenus</i> ,
	4-articulatis. Articulis ultimis palporum	{	inaequalibus	{	rotundato . . . . . ovatis { Thorace { postice truncato. Antennis { extrorsum crassioribus. Thorace filiformibus. Thorace	4. <i>Callicerus</i> . 11. <i>Oxyporus</i> . 1. <i>Staphylinus</i> . 7. <i>Omalium</i> .
			aequalibus	{	marginato . . . . . immarginato . . . . . cordato . . . . . quadrato . . . . .	10. <i>Tachinus</i> . 8. <i>Anthophagus</i> . 14. <i>Pinophilus</i> . 13. <i>Astrapaeus</i> .
			aequalibus	{	securiformibus . . . . . gracilibus { elongato . . . . . subuliformibus { brevi Tibiis { spinulosis. Thorace	2. <i>Lathrobium</i> . 5. <i>Aleochara</i> . 9. <i>Tachyporus</i> . 6. <i>Oxytelus</i> .

In opere suo posteriore, *Monogr. Micropt. Dom.* *Gravenhorst*, *Callicerum Aleocharae*, *Astrapaeum Staphylini* adiecit, et minus feliciter atque potius contra naturam *Pinophilum Lathrobii* subiunxit, nova autem tria genera addidit: *Lomechusam* ob thoracis formam ab *Aleochara* distinctam, *Euaesthetum*, structura thoracis cum *Oxytelo*, palporum et antennarum cum *Steno* convenientem, *Piestum*, corpore depresso, et pedibus brevibus, validis, tibiis serrato-ciliatis ab *Anthophago* discedentem. — „*Lathrobiorum*, *Paederorum*, *Stenorum* palporum structuram haud accuratissime vidisse *Dom. Gravenhorst*, — his insectis palpos maxillares articulis quatuor, ultimo tamen saepius ob exilitatem ope etiam microscopii vix conspicuo, compositos, — genera *Staphylinorum* ab illo auctore proposita naturalia, optima, sed characteribus novis, ex instrumentis cibariis praesertim depromendis fulcienda esse,” praestantissime iam monuit *Latreille* (*Gen. Crust. et Ins.* I. p. 289). Palpos *Calliceri* et *Paederorum* quadriarticulatos sese vidisse *Dom. Gravenhorst* in *Mon. affert*

Latreille in opusculo suo primario „Précis des Caractères génériques des Insectes” receptis generibus Fabricianis primus condidit genera *Stenus*, *Proteinus* et *Lesteua*, haec ultima dua autem a reliquis Staphylinorum generibus remotis, *Proteinum* Nitidulis, *Lesteuam* *Carabis* adiiciens. Magis iam provectum est systema, quod idem ille gravissimus auctor quinque annis post proposuit in *Hist. nat. Crust. et Ins.* vol. III, ubi Staphylini familiam peculiarem (Staphyliniae), inter *Necrophagos* et *Nitidularias* positam constituunt, genera 6 continentem; *Staphylinum* (*Staphylinus*, *Oxytelus*, *Aleochara*, *Omalium*, *Tachinus* et *Tachyporus*), *Lesteuam* (*Anthrophagus* Grav.), *Paederum* (*Lathrobium* et *Paederus* Gr.), *Stenum*, *Oxyporum*, *Astrapaeum*, omnia imprimis secundum palporum figuram distincta. Deinde in eodem opere (Vol IX et X) in speciebus enumerandis genera *Gravenhorstiana* *Lathrobium*, *Aleochara*, *Oxytelus*, *Omalium*, *Tachyporus* et *Tachinus* recepta sunt. *Proteinus* adhuc inter *Nitidularias* enumeratus. Nonnullis annis praeterlapsis in opere suo classico et summo studio elaborato, *Genera Crustaceorum et Insectorum* Latreille hanc familiam (Staphylinii) inter *Palpatores* (*Scydmaenum*) et *Necrophagos* collocatam, hoc modo in sectiones et genera dissolvit:

I. Palpi labiales articulo tertio sive terminali maiore, subtrigono fereque securiformi aut lunato.

*Oxyporus*, *Astrapaeus*.

II. Palpi labiales articulo tertio sive terminali haud maiore, nec trigono, nec lunato, saepius subcylindrico aut subconico, in nonnullis obsoleto.

1. Mandibulae ad latus internum processu aut saltem dente uno valido instructae.

*Staphylinus*, *Lathrobium*, *Paederus*, *Stenus*, *Oxytelus*.

2. Mandibulae dentibus nullis internis aut unico tantum vixque obvio,

A. Antennae ante oculos insertae, a secundo baseos articulo ad angulum fere rectum haud divaricatae, articulis nullis aut perpauca (3 — 4) transversis, plerisque obconicis aut subglobosis, terminali subovato, ad basin rotundato, non truncato.

a. Mandibulae fissura interne parva aut subunidentatae.

*Lesteua*, *Omalium*, *Proteinus*.

b. Mandibulae latere interno penitus edentulo.

*Tachinus*, *Tachyporus*.

B. Antennae oculorum margini interno et antico insertae, a secundo baseos articulo ad angulum fere rectum divaricatae (subfractae et antrorsum arcuatim inflexae), articulis 5 — 10 transversis, perfoliatis, terminali ovato-conico, basi truncato.

*Aleochara*.

Genera sunt pleraque *Gravenhorstiana*, at naturali modo disposita et characteribus certis circumscripta.

Nova disponendi methodo in hac familia, quam *Brachypterorum* (*Brachélytres*) nomine impositam secundam inter *Adephaga* et *Serricornes* collocavit Latreille in recentiore opere: „Familles naturelles du règne animal (1825)” usus est, ubi sectiones quatuor distinxit:

1. Fissilabri, labro profunde bilobo: *Oxyporus*, *Astrapaeus*, *Staphylinus*, *Xantholinus*, antennis supra labrum, *Pinophilus*, *Lathrobium*, antennis externe supra mandibularum basin insertis.





Sectiones tantum quatuor in Staphylinorum familia Dom. Stephens constituit: Tachyporidas, Aleocharinos et Tachyporinos continentem, Staphylinidas, Staphylininos cum Lathrobiis, Stenidas, Paederinos reliquos, Steninos, Oxyporum et Oxytelinos, Omalidas, Euaesthetum, Syntomium, Coprophilum, Omalinos et Proteininos continentes. Genera permulta proposuit auctor, quae vero habitualibus solis plerumque notis distincta, characteribus systematicis autem destituta, potius in tyronum usum quam in scientiae incrementum provocata videntur, et singula loco suo mihi recensenda sunt.

Ipse nova dispositione, iam in Coleopterorum Marchicorum descriptione tentata, usus, secundum stigmata prothoracica conspicua occultave, antennarum insertionem et coxarum anticarum posticorumque et trochanterum posticorum formam Tribus undecim, quae naturales mihi videntur, hoc modo distinxi:

Stigmata protho- racica	conspi- cua. An- tennae	in facie ad oculorum marginem interiorem			
		insertae . . . . .			<i>Aleocharini.</i>
		sub frontis margine laterali insertae . . . . .			<i>Tachyporini.</i>
	occulta. Coxae posticae	in frontis margine anteriore insertae . . . . .			<i>Staphylinini.</i>
		conicae. Pro- thorax spa- tio pone co- xas anticas	membraneo . . . . .		<i>Paederini.</i>
			corneo. An- tennae	sub frontis margine laterali insertae . . . . . in fronte insertae . . . . .	<i>Pinophilini.</i> <i>Stenini.</i>
	transversae. Trochanter postici	simplices. Coxae an- ticae		conicae, promi- nentes . . . . .	<i>Oxytelini.</i>
				globosae, haud pro- minentes . . . . .	<i>Piestini.</i>
		fulcrantes. Coxae an- ticae		conicae, { nulli . . . . . exsertae. { duo . . . . .	<i>Phloeocharini.</i> <i>Omalini.</i>
				Ocelli { subcylindricae haud exsertae . . . . .	<i>Proteinini.</i>

### Trib. I. Aleocharini.

Palpi maxillares articulo primo brevi, quarto minuto, subulato. Labrum integrum, simplex. Ocelli nulli. Elytra pectoris longitudine. Coxae anticae conicae, exsertae, posticae transversae. Trochanteres postici fulcrantes. Abdomen marginatum, segmento septimo saepius occulto.

4-articulati .	mutica, undique pubescens	antici 4-, posteriores 5-articulati, postici articulis 4 primis omnes 5-articulati .	inaequalibus. Ligula ad palporum labialium articulis 4 primis apicem usque porrecta { secundi apicem usque porrecta aequalibus .	<i>Aleochara.</i> <i>Myrmedonia.</i> <i>Falagria.</i> <i>Bolitochara.</i> <i>Phytosus.</i> <i>Calodera.</i> <i>Tachysa.</i> <i>Homalota.</i>
3 - articulati. Maxillarum mala interior	interne spinulosis ciliata. Paraglossae .	obsoletae, Tarsi	antici 4-, posteriores 5-articulati, postici articulis 4 primis omnes 5-articulati, postici articulis 4 primis omnes 4-articulati. { 11-articulatae 10-articulatae .	<i>Ocalea.</i> <i>Orypoda.</i> <i>Phloeopora.</i> <i>Hygronoma.</i> <i>Oligota.</i> <i>Dinarda.</i> <i>Lonechusa.</i> <i>Autalia.</i> <i>Euryusa.</i> <i>Pelusia.</i> <i>Gyrophacna.</i> <i>Placusca.</i> <i>Diglossa.</i> <i>Silusa.</i>
2-articulati, articulo secundo	unco terminata	primo crassitie aequali	recto. Tarsi postici aequalibus articulis 4 primis lunato { distantibus approximati	<i>Pronomaea.</i> <i>Myllaena.</i> <i>Gymnusa.</i>
exarticulati. Tarsi		omnes 4-articulati	antici 4-, posteriores 5-articulati, postici articulis 4 primis omnes 5-articulati	

## Trib. II. Tachyporini.

Labrum integrum, simplex. Mandibulae muticae, lacinia membranacea interiore connata. Ocelli nulli. Coxae anticae conicae, exsertae. Trochanteres postici fulcrantes. Tibiae saepius spinulosae. Abdominis segmentum septimum vix conspicuum.

Elytra	pectore longiora. Tarsi	{	4-articulati	{	filiformes. Palpi	{	subulati.	{	immarginatum	<i>Hypocyptus.</i>
			5-articulati.				Abdomen		marginatum	<i>Conurus.</i>
		Antennae	{	max.	{	filiformes	{		<i>Tachyporus.</i>	
				capillares					<i>Tachinus.</i>	
	pectoris longitudine. Tarsi	{	5-articulati.	Antennae	{	capillares	{		<i>Habrocerus.</i>	
									filiformes.	filiformes
		{	4-articulati		Palpi max.	{	subulati		<i>Boletobius.</i>	
								<i>Tanygnathus.</i>		

## Trib. III. Staphylinini.

Labrum plerumque bilobum, limbo membranaceo coriaceove. Mandibulae lacinia membranacea interiore libera. Antennae 11-articulati. Ocelli nulli. Elytra pectoris longitudine. Coxae anticae conicae, exsertae, posticae conicae. Trochanteres postici simplices. Tibiae saepius spinulosae. Tarsi 5-articulati. Abdomen marginatum, segmento septimo retractili.

Stigmata prothoracica	libera, detecta.	Anten-	approximatae	<i>Xantholinini.</i>
	nae basi		distantes	<i>St. genuini.</i>
	obtecta			<i>Oxyporini.</i>

### Subtrib. 1. Xantholinini.

Antennae inter se non magis quam ab oculis distantes.

Antennae	rectae. Paraglossae	{	nullae	{	filiformes	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
----------	---------------------	---	--------	---	------------	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

### Subtrib. 2. St. genuini.

Antennae inter se magis quam ab oculis distantes.

Maxillarum mala exterior	elongata. Tarsi posteriores	Palpi articulo ultimo teretes	deplanati.	acuminato.	distantes	<i>Huemotodes.</i>
			Palpi articulo ultimo teretes	Pedes intermedii truncato	approximati	<i>Cordylaspis.</i>
	interiori subaequali. Antennae	rectae. Palpi maxillares	filiformes. Ligula	biloba. Pedes intermedii	distantes	<i>Scariphaeus.</i>
				integra. Femora	approximati	<i>Palaestrinus.</i>
				antica et postica infra spinulosa		<i>Staphylinus.</i>
				omnia mutica.		<i>Ocyus.</i>
			subulati			<i>Belonuchus.</i>
						<i>Philonthus.</i>
						<i>Heterothops.</i>
						<i>Acylophorus.</i>



Stigmata prothoracica utrumque processu triangulari corneo membraceove oblecta.

Palpi	{	utrique securiformes . . . . .	{	<i>Astrapaenus.</i>
		maxillares filiformes, labiales		<i>Quedius.</i>
			{	<i>Euryporus.</i>
			{	<i>Oxyporus.</i>

Labrum lateribus saepius membranaceum. Mandibulae elongatae, falcatae, acutae, medio dentatae. Palpi maxillares magis minusve elongati, articulo primo brevi, quarto minimo, saepe vix observando, labiales breves. Antennae sub frontis margine laterali insertae, 11-articulatae, filiformes. Ocelli nulli. Elytra pectoris longitudine. Coxae anticae conicae, exsertae. Trochanteres postici fulcrantes. Tarsi 5-articulati. Abdomen marginatum, segmento septimo retractili, saepius vix conspicuo.

[illegible]

Labrum corneum, simplex. Mandibulae elongatae, falcatae, acutae, medio dentatae. Palpi maxillares elongati, articulo primo brevi, quarto reliquis haud minore, labiales breves. Antennae 11-articulatae, filiformes. Ocelli nulli. Elytra pectoris longitudine. Coxae anticae conicae, exsertae. Trochanteres postici fulcrantes. Tarsi 5-articulati. Abdomen segmento septimo vix conspicuo.

Abdomen	{	marginatum. Palpi maxillares	{	transverso . . . . .	<i>Pinophilus.</i>
		articulo quarto	{	securiformi . . . . .	<i>Taenodema.</i>
		immarginatum.	{	coniunctis in patellam dilatatis . . . . .	<i>Palaminus.</i>
				<div> <div> Tarsi antici ar- ticularum 4 primis </div> <div> { securiformi acuminato </div> </div>	<i>Oedichirus.</i> <i>Procirrus.</i>



**Trib. VIII. Piestini.**

Labrum corneum, apice utrinque lacinia membranacea auctum. Palpi filiformes. Antennae sub frontis margine laterali insertae, 11-articulatae, filiformes vel moniliformes. Ocelli nulli. Thorax infra totus corneus. Elytra pectoris longitudine. Tarsi 5-articulati. Abdomen segmento septimo omnino abscondito.

Abdomen	{	immarginatum. Mandibulae	{	dentatae . . . . .	<i>Leptochirus.</i>
			{	muticae . . . . .	<i>Lispinus.</i>
	{	marginatum. Mandibulae	{	dentatae . . . . .	<i>Piestus.</i>
				muticae. Palpi maxillares articulo quarto praecedente	<i>Prognatha, Isomalus, Hypotelus.</i>

**Trib. IX. Phloeocharini.**

Labrum corneum, integrum. Paraglossae distinctae. Antennae sub frontis margine laterali insertae, 11-articulatae, filiformes. Ocelli nulli. Thorax infra pone coxas anticas membraneus. Elytra pectoris longitudine. Tarsi 5-articulati. Abdomen late marginatum, apicem versus sensim angustatum, segmento septimo haud exserto.

Palpi maxillares	{	filiformes . . . . .	<i>Olisthaerus.</i>
	{	subulati . . . . .	<i>Phloeocharis.</i>

**Trib. X. Omalini,**

Labrum corneum integrum, apice saepius limbo membranaceo bilobo prominulo auctum. Paraglossae nullae. Palpi filiformes. Antennae sub frontis margine laterali insertae, 11-articulatae. Ocelli duo. Thorax pone coxas anticas membraneus. Elytra pectore longiora. Tarsi 5-articulati. Abdomen tenuiter marginatum, segmento septimo haud exserendo.

Mandibulae	{	dentatae. Palpi max. articulo quarto	{	praecedenti aequali . . . . .	<i>Anthophagus.</i>
			{	praecedente triplo longiore . . . . .	<i>Lestea.</i>
			{	muticae . . . . .	<i>Arpedium.</i>
	{	muticae. Tibiae articulis 2 primis aequalibus, sub-elongatis. Maxillae malis articulis 4 primis bus, aequalibus	{	spinulosae . . . . .	<i>Acidota.</i>
				corneis, elongatis . . . . .	<i>Olophrum.</i>
				membraneis. Tibiae { muticae . . . . .	<i>Lathrimaeum.</i>
				{ sae . . . . .	<i>Deliphrum,</i>
				{ simplicibus . . . . .	<i>Omalium.</i>
				{ dilatatis . . . . .	<i>Anthobium.</i>

**Trib. XI. Proteinini.**

Labrum corneum, integrum. Paraglossae distinctae. Palpi filiformes, articulo secundo incrassato. Antennae sub frontis margine laterali insertae. Ocelli nulli aut singulus. Thorax infra totus corneus. Elytra pectore longiora. Abdomen segmento septimo haud exserendo.





## Tribus prima.

### Aleocharini.

*Stigmata prothoracica conspicua.*

*Antennae in fronte ad oculorum marginem anteriorem insertae.*

Labrum integrum, corneum, laciniis membraneis nullis.

Mandibulae haud prominentes, subfalcatae, limbo membranaceo margini interiori affixo intus fimbriatae, apice libero, acutiusculo, muticae plerisque, medio subdentatae Phloeocharidi, ante apicem acute dentatae Gymnusae.

Maxillae cardine stipiteque corneis, malis iam membranaceis, iam corneis.

mala exteriore cornea, basi apiceque membranaceis, hoc barbatulo; interiore mutica, apice introrsum oblique submarginata, sinu spinulis ciliato, membranacea, margine exteriore corneo: sic in plerisque;

mala exteriore subelongata, interiore apice pubescente in Myrmedonia,

mala interiore unco corneo simplice in Dinarda,

aut unco duplici terminata, in Lomechusa;

mala exteriore ut in praecedentibus, interiore elongata, cornea, apice uncinata in Silusa;

mala utraque elongata, interiore cornea, apice uncinata, ante apicem denticulata in Pronomaea, Diglossa, Silusa, Gymnusa.

Palpi maxillares 4-articulati, articulo primo minuto, secundo tertioque elongatis, quarto iterum minuto, subulato.

Mentum corneum.

Labium membraneum, magis minusve exserendum.

Ligula membranacea, angusta, inter palporum articulos primos sita.

apice bifida plerisque;

apice bifida, laciniis iterum bifidis Autaliae;

integra Gyrophaenae, Peliusae, Placusae, Euryusae, Lomechusae, Silusae, Myllaenae.

Paraglossae nullae plerisque;  
 obsoletae Tachyusae;  
 ligulae breviores Autaliae, Falagriae, Bolitocharae, Caloderae,  
 Aleocharae, Placusae, Dinardae;  
 ligulae aequales Myrmedoniae.  
 Palpi labiales 3-articulati plerisque;  
 4-articulati Aleocharae;  
 2-articulati Autaliae, Gyrophaenae, Peliussae, Placusae, Euryusae;  
 exarticulati, setacei Silusae, Pronomaeae, Diglossae, Myllaenae, Gymnusae.

Antennae ad oculorum marginem anteriorem insertae, articulis tribus primis paulo longioribus, secundo tertioque obconicis,

11-articulatae, filiformes vel apicem versus sensim leviterque incrassatae plerisque;

10-articulatae, clavatae Oligotae.

Oculi parvi, rotundati vel ovoides, laterales.

Caput exsertum, interdum inflexum, saepe deflexum, plerumque porrectum, nonnunquam pedunculo tenui brevique thoraci affixum.

Thorax (Prothorax) margine posteriore coleopterorum basin attingens.

Prosternum triangulare, spatio pone coxas anteriores membraneo.

Pectus metathoracis epimeris magnis, retrorsum prominentibus.

Scutellum minutum, triangulare.

Elytra pectori longitudine aequalia.

Abdomen segmentis sex conspicuis, septimo plerumque omnino occulto, stylis analibus plerumque haud exsertis.

Pedes graciliores, coxis anticis exsertis, conicis, posticis transversis, fere parallelopipedis; trochanteribus posticis fulcrantibus; tibiis plerumque infermibus, spinulis apicalibus minutissimis, tarsis

omnibus 5-articulatis Ocaleae, Caloderae, Phloeopora, Oxypodae, Aleocharae, Gymnusae;

posterioribus 5-, anticis 4-articulatis Myrmedoniae, Autaliae, Falagriae, Bolitocharae, Tachyusae, Homalotae, Gyrophaenae, Peliussae, Placusae, Euryusae, Dinardae, Lomechusae, Silusae, Pronomaeae, Myllaenae;

omnibus 4-articulatis Hygronomae, Oligotae, Diglossae; articulis primis tarsorum anticorum aequalibus, posteriorum longitudine sensim decrescentibus, articulo primo posticorum elongato in plerisque;

tarsis posterioribus articulis primis aequalibus aut subaequalibus in Autalia, Hygronoma, Phloeopora, Homalota, Silusa;

articulo ultimo elongato, clavato plerisque;

reliquis aequali, depresso Hygronomae;

breviusculo, incrassato Diglossae;

unguiculis tenuibus, leviter falcatis, simplicibus plerisque;

brevibus, fortiter curvatis Diglossae.

Sexus differentia saepius obscura; interdum antennis longioribus validioribusque, interdum abdominis segmentis aut anterioribus aut posteriori-



bus tuberculatis, aut segmento ventrali sexto apice triangulariter productas distinguitur.

Tarsi antici nunquam dilatati.

#### O b s e r v a t i o n e s.

1. Corpus Aleocharinorum, minutum, agile, saepius lineare, habitu quodam peculiari, forma tamen varians, aliis (e. g. Falagriis) Paederos, aliis (e. g. Oxypodis, Myllaenis, Gymnusi) Tachyporos simulantibus. Antennis ad oculos marginem anteriorem insertis facilius distingui hanc tribum primus docuit Com. Mannerheim.

2. Nititur haec tribus numerosa, extricatu omnium difficillima, generibus Aleochara et Lomechusa, ingeniose a Gravenhorstio conditis. Com. Mannerheim secundum corporis habitum et antennarum tarsorumque posticorum structuram non sine successu in genera plura eam dissoluit, e quibus, oris et tarsorum structura quantum valui accuratissime examinata, alia recepi, alia mutavi, alia castigavi, alia nova iis addidi.

### Myrmedonia Er.

Er. Col. March. I. p. 286.

*Maxillae mala interiore mutica, apice undique subtiliter pubescent.*

*Ligula brevis bifida; paraglossae ei aequales, subacuminatae.*

*Palpi labiales 3-articulati, articulis longitudine subaequalibus.*

*Tarsi antici 4-, posteriores 5-articulati, postici articulo primo elongato.*

Corpus oblongum vel elongatum, minus convexum, aliis apterum aliis alatum. Caput exsertum, porrectum, aut thoracis latitudine, aut eodem paulo angustius, rotundatum, basi constrictum, oculis mediocribus, modice prominulis. Labrum latiusculum, truncatum, mandibularum apicem haud obtegens. Mandibulae muticae. Maxillae (t. III. f. 21.) mala exteriore elongata, palpo maxillari subaequali, cornea, apice pubescente; interiore dimidio brevior, membranea, margine exteriore corneo, apice oblique truncata, densius pubescente. Palpi maxillares modice elongati, articulo secundo tertio brevior, hoc leviter incrassato, quarto minuto, subulato. Labium (t. III. f. 1.) mento transversum, leviter emarginatum, ligula brevi, palpum labialium articulo primo brevior, bifida, paraglossis ligula longitudine subaequalibus, latis, subacuminatis, margine interiore haud ciliatis. Palpi labiales articulis tribus longitudine decrescentibus compositi. Antennae crassiusculae vel crassae, articulis tribus primis magis elongatis, secundo tertio brevior, ultimo saepe subacuminato. Thorax coleopteris angustior, depressiusculus. Scutellum triangulare. Elytra apice oblique introrsum truncata, ad angulum exteriorem leviter sinuata. Abdomen coleopterorum latitudine, supra planum, subtus convexum, vivis revolutum. Pedes modice elongati, intermedii basi distantes, tarsi antici 4-, posterioribus 5-articulatis, his elongatis, articulis 4 primis longitudine decrescentibus.

#### O b s e r v a t i o n e s.

1. Corpus aliis subtiliter punctatum, pube tenuissima sericea vestitum, aliis laevigatum, nitidum. Oris partibus respectis hoc genus facile distinguitur paraglossis ligulae aequalibus et maxillarum mala interiore pubescente, spinulis nullis ciliata. Praeterea ab aliis generibus tarsorum structura, ab aliis coxis intermedii maiore spatio distantibus distinguitur. Antennae validiores et abdomen revolutum notas leviores habituales praebent.

2. Com. Mannerheim plerasque species Bolitocharis adiungens e sola M. canaliculata constituit genus proprium: Drusilla, nomen a Leach pro-

positum, prius autem a Swainson (Zool. Illustr. I. tab. 11) Papilionum generi impositum, quam ob rem Stephens (Illustr. V. p. 106) nomen Astilbus, a Dom. Dillwyn propositum, recepit. Idem autor deinde *M. fulgidae* gen. Zyras (Illustr. V. p. 430), et *M. limbatae*, humerali et funestae gen. Pella (ibid. p. 434), condidit, quae omnia, cum characteribus vagis aut nullis constituta invenio, Myrmedoniae subiungere non dubito.

3. *Porus ochraceus* Hope in opere Roylii infra citato, depictus, petenti mihi cum oris partibus praeparatis a cel. Hope amicissime ad examinandum communicatus est. Oris structura, haud satis exacte cognita a Dom. Westwood, acutissimo ceterum pectore, (ligula praecipue haud acuminata sed bifida) omnino eum huic generi adiungit, a quo hoc insectum nonnisi tibiis spinulosis et spinis apicalibus longioribus intractis discedit, quae notae mihi haud sufficere videntur, ut ex hac singula specie genus peculiare constituatur.

4. Species indigenae, quantum observavi, in formicetis degere solent, inde nomen.

### \* A p t e r a e.

X \* 1. *M. canaliculata*: Punctatissima, rufo-testacea, capite abdominisque cingulo postico nigris, thoracē oblongo, canaliculato impressoque. — Long. 2 lin.

Er. Col. March. I. 290. 6.

*Staphylin. canaliculatus* Fab. Mant. Ins. I. 221. 29. Ent. Syst. I. II. 528. 42. Syst. El. II. 599. 22. — Payk. Mon. Staph. 31. 23. Faun. Suec. III. 385. 23. — Ross. Faun. Etr. Mant. 98. 219. (Faun. Etr. ed. Helw. p. 443) — Oliv. Ent. III. 42. 21. 25. t. 3. f. 31. — Panz. Faun. Germ. 27. 13. Ent. Germ. 356. 32. — Marsh. Ent. Brit. 507. 25.

*Aleochara canaliculata* Grav. Micr. 68. 1. Mon. 148. 1. — Latr. Hist. nat. Crust. et. Ins. IX. 378. 1. t. 80. f. 7. Gen. Crust. et Ins. I. 301. 2. — Gyll. Ins. Suec. II. 392. 14. — Zetterst. Faun. Lapp. I. 96. 1. Ins. Lapp. 75. 1. — Sahlb. Ins. Fenn. I. 356. 21. — Guér. Iconogr. Regn. Anim. Ins. t. 10. f. 5.

*Drusilla canaliculata* Mannerh. Brachelytr. 85. 1. — Boisd. et Lucord. Faun. Ent. Paris. I. 534. — Runde Brachel. Hal. 31. 1.

*Paederus impressus* Rossi Faun. Etr. I. 254. 627. (ed. Helw. p. 314)

Habitat in Europa, sub lapidibus.

Aptera. Corpus angustum, punctatissimum, densius pubescens, capite abdominisque segmentis quarto quintoque nigris. Antennae validiusculae, fuscae, basi testaceae, articulo secundo tertio paulo brevior. Thorax oblongus, coleopteris angustior, subtiliter canaliculatus, dorso in femina postice, in mare secundum totam longitudinem impresso. Elytra thorace breviora. Abdomen reflexum, dorso parcius punctato pubescenteque, segmento ultimo maris emarginato. Pedes pallide testacei.

Mas thoracis disco late excavato, et abdominis segmento dorsali sexto apice emarginato distinctus.

Obs. Vicinat, ni fallor, inter nidos Formicae flavae.

\* 2. *M. plicata*: Punctatissima, obscure testacea, capite thoraceque subquadrato piceis, abdomine supra ante apicem piceo, segmento quarto supra medio elevato-plicato. — Long. 2 lin.

Er. Col. March. I. 289. 5.

Habitat Berolini, Dom. Kirstein.

Aptera, punctatissima, densius pubescens, parum nitida, capite thorace-

que nigro-piceis. Antennae capite thoraceque paulo longiores, validiusculae, articulo secundo tertio paulo brevior, apicem versus haud incrassatae, rufo-testaceae. Caput thoracis prope latitudine, suborbiculatum, fronte leviter convexa. Thorax coleopteris vix angustior, latitudine haud brevior, basi leviter rotundatus, apice truncatus, lateribus anterieus subrotundatis, posterius subsinuatis, dorso depressiusculus, basi obsolete impressus. Scutellum punctatissimum, nigrum. Elytra thoracis longitudine, testacea. Abdomen obscure rufo-testaceum, apicem versus nigrum, supra segmento tertio medio tumidulo, quarto medio secundum totam longitudinem late abrupte elevato, segmentis tribus primis confertim punctatis, reliquis laeviusculis, nitidis. Pectus piceum. Pedes testacei.

Obs. 1. Praecedente haud longior, at paulo latior et antrorsum haud angustata. Elytra thorace haud breviora, alarum autem rudimenta tantummodo exigua sub elytris conspiciuntur.

2. Abdominis dorsum in altero sexu forte simplex: mihi nonnisi unicum, in collectione Kirsteiniana olim contentum specimen visum.

### \*\* A l a t a e !

\* 3. *M. limbata*: Subtilissime punctulata, fusco-testacea, capite, elytrorum disco abdomineque ante apicem fuscis, thorace subquadrato, maris late profundeque, feminae obsolete canaliculato. — Long.  $2\frac{1}{4}$  lin.

Er. Col. March. I. 288. 3.

*Staphyl. limbatus* Payk. Mon. Staph. 51. 39. Faun. Suec. III. 399. 41. — Marsh. Ent. Britt. 509. 30.

*Al. limbata* Gyll. Ins. Suec. II. 400. 22. — Zetterst. Faun. Lappon. I. 99. 10. Ins. Lappon. 76. 13. — Sahlb. Ins. Fenn. I. 365. 37.

Mas: Thorace medio longitudinaliter excavato, abdominis segmento ultimo dorsali emarginato.

*Al. limbata* Grav. Micr. 69. 2. — Latr. Hist. nat. Crust. et. Ins. IX. 378. 3.

*Bolitochara limbata* Mannerh. Brachelytr. 81. 39.

*Bolitochara limbata* var. (*foveicollis* Dej.) Boisd. et Lacord. Faun. Ent. Paris. I. 551. 17.

*Staph. divisus* Marsh. Ent. Britt. 510. 33.

Fem.: Thorace obsolete canaliculato, abdominis segmento ultimo dorsali integro.

*Al. laevis* Grav. Micr. 95. 39. Mon. 169. 52. — Latr. Hist. nat. Crust. et Ins. IX. 389. 39.

*Al. limbata* Germ. Faun. Ins. Europ. 13. 7.

*Bolitochara limbata* Boisd. et Lacord. Faun. Ent. Paris. I. 550. 17. — Runde Brachel. Hal. 31. 10.

Habitat in Europae campis, sub lapidibus, Formicae flavae, nisi memoria me fallit, hospes.

Corpus subtilissime punctulatum, pube tenuissima grisea sericans, maris dilutius, feminae obscurius fusco-testaceum, capite, elytrorum, plerumque etiam thoracis disco, abdominisque fascia indeterminata ante apicem fuscis. Antennae sat validae, testaceae, articulo secundo tertio dimidio fere brevior. Caput thorace paulo angustius, fronte obsolete impressa, oris partibus testaceis. Thorax latitudine paulo brevior, coleopterorum prope latitudine,



basin versus subangustatus, lateribus leviter rotundatus, maris longitudinaliter excavatus, feminae aequalis, obsolete canaliculatus. Elytra thorace paululum longiora. Abdomen supra punctulatum, segmento dorsali ultimo in mare profunde emarginato. Pedes testacei.

Sexus differentia supra notata.

Obs. Abdomine dorso punctulato a proxime sequentibus facile distinguenda.

- X \* 4. *M. humeralis*: Subtiliter punctata, fusca, elytrorum humeris, abdominis basi pedibusque rufo-testaceis, thorace subquadrato, depresso, inaequali. — Long.  $2\frac{1}{2}$ –3 lin.

*Er. Col. March. I. 287. 1.*

*Aleochara humeralis* Grav. *Micr.* 90. 33. *Mon.* 169. 50. — *Latr. Hist. nat. Crust. et Ins.* IX. 387. 33. — *Gyll. Ins. Suec. II.* 399. 21. — *Zetterst. Faun. Lapp. I.* 99. 10. *Ins. Lapon.* 76. 12. — *Sahlb. Ins. Fenn. I.* 64. 36.

*Bolitochara humeralis* Mannerh. *Brachelytr.* 81. 38. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 549. 16. — *Runde Bruchel. Hal.* 30. 9.

Habitat in Europae silvis, Formicae fuliginosae socia.

Subtiliter punctulata, pube tenuissima grisea subsericans, fusca. Antennae validae, brunneae, basi dilutiores, articulo secundo tertio dimidio brevior. Caput thorace multo angustius, fronte aliis aequali, aliis leviter impressa. Thorax coleopterorum fere latitudine, latitudine sua parum brevior, lateribus parum rotundatis, depressus, basi supra scutellum et utrinque ad marginem lateralem leviter impressus, aut totus fuscus, aut obscure rufo-testaceus, disco fusco. Elytra thorace vix longiora, aut fusca, humeris rufo-testaceis, aut rufo-testacea, disco fusco. Abdomen supra laevigatum, nigro-fuscum, basi rufo-testaceum, subtus rufo-testaceum, posterius minus infuscatum. Pectus iam fuscum, iam obscure rufo-testaceum. Pedes rufo-testacei.

Obs. Specimina fronte impressa forte mares; in nonnullis etiam thorax per totam longitudinem obsolete canaliculatus, at aliae sexus differentiae me fugiunt. Specimen fronte fortius impressa, thorace late et sat profunde canaliculato valde insigne, e Suecia a Dom. Schönherr sibi communicatum, Dom. Schüppel Museo Regio dedit.

- X \* 5. *M. funesta*: Subtiliter punctulata, nigra, nitida, antennis pedibusque brunneis, thorace subquadrato, basi leviter foveolato. — Long.  $2\frac{1}{2}$  lin.

*Er. Col. March. I. 287. 2.*

*Aleochara funesta* Grav. *Mon.* 168. 49. — *Gyll. Ins. Suec. IV.* 488. 22–23. — *Sahlb. Ins. Fenn. I.* 66. 38.

*Bolitochara funesta* Mannerh. *Brachelytr.* 81. 39.

*Var.*: Fusca, elytrorum litura humerali abdominisque segmentorum anteriorum margine fusco-testaceis.

Habitat in Europae silvis, in Formicae fuliginosae societate semper mihi obvia.

Statura omnino praecedentis et parum minor, subtilius minus dense punctulata, nigra, subnitida, pube tenuissime grisea subsericans. Antennae capitis thoracisque longitudine, validae, rufo-testaceae brunneaeve, articulo secundo tertio dimidio brevior. Caput thorace angustius, subtilissime parcius punctulatum, vel fere laeve. Thorax coleopterorum prope latitudine, latitu-

dine sua paulo brevior, lateribus leviter rotundatus, basin versus paulo magis quam antrorsum angustatus, subdepressus, aequalis, basi foveola minuta obsoletiora supra scutellum impressus. Elytra thorace vix longiora, confer-tim subtiliter punctata. Abdomen supra laevigatum. Pedes picei vel brunnei.

Mas differt thorace obsolete canaliculato et abdominis segmento dorsali ultimo submarginato.

Variat colore dilutiore, fusco, antennis pedibusque rufo-testaceis, elytrorum litura vel macula humerali, abdominis segmentorum anteriorum, nonnunquam etiam thoracis limbo brunneo-testaceis.

Obs. 1. A M. humerali praeter colorem capite fere laevi differt. Varietas descripta primo intuitu M. limbata simulat, abdomine autem supra laevigato haud diffcultur distinguitur.

2. Specimina Austriaca, nominibus Al. fuscicornis, laticollis, grandicollis et brevicornis a Dahlio accepta communicavit Dom. Schüppel. A. laticollis a specie genuina nullo modo, A. fuscicornis nonnisi elytris magis brunneis differunt. A. grandicollis cum varietate dilutiore convenit, A. grandicollis Knochii autem, secundum collectionem Hoffmannseggianam speciei genuinae mas, idemque est A. brevicornis Dahl.

6. *M. rigida*: *Confer-tim punctata, nigra, subnitida, tenuiter albido-pubescent, antennis tibiis tarsisque rufo-brunneis, thorace transversim subquadrato, postice obsolete impresso.* — Long. 2 lin.

Habitat in Sardinia, Mus. Reg. Taurin. A. Dom. Prof. Gené ad examinandum communicata.

Habitu praecedentium, at minor, M. canaliculatae aequalis licet minus angustata, nigra, nitidula, pube brevi subtili albida cinerascens. Antennae capite thoraceque multo longiores, validae, brunneo-rufae. Palpi rufi. Mandibulae rufo-piceae. Labrum subtiliter carinatum, apice rufescens. Caput thorace parum angustius, fronte dense sat distincte punctata, convexa, antice subretusum, supra os leviter excavatum sublaevigatumque. Thorax coleopteris paulo angustior, latitudine paulo brevior, lateribus leviter rotundatus, basin versus subangustatus, basi rotundatus, apice late emarginatus, angulis anterioribus acutiusculis, posterioribus obtusis, subelevatus, dorso depressiusculus, densius subtiliter licet sat distincte punctatus, supra scutellum late obsolete impressus. Elytra thorace parum longiora, dense punctata. Abdomen supra segmentis quarto quintoque parce, reliquis crebre subtiliter punctatis. Pedes sat elongati, tibiis tarsisque rufo-brunneis.

Obs. M. funestae proxima, at minor, angustior, cinerascens, abdomine dorso punctato facile distinguenda.

\* 7. *M. lugens*: *Subtiliter punctulata, fusca, pedibus testaceis, thorace brevi, transverso, basi leviter foveolato.* — Long. 2 lin.

Er. Col. March. I. 289. 4.

Al. *lugens* Grav. Micr. 95. 40. Mon. 169. 54. — Latr. Hist. nat. Crust. et Ins. IX. 389. 40.

Var.: *Elytris basi testaceis.*

Habitat in Germaniae silvis, intra Formicae fuliginosae societates.

Praecedentibus brevior, fusco-nigra, subtiliter dense punctulata, pube tenuissima grisea subsericans. Antennae capitis thoracisque longitudine,

crassae, articulo secundo tertio vix dimidio brevior, ultimo, ut in praecedentibus subcompresso-acuminato, brunneae, basi dilutiores. Palpi rufo-testacei. Caput thorace angustius, sublaeve, fronte nonnunquam obsolete impressa. Thorax coleopterorum fere latitudine, latitudine sua duplo fere brevior, lateribus leviter rotundatis, subdepressus, basi supra scutellum obsolete impressus. Elytra thorace paulo longiora, fusca, basi extrorsum plerumque testacea. Abdomen supra laevigatum, nigrum, nitidum, segmentis anterioribus testaceo-marginatis. Pedes testacei.

- \* 8. *M. fulgida*: Nigra, nitida, thorace transverso, vage punctato, basi foveolato, elytris parce profundeque punctatis antennisque rufo-testaceis, pedibus flavis. — Long.  $2-2\frac{1}{2}$  lin.

*Aleochara fulgida* Grav. Mon. 163. 37.

Var.: *Elytris angulo apicali exteriore late nigro.*

*Zyras Haworthi* Steph. Illustr. V. t. 26. f. 3.

Habitat in Austria, Hungaria, Dom. Schüppel.

Statura praecedentium, a quibus corpore glabro, nitido discedit. Antennae capitis thoracisque fere longitudine, extrorsum leviter incrassatae, rufo-testaceae, articulo secundo tertioque aequalibus. Caput thorace paulo angustius, lateribus parce grosseque punctatum, nitidissimum, nigrum, ore obscure testaceo. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus leviter rotundatus, basin versus subangustatus, subdepressus, vage punctis grossis piliferis impressus, basi supra scutellum foveola parva notatus, niger, pernitidus. Elytra thorace paulo longiora, parce profundeque punctata, rufo-testacea, nitida. Abdomen supra laevigatum, posterius punctis parvis maioribus sparsum, nitidum, obscure testaceum, segmentis singulis supra basi obscurioribus, ultimis duobus totis nigris.

Obs. Varietas indicata, cuius specimen Austriacum dedit Dom. Schüppel, genuinis duplo maior, praecipue latior, abdomine dorso minus picescente, elytrorum angulo apicali exteriore late et distincte nigro, facile speciei peculiaris speciem prae se ferat, at notae satis distinctivae desiderantur. Specimen secundum, huic omnino congruens, Lutetiae lectum, in collectione Dom. Aubé vidi.

- \* 9. *M. marginicollis*: Nigro-picea, nitida, antennis pedibusque rufis, thorace transversim quadrato, basi obsolete foveolato, margine omnino rufo, abdomine segmento dorsali penultimo utrinque longitudinaliter sulcato. — Long.  $2\frac{1}{2}$  lin.

Habitat ad Promont. Bon. Spei, Dom. Prof. Lichtenstein.

Statura fere *M. funestae*, nigro-picea, nitida, glabra. Antennae capitis thoracisque longitudine, minus crassae, articulo secundo tertioque longitudine aequalibus, rufo-testaceae. Caput thorace paulo angustius, subglobosum, posterius parce subtiliter punctulatum, nigerrimum, pernitidum, ore testaceo. Thorax latitudine dimidio brevior, lateribus leviter rotundatis, basin versus subangustatus, disco leviter convexo, margine laterali depresso, vage subtiliter punctatus, piceus, nitidus, margine omni dilutiore rufescente. Elytra thorace vix longiora, obsolete punctata, tenuiter pubescentia, picea, minus nitida. Abdomen nigro-piceum, apice testaceum, subtus densius punctatum, supra laevigatum, segmento penultimo utrinque stria longitudinali profunde impresso. Pedes rufo-testacei.



10. *M. obscura*: »*Nigra, nitida, thorace, elytris pedibusque obscure brunneis.*«

*Staph. obscurus* Fab. *Syst. El. II.* 595. 29.

Habitat in India orientali. Mus. Reg. Hafniense.

»Antennae nigrae, basi apiceque brunneae. Caput, atrum ore brunneo. Thorax et elytra subpunctata, fusco-brunnea, immaculata. Abdomen atrum, ano brunneo. Pedes brunnei.« Fab.

Obs. Hanc speciem in Museo citato vidi, at mihi quidem haud accuratius examinata, tamen de eius loco et affinitatibus indubius sum. Affinis certe sequenti, nisi forte omnino eadem sit. Color, a Fabricio pro fusco-brunneo descriptus, potius rufo-testaceus est.

\* 11. *M. spinigera*: »*Rufo-testacea, parce punctata, nitida, capite abdomineque nigris, hoc supra segmento primo apice laminato, utrinque bispinoso, secundo quintoque medio dentatis.* — Long. 3 lin.

Habitat in Aegypto, Dom. Prof. Ehrenberg.

Caput thoracis prope latitudine, nigrum, opacum, subtiliter vage punctatum, ore flavo. Antennae capite duplo longiores, crassae, brunneae, basi rufo-testaceae, articulo secundo tertio duplo brevior. Thorax coleopteris paulo angustior, latitudine parum brevior, basin versus vix angustatus, depressus, undique tenuiter marginatus, parcius punctatus, rufo-testaceus, nitidus. Elytra thorace parum longiora, testacea, profunde punctata. Abdomen nigrum, nitidum, vage punctatum, segmento primo secundique lateribus obscure rufis, segmento primo supra margine apicali in laminam, segmenti secundi basin obtegentem, apice utrinque spinis duabus elongatis, interiore exteriori imposita eamque tegente, terminatam producto, segmento secundo supra medio dente compresso, segmento quinto tuberculo acuto armatis. Pectus et pedes rufo-testacei.

\* 12. *M. armata*: »*Brunnea, opaca, elytris fuscis, capite abdomineque nigris, hoc nitido, segmento primo apice sublaminato, utrinque unispinoso.* — Long.  $2\frac{1}{2}$  lin.

Habitat in Africa australi, Dom. Drège.

Statura praecedentis. Antennae capitis thoracisque longitudine, crassae, subcompressae, articulo secundo tertio plus duplo minore, fuscae, articulo tertio brunneo, primis duobus testaceis. Mandibulae et palpi rufo-testacei, labrum flavum. Caput thoracis prope latitudine, nigrum, parum nitidum, parcius subtiliter punctulatum, fronte leviter impressa. Thorax coleopteris paulo angustior, latitudine paulo brevior, basin versus subangustatus, lateribus pone medium subsinuatis, parum convexus, subtiliter minus dense punctatus, interstitiis subtilissime alutaceis, linea tenuissima canaliculatus, brunneo-testaceus, vix nitidus. Scutellum brunneo-testaceum. Elytra subtilissime alutacea, minus dense subtiliter punctata, fusco-nigra, basi brunnea, opaca. Abdomen supra laeve, subtus parcius punctatum, nigrum, subtus basi brunneum, segmento primo dorsali margine apicali paululum producto, intra marginem spina singula, elongata, subarcuata, acuta armato. Pectus brunneo-testaceum. Pedes testacei.

- \* 13. *M. speciosa*: *Rufa, opaca, capite, elytris abdominisque cingulo lato apicali atris, thorace oblongo, aequali.* — Long. ultra 3 lin.

Habitat in Aegypto, Dom. Prof. Ehrenberg.

Praecedentibus paulo maior. Antennae breves, crassae, rufo-brunneae, medio obscuriores, articulo secundo tertio duplo minore. Caput thoracis fere latitudine, rotundatum, atrum, ore testaceo, fronte convexiuscula, obsolete punctata. Thorax latitudine longior, coleopteris paulo angustior, subdepressus, subtilissime vage punctatus, postice obsolete canaliculatus, testaceo-rufus, opacus. Scutellum rufo-testaceum. Elytra thoracis longitudine, subtiliter punctata, atra, opaca, humeris piceis. Pectus, pedes et abdomen rufa, hoc segmentis duobus penultimis atris, dorso laevigato.

- \* 14. *M. atricapilla*: *Rufo-testacea, parce subtilissime punctata, capite abdominisque apice nigris, elytris subtiliter granulatis, fusciscentibus.* — Long.  $2\frac{1}{4}$  lin.

Habitat ad Promont. Bon. Spei, Dom. Krebs.

Statura praecedentium. Antennae capitis thoracisque longitudine, crassiusculae, testaceae, medio fusciscentes, articulo secundo tertio dimidio brevior. Caput thoracis latitudine, nigerrimum, nitidum, ore flavo, fronte impressa. Thorax latitudine parum brevior, coleopteris paulo angustior, basin versus leviter angustatus, lateribus rectis, depressus, rufo-testaceus, opacus, subtiliter vage punctatus. Elytra thorace vix longiora, subtiliter granulata, fusciscentia, humeris dilutioribus, opaca. Abdomen rufo-testaceum, segmentis duobus penultimis nigris. Pectus rufo-testaceum. Pedes pallidi.

- \* 15. *M. frontalis*: *Rufo-testacea, nitida, parce punctata, fronte nigra, thorace basi lateribusque impresso.* — Long. 2 lin.

Habitat ad Prom. Bon. Spei, Dom. Krebs.

Statura *M. limbatae*. at paulo angustior, rufo-testacea, nitida, undique vage subtiliter punctata. Antennae capite thoraceque longiores, sat validae, articulo secundo tertio parum brevior. Frons convexiuscula, cum vertice, nigra. Thorax latitudine brevior, basi lateribusque leviter rotundatus, subdepressus, basi fovea transversa, prope marginem lateralem utrinque fovea maiore oblonga impressus. Elytra thorace sesqui longiora. Abdomen segmentis duobus penultimis medio fusciscentibus.

16. *M. boops*: *Brunneo-testacea, nitida, capite nigro, thorace latiore, fronte convexa, thorace brevissimo, elytris parce subtiliter punctatis.* — Long.  $2\frac{1}{4}$  lin.

Habitat prope Puerto Cabello Columbiae. Collect. Dom. Moritz.

Statura fere praecedentium, fusco-testacea, nitida, glabra. Antennae capitis thoracisque fere longitudine, validiusculae, articulo tertio secundo sesqui longiore. Caput thorace paulo latius, transversum, convexum, fronte utrinque obsolete punctulata, nigrum, ore testaceo, oculis sat magnis convexis. Thorax coleopteris angustior, latitudine duplo brevior, apice truncatus, basi lateribusque rotundatus, supra leviter convexus, disco sparsim obsoletius punctatus. Elytra thorace sesqui longiora, subdepressa, parcius

subtiliusque punctata, margine laterali apicem versus infuscato. Abdomen elongatum, subtus longitudinaliter convexum, supra longitudinaliter concavum, margine laterali sat elevato, acuto, laevigatum, subtus parce punctulatum parceque pilosellum. Pedes dilutius testacei.

\* 17. *M. collaris*: *Rufo-testacea, nitida, capite, elytris, pectore anoque nigris, thorace subquadrato, basi foveolato.* — Long. 2 lin.

*Er. Col. March. I. 290. 7.*

*Staph. collaris* Payk. *Faun. Suec. III. 401. 43.* — *Oliv. Ent. III. 42. 37. 54. t. II. f. 13.*

*Aleocho. collaris* Grav. *Mon. 164. 43.* — *Gyll. Ins. Suec. II. 385. 8.* — *Zetterst. Ins. Lappon. 75. 6.* — *Sahlb. Ins. Fenn. I. 348. 6.* — *Steph. Illust. V. t. 26. f. 2.*

*Bolitochara collaris* Mannerh. *Brachelytr. 75. 1.* — *Boisd. et Lacord. Faun. Ent. Paris. I. 541. 1.* — *Runde Brachel. Hal. 29. 1.*

Habitat in Europae silvis.

Antennae crassae, pilosae, fuscae, basi rufo-testaceae, articulis 4 ultimis flavescentibus, secundo tertioque subaequalibus. Caput thorace paulo angustius, nigrum, nitidum, nigro-pubescent, parce punctatum, fronte sublaevi. Thorax coleopteris nonnihil angustior, latitudine parum brevior, lateribus leviter rotundatus, parum convexus, parce pubescens, sparsim punctatus, basi medio foveolatus, rufo-testaceus, nitidus. Elytra thorace paulo longiora, parce pilosa, profunde parcus punctata, nigra, nitida. Abdomen supra fere impunctatum, rufo-testaceum, segmentis quinto sextoque nigris. Pectus nigrum. Pedes flavi.

18. *M. ochracea*: *Testacea, nitida, elytris profundius punctatis, antennis crassis, capite parum longioribus.* — Long. 3 lin.

*Porus ochraceus* Hope *Royle Himalaja. Zool. t. 9. f. 10.*

Habitat in Nepalia, Mus. Dom. Hope.

Corpus totum testaceum, nitidum. Antennae capite parum longiores, crassae, articulo secundo tertio brevior, 4—10 brevibus, perfoliatis, ultimo ovato. Caput thoracis latitudine, basi parum constrictum, fronte plana, parce subtiliterque punctatum. Thorax coleopteris parum angustior, latitudine dimidio brevior, basin versus subangustatus, lateribus basique leviter, angulis posterioribus fortiter rotundatis, parum convexus, parce subtilissimeque punctatus. Elytra thorace paulo longiora, minus dense profunde punctata. Abdomen parallelum. Pedes breves, crassiusculi, tibiis omnibus extus spinulis brevissimis, apice calcaribus duobus fortioribus munitis.

Obs. Conf. quod supra in generis descriptione notavi.

\* 19. *M. coriacea*: *Nigra, nitidula, punctata, thorace angustiore, basi impresso, elytris disco late impressis: antennis, pedibus anoque piceis, femoribus posticis basi albis.* — Long. 2½ lin.

Habitat in Brasilia: specimen alterum e collectione Virmondiana, alterum a def. Sellow prope San João del Rey lectum.

Piceo-nigra, nitidula, parce griseo-pubescent. Antennae capite thoraceque longiores, graciles, piceae, basi rufescentes, articulo secundo tertio vix



breviore. Caput minutum, rotundatum, fronte convexiuscula, obsolete punctulata. Thorax coleopteris duplo angustior, latitudine parum brevior, lateribus antice leviter rotundatus, subdepressus, crebrius punctatus, postice impressus. Elytra thorace paulo longiora, oblique punctata, medio leviter impressa. Abdomen revolutum, dorso impunctato, laevi, subtus subtiliter vage punctatum. Pedes elongati, graciles, nigro-picei, geniculis fuscis, femoribus posticis basi albidis.

20. *M. camura*: Ferruginea, nitidula, capite globoso, nigro, thorace angustiore, transversim subquadrato, convexo, elytris scabratiss, abdomine latiusculo. — Long.  $1\frac{1}{2}$  lin.

Habitat prope Puerto Cabello Columbiae. Collect. Dom. Moritz.

Habitu omnino praecedentis, at plus duplo minor. Antennae capitis, thoracis elytrorumque prope longitudine, apicem versus sensim subincrassatae, articulo secundo tertio paulo brevior, totae testaceae. Caput thoracis latitudine, globosum, basi profunde constrictum, fronte convexa, laevissimum, nigrum, ore ferrugineo. Thorax coleopteris plus dimidio angustior, latitudine paulo brevior, angulis anterioribus rotundatis, posterioribus rectis, convexus, laevis, ferrugineus, nitidus. Scutellum ferrugineum. Coleoptera thorace basi dimidio, apice duplo latiora, leviter convexa, punctis crebris elevatis asperata, ferruginea, nitida. Abdomen latiusculum, medio coleopteris fere latius, ferrugineum, nitidum, supra planum vel leviter excavatum, recurvum, laeve, subtus convexum, apicem versus obsolete punctulatum, et pilis breviusculis rigidis nigris hispidulum. Pectus ferrugineo-testaceum. Pedes elongati, pallidius testacei.

- \* 21. *M. labida*: Testacea, abdomine ante apicem elytrisque fuscis, his basi margineque apicali flavo-testaceis, capite thoraceque transverso, basin versus subangustato piceo-rufescentibus. — Long.  $1\frac{1}{3}$  lin.

Habitat in Guyana Venezuelensi, Dom. Moritz.

Statura fere Homalotae socialis, subdepressa, nitidula, pube brevissima subtilissimaque vestita. Antennae capitis thoracisque longitudine, articulis secundo tertioque aequalibus, ultimo oblongo, acuminato, testaceae, articulis basalibus pallidioribus. Caput thoracis latitudine, parce subtilissime punctulatum, piceo-rufo-testaceum, ore angustato, testaceo. Thorax coleopteris parum angustior, transversus, latitudine plus dimidio brevior, basin versus subangustatus, basi leviter rotundatus, apice truncatus, parum convexus, confertim subtilissime punctulatus, picescenti-testaceus. Scutellum rufo-piceum. Elytra confertim subtilissime punctulata, fusca, basi margineque apicali flavo-testaceis. Abdomen testaceum, segmentis tribus penultimis fuscis. Pectus picescenti-testaceum. Pedes testacei.

22. *M. munda*: Fusca, elytrorum humeris margineque apicali pedibusque flavis, antennis basi apiceque testaceis, thorace basin versus subangustato. — Long.  $1\frac{1}{3}$  lin.

Habitat in Americae meridionalis ins. St. Thomae. Coll. Dom. Moritz.

Habitu et statura omnino praecedentis, piceo-fusca, nitidula, omnium subtilissime pubescens. Antennae capite thoraceque paulo longiores, articulo

secundo tertio brevior, ultimo oblongo, acuminato, fuscae, articulis duobus primis ultimisque tribus flavis. Caput thoracis prope latitudine, subtilissime punctulatum, fronte planiuscula, ore angustato cum palpis flavescente. Thorax coleopteris paulo angustior, latitudine parum brevior, basin versus angustatus, lateribus pone medium subsinuatis, apice truncatus, leviter convexus, confertim subtilissime punctatus, rufescenti-piceus. Elytra confertim subtilissime punctata, fusco-picea, angulo humerali margineque apicali pallide flavescentibus. Abdomen supra laeve, fusco-piceum, segmentis secundo tertioque margine apicali testaceo. Pedes flavi.

23. *M. inscita*: Nigra, antennarum basi, tibiis tarsisque testaceis, elytris fuscis, humeris margineque apicali albidis, thorace transverso, basin versus subangustato. — Long.  $1\frac{1}{3}$  lin.

Habitat in Columbia ad montes prope Puerto Cabello. Collect. Dom. Moritz.

Statura et summa affinitas praecedentium. Nigra, nitidula, omnium subtilissime pubescens. Antennae capitis thoracisque longitudine, articulo secundo tertio vix brevior, ultimo oblongo, acuminato, nigrae, basi sordide testaceae. Caput thoracis fere latitudine, ore angustato, fusciscente. Thorax coleopteris parum angustior, latitudine duplo fere brevior, basin versus leviter angustatus, lateribus subrotundatis, parum convexus, laevis. Elytra thorace parum longiora, confertim subtilissime punctulata, fusca, nitida, macula transversa humerali margineque apicali albidis. Abdomen supra laeve. Pedes nigri, tibiis tarsisque rufo-testaceis.

24. *M. turbida*: Picea, antennarum basi, pedibus, elytrorum macula humerali, abdominisque segmentorum anteriorum margine apicali testaceis, thorace transverso, basin versus subangustato. — Long.  $1\frac{1}{2}$  lin.

Habitat in valle Araguensi Columbiae. Coll. Dom. Moritz.

Statura praecedentium, at paululum maior, fusco-picea, nitidula, subtiliter pubescens. Antennae capitis thoracisque longitudine, articulo secundo tertioque subaequalibus, ultimo oblongiusculo, pilosellae, piceae, articulis duobus primis testaceis. Caput thoracis latitudine, laeve vel parce subtilissime punctulato, ore angustato, piceo-testaceo. Thorax coleopteris paulo angustior, latitudine duplo fere brevior, lateribus subrotundatis, basin versus modice angustatus, leviter convexus, subtiliter punctulatus, margine laterali pilosello. Elytra thorace paulo longiora, confertim subtiliter punctulata, humeris sordide flavescentibus. Abdomen supra laeve, segmentis 2—4 margine posteriore testaceo. Pedes tenues, flavo-testacei.

25. *M. parilis*: Nigra, elytris fusco-piceis, antennarum basi pedibusque luteis, thorace subquadrato, basin versus subangustato, dorso longitudinaliter obsolete impresso. — Long. 1 lin.

Habitat ad Caracas Columbiae. Coll. Dom. Moritz.

Statura praecedentium, at duplo minor, nigra, nitidula, elytris obscure fusco-piceis, tota confertissime subtilissimeque punctulata, pube subtilissima grisea sericans. Antennae capite thoraceque longiores, articulis 2 et 3 aequalibus, 4—10 elongato-subobconicis, ultimo elongato, acuminato, nigrae,

articulo primo luteo. Caput thoracis fere latitudine, orbiculatum, fronte obsolete canaliculata, ore luteo. Thorax coleopteris angustior, latitudine vix brevior, basin versus sensim subangustatus, parum convexus, canalicula sat lata antrorsum evanescente minus profunde impressus. Elytra thorace paulo longiora, margine apicali obsolete pallescente. Abdomen subtus segmentis anterioribus margine apicali testaceo. Pedes lutei.

Obs. Species 5 ultimae inter se peraffines, Homalotarum quibusdam speciebus habitu simillima, facile autem distinguuntur tarsis posticis elongatis; a Tachyusis coxis intermediis distantibus differunt, et oris structura, quantum absque dissectione perspicere licet, huius generis esse mihi videtur.

\* 26. *M. tumidula*: Brevis, ferruginea, densius cinereo-pubescent, capite abdomineque ante apicem fuscis, pedibus flavescentibus, thorace transverso, brevi, convexo. — Long.  $1\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Brevis, crassiuscula, ferruginea, nitida, pube depressa grisea spissiore vestita. Antennae minus crassae, capite thoraceque paulo breviores, articulo tertio secundo vix brevior, at tenuiore, testaceae, ante apicem fusciscentes. Caput thorace angustius, at maiusculum, basi vix constrictum, omnium subtilissime punctulatum, fuscum, oris partibus testaceis. Thorax transversus, latitudine plus duplo brevior, basi coleopterorum prope latitudine, leviter bisinuatus, angulis posterioribus rectis, apicem versus modice angustatus, supra satis convexus, omnium subtilissime punctulatus. Elytra thorace sesqui longiora, apice truncata, minus depressa, evidentius punctulata, nonnunquam fusciscentia. Abdomen breviusculum, lateribus leviter rotundatum, supra parcius punctulatum, ante apicem fusciscent. Pedes flavescenti-testacei.

Obs. Non indubius hanc speciem ad hoc genus refero, habitus enim alienam quandam speciem prae se ferre videtur, et oris structura mihi haud perspecta. Differt autem ab Oxypodis et Aleocharis tarsis anticis 4-articulatis, ab Homalotis tarsis posticis articulo primo elongato: his notis et pedibus intermediis basi interstitio sat lato distantibus cum Myrmedoniis convenit, neque alii cuidam generi adscribi potest.

### ***Autalia* Leach.**

*Mannerh. Brachelytr. p. 87. — Er. Col. March. I. p. 291.*

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula elongata bifida, laciniis bifidis, lacinia interiore perbrevis, exteriori lineari; paraglossae parvae, angustae, acuminatae.*

*Palpi labiales biarticulati.*

*Tarsi anticis 4-, posteriores 5-articulati, postici articulis 4 primis aequalibus.*

Corpus minutum, minus depressum, alatum. Caput exsertum, porrectum, basi profunde constrictum, collo quasi tenui thoraci affixum, suborbiculatum, thorace paulo latius, oculis mediocribus, parum prominulis. Labrum latiusculum, semi-orbiculare. Mandibulae parvae, muticae. Maxillae mala exteriori cornea, apice membranacea pubescenteque, interiori exteriori longitudine aequali, extus cornea, intus membranacea, apice introrsum oblique emarginata, sinu spinulis brevibus ciliato. Palpi maxillares modice elon-



gati, articulo secundo tertioque subaequalibus, hoc apicem versus leviter incrassato, quarto minuto, subulato. Labium (t. III. f. 2.) mento sat magno, antrorsum leviter angustato, apice profunde inciso, lobis rotundatis; ligula angusta, elongata, apice bifida, lacinii iterum bifidis, laciniosis exterioribus longioribus extrorsum, interioribus brevioribus antrorsum vergentibus, paraglossis parvis, angustis, acuminatis, interne ciliatis. Palpi labiales biarticulati, articulo primo cylindrico, secundo hoc vix brevior, apice leviter incrassato. Antennae capitis thoracisque fere longitudine, crassiusculae, articulo primo cylindrico, secundo tertioque obconicis, magis elongatis, his inter se subaequalibus. Thorax coleopteris duplo fere angustior, subquadratus, angulis anterioribus rotundatis, apice attenuatus. Scutellum triangulare. Coleoptera sat ampla, abdomine paulo latiora, lateribus leviter rotundata, apice ad angulum exteriorem vix sinuata. Abdomen lineare, supra planum, subtus convexum, recurvum. Pedes mediores, intermedii basi leviter distantes, tarsi antici 4-, posteriores 5-articulati, parum elongati, articulis 4 primis subaequalibus.

#### Observationes.

1. Maxillae omnino ut in *Homalotis* (t. III. f. 22.), labium contra in hoc genere perquam singulare; paraglossae acuminatae affinitatem cum *Falagria* produnt, mentum autem profunde incisum et ligula elongata, apice quadrifida huic peculiaris. Palpi labiales certe biarticulati.

2. Habitus thorace apice attenuato et capite remoto *Falagria* affinis, discedit autem thorace impressionibus longitudinalibus plicato, et tarsis posticis haud elongatis, articulis 4 prioribus longitudine subaequalibus. Sunt autem quaedam *Homalotae* species (*H. didyma* et *bisulcata*), quae thoracis et capitis a thorace remoti forma, tarsorum structura et omni habitu cum *Autaliis* conveniunt, ut non nisi oris examine instituto ab iis distinguantur.

3. Sexus differentia mihi haud observata.

\* 1. *A. impressa*: Rufo-testacea, capite abdominisque postico nigricantibus, thorace sulculo antico foveisque quatuor posticis longitudinalibus impressis. — Long. 1 lin.

*Mannerh. Brachelytr.* 87. 2. — *Boisd. et. Lacord. Faun. Ent. Paris. I.* 558. — *Er. Col. March. I.* 292. 1.

*Staphyl. impressus* Oliv. *Ent. III.* 42. 23. 28. t. 5. f. 41.

*Aleoeh. impressa* Grav. *Micr.* 72. 7. *Mon.* 150. 4. — *Latr. Hist. nat. Crust. et Ins. IX.* 380. 7. *Gen. Crust. et Ins.* 302. 3. — *Gyll. Ins. Suec. II.* 381. 4. — *Sahlb. Fenn. I.* 346. 3.

◀Habitat in Europae boletis.

Obscure rufo-testacea, nitida, subtiliter pubescens. Caput nigricans, subtilissime punctatum. Thorax ante medium sulculo medio, postice sulculis duobus obliquis, abbreviatis, interioribus, et utrinque foveola exteriore rotundata impressus. Elytra fusco-testacea, subtilissime punctata, basi bifoveolata. Abdomen pone medium nigricans, summo apice dilute testaceo. Pedes testacei.

\* 2. *A. rivularis*: Nigra, nitida, antennis pedibusque piceis, thorace sulculo medio foveolisque quatuor basalibus impresso. — Long.  $\frac{5}{6}$  lin.

*Mannerh. Brachelytr.* 87. 1. — *Er. Col. March. I.* 293. 2.

*Aleoeh. rivularis* Grav. *Micr.* 73. 8. *Mon.* 150. 5. — *Latr. Hist. nat. Crust. et Ins. IX.* 380. 8. — *Gyll. Ins. Suec. II.* 382. 5. — *Sahlb. Ins. Fenn. I.* 347. 4.

Habitat in Europa.

Praecedente paulo minor, nigra, nitida, parce subtiliter pubescens, antennis pedibusque rufo-piceis. Thorax sulco medio, antice abbreviato, ad basin autem continuato, foveisque quatuor basalibus impressus, his iisdem ut in praecedente, licet profundioribus. Elytra basi bifoveolata, vix punctulata. Abdomen apice piceum.

### **Falagria Leach.**

*Mannerh. Brachelytr. p. 86. — Er. Col. March. 1. p. 293.*

*Maxillae mala inferiore mutica, intus spinulis ciliata.*

*Ligula elongata, linearis, apice bifida, paraglossae parvae, angustae, acuminatae.*

*Palpi labiales triarticulati, articulo secundo minore.*

*Tarsi antici 4-, posteriores 5-articulati, postici articulo primo elongato.*

Corpus minutum, minus depressum, alatum. Caput exsertum, porrectum, basi profunde constrictum, collo quasi tenui thoraci affixum, suborbiculatum, thoracis fere latitudine, oculis mediocribus, haud prominulis. Labrum mediocre, semiorbiculatum. Mandibulae parvae, muticae. Maxillae mala exteriore cornea, apice membranea, pubescenteque, interiore exteriore breviori, membranea, margine exteriore corneo, apice introrsum oblique emarginata, sinu spinulis brevibus ciliato. Palpi maxillares modice elongati, articulo secundo tertioque subaequalibus, hoc apicem versus sensim incrassato, quarto parvo, subulato. Labium (t. III. f. 3.) mento transversum, apice leviter emarginato, ligula angusta, haud ultra primum palporum articulum porrecta, apice bifida, paraglossis parvis acuminatis, interne ciliatis. Palpi labiales tri-articulati, articulo primo subcylindrico, secundo hoc duplo fere breviori, parum discretò, tertio paulo tenuiore, apice leviter incrassato. Antennae capiti thoracique longitudine aequales vel paulo longiores, crassiusculae, apicem versus plerumque parum incrassatae, articulis tribus primis magis elongatis, secundo tertioque obconicis, subaequalibus. Thorax basin versus angustatus, apice obtuse angulatus. Scutellum triangulare. Coleoptera abdominis latitudine, apice truncata, ad angulum exteriorem vix sinuata. Abdomen lineare, supra planum, subtus convexum, saepe recurvum. Pedes modice elongati, intermediis basi leviter distantes, tarsi antici 4-, posteriores 5-articulati, elongati, articulis 4 primis sensim decrecentibus.

### **Observationes.**

1. Maxillae omnino ut in *Homalota* (t. III. f. 22.). Paraglossae acuminatae ut in *Autalia* et *Bolitochara*. Cum *Bolitochara* summa similitudo in oris structura, at ligula minus elongata, palporum labialium articulum primum vix superans. Palpos labiales olim, licet non sine dubio quodam, biarticulatos descripsi, dein autem dissectione et examine reiteratis distincte triarticulatos vidi. Articulus secundus a primo minus distinctus.

2. Habitus huius generis gracilis, elegans, corpore plerumque subtilissime punctulato vel sublaevi, tenuiter subtilissimeque pubescente. Distinguitur facile a generibus affinibus thorace subcordato vel orbiculato, convexiusculo, et capite remoto. Thorax saepe, nonnunquam etiam scutellum canaliculata.

3. Differentia sexus in plerisque speciebus mihi non observata: in nonnullis mares fronte impressa cognoscuntur.

4. A Dom. Curtis loco infra apud F. thoracicam citato genus Falagria illustratum, at oris partes haud satis accurate observatae neque correcte delineatae. Ligula pictorem omnino fugit. Tarsos anticos elegantissimus auctor recte 4-articulatos vidit, posteriores autem observandos neglexit.

\* 1. *F. sulcata*: Nigro-picea, thorace subcordato scutelloque profunde canaliculatis, coleopteris quadratis, subtilissime punctulatis. — Long. 1 lin.

Mannerh. *Brachelytr.* 86. 1. — Boisd. et Lacord. *Faun. Ent. Paris.* I. 556. 2. — Runde *Brachel. Hal.* 31. 1.

*Staph. sulcatus* Payk. *Mon. Staph.* 32. 24. — Schrank *Faun. Boic.* I. 648. 888. — Oliv. *Ent.* III. 42. 23. 27. t. 6. f. 52. — Marsh. *Ent. Britt.* 507. 25.

*Al. sulcata* Grav. *Micr.* 73. 9. *Mon.* 150. 6 — Latr. *Hist. nat. Crust. et Ins.* IX. 380. 9. — Gyll. *Ins. Suec.* II. 378. 1. — Sahlb. *Ins. Fenn.* I. 345. 1.

*Falagria caesa* Er. *Col. March.* I. 295. 2.

Habitat in Europa.

Nigro-picea seu fusco-picea, nitida, pube subtili densiore grisea sericans. Antennae capitis thoracisque longitudine, crassiusculae, totae fuscae. Palpi fusi. Caput thoracis prope latitudine, fronte depressiuscula, omnium subtilissime punctulata, inter antennis obsolete bi-impressa. Thorax latitudine summa paene brevior, ante medium fortiter ampliatus, basin versus sensim angustatus, dorso subdepressus, subtilissime punctulatus, sulco profundo, apice summo abbreviato, basi dilatato canaliculatus. Scutellum bicarinatum. Coleoptera latitudine sua haud breviora, thorace longiora, subtilissime punctata. Abdomen confertim punctulatum, densius griseo-pubescent. Pedes fusco-testacei.

Obs. Nomen vulgatum restitui. Primo enim haec species a Paykul loco supra citato »antennis fusciscentibus« descripta est. *Staph. sulcatum* eiusdem auctoris in *Fauna Suecica* »antennis basi rufis« memoratum alteram speciem, *Al. sulcatulam* Grav., esse patet.

\* 2. *F. dissecta*: Nigro-picea, thorace subcordato scutelloque profunde canaliculatis, coleopteris transversis, subtiliter punctulatis. — Long. 1 lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Praecedenti simillima, nigro-picea, nitida, pube brevi subtili griseo-subsericans. Antennae capitis thoracisque longitudine, crassiusculae, nigro-piceae. Palpi picei. Caput thoracis latitudine, nigerrimum, subtilissime punctulatum, fronte convexiuscula. Thorax latitudine summa paulo brevior, ante medium fortiter rotundato-ampliatus, leviter convexus, subtiliter punctulatus, sulco profundo, apice abbreviato canaliculatus. Scutellum bicarinatum, inter carinas canaliculatum. Coleoptera latitudine sua paululum breviora, thorace paulo longiora, confertim subtiliter punctulata. Abdomen subtiliter punctulatum. Pedes picei, tarsi testacei, vel rufo-testacei, tarsi dilutioribus.

Obs. A praecedente aegerrime distinguenda, at revera specie distincta videtur. Differt fronte convexiuscula, thorace lateribus magis rotundato, sulco basi haud dilatato, elytris brevioribus, distinctius punctulatis, pedibus obscuris, tarsis solis testaceis.



- \* 3. *F. sulcatula*: *Picea*, *thorace subcordato, scutelloque profunde canaliculatis, coleopteris transversis, laevibus*. — *Long. vix 1 lin.*

*Aleochar. sulcatula* Grav. *Mon.* 151. 7.

*Staph. sulcatus* Payk. *Faun. Suec.* III. 385. 24.

*Aleochar. sulcata* var. *b* Gyll. *Ins. Suec.* II. 378. 1. — *Sahlb. Ins. Fenn.* I. 345. 1.

*Falagria sulcata* Er. *Col. March.* I. 294. 1.

*Fal. polita* Curtis *Britt. Ent.* X. f. 462. n. 4.

Habitat in Europa.

*Picea*, vel interdum rufo-picea, abdomine pone medium nigricante, nitida, parcissime subtilissimeque pubescens, haud sericans. Antennae capitis thoracisque prope longitudine, piceae, basi rufo-testaceae. Palpi picei. Caput thoracis latitudine, fronte depressiuscula, vix perspicue punctulata. Thorax latitudine summa haud brevior, antrorsum rotundato-amplius, basin versus sensim aequaliter angustatus, leviter convexus, vix perspicue punctulatus, sulco profundo longitudinali, apice summo abbreviato, basi subdilatato, exaratus. Scutellum bicarinatum, inter carinas canaliculatum. Coleoptera thorace haud longiora, latitudine sua distincte breviora, laevia. Abdomen dorso parce punctulatum. Pedes rufo-testacei.

Obs. Affinis certe *F. sulcatae*, at haud difficulter distinguenda statura strictiore, thorace longiore, coleopteris brevioribus, haud perspicue punctulatis, corpore haud sericante, magis rufo-piceo, antennis basi rufo-testaceis. De synonymis conf. quae apud illam ad calcem observavi.

- \* 4. *F. discisa*: *Nigra, nitida, elytris antennisque fuscis, his apice pedibusque testaceis, thorace subcordato scutelloque profunde canaliculatis*. — *Long. 1 lin.*

Habitat in Brasilia. Benevole dedit Dom. Melly.

Statura praecedentium, nigra, nitida, omnium subtilissime pubescens. Antennae capite thoraceque longiores, basi rufo-brunneae, medio fuscae, apice sensim testaceae. Caput subtiliter vage punctatum, vertice longitudinaliter impresso. Thorax antice valde rotundato-amplius, basin versus fortiter angustatus, ante basin leviter constrictus, latitudine summa vix brevior, parum convexus, laevissimus, sulco profundo canaliculatus. Scutellum canaliculatum. Elytra fusco-testacea, nitida, subtilissime parce punctulata. Abdomen supra subtilissime obsolete punctulatum. Pedes testacei.

- \* 5. *F. scissa*: *Rufo-picea, tenuiter pubescens, capite thoraceque fusciscentibus, hoc subcordato, punctatissimo, postice utrinque foveolato scutelloque canaliculatis*. — *Long. 1 $\frac{2}{3}$  lin.*

Habitat prope Puerto Cabello in Columbia, Dom. Moritz.

Praecedentibus paulo maior, rufo-picea, capite thoraceque fusciscentibus, subnitidis. Antennae capite thoraceque paulo longiores, crassiusculae, rufo-brunneae, ante medium fusciscentes. Caput subtiliter parcius punctatum, tenuissime pubescens, vertice obsolete impresso, fronte inter antennas leviter bi-impressa. Thorax latitudine summa paulo brevior, lateribus valde amplius, basin versus valde coarctatus, prope basin leviter constrictus, subdepressus, confertissime punctatus, subtilissime pubescens, canalicula profunda media foveolaque laterali utrinque pone medium impressus. Scutel-

lum canaliculatum. Elytra fusco-testacea, subtilissime vage punctulata, dense subtilissime pubescentia. Abdomen parce punctatum, apicem versus piceum. Pedes testacei.

- \* 6. *F. divisa*: *Picea, thorace subcordato, profunde canaliculato, rufo-piceo, elytrorum humeris, abdominis et antennarum basi pedibusque testaceis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Brasilia, Dom. Melly.

Praecedentibus paulo gracilior, *F. sulcata* paulo maior. Antennae longiores, piceae, basi testaceae (ultimi tres articuli desunt). Caput piceum, nitidum, laeve. Thorax latitudine summa longior, ante medium rotundato-amplius, rufo-piceus, nitidus, laevis, profunde canaliculatus, inter canaliculam subtiliter punctatus. Scutellum obsolete bicarinatum, punctatum, rufo-piceum. Elytra thorace parum longiora, vage subtilissime punctata, nitida, fusca, margine apicali humerisque sordide testaceis, pone scutellum impressione rotundata communi notata. Abdomen subtilissime densius punctatum pubescensque, obscure piceum, segmento primo secundoque testaceis, secundo obscuriore, subtus rufo-piceum, segmento primo flavo. Pedes testacei.

- \* 7. *F. concinna*: *Nigra, nitida, thorace subcordato, profunde canaliculato, piceo, elytris fuscis, basi apiceque pallide testaceis, abdominis segmento primo, antennarum basi apiceque pedibusque testaceis, his femoribus nigris.* — Long. 1 lin.

Habitat in Brasilia. Amice dedit Dom. Prof. Germar.

Affinis praecedenti, at minor, statura omnino *F. sulcatae*. Antennae capitis thoracisque longitudinem prope superantes, validiusculae, piceae, articulis tribus primis ultimoque testaceis. Caput thorace paululum latius, laevissimum, nigrum, nitidum. Thorax latitudine summa vix longior, antice rotundato-amplius, basin versus angustatus, convexiusculus, laevissimus, profunde canaliculatus, rufescenti-piceus, nitidus. Scutellum subtilissime bicarinatum, testaceum. Elytra thorace haud longiora, omnium subtilissime punctulata, pube subtilissima tenuissimaque grisea vestita, fusca, nitida, basi margineque apicali pallide testaceis, pone scutellum foveola rotundata communi impressa. Abdomen subtiliter punctulatum griseoque pubescens, nigrum, segmento primo dilute flavo-testaceo. Pectus nigrum. Pedes coxis anticis fuscis, posterioribus dilute flavo-testaceis, femoribus omnibus nigris, basi dilute testaceis, tibiis tarsisque dilute rufescenti-testaceis.

- \* 8. *F. stricta*: *Picea, nitida, antennis pedibusque piceo-testaceis, basi pallidis, thorace elongato, profunde canaliculato.* — Long.  $1\frac{1}{4}$  lin.

Habitat ad Caracas Columbiae, Dom. Moritz.

*F. sulcatae* magnitudine aequalis, at longior et gracilior, brunneo-picea, nitida, omnium subtilissime tenuissimeque cinereo-pubescens. Antennae capitis thoracisque longitudine, crassiusculae, piceo-testaceae, medio fusciscentes, basi apiceque pallidiores. Palpi piceo-testacei. Caput laeve, fronte inter oculos obsolete transversim impressa. Thorax coleopteris duplo angustior, latitudine sua summa multo longior, ante medium rotundato-amplius, basin versus angustatus, modice convexus, laevissimus, profunde canaliculatus, postice subcompressus, ante basin utrinque sulculo transverso

impressus. Scutellum obsolete bicarinatum. Coleoptera laevigata, infra scutellum impressa. Abdomen laeve, segmentis primis duobus apice testaceis. Pedes piceo-testacei, basi pallidi, tarsis flavescentibus.

- \* 9. *F. percisa*: *Fusco-picea, nitida, subtiliter pubescens, fronte thoraceque subcordato profunde canaliculatis, antennarum basi pedibusque testaceis.* — Long. vix ultra 1 lin.

Habitat in Brasilia, Dom. Prof. Germar.

*F. sulcata* perparum maior, at paulo magis depressa, fusco-picea, nitida, pube subtili densiore grisea vestita. Antennae capitis thoracisque longitudine, piceae, articulis duobus primis testaceis. Palpi testacei, articulo tertio piceo. Caput thorace paulo latius, subquadratum, angulis licet rotundatis, subdepressum, subtilissime obsolete punctulatum, per totam longitudinem canaliculatum, sulco in media fronte profundiore dilatatoque. Thorax coleopteris dimidio angustior, latitudine summa haud brevior, ante medium rotundato-amplius, basin versus modice angustatus, minus convexus, obsolete punctulatus, sulco profundo canaliculatus. Scutellum confertissime punctulatum, obsolete bicarinatum. Elytra thorace parum longiora, fusco-subtestacea, confertim subtilissime punctulata. Abdomen confertim punctulatum. Pedes rufo-testacei.

- \* 10. *F. thoracica*: *Rufo-picea, thorace subcordato, profunde canaliculato pedibusque rufis, abdomine nigro.* — Long.  $1\frac{1}{4}$  lin.

Curt. Britt. Ent. X. t. 462.

*Fal. lineolata* Boisd. et Lacord. Faun. Ent. Paris. I. 555. 1.

Habitat in Germania, Helvetia, Dom. Schüppel et Walth, in Gallia, Dom. Aubé, in Anglia, Curtis. Degit in fornicetis observante Dom. Walth.

Statura *F. sulcatae* at gracilior, subtilissime confertim punctata, pube tenuissima cinerea vestita. Antennae piceae, capitis thoracisque longitudine, basi apiceque summo rufo-testaceae. Palpi testacei. Caput piceum, fronte depressiuscula. Thorax latitudine summa haud longior, basin versus modice angustatus, sulco profundo canaliculatus, dilute rufus. Scutellum aequale, confertissime punctatum. Coleoptera thorace paulo longiora, quadrata, fusca. Abdomen vel totum nigrum, vel piceum, ante apicem nigrum. Pedes dilute rufo-testacei.

- \* 11. *F. denigrata*: *Nigerrima, nitida, subtiliter griseo-pubescens, capite, thorace elytrisque dense subtiliter punctatis, thorace canaliculato.* — Long.  $1\frac{2}{3}$ —2 lin.

Habitat in Mexico, Dom. Schleiden.

Statura omnino *F. sulcatae*, at duplo maior et ultra, tota nigra, pube subtili grisea sericeo-pubescens. Antennae capitis thoracisque longitudine, fortiusculae, nigrae. Palpi nigri. Caput creberrime minus subtiliter punctatum, obsolete canaliculatum. Thorax coleopteris plus dimidio angustior, ante medium rotundato-amplius, basin versus leviter angustatus, minus convexus, dense subtiliter, disco obsolete punctatus, profunde canaliculatus, canalicula basi impressione parva transversali terminata. Scutellum aequale,



confertim subtiliter punctatum. Elytra thorace sesqui fere longiora, dense subtiliter punctata. Abdomen parce subtiliter punctulatum. Pedes nigri, tarsis piceis.

- \* 12. *F. discreta*: Fusco-picea, nitidula, abdominis postico capiteque nigris, thorace subcordato, subtiliter punctulato profunde canaliculato. — Long.  $1\frac{1}{4}$  lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Praecedenti affinis, at dimidio minor. Antennae capitis thoracisque longitudine, crassae, fusco-piceae. Caput nigrum, nitidum, subtilissime obsolete punctulatum, fronte depressiuscula. Thorax antice rotundato-ampliat, basin versus fortiter angustatus, latitudine summa paulo brevior, subdepressus, dense subtilissime punctulatus, profunde canaliculatus, fusco-piceus, nitidus, subtilissime cinereo-pubescens. Scutellum subtilissime punctulatum, linea tenui longitudinali elevata notatum, fusco-piceum, subtilissime pubescens. Elytra thorace paulo longiora, lucide fusco-picea, subtilissime punctulata, subtiliter cinereo-pubescentia. Abdomen parce subtilissime punctulatum, griseo-pubescens, segmentis anterioribus tribus dilute fusco-piceis, posterioribus tribus nigro-piceis, nitidis. Pectus dilute piceum. Pedes piceo-testacei, tarsis flavescentibus.

- \* 13. *F. fissula*: Fusco-picea, nitida, capite abdominisque postico nigris, antennarum basi apiceque pedibusque testaceis, thorace cordato, subtiliter punctulato, profunde canaliculato. — Long.  $\frac{3}{4}$  lin.

Habitat in Guyana Venezuelensi, Dom. Moritz.

Habitu omnino *F. sulcatae*, at dimidio minor, *F. nigrae* aequalis, piceo-fusca, nitidula, subtilissime griseo-pubescens. Antennae capitis thoracisque longitudine, validiusculae, piceae, articulis duobus primis tertioque testaceis. Palpi testacei, articulo tertio fusco. Caput thoracis latitudine, obsoletissime punctulatum, nigrum, nitidum. Thorax latitudine summa parum brevior, antice valde rotundato-ampliat, basin versus fortiter angustatus, parum convexus, subtiliter punctulatus, sulco profundo exaratus. Scutellum aequale. Coleoptera thorace paulo latiora et longiora, vix perspicue punctulata, dilutius fusco-picea. Abdomen longius pubescens, segmentis penultimis duobus nigris. Pedes dilute flavo-testacei.

14. *F. soluta*: Picea, capite abdominisque apice nigris, thorace, abdominis basi pedibusque rufo-testaceis, thorace canaliculato. — Long. 1 lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Habitu omnino *F. obscurae*, at paululum minor. Antennae capitis thoracisque longitudine, nigrae, articulis duobus primis testaceis. Palpi testacei. Caput nigrum, nitidum, laeve. Thorax anterieus lateribus modice dilatatus, leviter convexus, laevis, profunde canaliculatus, rufo-testaceus, nitidus. Scutellum aequale, rufo-testaceum. Elytra laevia, picea, nitida. Abdomen laeviusculum, segmentis anterioribus tribus rufo-testaceis, posterioribus tribus nigris. Pectus piceum. Pedes testacei.

- ✕ \* 15. *F. obscura*: *Rufo-testacea, thorace subcordato, obsolete canaliculato, capite abdomineque obscuris.* — Long. 1 lin.

*Mannerh. Brachelytr.* 87. 2. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 556. 3. — *Runde Brachelytr. Hal.* 31. 2. — *Er. Col. March. I.* 295. 3.

*Aleochar. obscura* Grav. *Micr.* 74. 10. *Mon.* 151. 8. — *Latr. Hist. nat. Crust. et Ins.* IX. 381. 10. — *Gyll. Ins. Suec. II.* 379. 2. — *Sahlb. Ins. Fenn. I.* 346. 2.

Habitat in Europa.

Statura fere *F. sulcatae*, obscure rufo-testacea, nitida, densius et minus subtiliter pubescens. Antennae capitis thoracisque longitudine, apicem versus sensim leviter incrassatae, corpori concolores. Caput thoracis latitudine, orbiculatum, fronte convexa, nigrum, nitidum. Thorax coleopteris dimidio angustior, latitudine summa paululum brevior, basin versus angustatus, basi subacuminatus, lateribus rotundatis, convexus, omnium subtilissime punctulatus, dorso plerumque linea subtili longitudinali impressus. Scutellum basi subtiliter bicarinatum. Elytra thorace paulo longiora, subtilissime punctulata. Abdomen confertim subtiliter punctulatum, ante apicem nigricans. Pedes testacei.

Variat forte sexu vertice thoracisque disco longitudinaliter leviter impressis.

- \* 16. *F. nigra*: *Picea, subtiliter punctulata, thorace subgloboso, obsolete canaliculato vel fovea tantum postica impresso.* — Long.  $\frac{3}{4}$  lin.

*Mannerh. Brachelytr.* 87. 3. — *Runde Brachel. Hal.* 31. 3. — *Er. Col. March. I.* 296. 4.

*Aleochar. nigra* Grav. *Micr.* 75. 12. *Mon.* 152. 11. — *Latr. Hist. nat. Crust. et Ins.* IX. 381. 12. — *Gyll. Ins. Suec. II.* 380. 3.

*Aleochar. fracticornis* Grav. *Mon.* 152. 9.

*Aleochar. picea* Grav. *Micr.* 75. 11. *Mon.* 152. 11. — *Latr. Hist. nat. Crust. et Ins.* IX. 381. 11.

*Falagria picea* *Mannerh. Brachelytr.* 87. 4. — *Runde Brachel. Hal.* 32. 4.

Habitat in Europa.

Praecedente duplo minor, nigro-picea vel picea, subtilissime pubescens. Antennae capite thoraceque breviores, apicem versus sensim incrassatae, corpori concolores. Caput thoracis latitudine, suborbiculatum, fronte leviter convexa, vix perspicue punctulatum. Thorax coleopteris paulo angustior, latitudine brevior, undique rotundatus, basin versus magis angustatus, modice convexus, confertim subtilissime punctatus, obsolete canaliculatus, basi supra scutellum transversim impressus. Scutellum aequale, confertissime punctatum. Elytra thorace parum longiora, subtiliter punctata, apice dilutiora. Abdomen supra parcius punctulatum. Pedes picei.

Obs. *Al. picea* Grav. differt nonnisi thoracis canalicula omnino extincta, relicta foveola supra scutellum.

- \* 17. *F. gratilla*: *Subtiliter punctulata, picea, thorace subgloboso, basi foveolato, rufo-picea, elytrorum limbo basali et apicali, antennarum basi pedibusque testaceis.* — Long.  $\frac{3}{4}$  lin.

Habitat in Mesopotamia, Dr. Helfer.

Statura omnino et summa affinitas praecedentis. Antennae capitis tho-

racisque longitudine, apicem versus sensim incrassatae, brunneae, articulis duobus primis testaceis. Caput orbiculatum, thoracis latitudine, subtilissime punctulatum, nigrum, nitidum, oris partibus testaceis, palpis piceis. Thorax coleopteris plus dimidio angustior, suborbiculatus, modice convexus, crebre subtiliter punctulatus, basi foveola minuta impressus, piceo-rufescens, nitidus. Elytra thorace paulo longiora, confertim subtiliter punctata, tenuiter pubescentia, testacea, macula magna disci fusca, seu fusca, limbo omni, basali licet apicalique latiore, testaceo. Abdomen crebre punctulatum, griseo-pubescent. Pedes pallide testacei.

- \* 18. *F. naevula*: *Rufo-testacea, thorace suborbiculato, postice foveolato, elytris disco infuscatis, abdomine apice nigro.* — Long.  $\frac{5}{6}$  lin.

Habitat in Aegypto, Dom. Prof. Ehrenberg.

Statura *F. nigrae*. Subtilissime punctulata, omnium tenuissime pubescens, laete testacea. Antennae capite thoraceque breviores, apicem versus sensim modice incrassatae, corpori concolores. Caput thoracis latitudine, suborbiculatum, basi truncatum. Thorax coleopteris dimidio angustior, latitudine paulo brevior, undique rotundatus, basin versus magis angustatus, leviter convexus, posterius obsolete canaliculatus, basi leviter impressus. Scutellum aequale. Elytra thorace paulo longiora, flavo-testacea, macula magna discoidali fusciscente. Abdomen supra omnium subtilissime minus confertim punctulatum, segmentis ultimis nigris. Pedes flavo-testacei.

Variat forte sexu fronte thoracisque disco leviter impressis.

- \* 19. *F. munda*: *Nitida, nigra, elytris subaeneis, thorace, pedibus, abdominis antennarumque basi rufis, thorace subquadrato fronteque longitudinaliter fortiter impressis.* — Long. 1 lin.

Habitat in Brasilia, Dom. Prof. Germar.

Statura fere *F. sulcatae*. Antennae (mutilatae) piceae, articulo primo rufo-testaceo. Palpi testacei, articulo tertio fusco. Caput thoracis latitudine, orbiculatum, confertim subtilissime punctulatum, fronte medio sat fortiter longitudinaliter impressa, vertice obsoletius foveolato, nigrum, pernitidum, glabrum. Thorax coleopteris dimidio angustior, latitudine paulo brevior, lateribus leviter rotundatus, basin versus perparum angustatus, basi truncatus, angulis posterioribus obtusis, anterioribus late rotundatis, leviter convexus, subtilissime confertim punctatus, longitudinaliter sat late profundeque impressus, impressione antice sensim evanescente, rufus, pernitidus, glaber. Scutellum confertissime punctatum, aequale, nigrum, griseo-pubescent. Elytra thorace paulo longiora, subtiliter confertissime punctata, nigro-subaenea, nitida, subtiliter albidopubescentia, infra scutellum impressa. Abdomen confertim punctulatum, longius pubescens, segmentis anterioribus tribus rufis, posterioribus nigris. Pectus nigrum. Pedes antiqui flavi, posteriores rufo-testacei.

- \* 20. *F. venustula*: *Picea, punctulata, thorace suborbiculato, rufescente, elytris fuscis, humeris, abdominis basi pedibusque testaceis.* — Long.  $\frac{5}{6}$  lin.

Habitat in Pennsylvania, Dom. Zimmermann.



Statura fere *F. nigrae*, at paululum maior, pube subtilissima vestita. Antennae capitis thoracisque longitudine, graciliores, pilosellae, testaceae, ante apicem fuscae. Palpi fuscii. Caput thoracis latitudine, orbiculatum, fronte leviter convexa, confertissime minus subtiliter punctulatum, nigrum, vix nitidum. Thorax coleopteris paulo angustior, latitudine parum brevior, lateribus anterieus rotundatus, basin versus leviter angustatus, modice convexus, confertissime subtiliter punctatus, fusco-rufescens, vix nitidus. Scutellum aequale, fuscum. Elytra thorace paulo longiora, confertissime subtiliter punctata, fusca, angulo humerali rufo-testaceo. Abdomen supra parcius punctulatum, segmentis duobus primis pallide flavo-testaceis, sequentibus nigris, nitidis. Pectus nigrum. Pedes flavi.

- \* 21. *F. albicincta*: *Nigro-picea, subnitida, confertissime subtilissimeque punctata, antennarum articulo primo testaceo, elytris basi dilutioribus, margine apicali, tibiis tarsisque albidis. — Long. 1 lin.*

Habitat ad Caracas Columbiae, Dom. Moritz.

Statura omnino praecedentis, nigro-picea, subnitida, confertissime subtilissimeque punctulata, pube subtilissima cinerea subsericans. Antennae capite thoraceque paulo longiores, nigrae, articulo primo crassiusculo, testaceo. Caput thoracis latitudine. Thorax coleopteris dimidio angustior, latitudine paulo brevior, basin versus leviter angustatus, parum convexus, dorso pone medium leviter late foveolatus. Coleoptera latitudine communi breviora, margine basali dilutiore, apicali albido. Pedes picei, tibiis tarsisque albidis.

- \* 22. *F. deliciata*: *Nigra, subnitida, antennarum articulis duobus primis, tibiis tarsisque pallide testaceis, elytris fuscis, basi dilutioribus, margine apicali albido. — Long. 1 lin.*

Habitat in Brasilia, Dom. Prof. Germar.

Praecedenti simillima, nigra, subnitida, confertissime subtilissimeque punctulata, pube subtilissima cinerea subsericans. Antennae capite thoraceque paulo longiores, graciles, pilosellae, nigrae, articulis duobus primis dilute flavo-testaceis, tertio piceo. Caput thoracis latitudine, suborbiculatum, fronte leviter convexa. Thorax coleopteris paulo angustior, latitudine dimidio fere brevior, undique rotundatus, basin versus subangustatus, satis aequaliterque convexus. Scutellum aequale. Coleoptera thorace paulo longiora, latitudine communi breviora, subtilissime punctulata, basi subtestacea, margine apicali albido. Abdomen supra confertim subtilissime punctulatum, nigrum, segmento primo obscure piceo. Pedes nigri, tibiis tarsisque pallide testaceis.

Obs. A praecedente praecipue thorace brevior, magis rotundato, aequaliter convexo, antennarumque articulo secundo etiam testaceo praecipue distincta.

- \* 23. *F. albo-guttata*: *Nigra, nitida, subtiliter punctulata, capite thoraceque rufo-piceis, elytris brunneis abdominisque basi albo-guttatis, femoribus posticis basi, geniculis tarsisque albidis. — Long. 1 lin.*

Habitat in Columbia, Dom. Moritz, in Brasilia, Dom. Prof. Germar.

*F. venustula* paulo gracilior. Antennae capite thoraceque multo lon-

giores, articulis sex primis testaceis, reliquis nigris. Palpi picei. Caput thoracis latitudine, laeve, dilute rufo-piceum. Thorax coleopteris dimidio angustior, basin versus sensim subangustatus, latitudine haud brevior, dorso subdepressus, subtiliter punctatus, pube brevi erecta fusca vestitus, rufo-piceus, marginibus dilutioribus. Scutellum nigrum. Elytra latitudine communi breviora, confertim subtiliter punctata, subtilissime cinereo-pubescentia, fusco-picea, nitida, macula transversali basali guttae in angulo apicis albis. Abdomen nigrum, nitidum, segmentis duobus primis subtus totis albis, supra nigris, singulo fascia lata, medio interrupta albida. Pectus nigrum. Pedes picei, postici trochanteribus femorumque basi, omnes geniculis tarsisque albidis, his longis, tenuibus.

### **Bolitochara Mannerh.**

*Mannerh. Brachelytr. p. 75. — Er. Col. March. I. p. 296.*

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula elongata, linearis, apice bifida; paraglossae parvae, angustae, acuminatae.*

*Palpi labiales triarticulati, articulo secundo brevior.*

*Tarsi antici 4-, posteriores 5-articulati, postici articulo primo elongato.*

Corpus minus depressum, alatum. Caput exsertum, porrectum, rotundatum, basi profunde constrictum, a thorace remotum eodemque paulo angustius, oculis mediocribus, modice prominulis. Labrum sat magnum, latiusculum, apice leviter rotundatum. Mandibulae sub labro occultae, muticae. Maxillae mala exteriore cornea, basi apiceque membranacea, portione membranacea apicali dense pubescente, interiore exteriore parum brevior, membranacea, margine exteriore corneo, apice introrsum oblique emarginata, sinu spinulis brevibus ciliato. Palpi maxillares modice elongati, articulo secundo tertioque longitudine subaequalibus, hoc parum incrassato, quarto minuto, subulato. Labium (t. III. f. 4.) mento latitudine paulo brevior, apicem versus angustato, apice leviter emarginato, ligula angusta, ad palporum labialium articuli secundi apicem usque porrecta, apice bifida, paraglossis parvis, acuminatis, interne ciliatis. Palpi labiales triarticulati, articulo primo secundoque cylindricis, hoc dimidio brevior, tertio tenuior, apicem versus leviter incrassato, longitudine primo aequali. Antennae capitis thoracisque fere longitudine, crassiusculae, apicem versus incrassatae, articulis tribus primis subelongatis, secundo tertioque obconicis. Thorax coleopteris angustior, latitudine brevior, antrorsum angustatus, angulis anterioribus rotundatis. Scutellum triangulare. Elytra apice truncata, ad angulum exteriorem distincte sinuata. Abdomen coleopterorum latitudine, supra planum, subtus convexum, saepe recurvum. Pedes modice elongati, intermedii basi leviter distantes, tarsi antici 4-, posteriores 5-articulati, elongati, articulis 4 primis sensim decrecentibus.

### **O b s e r v a t i o n e s.**

1. Maxillae omnino ut in *Homalota* (t. III. f. 22.). Paraglossae omnino ut in *Falagria*, ligula autem magis elongata.

2. Habitu quibusdam *Oxypodarum* Fam. II. speciebus (e. g. *O. tereti*) non dissimiles, discedunt autem capite basi constricto, remoto, et tarsis anticis 4-articulatis. Corpus rudius punctatum, parcius pubescens.

3. Genus *Bolitochara* a Com. Mannerheim quidem conditum, at a me re-  
restrictum, plurimis a celeberrimo auctore enumeratis speciebus *Homalotae*, aliis  
generibus aliis adscriptis.

4. Victus in fungis.

5. Mares abdominis segmento sexto dorsali aut tuberculato aut granulato in-  
signes.

\* 1. *B. lucida*: *Rufo-testacea, elytrorum disco posteriore abdominisque*  
*segmentis posterioribus basi fuscis, thorace suborbiculato, antrorsum*  
*angustato. — Long. 2 lin.*

*Aleochara lucida* Grav. *Micr.* 70. 4. *Mon.* 167. 47. — *Latr. Hist. nat. Crust. et Ins.*  
*IX.* 379. 4.

*Aleochara lunulata* var. *Gyll. Ins. Suec. II.* 386. 9.

Habitat in Germania, Suecia.

Rufo-testacea, nitida, pubescens. Antennae capitis thoracisque prope  
longitudine, concolores, basi apiceque paulo dilutiores. Caput thoracis basi  
paulo angustius, orbiculatum, fortius punctulatum. Thorax coleopterorum  
basi dimidio angustior, latitudine baseos paulo brevior, lateribus rotundatus,  
antrorsum angustatus, apice duplo quam basi angustior, modice convexus,  
fortius punctulatus, basi foveola transversali impressus. Elytra thorace ses-  
qui longiora, sat fortiter punctata, disco posteriore exteriori et spatio com-  
muni scutellari infuscatis. Abdomen supra disperse minus subtiliter puncta-  
tum, segmentis quinto sextoque basi infuscatis. Pectus fuscum. Pedes di-  
lutius testacei.

Mas elytris plicula longitudinali pone medium prope suturam, abdomi-  
nisque segmento sexto superiore carina longitudinali elevatis.

Obs. Sequenti simillima, differe tamen videtur capite testaceo, thorace antror-  
sum angustato, nec non statura sesqui maiore.

\* 2. *B. lunulata*: *Rufo-testacea, capite, antennarum medio, elytris, ex-*  
*ceptis humeris, abdomineque ante apicem fuscis, thorace suborbiculato.*  
*— Long. 1½—2 lin.*

*Mannerh. Brachelytr.* 75. 2. — *Er. Col. March. I.* 297. 1.

*Staph. lunulatus* Payk. *Mon. Staph.* 58. 41. *Faun. Suec. III.* 415. 61. — *Walk.*  
*Faun. Paris. I.* 272. 18.

*Aleochara lunulata* *Gyll. Ins. Suec. II.* 386. 9. — *Zetterst. Faun. Lappon. I.* 97. 5.  
*Ins. Lappon.* 76. 7. — *Sahlb. Ins. Fenn. I.* 348. 7.

*Aleochara cincta* Grav. *Mon.* 166. 46.

*Aleochara pulchra* Grav. *Mon.* 164. 44.

*Bolitochara pulchra* *Boisd. et Lacord. Faun. Ent. Paris. I.* 541. 2.

Habitat in Europa.

Rufo-testacea, subnitida, flavescens-pubescens. Antennae capitis thora-  
cisque prope longitudine, fuscae, basi rufo-testaceae, articulo ultimo dilute  
testaceo. Palpi testacei. Caput thoracis prope latitudine, nigrum, nitidum,  
profundius punctatum. Thorax coleopteris dimidio angustior, latitudine pa-  
rum brevior, lateribus rotundatus, antrorsum vix angustatus, leviter con-  
vexus, crebre punctatus, basi foveola transversa impressus. Elytra thorace  
sesqui longiora, crebre sat fortiter punctata, fusco-nigra, humeris margine-  
que summo apicali rufo-testaceis, vel rufo-testacea, disco posteriore magis



minusve fusco. Abdomen profunde vage punctatum, segmentis quinto toto et sexto basi nigris. Pedes pallidius testacei.

Mas elytris plicula longitudinali pone medium prope suturam abdominisque segmento dorsali sexto carina acuta longitudinali elevatis.

\* 3. *B. obliqua*: Fusca, elytrorum vitta obliqua pedibusque obscure ferrugineis. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 298. 2.

Bol. cincta Boisd. et Lacord. Faun. Ent. Paris. I. 542. 3.

Habitat in Germania.

Statura praecedentis, at paulo minor, fusca, parcius pubescens. Antennae capite thoraceque paulo breviores, crassiusculae, fuscae, apice concolores, basi obscure ferrugineae. Caput thorace parum angustius, nigrum, profundius punctulatum. Thorax coleopteris dimidio angustior, latitudine paulo brevior, lateribus rotundatus, basin versus subangustatus, leviter convexus, dense punctatus, basi foveola transversa profundiore impressus. Elytra thorace sesqui longiora, dense fortiter punctata, fusca, vitta obliqua, a humero ad angulum apicis internum percurrente, angusta, ad humerum dilatata, nonnunquam etiam margine laterali ferrugineis. Abdomen nigrum, segmentis anterioribus margine rufescentibus, supra parce obsoleteque punctulatum. Pedes obscure ferruginei.

Mas abdominis segmento dorsali sexto granulato.

\* 4. *B. varia*: Nigra, capite thoraceque rufo-testaceis, antennarum basi, pedibus coleopterisque testaceis, his maculis tribus nigris. — Long.  $1\frac{1}{2}$  lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura omnino praecedentium. Antennae capite thoraceque paulo breviores, apicem versus incrassatae, basi testaceae, medio rufo-testaceae, apice fuscae. Caput thoracis prope latitudine, parce subtilissime punctulatum, saturate rufo-testaceum, vel saepius rufo-piceum, nitidum. Thorax coleopteris plus dimidio angustior, latitudine dimidio brevior, lateribus ultra medium subrectis, ante medium rotundatis, et antrorsum angustatis, parum convexus, subtiliter obsoleteque punctulatus, supra scutellum late et parum profunde impressus, subtilissime pubescens, laete rufo-testaceus, nitidus. Elytra thorace sesqui fere longiora, crebre fortiterque punctata, densius pubescentia, flavo-testacea, macula magna transversa exteriore ante apicem, aliaque parva scutellari communi nigris. Abdomen supra parce subtiliterque punctatum, parcius pubescens, nitidum, vel totum nigrum, vel segmentis anterioribus tribus testaceis, medio magis minusve nigris piceisve, plerumque nigris, singulis margine omni testaceo. Pectus nigrum. Pedes flavi vel testacei, femoribus posticis fuscis.

Mas abdominis segmento dorsali sexto parcius granulato.

Variat colore obscuriore, capite piceo, thorace rufo, elytris piceo-testaceis, antennarum basi pedibusque rufo-testaceis.

Obs. *B. obliquae* proxima, differt autem colore, capite, thorace abdomineque parcius obsoleteque, elytris minus confertim punctatis.

Exempla numerosa misit cel. Prof. Gené.

**Ocalea Er.***Er. Col. March. I. p. 298.**Maxillae mala interiore mutica, apice spinulis ciliata.**Ligula elongata, linearis, apice bifida; paraglossae nullae.**Palpi labiales 3-articulati, articulis longitudine subaequalibus, ultimo tenuiore.**Tarsi omnes 5-articulati, postici articulo primo elongato.*

Corpus minus depressum, alatum. Caput exsertum, porrectum, rotundatum, basi constrictum, oculis mediocribus, parum prominulis. Labrum sat magnum, apice leviter rotundatum. Mandibulae sub labro reconditae, muticae. Maxillae mala exteriore cornea basi apiceque membranea, portione membranea apicali dense pubescente, interiore exteriore parum brevior, membranea, margine exteriore corneo, apice introrsum oblique emarginata, sinu spinulis brevibus ciliato. Palpi maxillares modice elongati, articulo secundo tertioque longitudine subaequalibus, tertio leviter incrassato, quarto minuto, subulato. Labium (t. III. f. 5.) mento transverso, apice leviter emarginato, ligula angusta, ad palporum labialium articuli secundi apicem usque porrecta, apice bifida, paraglossis omnino nullis. Palpi labiales triarticulati, articulo secundo paulo brevior, tertio secundo longitudine subaequali, at paulo tenuiore. Antennae minus crassae, articulis tribus primis longioribus, secundo tertioque subaequalibus. Thorax coleopteris paulo angustior, suborbiculatus. Scutellum triangulare. Elytra apice truncata, ad angulum exteriorem leviter sinuata. Abdomen coleopterorum latitudine, supra planum, subtus convexum. Pedes modice elongati, intermedii basi approximati, tarsis omnibus 5-articulatis, elongatis, articulis 4 primis sensim decrescenibus.

**Observationes.**

1. Maxillae omnino ut in *Homalota* (t. III. f. 22.). Ligula bifida valde elongata omnino ut in *Bolitochara*, paraglossae autem omnino nullae.

2. Characteribus omnino genus hocce cum *Oxypoda* convenit, ut vix nisi ligula elongata distinguatur; habitus autem magis differt, *Caloderæ* et *Tachysae* similis, ab hac tarsis anticis 5-articulatis, ab illa, ut a plerisque aliis huius tribus generibus, palporum labialium articulo tertio secundo haud longiore distinctum.

3. Sexus differentiam peculiarem non observavi.

\* 1. *O. castanea: Fusco-castanea, subtiliter punctulata, thorace sub-orbiculato, basi foveolato, pedibus pallidis.* — Long. 2 lin.

*Er. Col. March. I. 299. 1.*

Habitat in Germania. Sub musco humido in rivulorum ripis a def. Weber copiose lecta.

Statura fere *Myrmedoniae* collaris. Obscure brunnea, thorace elytris-que obscure castaneis, antennarum basi palpisque dilute rufo-piceis, pedibus pallide testaceis; capite, thorace elytris-que parce subtiliter-que vel fere obsolete punctatis, pube parca tenui fulva vestitis. Antennae capitis thoracis-que longitudine, apicem versus parum incrassatae. Caput thorace paulo angustius, basi fortiter constrictum, inter antennas leviter transversim impressum. Thorax latitudine parum brevior, coleopteris paulo angustior, suborbicularis, leviter convexus, postice fovea parum profunda impressus,

lateribus pilosellus. Coleoptera subquadrata, thorace paulo longiora. Abdomen fere nigrum, segmentorum marginibus, praecipue infra, rufo-brunneis, supra nitidum, subtiliter vage punctatum, pilis parcis depressis fulvis vestitum.

- \* 2. *O. badia*: *Rufo-picea, fortius punctulata, thorace orbiculato, postice foveolato.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 300. 2.*

Habitat Berolini, ipse, in Austria, Dom. Schüppel.

Praecedenti affinis, at minor, gracilior, rufo-brunnea, nitidula, pube parca flavescenti vestita. Antennae capitis thoracisque longitudine, apicem versus leviter incrassatae, rufo-testaceae, ante apicem obscuriores. Palpi rufo-testacei. Caput thorace paulo angustius, basi profunde constrictum, minus subtiliter at parum profunde vage punctatum. Thorax coleopteris paulo angustior, latitudine haud brevior, suborbiculatus, convexiusculus, densius fortiusque punctatus, basi foveola levi impressus. Coleoptera subquadrata, dense profundeque punctata. Abdomen brunneum, apice dilutius, supra vage punctatum, pilis parcis depressis flavescens alisque erectis nigris vestitum.

- \* 3. *O. spadicea*: *Rufo-picea, fortius punctata, thorace transverso, antice truncato, aequali.* — Long.  $1\frac{2}{3}$  lin.

*Er. Col. March. I. 300. 3.*

Habitat Berolini.

Rufo-picea, pilis depressis flavescens parcius vestita. Antennae vix capitis thoracisque longitudine, apicem versus parum incrassatae, fusco-piceae. Caput thorace paulo angustius, basi perparum constrictum, subtilissime vage punctatum. Thorax latitudine dimidio brevior, coleopterorum fere latitudine, basi lateribusque leviter rotundatus, antice truncatus, angulis deflexis, supra parum convexus, confertim punctatus, fovea basali nulla. Coleoptera thorace haud longiora, confertim punctata, punctis obliquis. Abdomen supra vage subtiliter punctatum, nigro-piceum, apice brunneum.

- \* 4. *O. procera*: *Picea, fortius punctata, thorace elytrisque castaneis, thorace transverso, antice truncato, basi impresso.* — Long.  $2\frac{1}{2}$  lin.

Habitat in Germania, Dom. Schüppel.

*O. castanea* dimidio maior et magis oblonga, statura elongata *O. badiae*, at duplo maior. Antennae capitis thoracisque prope longitudine, apicem versus sensim subincrassatae, articulo tertio secundo paulo longiore, 4—10 sensim paulo crassioribus, subobconicis, ultimo praecedente sesqui longiore, ovato, apice acuminato, rufae, griseo-pubescentes. Caput thorace paulo angustius, rotundatum, basi haud constrictum, convexum, parce subtiliter punctatum, parce pubescens, nigro-piceum, nitidum, ore rufo. Thorax coleopteris paululum angustior, latitudine dimidio brevior, lateribus modice, basi leviter rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, leviter transversim convexus, densius minus subtiliter punctatus, basi leviter transversim impressus, rufo-piceus seu rufo-castaneus, niti-



dus, pube griseo-fulvescente densiore vestitus. Elytra thoracis longitudine, dense fortius punctata, castaneo-rufa, subnitida, pube longiore depressa fulvescente densius vestita. Abdomen elongatum, segmentis singulis basi crebre, apice parce punctatis, nigro-piceum, nitidum, apice segmentorumque singulorum margine posteriore supra saturatius, subtus dilutius rufo-piceis, parcius fulvescenti-pubescens. Pectus nigro-piceum. Pedes sat elongati, rufi.

Obs. Praecedenti proxima, statura maiore, capite magis rotundato, thorace paulo brevior, basi impresso, angulis anterioribus magis rotundatis, abdominis segmentis basi crebrius punctatis praecipue distinguenda.

\* 5. *O. murina*: *Nigra, cinereo-pubescens, antennis pedibusque piceis, geniculis tarsisque flavis, thorace subquadrato, apice truncato, basi foveolato. — Long. 1½ lin.*

Habitat in Austria et in Istria, Dom. Schüppel, in Sardinia, Dom. Prof. Gené.

Statura fere Homalotae graminicolae, praecedentibus paulo magis depressa, *O. spadiceae* capite basi minus constricto et thorace apice truncato affinis, nigra, subnitida, pube albedo-cinerea densius vestita. Antennae capitulis thoracisque longitudine, graciles, apicem versus perparum incrassatae, piceae. Palpi flavescentes, articulo tertio piceo. Caput thorace paulo angustius, basi leviter constrictum, obsoletius punctulatum. Thorax coleopteris paulo angustior, latitudine paulo brevior, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus deflexis rotundatis, leviter convexus, dense fortius punctatus, basi fovea parum profunda supra scutellum notatus. Elytra thorace paulo longiora, densius fortiusque punctata. Abdomen dorso crebrius punctatum. Pedes picei, geniculis tarsisque flavo-testaceis.

Mas antennis longioribus, articulo ultimo elongato, ovato, femina antennis brevioribus, (capite sesqui longioribus), rufis.

Obs. Habitus fere Aleocharae genuinae, at secundum oris structuram non nisi huic generi adscribenda, quorum reliqui characteres omnino conveniunt.

6. *O. decumana*: *Nigra, cinereo-pubescens, antennis piceis, tibiis tarsisque testaceis, thorace transversim subquadrato, apice truncato, basi foveolato. — Long. 2 lin.*

Habitat in Gallia? Mus. Dom. Chrevrolat.

Praecedenti affinis, at paulo maior, paulo magis depressa et subtilius punctata, nigra, subnitida, pube albida cinerascens. Antennae capitulis thoracisque longitudine, graciles, piceae, basi apiceque rufo-brunneae. Palpi flavescentes, apice piceae. Mandibulae dilute rufae, basi nigrae. Caput thorace angustius, basi perparum constrictum, subtiliter punctulatum. Thorax coleopteris dimidio prope angustior, latitudine dimidio brevior, lateribus ultra medium subrectis, ante medium rotundatis, apicemque versus angustatis, angulis anterioribus rotundatis, subdeflexis, apice truncatus, basi rotundatus, parum convexus, parce subtiliterque punctatus, supra scutellum late et parum profunde impressus. Elytra thorace paulo longiora, parcius subtiliter punctata. Abdomen supra parce punctulatum. Pedes nigri, antici coxis apice femoribusque piceo-testaceis, tibiis tarsisque omnibus testaceis.

Species mihi invisae, forte huius generis.

7. *O. (?) proluxa*: *Rufo-picea, subnitida, pubescens, antennis, elytris anoque dilutioribus, pedibus pallide testaceis.*

*Aleoch. proluxa* Gyll. *Ins. Suec. IV.* 484. 9—10.

*Bolitochara proluxa* Mannerh. *Brachelytr.* 75. 3.

Habitat in Scania, ad Abusa, rarissime.

Statura, magnitudo et affinitas *Bol. lunulatae*, sed color alius, antennae paulo longiores etc. Caput latitudine thoracis, rotundatum, nigrum, nitidum, pubescens, subtilissime punctulatum, ore rufo-piceo. Antennae longitudine capitis cum thorace, crassiusculae, basi apiceque rufo-ferrugineae, articulis mediis 3—6 obscurioribus infuscatis. Thorax latitudine antica vix longior, apice truncatus, basi rotundatus, supra convexus, nigro-fuscus, nitidus, tenuiter pubescens et pilosus, obsolete punctulatus, versus margines dilutior, ante medium baseos foveola parva obsoleta impressus. Scutellum parvum, triangulare, fuscum. Elytra thorace paulo latiora et longiora, apice truncata, supra modice convexa, fusco-testacea, iuxta scutellum obscuriora, undique confertim evidentius punctata, longius pubescentia. Abdomen pone elytra longitudine fere reliqui corporis, crassum, supra nigro-piceum, nitidum, vix punctulatum, ano testaceo. Corpus subtus dilutius rufo-testaceum, nitidum, obsolete punctulatum, segmentis ventralibus basi obscurioribus, nigro-piceis. Pedes elongati, tenues, pallide rufo-testacei. (Gyll.)

8. *O. (?) rufilabris*: *Fusco-picea, subnitida, pubescens, ore, antennis, elytris, marginibus segmentorum abdominis anoque rufescentibus, pedibus pallidis, thorace canaliculato.* — Long.  $1\frac{1}{2}$  lin.

*Aleoch. rufilabris* Sahlb. *Ins. Fenn. I.* 349. 8.

Habitat in Fennia australi, rarissime.

*O. (?) proluxae* similis, sed differt magnitudine minore, thorace evidentius canaliculato, antennis extrorsum minus incrassatis, elytrisque subtilius punctulatis. Caput thorace paulo angustius, rotundatum, fusco-nigrum, nitidum, subpubescens, subtilissime punctulatum, ore palpisque rufescentibus. Antennae longitudine thoracis, crassiusculae, ferrugineae, basi pallidiores. Thorax longitudine paulo latior, antice posticeque truncatus, lateribus vix dilatatus, angulis rotundatis, supra convexus, piceo-brunneus, subtiliter punctulatus, parce pubescens, canalicula sat evidenter impressa. Scutellum triangulare, piceo-ferrugineum, subtilissime punctulatum. Elytra thorace paulo longiora et latiora, apice truncata, iuxta angulum externum subsinuato-emarginata, supra subdepressa, subtiliter punctulata, subpubescentia, brunneo-rufescentia. Abdomen pone elytra longitudine corporis reliqui, nigro-piceum, obsolete subtilissime punctulatum, segmentorum marginibus et ano rufescentibus. Corpus subtus fusco-piceum, subnitidum, subtiliter punctulatum, pubescens, ano et marginibus segmentorum rufescentibus. Pedes elongati, toti pallide testacei. (Sahlb.)

**Calodera Mannerh.**

*Mannerh. Brachelytr. p. 85. — Er. Col. March. I. p. 301.*

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula brevis, bifida; paraglossae breves.*

*Palpi labiales 3-articulati, articulo secundo brevior, tertio tenuiore.*

*Tarsi omnes 5-articulati, postici articulo primo elongato.*

Corpus elongatum, minus depressum, alatum. Caput exsertum, porrectum, thoracis fere latitudine, rotundatum, basi magis minusve constrictum, oculis medioeribus, parum prominulis. Labrum maiusculum, lateribus rotundatum, apice truncatum. Mandibulae muticae. Maxillae mala exteriore cornea, basi apiceque membranacea, portione membranacea apicali dense pubescente, interiore exteriore paulo brevior, membranacea, margine exteriore corneo, apice introrsum oblique emarginata, sinu spinulis brevibus ciliato. Palpi maxillares modice elongati, articulo secundo tertioque longitudine subaequalibus, hoc modice incrassato, quarto minuto, subulato. Labium (t. III. f. 6.) mento perbrevis, antrorsum angustato, apice leviter emarginato, ligula breviuscula, non ultra articuli primi apicem producta, bifida, paraglossis ligula brevioribus, latis, rotundatis, margine haud ciliatis. Palpi labiales triarticulati, articulo secundo duplo fere brevior, tertio tenuiore, primo longitudine subaequali. Antennae crassiusculae vel crassae, articulis tribus primis magis elongatis, secundo tertioque obconicis, 4—10 plerumque transversis, ultimo oblongo-ovato. Thorax coleopteris angustior, subquadratus aut subovatus. Scutellum triangulare. Elytra apice truncata, ad angulum exteriorem profundius sinuata. Abdomen coleopterorum latitudine, lineare, supra planum, subtus convexum. Pedes modice elongati, intermedii basi subapproximati, tarsis omnibus 5-articulatis, his elongatis, articulis 4 primis longitudine de-crescentibus.

**Observationes.**

1. Maxillae omnino ut in *Homalota* (t. III. f. 22.). Paraglossae latae, rotundatae ut in *Aleochara*, at ligula non adeo brevis.

2. Antennae in plerisque huius generis speciebus sat longae et fortes, ceterum habitu *Tachyus* similis, a quibus praeter paraglossas praecipue tarsis anticis 5-articulatis differunt. Corpus plerisque confertissime subtilissimeque punctatum, pube brevissima subtilissimaque pruinae instar vestitum et sericans.

3. Tarsi postici in hoc genere satis elongati, articulis inaequalibus, primo fortiter elongato, nescio igitur quo errore Com. Mannerheim ei tarsos »articulis aequalibus« tribuit. Ceterum sine dubio illius *Calodera* cum nostra congruit. Addidi *C. nigricollis*, quippe quae licet habitu quodammodo discrepat, tarsis anticis 5-articulatis, et oris etiam structura omni modo convenit.

4. Victus plerisque speciebus in udis.

5. Sexus differentiam peculiarem haud observavi.

\* 1. *C. nigricollis*: *Rufopicea, opaca, creberrime punctata, tenuiter pubescens, thorace subquadrato, fusco. — Long. 1½—3 lin.*

*Er. Col. March. I. 302. 1.*

*Staph. nigricollis Payk. Faun. Suec. III. 400. 42.*

*Aleochar. nigricollis Grav. Micr. 84. 24. Mon. 163. 36. — Latr. Hist. nat. Crust. et Ins. IX. 385. 24. — Gyll. Ins. Suec. II. 384. 7.*

*Bolitoch. nigricollis Boisd. et. Lacord. Faun. Ent. Paris. I. 545. 5.*



Habitat in Europae silvis.

Statura fere Myrmedoniae canaliculatae, at plerumque maior. Antennae rufae, validae, capite thoraceque longiores, articulo secundo tertio paulo brevior. Caput, thorax et elytra punctatissima, pube densiore flavescens vestita. Caput obscure rufum, postice parum constrictum. Thorax coleopteris paulo angustior, subquadratus, parum convexus, aequalis, nigricans. Elytra rufa. Abdomen parcius punctatum, supra subnitidum, piceum, apice rufo. Pedes dilute rufo-testacei.

Obs. Punctura fortiore, profunda et pube haud subtilissima primo intuitu in hoc genere aliena videtur.

\* 2. *C. forticornis*: *Rufa castanea, subtiliter punctulata, thorace suborbiculato, aequali, elytris fortius punctatis.* — Long. 2 lin.

*Bolitoch. forticornis* Boisd. et Lacord. *Faun. Ent. Paris. I.* 543. 4.

Habitat in Germania, Austria, Dom. Schüppel, Lutetiae, Mus. Dr. Aubé.

Rufa-castanea, nitida, pube minus subtili grisea parcius vestita. Antennae capite thoraceque breviores, crassae, articulo secundo tertioque subaequalibus, ultimo breviter ovato, obtuso. Caput subtilissime parce punctatum, basi profunde constrictum. Thorax suborbiculatus, basi, lateribus et angulis anterioribus rotundatis, angulis posterioribus obtusis, basi coleopterorum prope latitudine, antrorsum leviter angustatus, apice truncatus, latitudine baseos paulo brevior, leviter convexus, subtiliter parcius punctatus. Coleoptera thoracis longitudine, subquadrata, profundius punctata, punctis obliquis. Abdomen elongatum, supra laeve, segmentis basi summa transversim impressis ibique punctatis. Pedes rufo-testacei.

\* 3. *C. nigrita*: *Atra, subtilissime punctulata, subtilissime dense cinereo-pubescent, thorace subdepresso, obsolete canaliculato, ore tarsisque flavis.* — Long. 2 lin.

*Mannerh. Brachelytr.* 86. 1. — *Er. Col. March. I.* 303. 2.

Habitat in Germania, locis uliginosis.

Atra, vix nitida, pube subtili, densa cinereo-sericea. Antennae capite thoraceque longiores, fuscae, articulis primis sub-testaceis, articulo secundo tertio sesqui longiore, ultimo ovato. Caput, thorax et elytra omnium subtilissime confertissimeque punctata. Caput basi fortiter constrictum, fronte obsolete impressa, ore flavescens. Thorax subquadratus, vel latitudine paulo longior, basin versus haud angustatus, lateribus ante medium leviter rotundatus, apice leviter angustatus, subdepressus, basi fovea transversa, dorso canalicula obsolete impressus. Coleoptera thorace paulo longiora et sesqui latiora. Abdomen supra confertim subtiliter punctulatum, planum et aequale. Pedes nigri, geniculis, tibiis apice tarsisque flavescentibus.

\* 4. *C. humilis*: *Depressa, atra, subtilissime punctulata, subtiliter dense cinereo-pubescent, thorace subquadrato, obsolete canaliculato, abdomine supra confertissime punctato, ore, antennarum basi tarsisque flavis.* — Long. 1½ lin.

*Er. Col. March. I.* 303. 3.

Habitat in Germania boreali, ipse, in Austria, Dom. Schüppel.

Praecedente minor et magis depressa, atra, pube tenui subtili dense cinereo-sericea, tota subtilissime confertissimeque punctulata. Antennae capitis thoracisque longitudine, fuscae, basi dilutiores, articulo secundo tertio duplo maiore, ultimo ovato. Caput basi profunde constrictum, infra antennas flavum. Thorax subquadratus, latitudine haud longior, basin versus subangustatus, lateribus ante medium rotundatis, depressus, obsoletissime canaliculatus, basi foveola transversa impressus. Coleoptera thorace paulo latiora et vix longiora, depressa. Abdomen reliquo corpore non distinctius punctatum, dorso planum et aequale. Pedes ut in praecedente.

- \* 5. *C. longitarsis*: *Atra, subtilissime punctulata, dense subtilissime cinereo-pubescent, thorace subdepresso, basi foveolato, abdomine supra confertissime punctato, palpis, antennarum basi tarsisque testaceis. — Long. 2 lin.*

*Er. Col. March. I. 698. 3. a.*

Habitat in Germania, Anglia, Gallia, Italia.

Habitu omnino proxime praecedentis, at paulo maior, *C. nigrityae* aequalis, atra, tota subtilissime confertissimeque punctata, pube subtili cinereo-sericea dense vestita. Antennae capite thoraceque parum longiores, apicem versus sensim incrassatae, fuscae, basi dilutiores, articulo primo testaceo, secundo tertioque aequalibus, ultimo maris elongato, antecedentibus duobus coniunctis aequali, feminae ovato, praecedenti sesqui longiore. Palpi fusco-testacei. Labrum piceum. Caput thorace paulo angustius, basi profundius constrictum. Thorax subquadratus vel latitudine sublongior, basin versus haud angustatus, lateribus ante medium leviter rotundatis, apice angustatus, angulis anterioribus rotundatis, deflexis, parum convexus, basi foveola transversa impressus, certo situ obsolete canaliculatus. Coleoptera thorace sesqui latiora et paulo longiora. Abdomen aequè subtiliter ac reliquum corpus punctatum, apice attenuatum, dorso segmentis 4 primis summa basi transversim depressis. Pedes nigri, femoribus summo apice, tibiis tarsisque totis testaceis.

Variat minus matura colore corporis fusco.

Obs. 1. Praecedentibus primo intuitu simillima, at antennarum articulis secundo tertioque aequalibus, sequentibus sensim crassioribus, haud transversis facile distinguenda. Hac structura *Tachysa* simulat et cum *T. umbratica* facile confundi possit, at tarsi antici distincte 5-articulati.

2. A. cel. Dom. Spence ex Anglia missa nomine *Aleocharae longitarsis* Kirby. Duo huius speciei exempla Gallica in collectione Dom. Aubé vidi, alterum *Bolitocharae cinereae* Dej., alterum *B. perplexae* Dej. nomine signatum. Specimina in Sardinia lecta ad examinandum communicavit Dom. Prof. Genève.

- \* 6. *C. rubicunda*: *Rufa, subtilissime punctulata, subtiliter cinereo-pubescent, capite abdominisque segmentis duobus penultimis fuscis, thorace subovato, basi leviter foveolato, abdomine supra subtilissime punctato. — Long. 2 lin.*

*Tachysa rubicunda Er. Col. March. I. 309. 4., 639. 4.*

Habitat in Germania: Berolini Dom. Schüppel et Weber, in Austria, Dom. Schüppel.

Statura, habitus et summa similitudo praecedentis. Corpus iam rufo-

testaceum iam saturatius rufum vel rufo-piceum, parum nitidum, pube subtili sericante cinerea vestitum. Antennae eadem structura ut in praecedente, at paulo validiores, capite thoraceque longiores, apicem versus sensim modice incrassatae, rufae, basi testaceae, articulo secundo tertioque aequalibus, sequentibus sensim crassioribus, crassitie haud brevioribus, ultimo magno ovato. Palpi dilute testacei. Caput thorace paulo angustius, breviter ovatum, basi sat fortiter constrictum, subtilissime confertim punctatum, piceum, nitidum. Thorax coleopteris paululum angustior, latitudine summa vix longior, lateribus ante medium cum angulis anterioribus rotundatus, basin versus subangustatus, satis convexus, confertim subtilissime punctatus, basi foveola supra scutellum impressus. Elytra thorace vix longiora, subdepressa, confertim subtiliter punctata. Abdomen aequè subtiliter ac reliquum corpus punctatum, segmentis 4 anterioribus summa basi transversim depressis, penultimis duobus nigricantibus.

Obs. Antennarum structura praecedenti associanda, a qua praeter colorem thorace convexiore, basin versus paululum angustato coleopterisque thorace vix longioribus et parum latioribus at distinctius punctatis specie distinguitur. Ipsum me olim habitus fefellit et antennarum structura aliena, ut *Tachyusa* e hanc speciem adscripserim: iam alteris duobus speciminibus examinatis tarsos anticos distincte 5-articulatos cognovi.

\* 7. *C. rubens*: *Rufo-picea, subtilissime punctulata, subtiliter cinereo-pubescent, thorace quadrato, subdepresso, basi obsolete foveolato, abdomine supra subtilissime confertissimeque punctato. — Long. 2 lin.*

*Er. Col. March. I. 304. 5.*

Habitat in Germania. Unicum specimen Berolini legi.

Statura omnino praecedentis, obscure rufa, opaca, aequaliter confertissime subtilissimeque punctata, pube densa, subtili, cinereo-sericea obducta. Antennae capite thoraceque paulo longiores, validae, rufae, articulis duobus primis rufo-testaceis, tertio secundo paulo brevior, sequentibus omnibus transversis, ultimo oblongo-ovato. Caput thorace paulo angustius, ovatum, basi parum constrictum, fusco-piceum, ore testaceo. Thorax subquadratus, lateribus ultra medium rectis, ante medium leviter rotundatis, apice angustatus, angulis anterioribus rotundatis, deflexis, latitudine haud longior, leviter convexus, dorso late at obsoletissime canaliculatus, basi parum profunde impressus. Coleoptera thorace paululum latiora et longiora. Abdomen aequè subtiliter ac reliquum corpus punctatum, dorso plano et aequali, ante apicem nigricans. Pedes testacei.

Obs. Affinis *C. nigritae*, at paulo brevior, antennae crassiores, articulo tertio maiore, thorax latior. A sequentibus abdomine supra plano discedit.

\* 8. *C. uliginosa*: *Atra, subtilissime punctulata, parcius griseo-pubescent, thorace subovato, convexiusculo, basi foveolato, abdomine supra parce punctato, nitidulo, ore flavo, antennis tarsisque piceis. — Long. 1 $\frac{3}{4}$  lin.*

*Er. Col. March. I. 305. 6.*

*Aleoch. aethiops Gyll. Ins. Suec. II. 383. 6.*

Habitat in Germania.

Atra, nitidula, pube griseo-sericea tenuiter vestita. Antennae piceae,



capite thorace paulo longiores, articulo tertio secundo aequali, sequentibus vix transversis, ultimo oblongo-ovato. Caput, thorax et elytra subtilissime confertim punctata. Caput thorace minus, rotundatum, basi fortiter constrictum, ore obscure flavo. Thorax latitudine paulo longior, coleopteris angustior, ante medium leviter rotundatus, basin versus subangustatus, angulis anterioribus rotundatis, deflexis, convexus, basi foveola minuta transversa impressus. Abdomen supra inaequale, parcius subtiliter punctatum, dorso segmentis 4 primis singulis convexis, basi fortiter transversim depressis. Pedes nigri, tarsis piceis.

- \* 9. *C. riparia*: *Picea*, subtilissime punctulata, griseo-pubescent, thorace suborbiculato, convexiusculo, basi foveola gemina impresso, abdomine supra parce punctulato, antennis pedibusque rufis. — Long.  $1\frac{1}{4}$  lin.

*Er. Col. March. I.* 305. 7.

Habitat in Germania.

Praecedenti affinis, at plus duplo minor, vel nigro-picea, vel thorace elytris dilutioribus, vel interdum tota rufo-testacea, nitidula, confertissime subtiliter punctata, tenuiter griseo-pubescent. Antennae vel rufo-piceae, vel testaceae, capitis thoracisque longitudine, validae, articulo tertio secundo dimidio brevior, sequentibus transversis, ultimo ovato. Caput thorace dimidio minus, rotundatum, basi profunde constrictum, ore rufo. Thorax latitudine summa haud longior, coleopteris dimidio angustior, lateribus ante medium et angulis anterioribus rotundatus, basin versus vix angustatus, convexiusculus, basi foveolis duabus rotundatis approximatis vel contiguis impressus. Elytra thorace parum longiora. Abdomen supra parce punctatum, segmentis 4 primis singulis leviter convexis, basi depressis. Pedes rufo-picei.

- \* 10. *C. aethiops*: *Picea*, subtiliter punctulata, griseo-pubescent, thorace suborbiculato, convexiusculo, basi foveola transversa impresso, abdomine supra confertissime punctato, antennis pedibusque piceis. — Long. 1 lin.

*Er. Col. March. I.* 306. 6.

*Aleochar. aethiops* Grav. *Micr.* 77. 15. *Mon.* 153. 14. — *Latr. Hist. nat. Crust. et Ins.* IX. 382. 15.

*Calodera testacea* Mannerh. *Brachelytr.* 86. 3.

*Aleochar. testacea* Sahlb. *Ins. Fenn. I.* 347. 5.

Habitat in Germania.

Praecedenti simillima, at minor, plerumque nigro-picea, rarius rufo-picea, vel testacea, nitidula, confertissime subtiliter punctata, pube densa griseo-sericea vestita. Antennae capitis thoracisque longitudine, validae, piceae, articulo secundo tertio duplo fere longiore, sequentibus transversis, ultimo ovato. Caput thorace dimidio minus, suborbiculatum, basi profunde constrictum, ore flavescente. Thorax coleopteris dimidio angustior, lateribus ante medium et angulis anterioribus rotundatus, basin versus subangustatus, latitudine haud brevior, convexiusculus, basi foveola transversa impressus. Elytra thorace distincte longiora. Abdomen supra confertissime subtiliter punctatum, opacum, pube aequae densa ac reliquum corpus vestitum, apice

rufo-piceo, dorso segmentis 4 primis singulis leviter convexis, basi profunde transversim impressis. Pedes picei.

Obs. A praecedente antennarum articulo tertio brevior et abdomine confer-tim punctato opacoque distinguenda.

\* 11. *C. umbrosa*: *Subdepressa, subtiliter punctata, cinereo-pubescent, nitidula, nigra, elytris fuscis, antennarum basi pedibusque flavis, thorace subquadrato, aequali.* — Long.  $1\frac{1}{4}$  lin.

*Er. Col. March. I.* 304. 4.

Habitat in Germania: Berolini a Dom. Schüppel lecta.

Statura elongata, depressiuscula Homalotae, Hom. elongatulae non dissimilis, et minimis eius individuis aequalis, a plerisque huius generis punctura minus conferta discedens, C. aethiope perparum maior et paulo magis depressa, pube longiore minus subtili grisea vestita. Antennae capite thoraceque paululum breviores, crassiusculae, nigrae, articulis tribus primis testaceis, tertio secundo paulo brevior, sequentibus transversis, ultimo oblongo-ovato, apice obtuse rotundato. Palpi testacei. Caput thorace parum angustius, suborbiculatum, basi profunde constrictum, parce subtiliterque obsolete punctatum, nigrum, nitidum. Thorax coleopteris dimidio fere angustior, lateribus perparum rotundatus, neque basin neque apicem versus angustatus, latitudine paululum brevior, subdepressus, aequalis, minus subtiliter punctatus, piceus, nitidus. Elytra thorace paulo longiora, crebre minus subtiliter punctata, piceo-testacea, nitida. Abdomen nigrum, nitidulum, parcius subtiliter punctulatum, segmentis tribus primis piceis, singulis dorso levissime convexis, basi distincte transversim impressis. Pedes testacei.

Species mihi invisa:

12. *C. protensa*: *Linearis, nigra, griseo-pubescent, thorace elongato, canaliculato.*

*Mannerh. Brachelytr.* 86. 2.

Habitat in Suecia.

C. uliginosa (Al. aethiope Gyll.) duplo minor, magis linearis, minus pubescens, nec non thorace longiore canaliculato diversa.

### **Tachyusa Er.**

*Er. Col. March. I.* p. 307.

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula brevis, bifida; paraglossae vix ullae.*

*Palpi labiales 3-articulati, articulo secundo brevior.*

*Tarsi antici 4-, posteriores 5-articulati, articulo primo elongato.*

Corpus elongatum, minus depressum, alatum. Caput exsertum, porrectum, thoracis latitudine, rotundatum, basi constrictum, oculis mediocribus, parum prominulis. Labrum transversum, apice truncatum, mandibulas haud obtegens. Mandibulae muticae. Maxillae mala exteriori cornea, basi apiceque membranacea, portione membranacea apicali pubescente, interiori exteriori paulo brevior, membranacea, margine exteriori corneo, apice introrsum obli-

que emarginata, sinu spinulis brevibus ciliato. Palpi maxillares modice elongati, articulo secundo terdoque longitudine aequalibus, hoc leviter incrassato, quarto minuto, subulato. Labium (t. III. f. 7.) breve, mento transverso, apice leviter emarginato, ligula brevi, palporum labialium articulum primum haud excedente, bifida, paraglossis obsoletis. Palpi labiales triarticulati, articulo secundo primo minore, tertio hoc tenuiore at paulo longiore. Antennae longiores, graciles, articulis tribus primis magis elongatis, inter se longitudine aequalibus, primo crassiore. Thorax coleopteris paulo angustior. Scutellum triangulare. Elytra apice introrsum oblique truncata, ad angulum anteriorem sinuata. Abdomen supra planiusculum, subtus convexum, vivis saepe elevatum. Pedes elongati, graciles, intermedii basi approximati, tarsis anticis 4-, posterioribus 5-articulatis, his elongatis, articulis 4 primis longitudine decrescentibus.

### Observationes.

1. Maxillae omnino ut in *Homalota* (t. III. f. 22.). Ligula fere ut in *Calodera*, at paraglossae valde obsoletae, nonnisi angulo prominulo indicatae.

2. Habitu gracili *Tachysae* *Caloderis* et quibusdam *Homalotis* similes, ab illis antennis gracilioribus et tarsis anticis 4-articulatis, ab his pedibus magis elongatis et tarsis posticis longis, articulo primo elongato distinguuntur. A *Palagris*, quibuscum habitu et tarsorum structura conveniunt, paraglossis obsoletis, coxis intermediis magis approximatis et capite minus remoto differunt.

3. Abdomen in hoc genere saepius insigne segmentis inaequalibus, anterioribus longioribus: his plerumque abdomen basi magis minusve coarctatum. Aliis abdominis segmenta magnitudine aequalia. Forte in posticum, ubi specierum numerus satis increvit, hac nota ad sectiones in hoc genere instituendas utendum erit, quibus, speciebus iam cognitis parum numerosis, nondum opus esse videtur. Nihilominus species abdominis segmentis inaequalibus et aequalibus haud commiscui.

4. Nomen *Ischnopoda* Stephens (Illustr. V. p. 430.) tribuit *Aleocharis* pedibus elongatis: species ab eo enumeratae partim ad hoc, partim ad praecedens genus pertinere videntur. Quodsi neque satis distinctum, neque nulla descriptione illustratum invenio istud genus, nomen citatum omnino negligendum esse non dubitavi.

5. Sexus differentias, si quas observavi, apud singulas species notavi.

6. Victus plerisque in ripis. Nonnullae (*T. constricta* et *coarctata*) in arena humida riparia summa agilitate circumvagantur, et territae abdomine elevato velocissime fugiunt.

\* 1. *T. constricta*: *Picea*, subtiliter pubescens, antennis pedibusque rufis, thorace subovato, abdomine basi fortiter angustato. — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I.* 307. 1.

Habitat in Germania, Gallia.

Corpus eximie gracile, nigro-piceum, nitidulum, subtilissime punctatum, tenuissime griseo-pubescens. Antennae capitis thoracisque longitudine, testaceae, apicem versus leviter incrassatae, piceae. Caput rotundatum, basi profunde constrictum. Thorax longitudine vix angustior, lateribus ante medium cum angulis anterioribus rotundatus, basin versus leviter angustatus, convexiusculus, medio subdepressus. Coleoptera thorace paulo longiora et latiora, lateribus subrotundata, leviter convexa, apice simul emarginata. Abdomen elongatum, basi coleopteris duplo angustius, apicem versus sensim latius, apice coleopteris latitudine subaequale, segmentis inaequalibus, ante-



rioribus transversis, supra parce subtilissime punctulatum. Pectus rufo-piceum. Pedes rufo-testacei.

- \* 2. *T. coarctata*: Nigra, subtiliter pubescens, antennis pedibusque piceis, thorace subovato, abdomine basi angustato. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 308. 2.

Habitat in Germania, Gallia, Italia.

Praecedenti affinis at non adeo gracilis. Nigra, nitidula, subtilissime confertim punctata, pube tenui densius vestita. Antennae capite thoraceque vix breviores, apicem versus parum incrassatae, piceae. Caput rotundatum, basi profunde constrictum, fronte maris medio subimpressa. Thorax coleopteris paulo angustior, latitudine paululum brevior, lateribus ante medium leviter rotundatus, basin versus haud angustatus, angulis anterioribus deflexis rotundatis, leviter convexus, supra scutellum obsolete impressus. Elytra thorace longiora, parum convexa, lateribus subrotundata, apice simul emarginata. Abdomen basi coleopteris dimidio angustius, apicem versus sensim incrassatum, ibique coleopteris aequale, segmentis inaequalibus, anterioribus tribus longioribus, licet latitudine sua brevioribus, supra parce subtilissimeque punctatum. Pedes picei, tarsis flavescentibus.

Obs. Variat saepius corpore subcoerulescente: Al. coeruleam Sahlb. eandem putarem, ni notis pluribus, praecipue thoracis abdominisque structura, nimium discedere videatur.

- \* 3. *T. balteata*: Nigra, nitidula, subtiliter pubescens, thorace subovato, basi foveolato, abdomine basi angustato, segmentis anterioribus duobus rufis, pedibus piceis. — Long.  $1\frac{1}{2}$  lin.

Habitat in Austria, Helvetia, Italia.

Species insignis, statura omnino praecedentis. Subtiliter punctata, pube subtili depressa grisea vestita, nigra, capite, thorace elytrisque supra fusco-nigris, nitidulis. Antennae capite thoraceque vix longiores, apicem versus leviter incrassatae, nigro-piceae, basi nonnunquam dilutiores. Palpi picei. Caput thoracis prope latitudine, orbiculatum, basi profunde constrictum. Thorax coleopteris dimidio angustior, latitudine haud brevior, lateribus ante medium cum angulis anterioribus rotundatus, basin versus subangustatus, modice convexus, basi foveola minuta supra scutellum impressus. Coleoptera thorace paulo longiora, parum convexa, lateribus subrotundata, apice simul leviter emarginata. Abdomen basi angustatum, coleopteris dimidio angustius, apicem versus sensim leviter incrassatum, apice coleopteris latitudine subaequale, segmentis inaequalibus, anterioribus longioribus, licet latitudine sua brevioribus, supra parce subtilissime punctatum, segmentis duobus anterioribus rufis, posterioribus nigerrimis. Pedes picei, antici rufescentes, tarsi omnes rufo-testacei.

- \* 4. *T. ferialis*: Fusca, nitida, subtilissime pubescens, antennis basi pedibusque testaceis, thorace subovato, rufo-testaceo, abdomine basi angustato, nigro, segmentis anterioribus duobus rufis. — Long.  $1\frac{1}{2}$  lin.

Habitat in Sardinia, Dom. Prof. Gené.

Animulculum elegans, statura omnino praecedentis. Corpus totum

parcius subtilissime griseo-pubescens. Antennae capitis thoracisque longitudine, apicem versus sensim leviter incrassatae, piceae, basi testaceae. Palpi picei. Caput thoracis latitudine, orbiculatum, basi profunde constrictum, omnium subtilissime punctulatum, nigrum, nitidum. Thorax coleopteris vix dimidio angustior, latitudine vix longior, lateribus ante medium cum angulis anterioribus fortius rotundatis, basin versus angustatus, leviter convexus, dense subtiliter punctatus, supra scutellum foveola notatus, laete testaceo-rufus, nitidus. Coleoptera thorace vix longiora, lateribus subrotundata, apice simul leviter emarginata, minus confertim subtiliter punctata, nitida, fusca, humeris margineque apicali subtestaceis. Abdomen basi angustatum, coleopteris dimidio fere angustius, apicem versus sensim leviter incrassatum, apice coleopteris latitudine aequale, dorso segmentis anterioribus tribus laevigatis, posterioribus parce obsoleteque punctulatis, segmentis inaequalibus ut in proximis praecedentibus, anterioribus duobus rufis, sequentibus nigerrimis. Pectus fuscum. Pedes testacei.

Mas vertice thoracisque disco longitudinaliter leviter impressis.

- \* 5. *T. scitula*: *Nigra, subtiliter pubescens, antennarum basi, palpis tarsisque testaceis, thorace subovato, abdomine subangustato. — Long. 1 $\frac{1}{3}$  lin.*

*Er. Col. March. I. 308. 3.*

Habitat Berolini, Dom. Schüppel et ipse.

Praecedentibus minus gracilis, nigra, subtilissime confertim punctata, pube tenui griseo-sericea vestita. Antennae capite thoraceque breviores, apicem versus vix incrassatae, piceae, basi testaceae. Palpi testacei. Caput thorace parum angustius, orbiculatum, basi profunde constrictum. Thorax coleopteris perparum angustior, latitudine vix brevior, ante medium leviter rotundatus, basin versus subangustatus, convexiusculus, supra scutellum foveola minuta impressus. Elytra thorace paululum longiora, subdepressa, apice truncata. Abdomen basi coleopteris paulo angustior, lineare, segmentis subaequalibus, dorso aequali modo ac reliquum corpus punctatum. Pedes picei, tibiis tarsisque omnibus testaceis, illis medio fusciscentibus.

- \* 6. *T. exarata*: *Testacea, subtiliter pubescens, abdomine pone medium capiteque nigris, thorace subquadrato, late canaliculato, rufo. — Long. 2 lin.*

*Drusilla exarata* Mannerh. *Brachelytr.* 85. 2.

Habitat in Austria, Dom. Ullrich.

Statura omnino *T. umbraticae*, at sesqui maior, tota confertissime subtiliter punctata, pube brevissima sericante fulvescenti-grisea dense vestita. Antennae capite thoraceque parum longiores, apicem versus subincrassatae, testaceae. Caput thoracis latitudine, fronte leviter impressa, nigrum, oris partibus testaceis. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus ante medium leviter rotundatis, basi apiceque truncatus, parum convexus, dorso sat late longitudinaliter impressus, testaceo-rufus. Scutellum fuscum. Elytra thorace paulo longiora, fusco-testacea, basi dilutiora. Abdomen lineare, testaceum, segmentis aequalibus, duobus antepenultimis fusco-nigris. Pectus rufescenti-testaceum. Pedes dilute testacei.

- \* 7. *T. laesa*: *Nigro-fusca, opaca, antennarum basi pedibusque testaceis, abdomine lineari, nigro, nitido, thorace subquadrato, maris longitudinaliter excavato, feminae canaliculato. — Long. 1½ lin.*

Habitat in Sardinia, Dom. Prof. Gené.

Statura omnino *T. umbraticae*, eaque vix maior, thoracis structura praecedenti affinis, nigro-fusca, vix nitida, subtilissime tenuiter griseo-pubescentis. Antennae sat elongatae, capite thoraceque longiores, apicem versus perparum incrassatae, fuscae, articulo primo laete, sequentibus duobus obscure testaceis. Caput thoracis latitudine, suborbiculatum, densius subtilissime punctatum, ore subtestaceo, vertice maris longitudinaliter impresso. Thorax latitudine summa haud brevior, basin versus subangustatus, basi coleopteris dimidio angustior, truncatus, apice cum angulis anterioribus rotundatus, lateribus rectis, subdepressus, confertissime subtilissimeque punctatus, feminae fortiter canaliculatus, maris sat profunde lateque longitudinaliter excavatus. Elytra thoracis latitudine, subdepressa, dense subtiliterque punctata. Abdomen lineare, nigrum, nitidum, supra parce subtilissime punctatum, segmentis aequalibus. Pedes testacei.

Marem vertice impresso, thorace late longitudinaliter excavato, feminam thorace canaliculato, vertice integro differe in descriptione relatum.

- \* 8. *T. atra*: *Atra, opaca, subtilissime cinereo-pubescentis, thorace quadrato, leviter canaliculato, abdomine apicem versus sub-angustato, pedibus piceis, tarsis flavescens. — Long. 1½ lin.*

*Er. Col. March. I. 309. 5.*

*Aleochar. atra Grav. Mon. 162. 35. — Gyll. Ins. Suec. II. 395. 17. — Sahlb. Ins. Fenn. I. 358. 24.*

*Bolitochara atra Mannerh. Brachelytr. 79. 26. — Runde Brachel. Hal. 30. 8.*

*Staph. leucopus Marsh. Ent. Britt. 506. 22.*

Habitat in Europa.

*Atra*, opaca, vix perspicue punctata, pube tenuissima cinereo-sericea vestita. Antennae capite thoraceque longiores, tenues, apicem versus haud incrassatae, nigrae. Caput thorace vix angustius, orbiculatum, basi constrictum, fronte obsolete impressa. Thorax coleopteris paulo angustior, latitudine paululum brevior, neque basin neque apicem versus angustatus, lateribus ante medium rotundatus, angulis anterioribus obtusiusculis, subdeflexis, depressus, leviter canaliculatus. Coleoptera thorace paulo longiora, apice truncata. Abdomen minus elongatum, apicem versus subangustatum. Pedes graciles, picei, summa basi, geniculis tarsisque flavescens.

- \* 9. *T. umbratica*: *Nigra, nitidula, subtilissime griseo-pubescentis, thorace quadrato, obsolete canaliculato, abdomine lineari, antennis pedibusque rufo-piceis. — Long. 1½ lin.*

*Er. Col. March. I. 310. 6.*

Habitat in Germania, Gallia.

Praecedenti aequalis, nigra, nitidula, capite, thorace elytrisque fusconitentibus, subtilissime punctata, pube tenui griseo-sericea obducta. Antennae capite thoraceque paulo longiores, apicem versus levissime incrassatae,



rufo-piceae. Caput thoracis prope longitudine, suborbiculatum, basi constrictum, fronte obsolete impressa, ore piceo-testaceo. Thorax coleopteris dimidio angustior, latitudine haud brevior, neque basin neque apicem versus angustatus, lateribus ante medium et basi subrotundatus, apice truncatus, subdepressus, canalicula parum profunda, antrorsum obsolescente impressus. Coleoptera thorace paulo longiora. Abdomen parallelum, segmentis aequalibus. Pedes rufescenti-picei, tarsis flavescentibus.

10. *T. fugax*: *Nigra, nitidula, subtilissime griseo-pubescent, thorace elytrisque fuscis, antennarum basi pedibusque testaceis, thorace subquadrate, obsolete canaliculato, abdomine parallelo.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Sarmia, Mus. Reg. Taurin. Dom. Prof. Gené specimen singulum examandum communicavit.

Praecedenti aequalis, at minus gracilis, nigra, nitidula, subtilissime griseo-pubescent. Antennae capite thoraceque breviores, articulo tertio secundo parum minore, 4—10 transversis. ultimo maiusculo, praecedentibus ambobus longitudine aequali, ovato, subacuminato, fuscae, articulis duobus primis testaceis. Palpi testacei, maxillares articulo tertio fusco. Caput thoracis fere latitudine, suborbiculatum, confertissime subtiliter punctatum, nigrum, nitidum. Thorax coleopteris dimidio prope angustior, latitudine haud brevior, basi truncatus, apice cum angulis anterioribus et lateribus ante medium rotundatus, his dein fere rectis, basin versus subangustatus, leviter convexus, obsoletius canaliculatus, basi leviter transversim impressus, confertissime subtiliter punctatus, fuscus, nitidulus. Elytra thorace paululum longiora, apice leviter sinuata, confertissime minus subtiliter punctata, fusca, nitidula. Abdomen coleopterorum latitudine, parallelum, segmentis aequalibus, supra confertim subtiliter punctatum. Pedes testacei.

Obs A. praecedente antennarum structura facile distinguitur, a *T. scitula*, cui antennarum structura magis convenit, abdominis segmentis omnino aequalibus differt.

Species mihi invisae, forte huius generis.

11. *T. (?) flavitarsis*: *Nigra, subnitida, griseo-pubescent, antennis subfiliformibus, thorace longioribus, fusco-piceis, tarsis apiceque tibiarum testaceis, thorace longiore, postice foveola impresso.* — Long.  $1\frac{1}{4}$  lin.

*Aleochar. flavitarsis* Sahlb. Ins. Fenn. I. 349. 9.

Habitat in Fennia australi, rarissime.

Calodera nigrita paulo minor, structura antennarum ab illa imprimis distincta. Caput thoracis fere latitudine, orbiculatum, fronte subimpressa, nigrum, nitidum, subtilissime punctulatum, tenuiter pubescens, ore palpisque piceo-ferrugineis. Antennae thorace paulo longiores, subfiliformes, tenuiores, fusco-ferrugineae, articulis tribus baseos longis, primo subcylindrico, duobus insequentibus crassiore, 4—10 brevioribus, et extrorsum valde parum incrassatis, ultimo ovato, obtuso. Thorax latitudine fere longior, antice rotundatus, lateribus ante medium dilatatus, posterius paulo angustior, basi truncatus, angulis obtusis, supra niger, subnitidus, subtilissime punctulatus, tenuiter pubescens, canalicula longitudinali obsoletius, foveola autem ante scutellum evidentius impressa. Scutellum parvum, triangulare, nigrum, pu-

bescens. Elytra thorace latiora et paulo longiora, apice truncata, supra convexa, nigra, subnitida, subtilissime punctulata, pube griseo-sericea adspersa. Abdomen pone elytra longitudine reliqui corporis, nigrum, nitidum, parce pubescens, subtilissime punctulatum. Corpus subtus nigrum, nitidum, punctulatum, pubescens. Pedes elongati, tenues, pubescentes, nigri, tarsis apiceque tibiaram ferrugineo-testaceis. (Sahlb.)

12. *T. (?) coracina*: Nigra, nitida, subpubescens, tibiis picco-ferrugineis, tarsis pallidis, antennarum articulo ultimo elongato, thorace subquadrato, foveola postice impresso. — Long.  $1\frac{1}{2}$  lin.

Sahlb. Ins. Fenn. I. 350. 10. — Zetterst. Ins. Lappon. 76. 8.

Habitat in Lapponia, rarissime.

Magnitudine et statura *T. aterrimae* valde similis, sed differre videtur forma thoracis magis quadrata, et articulo antennarum ultimo magis elongato. Caput thorace angustius, orbiculatum, fronte impressa, nigrum, nitidum, parce pubescens, ore et palpis concoloribus. Antennae thorace longiores, apicem versus parum incrassatae, nigrae, subpubescentes, articulo primo longo, incrassato, secundo et tertio etiam elongatis, insequentibus brevibus, sensim crassioribus, ultimo duos antecedentes simul sumtos longitudine aequante. Thorax subquadratus, longitudine paulo latior, antice truncatus, angulis deflexis, lateribus parum rotundatus, angulis posticis rotundatis, basi subtruncatus, supra convexiusculus, niger, nitidus, pube rariore conspersus, subtilissime punctulatus, foveola ante scutellum impressa. Scutellum triangulare, nigrum, subtilissime punctulatum. Elytra thorace multo latiora et paulo longiora, apice truncata, ad angulum externum vix sinuata, supra subdepressa, nigra, nitida, pube rara conspersa, crebre et evidenter punctulata. Abdomen pone elytra longitudine reliqui corporis, nigrum, nitidum, parce et subtilissime punctulatum, subpubescens. Corpus subtus nigrum, nitidum, paulo profundius et crebrius punctatum, parce pubescens. Pedes elongati, femoribus nigris, tibiis ferrugineo-piceis, tarsis pallidis. (Sahlb.)

13. *T. (?) carbonaria*: Atra, subnitida, griseo-pubescens, thorace subquadrato, postice foveolato, antennarum articulo ultimo oblongo-ovato, pedibus elongatis, tarsis ferrugineo-testaceis. — Long.  $1\frac{1}{2}$  lin.

*Bolitochara carbonaria* Mannerh. *Brachelytr.* 75. 4.

*Aleochara carbonaria* Sahlb. Ins. Fenn. I. 351. 11.

Habitat in Fennia australi, rarissime.

A praecedente magnitudine minore, pubescentia densiore et articulo antennarum ultimo magis ovato, a *T. atra* antennis extrorsum crassioribus et a *T. (?) flavitarsi* thorace breviori differt. Caput thorace paulo angustius, suborbiculatum, nigrum, nitidum, vix punctulatum, palpis fusco-piceis. Antennae thorace longiores, nigrae, versus apicem parum incrassatae, articulis ut in praecedente, ultimo tamen magis ovato. Thorax longitudine vix latior, subquadratus, lateribus angulisque parum rotundatis, antice et postice truncatus, supra subconvexus, niger, subnitidus, tenuiter griseo-pubescens, vix punctulatus, foveola ante scutellum obsoletius impressa. Scutellum breve, triangulare, nigrum. Elytra, abdomen pone haec corpusque

subtus ut in *T. atra*. Pedes elongati, nigri, genubus anguste, tibiis apice et tarsis totis pallide ferrugineis. (Sahlb.)

11. *T. (?) coerulea*: *Coerulescens, subnitida, sericeo-pubescent, antennis nigro-fuscis, subfiliformibus, thorace subquadrato, foveola ante scutellum impresso. — Long.  $1\frac{1}{4}$  lin.*

*Sahlb. Ins. Fenn. I. 351. 12.*

Ad Helsingforsiam semel tantum capta.

*T. atrae* magnitudine et statura similis. Caput thoracis fere latitudine, rotundatum, nigro-coerulescens, nitidum, palpis concoloribus. Antennae thoracis longitudine, subfiliformes, versus apicem admodum parum incrassatae, articulo ultimo oblongo-ovato, fusco-nigrae, subpubescentes. Thorax longitudine vix lator, antice truncatus, lateribus ante medium parum dilatatus, postice subtruncatus, angulis omnibus rotundatis, supra convexiusculus, foveola ante scutellum impressa, coerulescens, sericeo-pubescent, subnitidus. Scutellum minutum, triangulare, nigro-coerulescens. Elytra thorace latiora et paulo longiora, postice truncata, supra convexiuscula, coerulescentia, sericeo-pubescentia, subnitida. Abdomen pone elytra reliqui corporis longitudine corqusque subtus nigro-coerulescentia, tenuiter sericeo-pubescentia. Pedes elongati, nigro-coerulescentes, tarsis piceo-ferrugineis.

Obs. Conf. quod apud *T. coarctatam* ad calcem notavi.

## **Phloeopora Er.**

*Er. Col. March. I. p. 311.*

*Maxillae mala interiore mutica, apice spinulis ciliata.*

*Ligula brevis, apice bifida; paraglossae nullae.*

*Palpi labiales 3-articulati, articulo secundo brevior.*

*Tarsi omnes 5-articulati, postici articulo primo sequente paulo longiore.*

Corpus elongatum, lineare, subdepressum, alatum. Caput exsertum, porrectum, thoracis latitudine, rotundatum, basi constrictum, oculis mediocribus, leviter prominulis. Labrum breve, truncatum, mandibulas non omnino obtegens. Mandibulae muticae. Maxillae mala exteriore cornea, basi apiceque membranea, portione membranea apicali pubescente; mala interiore exteriore paulo brevior, membranea, margine exteriore corneo, apice oblique introrsum emarginata, sinu spinulis brevibus ciliato. Palpi maxillares minus elongati, articulo tertio secundo paulo longiore, leviter incrassato, quarto parvo, subulato. Labium mento transversum, apice leviter emarginato, ligula brevi, palporum labialium articulum primum haud excedente, bifida, paraglossis nullis. Palpi labiales triarticulati, articulo secundo primo duplo fere minore, tertio tenuiore at primo parum brevior. Antennae breviusculae, articulis tribus primis leviter elongatis, secundo tertioque obconicis, inter se subaequalibus, ultimo apice obtusiusculo. Thorax coleopteris parum angustior, subquadratus, subdepressus. Elytra apice introrsum oblique truncata, extrorsum ad angulum exteriorem leviter sinuata. Abdomen elongatum, lineare. Pedes breves, intermediis basi approximatis, tarsis omnibus 5-articulatis, anterioribus articulis 4 primis brevibus, subaequalibus, postici articulo primo vix elongato.



## O b s e r v a t i o n e s .

1. Labium et maxillae omnino ut in Homalota (t. III. f. 8. et 22.).
2. Habitus linearis, subdepressus Homalotarum plurimarum, a quibus nisi tarsis anticis 5-articulatis vix distinguatur. Cum Oxypodis quoque hoc genus plurimis characteribus convenit, at habitus alius, caput porrectum, subremotum, pedes breves, et tarsi postici minus elongati, tibia breviores, articulus primus sequentibus perparum longior, ultimo brevior.
3. Sexus differentia mihi haud observata.
4. Victus sub arborum cortice. Species indigenae sub cortice Pinus degunt.

\* 1. *Ph. reptans*: *Atra, ore, antennis, elytris, pedibus anoque rubris, thorace transversim quadrato, angulis anterioribus obtusis.* — Long.  $1\frac{1}{3}$  lin.

*Er. Col. March. I. 311. 1.*

*Aleoch. reptans* Grav. Mon. 154. 19. — *Gyll. Ins. Suec. II. 389. 12.* — *Sahlb. Ins. Fenn. I. 354. 17.* — *Zetterst. Faun. Lappon. I. 98. 8. Ins. Lappon. 76. 11.*

*Bolitoch. reptans* Mannerh. *Brachelytr. 76. 10.* — *Runde Brachel. Hal. 30. 5.*

Habitat in Europa temperata et septentrionali sub cortice Pinus.

Fusco-atra, opaca, subtilissime dense punctata, pube tenuissima flavescenti-grisea obducta. Os rufo-testaceum. Antennae capite paulo longiores, articulo ultimo breviter ovato, apice subacuminato, testaceae, nonnunquam apicem versus fuscесcentes. Thorax coleopteris vix angustior, latitudine paulo brevior, lateribus subrectis, apice truncatus, angulis anterioribus obtusis, basi interdum obsolete impressus. Elytra confertim evidentius punctata, rubra, basi plerumque fuscесcentia. Abdomen supra crebrius punctulatum, segmentis piceo-marginatis, apice rubrum. Pedes rubri.

\* 2. *Ph. corticalis*: *Nigra, antennis pedibusque piceis, elytris fuscis, apice rufescentibus, thorace subquadrato, angulis anterioribus rotundatis.* — Long.  $1\frac{1}{4}$  lin.

*Er. Col. March. I. 312. 2.*

*Aleoch. corticalis* Grav. *Micr. 76. 13. Mon. 152. 12.* — *Latr. Hist. nat. Crust. et Ins. IX. 381. 13.*

*Aleoch. tenuis* Grav. *Micr. 84. 23.* — *Latr. Hist. nat. Crust. et Ins. IX. 384. 23.*

*Aleoch. teres* Grav. *Micr. 79. 19. Mon. 155. 20.* — *Latr. Hist. nat. Crust. et Ins. IX. 383. 19.*

Habitat in Europa sub cortice Pinus.

Praecedenti simillima, differt autem statura graciliore, colore magis obscuro, et thoracis structura. Nigra, minus confertim subtiliter punctulata, parum nitida, pube tenui minus densa vestita. Antennae capite sesqui longiores, articulo ultimo breviter ovato, apice obtusiusculo, piceae, articulis tribus primis rufis vel rufo-piceis. Caput thoracis latitudine, suborbiculatum, subtiliter punctulatum, fronte medio subtiliter obsolete canaliculata, ore piceo. Thorax coleopteris paulo angustior, latitudine non brevior, basin versus subangustatus, lateribus ante medium cum angulis anterioribus rotundatus, basi plerumque foveola transversa, magis minusve obsoleta impressus. Elytra fusca, apice magis minusve rufescentia. Abdomen pilosellum, supra crebrius punctatum, apice interdum piceo. Pedes picei.

- \* 3. *Ph. latens*: *Nigra, nitida, antennarum basi pedibusque testaceis, elytris rufis, basi fuscis, thorace subquadrato, angulis anterioribus rotundatis.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Pensylvania, Dom. Zimmermann; dedit Dom. Schüppel.

Statura et summa affinitas praecedentis. Nigra, nitida, griseo-pubescentis. Antennae capite sesqui longiores, articulo ultimo breviter ovato, apice obtusiusculo, nigro-piceae, articulis duobus primis testaceis, apicali brunneo. Palpi testacei. Caput thorace paulo minus, licet vix angustius, suborbiculatum, parce punctulatum, tenuiter subtiliterque pubescens, fronte subtiliter leviterque medio canaliculata. Thorax latitudine laud brevior, coleopteris paulo angustior, basin versus subangustatus, lateribus ante medium cum angulis anterioribus rotundatus, parcius subtiliter punctatus, tenuiter griseo-pubescentis. Elytra thorace paulo longiora, dense evidentius punctata, nitida, rufa, basi ultra tertiam partem fusca, densius pubescentia. Abdomen parcius punctulatum, pube longiore et minus subtili parcius vestitum, ano concolore. Pedes testacei,

Obs. Differt a praecedente praecipue corpore nitido et punctura longe parciore.

- \* 4. *Ph. subtilusa*: *Nigra, ore, antennis, pedibus anoque testaceis, elytris fusco-testaceis, thorace fusco, subquadrato, obsolete canaliculato, basi foveola transversa impresso.* — Long. 1 lin.

Habitat in Americae insula St. Johannis, Dom. Moritz.

Praecedenti affinis, at dimidio minor. Nigra, opaca, subtilissime confertim punctata, pube brevissima tenuissimaque vestita. Antennae rufo-testaceae, basi testaceae. Caput fronte obsolete canaliculata, ore testaceo. Thorax fuscus, coleopteris angustior, latitudine non brevior, angulis anticis rotundatis, depressus, obsoletius canaliculatus, basi foveola transversa impressus. Elytra fusco-testacea. Abdomen parcius punctatum, nigrum, nitidum, ano testaceo. Pedes testacei.

- \* 5. *Ph. colubrina*: *Brunnea, abdomine ante apicem capiteque nigricantibus, antennis, pedibus anoque testaceis, thorace obsolete canaliculato.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Brasilia, Dom. Thorey, in valle Araguensi Columbiae, Dom. Moritz.

Praecedentibus maior, habitu fere Lithocharidis. Corpus lineare, subdepressum, tenuiter pubescens. Antennae capite dimidio longiores, flavo-testaceae. Caput thoracis latitudine, basi profunde constrictum, depressum, nigro-piceum, ore testaceo. Thorax fuscus, latitudine haud brevior, basin versus sensim angustatus, apice, praecipue angulis anticis rotundatus, depressus, late obsolete canaliculatus, basi medio transversim impressus, utrinque puncto impresso notatus. Coleoptera dilute brunnea, vel fusco-testacea. Abdomen coleopteris angustius, elongatum, supra parce subtiliter punctatum, brunneum, ante apicem nigrum, apice testaceum. Pedes flavo-testacei.

6. *Ph. religiosa*: *Flavo-testacea, abdomine ante apicem nigro, capite nigricante, thorace basi foveola transversa impresso. — Long. vix ultra 1 lin.*

Habitat in Americae mer. ins. Puerto-Rico, Dom. Moritz.

Habitus omnino staturaque linearis depressa et summa affinitas praecedentis, at triplo minor, confertim subtiliter punctata, tenuiter cinereo-pubescentis, laete testacea, parum nitida. Antennae capite sesqui longiores, apicem versus sensim incrassatae. Caput thoracis latitudine, depressum, fronte obsolete canaliculata, piceum, ore testaceo. Thorax coleopterorum prope latitudine, latitudine paulo brevior, lateribus rectis, angulis posterioribus obtusis, anterioribus rotundatis, depressus, basi foveola angusta transversa impressus. Scutellum flavum. Elytra depressa, macula triangulari communi circa scutellum, aliaque valde obsoleta in medio margine laterali fusciscentibus. Abdomen supra parcius obsolete punctulatum, testaceum, apice fulvum, ante apicem, segmento penultimo dimidio et antepenultimo toto, nigrum. Pedes pallide testacei.

### **Hygronoma Er.**

*Er. Col. March. I. p. 312.*

*Homalota Curtis Britt. Ent. XI. t. 514.*

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula brevis, bifida; paraglossae nullae.*

*Palpi labiales 3-articulati, articulo secundo brevior.*

*Tarsi omnes 4-articulati, articulis brevibus depressis.*

Corpus elongatum, lineare, depressum, alatum. Caput exsertum, porrectum, thoracem latitudine paululum superans, subrhombicum, basi modice constrictum, oculis mediocribus, leviter prominulis. Labrum transversum, truncatum, mandibularum apicem haud obtegens. Mandibulae muticae. Maxillae mala exteriore cornea, basi apiceque membranacea, portione membranacea apicali pubescente, mala interiore exteriore paulo brevior, membranacea, margine exteriore corneo, apice oblique introrsum emarginata, sinu spinulis brevibus ciliato. Palpi maxillares breviusculi, articulo secundo tertioque longitudine subaequalibus, hoc modice incrassato, quarto minuto, subulato. Labium mento transversum, apice leviter emarginato, ligula brevi, palporum articulum primum haud excedente, bifida, paraglossis nullis. Palpi labiales triarticulati, articulo secundo primo duplo fere minore, tertio hoc tenuiore, at paulo longiore. Antennae filiformes, articulis tribus primis leviter elongatis, inter se longitudine subaequalibus, secundo tertioque obconicis, ultimo subacuminato. Thorax coleopteris vix angustior, basin versus subangustatus. Scutellum triangulare. Elytra apice truncata. Abdomen lineare. Pedes breves, intermedii basi approximati, tarsi omnibus breviusculis, depressis, 4-articulatis, articulo primo reliquis paululum longiore, triangulari, ultimo s. unguiculari etiam depresso, praecedentibus aequali.

### **Observationes.**

1. Labium et maxillae omnino ut in *Homalota* (t. III. f. 8. et 22.).

2. Habitus omnino *Homalotarum* quarundam linearium et maxime depressarum, pedibus brevibus autem et tarsis omnibus 4-articulatis, depressis, articulo ultimo non elongato hoc genus satis distinctum.



3. Dom. Curtis in splendido suo opere supra citato hoc insectum pro typo generis *Homalotae* depinxit, at Com. Mannerheim hoc genus ex *Aleochara plana* condidit, cui tarsi posteriores 5-articulati sunt, articulo ultimo elongato, reliquis coniunctis aequali, quem cel. Mannerheim generis sui characterem affert. Dom. Curtis ipse dubius *H. planam* speciei icone illustratae associat. Nomen quidem huic satis convenit, et magis quam generi sequenti, at typo primitivo servandum est. Ceterum generis huius (*Hygronomae*) characteres Dom. Curtis recte exhibuit.

4. Sexus differentia latet.

5. Victus in lacuum ripis et in paludosis, in arundinetis. Cursitat in arena humida inter arundines. Hybernatur intra arundinum culmos.

\* 1. *H. dimidiata*: Nigra, antennis basi pedibusque rufis, elytris dimidiato-flavis. — Long.  $1\frac{1}{4}$  lin.

Er. Col. March. I. 313. 1.

*Aleochara dimidiata* Grav. Mon. 149. 3.

*Homalota dimidiata* Curtis Britt. Ent. XI. t. 514.

Habitat in Germania.

Nigra, opaca, pube tenuissima flavescenti-grisea vestita. Antennae rufae, apicem versus fusciscentes. Frons interdum obsolete impressa. Thorax capitis vix latitudine, latitudine paulo brevior, apicem versus subangustatus, basi leviter canaliculatus. Elytra flava, basi late nigra. Pedes rufi.

## ***Homalota Mannerh.***

*Mannerh. Brachelytr. p. 73. — Er. Col. March. I. p. 314.*

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula brevis, bifida; paraglossae nullae.*

*Palpi labiales triarticulati, articulo secundo brevior.*

*Tarsi antici 4-, posteriores 5-articulati, postici articulis 4 primis aequalibus.*

Corpus vel elongatum, vel oblongum, saepe depressum, plerumque alatum. Caput exsertum, porrectum, basi magis minusve constrictum, iam thoracis latitudine, jam eodem angustius, oculis mediocribus, parum prominulis. Labrum apice rotundatum, mandibulas obtegens. Mandibulae muticae. Maxillae (t. III. f. 22.) mala exteriori cornea, basi apiceque membranea, portione membranea apicali pubescente, mala interiori exteriori paulo brevior, membranea, margine exteriori corneo, apice oblique introrsum emarginata, sinu spinulis brevibus ciliato. Palpi maxillares leviter elongati, articulo secundo tertioque longitudine subaequalibus, hoc leviter incrassato, quarto minuto, subulato. Labium (t. III. f. 8.) mento transversum, apice leviter emarginato, ligula brevi, palporum labialium articulum primum haud excedente, bifida, paraglossis nullis. Palpi labiales triarticulati, articulo primo maiore, secundo hoc duplo fere minore, tertio tenuiore at praecedente longiore. Antennae articulis tribus primis reliquis magis elongatis, secundo tertioque inter se aequalibus. Thorax iam lateribus parallelis, iam antrosum angustatus. Scutellum triangulare. Elytra apice truncata. Abdomen iam lineare, iam apicem versus angustatum. Pedes leviter elongati, tarsis anticis 4-, posterioribus 5-articulatis, illis articulis tribus, his articulis quatuor primis brevibus, inter se aequalibus.

## Observationes.

1. Com. Mannerheim ex unica specie, *Aleocho. plana* Gyll., proposuit hoc genus, tarsorum posticorum structura bene observata. Reliquas plerasque sub *Bolitochara* enumerat illust. auctor, quamquam tarsorum posticorum articulis 4 primis aequalibus cum *H. plana* omnino congruunt.

2. *Homalotae*, generis recte descripti, nomen temere mutare nolui, quamvis multis speciebus aegre quadrat. Occurrunt in hoc genere, sicut a me constitutum est, numerosissimo plerorumque aliorum huius tribus formae, ut saepenumero, nisi characteribus genericis acerrime observandis, de genere facile decipiari. Simulat e. g. *Hom. didyma* Autaliam, *H. trimaculata* *Bolitochara*, *H. velata* *Tachyusam*, *H. callicera* *Caloderam*, *H. lividipennis* *Oxypodam*, *H. cinnamomea* *Aleocharam*, *H. notha* *Gyrophaenam*, etc.

3. Sunt quaedam in hoc genere species, quae tarsorum posticorum articulo primo paululum elongato a ceteris discedere videntur, e. c. *H. concolor*, livida et ambigua, characteres autem reliqui huius generis in iis observantur, ut nulli alii generi adscribi possint, nisi *Oxypodae*, cuius habitum magis minusve distincte prae se ferunt, quacum autem tarsorum posticorum structura minus conveniunt, quam tarsis anticis 4-articulatis ab ea distinguuntur.

4. Mas plerumque antennis validioribus instructus, in quibusdam etiam abdominis structura, in aliis speciebus alio modo distinguitur, quas differentias in singulis speciebus memoratas invenies.

Secundum corporis totius et thoracis formam sectiones quatuor in hoc genere distinguam.

\* *Lineares*, thorace vel quadrato vel subquadrato.

\* 1. *H. graminicola*: *Nigra, nitida, elytris nigro-fuscis, pedibus piceis, thorace coleopteris angustiore, transversim subquadrato, postice late foveolato, abdomine supra parce obsoleteque punctato. — Long. 1½ lin.*

*Aleocho. graminicola* Grav. Mon. 176. 75.

*Aleocho. linearis* Gyll. Ins. Suec. II. 292. 15. — Sahlb. Ins. Fenn. I. 357. 22.

*Bolitochara linearis* Mannerh. Brachelytr. 79. 24.

*Aleocho. longiuscula* Gyll. Ins. Suec. IV. 485. 12–13. — Sahlb. Ins. Fenn. I. 355. 18.

*Bolitochara longiuscula* Mannerh. Brachelytr. 17. 13.

*Aleocho. longicornis* var. *b* Gyll. Ins. Suec. II. 405. 27.

*Aleocho. moesta* Zetterst. Faun. Lapp. I. 101. 15. Ins. Lappon. 77. 20.

*Bolitochara fracticornis* Boisd. et Lacord. Faun. Ent. Paris. I. 544. 7. (forte)

Habitat in Europa temperata et boreali.

*Nigra, nitida, parce tenuissime pubescens. Antennae capitis thoracisque longitudine, graciles, apicem versus parum incrassatae, nigrae, articulo primo piceo, 4–10 subobconicis, ultimo elongato-ovato, praecedente duplo prope longiore, apice acuminato. Caput thorace angustius, basi constrictum, parce subtilissime punctatum, fronte medio interdum obsolete impressa. Thorax coleopteris paulo angustior, latitudine dimidio fere brevior, lateribus et basi leviter rotundatus, apice truncatus, basin versus haud angustatus, angulis posterioribus obtusis, subdepressus, parce subtilissime punctatus, basi fovea lata, antrorsum saepe ultra medium producta, interdum etiam parva, raro fere nulla, impressus. Elytra thorace paulo longiora, obscure fusca, dense evidenter punctata. Abdomen supra laevigatum, segmentis anterieus punctis*

raris, posterius vix conspicuis sparsis. Pedes longiusculi, picei, tibiis tarsisque rufescentibus.

Mas differt fronte thoraceque magis impressis, et abdominis segmento superiore terminali punctis elevatis asperato.

Obs. 1. Thoracis structura ambigua inter hanc et sequentem sectionem.

2. Antennarum basis et pedes nonnunquam dilutiores: talia exempla cel. Gyllenhal nomine *Al. linearis* descripsisse puto: specimina Suecica non vidi, specimen tamen arcticum, ad Promont. boreale (Nordcap) lectum Dom. Schüppel dedit.

3. *Al. linearis* Gyll. magis ad marem, *Al. longiuscula* magis ad feminam spectare videtur. Vidi in collectione Dom. Prof. Germar specimen nom. *Al. aterrimae* e Fennia a cel. Prof. Sahlberg missum, quod cum feminis indigenis omnino congruit, et quocum descriptio *Al. longiusculae* apud Gyllenhal omnino convenit. Sub nomine *Al. aterrimae* a Sahlbergio plures species confusae esse videntur, qua de re v. quod ad *H. socialem* referam.

\* 2. *H. umbonata*: *Nigra, nitida, elytris pedibusque fusco-testaceis, thorace coleopteris angustiore, transversim subquadrato, postice leviter foveolato, abdomine supra parce subtiliter punctato, sublaevigato.* — Long.  $1\frac{3}{4}$  lin.

Habitat in Lusitania, Com. Hoffmannsegg, in Hispania meridionali, in Istria et in Austria Dom., Schüppel, in Sardinia, Dom. Prof. Gené, in Anglia, Dom. Rudd.

Statura et summa similitudo praecedentis. Nigra, nitida, tenuiter subtiliterque griseo-pubescens. Antennae capite thoraceque paulo longiores, minus graciles, apicem versus vix incrassatae, articulis 4—10 subcylindricis, crassitie haud brevioribus, ultimo oblongo-ovato, praecedente plus sesqui longiore, apice subacuminato, totae nigrae, unicolores. Palpi nigro-picei. Caput thorace paulo angustius, parce subtilius punctatum, fronte utrinque leviter impressa. Thorax coleopteris paulo angustior, latitudine dimidio fere brevior, lateribus et basi leviter rotundatus, apice truncatus, basin versus haud angustatus, angulis anterioribus subrotundatis, posterioribus obtusis, subdepressus, minus dense subtiliter punctatus, basi medio obsolete impressus. Elytra thorace sesqui prope longiora, depressa, confertim subtiliter punctata, testacea, margine exteriori fusciscente. Abdomen supra vage subtiliter punctatum, ceterum laevigatum. Pedes fusco-testacei, femoribus picescentibus.

Mas abdominis segmento dorsali secundo medio tuberculo minuto elevato, ultimo punctis elevatis subtiliter asperato.

\* 3. *H. rigidicornis*: *Fusca, subnitida, tenuiter griseo-pubescens, antennis validis, elytris, ano pedibusque testaceis, thorace coleopteris angustiore, basin versus leviter angustato, abdomine supra parce subtilius punctato.* — Long. 2 lin.

Habitat Genevae, Dom. Schüppel.

Praecedentibus maior, antennis validis et statura Myrmedoniam facile simulans, at huius generis. Antennae capite thoraceque fere longiores, crassae, articulo secundo tertio paulo minore, his obconicis, sequentibus seu articulis 4—10 spissis, cylindricis, crassitie haud brevioribus, ultimo praecedente duplo longiore, cylindrico, apice rotundato vel obtuse acuminato, totae testaceo-rufae. Palpi rufo-testacei, articulo tertio fortius incras-



sato. Labrum testaceum. Mandibulae rufo-piceae. Caput thorace angustius, suborbiculatum, basi constrictum, subtiliter punctatum, parum pubescens, fusco-nigrum, minus nitidum, fronte medio depressiuscula. Thorax coleopteris paulo angustior, latitudine summa dimidio fere brevior, lateribus ante medium fortiter rotundatis, basin versus modice angustatus, basi subrotundatus, apice levissime emarginatus, angulis anterioribus obtusis, subdeflexis, posterioribus obtusiusculis, leviter prominulis, levissime convexus, obsolete latius canaliculatus, confertim subtiliter punctatus, densius subtiliter griseo-pubescens, fusco-niger, parum nitidus. Scutellum dense punctatum, fuscum. Elytra thorace paulo longiora, confertim subtiliter punctata, rufo-testacea, parum nitida, fulvo-pubescentia. Abdomen nigrum, nitidulum, parcius pubescens, supra minus dense subtiliter punctulatum, segmento apicali piceo-testaceo, subtus confertim subtiliter punctatum, densius pubescens, segmentorum marginibus rufo-piceis. Pectus nigrum. Pedes flavo-testacei.

Specimen unicum, abdominis segmento dorsali secundo medio tuberculo minuto prominulo instructum, forte masculum.

- \* 4. *H. occulta*: *Linearis, nigra, opaca, elytris fuscis, pedibus fusco-testaceis, thorace transversim subquadrato, coleopteris angustiore, abdomine nitido, supra basi subtiliter parce punctato, apice laevi.* — Long.  $1\frac{2}{3}$  lin.

*Er. Col. March. I. 317. 4.*

Habitat in Germania.

Statura *H. elongatulae*. Corpus nigrum, pube tenuissima grisea vestitum. Antennae capitis thoracisque longitudine, apicem versus non incrassatae, articulis 4—10 subobconicis, longitudine haud crassioribus, ultimo ovato, apice acuminato, totae nigrae. Caput et thorax subtilissime alutacea, opaca, parce subtilissime punctata. Caput thoracis latitudine, basi constrictum, fronte maris leviter impressa, feminae subconvexa, nitidula. Thorax coleopteris subangustior, latitudine sesqui prope brevior, lateribus leviter rotundatus, basin versus subangustatus, subdepressus, basi foveolatus, interdum obsolete canaliculatus. Elytra thorace sesqui longiora, subtilissime alutacea, dense subtilius punctata, fusca. Abdomen nitidum, supra segmentis quatuor primis parce punctatis, ultimis fere laevibus. Pedes fusco-testacei.

Mas abdominis segmento superiore sexto laevi, apice emarginato, margine laterali elevato.

- \* 5. *H. pagana*: *Nigra, subnitida, thorace elytrisque fuscis, antennarum basi pedibusque testaceis, thorace subquadrato, coleopteris angustiore, abdomine supra parce obsoleteque punctato.* — Long.  $1\frac{2}{3}$  lin.

Habitat Lutetiae, Dom. Dr. Aubé.

Statura omnino *H. graminicolae*. Antennae capite thoraceque paulo longiores, apicem versus vix incrassatae, fuscae, articulis 2 vel 3 primis rufo-testaceis, sequentibus subcylindricis, crassitie paulo longioribus, ultimo ovato, acuminato. Palpi testacei. Caput thorace angustius, subovatum, fronte parum convexa, subtilius punctatum, linea media longitudinali laevi, tenuiter pubescens, nigrum, subnitidum, ore rufescente. Thorax coleopteris

paululum angustior, latitudine sesqui prope brevior, lateribus subrotundatus, basin versus haud angustatus subdepressus, aequalis vel nonnunquam obsolete canaliculatus, griseo-pubescent, crebre subtilius punctatus, fuscus. Elytra thorace parum longiora, depressa, crebre subtilius punctata, tenuiter griseo-pubescentia, testaceo-fusca. Abdomen nigrum, nitidum, supra parce griseo-pubescent, parce obsoleteque punctatum, subtus obsoletius punctulatum, segmentorum marginibus anoque ferrugineis. Pectus nigro-fuscum. Pedes testacei.

Mas: Abdominis segmento dorsali sexto punctis elevatis asperato.

Obs. Sequenti proxima et simillima, distinguenda autem thorace brevior, basin versus minime angustato, antennis longioribus et abdominis segmento dorsali sexto maris asperato. Ad *H. graminicolam* etiam proxime accedit, differt autem antennis basi testaceis, corporis colore opaco, etc

\* 6. *H. vestita*: Nigra, subnitida, elytris fuscis, antennarum basi pedibusque testaceis, thorace coleopteris angustiore, basin versus subangustato, obsolete canaliculato, abdomine supra parce obsoleteque punctato. — Long.  $1\frac{2}{3}$  lin.

*Paederus vestitus* Grav. Mon. 140. 6. — Oliv. Encycl. meth. Ins. VIII. 629. 11.

*Aleochara quisquiliarum* Gyll. Ins. Suec. II. 398. 20. — Sahlb. Ins. Fenn. I. 361. 30. — Zetterst. Faun. Lappon. I. 97. 4. Ins. Lappon. 75. 5.

*Bolitoch. quisquiliarum* Mannerh. Brachelytr. 80. 32.

Habitat in Germania, Suecia.

Statura praecedentium, nigra, nitidula, pube grisea sat conspicua vestita. Antennae maris capitis thoracisque longitudine, feminae iisdem paulo breviores, apicem versus vix incrassatae, rufae vel rufo-piceae, articulis tribus primis rufo-testaceis, sequentibus obconicis, longitudine haud crassioribus, ultimo oblongo-ovato. Os cum palpis rufo-testaceum. Caput thorace paulo angustius, subovatum, basi leviter constrictum, subdepressum, fronte utrinque crebrius subtiliter punctata. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus ante medium leviter rotundatus, pone medium subsinuatus, basin versus leviter angustatus, apice truncatus, subdepressus, densius subtiliter punctatus, obsolete subtiliterque canaliculatus, basi obsolete impressus, fusco-niger, lateribus saepe rufescentibus. Elytra thorace longiora, depressa, subtiliter dense punctata, fusca vel fusco-testacea. Abdomen nigrum, nitidum, apice rufo-piceum, supra parce obsoleteque punctatum, fere laevigatum. Pedes flavo-testacei.

Mas nonnisi antennis paulo longioribus et validioribus distinguendus.

\* 7. *H. callicera*: Linearis, subdepressa, fusca, thorace subquadrato, coleopteris angustiore, basi foveolato, abdomine supra laevi, antennis articulis duobus ultimis elongatis. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 319. 7.

*Aleochara callicera* Grav. Mon. 153. 17.

*Callicera obscura* Grav. Micr. 66. 1.

*Callicerus Spencei* Curtis Britt. Ent. X. t. 443.

Habitat in Germania, Anglia.

Statura fere proxime sequentium, ut paulo minor, fusca, opaca, pube

subtili cinerea leviter incana. Antennae in utroque sexu capite thoraceque longiores, validiusculae, articulis tribus primis nigro-, reliquis rufo-fuscis, articulo secundo tertioque obconicis, sequentibus omnibus cylindricis, antepenultimis duobus vel tribus transversis, feminae penultimo crassitie paulo longiore, ultimo hoc duplo fere longiore, apice obtuse acuminato, maris articulo penultimo crassitie duplo fere longiore, ultimo hoc sesqui longiore. Os testaceum, palpi maxillares articulo tertio tumido, ovato, fusciscente. Caput thorace paulo angustius, basi modice constrictum, suborbiculatum, vix perspicue punctulatum, nigricans, opacum, fronte subimpressa, vertice leviter canaliculato. Thorax latitudine haud brevior, basin versus subangustatus, basi coleopteris dimidio angustior, lateribus ante medium subangulato-rotundatus, apice fortiter angustatus, subdepressus, basi subcanaliculatus impressusque, vix perspicue punctulatus, opacus, nigro-fuscus. Elytra thorace paulo longiora, confertim subtiliter punctata, fusca, opaca. Abdomen lineare, parcius subtilissime pubescens, nigro-piceum, nitidulum, supra laevigatum, subtus subtilissime punctulatum. Pedes testacei, coxis femoribusque fuscis.

Articuli ultimi duo coniuncti in mare tertiae, in femina quartae fere antennarum parti-aequales.

Obs. 1. Antennarum structura in hac specie singularis, quare genus peculiare, *Callicerus*, primo a cel. Gravenhorst propositum, dein ab eodem auctore apte deletum, ab Anglis auctoribus restitutum est, quod autem, antennarum structura in hoc genere tam varia et inconstante, neque hoc caractere satis determinatum, neque alia quadum nota confirmatum videtur.

2. Cave ne corporis habitu et antennarum structura insignem hanc speciem pro *Calodera* habeas, sunt ei enim tarsi antici 4-articulati, et postici articulis 4 primis aequalibus huius generis.

3. Entomologiae Britannicae nitidus auctor loco supra citato nomine *Calliceri Spencei* marem depinxit, *Calliceri* hybridi feminam descripsit.

4. Idem auctor tarsos 5-articulatos affert, et pedem anticum tarso 5-articulato delineavit, quod nonnisi errore fieri potuit. Equidem distinctissime tarsos anticos 4-articulatos observavi.

\* 8. *H. pavens*: *Linearis, subdepressa, nigra, subtiliter sericeo-pubescens, elytris fuscis, antennarum basi pedibusque testaceis, thorace subquadrato, coleopteris angustiore, obsolete canaliculato, abdomine supra parcius subtiliterque punctulato.* — Long.  $1\frac{3}{4}$  lin.

*Er. Col. March. I. p. 689.*

*Homalota quisquiliarum Er. Col. March. I. 317. 5.*

Habitat in Germania.

Statura fere *H. elongatulae*, et maioribus eius individuis aequalis, eiusdem coloris, at densius punctata ideoque magis opaca. Antennae capite thoraceque fere longiores, apicem versus leviter incrassatae, piceae, basi testaceae, articulis 4—10 obconico-cylindricis, sensim brevioribus, primis eorum subelongatis, decimo crassitie paulo brevior, ultimo ovato, apice obtuse acuminato. Caput, thorax et elytra subtilissime confertim punctata, opaca, pube tenuissima griseo-sericea densius vestita. Caput thorace paulo angustius, rotundatum, basi constrictum, ore cum palpis rufo-testaceo. Thorax coleopteris paulo angustior, latitudine haud brevior, basin versus subangustatus, lateribus ante medium cum angulis anterioribus rotundatus, sub-



depressus, obsolete canaliculatus. Elytra thorace paulo longiora, fusca vel fusco-testacea. Abdomen nitidulum, supra parce pubescens, subtilissime minus confertim punctatum, segmento quinto laevigato glabroque. Pedes testacei.

Mas abdominis segmento dorsali quinto medio carinula apice acuminata munito et fronte leviter canaliculata distinctus.

- \* 9. *H. languida*: *Linearis, subdepressa, nigra, elytris anoque fuscis, antennarum basi pedibusque flavis, thorace subquadrato, coleopteris angustiore, basi foveolato, abdomine subtiliter confertissime punctato, opaco.* — *Long.*  $1\frac{3}{4}$ —2 lin.

*Er. Col. March. I.* 318. 6.

Habitat in Germania; in ins. Coreyra, Dom. Schüppel.

Statura omnino praecedentis. Corpus nigrum, abdomine excepto parce pubescens. Antennae capite thoraceque fere longiores, piceae, basi testaceae, articulis 4—10 subobconicis, omnibus subelongatis, licet sensim paulo brevioribus, et primis eorum paulo gracilioribus, decimo vero longitudine haud crassiore, apicali elongato-ovato, apice modice acuminato. Caput thorace angustius, rotundatum, basi constrictum, nigrum, nitidulum, subtilissime punctatum. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus subrotundatus, basin versus haud angustatus, angulis anterioribus fortiter rotundatis, subdepressus, confertim subtiliter punctatus, basi foveola obsoletiore impressus, plerumque piceus, subnitidus. Elytra thorace paulo longiora, fusco-testacea, confertissime subtiliter punctata. Abdomen nigrum, apice fuscescente, undique confertissime subtilissime punctatum, pube tenuissima densa obductum, segmentis duobus ultimis glabris, parce punctatis. Pedes testacei.

Mas abdominis segmento quinto dorsali tuberculo minuto acutiusculo munito distinctus.

Variat minus maturus corpore testaceo.

Obs. Affinis certe praecedenti abdomine subtilissime punctato, differt autem punctis confertissimis, pube brevissima densa occultis, et thorace haud canaliculato, basin versus nullo modo angustato.

- \* 10. *H. gracilicornis*: *Depressa, nigra, subnitida, confertim subtiliter punctata, densius griseo-pubescens, antennis tenuibus concoloribus, pedibus testaceis, elytris fuscis, thorace subquadrato, coleopteris angustiore, basi foveolato, abdomine supra confertim punctato.* — *Long.*  $1\frac{1}{2}$  lin.

Habitat in Austria, Dom. Ullrich.

Oblonga, depressa, nigra, subnitida, sat dense minus subtiliter griseo-pubescens, pube minus sericante. Antennae capite thoraceque paulo longiores, tenues, apicem versus haud incrassatae, nigrae, articulis 2—10 elongato-obconicis, inter se subaequalibus, ultimo elongato-ovato. Caput thorace paulo angustius, basi constrictum, subtiliter punctulatum, fronte medio leviter foveolata. Thorax coleopteris angustior, longitudine vix latior, basi lateribusque leviter rotundatus, basin versus haud angustatus, apice truncatus, confertim subtiliter punctatus, basi foveola minus profunda impressus.

Scutellum nigrum. Elytra sat lata, depressa, fusca, subtilissime confertissimeque punctata, pube densiore et magis sericante quam reliquum corpus vestita. Abdomen latiusculum, lateribus subrotundatum, supra minus subtiliter dense punctatum. Pedes testacei, femoribus picescentibus.

Obs. Praecedentibus affinis, at paulo brevior; magis depressa, antennis tenuibus Tachysae atrae, tarsis posticis autem huius generis.

- \* 11. *H. debilicornis*: *Depressa, nigra, subnitida, omnium subtilissime punctata, densius pube brevissima grisea vestita, antennis tenuibus elytrisque fuscis, thorace coleopteris angustiore, subquadrato, basi foveolato, abdomine supra parce subtilissime punctato.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura omnino et summa affinitas praecedentis. Nigra, subnitida, pube brevissima densa griseo-subsericans. Antennae capite thoraceque paulo longiores, graciles, apicem versus haud incrassatae, articulis 2—10 elongato-obconicis, secundo tertioque reliquis longioribus, ultimo elongato-ovato, fuscae, articulo primo subtestaceo. Palpi testacei. Caput thorace paulo angustius, basi leviter constrictum, parce subtilissime punctatum, depressum, fronte leviter impressa. Thorax latitudine haud brevior, coleopteris dimidio fere angustior, basin versus subangustatus, lateribus ante medium et basi leviter rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, subdepressus, confertim subtilissime punctatus, basi foveola obsoletius impressus. Elytra thorace paulo longiora, depressa, confertim subtilissime punctata, fusca, parum nitida. Abdomen parce subtilissime punctatum, segmentis quarto quintoque laevigatis, parce longius pubescens, nigrum, nitidum. Pedes testacei.

Mas abdominis segmento dorsali quinto medio tuberculo minuto compresso acutiusculo notatus.

- \* 12. *H. gregaria*: *Linearis, subdepressa, fusco-nigra, subnitida, sericeo-pubescens, elytris fuscis, pedibus testaceis, thorace subquadrato, basi foveolato, abdomine supra segmentis 4 anterioribus crebre subtiliter punctatis.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Aegypto, frequens, Dom. Prof. Ehrenberg, in Sardinia, Dom. Prof. Gené, in Austria, Dom. Ullrich, Berolini, ipse.

Habitus et summa affinitas *H. elongatae*, subnitida, nigra, capite thoraceque plerumque nigro-fuscis, vel fuscis, elytris dilutius fuscis, vel fusco-testaceis, abdomine toto nigro, nitidulo. Antennae capite thoraceque parum longiores, apicem versus parum incrassatae, articulis 4—10 subcylindricis, crassitie subbrevioribus, ultimo oblongo-ovato, totae rufo-brunneae. Palpi testacei. Caput thoracis prope latitudine, orbiculatum, basi sat fortiter constrictum, subtilissime punctulatum. Thorax coleopteris antice parum angustior, basin versus subangustatus, latitudine vix brevior, subtilissime punctulatus, subdepressus, basi supra scutellum impressus. Elytra thorace paulo longiora depressa, confertim subtiliter punctata. Abdomen supra segmentis 4 anterioribus confertim subtiliter punctatis, quinto laevi, sexto iterum punctulato. Pedes flavo-testacei.

Variat thoracis dorso late longitudinaliter impresso.

Obs. Ab II. elongatula praecipue thorace paulo longiore et abdominis segmento etiam quarto confertius punctulato distinguenda.

- \* 13. *H. ravilla*: *Linearis, nigra, opaca, subtilissime sericeo-pubescent, tibis tarsisque testaceis, thorace subquadrato, coleopteris angustiore, basi canaliculato, abdomine supra sublaevigato.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Austria, Dom. Ullrich.

Statura omnino II. elongatulae, et minutis eius individuis aequalis, nigra, minus nitida, abdomine supra nitidulo. Antennae capite thoraceque paulo longiores, apicem versus sensim subincrassatae, articulis penultimis vix transversis, ultimo magno, ovato, praecedente triplo fere longiore, corpori concolores. Caput thoracis latitudine, rotundatum, basi constrictum, omnium subtilissime punctulatum, fronte subimpressa. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus rectis, basi apiceque truncatus, angulis omnibus rotundatis, parum convexus, omnium subtilissime punctulatus, basi ad medium usque late et parum profunde canaliculatus. Elytra thorace paulo longiora, confertim subtiliter punctata, concolora. Abdomen lineare, supra segmentis anterioribus duobus parce subtiliter obsoleteque punctulatis, parceque pubescentibus, reliquis laevigatis. Pedes nigri, tibiis tarsisque picescenti-testaceis,

Obs. Colore nigro et abdomine supra fere laevigato nec non antennarum structura ab affinis satis distincta videtur.

- \* 14. *H. labilis*: *Nigra, nitida, tenuiter griseo-pubescent, antennarum basi pedibusque rufo-piceis, thorace subquadrato, leviter convexo, abdomine supra creberrime subtilius punctato.* — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 699. 7. a.

Habitat in Austria, Dom Ullrich, Berolini, Dom. Schüppel, Lutetiae Mus. Dom. Aubé, in Sardinia, Mus. Reg. Taurin.

Statura omnino, facies et summa similitudo Tachysae umbraticae, at huius generis, nigra, nitida, pube minus densa parum sericante grisea vestita. Antennae capite thoraceque paulo longiores, tenuiusculae, apice leviter incrassatae, nigrae, articulis 3 vel 4 primis rufescenti-piceis, sequentibus elongato-obconicis, penultimis solis crassitie vix longioribus, ultimo oblongo-subacuminato. Caput thorace paulo angustius, rotundatum, basi constrictum, subtiliter punctulatum. Thorax coleopteris angustior, latitudine vix brevior, lateribus anterioribus subrotundatus, basin versus subangustatus, leviter convexus, subtilissime punctulatus, basi foveola minuta angusta transversa impressus. Elytra thorace paulo longiora, sat crebre subtiliter punctata, concolora. Abdomen parallelum, supra confertim subtilius punctatum. Pedes rufo-picei, femoribus nigricantibus.

Mas fronte impressa et antennis paulo longioribus distinctus.

- \* 15. *H. velata*: *Nigra, sericeo-pubescent, thorace transversim subquadrato, abdomine toto subtilissime punctulato, antennis fuscis basi pedibusque pallide flavis.* — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 319. 8.

Habitat in Germania, Gallia.



Statura et habitu fere Tachyusae. Antennae capite thoraceque breviores, apicem versus vix incrassatae, articulis penultimis longitudine paulo crassioribus, ultimo ovato, fuscae basi flavae, vel flavae apice subinfuscatae. Corpus atrum, parum nitidum, totum pube tenuissima densa cinereo-sericeum. Oris partes testaceae. Caput thorace paulo angustius, suborbiculatum, basi leviter constrictum. Thorax latitudine dimidio fere brevior, coleopteris paulo angustior, basin versus subangustatus, lateribus ante medium cum angulis anterioribus rotundatus, subdepressus, basi foveola punctiformi notatus. Elytra thorace sesqui longiora. Abdomen sub pube omnium subtilissime punctulatum. Pedes toti dilute flavi.

Obs. A praecende pube sericea conferta, thorace antennisque brevioribus et pedibus totis pallide flavo-testaceis distincta.

\* 16. *H. luteipes*: *Linearis, nigra, subnitida, antennis longioribus piceis, pedibus laete testaceis, thorace subquadrato, leviter convexo, abdomine supra subtilissime confertim punctato.* — Long.  $1\frac{1}{3}$  lin.

Er. Col. March. I. 320. 9.

Habitat Berolini.

Statura fere sequentis. Nigra, subnitida, pube tenerrima vix conspicua sericea. Antennae elongatae, capite thoraceque longiores, graciliores, apicem versus leviter incrassatae, nigro-piceae, basi apiceque plerumque fusco-piceae, articulis 4—10 subobconicis, crassitie haud brevioribus, terminali oblongo-ovato. Caput thorace paulo angustius, basi leviter constrictum, subtiliter densius punctatum, oris partibus testaceis. Thorax coleopteris vix angustior, latitudine haud brevior, basin versus haud angustatus, angulis anterioribus late rotundatis, supra leviter convexus, basi interdum obsolete foveolatus. Elytra thorace paulo longiora, subtilissime dense punctata. Abdomen supra anterieus subtiliter confertissime, posterius subtilissime dense punctatum. Pedes laete testacei.

Obs. A sequente praecipue thorace antennisque longioribus et abdomine confertim punctato distincta.

\* 17. *H. elongatula*: *Linearis, subdepressa, nigra, subnitida, sericeo-pubescentis, elytris fuscis, antennarum basi pedibusque testaceis, thorace transversim subquadrato, abdomine supra segmentis anterioribus 3 crebre punctulatis.* — Long.  $1\frac{1}{4}$ —2 lin.

Var. a: *Thorace aequali.*

*Aleoch. terminalis* Grav. Mon. 160. 29. — Gyll. Ins. Suec. II. 397. 19. — Sahlb. Ins. Fenn. I. 360. 27.

*Bolitoch. terminalis* Mannerh. Brachelytr. 80. 30.

Var. b: *Thorace basi foveolato.*

*Aleoch. elongatula* Grav. Micr. 79. 18. Mon. 153. 18. — Latr. Hist. nat. Crust. et Ins. IX. 383. 18. — Gyll. Ins. Suec. II. 398. 18. — Sahlb. Ins. Fenn. I. 359. 25. — Zetterst. Faun. Lappon. I. 97. 3. Ins. Lappon. 75. 3.

*Aleoch. teres* Gyll. Ins. Suec. II. 390. 13. — Sahlb. Ins. Fenn. I. 355. 19. — Zetterst. Faun. Lappon. I. 98. 7. Ins. Lappon. 76. 10.

*Bolitoch. teres* Mannerh. Brachelytr. 76. 7.

Var. c: *Thorace obsolete canaliculato.*

*Bolitoch. complana* Mannerh. *Brachelytr.* 79. 29.

*Aleoch. complana* Sahlb. *Ins. Fenn.* I. 359. 26.

*Bolitoch. oblonga* Boisd. et Lacord. *Faun. Ent. Paris.* I. 548. 15.

*Bolitoch. exilis* Mannerh. *Brachelytr.* 80. 31.

*Aleoch. exilis* Sahlb. *Ins. Fenn.* I. 360. 28.

*Bolitoch. elongatula* Boisd. et Lacord. *Faun. Ent. Paris.* I. 548. 14.

*Bolitoch. planiuscula* Mannerh. *Brachelytr.* 80. 33.

*Bolitoch. depressiuscula* Mannerh. *Brachelytr.* 80. 34.

*Aleoch. depressiuscula* Sahlb. *Ins. Fenn.* I. 362. 32.

Habitat in Europa.

Linearis, subdepressa, nigra, subnitida, dense subtilissime punctata, pube tenuissima griseo-sericea vestita. Antennae capitis thoracisque longitudine, apicem versus vix incrassatae, articulis 4—10 subobconicis, crassitie haud brevioribus, ultimo oblongo-ovato, apice acuminato, nigro-piceae, basi obscure testaceae, raro piceae. Caput thorace paulo minus, basi constrictum, subdepressum, subtilissime punctatum, oris partibus testaceis. Thorax coleopterorum fere latitudine, latitudine dimidio fere brevior, basin versus haud angustatus, subdepressus, confertissime punctatus, vel basi foveolatus, vel dorso obsolete canaliculatus, vel interdum omnino aequalis. Elytra thorace dimidio longiora, fusca, opaca. Abdomen supra nitidulum, segmentis 3 primis crebrius subtiliter, 4 et 5 vage, ultimo iterum dense punctatis, apice iam testaceo, iam piceo, iam concolore. Pedes testacei.

Species vulgaris, quodammodo instabilis, statura corporis lineari subdepressa, et abdominis punctura semper cognoscenda. Forte species complures sub hoc nomine coniungo, quarum autem differentiae, quanto magis eas petii, tanto magis me fugerunt. Varietatum agmen hoc modo coegi:

I. Antennae articulis 4—10 crassitie haud brevioribus.

A. Abdomen apice piceo. Long.  $1\frac{1}{3}$ — $1\frac{2}{3}$  lin.

a. Thorace basi foveolato vel omnino aequali. — Apud non frequentissima.

b. Thorace canaliculato. — Sat frequens.

Huc *Bolitochara complana* et *depressiuscula* Mannerh. et *Bolitochara oblonga* Lacord. pertinent.

B. Abdominis apice testaceo segmentorumque singulorum marginibus rufis. Antennae plerumque totae rufae. Long.  $1\frac{2}{3}$ — $1\frac{3}{4}$  lin. — Haud infrequens.

a. Thorace basi foveolato. Hanc pro *Aleochara elongatula* sua Dom. Gyllenhal Dom. Schüppel communicavit.

b. Thorace canaliculato. — Huius specimen a Gyllenhal nomine A. terminalis acceptum Dom. Schüppel communicavit. *Bolitochara exilis* et *planiuscula* Mannerh. et *Bolitochara elongatula* Lacord. ad hanc revocandae.

C. Abdomine nigro, ano etiam concolore.

a. Abdominis segmentis dorsalibus tribus primis crebre profundiusque punctatis. Long. 2 lin. — Reliquis maior et robustior, ceterum haud distincta. Rarior.

b. Abdominis segmentis dorsalibus tribus primis crebrius punctulatis.

Long.  $1\frac{1}{2}$  lin. — Haud infrequens, a typo nonnisi ano concolore divergens. Huic adscribenda Bolitochara oblonga Mannerh. Brachelytr. 79. 26. — Aleochoch. elongatula var. b Gyll. Ins. Suec. II. p. 396. Sahlb. Ins. Fenn. I. p. 359. Zetterst. Ins. Lappon. p. 75.

c. Abdominis segmentis dorsalibus tribus primis parce subtiliterque punctatis. Long.  $1\frac{1}{3}$  lin. — Reliquis paulo minor et abdominis anteriore etiam dorso sublaevigato a reliquis discedens. Huius exempla 4 Dom. Schüppel communicavit, unum Berolini lectum, reliqua a Gyllenhal nomine Al. teretis e Suecia accepta.

II. Antennae articulis 4—10 crassitie paulo brevioribus, leviter transversis, totae nigrae. Anus etiam concolor. Reliquis minor, long.  $1\frac{1}{4}$  lin., ceterum vix distinguenda. Haud infrequens cum reliquis occurrit.

Variat praeterea pallidior, rufo-picea, capite abdominisque cingulo ante apicem nigricantibus. Huc pertinet forte Bolitochara longiuscula Boisd. et Lacord. Faun. Ent. Paris. I. 545. 8.

\* 18. *H. linearis*: *Linearis, nigra, antennis basi apiceque, pedibus anoque testaceis, fronte thoraceque subquadrato canaliculatis, elytris thoracis longitudine, rufo-brunneis, abdomine supra parce subtiliter punctulato.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 322. 12.*

*Aleochoch. linearis Grav. Micr. 69. 2. Mon. 148. 2. — Latr. Hist. nat. Crust. et Ins. IX. 378. 2.*

Habitat in Germania.

Statura fere praecedentium. Antennae capitis thoracisque longitudine, apicem versus leniter incrassatae, articulis penultimis transversis, ultimo leviter ovato, acuminato, rufo-testaceae, ante apicem fuscae. Corpus nigrum, subnitidum, tenuiter griseo-pubescens. Caput thorace paulo angustius, basi constrictum, subtilissime punctatum, minus nitidum, fronte subdepressa, parce subtiliterque punctata, parum profunde canaliculata, ore testaceo. Thorax coleopterorum fere latitudine, latitudine dimidio brevior, lateribus parum rotundatis, apice truncatus, angulis anterioribus obtusis, subdeflexis, depressus, omnium subtilissime punctulatus, in utroque sexu late canaliculatus. Elytra thorace vix longiora, depressa, confertim subtiliter punctata, rufo-brunnea. Abdomen nigrum, nitidum, apice testaceum, segmentis anterioribus margine posteriore rufo-piceo, supra parce subtiliter punctatum, quarto quintoque in utroque sexu sublaevigatis. Pedes testacei.

Mas nonnisi thoracis frontisque canalicula latiore distinctus.

Obs. Elytris thoraci longitudine subaequalibus ab affinibus discedit.

\* 19. *H. angustula*: *Linearis, subdepressa, nigra, antennarum basi, elytris pedibusque testaceis, thorace quadrato, canaliculato, abdomine confertissime subtiliter punctulato.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 322. 13.*

*Aleochoch. angustula Gyll. Ins. Suec. II. 393. 16. — Sahlb. Ins. Fenn. I. 357. 23. — Zetterst. Faun. Lappon. I. 97. 2. Ins. Lappon. 75. 2.*

*Bolitoch. angustula Mannerh. Brachelytr. 79. 25. — Boisd. et Lacord. Faun. Ent. Paris. I. 547. 13. — Runde Brachel. Hal. 30. 7.*

*Aleochoch. linearis var. Grav. Mon. 149. 2.*



Habitat in Europa.

Angusta, linearis, subdepressa, nigra, subnitida, subtilissime dense punctata, tenuiter dense griseo-pubescent. Antennae capitis thoracisque longitudine, apicem versus subincrassatae, articulis penultimis longitudine paulo crassioribus, rufo-piceae, basi testaceae. Caput thoracis latitudine, basi constrictum, feminae medio obsolete canaliculatum, maris late impressum: oris partibus testaceis. Thorax coleopterorum latitudine, latitudine vix brevior, lateribus ante medium leviter rotundatus, depressus, in mare late, in femina subtilius canaliculatus, opacus, niger, interdum piceus. Elytra thorace vix longiora, piceo-testacea, circa scutellum picea. Abdomen supra confertissime subtiliter punctatum, segmento sexto in mare supra granulato. Pedes testacei.

Mas fronte impressa, thorace late canaliculato et abdominis segmentis dorsalibus quinto sextoque granulatis distinctus.

\* 20. *H. aequata*: *Linearis, depressa, nigra, opaca, elytris fuscis, antennarum basi pedibusque rufis, thorace subquadrato, late canaliculato, abdomine nitidulo, parcius punctulato.* — Long.  $1\frac{2}{3}$  lin.

Er. Col. March. I. 323. 14.

Habitat in Germania.

Praecedenti affinis, at paulo magis elongata depressaque. Antennae capite sesqui longiores, articulis 6—10 transversis, ultimo subconico, piceae, articulis duobus primis rufis. Caput thoracis fere latitudine, nigrum, opacum, obsolete punctatum, fronte feminae plana, maris leviter impressa: oris partibus rufis. Thorax latitudine vix brevior, coleopteris paululum angustior, lateribus aequaliter leviter rotundatus, depressus, in utroque sexu late canaliculatus, subtilissime confertissimeque punctatus, niger, opacus. Coleoptera thorace vix longiora, depressa, subtiliter dense punctata, fusca, apice rufescentia. Abdomen nigrum, segmentis primis apice obscure rufo-piceis, supra subtiliter minus dense punctatum, nitidulum. Pedes rufo-testacei.

Mas fronte impressa et abdominis segmento dorsali quinto granulato distinctus.

Obs. Proxima praecedenti, a qua praecipue colore capitis, thoracis elytrorumque magis opaco, fronte maris minus impressa, feminae planata, et abdomine minus subtiliter et minus confertim punctato discedit.

\* 21. *H. nigella*: *Linearis, depressa, nigra, antennis basi piceo-rufis, thorace subquadrato, leviter canaliculato, abdomine supra crebre fortius punctato.* — Long.  $1\frac{1}{4}$  lin.

Er. Col. March. I. 323. 15.

Habitat Berolini.

Statura praecedentium, at paulo minor. Corpus depressum, nigrum, unicolor, tenuissime parce griseo-pubescent. Antennae capite thoraceque breviores, apicem versus sensim incrassatae, articulis penultimis transversis, ultimo ovato, basi truncato, apice obtuso, articulis duobus primis obscure rufis. Palpi picei. Caput thoracis latitudine, basi constrictum, depressum,

dense minus subtiliter punctatum, opacum, fronte feminae plana, maris leviter impressa. Thorax coleopteris vix angustior, latitudine parum brevior, basin versus subangustatus, lateribus ante medium leviter rotundatus, depressus, opacus, confertim subtiliter punctatus, late et parum profunde canaliculatus. Elytra thorace paulo longiora, depressa, dense minus subtiliter punctata, opaca. Abdomen supra nitidulum, segmentis 4 anterioribus crebre fortiusque, ultimis tribus parcius subtiliusque punctatis. Pedes nigri, geniculis tarsisque piceis.

Mas abdominis segmento dorsali sexto subtiliter granulato distinctus.

- \* 22. *H. arcana*: *Linearis, subdepressa, nigra, antennis concoloribus, pedibus testaceis, thorace subquadrato fronteque late longitudinaliter impressis, abdomine supra parce subtiliterque punctato, nitidulo.* — Long.  $1\frac{1}{3}$  lin.

Habitat in saltu Thuringensi, Dom. Kellner.

Statura omnino *H. nigellae*, nigra, parum nitida, subtiliter griseo-pubescent. Antennae capitis thoracisque prope longitudine, apicem versus vix incrassatae, articulo secundo tertioque aequalibus, 4—10 longitudine haud vel parum crassioribus, ultimo praecedentibus ambobus longitudine aequai, ovato, subacuminato, totae nigrae. Palpi nigri. Caput thorace parum angustius, subquadratum, obsolete punctulatum, fronte longitudinaliter impressa. Thorax coleopteris parum angustior, latitudine tertia fere parte brevior, lateribus subrectis, basi rotundatus, apice truncatus, angulis anterioribus obtusiusculis, deflexis, depressus, crebre subtilius punctatus, dorso secundum totam longitudinem late et parum profunde licet medio paulo profundius impressus, margine laterali pilosello. Elytra thorace sesqui prope longiora, confertim subtilius punctata, piceo-nigra. Abdomen supra nitidulum, parce subtiliter obsoleteque punctatum, segmento quinto sublaevi, sexto apice emarginato, septimo conspicuo ut in *Oxytelinis*. Pedes testacei.

- \* 23. *H. plana*: *Linearis, depressa, nigra, antennis piceis, pedibus anoque testaceis, fronte thoraceque transversim subquadrato canaliculatis, elytris thorace sesqui longioribus, brunneis, abdomine supra parce subtiliter punctato.* — Long.  $1\frac{1}{4}$  lin.

*Mannerh. Brachelytr.* 73. 1. — *Er. Col. March.* I. 700. 15. a.

*Aleoch. plana* Gyll. *Ins. Suec.* II. 402. 24. — *Sahlb. Ins. Fenn.* I. 367. 40.

Habitat in Germania, Suecia.

*H. lineari* proxima eidemque simillima, at dimidio minor, fronte minus subtiliter punctata et elytris longioribus distincta, nigra, parum nitida, subtiliter tenuiterque pubescens. Antennae capite sesqui longiores, apicem versus leviter incrassatae, articulis 5—10 subtransversis, ultimo breviter ovato, obtuse acuminato, pilosellae, iam dilutius, iam saturatius rufo-piceae. Palpi rufo-picei. Caput thorace parum angustius, suborbiculatum, basi parum constrictum, depressum, late parum profunde canaliculatum, crebre fortius punctatum, apice laeve, ore piceo. Thorax coleopteris paulo angustior, latitudine dimidio prope brevior, lateribus ante medium leviter rotundatus, basin versus subangustatus, depressus, omnium subtilissime confertim punctatus, late canaliculatus, canalicula basi profundiore, apice evanescente.

Elytra thorace sesqui longiora, depressa, dense subtiliter punctata, iam fusco-, iam rufo-testacea. Abdomen nitidum, segmentis tribus anterioribus margine postico piceo, dorso parce subtiliterque punctatis, sequentibus laevigatis, apice testaceum. Pedes testacei, tibiis tarsisque dilutioribus.

Obs. Habitu, capite thoraceque canaliculatis et fronte fortius punctata *H. cuspidatae* affinis, differt autem statura maiore, ano mutico, etc.

\* 24. *H. debilis*: *Linearis, subdepressa, fusca, ore, antennis, elytris, pedibus anoque testaceis, thorace subquadrato, leviter canaliculato, abdomine supra subtiliter punctulato.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 321. 11.*

Habitat Berolini.\*

Praecedente gracilior, fusca, tenuiter griseo-pubescent. Antennae capitis thoracisque longitudine, apicem versus parum crassiores, articulis 4—10 obconicis, crassitie haud brevioribus, ultimo ovato, subacuminato, totae testaceae. Caput thoracis latitudine, basi constrictum, subtiliter punctatum, nitidulum, medio obsolete canaliculatum, ore testaceo. Thorax coleopteris vix angustior, basin versus subangustatus, latitudine parum brevior, subnitidus, dense subtiliter punctatus, subdepressus, tenuiter canaliculatus. Elytra thorace paulo longiora, subtiliter dense punctata, fusco-testacea. Abdomen supra nitidulum, segmentis primis parcius subtiliter, quinto sparsim punctatis, sexto in utroque sexu laevi, septimo densius punctato, testaceo. Pedes pallide testacei.

Mas nonnisi antennis paulo longioribus, apicem versus distinctius incrassatis, articulo ultimo oblongo-ovato distinguitur.

Obs. *H. elongatulae* affinis, at gracilior, antennis laete testaceis, thorace longiore, basin versus subangustato et abdomine confertius punctatis differt.

25. *H. gracilentata*: *Elongata, linearis, nigra, nitida, antennis, ore, pedibus anoque testaceis, elytris fusco-testaceis, thorace subquadrato, aequali, abdomine supra laevigato.* — Long.  $1\frac{1}{3}$  lin.

Habitat Lutetiae, Mus. Dr. Aubé.

Statura omnino *H. elongatulae* et minimis eius individuis aequalis, nigra, nitida, subtiliter pubescens. Antennae capitis thoracisque fere longitudine, apicem versus sensim incrassatae, articulis penultimis transversis, ultimo ovato, subacuminato, totae testaceo-flavae. Caput thoracis prope latitudine, basi profundius constrictum, subtilissime punctulatum, ore flavo-testaceo. Thorax coleopterorum prope latitudine, latitudine paululum brevior, subdepressus, aequalis, sat crebre subtiliter punctulatus. Elytra thorace sesqui longiora, parce obsoleteque subtiliter punctulata, fusco-testacea (basin versus magis fusca, apicem versus magis testacea). Abdomen supra laevigatum, apice testaceum. Pedes testaceo-flavi.

Obs. *H. debili* coloribus similis, statura autem paulo minore, minus depressa, thorace haud canaliculato, abdomine supra laevigato distincta.



- \* 26. *H. macella*: *Linearis, depressa, rufo-testacea, dense subtiliter griseo-pubescent, ore, antennis pedibusque testaceis, thorace quadrato fronteque obsolete canaliculatis, abdomine supra subtilissime punctulato.* — Long. 1 lin.

Habitat Passaviae in Bavaria meridionali, Dom. Walzl.

Habitus fere Lithocharidis, *H. debili* duplo minor, paulo angustior et multo magis depressa, rufo-testacea, opaca, pube brevi albida subsericans, omnium subtilissime et vix perspicue punctata. Antennae capitis thoracisque prope longitudine, apicem versus parum incrassatae, articulis penultimis subtransversis, ultimo oblongo-ovato, totae flavo-testaceae. Caput thoracis latitudine, basi fortiter constrictum, subquadratum, depressum, obsolete canaliculatum, ore testaceo. Thorax coleopteris perparum angustior, latitudine haud brevior, lateribus rectis, basi apiceque vix rotundatus, depressus, obsolete canaliculatus. Elytra thorace paulo longiora. Abdomen supra segmentis singulis margine apicali dilute testaceis, penultimis basi piceis, apice toto testaceo. Pectus piceum. Pedes flavo-testacei.

- \* 27. *H. brevicornis*: *Elongata, linearis, subdepressa, nigra, nitida, elytris pedibusque fusco-testaceis, antennis capitis longitudine, abdominis segmentis 2 et 3 supra bicarinatis.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Brasilia.

Corpus admodum elongatum, lineare, subdepressum, nigrum, nitidum, subtiliter parce punctatum, tenuissime pubescens. Antennae capitis longitudine, apicem versus incrassatae, articulis 5—10 fortiter transversis, ultimo brevior, acuminato, piceae, basi rufescentes. Caput thoracis latitudine, transversim subquadratum, vertice transversim leviter impresso, fronte medio obsolete foveolata. Thorax subquadratus, latitudine paulo brevior, lateribus rectis, basi leviter rotundatus, apice leviter bisinuatus, subdepressus, aequalis. Elytra thorace paulo longiora, fusca, disco fusco-testacea, evidentius punctata. Abdomen supra laevigatum, segmento secundo et tertio utrinque acute carinato, sexto apice carinulis duabus elevatis, ultimo apice tricuspi. Pedes breves, fusco-testacei.

Obs. Habitu quibusdam *Phloeoporis* non dissimilis, at tarsi antici mihi quadriarticulati videntur.

- \* 28. *H. illota*: *Depressa, nigra, tibiis, tarsis elytrisque luteis, his circa scutellum infuscatis, capite fortius punctato thoraceque basin versus subangustato late canaliculatis, abdomine supra parce punctato.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Brasilia.

Habitu fere *H. elongatulae*, at brevior et paulo magis depressa, nigra, parum nitida. Antennae capitis thoracisque longitudine, apicem versus sensim subincrassatae, articulis 5—10 leviter obconicis, longitudine vix crassioribus, ultimo ovato, pilosellae, nigrae, articulis tribus primis piceis. Palpi maxillares picei, labiales pallidi. Caput thoracis latitudine, basi modice constrictum, planatum, dense fortius punctatum, obsolete canaliculatum, antice laeve. Thorax coleopteris angustior, latitudine paululum brevior, lateribus leviter rotundatis, basin versus subangustatus, depressus, longitudina-

- liter late at parum profunde impressus, confertissime subtiliter punctatus, uti caput pube densa, erecta, brevissima, nigra vestitus. Elytra thorace paulo longiora, depressa, subtiliter punctulata, tenuiter griseo-pubescentia, lutea, macula magna communi triangulari circa scutellum margineque laterali fuscis. Abdomen supra parce subtiliter punctatum parceque cinereo-pubesceus, nitidum. Pedes nigri, tibiis tarsisque rufo-testaceis.

- \* 29. *H. intrusa*: *Linearis, subdepressa, nigra, opaca, elytris, antennarum basi pedibusque fusco-testaceis, thorace coleopteris angustiore, subquadrato, medio late impresso, abdomine nitidulo, antice crebre punctato, postice laevigato.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Brasilia.

Habitu fere *H. elongatulae*, at minor, nigra, opaca, pube tenui vestita. Antennae capitis thoracisque longitudine, validiusculae, apicem versus sensim leviter incrassatae, articulis penultimis transversis, ultimo oblongo-ovato, nigrae, basi obscure testaceae. Caput thoracis fere latitudine, depressum, basi parum constrictum, subtiliter punctulatum, fronte plana. Thorax subquadratus, coleopteris plus dimidio angustior, latitudine vix brevior, lateribus ante medium rotundatis, depressus, subtiliter dense punctulatus, fovea lata longitudinali media per totam fere longitudinem excavatus. Elytra thorace paulo longiora, depressa, dense subtiliter punctata, fusco-testacea. Abdomen nitidum, supra segmentis anterioribus tribus crebre punctatis, posterioribus laevigatis, apice testaceum. Pedes testacei.

- \* 30. *H. cuspidata*: *Linearis, depressa, fusca, capite thoraceque obsolete canaliculatis, abdomine apice testaceo, mucronato.* — Long.  $\frac{3}{4}$  lin.

*Er. Col. March. I. p. 689.*

*Homalota plana Er. Col. March. I. 324. 16.*

Habitat in Europa, sub arborum cortice.

Minuta, valde depressa, linearis, fusca. Antennae capite sesqui longiores, apicem versus incrassatae, articulis duobus primis tumidulis, penultimis transversis, ultimo subgloboso. Caput thoraci fere aequale, subquadratum, depressum, minus dense fortius punctatum, linea media laevi, plerumque subimpressa, ore testaceo. Thorax coleopteris non angustior, latitudine parum brevior, lateribus subrectis, depressus, subtiliter punctatus, leviter canaliculatus. Elytra thorace paulo longiora, subtiliter punctata, fusco-testacea. Abdomen basi fuscum, dein nigricans, apice testaceum, supra subtilius punctatum, segmento ultimo apice acute mucronato. Pedes testacei.

Individuis nonnullis, forte masculis abdominis segmenta dorsalia intermedia utrinque tuberculo minuto instructa.

Variat colore saepius in testaceum, rarius in nigrum vergente.

Obs. Statura pusilla et abdomine apice mucronato facile cognoscenda.

- \* 31. *H. immersa*: *Linearis, depressa, nigra, nitida, elytris fuscis, antennarum basi pedibusque testaceis, fronte thoraceque subquadrato late impressis, abdomine supra parce punctato.* — Long.  $\frac{3}{4}$  lin.

*Er. Col. March. I. 324. 17.*

Habitat in Germania. Unicum specimen Berolini legi.

Praecedente latior, plana, nigra, nitida, tenuissime parce pubescens. Antennae capitis thoracisque longitudine, articulis 4 — 10 aequalibus, transversim obconicis, ultimo maiore, ovato, pilosellae, piceae, basi pallide testaceae. Caput thoracis latitudine, transversim subquadratum, planum, fronte tota impressa, subtilissime parce punctata. Thorax antice coleopteris parum angustior, et latitudine vix brevior, basin versus subangustatus, subtiliter parce punctulatus, obsolete canaliculatus, a basi ultra medium late parum profunde impressus. Elytra thorace sesqui prope longiora, crebre distinctius punctata, fusco-testacea. Abdomen supra vage subtiliter punctatum, apice piceo-testaceum. Pedes dilute fusco-testacei.

- \* 32. *H. vilis*: *Linearis, subdepressa, fusca, pedibus testaceis, thorace subquadrato, obsolete foveolato, abdomine nigro, supra confertim subtiliter punctato, ano piceo.* — Long. 1 lin.

*Er. Col. March. I. 325. 18.*

Habitat in Germania. Singulum exemplum Berolini legi.

Habitu fere *H. elongatulae*, at multo minor, angusta, linearis, subdepressa, abdomine excepto fusca, opaca, subtilissime punctata, omnium tenuissime sericeo-pubescens. Antennae capitis thoracisque longitudine, apicem versus parum incrassatae, articulis 4 — 10 longitudine vix crassioribus, ultimo ovato, rufo-piceae. Caput thoracis fere latitudine, basi leviter constrictum, aliquantulum distinctius punctatum, fronte medio obsolete canaliculata, oris partibus testaceis. Thorax coleopteris non angustior, latitudine paulo brevior, lateribus subrectis, angulis omnibus rotundatis, subdepressus, basi foveola minuta obsolete impressus. Elytra thorace longiora, dense subtiliter punctata, apice subdilutiora. Abdomen distinctius griseo-pubescens, undique confertim subtiliter punctatum, nitidum, nigrum, ano rufo-piceo. Pedes testacei.

- \* 33. *H. caesula*: *Linearis, fusca, antennis pedibusque testaceis, thorace subquadrato, fortiter canaliculato, abdomine nigro, parce subtilissime punctato, ano piceo.* — Long.  $\frac{3}{4}$  lin.

Habitat in Germania. Singulum specimen Dom. Schüppel dedit.

Praecedenti habitu affinis, at paulo minor et minus depressa, capite basi haud constricto et thorace profunde canaliculato satis distincta. Nigro-fusca, subnitida, parce tenuissimeque griseo-pubescens, abdomine solo nigro. Antennae capitis thoracisque longitudine, apicem versus sensim modice incrassatae, articulis penultimis longitudine crassioribus, ultimo oblongo-ovato, pilosellae, totae obscure testaceae. Caput thoracis latitudine, basi haud constrictum, parce subtilissime punctulatum, ore testaceo. Thorax coleopterorum fere latitudine, latitudine vix brevior, lateribus subrectis, apice truncatus, angulis anterioribus obtusis, deflexis, posterioribus cum basi rotundatis, modice transversim convexus, subtilissime punctatus, canalicula lata sat profunda exaratus. Elytra thoracis longitudine, depressa, distinctius punctata, fusca. Abdomen nigrum, nitidum, apice piceum, supra segmentis tribus anterioribus parce subtiliterque punctatis, posterioribus laevigatis. Pedes testacei.



\* 34. *H. circellaris*: Aptera, rufo-testacea, subnitida, capite abdominisque cingulo piceis, thorace subquadrato, basi foveolato. — Long. 1 lin.

Er. Col. March. I. 315. 1.

*Al. circellaris* Grav. Mon. 155. 22. — Gyll. Ins. Suec. II. 388. 10. — Sahlb. Ins. Fenn. I. 353. 14. — Zett. Faun. Lapp. I. 98. 6. Ins. Lapp. 76. 9.

*Bolitochara circellaris* Mannerh. Brachelytr. 75. 5. — Runde Brachelytr. Hal. 30. 2.

Habitat in Europa.

Angusta, linearis, obscure rufo-testacea, subnitida, subtiliter griseo-pubescent. Antennae capitis thoracisque longitudine, apicem versus leviter incrassatae, basi dilutiores. Caput thorace angustius, oblongum, basi profunde constrictum, subtilissime punctatum, piceum, ore rufo-testaceo. Thorax coleopterorum latitudine, longitudine non lator, lateribus leviter, apice parum, basi fortiter rotundatus, supra leviter convexus, basi foveolatus. Elytra depressa, brevia, confertim punctata. Abdomen ante apicem nigro-piceum, apice ipso ferrugineo, supra segmentis anterioribus confertim, posterioribus vage punctatis. Pedes testacei.

Mas differt elytris basi uni-tuberculatis, abdominis segmento penultimo supra ante apicem denticulo prominulo armato.

Obs. *Bolitochara inguinalis* Mannerh. Brachelytr. 76. 6. Aleochar. inguinalis Sahlb. Ins. Fenn. I. 353. 14., quam thorace longiore, foveola ante scutellum et duabus longitudinalibus impresso, antennis brevioribus, extrorsum multo magis incrassatis ut et colore fusco-ferrugineo, capite abdomineque obscurioribus distinctam esse Dom. Com. Mannerheim affert, a Dom. Prof. Sahlberg pro *H. circellaris* varietate habetur; specimen eius quoque nonnisi singulum observatum, et impressiones thoracis longitudinales facile fortuitae, quam ob rem differentiae eius specificae in praesens dubiae videntur.

\* 35. *H. procidua*: Aptera, nigra, fortius punctata, thorace subquadrato, convexiusculo, pedibus piceis. — Long. 1 lin.

Er. Col. March. I. 315. 2.

Habitat in Germania. Specimen unicum Berolini legi.

Statura fere praecedentis, at paulo crassior, nigra, parum nitida, dense profundius punctata. Antennae capitis thoracisque longitudine, apicem versus subincrassatae, nigrae. Caput thoraci latitudine fere aequale, basi leviter constrictum, labro piceo, palpis nigris. Thorax latitudine paululum lator, coleopteris subangustior, convexus, lateribus subdeflexis praecipue ante medium rotundatis. Coleoptera thoracis longitudine. Abdomen crassiusculum, supra subtilius aequaliter punctatum. Pedes picei.

Obs. Species minuta, corpore crassiusculo et punctura forti profunda in solita insignis.

\*\* Lineares; thorace transverso.

✕ \* 36. *H. brunnea*: Depressa, testacea, nitida, capite abdominisque segmentis penultimis piceis, thorace transverso, postice late subimpresso, elytris parcius subtiliter punctatis, abdomine supra crebre fortius punctato, ante apicem laevigato. — Long. 2 lin.

Er. Col. March. I. 326. 19.

*Staph. brunneus* Fab. Ent. Syst. Suppl. 180. 44. Syst. El. II. 600. 55.

*Aleoeh. depressa* Gyll. *Ins. Suec. II.* 401. 23. — *Sahlb. Ins. Fenn. I.* 366. 39.  
*Bolitochara depressa* Mannenh. *Brachelytr.* 81. 41 — *Runde Bruchel. Hal.* 31. 11.  
*Staph. nigriceps* Marsh. *Ent. Britt.* 515. 51.

Habitat in Europa temperata et septentrionali.

Corpus oblongum, depressum, testaceum, nitidum, pube brevi erecta grisea et pilis parvis erectis nigris vestitum. Antennae capite thoraceque breviores, tenuiores, articulo secundo tertio paulo brevior, articulis 6—10 transversis, ultimo praecedentibus ambobus longitudine aequali, oblongo-ovato, acuminato, testaceae, apice plerumque ultra medium fuscae nigraeve. Caput thorace multo angustius, basi haud constrictum, parce subtilius punctatum, fronte depressa, medio late subimpressa, piceum, nitidum, ore testaceo. Thorax coleopteris perparum angustior, latitudine dimidio brevior, basi leviter, lateribus fortius rotundatus, apice truncatus, angulis posterioribus rotundatis, subdepressus, dorso posterior late leviter impressus, parce subtilius punctatus, pilis raris erectis nigris sparsus. Elytra thorace paulo longiora, depressa, minus dense subtilius punctata, pilis raris erectis nigris sparsa, circa scutellum nonnunquam infuscata. Abdomen depressum, supra segmentis 4 primis crebre profundius punctatis, quarto sparsum punctato, quinto laevi, sexto iterum crebre punctato, apice leviter emarginato, pube brevi erecta parciore grisea et pilis raris erectis nigris vestitum, subtus aequaliter densius punctatum et pube longiore depressa fulvescenti-grisea vestitum, testaceum, segmentis quarto et quinto nigris, praecedentibus nonnunquam basi piceis. Pedes dilute testacei.

Mas nonnisi antennis paulo longioribus et validioribus distinguuntur.

\* 37. *H. depressa*: *Depressa, testacea, capite abdominisque segmentis pen-ultimis piceis, thorace transverso, postice leviter longitudinaliter impresso, elytris abdomineque punctatis, hoc apicem versus sublaevi.* — *Long.*  $1\frac{2}{3}$ —2 lin.

*Aleoeh. depressa* Grav. *Micr.* 100. 49. *Mon.* 176. 74. — *Latr. Hist. nat. Crust. et Ins. IX.* 391. 49.

*Bolitoch. depressa* Boisd. et Lacord. *Faun. Ent. Paris. I.* 551. 18.

Habitat in Germania.

Statura omnino praecedentis, at paulo angustior, dilutius testacea, minus nitida, pube fulvescenti-grisea subdepressa et pilis raris erectis nigris vestita. Antennae capite thoraceque paulo longiores, tenuiores, apicem versus vix crassiores, articulo secundo tertio paulo brevior, sequentibus inter se subaequalibus, subobconicis, longitudine haud crassioribus, ultimo maiusculo, praecedentibus ambobus longitudine aequali, oblongo-ovato, apice acuminato, fuscae, articulis 3 vel 4 primis pallidius testaceis. Caput thorace multo angustius, basi vix constrictum, depressum, fronte antice subimpressa, utrinque parce obsoleteque punctata, piceum, vel piceo-testaceum, ore testaceo, saepius etiam totum testaceum. Thorax coleopteris tertia parte angustior, longitudine plus dimidio brevior, lateribus et basi leviter rotundatus, apice truncatus, angulis posterioribus obtusis, subdepressus, posterius aut leviter canaliculatus aut latius longitudinaliter leviter impressus, parce subtiliterque punctatus, pilis erectis nigris raris instructus. Elytra thorace sesqui longiora, depressa, minus dense subtilius punctata, pilis erectis nigris

raris obsita, circa scutellum saepius infuscata. Abdomen depressum, supra segmentis anterioribus tribus parce subtilius punctatis, posterioribus sublaevibus, pube parca grisea et pilis raris erectis nigris vestitum, subtus crebrius subtilius punctatum, pube longiore depressa fulvescenti-grisea vestitum, testaceum, segmentis penultimis duobus nigro-piceis. Pectus piceo-testaceum. Pedes flavo-testacei.

Mas abdominis segmento dorsali quinto apice leviter tuberculato, sexto apice quadridentato insignis.

Obs. Praecedenti simillima atque mihi olim cum ea confusa et pro eius mare habita. Differt statura paulo graciliore, colore minus nitido, antennis longioribus, articulis penultimis haud transversis, thorace lateribus minus rotundato, abdomine supra parcius et subtilius punctato et differentia sexuali. Specimina Brunsvicensia a Gravenhorstio descripta ad hanc pertinent.

\* 38. *H. nigrifrons*: Depressa, testacea, capite abdominisque segmentis penultimis nigris, thorace transversim subquadrato, leviter longitudinaliter impresso, elytris dense subtiliter punctatis, abdomine supra crebre fortius punctato, ante apicem laeviore. — Long.  $1\frac{2}{3}$  — 2 lin.

Habitat in Gallia, Italia.

Statura omnino praecedentis, testacea, minus nitida, pube depressa griseo-fulvescente densius vestita. Antennae capitis thoracisque longitudine, tenuiores et apicem versus haud incrassatae, articulo tertio secundo vix longiore, 4—10 inter se fere aequalibus, subobconicis, crassitie sublongioribus, ultimo praecedente sesqui longiore, oblongo-ovato, acuminato, pilosellae, fuscae, articulis tribus primis testaceis. Caput thorace multo angustius, basi vix constrictum, depressum, parcius obsolete punctatum, fronte obsolete canaliculata, supra nigrum, ore testaceo, subtus testaceum. Thorax coleopteris paululum angustior, latitudine tertia parte brevior, lateribus et basi leviter rotundatus, apice truncatus, angulis posterioribus obtusis subrotundatis, subdepressus, longitudinaliter late subimpressus, crebrius subtiliter punctatus, rufo-testaceus, pube densiore et pilis raris erectis vestitus. Scutellum rufo-testaceum. Elytra thorace paulo longiora, depressa, dense subtiliter punctata, densius pubescentia, fusco-testacea. Abdomen depressum, supra pube depressa parciore et pilis erectis nigris raris vestitum, segmentis tribus primis crebre punctatis, piceis, margine testaceis, quarto quintoque parce vageque punctatis, nigris, hoc apice testaceo, sexto iterum crebrius punctato, testaceo, subtus crebre punctatum, pube longiore et densiore vestitum, basi apiceque testaceum, medio piceum, ante apicem nigrum. Pectus piceo-testaceum. Pedes flavo-testacei.

Obs. 1. *H. brunneae* et *depressae* perquam affinis, cum illa abdominis punctura, cum hac antennarum structura convenit; differt ab utraque elytris dense punctatis, densius pubescentibus, at pilis erectis nullis instructis, et thorace longiore. Secundum thoracis structuram forte sectioni praecedenti adscribenda fuisset, ni summa affinitate cum speciebus duabus modo laudatis iungeretur.

2. Vidi hoc insectum in collectione Dom. Aubé nomine *Bolitochara nigriceps* Dej. signatum, quod autem, cum idem nomen specificum a Marshamio *H. brunneae* tributum sit, mutandum censui,



39. *H. atricilla*: *Depressa, testacea, nitida, capite abdominisque segmento antepenultimo nigro-piceis, thorace brevi, transverso, aequali, elytris abdomineque supra parcius subtiliter punctatis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Sardinia, Mus. Reg. Taurin.

Praecedentibus affinis, at minor, praecipue brevior, testacea, nitida, pube longiore subdepressa grisea vestita. Antennae capitis thoracisque longitudine, apicem versus sensim subincrassatae, articulo tertio secundo paulo longiore et tenuiore, 4—10 subobconicis, sensim paulo maioribus, longitudine haud crassioribus, ultimo praecedentibus ambobus longitudine subaequali, oblongo-ovato, subacuminato, pilosellae, fuscae, basi testaceae. Palpi testacei. Caput thorace angustius, suborbiculatum, minus depressum, parce subtiliter obsoleteque punctatum, nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine duplo prope brevior, basi et lateribus leviter rotundatus, antrorsum subangustatus, apice truncatus vel levissime emarginatus, angulis posterioribus modice obtusis, nullo modo rotundatis, subdepressus, aequalis, densius subtiliter punctatus, rufo-testaceus, margine laterali parce pilosello. Elytra thorace sesqui longiora, depressa, minus confertim subtiliter punctata, luteo-testacea, ante apicem extrorsum fuscescencia. Abdomen subdepressum, supra parce subtilius punctatum, rufo-testaceum, segmento tertio piceo-testaceo, quarto nigro-piceo, quinto basi piceo, apice sextoque toto dilute testaceis. Pectus piceo-testaceum. Pedes dilute testacei.

Obs. Thorace brevi a praecedentibus facile distinguitur.

- \* 40. *H. oblonga*: *Depressa, picea, capite nigro, elytris pedibusque testaceis, fronte fortius punctulata, thorace transversim subquadrato, dorso late subimpresso, abdomine antice parce subtiliterque punctato, postice laevigato.* — Long.  $1\frac{2}{3}$  lin.

*Aleoch. elongatae* var. Grav. Mon. p. 154.

Habitat in Germania.

Statura omnino *H. depressae*, at paulo minor, picea, subnitida, pube depressa grisea vestita. Antennae capitis thoracisque longitudine, apicem versus vix incrassatae, articulo tertio secundo paulo longiore, 4—10 subcylindricis, 4—7 crassitie haud brevioribus, 8—10 leviter transversis, ultimo maiusculo, praecedentibus ambobus longitudine aequali, oblongo-subovato, apice obtuse acuminato, totae piceae. Palpi piceo-testacei. Caput thorace angustius, basi parum constrictum, depressum, fronte crebrius sat fortiter punctata, nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine tertia parte fere brevior, basi et lateribus leviter rotundatus, antrorsum subangustatus, apice truncatus, angulis posterioribus obtusis, subdepressus, parce subtiliterque punctatus, dorso pone medium late subimpressus, nigro-piceus, nitidus, versus latera magis brunneus. Elytra thorace tertia parte longiora, depressa, densius subtiliter punctata, fusco-testacea, circa scutellum infuscata, subnitida. Abdomen depressum, parce pubescens, supra segmentis tribus primis parce subtiliterque punctatis, posterioribus tribus laevigatis, subtus parce subtiliterque punctatum, nigro-piceum, ante apicem nigrum. Pectus piceum. Pedes testacei.

Obs. Thoracis structura propius ad sectionem praecedentem aggreditur, statura corporis autem omnino cum praecedentibus convenit, licet colore satis discedit.

Frons fortius punctata huic speciei peculiaris videtur, cuius unicum specimen solum in collectione Hoffmannseggiana nomine supra citato signatum mihi hucusque obvium fuit.

- \* 41. *H. hepatica*: Nigra, antennis elytrisque brunneis, pedibus testaceis, thorace depresso, abdomine supra parce punctato, nitidulo, segmento sexto apice pliculis duabus conniventibus. — Long.  $1\frac{1}{4}$  lin.

Habitat in Germania. Exemplum unicum in Pomerania legi.

Statura praecedentis, *H. sociali* propior, antennis rufo-brunneis et abdominis structura praecipue distincta, piceo-nigra, minus nitida, crebrius punctata, pube minus tenui depressa flavescens vestita. Antennae capitis thoracisque longitudine, validiores, apicem versus leviter sensim incrassatae, articulis 4—10 cylindricis, penultimis vix crassitie brevioribus ultimo praecedentibus ambobus longitudine aequali, elongato-conico, acuminato, rufae, basi rufo-piceae. Oris partes rufo-piceae. Caput thorace angustius, fronte subdepressa. Thorax latitudine dimidio brevior, coleopteris vix angustior, lateribus ante medium rotundatis, dein subrectis, parum convexus, dorso longitudinaliter depressus, sat crebre et parum subtiliter punctatus. Elytra thorace paulo longiora, depressa, sat crebre minus subtiliter punctata, rufo-brunnea. Abdomen nigrum, apice testaceum, segmentorum marginibus brunneis, supra nitidum, parce subtiliter punctatum, segmentis posterioribus tribus laevigatis. Pedes testacei.

Specimen singulum, segmento quinto apice pliculis duabus elevatis approximatis postice conniventibus instructo: forte mas.

- \* 42. *H. aemula*: Nigra, nitida, antennarum basi, elytris pedibusque testaceis, thorace subdepresso, basi leviter foveolato, obsolete canaliculato, abdomine supra laevigato, segmentis anterioribus subtiliter vage punctatis. — Long.  $1\frac{1}{2}$  lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Statura et summa similitudo sequentis et maximis eius individuis (var. a) aequalis, nigra, nitidissima, pube tenui raraque vestita. Antennae capitis thoracisque longitudine, apicem versus leviter incrassatae, articulis penultimis leviter transversis, ultimo oblongo-ovato, subacuminato, nigrae, basi testaceae. Palpi testacei. Caput thorace angustius, rotundatum, basi modice constrictum, parce obsoleteque punctatum. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus et basi leviter rotundatis, apice truncatus, angulis omnibus obtusis, subtiliter minus dense punctatus, parum convexus, obsolete canaliculatus foveolaeque parum profunda basi impressus. Elytra thorace sesqui longiora, confertim evidentius punctata, testacea, nitida, lateribus fuscis. Abdomen ut in *H. sociales* var. a. Pedes testacei.

- X \* 43. *H. socialis*: Nigra, tenuiter pubescens, elytris pedibusque testaceis, thorace subdepresso, basi obsolete foveolato, abdomine supra anterieus parce subtiliterque punctato. — Long.  $1\frac{1}{3}$ —2 lin.

Er. Col. March. I. 327. 21.

*Staphyl. socialis* Payk. Mon. Staph. 60. 43. Faun. Suec. III. 407. 51. — Oliv. Ent. III. 42. 37. 53. t. 3. f. 25.

*Aleochar. socialis* Gyll. Ins. Suec. III. 406. 28. — Sahlb. Ins. Fenn. I. 370. 46. — Zetterst. Faun. Lappon. I. 99. 11. Ins. Lapp. 76. 14.

*Bolitochara socialis* Mannerh. *Brachelytr.* 77. 17. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 546. 10.

*Aleoch. Boleti* Grav. *Micr.* 80. 21. *Mon.* 156. 23. — *Latr. Hist. nat. Crust. et Ins.* IX. 384. 21.

*Staphyl. sordidus* Marsh. *Ent. Britt.* 514. 48.

*Var. a.*

*Aleoch. sericans* Grav. *Mon.* 159. 28. — *Gyll. Ins. Suec. II.* 404. 26. — *Sahlb. Ins. Fenn. I.* 367. 14. — *Zetterst. Faun. Lappon. I.* 100. 13. *Ins. Lapp.* 77. 18.

*Bolitochara sericans* Mannerh. *Brachelytr.* 77. 14.

*Homalota sericans* Er. *Col. March.* 1. 326. 20.

*Bolitochara castanoptera* Mannerh. *Brachelytr.* 77. 16.

*Aleoch. castanoptera* Sahlb. *Ins. Fenn. I.* 69. 45.

*Var. b.*

*Aleoch. longicornis* Gyll. *Ins. Suec. II.* 405. 27. — *Sahlb. Ins. Fenn. I.* 386. 43. *Zetterst. Faun. Lappon. I.* 100. 14. *Ins. Lappon.* 77. 19.

*Bolitochara Boleti* *Boisd. et Lacord. Faun. Ent. Paris. I.* 545. 9.

*Var. c.*

*Aleochara nigrifula* Grav. *Micr.* 85. 25. — *Latr. Hist. nat. Crust. et Ins. IX.* 385. 25. — *Gyll. Ins. Suec. II.* 408. 29. — *Sahlb. Ins. Fenn. I.* 372. 49.

*Bolitochara nigrifula* Mannerh. *Brachelytr.* 77. 18.

Habitat in Europa.

Magnitudine valde variabilis, subdepressa, nigra, subnitida, tenuiter griseo-pubescent, vix sericans, thorace abdomineque lateribus parce nigro-pilosellis. Antennae capitis thoracisque fere longitudine, validiusculae, articulo secundo tertio paululum brevior, 5—10 longitudine paulo crassioribus, ultimo praecedentibus ambobus longitudine aequali vel paulo longiore, oblongo, subacuminato. Caput thorace paulo angustius, basi modice constrictum, rotundatum, subtiliter punctatum. Thorax coleopteris parum angustior, latitudine dimidio brevior, basi leviter rotundatus, lateribus postice subrectis, ante medium rotundatis, apice truncatus, angulis anterioribus deflexis, minus obtusis, subdepressus, confertim subtiliter punctatus, basi plerumque late et parum profunde impressus. Elytra thorace sesqui longiora, apice truncata, depressa, confertim subtiliter punctata, lutea testaceave, circa scutellum et angulo apicali exteriori plerumque infusca. Abdomen parallelum, subtus crebrius punctulatum, apice segmentorumque ventralium marginibus plerumque piceis vel rufo-piceis. Pedes testacei.

Ne temere species notis vacillantibus diagnosendas constituam timens, sub hoc nomine coniunxi species forte vere distinctas, quas saepe sedulo examinatas sufficientibus characteribus separare diu frustra conatus sum. Varietates praecipuas sequentes distinx:

a. Maior, H. graminicola vix parum minor, nitidior, antennis crassioribus nigris piceisve, palpis piceis, thorace lateribus fortius rotundato, abdomine dorso toto laevigato, vel segmentis anterioribus subtilissime punctatis. Hanc a Gyllenhal nom. *Al. sericantis*, *longicornis* et *socialis* Dom. Schüppel missam communicavit.

b. Praecedente parum minor, at angustior et magis elongata, thorace lateribus minus rotundato, antennis paulo gracilioribus, rufo-testaceis, apice



saepius fusciscentibus, palpis rufo-testaceis, abdomine segmentis tribus anterioribus parce subtiliterque punctatis.

c. Corporis habitu var. a., at duplo minor, antennis crassioribus, rufo-piceis, palpis rufo-testaceis, abdomine segmentis tribus primis parce subtiliterque punctatis. Haec apud nos frequentissime obvia.

d. Magnitudine praecedentis, at magis linearis depressaque, sub-opaca, abdomine toto parce subtiliterque punctato.

Differentia sexus in omnibus his varietatibus eadem: mas abdominis segmento sexto superiore margine apicali crenato, utrinque denticulo magis prominulo terminato, inferiore apice magis triangulariter producto distinctus.

Obs. Synonymia huius speciei valde intricata, et dubito ane saepius sub eodem nomine plures species confusae sint. *H. longicornem* pro *Al. sericante* saepius miserunt Entomologi Sueci. *Aleochara aterrima* Sahlb. Ins. Fenn. I. 368. 44. *Bolitochara aterrima* Mannerh. Brachelytr. 78. 21. certe e compluribus speciebus conflata: Prof. Sahlberg ipse sub hoc nomine *Hom. graminicolam* et *H. atramentariam* D. Prof. Germaro communicavit. Apud utrumque autorem citatum var. a. ad *H. graminicolam*, var. b. ad *H. atramentariam*, var. c. ad *H. socialem* c forte pertinet. *Bolitochara castanoptera* Mannerh. supra ad hanc speciem allegata, antennarum articulo ultimo magis elongato, ad sexum masculum spectare videtur.

\* 44. *H. sodalis*: *Nigra, nitida, antennis elytrisque castaneis, pedibus ferrugineis, thorace coleopteris angustiore, basi impresso, abdomine supra anterieus parce punctato, apice laevi.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 328. 22.*

Habitat in Germania.

*H. sociali* proxima et minoribus eius individuis (var. c.) aequalis, thorace angustiore nitido praecipue distincta. Corpus nigrum, nitidum, parce subtiliter pubescens. Antennae capitis thoracisque longitudine, crassiusculae, articulis penultimis leviter transversis, ultimo praecedentibus duobus longiore, sat magno, subovato, apice acuminato, rufo-brunneae, basi dilutiores. Palpi rufo-testacei. Caput thorace parum angustius, laeve. Thorax coleopteris dimidio angustior, latitudine dimidio brevior, basi leviter, lateribus praecipue ante medium fortius rotundatus, apice truncatus, angulis anterioribus rotundatis, leviter convexus, vage subtiliter punctatus, basi obsolete impressus, niger, nitidus. Elytra thorace sesqui longiora, dense subtiliter punctata, castanea, nitida. Abdomen supra segmentis tribus anterioribus fortius punctatis, subtus crebrius punctatum, segmentis ventralibus margine anoque piceis. Pedes rufo-testacei.

\* 45. *H. ochracea*: *Testacea, capite abdominisque cingulo fuscis, thorace antrorsum subangustato, convexiusculo, abdomine supra parce punctato.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 329. 24.*

Habitat in Germania, Gallia, Anglia.

Statura omnino praecedentis, testacea, pube depressa flavescenti-grisea subtiliter vestita, capite abdominisque segmentis duobus penultimis fuscis. Antennae capitis thoracisque longitudine, apicem versus incrassatae,

articulo secundo tertioque subaequalibus, quarto parvo globoso, 5—10 fortiter transversis, crassis, longitudine duplo crassioribus, ultimo praecedentibus ambobus vix aequali, conico, fuscae, basi testaceae. Caput thorace angustius, subtiliter punctatum. Thorax basi coleopteris paulo angustior, latitudine dimidio brevior, antrorsum subangustatus, basi rotundatus, utrinque subsinuatus, angulis posterioribus prope rectis, lateribus et angulis anterioribus rotundatis, apice truncatus, leviter, convexus, subtiliter dense punctatus. Elytra thorace sesqui longiora, apice ad angulum exteriorem subsinuata, subtiliter confertim punctata. Abdomen supra distincte, vage, segmentis posterioribus vix parcius quam anterioribus, punctatum.

- ✕ \*46. *H. trimaculata*: Nigra, antennarum basi apiceque, pedibus coleopterisque testaceis, his fusco-trimaculatis, thorace basi foveolato, lateribus piceo, abdomine basi rufo-piceo, supra fortius punctato. — Long  $1\frac{1}{4}$  lin.

Habitat in Pensylvania, Dom. Zimmermann.

Praecedentibus maior, *H. brunneae* fere aequalis, ad minus depressa, statura omnino Bolitocharae lunulatae. Corpus nigrum, nitidulum, subtiliter flavescenti-pubescent. Antennae capitis thoracisque longitudine, crassiusculae, articulis 5—10 transversis, ultimo praecedentibus ambobus prope aequali, breviter ovato, apice acuminato, nigro-fuscae, basi apiceque testaceae. Caput thorace angustius, parcius punctatum, ore testaceo. Thorax coleopteris angustior, longitudine plus dimidio latior, basi fortius, lateribus leviter rotundatus, apice truncatus, convexusculus, crebre profundius punctatus, basi transversim impressus, lateribus piceus. Coleoptera thorace sesqui longiora, fortius crebre punctata, dilute testacea, maculis tribus rotundatis fuscis, prima communi circa scutellum, reliquis duabus in angulo utroque exteriore. Abdomen segmentis 4 primis rufo-piceis, basi obscurioribus, reliquis nigro-piceis, supra crebrius fortiusque punctatum. Pedes testacei.

- \*47. *H. lunata*: Nigra, antennarum basi, pedibus, elytrorum fascia lunata basali, abdominis basi fasciaque ante apicem pallide testaceis, thorace coleopteris angustiore, transverso, subaeneo. — Long.  $1\frac{1}{2}$  lin.

Habitat in Brasilia, Dom. Thorey.

Antennae capite thoraceque longiores, testaceae, apice nigrae, articulis tribus primis elongatis, quarto quintoque crassitie duplo longioribus, teretibus, 6—10 subobconicis, sensim paulo crassioribus, ultimo elongato, acuminato. Caput thoracis latitudine, suborbiculatum, basi modice constrictum, nigrum, opacum, ore piceo, palpis testaceis, fronte depressa, sparsim obsolete punctata. Thorax coleopteris angustior, latitudine dimidio brevior, lateribus, basi et angulis posterioribus rotundatis, minus convexus, subtiliter punctatus, puncto maiore in medio baseos impressus, nigro-subaeneus, subnitidus, margine parce pilosellus. Elytra thorace paulo longiora, dense subtiliter punctata, nigra, fascia sublunata baseos, ab humero suturae medium petente, pallide rufescenti-flava. Abdomen apice subangustatum, segmentis 3 primis dilute rufo-testaceis, secundo tertioque supra macula oblonga media nigra notatis, reliquis nigris, sexto fascia apicali testacea. Pectus nigrum. Pedes pallide flavi.

- \* 48. *H. varia*: *Nigra, ore, antennis, pedibus, thorace, elytrorum macula humerali abdomineque testaceis, hoc maculis dorsalibus fasciaque ante apicem nigris.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Brasilia.

Statura omnino praecedentis. Antennae capite thoraceque paulo longiores, graciliores, apicem versus parum crassiores, articulis 4—10 subobconicis, crassitie haud longioribus, ultimo elongato, cylindrico, testaceae, apice subfuscescentes. Caput thoracis prope latitudine, laevissimum, inter oculos obsolete bifoveolatum, subaeneo-nigrum, pernitidum, ore testaceo. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus subrotundatus, basin versus subangustatus, parum convexus, aequalis, laevissimus, testaceus, nitidus, apice fuscescens. Elytra thorace sesqui fere longiora, confertim subtiliter punctata, nitida, aeneo-nigra, basi magis picea, fascia basali, ad suturam obsoleta, testacea. Abdomen supra laeve, subtus parce punctulatum parceque pubescens, nitidum, dilute testaceum, segmento secundo tertioque macula dorsali media nigra, quarto nigro, basi testaceo, quinto toto nigro. Pectus nigrum. Pedes testacei.

- \* 49. *H. picta*: *Testacea, elytrorum macula magna angulari, pectoris lateribus, abdominis punctis dorsalibus fasciaque subtus interrupta ante apicem nigris.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Brasilia, Dom. Prof. Reich.

Habitu omnino praecedentium, at paululum minor, capite, thorace elytrisque subtilissime confertim punctatis, parum nitidis. Antennae capitis thoracisque longitudine, graciliores, apicem versus leviter incrassatae, articulis 4—9 subobconicis, crassitie paulo longioribus, penultimo transverso, ultimo maiusculo, oblongo-conico, testaceae, articulo apicali fuscescente. Caput thoracis prope latitudine, testaceum. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus anterieus leviter rotundatis, posterius subsinuatis, basi leviter rotundatus, parum convexus, aequalis, testaceus. Scutellum testaceum. Elytra pube subtilissima fulva sericantia, nigra, opaca, basi suturaque late testaceis. Abdomen parcius subtiliter punctatum, supra nitidum, segmentis 2—4 dorso in medio baseos puncto nigro, quinto supra toto, subtus lateribus nigro. Pectus piceo-testaceum, lateribus nigrum. Pedes flavi.

- \* 50. *H. marcida*: *Obscure testacea, capite abdominisque cingulo postico piceis, pedibus pallidis, thorace transverso, subdepresso, basi obsolete impresso, abdomine supra parce subtiliterque punctato.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 328. 23.*

Habitat in Germania.

Statura *H. socialis*. Antennae capite thoraceque longiores, praecipue in mare, parum incrassatae, articulis 5—10 longitudine haud crassioribus, ultimo elongato-ovato, pilosellae, piceae, basi pallide testaceae. Corpus obscure testaceum. Caput thorace paulo angustius, rotundatum, basi leviter constrictum, piceum, nitidum, obsolete punctulatum. Thorax coleopteris dimidio prope angustior, latitudine dimidio brevior, basi, lateribus ante medium et



angulis omnibus rotundatis, apice truncatus, supra subdepressus, dense subtiliterque punctulatus, basi medio obsolete impressus. Elytra dense subtiliterque punctata, sordide testacea, apicem versus fusciscentia. Abdomen supra laevigatum vel parce subtiliter obsoleteque punctatum, nitidum, ante apicem nigrum. Pedes pallide flavi.

Mas antennis longioribus differt.

Obs. Antennis longioribus gracilioribusque et elytris longioribus ab *H. sericante*, abdomine supra laevigato praeterea a sequentibus discedit.

\*51. *H. spelaea*: Nigra, parum nitida, antennarum basi elytris pedibusque testaceis, thorace basi foveolato, abdomine supra crebre punctato, basi brunneo. — Long.  $1\frac{1}{2}$  lin.

Habitat in specu Adelsbergensi Carinthiae, Dom. Schmidt.

Statura omnino *H. elongatulae* et maioribus eius individuis aequalis, nigra, parum nitida, tenuissime parce pubescens. Antennae capitis thoracisque longitudine, apicem versus subincrassatae, articulo tertio secundo dimidio longiore, 5—10 longitudine vix crassioribus, ultimo praecedentibus ambobus longitudine subaequali, subovato, apice acuminato, piceae, basi testaceae. Palpi testacei. Caput thorace angustius, rotundatum, subtilissime punctulatum, fronte medio obsolete longitudinaliter impressa. Thorax coleopteris paulo angustior, latitudine dimidio brevior, basi et lateribus parum rotundatus, apice truncatus, angulis anterioribus obtusis, deflexis, perparum convexus, medio longitudinaliter depressus, basi foveola gemina impressus, confertim subtiliter punctatus, piceus, medio nigricans. Elytra thorace parum longiora, fusco-testacea, humeris dilutiora, confertim punctata. Abdomen anterieus rufo-brunneum, posterius nigrum, supra confertim, apice minus crebre punctatum. Pedes testacei.

Obs. *H. elongatulae* proximâ, at thorace brevior etc. satis distincta, ab *H. sociali* statura magis lineari depressaque elytrisque brevioribus praecipue discedens.

\*52. *H. dichroa*: Nigra, subnitida, antennis gracilioribus basi, elytris, pedibus anoque luteis, thorace subtiliter canaliculato, basi obsolete foveolato, abdomine supra confertim punctato. — Long.  $1\frac{1}{2}$  lin.

*Al. dichroa* Grav. Micr. I. 186. 1. Mon. 158. 25.

Habitat in America septentrionali.

Statura omnino *H. elongatulae*. Corpus nigrum, subnitidum, pube tenuissima subsericeum, lateribus parce pilosellum. Antennae capite thoraceque fere longiores, graciliores, apicem versus subincrassatae, articulo tertio secundo paulo longiore, 5—10 subaequalibus, longitudine haud crassioribus, ultimo praecedentibus duobus aequali, apice subacuminato, nigrae, basi testaceae. Palpi testacei. Caput thorace paulo angustius, suborbiculatum, basi constrictum, laeve, fronte subdepressa. Thorax coleopteris non angustior, latitudine plus dimidio brevior, basi et lateribus leviter rotundatus, apice leviter emarginatus, angulis anterioribus acutiusculis, subdepressus, subtiliter canaliculatus, basi foveola obsolete impressus, subtilissime punctatus, nigropiceus, subopacus. Elytra thorace paulo longiora, subtiliter punctata, opaca, lutea. Abdomen supra nitidulum, confertim sat distincte punctatum, apice luteo. Pedes lutei.

- \* 53. *H. propinqua*: *Linearis, subdepressa, nigra, elytris anoque fusco-testaceis, antennarum basi pedibusque testaceis, thorace transversim subquadrato, basi foveolato, abdomine supra densius punctulato.* — *Long.*  $1\frac{1}{2}$  lin.

Habitat in Americae insulis St. Thomae et St. Johannis, Dom. Moritz.

*H. elongutalae* similis, thorace brevior autem diffinis, subdepressa nigra, subtilissime punctata, tenuissime pubescens. Antennae capitis thoracisque longitudine, apicem versus leviter incrassatae, articulo tertio secundo paulo brevior, penultimis leviter transversis, ultimo praecedentibus ambobus longiore, apice acuminato, nigrae, basi testaceae. Caput thorace paulo angustius, rotundatum, confertim subtiliter punctatum, ore testaceo. Thorax coleopteris latitudine vix aequalis, hac dimidio brevior, basi et lateribus leviter rotundatus, apice truncatus, depressus, basi fovea oblonga impressus, nigro-piceus. Elytra thorace paulo longiora, fusco-testacea, circa scutellum infuscata. Abdomen nigrum, segmentorum anteriorum marginibus apiceque obscure testaceis, supra totum subtiliter crebrius punctatum. Pedes testacei.

Variat saepius corpore toto ferrugineo.

- \* 54. *H. conformis*: *Linearis, subdepressa, nigra, elytris fuscis, antennarum basi apiceque pedibusque testaceis, thorace transversim subquadrato, basi foveolato, abdomine supra densius punctulato.* — *Long.*  $1\frac{1}{2}$  lin.

Habitat prope Caracas Columbiae, Dom. Moritz.

Statura omnino et summa affinitas praecedentis. Antennae capitis thoracisque longitudine, apicem versus leviter incrassatae, piceae, articulo primo ultimoque laete testaceis, hoc longiore, acuminato. Palpi piceo-testacei. Caput thorace parum angustius, nigrum, subnitidum, omnium subtilissime punctatum, subtilissime pubescens. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus subrectis, subdepressus, omnium subtilissime pubescens, basi foveola transversa impressus, nigro-piceus, minus nitidus. Elytra thorace paulo longiora, confertim subtiliter punctulata, subtiliter griseo-pubescentia, opaca, fusca, angulo humerali rufescenti-testaceo. Abdomen supra totum subtiliter crebrius punctulatum, parcius tenuiter pubescens, nigrum, nitidum, basi saturate piceum, subtus piceum, segmentis singulis apice dilutioribus. Pectus nigrum. Pedes flavi.

- \* 55. *H. lurida*: *Nigra, subnitida, antennarum articulo primo elytrisque fusco-testaceis, pedibus pallidis, thorace transversim quadrato, aequali, abdomine supra crebrius punctulato.* — *Long.*  $1\frac{1}{4}$  lin.

Habitat in Brasilia, Dom. Prof. Reich.

Habitu omnino praecedentium, at paululum minor, nigra, subnitida, subtilissime pubescens, abdomine apice pilosello. Antennae capite thoraceque paulo longiores, apicem versus subincrassatae, articulis penultimis leviter transversis, ultimo praecedentibus ambobus paulo longiore, apice acuminato, nigrae, articulo primo obscure testaceo. Caput thorace paulo angustius, suborbiculatum, subtilissime obsolete punctulatum. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus subrectis, apice truncatus, parum convexus, aequalis, subtilissime obsolete punctulatus. Elytra

thorace paulo longiora, depressa, confertissime subtiliter punctulata, fuscescenti-testacea, circa scutellum infuscata. Abdomen supra nitidulum, crebrius subtiliter punctulatum. Pedes pallide flavescenti-testacei.

- \* 56. *H. semilutea*: Fortiter punctata, testacea, elytris dilutioribus, pectore abdomineque nigris, hoc basi obscure testaceo, thorace leviter convexo, aequali. — Long.  $1\frac{2}{3}$  lin.

Habitat in Brasilia, Dom. Prof. Germar.

Statura linearis, subdepressa *H. elongatae* et maioribus eius individuis aequalis. Antennae articulis duobus primis flavo-testaceis (reliqui desunt). Caput thorace paulo angustius, basi parum constrictum, fronte depressiuscula, sat crebre fortiterque punctatum, rufo-testaceum, nitidum, ore dilutiore. Thorax coleopteris parum angustior, latitudine dimidio brevior, basi modice, lateribus vix rotundatus, apice truncatus, leviter transversim convexus, aequalis, crebre fortiter grossiusque punctatus, testaceus, nitidus. Scutellum nigrum. Elytra thorace sesqui longiora, subdepressa, crebre fortiter profundeque punctata, flava, nitida, parce pubescentia. Abdomen sat elongatum, lineare, crebrius minus subtiliter punctatum, pube minus subtili fulvescenti-grisea vestitum, nigrum, nitidum, supra segmentis duobus primis piceo-testaceis, margine laterali testaceo, sexto margine posteriore piceo-testaceo, subtus segmentis omnibus margine posteriore, ultimo apice latiore, piceo-testaceo. Pectus nigrum. Pedes flavi.

- \* 57. *H. laesicollis*: Picea, opaca, capite abdominisque segmentis 3 et 4 nigris, thorace basi foveola magna longitudinali impresso, abdomine supra confertim, segmentis duobus ultimis parce punctato. — Long.  $1\frac{1}{3}$  lin.

Habitat in Brasilia.

Statura fere praecedentium. Obscure rufo-picea, opaca, pube tenui vestita. Antennae capitis thoracisque longitudine, crassiusculae, articulis penultimis transversis, ultimo praecedentibus tribus conjunctis longitudine aequali, oblongo, apice acuminato, nigrae, basi obscure testaceae. Caput thoracis latitudine, crebre subtiliter punctatum, piceo-nigrum, ore obscure testaceo, fronte subdepressa. Thorax coleopteris angustior, latitudine plus dimidio brevior, lateribus et basi perparum rotundatus, apice truncatus, subdepressus, subtilissime confertissimeque punctatus, fovea lata, a basi ultra medium adscendente excavatus. Elytra thorace dimidio longiora, subtilissime punctata. Abdomen supra segmentis 5 primis confertim, ultimis parce subtilius punctatis, tertio quartoque nigris, sexto rufo-testaceo. Pedes testacei.

- \* 58. *H. incana*: Elongata, subdepressa, nigra, opaca, dense cinereo-pubescentis, antennis basi tarsisque rufis, thorace coleopteris angustiore, medio depresso, abdomine supra segmentis ultimis subtiliter granulatis. — Long.  $1\frac{1}{2}$  —  $1\frac{2}{3}$  lin.

Er. Col. March. I. 329. 25.

Habitat in Germania septentrionali.

*H. sociali* paulo magis elongata, depressaque, nigra, opaca, pube tenui



cinerea dense vestita. Antennae breviusculae, apicem versus leviter incrassatae, articulis penultimis vix transversis, ultimo ovato, subacuminato, 5 basalibus pallide rufis, reliquis nigris. Caput thoracis prope latitudine, subtiliter parce punctatum, fronte depressa, medio obsolete foveolata, palpis piceis. Thorax coleopteris dimidio angustior, latitudine dimidio brevior, lateribus leviter rotundatus, parum convexus, dorso longitudinaliter depressus. Elytra thorace sesqui longiora, dense subtiliter punctata. Abdomen supra parcius punctatum, subnitidum, segmentis 3 ultimis subtiliter granulatis. Pedes nigri, trochanteribus, geniculis tarsisque rufescentibus.

Variat thoracis dorso longitudinaliter impresso.

\* 59. *H. excavata*: Nigra, subnitida, elytris fuscis, pedibus testaceis, femoribus fusciscentibus, thorace fovea magna media excavato, abdomine supra parce subtiliterque punctato. — Long. 1 lin.

*Aleochara excavata* Gyll. Ins. Suec. IV. 490. 30—31. — Sahlb. Ins. Fenn. I. 374. 52. *Boletochara excavata* Mannerh. Brachelytr. 78. 22. — Boisd. et Lacord. Faun. Ent. Paris. I. 547. 12.

Habitat in Suecia, Dom. Schüppel, in Germania meridionali, Dom. Dr. Walzl.

Habitu fere *H. elongatulae*, at *H. anali* vix maior, nigra, subnitida, confertim subtilissime punctata, subtilissime griseo-pubescentis. Antennae capite thoraceque paulo longiores, apicem versus sensim leviter incrassatae, articulo secundo tertioque subaequalibus, 4—10 sensim paulo crassioribus, penultimis crassitie paulo brevioribus, ultimo praecedente sesqui longiore, ovato, subacuminato, totae nigrae. Caput thorace paulo angustius, suborbiculatum, fronte medio impressa. Thorax coleopteris paulo angustior, latitudine dimidio brevior, basin versus subangustatus, basi rotundatus, medio fovea magna sat profunde impressus, margine laterali parce pilosello. Elytra thorace sesqui longiora, iam obscurius iam dilutius fusca. Abdomen lineare, supra parce subtiliterque punctatum, segmentis quarto quintoque sublaevigatis, apice fuscum. Pedes testacei, femoribus fusciscentibus.

Obs. Variat fovea thoracis in duas divisa, altera maiore rotundata in medio disci, altera minuta in medio baseos, sec. Gyll.

60. *H. deplanata*: Depressa, thorace brevissimo, dorso longitudinaliter late subimpresso, fronte late leviter excavata, abdomine supra laevigato. — Long. 1 lin.

*Aleochara deplanata* Grav. Micr. 100. 50. Mon. 177. 76. — Latr. Hist. nat. Crust. et Ins. IX. 391. 50.

Habitat in Germania, Mus. Univ. Wratislav.

*H. anali* vix longior, at minus angusta, fortiter depressa. Antennae capite vix sesqui longiores, articulo secundo crassiusculo tertioque hoc plus duplo minore obconicis, quarto transverso, sequentibus paulo angustiore, 5—10 inter se aequalibus, fortiter transversis, crassitie triplo brevioribus, fere perfoliatis, ultimo subgloboso, apice acuminato. Caput thorace paulo angustius, rotundatum, parce obsoleteque punctatum, fronte late leviter excavata, excavatione laevigata. Thorax coleopteris vix angustior, latitudine plus duplo brevior, basi et angulis posterioribus rotundatus, lateribus sub-

rectis, depressus, dorso late obsolete impressus, parce obsoleteque punctulatus, tenuissime griseo-pubescens. Elytra thorace duplo prope longiora, parce subtiliterque punctulata, subtiliter pubescentia. Abdomen lineare, supra laevigatum.

Obs. Specimen unicum, examinandum mihi benevole a Dom. Prof. Gravenhorst communicatum, minus maturum mihi videtur. Sunt ei caput, thorax, pectus et abdomen rufo-testacea, nitidula, os antennae et elytra testacea, pedes pallide flavi, abdomen ante apicem nigrum. Antennarum brevium crassarumque structura, thorace brevissimo, elytris parce punctatis minuta haec species facile cognoscatur.

\*61. *H. viduata*: Nigra, parum nitida, antennis pedibusque rufo-piceis, thorace transversim suborbiculato, convexiusculo, basi obsolete foveolato, abdomine supra confertissime punctulato. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 330. 26.

Habitat Berolini.

Statura fere *H. longicornis*, at minus depressa, nigra, perparum nitida, pube subtilissima subsericea vestita. Antennae capite thoraceque paulo longiores, apicem versus leviter incrassatae, articulis 4—10 subcylindricis, penultimis crassitie haud longioribus, ultimo oblongo-ovato, apice acuminato, rufo-piceae, basi dilutiores. Caput thorace angustius, suborbiculatum, basi modice constrictum, subtilissime parce punctatum, palpis rufo-piceis. Thorax coleopteris vix angustior, latitudine dimidio brevior, lateribus, basi angulisque omnibus rotundatis, leviter convexus, subnitidus, subtilissime dense punctatus, basi obsolete foveolatus. Elytra thorace paulo longiora, subtilissime dense punctata, opaca. Abdomen supra anterieus confertissime, posterius confertim at minus subtiliter punctatum, apice piceum. Pedes rufo-picei.

\*62. *H. atramentaria*: Nigro-subaenea, subnitida, antennis basi palpisque nigris, thorace coleopteris angustiore, basi foveolato, abdomine nigririmo, nitido, supra parcius punctato. — Long.  $1\frac{1}{2}$  lin.

*Aleochara atramentaria* Gyll. Ins. Suec. II. 408. 30. — Sahlb. Ins. Fenn. I. 373. 51. — Zetterst. Faun. Lappon. I. 100. 12. Ins. Lapp. 77. 15.

*Bolitochara atramentaria* Mannerh. Brachelytr. 78. 20. — Boisd. et Lacord. Faun. Ent. Paris. I. 546. 11.

*Aleochara aenescens* Zetterst. Ins. Lappon. 77. 16.

Habitat in Suecia, Germania septentrionali, Anglia, Sardinia.

Statura fere *H. socialis*. Corpus nigrum, nitidum, capite, thorace elytrisque subaeneis, modice nitidulis, tenuissime pubescens. Antennae vix capitis thoracisque longitudine, tenuiores, apicem versus parum incrassatae, articulis penultimis crassitie vix longioribus, ultimo praecedentibus ambobus prope brevior, apice acuminato, piceae, basi nigrae. Palpi nigri. Caput thorace paulo angustius, obsolete parce punctatum, fronte subdepressa, palpis nigris. Thorax coleopteris dimidio angustior, latitudine dimidio brevior, basi et lateribus leviter rotundatus, apice truncatus, parum convexus, basi foveolatus, crebre subtiliter punctatus. Elytra thorace sesqui longiora, subtilius punctata. Abdomen nigrum, nitidum, supra parce punctatum, lateribus pilosellum. Pedes picei.

Obs. I. Insectum forte maritimum, mihi in littore maris Baltici sub fucis lectum. Dubito an revera Lutetiae lectum sit, commutatis et commixtis enim specie-

bus similibus autores Faunae Ent. Parisiensis saepius specimina collectionis Dejeaniana extranea descripsisse mihi videntur, et in collectione Dom. Aubé, qui Coleoptera Parisiensia diligentissime collegit, hanc speciem non inveni.

2. *Al. aenescens* Zett „simillima atramentariae, sed differt thorace elytris-que subaenescens, glaberrimis, pedibus nigris, tarsis tantum testaceis, antennis pubescentibus, abdomine pilis rarioribus obsito, foveola antescutellari parva obsoleta in thoracis postico impressa” non satis distinctam puto. Color subaeneus *H. atramentariae* peculiaris videtur.

\* 63. *H. festinans*: *Linearis, subdepressa, nigra, nitida, elytris fuscis, antennarum basi pedibusque testaceis, thorace basi obsolete canaliculato, abdomine supra segmentis anterioribus crebrius, posterioribus parcius punctatis.* — *Long.*  $1\frac{1}{2}$  lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Statura fere *H. elongatulae*, et minoribus eius individuis aequalis, nigra, nitida, tenuissime parce pubescens. Antennae capite thoraceque breviores, apicem versus vix incrassatae, articulis 4—10 subaequalibus, haud transversis, ultimo praecedentibus ambobus vix aequali, apice acuminato, piceae, articulis primis testaceis. Caput thoracis latitudine, fronte subconvexa, parce subtiliter punctata. Thorax coleopteris vix angustior, longitudine dimidio latior, basi et lateribus rotundatus, apice truncatus, transversim convexus, parce subtiliter punctatus, canalicula parum profunda, brevi, vix medium superante impressus. Elytra thorace paulo longiora, picea, parce punctata. Abdomen nitidissimum, supra segmentis anterioribus densius, posterioribus vage punctatum. Pedes pallide testacei.

\* 64. *H. vestigialis*: *Linearis, subdepressa, nigra, subnitida, elytris fuscis, pedibus testaceis, thorace transverso, aequali, abdomine supra parce punctato.* — *Long.*  $1\frac{1}{3}$  lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Praecedenti affinis, differt autem corpore minus nitido, antennis totis nigris, elytris densius, abdomine aequaliter parce punctatis. Corpus nigrum, subnitidum, pube tenui densius vestitum. Antennae capitis thoracisque longitudine, apicem versus sensim incrassatae, articulis 5—10 transversis, longitudine duplo crassioribus, ultimo oblongo, praecedentibus ambobus paulo longiore, apice acuminato, totae nigrae. Palpi picei. Caput thorace vix angustius, fronte subdepressa, parce subtiliter punctata. Thorax coleopteris latitudine aequalis, lateribus subrectis, apice truncatus, subdepressus, subtiliter parcius punctatus, basi haud impressus. Elytra thorace sesqui longiora, densius punctata, fusca. Abdomen supra aequaliter subtilissime parce punctatum, segmentorum anteriorum margine apiceque piceis. Pedes testacei.

\* 65. *H. gemina*: *Linearis, subdepressa, nigra, elytris fuscis, antennarum basi, pedibus anoque testaceis, thorace transverso, basi obsolete bifoveolato.* — *Long.* 1 lin.

*Er. Col. March. I.* 330. 27.

Habitat in Germania.

Statura *H. analis* et maximis eius speciminibus vix maior, nigra, subnitida, subtilissime pubescens, dense subtiliterque punctata. Antennae capi-



tis thoracisque longitudine, apicem versus subincrassatae, articulis 4 — 10 longitudine parum crassioribus, ultimo oblongo-ovato, apice acuminato, fuscae, basi testaceae. Caput rotundatum, thorace angustius, ore testaceo, fronte medio canalicula brevi obsolete impressa. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus et basi rotundatus, antice leviter bisinuatus, angulis anterioribus rotundatis, posterioribus obtusis, parum convexus, basi punctis duobus maioribus approximatis, magis minusve distinctis impressus. Elytra thorace sesqui prope longiora, apice medio coniuncta leviter emarginata, singula ad angulum anteriorem subsinuata, fusca, thorace parum fortius punctata. Abdomen nigrum, apice testaceum, supra antierius confertim, posterius parcius punctatum. Pedes testacei.

Obs. *H. elongatae* perquam affinis est haecce species et dubito, an rite sit distincta, differre mihi autem videtur statura minore thoraceque paulo brevior.

\* 66. *H. autumnalis*: *Linearis, nigra, nitidula, antennarum basi pedibusque testaceis, elytris fuscis, thorace postice canaliculato foveolatoque, abdomine supra creberrime punctato.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Germania meridionali, Dom. Dr. Walzl.

Statura lineari subdepressa fere *H. elongatae* et minoribus eius individuis aequalis, nigra, nitidula, subtiliter pubescens. Antennae capite thoraceque breviores, apicem versus sensim leviter incrassatae, articulis 7—10 leviter transversis, ultimo maiusculo, breviter ovato, subacuminato, fuscae, basi testaceae. Palpi testacei. Caput thorace angustius, rotundatum, subtiliter punctulatum, fronte obsolete impressa. Thorax coleopteris dimidio angustior, latitudine plus dimidio brevior, basi fortius, lateribus leviter rotundatus, apice truncatus, angulis anterioribus rotundatis, posterioribus obtusis, parum convexus, subtiliter punctulatus, basi foveola maiore antrorsum in canaliculam elongata impressus. Elytra thorace sesqui longiora, confertim subtiliter punctata, nigro-fusca. Abdomen parallelum, supra minus subtiliter sat crebre punctatum, segmentis ultimis duobus parcius punctatis, punctis autem exasperatis. Pedes testacei.

\* 67. *H. oblita*: *Linearis, subdepressa, nigra, parum nitida, pedibus testaceis, elytris fuscis, thorace canaliculato, abdomine supra antierius subtiliter punctulato.* — Long.  $1\frac{1}{4}$  lin.

Habitat Lutetiae, Dom. Dr. Aubé, in Germania meridionali, Dom. Dr. Walzl.

Habitu *H. socialis*, at minor, *H. clanculae* aequalis, nigra, opaca, tenuiter pubescens. Antennae capite thoraceque paulo breviores, extrorsum leviter incrassatae, articulis 5 — 10 transversis, ultimo praecedentibus duobus aequali, apice acuminato, fusco-nigrae. Palpi picei. Caput thorace vix angustius, subtilissime obsoleteque punctatum, fronte depressa. Thorax coleopteris latitudine subaequalis, longitudine dimidio lator, basi fortius, lateribus leviter rotundatus, apice truncatus, angulis omnibus obtusis, supra subdepressus, subtiliter punctatus, canalicula plerumque antrorsum abbreviata impressus. Elytra thorace paulo longiora, crebre punctata, fusca. Abdomen nigrum, nitidum, supra punctulatum, apice laevigatum. Pedes testacei.

Obs. *H. clanculae* proxima, antennarum autem structura abdomineque parallelo facile distinguenda.

- \* 68. *H. clancula*: *Brevis, nigra, subnitida, elytris fuscis, antennarum basi pedibusque testaceis, thorace transverso, leviter convexo, basi obsolete canaliculato, abdomine supra aequaliter punctato, margine laterali subrotundato.* — Long. 1 lin.

*Er. Col. March. I.* 331. 28.

Habitat Berolini.

*Brevis*, crassiuscula, nigra, subnitida, confertissime subtiliter punctata, tenuissime griseo-pubescens. Antennae capitis thoracisque fere longitudine, apicem versus subincrassatae, articulis 4 — 10 subturbinatis, crassitie vix brevioribus, ultimo ovali, fuscae, basi testaceae. Palpi testacei. Caput thorace parum angustius, basi leviter tantum constrictum, fronte convexiuscula. Thorax coleopteris vix angustius, latitudine duplo brevior, lateribus subrectis, angulis anterioribus fortiter rotundatis, leviter convexus, basi obsolete canaliculatus. Elytra thorace sesqui longiora, apice coniunctim emarginata, ad marginem distincte sinuata, confertim evidentius punctata, fusca. Abdomen supra aequaliter densius punctatum, pube longiore vestitum, margine laterali lato, elevato, rotundato. Pedes testacei.

- \* 69. *H. luteola*: *Brevis, flavo-testacea, abdomine postice nigricante, thorace transverso, basi obsolete foveolato, abdomine supra confertissime punctato.* — Long. 1 lin.

Habitat in America septentrionali, Dom. Zimmermann.

Parva, brevis, statura et habitu fere praecedentis, flavo-testacea, tenuiter pubescens. Antennae capite thoraceque breviores, extrorsum parum incrassatae, articulis penultimis subtransversis, ultimo oblongo, apice acuminato. Caput thorace angustius, basi vix constrictum, crebre profundius punctatum, interdum nigricans, nitidum. Thorax coleopteris latitudine aequalis, longitudine duplo latior, basi et lateribus leviter, angulis anterioribus fortiter rotundatis, posterioribus obtusis, subdepressus, confertim subtiliter punctatus, basi foveola obsoleta transversa notatus. Elytra thorace paulo longiora, confertim evidentius punctata. Abdomen segmentis anterioribus duobus exceptis nigro-piceum, supra nitidum, crebre minus subtiliter punctatum. Pectus lateribus piceum. Pedes et elytra magis flavescunt.

Mas antennis paulo longioribus, articulo ultimo magis ovato, obtusiore distinctus.

- ✕ \* 70. *H. analis*: *Linearis, subdepressa, nigra, thorace elytris piceis, antennis, pedibus anoque testaceis, thorace suborbiculato, leviter convexo, basi foveola transversa impresso.* — Long.  $\frac{3}{4}$  — 1 lin.

*Er. Col. March. I.* 332. 30.

*Aleochara analis* Grav. *Micr.* 76. 14. *Mon.* 153. 13. — *Latr. Hist. nat. Crust. et Ins.* IX. 382. 14.

*Bolitochara analis* Runde *Brachel. Hal.* 30. 4.

*Bolitochara evanescens* Mannerh. *Brachelytr.* 81. 37.

*Aleochara evanescens* Sahlb. *Ins. Fenn. I.* 364. 35.

*Bolitochara tages* Runde *Brachel.* 30. 3.

*Var.: Thorace elytrisque rufo-piceis.*

*Bolitochara bifoveolata* Mannerh. *Brachelytr.* 79. 23.

*Aleochara bifoveolata* Sahlb. *Ins. Fenn.* 1. 374. 53.

Habitat in Europa frequens: in America septentrionali legit Dom. Zimmerm.

Angustula, subdepressa, linearis, subtilissime griseo-pubescent. Antennae capite thoraceque breviores, apicem versus vix incrassatae, articulis 4—10 subturbinatis, longitudine parum crassioribus, ultimo maiusculo, ovato, apice subacuminato, testaceae. Caput thoracis fere latitudine, rotundatum, basi profunde constrictum, ut pedunculo quasi brevi thoraci insertum videatur, confertissime subtiliter punctatum, nigrum vel piceum, ore testaceo. Thorax coleopteris haud angustior, latitudine dimidio brevior, lateribus, basi et angulis omnibus rotundatus, supra leviter convexus, confertissime subtiliter punctatus, basi foveola transversa profunde impressus, piceus. Elytra thorace parum longiora, subtiliter dense punctata, picea. Abdomen subtiliter dense punctatum, nigrum, apice testaceum. Pedes testacei.

Variat thorace elytrisque piceo-testaceis.

\* 71. *H. exilis*: *Linearis, depressa, nigra, tenuiter sericeo-pubescent, elytris fuscis, antennarum basi, pedibus anoque testaceis, thorace transversim subquadrato, abdomine supra confertim punctulato.* — Long.  $\frac{3}{4}$  lin.

*Er. Col. March.* I. 333. 32.

*Al. analis* var. (*exilis* Kn.) *Grav. Mon.* 153. 13.

Habitat in Germania.

H. anali plerumque minor, angustior, depressa, nigro-picea, parum nitida, subtilissime punctata, pube tenui grisea vestita. Antennae capite vix longiores, apicem versus fortiter incrassatae, articulis 4—10 valde transversis, fere perfoliatis, ultimo maiusculo, breviter ovato, obtusiusculo, fuscae, basi testaceae, Caput thoracis latitudine, subquadratum, obsolete canaliculatum, antice leviter impressum, ore testaceo. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus subrectis, perparum convexus, medio longitudinaliter depressus. Elytra thorace paulo longiora, minus subtiliter punctata, fusca. Abdomen nigrum, apice testaceum, minus subtiliter confertim punctatum. Pedes testacei.

Variat saepe corpore testaceo.

\* 72. *H. palleola*: *Linearis, subdepressa, testacea, elytris fusco-testaceis, capite abdominisque segmentis penultimis piceis, thorace transversim suborbiculato, leviter convexo, abdomine supra parce punctato.* — Long.  $\frac{3}{4}$  lin.

*Er. Col. March.* I. 333. 31.

Habitat in Germania.

Statura H. analis et minoribus eius individuis magnitudine aequalis, testacea, parce subtiliter pubescens. Antennae capitis thoracisque longitudine, apicem versus sensim incrassatae, articulis 4—10 transversis, ultimo breviter ovato, subacuminato, pilosellae, fuscae, basi testaceae. Caput thorace parum angustius, rotundatum, basi leviter constrictum, laeve, nitidum,



piceum, ore testaceo. Thorax coleopteris dimidio angustior, latitudine dimidio brevior, lateribus, basi angulisque omnibus rotundatis, apice truncatus, leviter convexus, subtilissime vage punctatus. Elytra thorace plus sesqui longiora, subtiliter densius punctata, fusco-testacea. Abdomen testaceum, nitidum, supra antrorsum vage punctatum, apicem versus laeve, segmentis duobus antepenultimis nigris. Pectus piceum. Pedes flavi.

- \* 73. *H. inconspicua*: *Linearis, nigra, opaca, fortius punctulata, tenuiter sericeo-pubescentis, tarsis flavescentibus, thorace transverso, obsolete canaliculato*. — Long.  $\frac{3}{4}$  lin.

Habitat in Germania, Dom. Dr. Waltl, Lutetiae, Dom. Dr. Aubé.

Statura *H.* exilis, nigra, opaca, confertim fortius punctata, pube tenuissima vestita. Antennae capitis thoracisque longitudine, apicem versus subincrassatae, articulo ultimo ovato, corpori concolores. Caput thorace paulo angustius, basi leviter constrictum. Thorax coleopteris parum angustior, latitudine plus dimidio brevior, lateribus et angulis omnibus rotundatis, leviter convexus, obsoletissime canaliculatus. Elytra thorace duplo fere longiora. Abdomen supra nitidulum, anteriùs fortius, apicem versus obsoletius punctulatum. Pedes concolores, geniculis piceis, tarsis flavescentibus.

- \* 74. *H. inquinula*: *Linearis, subdepressa, nigra, tenuiter sericeo-pubescentis, elytris fuscis, pedibus obscure testaceis, thorace transversim subquadrato*. — Long.  $\frac{1}{2}$  lin.

*Er. Col. March. I.* 334. 33.

*Al. inquinula Grav. Micr.* 78. 16. *Mon.* 153. 15. — *Latr. Hist. nat. Crust. et. Ins.* IX. 382. 16.

Habitat in Germania.

Forma praecedentium, at minutissima, angusta, subdepressa, nigra, parum nitida, pube tenui grisea vestita. Antennae capitis thoracisque longitudine, apicem versus subincrassatae, articulis 4—10 leviter transversis, ultimo breviter ovato, subacuminato, nigricantes. Caput thoracis prope latitudine, subtilissime punctatum, fronte planiuscula, medio subimpressa. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus rectis, basi rotundatus, subdepressus, confertissime subtilissimeque punctatus. Elytra thorace sesqui longiora, dense subtiliter punctata, fusca. Abdomen parcius punctatum, apice piceum. Pedes fusco-testacei.

- \* 75. *H. didyma*: *Testacea, capite exserto, fronte antice transversim impressa, thorace transverso, antice rotundato, basi transversim impresso foveolatoque*. — Long. 1 lin.

Habitat in Columbia, Dom. Moritz.

Habitus omnino *Autaliae impressae*, at duplo fere minor. Corpus testaceum, nitidulum, subtilissime punctulatum, dense subtilius pubescens. Antennae capitis thoracisque longitudine, crassiusculae, apicem versus distincte crassiores, articulis 6—10 fortiter transversis, ultimo ovato, subacuminato. Caput thoracis fere latitudine, rotundatum, pedunculo angusto thoraci insertum, fronte convexiuscula, supra antennarum insertionem transversim leviter impressa. Thorax coleopteris tertia parte angustior, longitudine duplo fere

lterior, basi truncatus, apice cum angulis anterioribus rotundatus, supra leviter convexus, basi foveolis quatuor distantibus impressis, exterioribus obsoletioribus, interioribus sulculo transverso connexis. Abdomen segmentis dorsalibus depressis, subglabris, vix punctatis. Anus et pedes pallidiores. Caput interdum piceum.

Obs. Adeo haec species habitu Autaliam simulat, ut nonnisi ex oris structura examinata locum eius naturalem cognoverim.

\*76. *H. bisulcata*: *Obscure rufo-testacea, capite exserto, abdominis segmento quinto antennisque nigris, his basi testaceis, thorace transverso, dorso bisulcato.* — Long. 1 lin.

Habitat in Columbia, Dom. Moritz.

Statura praecedentis. Corpus obscure rufo-testaceum, totum confertim subtilissime punctulatum, pube tenuissima densius obductum, opaculum. Antennae capitis thoracisque longitudine, apicem versus parum incrassatae, articulis 6—10 longitudine paulo crassioribus, ultimo articulis tribus praecedentibus coniunctis vix brevior, subcylindrico, apice subacuminato, fuscae, basi testaceae. Caput pedunculo brevi thoraci insertum, rotundatum, piceum, nitidulum, supra antennis transversim subimpressum. Thorax coleopteris angustior, longitudine duplo latior, basi apiceque truncatus, lateribus rotundatus, angulis anticis obtusis, subrotundatis, supra leviter convexus, sulcis duobus dorsalibus latis, postice confluentibus, antice abbreviatis, et ante apicem lineola minuta longitudinali impressus. Abdomen supra subtiliter punctulatum, segmento quinto nigro-piceo, ano pallidiore. Pectus piceum. Pedes testacei.

\*77. *H. ustulata*: *Testacea, capite abdominisque segmento quarto quinque nigris, elytris fuscis, humeris pallidioribus, antennis nigricantibus, basi apiceque testaceis, thorace transverso, basi obsolete foveolato.* — Long. 1 lin.

Habitat in Columbia, Dom. Moritz.

Statura fere praecedentium. Corpus totum subtilissime pubescens. Antennae capitis thoracisque longitudine, apicem versus leviter incrassatae, articulis penultimis transversis, ultimo maiore, ovato, subacuminato, nigricantes, basi summoque apice testaceis. Caput thorace parum angustius, suborbiculare, subtiliter punctatum, nitidum, nigrum, ore piceo, palpis pallidis. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus et basi leviter rotundatus, apice truncatus, supra parum convexus, basi foveola parva obsoleta impressus, confertissime subtiliter punctulatus, obscure rufo-testaceus, parum nitidus. Elytra thorace parum longiora, subtilius confertissime punctata, opaca, dilute fusca, macula humerali testacea. Abdomen dorso parcius distinctiusque punctulatum, rufo-testaceum, segmentis quarto quinque nigris. Pectus nigrum. Pedes flavi.

\*78. *H. melanura*: *Rufo-testacea, capite, ano antennisque nigris, his basi testaceis, elytris flavis, thorace transverso, basi profunde foveolato.* — Long. 1 lin.

Habitat in Americae insulis St. Johannis et Puerto Rico, Dom. Moritz.

Statura omnino praecedentis, dense subtiliter punctata, tenuiter flavescenti-pubescent. Antennae capitis thoracisque longitudine, apicem versus incrassatae, articulis 5—10 longitudine paulo crassioribus, ultimo maiore, ovato, apice vix acuminato, basi testaceae. Caput thorace parum angustius, subnitidum, nigrum, ore rufescente. Thorax coleopteris parum angustior, latitudine duplo brevior, lateribus et basi parum rotundatus, subdepressus, basi foveola profunda impressus, rufo-testaceus. Elytra thorace sesqui longiora, obscure flava. Abdomen parcius subtiliter punctatum, segmentis quatuor anterioribus rufo-testaceis, reliquis nigris. Pectus testaceum, lateribus piceum. Pedes flavi.

\* 79. *H. flavipennis*: *Nigra, nitida, antennarum basi, elytris pedibusque flavis, thorace transverso, basi obsolete foveolato, abdomine supra parcius punctulato.* — Long. 1 lin.

Habitat in Americae ins. St. Johannis, Dom. Moritz.

Statura fere praecedentium, nigra, tenuissime pubescens, subtilissime confertissimeque punctata, parum nitida. Antennae capitis thoracisque longitudine, crassiusculae, articulis 5—10 fortius transversis, ultimo maiore, ovato, subacuminato, nigrae, basi testaceae, articulo primo flavo. Caput thoracis fere latitudine, fronte latiuscula, plana. Thorax coleopteris dimidio prope angustior, latitudine fere duplo brevior, lateribus ante medium rotundatis, basin versus subangustatis, angulis posterioribus minus obtusis, supra parum convexus, basi foveola transversa obsolete impressus. Elytra thorace paulo longiora, flava. Abdomen nitidulum, minus confertim punctatum, praecipue apicem versus. Pedes flavi.

\* 80. *H. obsoleta*: *Nigra, nitida, elytris fuscis, humeris, antennarum basi pedibusque testaceis, thorace transverso, convexiusculo, basi obsolete foveolato, abdomine supra parcius punctulato.* — Long. 1 lin.

Habitat in Columbia, Dom. Moritz.

Statura exacte praecedentis. Antennae capitis thoracisque longitudine, crassiusculae, articulis secundo tertioque obconicis, hoc minore, 4—10 sat fortiter transversis, ultimo maiore, ovato, subacuminato, nigrae, articulis tribus primis testaceis, primo dilutiore. Corpus nigrum, parum nitidulum, subtilissime punctulatum, pube subtilissima vestitum. Caput thoracis latitudine, basi parum constrictum, fronte leviter convexa. Thorax coleopteris parum angustior, latitudine duplo brevior, basi lateribusque parum rotundatus, apice truncatus. Elytra thorace paulo longiora, nigro-fusca, humeris transversim testaceis. Abdomen supra parcius punctulatum. Pedes testacei, femoribus basi piceis.

\* 81. *H. curticollis*: *Nigra, nitida, elytris fuscis, humeris apiceque, antennarum basi pedibusque obscure testaceis, thorace transverso, convexiusculo, basi profunde foveolato, abdomine supra crebrius punctulato.* — Long. 1 lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Statura et summa affinitas praecedentis. Nigra, nitidula, minus subtiliter griseo-pubescent. Antennae capitis thoracisque vix longitudine, ab ar-



ticulo quarto inde crassae, articulis 5—10 longitudine triplo fere crassioribus, ultimo maiore, ovali, apice obtuso, pilosellae, nigrae, basi testaceae. Caput thoracis latitudine, basi parum constrictum. Thorax coleopteris parum angustior, latitudine duplo prope brevior, lateribus parum rotundatus, supra convexiusculus, crebre subtiliter punctatus, basi foveola profunda transversa impressus. Elytra thorace paulo longiora, crebre profundius punctata, fusca, humeris rufescentibus, margine apicali flavescente. Abdomen supra crebrius punctulatum. Pedes obscure testacei.

Obs. A praecedente thorace profunde foveolato, elytris fortius punctatis, margineque eorum apicali pallido facile distinguenda.

\* 82. *H. decipula*: *Ferruginea, nitida, antennis medio nigricantibus, apice ferrugineis, thorace transverso, basin versus subangustato, basi foveolato, elytris fortius punctulatis, abdomine supra laevi.* — Long. 1 lin.

Habitat in Americae merid. ins. Puerto Rico, Dom. Moritz.

Habitus omnino Oxypodae cuiusdam sectionis II (prolixae), at huius generis, linearis, piceo-rufa, nitidula, tenuiter fulvescenti-pubescent. Antennae capite thoraceque paulo longiores, apicem versus sensim incrassatae, articulis penultimis transversis, ultimo maiore, ovato, dense longius pubescentes, basi rufo-piceae, medio fuscae, articulis ultimis duobus dilute testaceis. Caput thorace paulo angustius, basi leviter constrictum, confertim subtiliter punctatum, ore subattenuato. Thorax coleopteris vix angustior, latitudine dimidio brevior, basin versus subangustatus, lateribus anterioribus leviter rotundatis, posterius subsinuatis, supra leviter convexus, subtilissime punctulatus, basi foveola media transversa sat distincta impressus. Elytra thorace paulo longiora, sat crebre minus subtiliter punctulata. Abdomen lineare, dorso laeve, subtus parce obsoleteque punctulatum. Pedes testacei.

\* 83. *H. nigripennis*: *Rufo-testacea, nitida, elytris nigris, pectore antennisque nigricantibus, his basi apiceque testaceis.* — Long. 1 lin.

Habitat in Americae insula St. Johannis, Dom. Moritz.

Statura fere *H. analis*. Laete rufo-testacea, nitida, subtilissime pubescens. Antennae fere capitis thoracisque longitudine, apicem versus leviter incrassatae, articulis penultimis transversis, ultimo maiusculo, ovato, nigrae, basi rufo-testaceae, articulo ultimo dilute testaceo. Corpus rufo-testaceum. Caput thorace paulo angustius, basi leviter constrictum, subtiliter parce punctatum. Thorax coleopteris vix angustior, latitudine duplo brevior, lateribus et basi leviter rotundatus, apice late leviter emarginatus, subdepressus, subtiliter parcius punctatus. Elytra thorace haud longiora, nigerrima, distinctius confertim punctata. Abdomen supra subtiliter vage punctatum. Pectus piceum.

\* 84. *H. alternata*: *Nigra, elytris basi, abdomine et antennis basi apiceque pedibusque flavis.* — Long. 1 lin.

Habitat in Americae ins. St. Thomae, Dom. Moritz.

Statura praecedentis. Antennae capitis thoracisque fere longitudine, apicem versus leniter incrassatae, articulis penultimis transversis, ultimo ovato, subacuminato, nigrae, basi testaceae, articulo ultimo laete testaceo.

Thorax coleopteris vix angustior, latitudine dimidio brevior, lateribus et basi leviter rotundatus, parum convexus, laevis, niger. Elytra thorace parum longiora, subtilissime punctata, parte dimidia anteriore pallide flava, posteriore nigra. Abdomen laeve, laete testaceum vel flavum, segmentis quarto quintoque nigris. Pectus nigrum. Pedes flavi.

- \* 85. *H. seminigra*: *Linearis, dilute rufo-testacea, pectore abdomineque nigris, hoc basi testaceo, antennis nigris, basi flavis, thorace transverso, leviter convexo.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Habitu omnino praecedentis, at sesqui maior, subtilissime punctulata pubescensque. Antennae capite sesqui longiores, apicem versus leviter incrassatae, articulis 4 — 10 transversis, ultimo ovato, nigrae, articulis 4 primis pallide flavis. Caput thorace minus et angustius, testaceum. Thorax coleopteris parum angustior, latitudine dimidio brevior, basi leviter rotundatus, lateribus subrectis, leviter transversim convexus, aequalis, testaceus. Scutellum testaceum. Elytra thorace paulo longiora, flavo-testacea. Abdomen sat elongatum, supra laevigatum, subtus segmentis anterioribus subtiliter punctulatis, segmento primo testaceo, secundo piceo, reliquis nigerimis, nitidis, ano picescente. Pectus posterius nigrum. Pedes flavi.

- \* 86. *H. stigma*: *Linearis, flavo-testacea, abdomine segmento quarto macula magna dorsali atra, thorace transverso, leviter convexo, aequali.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Brasilia, Dom. Prof. Germar.

Statura omnino *H. seminigrae*, tota pallide testacea, elytris pectoreque magis flavescentibus, nitidula, tenuiter subtiliterque griseo-pubescens. Antennae capite sesqui longiores, apicem versus sensim incrassatae, articulis duobus primis maiusculis, subaequalibus, secundo tertioque, hoc dimidio minore, obconicis, 4 — 10 longitudine subaequalibus, sensim crassioribus et fortius transversis, penultimis longitudine duplo crassioribus, ultimo breviter ovato. Caput thorace paulo angustius, suborbiculatum, basi vix constrictum, fronte depressa, confertim minus subtiliter punctatum. Oculi nigri. Thorax coleopteris parum angustior, latitudine plus dimidio brevior, basi et lateribus modice rotundatus, apice truncatus, angulis omnibus obtusis, leviter transversim convexus, aequalis, crebre fortius punctatus. Elytra thorace sesqui longiora, subdepressa, sat crebre fortius punctata. Abdomen lineare, minus confertim subtilius punctatum, supra segmento quarto macula magna media aterrima, subtus immaculatum. Pedes pallidi.

- \* 87. *H. silacea*: *Testacea, antennis apice abdomineque ante apicem nigris, thorace transverso, aequali, abdomine dorso subtiliter punctulato.* — Long.  $\frac{4}{5}$  lin.

Habitat in America septentrionali, Dom. Zimmermann.

Statura omnino et summa affinitas proxime praecedentium, at paulo minor, flavo-testacea, parum nitida, subtilissime griseo-pubescens. Antennae capite parum longiores, apicem versus sat distincte sensim incrassatae, articulis penultimis transversis, ultimo subgloboso, testaceae, articulis ultimis 4

nigris. Caput thorace paulo angustius, basi leviter constrictum, subtilissime punctulatum, fronte planiuscula. Thorax coleopterorum prope latitudine, latitudine duplo brevior, basi lateribusque leviter rotundatus, confertim subtiliter punctatus, dilutius flavo-testaceus. Elytra thorace plus sesqui longiora, confertim subtiliter punctata, subfuscentia. Abdomen supra subtiliter punctulatum, segmentis antepenultimis duobus totis, penultimo basi nigris. Pedes pallide flavo-testacei.

- \*88. *H. cinctella*: Testacea, pectore abdominisque segmentis tribus penultimis nigris, elytris nigricantibus, basi testaceis, antennis apice nigris. — Long.  $\frac{3}{4}$  lin.

Habitat in Columbia, Dom. Moritz.

Statura omnino praecedentis, subtiliter punctulata, subtilissime pubescens. Antennae capite paulo longiores, apicem versus sensim incrassatae, articulis penultimis transversis, ultimo maiusculo, subglobozo, testaceae, apice fuscae. Caput thorace paulo angustius, testaceum. Thorax coleopteris parum angustior, latitudine duplo brevior, basi et lateribus aequaliter rotundatus, leviter transversim convexus, aequalis, dilute testaceus. Elytra thorace sesqui prope longiora, piceo-nigra, basi dilutiora, humeris testaceis. Abdomen supra subtiliter punctulatum, segmentis duobus primis testaceis, tercio, quarto quintoque nigris, hoc margine apicali sextoque toto iterum testaceis. Pectus posterius nigrum. Pedes pallide testacei.

- \*89. *H. atratula*: Linearis, atra, parum nitida, antennarum basi, tibiis tarsisque rufo-testaceis, thorace transverso, convexiusculo. — Long. 1 lin.

Habitat in Columbia, Dom. Moritz.

Statura H. analis. Corpus lineare, atrum, parum nitidum, vix pubescens. Antennae capite parum longiores, apicem versus subincrassatae, articulis penultimis transversis, ultimo ovato, subacuminato, pilosellae, nigrae, articulis 4 primis pallide rufo-testaceis. Caput profunde crebreque punctatum, rotundatum, thorace paulo angustius, feminae fronte convexiuscula, maris late profundeque excavata. Thorax coleopteris paulo angustior, latitudine duplo fere brevior, lateribus et basi parum, angulis anterioribus fortiter rotundatis, creberrime punctatus, feminae convexiusculus, aequalis, maris profunde lateque canaliculatus. Elytra thorace plus sesqui longiora, crebre punctata. Abdomen supra parcius punctatum, pilis raris depressis vestitum, segmento ultimo maris supra plano, subtiliter granulato. Pedes nigri, tibiis tarsisque rufis, tibiis posterioribus medio saepe infuscatis.

Mas fronte excavata, thorace late canaliculato, abdominis segmento ultimo supra granulato distinctus.

- \*90. *H. fervidula*: Nigro-picea, capite thoraceque transverso rufis, pedibus antennisque testaceis, his apice nigris. — Long.  $\frac{3}{4}$  lin.

Habitat in Columbia, Dom. Moritz.

Statura H. analis. Corpus lineare, tenuissime pubescens. Antennae capite paulo longiores, apicem versus sensim incrassatae, articulis penulti-



mis transversis, ultimo maiusculo, subgloboso, testaceae, apice nigrae. Caput thoracis latitudine, pedunculo brevi thoraci insertum, profunde punctatum, rufum. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus et basi parum rotundatis, angulis anterioribus fortiter rotundatis, supra parum convexus, aequalis, subtiliter creberrime punctulatus, laete rufus. Elytra thorace sesqui longiora, crebre punctata, picea. Abdomen supra parcius punctulatum, nigro-piceum, nitidulum. Pectus nigrum. Pedes testacei.

\* 91. *H. cauta*: *Nigra, nitidula, pedibus fuscis, thorace transverso, leviter convexo, basi obsolete canaliculato abdomineque supra basi confertim, postice parce punctato margine pilosellis.* — Long. 1 lin.

*Er. Col. March. I. 334. 34.*

*Homalota pulicaria Er. Col. March. I. 340. 45.*

Habitat in Germania.

Forma fere *H. socialis*, at multo minor, nigra, nitidula, pube tenuissima parce vestita, thoracis abdominisque lateribus pilosellis. Antennae capitis thoracisque longitudine, apicem versus vix incrassatae, articulis 4 — 6 subglobosis, 7 — 10 subturbinatis, longitudine haud crassioribus, ultimo elongato-ovato, apice subacuminato, totae nigrae. Caput thorace angustius, rotundatum, crebre minus subtiliter punctatum, basi leviter constrictum. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus et basi leviter rotundatus, apice truncatus, angulis anterioribus fortiter rotundatis, deflexis, leviter transversim convexus, subtilissime punctatus, basi obsolete canaliculatus. Elytra thorace paulo longiora, confertissime punctata, minus nitida, fusco-nigra. Abdomen nitidulum, supra segmentis quatuor primis confertim, reliquis parce punctatis. Pedes fusci vel fusco-testacei, tarsi dilutioribus.

Obs. *Hom. pulicaria* secundum singulum specimen descripta, accuratius examinata ab hac specie distinguì non debet.

\* 92. *H. celata*: *Nigra, subopaca, pedibus fusco-testaceis, thorace transverso, parum convexo, basi obsolete canaliculato, abdomine supra toto confertim punctato.* — Long. vix 1 lin.

*Er. Col. March. I. 335. 35.*

Habitat in Germania.

Praecedenti simillima, nigra, subopaca. Antennae vix capitis thoracisque longitudine, apicem versus subincrassatae, articulis 4 — 10 longitudine vix crassioribus, ultimo oblongo-ovato, nigrae. Caput thorace paulo angustius, rotundatum, basi leviter constrictum, confertissime punctatum, fronte supra canalicula abbreviata leviter impressa. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus et basi leviter rotundatus, apice truncatus, angulis anterioribus fortiter rotundatis, parum convexus, confertissime punctatus, basi obsoletius canaliculatus. Elytra thorace sesqui fere longiora, confertissime punctata, fusco-nigra. Abdomen supra aequaliter confertim punctatum, lateribus vix pilosellum. Pedes fusci, tarsi flavescens, interdum toti fusco-testacei.

Obs. Praecedenti proxima et aegre distinguenda, differe autem videtur thorace lateribus haud pilosello, abdomine supra toto confertim punctato et colore magis opaco.

- \* 93. *H. recondita*: *Nigra, nitidula, tarsis piceis, thorace transverso, leviter convexo abdomineque supra aequaliter densius punctato margine pilosellis.* — Long. 1 lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Praecedentibus proxima, nigra, parce et omnium subtilissime pubescens, thoracis abdominisque margine laterali pilosello. Antennae capitis thoracisque longitudine, apicem versus vix incrassatae, articulis penultimis vix transversis, ultimo oblongo, apice subacuminato, totae nigrae. Caput thorace angustius, basi parum constrictum, parce punctatum. Thorax coleopteris vix angustior, latitudine duplo brevior, lateribus et basi leviter rotundatus, leviter transversim convexus, subtiliter punctulatus, basi interdum obsolete canaliculatus. Elytra thorace paulo longiora, dense subtilius punctata, apice medio conjunctim emarginata. Abdomen supra totum dense minus subtiliter punctatum. Pedes nigri, tarsis vel etiam tibiis anticis piceo-testaceis.

Obs. Statura *H. luteolae* proxima, differt autem thoracis abdominisque margine laterali pilosello. Utraque habitu quibusdam sectionis quartae speciebus (e. g. *H. fungi*) appropinquata, sunt autem iis neque thorax neque abdomen apicem versus angustata.

- \* 94. *H. subida*: *Nigra, nitidula, elytris fuscis, tibiis tarsisque testaceis, thorace transverso, aequali, leviter convexo, abdomine supra parcius aequaliter punctato.* — Long. vix 1 lin.

Habitat in Brasilia, Dom. Prof. Germar.

Statura praecedentium, nigra, nitidula, thorace abdomineque margine haud pilosellis. Antennae capite thoraceque paulo longiores, apicem versus leviter incrassatae, articulis penultimis subtransversis, ultimo ovato, totae nigrae. Caput thorace parum angustius, rotundatum, basi modice constrictum, subtilissime parce punctatum. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus et basi leviter rotundatus, apice truncatus, angulis anterioribus rotundatis, leviter convexus, subtilissime punctatus. Elytra thorace paulo longiora, nigro-fusca, densius evidenter punctata. Abdomen supra totum aequaliter parce subtiliter punctatum, apice pilosellum. Pedes testacei, femoribus fusciscentibus.

- \* 95. *H. sordidula*: *Nigra, opaca, elytris anoque fuscis, pedibus pallidis, fronte impressa, thorace transversim subquadrato, subtiliter canaliculato, abdomine supra toto confertim punctato.* — Long.  $\frac{2}{3}$  lin.

Er. Col. March. I. 335. 36.

Habitat Berolini, Dom. Schüppel.

Forma praecedentium, at minor, nigra, opaca. Antennae capite thoraceque longiores, apicem versus parum incrassatae, articulis 4 — 10 longitudine haud crassioribus, ultimo elongato, apice subacuminato, nigrae. Caput thorace parum angustius, rotundatum, basi leviter constrictum, fronte late canaliculata. Thorax coleopteris paulo angustior, latitudine dimidio brevior,

lateribus subrectis, basi rotundatus, apice truncatus, angulis anterioribus obtusiusculis, parum convexus, subtiliter canaliculatus. Elytra thorace paulo longiora, nigro-fusca. Abdomen dense punctatum, apice piceum. Pedes flavescences, femoribus fusciscentibus.

Obs. Antennis longioribus, fronte canaliculata, colore opaco et statura minima a praecedentibus praecipue discedens.

\*96. *H. luteicornis*: *Linearis, subdepressa, distinctius punctulata, nigra, elytris fuscis, ore, antennis pedibusque testaceis, abdomine supra crebre, apice parcius punctato.* — Long. 1 lin.

*Er. Col. March. I. 332. 29.*

Habitat in Germania: Berolini singulum specimen cepi.

Corpus minutum, lineare. H. anali magnitudine aequalis, nigra, parum nitida, tenuiter griseo-pubescent. Antennae capite sesqui longiores, apicem versus sensim leviter incrassatae, articulo secundo crassiusculo, tertio hoc duplo minore, 4 — 10 brevibus, sensim paulo crassioribus et simul fortius transversis, penultimis longitudine duplo crassioribus, ultimo maiusculo, breviter ovato, totae luteo-testaceae. Caput thoracis prope latitudine, subdeflexum, confertim minus subtiliter punctatum, ore testaceo. Thorax coleopterorum latitudine, lateribus et basi parum rotundatus, apice truncatus, angulis subrectis, anterioribus deflexis, leviter transversim convexus, aequalis, confertim minus subtiliter punctatus. Elytra thorace paulo longiora, confertim minus subtiliter punctata, obscure fusca, apice ferruginea, ad angulum anteriorem subsinuata. Abdomen confertim minus subtiliter punctulatum, longius pubescens, segmento quinto parce punctato, apice piceum, segmentis ventralibus margine postico picescente. Pedes testacei.

Obs. Capite deflexo, basi nullo modo constricto ad Oxypodas accedit, at tarsi omnino huius generis.

✕ \*97. *H. flavipes*: *Nigra, nitida, subtilius punctulata, antennis basi apiceque pedibusque testaceis, elytris rufo-piceis, thorace aequali, angulis posterioribus acutiusculis, abdomine supra sublaevigato.* — Long. prope 2 lin.

*Aleochara flavipes* Grav. Mon. 161. 32. — Gyll. Ins. Suec. II. 421. 43. — Sahlb. Ins. Fenn. I. 386. 73.

*Bolitochara flavipes* Mannerh. Brachelytr. 82. 42.

Habitat Lutetiae, Dom. Dr. Aubé.

Maximis H. socialis individuis longitudine aequalis, at latior crassiorque, nigra, nitida, pube grisea sericante vestita. Antennae capitis thoracisque fere longitudine, validiusculae, articulo secundo tertioque subaequalibus, 4—10 inter se itidem aequalibus, longitudine haud crassioribus, ultimo elongato, praecedentibus tribus coniunctis prope aequali, acuminato, rufae, medio fuscae. Caput thorace paulo angustius, globosum, fronte satis convexa, parce subtilissime punctata, oris partibus rufis. Thorax latitudine duplo brevior, basi coleopterorum prope latitudine, antrorsum subangustatus, angulis anterioribus fortiter rotundatis, basi rotundatus, utrinque sinuatus, angulis posterioribus acutiusculis, parum convexus, aequalis, subtilius crebriusque punctulatus. Scutellum punctulatum, nigrum. Elytra thorace sesqui longiora, con-



fertum subtilius punctulata, rufo-brunnea, circa scutellum et versus latera nigricantia, apice ad angulum anteriorem leviter sinuata. Abdomen supra parce subtiliter punctatum, fere laevigatum, segmentis anterioribus penultimoque margine apicali, ultimoque toto, subtus segmentorum omnium marginibus anoque rufo-piceis. Pedes rufo-testacei.

- \* 98. *H. subsinuata*: Nigra, nitidula, subtiliter punctata, pedibus piceo-testaceis, thorace leviter convexo, basi fortiter rotundato, utrinque subsinuato, angulis posterioribus prominulis, abdomine supra parce subtiliter punctato, nitido. — Long.  $1\frac{1}{2}$  lin.

Habitat in Germania meridionali, Dr. Walth.

Praecedente duplo minor, angustior, minus crassa, sequentibus aequalis, nigra, nitida, subtiliter tenuiterque griseo-pubescent. Antennae capitis thoracisque longitudine, apicem versus sensim incrassatae, articulo secundo tertioque aequalibus, 4—10 subobconicis, sensim paulo crassioribus, penultimis duobus transversis, ultimo maiusculo, praecedentibus ambobus sublongiore, oblongo-ovato, totae nigrae. Palpi picei. Caput minutum, thorace multo minus, rotundatum, basi haud constrictum, fronte leviter convexa, subtiliter minus dense punctatum. Thorax coleopteris dimidio brevior, basi coleopterorum latitudine, antrorsum subangustatus, apice truncatus, lateribus leviter rotundatus, basi medio late et sat fortiter rotundatus, utrinque subsinuatus, angulis posterioribus prominulis, subrectis, at minutis, anterioribus rotundatis, leviter transversim convexus, densius subtiliter punctatus, basi supra scutellum foveola minuta leviter impressus. Elytra thorace paulo longiora, apice ad angulum anteriorem subsinuata, confertim subtiliter punctata, minus nitida. Abdomen lineare, nigrum, nitidum, segmentis tribus primis parcius, sequentibus duobus parce, ultimo iterum crebrius subtiliter punctatis, hoc piceo. Pedes piceo-testacei.

- \* 99. *H. anceps*: Fusco-nigra, opaca, punctatissima, pedibus luteis, thorace plano, canaliculato, angulis posticis subrectis, abdomine supra subtiliter punctulato, nitidulo, ano piceo. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 336. 37.

Habitat Berolini, rarissime, Lutetiae, Dom. Aubé.

Statura fere *H. socialis* et minoribus eius individuis magnitudine aequalis, fusco-nigra, opaca, tenuissime pubescens. Antennae capitis thoracisque vix longitudine, graciliores, apicem versus parum incrassatae, articulis 4—10 longitudine haud crassioribus, ultimo praecedentibus ambobus longitudine aequali, oblongo-ovato, apice subacuminato, fuscae, basi obscure luteae. Caput thorace angustius, basi vix constrictum, confertim punctatum. Thorax coleopteris paulo angustior, latitudine plus dimidio brevior, antrorsum subangustatus, lateribus leviter rotundatus, basi utrinque subsinuatus, angulis posterioribus subrectis, supra depressus, confertissime punctatus, leviter canaliculatus. Elytra thorace sesqui longiora, confertissime punctata, subrugulosa, apice ad angulum anteriorem sinuata. Abdomen parallelum, nigrum, nitidulum, apice piceum, supra subtiliter confertissime punctatum. Pedes lutei.

Obs. 1. Tarsi postici articulis 4 primis longitudine aequalibus.

2. A. Dom. Aubé in societate Formicae rufae lecta.

- \* 100. *H. concolor*: *Nigra, parum nitida, pube cinerascenti sericea, tarsis piceis, thorace subdepresso, supra scutellum foveolato, basi utrinque sinuato, angulis posticis acutiusculis.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Germania meridionali, Dom. Dr. Walzl.

Statura omnino praecedentis, nigra, minus nitida, pube subtilissima densa cinereo-sericea, omnium subtilissime confertissimeque punctata. Antennae capite thoraceque breviores, tenuiores, apicem versus haud incrassatae, articulo secundo tertio paulo minore, 4 — 10 subaequalibus, ultimo praecedentibus ambobus longitudine aequali, apice acuminato, totae nigrae. Palpi nigri. Caput thorace paulo angustius, rotundatum, basi leviter constrictum. Thorax latitudine dimidio prope brevior, basi coleopteris vix angustior, antrosum parum angustatus, basi leviter rotundatus, ad angulum utrinque sinuatus, lateribus leviter rotundatus, apice subemarginatus, angulis anterioribus leviter rotundatis, posterioribus acutiusculis, subdepressus, basi foveolatus. Elytra thorace paulo longiora, apice ad angulum anteriorem subsinuata. Abdomen apicem versus subangustatum. Pedes nigri, tarsis testaceis.

Obs. Tarsi postici articulis 4 primis longitudine sensim decrescentibus, nihilominus huius generis videtur, summa affinitate cum praecedentibus coniuncta.

### \*\*\* R o t u n d a t a e.

- \* 101. *H. notha*: *Brevis, nigra, nitida, griseo-pubescens, antennarum basi pedibusque rufis, thorace brevi, convexo, abdomine lateribus rotundato.* — Long. 1 lin.

Habitat in Carinthia, Dom. Schmidt.

Statura brevi crassulaque omnino sequentis, at duplo minor, pariter Gyrophanae simulans. Antennae capite thoraceque vix longiores, extrorsum sensim leviter incrassatae, articulis penultimis transversis, ultimo breviter ovato, acuminato, fuscae. Corpus nigrum, nitidum, griseo-pubescens, lateribus pilosellum. Caput subdeflexum, basi parum constrictum, fronte utrinque vage punctata, palpis rufis. Thorax coleopteris parum angustior, latitudine duplo brevior, lateribus leviter rotundatus, antice truncatus, basi rotundatus, supra undique sat convexus, vage subtilissime punctatus. Elytra thorace vix longiora, dense punctata, densius longiusque pubescentia. Abdomen breve, rotundatum, supra obsolete punctatum, pubescens, planum, margine elevato, subtus convexum. Pedes rufi, femoribus posterioribus medio piceis.

Variat corpore brunneo, capite abdominisque apice nigris.

- \* 102. *H. hirta*: *Brevis, ferruginea, griseo-hirta, antennis fuscis, basi apiceque testaceis, thorace brevi, convexo, abdomine supra subglabro, parce punctato.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Brasilia.

Ferruginea, nitida, pilis densis erectis griseis hirta. Antennae graciles, capitis thoracisque longitudine, articulis 5—10 subobconicis, longitudine haud crassioribus, ultimo ovato, acuminato, basi apiceque testaceae, medio fuscae. Caput deflexum, thorace angustius, basi vix constrictum. Thorax coleopteris an-

gustior, latitudine plus duplo brevior, valde convexus, subtiliter vage punctatus. Elytra thorace paulo longiora, apice ad angulum exteriorem leviter sinuata, fortius minus dense punctata. Abdomen subtus convexus, supra pilis depressis raris vestitum, minus crebre punctatum, ante apicem piceum, apice testaceum. Pedes testacei.

Obs. Species singularis in hoc genere, a reliquis habitu discedens, Gyrophaenam simulans, at neque corpus grabrum neque pedes intermedii basi spatio maiore distantes; capite deflexo et corpore puberulo Aleocharam etiam facile mentiens, ab utroque genere autem tarsorum structura remota.

103. *H. importuna*: Brevis, testacea, nitida, glabra, capite thoraceque nigris, hoc parvo, brevi, leviter convexo, abdomine lateribus rotundato. — Long.  $1\frac{1}{3}$  lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Habitu fere Gyrophaenae, ad huius generis videtur. Antennae capitis thoracisque prope longitudine, apicem versus sensim incrassatae, testaceae, articulis duobus primis flavis. Caput thoracis latitudine, suborbiculatum, fere transversum, basi profunde constrictum, fronte leviter convexa, laevissimum, utrinque subtiliter nigro-pilosellum, nigrum, nitidum, oris partibus flavis. Thorax coleopteris duplo angustior, latitudine dimidio brevior, basi lateribusque eodem arcu rotundatus, apice truncatus, leviter convexus, basi distinctius marginatus, laevissimus, niger, nitidus, vix pubescens. Scutellum testaceum. Elytra thorace paulo longiora, minus depressa, parce obsoletissime punctulata, testacea, nitida. Abdomen rufo-testaceum, supra concavum, margine laterali sat elevato, acuto, modice rotundato. Pectus rufo-testaceum. Pedes flavi, intermedii basi spatio lato distantes, posteriores tarsi sat elongatis, postici tibiis elongatis, leviter curvatis.

#### \*\*\*\*\* F u s i f o r m e s .

- \* 104. *H. cinnamomea*: Rufo-brunnea, subtilissime punctulata, thorace transverso, leviter convexo abdomineque nigro, supra polito apicem versus angustatis. — Long. 2 lin.

Er. Col. March. I. 336. 38.

*Al. cinnamomea* Grav. Micr. 88. 30. Mon. 160. 30. — Latr. Hist. nat. Crust. et Ins. IX. 386. 30. — Gyll. Ins. Suec. II. 412. 34. — Sahlb. Ins. Fenn. I. 382. 65. *Bolitochara cinnamomea* Mannerh. Brachelytr. 82. 43. — Boisd. et Lacord. Faun. Ent. Paris. I. 552. 19.

Habitat in Europa.

Statura fere Aleocharae. Rufo-brunnea, opaca, subtilissime punctata, tenuiter pubescens, lateribus pilosella. Antennae vix capitis thoracisque longitudine, apicem versus incrassatae, articulo secundo tertio dimidio brevior, 5 — 10 transversis, sensim paulo crassioribus, ultimo praecedentibus tribus coniunctis parum brevior, acuminato, pilosellae, fuscae, articulis tribus primis et ultimo apice testaceis. Palpi testacei. Caput thorace angustius, basi vix constrictum, fronte parum convexa. Thorax basi coleopteris paululum angustior, latitudine duplo brevior, antrorsum leviter angustatus, lateribus praecipue ante medium et basi rotundatus, apice truncatus vel subemarginatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, leviter con-



vexus, aequalis. Elytra thorace paulo longiora, apice truncata, ad angulum exteriorem subsinuata, sutura basi elevata, carinata. Abdomen nigrum, nitidum, apice segmentorumque margine rufo-piceis, infra convexum, parcius punctatum, supra planum, laevigatum, recurvum. Pedes testacei.

Mas abdominis segmento dorsali sexto tuberculo minuto munito distinctus.

105. *H. semirufa*: Nigra, thorace, elytris, antennis pedibusque rufis, thorace transverso, leviter convexo abdomineque supra parce punctato apicem versus subangustatis. — Long.  $1\frac{1}{3}$  lin.

Habitat in Gallia? Mus. Dom. Chevrolat.

Habitu omnino praecedentis, at triplo minor. Antennae capite thoraceque breviores, apicem versus incrassatae, articulo secundo tertioque subaequalibus, 5—10 transversis, sensim paulo crassioribus, ultimo praecedentibus ambobus paulo longiore, acuminato, pilosellae, rufae, medio fuscescentes. Caput thorace paulo angustius, deflexum, basi vix constrictum, fronte leviter convexa, parce punctulatum, griseo-pubescentia, nitidum, piceum, ore rufescente. Thorax latitudine duplo prope brevior, basi coleopteris vix angustior, apicem versus vix angustatus, lateribus ante medium et basi fortius rotundatus, basi utrinque subsinuatus, apice truncatus, angulis obtusis, subrotundatis, anterioribus deflexis, leviter convexus, supra scutellum leviter impressus, subtiliter punctulatus, rufus, nitidus, griseo-pubescentia. Elytra thorace paulo longiora, apice truncata, ad angulum exteriorem leviter sinuata, minus dense distinctius punctata, rufa, nitida, griseo-pubescentia. Abdomen apicem versus subangustatum, supra parcius punctulatum, nigrum, nitidum, longius griseo-pubescentia. Pectus parce punctatum, nigropiceum. Pedes rufi, coxis omnibus concoloribus.

Mas abdominis segmento quinto dorsali carinula acuta elevata instructo distinctus.

Obs. Habitus fere Silusae, at palpi labiales distincte triarticulati.

106. *H. discors*: Nigra, nitida, confertim punctata, antennarum basi pedibusque testaceis, elytris brevibus abdomineque rufis, illis angulo exteriore, hoc ante apicem nigris, thorace brevissimo, basi foveolato. — Long. 2 lin.

Habitat ad Caracas Columbiae, Dom. Moritz.

Habitu fere Aleocharae, at huius generis, *H. cinnamomeae* aequalis, licet paulo latior, nigra, nitidula, tenuiter pubescens. Antennae capite thoraceque paulo breviores, apicem versus sensim leviter incrassatae, articulis 5—10 transversis, ultimo sat magno, compresso, nigrae, articulis 3 primis testaceis, ultimo summo apice ferrugineo. Palpi testacei. Caput basi haud constrictum, thoraci immersum, breviusculum, subtiliter punctulatum. Thorax coleopterorum prope latitudine, antice paulo angustior, latitudine plus duplo brevior, lateribus rotundatus, supra modice convexus, crebre subtiliter punctatus, basi supra scutellum foveola transversa sat profunda impressus. Elytra thorace sesqui longiora, sat crebre punctata, prope basin medio leviter elevata, apice ad angulum exteriorem leviter sinuata, testaceo-rufa, angulo exteriore apicali nigro. Abdomen apicem versus leviter attenuatum, supra

sublaevigatum, rufum, segmentis duobus antepenultimis nigris. Pedes rufo-testacei.

- \* 107. *H. longicornis*: Oblonga, subdepressa, nigra, nitida, creberrime punctulata, elytris, pedibus anoque testaceis, thorace suborbiculato, abdomine supra confertissime punctato. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 337. 40.

*Al. longicornis* Grav. Micr. 87. 29. — Latr. Hist. nat. Crust. et Ins. IX. 386. 89.

*Bolitochara longicornis* Mannerh. Brachelytr. 83. 50.

*Aleochara validicornis* var. *b* Sahlb. Ins. Fenn. I. 369. 44.

Var. *Fusco-testacea, immatura.*

*Bolitochara validicornis* Mannerh. Brachelytr. 83. 51.

*Aleochara validicornis* Sahlb. Ins. Fenn. I. 369. 44.

Habitat in Europa.

Praecedente magis depressa et paulo minor, nigra, nitida, pube depressa parce vestita, lateribus pilosella. Antennae capite thoraceque longiores, crassiusculae, articulo secundo tertio paulo minore, 5 — 10 crassitie paulo longioribus, cylindricis, ultimo praecedentibus ambobus paulo brevior, elongato-ovato, piceae, basi interdum rufescentes. Palpi fusco-testacei. Caput parvum, rotundatum, subtiliter parce punctatum, fronte parum convexa vel subdepressa. Thorax coleopterorum prope latitudine, antrorsum subangustatus, longitudine parum brevior, lateribus et basi rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, posterioribus subrotundatis, parum convexus, dense subtiliter punctatus, interdum obsolete canaliculatus. Elytra thorace parum longiora, apice truncata, depressa, confertissime punctata, fusco-testacea. Abdomen nitidulum, nigrum, apice et segmentorum singulorum marginibus piceo-testaceis, undique dense punctatum. Pedes testacei.

Obs. 1. Tibiae setis longis instructae et spinulis apicalibus longioribus munitae affinitatem quandam cum Aleocharis genuinis indicare videntur. Thoracis forma ambigua inter hanc et secundam sectionem est haec species, *H. sociali* facie simillima, abdomen autem distincte apicem versus attenuatum.

2. Typus Gravenhorstianus perditus, inde in Mon. emissa. Descriptio citata huic speciei satis congruit. Ab auctoribus Suecis saepius cum *H. sociali* confusa esse videtur, nam specimen pro *Al. sericante* Gyll. a Dom. Schönherr missum Dom. Schüppel communicavit.

- \* 108. *H. lividipennis*: Nigra, opaca, dense subtiliter punctata, sericeo-pubescentis, antennis longis, crassis, piceis, pedibus elytrisque testaceis, his circa scutellum infuscat. — Long.  $1\frac{2}{3}$  lin.

*Oxyopoda lividipennis* Mannerh. Brachelytr. 70. 2.

*Aleochara lividipennis* Sahlb. Ins. Fenn. I. 388. 77.

*Homalota livida* Er. Col. March. I. 337. 39.

Habitat in Europa, America septentrionali.

Statura fere *Oxyopodae* opacae et parum minor. Caput nigrum, opacum, confertissime subtiliter punctatum, pube tenui griseo-sericea obductum. Antennae capite thoraceque longiores, crassae, articulo tertio secundo paulo longiore, 4 — 10 longitudine paulo crassioribus, cylindricis, ultimo praece-

dentibus ambobus longiore, elongato-cylindrico, acuminato, totae nigro-vel fusco-piceae. Caput thorace multo angustius, rotundatum, deflexum, fronte convexa, palpis fuscis. Thorax basi coleopterorum latitudine, antrorsum angustatus, latitudine postica dimidio brevior, basi et lateribus rotundatus, apice subemarginatus, angulis omnibus obtusis, anterioribus deflexis, leviter convexus, basi leviter transversim impressus, obsolete canaliculatus. Elytra thorace vix longiora, apice ad angulum anteriorem leviter sinuata, dilute testacea, macula communi triangulari fuscescente circa scutellum. Abdomen apicem versus sensim attenuatum, margine pilosellum, apice et segmentorum singulorum marginibus obscure testaceis. Pedes testacei.

Variat elytris testaceis, unicoloribus, nonnunquam etiam thoracis lateribus obscure testaceis.

\* 109. *H. terminata*: *Nigra, subnitida, antennarum articulo ultimo ferrugineo, pedibus piceis, thorace transverso, abdomine basi parcius, apice crebrius punctato.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Puerto Rico, Dom. Moritz.

Corpus nigrum, nitidulum, pube parca subtili depressa vestitum. Antennae capite thoraceque paulo longiores, apicem versus leviter incrassatae, articulis 5—10 transversis, sensim paulo crassioribus, nigrae, articulo ultimo ferrugineo. Caput thorace multo angustius, rotundatum, parce punctatum, ore subangustato, rufo-piceo. Thorax latitudine duplo fere brevior, basi coleopterorum latitudine, antrorsum paulo angustior, lateribus, et basi rotundatus, apice truncatus, angulis omnibus obtusis, posterioribus subrotundatis, anterioribus deflexis, parum convexus, dense subtiliter punctatus. Elytra thorace paulo longiora, apice ad angulum anteriorem subsinuata, confertissime subtiliter punctata. Abdomen apicem versus leviter angustatum, supra laevigatum, segmentis singulis postice vage punctatis, sexto solo dense punctato. Pedes picei, geniculis tarsisque dilutioribus.

Mas distinctus elytris angulo apicali interiore tuberculo elevato, abdominis segmento tertio, sextoque carinatis.

Variat interdum colore corporis dilutiore, ferrugineo.

110. *H. rubicula*: *Rufa, nitida, antennarum medio pectoreque piceis, elytris abdominisque segmentis duobus penultimis nigris, thorace transverso, basi subfoveolato.* — Long. vix 1 lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Habitu fere praecedentis, at triplo prope minor. Antennae validae, capite thoraceque longiores, apicem versus sensim incrassatae, testaceae, articulis intermediis fuscis. Caput thorace angustius, basi parum constrictum, rufum, nitidum. Thorax latitudine duplo brevior, basi coleopteris paululum angustior, apicem versus leviter angustatus, lateribus leviter rotundatus, modice convexus, vix nisi obsolete punctulatus, basi obsolete foveolatus, rufus, nitidus, pube tenui erecta grisescente parce vestitus. Scutellum piceum. Elytra thorace duplo fere longiora, fortius crebriusque punctata, longius pubescentia, nigra, nitida, callo humerali, sutura margineque apicali piceis. Abdomen apicem versus subangustatum, parce pubescens, rufum, nitidum, segmentis duobus penultimis nigris. Pectus piceum. Pedes flavo-testacei.



- \*111. *H. testudinea*: *Breviuscula, nigra, nitida, creberrime punctulata, elytris pedibusque testaceis, thorace convexo, abdomine crebrius subtiliter punctato*. — Long.  $1\frac{1}{4}$  lin.

Habitat in Germania: specimen singulum Dom. Schüppel prope Berolinum legit.

*H. longicorni* parum minor, statura fere *H. lividipennis*, at brevior, fortius convexa, apicem versus magis attenuata, habitu fere *Tachypori*, nigra, nitida, pube brevissima tenuissimaque grisea parcius vestita. Antennae capitis thoracisque longitudine, tenuiores, apicem versus haud incrassatae, articulo secundo tertio vix brevior, 4—10 subaequalibus, crassitie haud brevioribus, ultimo praecedentibus ambobus longitudine aequali, subcylindrico, apice acuminato, nigrae, articulis tribus primis obscure piceis. Caput minusculum, thorace autem multo minus et angustius, rotundatum, fronte leviter convexa, subtilissime punctata. Thorax latitudine postica tertia parte brevior, basi coleopteris fere latior, antrorsum angustatus, apice dimidio angustior, basi leviter, lateribus fortius rotundatus, angulis omnibus deflexis, anterioribus subrotundatis, posterioribus obtusis, fortiter transversim convexus, confertissime subtiliter punctatus, aequalis. Elytra thorace paulo longiora, apice ad angulum exteriorem subsinuata, confertim subtilius punctata, piceo-testacea, circa scutellum fusciscentia. Abdomen apicem versus sensim at fortiter attenuatum, acuminatum, confertim subtiliter punctatum, lateribus pilosellum, apice piceum. Pedes testacei.

- \*112. *H. fungi*: *Nigra, nitidula, creberrime punctulata, antennis pedibusque rufis, thorace transverso, convexiusculo, aequali*. — Long.  $1\frac{1}{4}$  lin.

*Er. Col. March. I.* 338. 41.

*Al. fungi* Grav. Mon. 157. 24. — *Gyll. Ins. Suec. II.* 410. 32. — *Sahlb. Ins. Fenn. I.* 378. 58. — *Zetterst. Faun. Lappon. I.* 102. 17. *Ins. Lapp.* 78. 25.

*Bolitochara fungi* Mannerh. *Brachelytr.* 83. 52. — *Runde Brachelytr. Hal.* 32. 12

Habitat in Europa.

*Breviuscula*, utrinque subangustata, nigra, nitidula, parce subtilissime pubescens, confertim punctata. Antennae capitis thoracisque longitudine, graciles, apicem versus parum incrassatae, articulo primo crassiusculo, penultimis longitudine crassioribus, ultimo praecedentibus ambobus aequali, oblongo-ovato, acuminato, pilosellae, rufae, apice interdum fusciscentes. Caput thorace multo angustius, rotundatum, deflexum, fronte leviter convexa, ore piceo, palpis testaceis. Thorax basi coleopterorum latitudine, antrorsum leviter angustatus, latitudine postica dimidio brevior, lateribus et basi leviter rotundatis, angulis omnibus obtusis, subrotundatis, anterioribus deflexis, supra transversim convexus, aequalis. Elytra thorace paulo longiora, densius et minus subtiliter punctata, minus nitida. Abdomen supra confertim, postice parcius punctatum, margine pilosellum. Pedes rufo-testacei.

Variat thorace elytris rufo-piceis, vel corpore toto rufo.

- \*113. *H. vernacula*: *Nigra, nitidula, confertissime punctulata, thoracis lateribus elytrisque ferrugineis, antennarum basi pedibusque testaceis, thorace transverso, convexiusculo, subtiliter canaliculato*. — Long.  $1\frac{1}{4}$  lin.

*Er. Col. March. I. 339. 42.*

*Bolitochara fungi* Boisd. et Lacord. Faun. Ent. Paris. I. 553. 22.

Habitat in Europa.

Statura omnino praecedentis, nigra, nitidula, confertim punctata, subtilissime parce pubescens. Antennae capite thoraceque paulo longiores, apicem versus leviter incrassatae, articulo primo crassiusculo, penultimis longitudine vix crassioribus, ultimo praecedentibus ambobus prope aequali, oblongo-ovato, subacuminato, fuscae, basi testaceae. Caput thorace angustius, rotundatum, parcius subtiliter punctatum, nigrum, nitidum, ore concolor, palpis flavo-testaceis. Thorax basi coleopterorum prope latitudine, antrorsum subangustatus, latitudine dimidio brevior, lateribus et basi rotundatus, apice truncatus, angulis posterioribus obtusis, anterioribus rotundatis, leviter transversim convexus, subtiliter canaliculatus, basi foveola impressus, confertim et minus subtiliter punctatus, nigro-piceus, lateribus rufescentibus. Elytra thorace paulo longiora, dense minus subtiliter punctata, rufo-ferruginea, circa scutellum et lateribus leviter infuscata. Abdomen nigrum, apice piceum, segmentis singulis piceo-marginatis, supra dense punctatum, margine pilosellum. Pedes testacei.

Mas abdominis segmento dorsali sexto apice 4-dentato distinctus.

Obs. 1. Statura praecedenti simillima, differt autem colore, punctura minus subtili, thorace canaliculato, basi impresso, angulis posterioribus haud rotundatis.

2. *Bolitochara patruelis* Boisd. et Lacord. Faun. Ent. Paris. I. 553. 23. „huic paulo maior (Long. 1 lin. lat.  $\frac{1}{2}$  lin.) ceterum forma et colore simillima, elytris fuliginosis, macula magna obscure ferruginea, pedibus rufescenti-brunneis,” mihi obscura et dubia, num ab ea vere distincta sit. — *Oxypoda obfuscata* Boisd. et Lacord. Faun. Ent. Paris. I. 539. 4. quoque ad *H. vernaculam* pertinere videtur.

\* 114. *H. orbata*: Nigra, nitidula, creberrime punctulata, elytris rufo-brunneis, pedibus testaceis, thorace transverso, convexiusculo, aequali. — Long. 1 lin.

*Er. Col. March. I. 339. 43.*

*Bolitochara agaricola* Mannerh. *Brachelytr.* 84. 53. (forte)

*Aleochara agaricola* Sahlb. *Ins. Fenn. I. 379. 59. (forte)*

Habitat in Germania.

*H. fungi* simillima, nigra, nitidula, confertim punctata, subtilissime parce pubescens. Antennae capite thoraceque paulo longiores, apicem versus vix incrassatae, articulo primo crassiusculo, penultimis longitudine haud crassioribus, ultimo penultimis ambobus paulo longiore, acuminato, totae nigro-fuscae. Caput thorace paulo angustius, orbiculatum, deflexum, parce subtiliterque punctulatum, ore piceo, palpis testaceis. Thorax amplus, latitudine tertia parte brevior, basi coleopterorum latitudine, antrorsum parum angustatus, lateribus et basi leviter rotundatus, angulis omnibus obtusis, subrotundatis, transversim convexus, aequalis, confertim subtiliter punctatus. Elytra thorace paulo longiora, confertim minus subtiliter punctata, rufo-brunnea. Abdomen supra conferim, apice parcius punctatum, margine pilosellum, ano piceo. Pedes testacei.

Obs. Ab *H. fungi* antennis obscuris et thorace paulo longiore distinguitur.

- \* 115. *H. clientula*: *Nigra, nitidula, subtiliter punctulata, elytris fortiter punctatis pedibusque ferrugineis, thorace transverso, convexiusculo, abdomine supra parce punctato.* — Long. 1 lin.

Habitat in ins. Corcyra, Dom. Parreyss.

Praecedentibus angustior, nigra, nitida, pube tenui depressa parcius vestita. Antennae capitis thoracisque longitudine, apicem versus subincrassatae, articulo primo crassiusculo, penultimis longitudine haud crassioribus, ultimo praecedente sesqui longiore, ovato, fuscae. Palpi testacei. Caput thorace multo minus et angustius, deflexum, rotundatum, subtilissime vage punctatum. Thorax latitudine dimidio fere brevior, basi coleopteris vix angustior, antrorsum distinctius angustatus, lateribus et basi leviter rotundatus, apice truncatus, angulis omnibus obtusis, subrotundatis, leviter convexus, subtiliter punctatus, aequalis. Elytra thorace paulo longiora, apice ad angulum anteriorem distinctius sinuata, confertim fortius punctata, saturate testacea. Abdomen apicem versus distincte attenuatum, supra parcius subtiliusque punctatum, ano piceo. Pedes testacei.

Varietatem antennis, thorace elytrisque rufo-testaceis in Aegypto collegit Dom. Prof. Ehrenberg.

Obs. Affinis praecedentibus, statura autem angustiore, thorace lateribus minus rotundato, elytris fortiter punctatis et apice sinuatis differt.

- \* 116. *H. orphana*: *Nigra, nitidula, confertim punctulata, antennarum basi pedibusque obscure testaceis, thorace transverso, convexiusculo, aequali.* — Long.  $\frac{3}{4}$  lin.

Er. Col. March. I. 340. 44.

Habitat in Germania.

*H. aterrima* paulo minor, ceterum habitu praecedentium, nigra, distincte punctulata, subtiliter parce pubescens, nitidula. Antennae capitis thoracisque longitudine, apicem versus leviter incrassatae, articulo primo crassiusculo, articulis 6—10 longitudine paulo crassioribus, ultimo praecedentibus ambobus aequali, oblongo-ovato, subacuminato, pilosellae, nigricantes, articulo primo, interdum etiam secundo obscure testaceo. Caput thorace angustius, orbiculatum, parce subtiliter punctatum, ore piceo, palpis testaceis. Thorax basi coleopterorum latitudine, antrorsum subangustatus, latitudine dimidio brevior, basi rotundatus, utrinque subsinuatus, angulis posterioribus obtusis, lateribus angulisque anterioribus rotundatis, apice truncatus, leviter transversim convexus, aequalis, confertim subtiliter punctatus. Elytra thorace sesqui longiora, apice ad angulum anteriorem leviter sinuata, dense minus subtiliter punctata, fusco-nigra. Abdomen nitidulum, crebrius distinctiusque punctulatum, margine pilosellum. Pedes fusco-testacei.

- \* 117. *H. aterrima*: *Nigra, subopaca, tenuiter sericeo-pubescens, pedibus flavescentibus, thorace transverso, convexiusculo, aequali.* — Long. 1 lin.

Er. Col. March. I. 341. 46.

*Aleochara aterrima* Grav. Micr. 83. 22. Mon. 162. 33.

*Bolitochara fuscula* Mannerh. Brachelytr. 84. 54.

*Aleochara fuscula* Sahlb. Ins. Fenn. I. 379 60 — Zetterst. Ins. Lappon. 78. 24



*Var. a: Elytris brunneis.*

*Aleocho. pygmaea* Grav. Micr. 36. 27. Mon. 163. 39. — Latr. Hist. nat. Crust. et Ins. IX. 385. 27.

*Aleocho. umbrata* Grav. Micr. 90. 32. Mon. 163. 42. — Latr. Hist. nat. Crust. et Ins. IX. 387. 32.

*Bolitochara minutissima* Boisd. et Lacord. Faun. Ent. Paris I. 553. 21. (forte)

*Var. b: Thoracis lateribus elytrisque brunneis.*

*Aleocho. obfuscata* Grav. Micr. 87. 28. Mon. 163. 40. — Latr. Hist. nat. Crust. et Ins. IX. 386. 28. — Gyll. Ins. Suec. II. 418. 40. — Sahlb. Ins. Fenn. I. 385. 71.

*Oxyopoda obfuscata* Mannerh. Brachelytr. 72. 12.

Habitat in Europa.

Statura fere *H. fungi*, at paulo minor, angustior, subopaca, subtilissime dense punctata, pube subtili griseo-sericea dense vestita. Antennae capite thoraceque paulo longiores, apicem versus parum incrassatae, articulo primo haud incrassato, penultimis longitudine vix crassioribus, articulo ultimo oblongo-ovato, subacuminato, nigrae, articulis primis saepe flavescentibus. Palpi fusciscentes. Caput thorace angustius, rotundatum, ore paulum attenuato, deflexum, fronte convexa, nigrum. Thorax coleopteris basi haud angustior, antrorsum leviter angustatus, latitudine postica plus dimidio brevior, basi et lateribus rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus leviter deflexis, subrotundatis, leviter transversim convexus, aequalis, niger vel fuscus, lateribus saepe dilutioribus. Elytra thoracis sesqui longiora, apice ad angulum anteriorem subsinuata, nigra, apicem versus saepe brunnea, vel brunnea, basi magis minusve nigricantia. Abdomen apicem versus attenuatum, supra confertissime subtiliter punctatum, minus opacum, nigrum, apice piceum, parce pilosellum. Pedes obscure flavi, coxis anticis dilutioribus.

Individuis praecocibus thorax et elytra vel corpus totum rufo-testacea.

\* 118. *H. ambigua*: Nigra, opaca, sericeo-pubescent, thorace elytrisque fusco-testaceis, antennarum basi pedibusque flavis, thorace convexiusculo. — Long. 1 lin.

Habitat in America septentrionali, Dom. Zimmermann.

Affinis praecedenti, at angustior, sublinearis, fusco-nigra, opaca, subtilissime confertim punctata, pube tenuissima griseo-sericea dense vestita. Antennae capitis thoracisque longitudine, apicem versus sensim leviter incrassatae, articulo secundo tertio sesqui maiore, penultimis leviter transversis, ultimo maiusculo, oblongo-ovato, apice subacuminato, fuscae, basi flavae. Caput thorace parum angustius, rotundatum, nigrum. Thorax latitudine parum brevior, lateribus et basi rotundatus, apice truncatus, angulis posterioribus obtusis, anterioribus subrotundatis, leviter convexus, aequalis, nigro-piceus. Elytra thoracis latitudine, fusco-testacea. Abdomen supra totum subtiliter confertissime punctatum, nigrum. Pedes flavi, tarsis posticis articulo primo sequentibus paulo longiore.

Species mihi invisae, forte huius generis.

\*

119. *H. (?) crassicornis*: Brevior, nigra, pedibus testaceis, elytris brunneis, thorace amplo, suborbiculato, antennis longis, crassis, totis, nigris. — Long.  $1\frac{1}{2}$  lin.

*Aleocho. crassicornis* Gyll. Ins. Suec. IV. 486. 20–21. — Sahlb. Ins. Fenn. I. 362. 31.

*Bolitochara crassicornis* Mannerh. *Brachelytr.* 77. 12.

Habitat in Suecia, in succo arborum, fungis et quisquiliis, passim.

Statura brevior, latior et crassior quam *H. elongatulae*, *H. sociali* (longicorni) propinqua, sed thorax multo maior, longior, suborbiculatus. Caput thorace paulo minus, rotundatum, convexum, nigrum, nitidum, punctulatum, ore palpisque nigro-piceis. Antennae capite cum thorace multo longiores, crassae, nigrae, articulo primo longo, clavato, basi piceo, 2, 3 longiusculis, obconicis, 4—10 cyathiformibus, sensim crassioribus, ultimo adhuc maiore, oblongo-ovato, acuminato. Thorax amplius, suborbiculatus, vel longitudine perparum latior, supra subdepressus, minus saturate niger, subtiliter punctulatus, tenuiter pubescens, in medio baseos foveola subrotunda et ante illam impressiones obsoletae in medio dorsi. Scutellum parvum, triangulare, concolor. Elytra thoracis medio non latiora, nec illo longiora, apice subsinuata, supra subdepressa, conferunt subtilissime punctulata, aut saturatius aut dilutius brunnea, subopaca. Abdomen pone elytra illis duplo longius, nigrum, nitidum, apice obtusum. Corpus subtus nigro-piceum, nitidum, punctulatum, pubescens, segmentorum ventralium marginibus anoque testaceis. Pedes mediocres, toti testacei. (Gyll.)

120. *H. (?) annularis*: *Linearis, subnitida, rufo-testacea, capite abdominisque cingulo piceis, thorace aequali, antennis validioribus, brevibus, antrosum parum incrassatis.* — *Long. vix 1 lin.*

*Bolitochara annularis* Mannerh. *Brachelytr.* 76. 8.

*Aleochara annularis* Sahlb. *Ins. Fenn. I.* 353. 15.

Habitat in Fennia australi, rarius.

*H. circellari* similis, sed differt corpore magis lineari, thorace brevior et antennis validioribus brevioribusque. Caput thorace paulo angustius, rotundatum, convexum, ferrugineo-piceum, nitidum, vix punctulatum. Os et palpi testacea. Antennae thorace breviores, basi pallide testaceae, extrorsum crassiores, fusco-ferrugineae. Thorax longitudine latior, antice truncatus, lateribus parum rotundatus, postice subtruncatus, supra modice convexus, rufo-testaceus, vix punctulatus, pube sericante obductus. Scutellum minus, triangulare, rufo-testaceum. Elytra thoracis vix latiora, sed eodem paulo longiora, supra rufo-testacea, depressa, subtilissime punctulata. Abdomen pone elytra longitudine reliqui corporis, fusco-piceum, segmentis ultimis testaceis. Corpus subtus ferrugineo-testaceum, segmentis abdominis anterioribus piceis, marginibus tamen ferrugineis. Pedes elongati, pallide testacei. (Sahlb.)

121. *H. (?) haemorrhoea*: *Nigra, subnitida, longius pubescens, antennarum basi, pedibus, apice elytrorum anoque rufis, articulo antennarum ultimo magno, subgloboso.* — *Long.  $\frac{2}{3}$  lin.*

*Bolitochara haemorrhoea* Mannerh. *Brachelytr.* 76. 11.

*Aleochara haemorrhoea* Sahlb. *Ins. Fenn. I.* 353. 20.

Habitat in Fennia australi.

*Phloeopora reptanti* statura simillima, sed duplo minor est, longius pubescens, et aliter colorata, antennisque gaudet brevioribus. Caput thorace

angustius, convexiusculum, nigrum, subnitidum, vix subtilissime punctulatum, pubescens; os et palpi testacea. Antennae thorace breviores, validiusculae, versus apicem incrassatae, fusco-ferrugineae, basi pallidiores, articulo ultimo magno, subgloboso. Thorax longitudine paulo latior, antice truncatus, angulis deflexis, lateribus parum dilatatus, postice subtruncatus, angulis obtusis, supra convexus, niger, subnitidus, subtilissime punctulatus, pube longiore grisescente parcius adpersus. Scutellum brevissimum, nigrum. Elytra thorace paulo latiora et longiora, apice subtruncata, supra nigra, versus apicem rufescentia, subnitida, longius pubescentia, crebre et evidentius punctulata. Abdomen pone elytra longitudine corporis reliqui, nigrum, subnitidum, pubescens, punctulatum, ano rufescente. Corpus subtus nigrum, subnitidum, pubescens, segmentorum marginibus anoque rufis. Pedes elongati, pallidi. (Sahlb.)

Obs. Dubito ane potius ad *Phloeopora* genus revocanda sit, *Phl. corticali* longe minor esse videtur.

122. *H. (?) angustata*: *Linearis, elongata, nigro-picea, ore, antennarum basi, elytris, ano pedibusque testaceis, thorace longiore, postice angustiore, basi foveola impresso.* — Long.  $\frac{3}{4}$  lin.

*Aleochara angustata* Sahlb. *Ins. Fenn. I.* 363. 33.

*Bolitochara compressa* Mannerh. *Brachelytr.* 80. 35.

Habitat in Fennia australi, rarissime.

*H. elongatulae* minoribus individuís (Al. depressiusculae) magnitudine et statura similis, sed differt imprimis thoracis forma et antennis paulo crassioribus. Caput basi truncatum, antice rotundatum, nigrum, subtiliter punctulatum, ore et palpis testaceis. Antennae thoracis longitudine, basi testaceae, versus apicem nigro-fuscae et incrassatae, articulo primo incrassato, clavato, ultimo ovato, obtuso. Thorax longitudine parum latior, antice truncatus, angulis anterioribus obtusis, basi subtruncatus, supra depressus, nigro-piceus, subopacus, subtiliter punctulatus, subpubescens, basi foveola obsoleta impressus. Scutellum minutissimum, nigrum. Elytra thorace latiora et longiora, subquadrata, apice truncata, supra depressa, testacea, subtiliter punctulata, subpubescentia. Abdomen pone elytra reliquo corpore brevius, corpusque subtus nigro-picea, subpubescentia, ano testaceo. Pedes mediocres, toti testacei. (Sahlb.)

123. *H. (?) tenella*: *Fusco-picea, linearis, antennis longis, extrorsum vix crassioribus, thorace subquadrato, canaliculato, tibiis tarsisque pallidioribus.* — Long.  $\frac{2}{3}$  lin.

*Bolitochara tenella* Mannerh. *Brachelytr.* 81. 36.

*Aleochara tenella* Sahlb. *Ins. Fenn. I.* 364. 34.

Habitat in Fennia australi, rarius.

Praecedente (angustata) plus quam duplo minor, minus depressa, thorace latiore subquadrato et colore diversa. (Mannerh.)

124. *H. (?) luridipennis*: *Linearis, nigro-picea, opaca, pubescens, antennarum basi, elytris, pedibus anoque livido-testaceis, thorace subquadrato, postice foveolato.*

*Bolitochara luridipennis* Mannerh. *Brachelytr.* 77. 15.

Habitat Petropoli.



H. sociali triplo maior, magis pubescens, opaca, thorace longiore, antennis medio haud incrassatis diversa. (Mannerh.)

Obs. Forte H. elongatulae var.

\*\*\*

125. *H. (?) macilenta*: *Creberrime punctulata, subnitida, fusco-nigra, elytris quadratis pedibusque fusco-testaceis, thorace obsolete canaliculato.*

*Aleoch. macilenta* Sahlb. *Ins. Fenn. I.* 371. 47. — *Zetterst. Ins. Lapp.* 77. 21.

*Aleoch. pygmaea* Zetterst. *Faun. Lappon. I.* 101. 16.

Habitat in Lapponia, sub cortice Betulae mense Augusto capta.

Pusilla, H. sociali minor, H. fungi subaequalis. Caput thorace paulo angustius, subtilissime punctulatum, nitidum, nigrum, ore fusco-testaceo. Antennae thorace parum longiores, pubescentes, graciles, extrorsum paulo crassiores, totae nigrae. Thorax latitudine paulo latior, supra depressus, niger, parum nitidus, vix pubescens, creberrime punctulatus, in medio baseos foveola parva obsoleta. Elytra subquadrata, tenuiter pubescentia, subtilissime punctulata, testacea. Corpus fusco-nigrum, pubescens. Pedes pallide testacei. (Zetterst.)

Obs. H. oblitae affinis videtur.

126. *H. (?) fusca*: *Brunneo-fusca, nitida, crebre punctulata, sericeo-pubescent, capite abdomineque nigris, thorace obsolete canaliculato, antennis longis, filiformibus, testaceis.* — *Long.* 1 $\frac{1}{6}$  lin.

*Aleoch. fusca* Sahlb. *Ins. Fenn. I.* 371. 48. — *Zetterst. Ins. Lappon.* 78. 22.

Habitat in Lapponia, rarissime, unicum specimen invenit F. Sahlberg.

H. sociali (longicorni) statura satis similis, sed differt colore, antennis tenuioribus et thorace magis depresso. Caput thorace angustius, rotundatum, nigrum, nitidum, subtilissime punctulatum, subpubescens, ore palpisque fusco-brunneis. Antennae thorace longiores, fuscescentes, filiformes, articulo ultimo oblongo-ovato, subacuminato. Thorax transversus, antice truncatus, angulis compressis, lateribus et postice rotundatus, supra subdepressus, obsoletius canaliculatus, brunneo-fuscus, ad latera parum pellucidus, crebre punctulatus, sericeo-pubescent. Scutellum minutum, triangulare, punctulatum, pubescens. Elytra thorace vix vel parum latiora et longiora, apice truncata, supra subdepressa, crebre punctulata, pubescentia, brunneo-fusca. Abdomen pone elytra longitudine reliqui corporis, ovatum, nigrum, nitidum, tenuiter pubescens, vix punctulatum, segmento sexto piceo-ferrugineo. Corpus subtus brunneo-fuscum, nitidum, punctulatum, pubescens, segmentis ventralibus piceo-nigris, sexto piceo-ferrugineo. Pedes toti pallide testacei. (Sahlb.)

Obs. Staphyl. fuscus Marsh. Ent. Britt. 514. 49. quoque ad hoc genus pertinet, mihi autem non extricatus.

127. (*H.*) *scapularis*: *Piceo-ferruginea, nitida, capite, abdominis medio elytrisque nigro-fuscis, humeris, basi antennarum, ano et pedibus pallide rufescentibus, antennis brevibus, apice incrassatis.* — *Long.* vix 1 lin.

*Aleoch. scapularis* Sahlb. *Ins. Fenn. I.* 372. 50.

*Bolitochara axillaris* Mannerh. *Brachelytr.* 78. 19.

Habitat in Fennia australi, rarissime.

H. sociali var. c. (nigritulae) similis, sed differt thorace paulo longiore et convexiore. Caput thorace angustius, rotundatum, basi truncatum, nigro-fuscum, nitidum, subtilissime punctulatum, foveola frontis impressa. Antennae thorace paulo breviores, ferrugineo-piceae, basi pallide rufescentes, articulo primo elongato, incrassato, secundo et tertio itidem longioribus, sequentibus brevibus, versus apicem sensim incrassatis, ultimo oblongo-ovato, apice obtusiusculo. Thorax subtransversus, antice truncatus, angulis depressis, lateribus et postice rotundatus, supra subconvexus, piceo-ferrugineus, nitidus, crebre punctulatus, pubescens. Scutellum minutum, triangulare, nigro-fuscum, punctulatum. Elytra thorace latiora et paulo longiora, apice truncata, supra depressa, nigro-fusca, humeris rufescentibus, crebre punctulata, pubescentia. Abdomen pone elytra vix longitudine reliqui corporis, piceo-ferrugineum, nitidum, segmentis 4—6 nigro-fuscis, ano pallide rufescente, vix punctulatum. Corpus subtus piceo-ferrugineum, nitidum, subtilissime punctulatum, pubescens, segmentis 4—6 ventralibus nigro-fuscis, ano pallide rufescente. Pedes toti pallide rufescentes. (Sahlb.)

128. *H. (?) borealis*: *Nigra, nitida, subtiliter punctulata, pubescens, tarsis piceo-ferrugineis, thorace postice subangulato-rotundato.* — Long.  $1\frac{1}{4}$  lin.

*Aleocho. borealis* Sahlb. *Ins. Fenn. I.* 361. 29. — *Zetterst. Ins. Lappon.* 74. 4.

Habitat in Lapponia, rarissime.

Magnitudine et statura *Tachyusae atrae*, sed ab hac et ceteris thoracis forma imprimis distincta. Caput thorace angustius, suborbiculatum, nigrum, nitidum, subtilissime punctulatum, subpubescens, fronte obsolete foveolata. Antennae thorace longiores, versus apicem incrassatae, articulo ultimo oblongo-ovato. Thorax longitudine latior, antice truncatus, lateribus longe ultra medium subrectus, vel parum tantum rotundatus, iuxta basin vero subangulatum angustatus, basi truncatus, supra depressus, niger, subtilissime punctulatus, pubescens. Scutellum triangulare, nigrum. Elytra thorace latiora et longiora, apice truncata, supra subdepressa, nigra, parum aenescens, subtiliter punctulata. Abdomen pone elytra longitudine reliqui corporis, nigrum, nitidissimum, punctulatum, parce pubescens. Corpus subtus nigrum, nitidum, pube depressa longiore parce obsitum. Pedes nigropicei, tarsis piceo-ferrugineis. (Sahlb.)

Obs. H. ancipiti et concolori affinis videtur.

\*\*\*\*\*

129. *H. (?) marginalis*: *Nigra, nitida, punctulata, pubescens, ore, antennis, thoracis lateribus, elytris, pedibus atque rufo-testaceis, elytrorum angulo apicis fusco.* — Long.  $1\frac{1}{2}$  lin.

*Aleocho. marginalis* Grav. *Mon.* 161. 31. — *Gyll. Ins. Suec. II.* 420. 42.

Habitat in Gallia, Suecia.

Oxypoda alternante paulo latior et crassior, sed non longior. Caput thorace parum angustius, subtransversum, nigrum, nitidum, crebre

punctulatum, versus os angustatum, ore palpisque rufo-testaceis; oculi magni, oblongi, brunnei. Antennae vix longitudine thoracis, crassiusculae, in medio fusco-ferrugineae, basi apiceque dilutius rufo-testaceae. Thorax brevis, transversus, antice truncatus, angulis deflexis, valde compressis, unde ibi angustior apparet, lateribus parum rotundatus, basi truncatus, angulis obtusis, supra convexus, niger, nitidus, subtilissime crebre punctulatus, tenuiter pubescens, angulis posticis late rufo-testaceis, qui color iuxta marginem et basin anguste extenditur. Scutellum breve, triangulare, fuscum, punctulatum. Elytra thoracis basi vix latiora, sed illo dimidio longiora, subquadrata, supra depressa, rufo-testacea, nitida, confertissime et evidentius rugoso-punctata, tenuiter pubescentia, angulo exteriori apicis oblique infusato. Abdomen pone elytra longitudine reliqui corporis, crassum, obtusum, nigrum, nitidum, creberrime punctulatum, ano rufo-testaceo. Corpus subtus nigrum, nitidum, crebre punctulatum, segmentorum ventralium marginibus anoque late rufo-testaceis. Pedes mediocres, toti rufo-testacei. (Gyll.)

Obs. *H. cinnamomeae* affinis videtur.

130. *H. (?) Mannerheimii*: *Brevis, lata, nigra, subnitida, subtilissime punctulata, pubescens, pedibus fusco-testaceis, antennis thorace canaliculato multo longioribus.* — *Long*  $\frac{2}{3}$  lin.

*Aleoch. Mannerheimii* Sahlb. *Ins. Fenn. I.* 330. 61.

Habitat in Fennia, sub lapidibus, rarissime.

Media inter *H. aterrimam* et sequentem, ab illa thorace canaliculato et minus convexo, ab hac antennis multo longioribus, ab utraque corpore paulo latiore distincta. Caput thorace angustius, rotundatum, basi truncatum, nigrum, nitidum, subtilissime punctulatum. Antennae thorace multo longiores, extrorsum parum incrassatae, nigrae, articulo ultimo oblongo-conico, acuminato. Thorax latitudine plus quam dimidio latior, antice paulo angustior, truncatus, angulis compressis, lateribus et postice rotundatus, supra subconvexus, niger, subnitidus, subtilissime punctulatus, tenuiter sericeo-pubescens, obsolete canaliculatus. Scutellum triangulare, nigrum, subtilissime punctulatum. Elytra thorace vix latiora vel longiora, apice truncata, supra depressa, nigra, subtilissime punctulata, densius pubescentia, subopaca. Abdomen pone elytra reliquo corpore brevius, corpusque subtus nigra, nitida, pubescentia. Pedes fusco-testacei. (Sahlb.)

131. *H. (?) parva*: *Nigra, nitida, vix pubescens, antennis brevibus, tenuioribus, extrorsum vix incrassatis, thorace canaliculato, elytris piceis, pedibus pallidis.* — *Long.* vix  $\frac{2}{3}$  lin.

*Aleoch. parva* Sahlb. *Ins. Fenn. I.* 380. 62.

*Bolitochara parvula* Mannerh. *Brachelytr.* 84. 55.

Habitat in Fennia australi.

Antennis brevioribus, extrorsum vix incrassatis et thorace canaliculato ab *H. aterrima* (fuscula) differt. Caput ut in praecedente, sed palpi piceo-testacei. Antennae longitudine thoracis, totae nigrae, extrorsum vix vel parum omnino incrassatae, articulo ultimo ovato. Thorax transversus, longitudine dimidio latior, antice angustior, truncatus, angulis compressis,



lateribus et postice rotundatus, supra convexus, niger, nitidus, subtilissime punctulatus, canaliculatus. Scutellum triangulare, nigrum, subtilissime punctulatum. Elytra thorace vix latiora, sed eodem paulo longiora, apice truncata, supra depressa, nigro-picea, nitida, crebre punctulata, vix pubescentia. Abdomen pone elytra vix reliqui corporis longitudine, corpusque subtus nigra, nitida, subtilissime punctulata, ano piceo. Pedes testacei. (Sahlb.)

132. *H. (?) pallidula*: Supra luteo-ferruginea, subnitida, oculis nigris, abdominisque cingulo fusco, subtus pallidior, thorace brevi, valde convexo, basi foveolato. — Long. vix  $\frac{1}{2}$  lin.

*Bolitochara pallidula* Mannerh. *Brachelytr.* 84. 56.

*Aleochara pallidula* Sahlb. *Ins. Fenn. I.* 381. 63.

Habitat in Fennia australi.

Praecedente adhuc duplo minor, colore et thorace multo convexiore ab eadem distincta, *H. (?) annulari* quoad colorem affinis, sed differt thorace brevior et latior. Corpus totum, praeter oculos nigros et segmenta abdominis 3 et 4 infusca, supra luteo-ferrugineum, subtus pallidius, subnitidum, tenuiter pubescens. Caput thorace angustius, rotundatum, basi truncatum. (Antennae in typo desunt.) Thorax transversus, longitudine fere duplo latior, antice angustior, truncatus, angulis compressis, lateribus parum rotundatus, basi subtruncatus, supra valde convexus, foveola supra scutellum obsoleta impressus. Scutellum triangulare. Elytra thorace vix longiora vel latiora, apice truncata, supra depressa. Abdomen pone elytra longitudine reliqui corporis. (Sahlb.)

133. *H. (?) tenera*: Nigra, nitida, creberrime punctulata, palpis, segmentorum ventralium marginibus, ano pedibusque piceo-testaceis, elytris nigro-piceis, apice submarginatis, antennis subfiliformibus. — Long. vix 1 lin.

*Aleochara tenera* Sahlb. *Ins. Fenn. I.* 386. 74.

Habitat in Fennia australi, rarissime.

Magnitudine et statura *Myllaenae* dubiae, a qua tamen multis notis differt, ab *H. flavipede*, cui etiam similis est, thorace longiore et angulis eius posterioribus haud acutis satis diversa. Caput thorace multo minus, subtransversum, nigrum, nitidum, subtilissime punctulatum, subpubescens, palpis piceo-testaceis. Antennae thorace paulo longiores, subfiliformes, fusco-nigrae, pubescentes, articulis tribus baseos elongatis, primo incrassato, 2 et 3 obconicis, illo tamen hoc paulo brevior, insequentibus brevibus, subaequalibus, ultimo oblongo-conico, acuminato. Thorax transversus, longitudine dimidio latior, antice angustior, truncatus, angulis deflexis, lateribus rotundatus, basi subtruncatus, angulis posterioribus obtusis, supra convexus, niger, nitidus, creberrime et subtilissime punctulatus, tenuiter pubescens. Scutellum triangulare, nigrum, subtilissime punctulatum. Elytra thoracis basi vix latiora, sed eodem paulo longiora, apice intra angulum exteriorem sinuato-submarginata, supra depressa, nigro-picea, nitida, creberrime punctulato-rugosa, tenuiter pubescentia. Abdomen pone elytra reliquo corpore paulo brevius, nigrum, nitidum, vix punctulatum. Corpus subtus nigrum,

nitidum, subtilissime punctulatum, pubescens, segmentorum ventralium marginibus anoque piceo-testaceis. Pedes mediocres, piceo-testacei. (Sahlb.)

134. *H. (?) impressifrons*: *Nigra, nitida, pedibus piceo-testaceis, capite thorace vix angustiore, vertice obsolete foveolato, thorace brevi, rotundato, antennis thorace longioribus, hirsutis, extrorsum paulo crassioribus.* — Long.  $\frac{3}{4}$  lin.

*Var. Antennarum basi et elytris fusco-testaceis, pedibus pallidioribus.*

*Bolitochara impressifrons* Mannerh. *Brachelytr.* 84. 57.

*Aleoeh. impressifrons* Sahlb. *Ins. Fenn. I.* 387. 75.

Habitat in Fennia australi, rarius.

Statura ad Staphylini genus aliquatenus accedit. Caput thorace vix angustius, rotundatum, basi truncatum, nigrum, nitidum, vix subtilissime punctulatum, vertice foveola plus minusve conspicua impresso. Os et palpi piceo-testacea. Antennae thorace paulo longiores, fusco-nigrae, dense pilosae, articulis baseos tribus piceo-testaceis, elongatis, primo insequentibus crassiore, secundo subcylindrico, tertio obconico, 4 — 10 brevibus, extrorsum sensim crassioribus, ultimo magno, ovato, subacuto. Thorax longitudine latior, antice truncatus, lateribus et postice rotundatus, supra subdepressus, niger, vix subtilissime punctulatus, subpubescens, ante scutellum obsolete foveolatus. Scutellum minutum, triangulare, nigrum. Elytra thorace latiora et nonnihil longiora, subquadrata, apice truncata, supra depressa, nigra, subtilissime punctulata, tenuiter pubescentia. Abdomen pone elytra reliquo corpore brevius, corpusque subtus nigra, nitida, pubescentia. Pedes mediocres, toti piceo-testacei.

*Var. colore pallidior tantum differt. Antennae fuscescentes, basi pallidiores. Thorax nigro-fuscus. Elytra fusco-testacea, et pedes toti pallidi.*

*Obs. Sub diversa sectione, thorace capite vix latiore, suborbiculato et abdomine sublineari distincta, hanc speciem proponit Com. Mannerheim. — (Sahlb.)*

## **Oxypoda Mannerh.**

*Mannerh. Brachelytr. p. 69. — Er. Col. March. I. p. 342.*

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula apice bifida; paraglossae nullae.*

*Palpi labiales 3-articulati, articulo secundo brevior.*

*Tarsi omnes 5-articulati, postici articulo primo elongato.*

Corpus vel oblongum, vel elongatum, minus depressum, alatum. Caput exsertum, saepe deflexum, basi haud constrictum, oculis mediocribus, vix prominulis. Labrum apice leviter rotundatum, mandibulas obtegens. Mandibulae muticae. Maxillae mala exteriori cornea, basi apiceque membranacea, portione membranacea apicali pubescente, mala interiore exteriori paulo brevior, membranacea, apice oblique introrsum emarginata, sinu spinulis brevibus ciliato. Palpi maxillares modice elongati, articulo secundo tertioque longitudine subaequalibus, hoc leviter incrassato, quarto minuto, subulato. Labium mento transverso, apice leviter emarginato, ligula minus brevi, palporum labialium articulum primum excedente, apice bifida, paraglossis nullis. Palpi labiales tri-articulati, articulo primo maiore, secundo hoc duplo minore, tertio te-

nuiore, at primo parum brevior. Antennae articulo secundo tertioque reliquis magis elongatis, inter se aequalibus. Thorax iam antrorsum angustatus, convexus, iam lateribus parallelis. Scutellum triangulare. Elytra apice introrsum oblique truncata, extrorsum ad angulum anteriorem sinuata. Abdomen iam apicem versus angustatum, iam lineare. Pedes leviter elongati, intermedii basi approximati, tarsi omnibus 5-articulatis, postici articulo primo elongato.

#### O b s e r v a t i o n e s.

1. Oris structura omnino ut in Homalota (v. t. III. f. 8. et f. 22.) Distinguitur ab hoc genere tarsi postici articulo primo elongato et imprimis tarsi anterioris 5-articulatis.

2. Genus *Sphenoma* Mannerh. *Brachelytr.* p. 68. ab *Oxypoda* nullo modo distinctum. Auctor illustrissimus errore quodam tarsorum articulos aequales observavit.

3. Mas saepius antennarum validioribus, et apice magis incrassatis, nonnunquam etiam abdominis segmento inferiore sexto apice medio producto distinguitur.

Duas in hoc genere sectiones condidi:

Sectio prima habitu *Tachyporini*, i. e. thorace amplo, convexo, elytrorum basin et caput amplectente, abdomine apicem apicem versus attenuato, apice saepius pilosello, corpore plerumque dense subtiliterque punctato, pube brevissima subtilissimaque sericante obducto. — Victus in fungis et sub foliis deciduis, musco, et in quisquiliis.

Sectio altera corpore lineari, elongato, thorace subquadrato, capite magis exserto, abdomine lineari, punctura et pubescentia minus confertis subtilibusque. — Victus sub arborum cortice.

#### \* F u s i f o r m e s.

\* 1. *O. ruficornis*: *Nigra, nitida, antennis, pedibus, elytris anoque testaceis, thorace parum convexo, postice foveolato obsoleteque canaliculato.* — Long.  $2\frac{1}{2}$  lin.

*Mannerh. Brachelytr.* 69. 1.

*Aleochar. ruficornis* Gyll. *Ins. Suec.* II. 422. 44.

*Oxypoda luteipennis* var. *Er. Col. March.* I. 343. 1.

Habitat in Germania septentrionali, Suecia, Dom. Schüppel et Weber, in Gallia. Mus. Dom. Aubé.

Procera in hoc genere, sequente plus duplo maior, nigra, nitida, pube subtili griseo-sericea vestita. Antennae capite thoraceque paulo longiores, graciles, articulo secundo tertio vix brevior, ultimo oblongo, acuminato, totae rufo-testaceae. Palpi rufo-testacei. Caput deflexum, subtiliter punctatum. Thorax basi coleopteris haud angustior, latitudine dimidio brevior, basi et lateribus leviter rotundatus, antrorsum subangustatus, apice truncatus, angulis omnibus obtusiusculis, anterioribus deflexis, dorso leviter convexus, densius subtiliter punctatus, postice obsolete canaliculatus, supra scutellum profundius impressus, margine laterali planiusculo, rufo-piceo. Elytra thorace longiora, confertim subtiliter punctata, testacea. Abdomen apicem versus attenuatum, confertissime subtilissimeque punctatum, ano segmentorumque singulorum marginibus rufo-testaceis. Pedes dilute testacei, tarsi postici articulis intermediis elongatis, articulo primo sequentibus ambobus aequali.



Mas abdominis segmento ventrali sexto apice triangulariter producto distinctus.

Obs. 1. Revera distincta videtur a sequente, statura multo maiore, thorace minus convexo, antrorsum minus angustato, margine laterali planiusculo, antennis elytrisque unicoloribus, et tarsorum posticorum articulis intermediis elongatis.

2. Aleocharam ruficornem Grav. in genere proxime sequente quaeras.

\* 2. *O. luteipennis*: Nigra, sericeo-pubescent, antennarum basi, pedibus elytrisque testaceis, his circa scutellum lateribusque infuscatis, thorace leviter convexo, postice foveolato obsoleteque canaliculato. — Long.  $1\frac{3}{4}$  lin.

Er. Col. March. I. 343. 1.

*Aleochara ruficornis* var. *b* Gyll. Ins. Suec. II. 422.

*Aleochara ruficornis* Sahlb. Ins. Fenn. I. 388. 76.

Habitat in Germania, Suecia.

O. opaca paulo maior et minus convexa, nigra, parum nitida, pube tenui cinereo-sericea dense vestita. Antennae capitis thoracisque longitudine, apicem versus subincrassatae, articulo secundo tertioque longitudine aequalibus, ultimo apice abrupte compresso, articulo primo testaceo, proximis rufo-piceis, reliquis nigris. Palpi testacei. Caput deflexum, confertum subtiliter punctatum. Thorax basi coleopteris paululum angustior, latitudine dimidio brevior, antrorsum angustatus, basi leviter, lateribus fortius rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, supra leviter aequaliterque convexus, confertim subtiliter punctatus, obsolete canaliculatus, basi foveolatus, limbo laterali rufo-piceo. Elytra thorace longiora, confertim subtiliter punctata, testacea, circa scutellum lateribusque infuscata. Abdomen apicem versus attenuatum, confertissime subtilissimeque punctatum, apice segmentorumque marginibus rufo-testaceis. Pedes testacei, tarsis posticis articulo primo sequentibus tribus coniunctis longitudine aequali.

Mas abdominis segmento ventrali sexto apice utrinque sinuato, medio triangulariter producto distinguendus.

\* 3. *O. prospera*: Subdepressa, rufo-testacea, opaca, capite, pectore abdomineque piceis, hoc apice segmentorumque marginibus rufo-testaceis, thorace parum convexo, basi obsolete foveolato. — Long.  $1\frac{3}{4}$  lin.

Habitat in Germania boreali, Dom. Schüppel.

Statura ab affinibus paululum discedens, Homalotae similis, H. brunnea paulo brevior et minus depressa, at huius generis. Corpus rufo-testaceum, pube depressa grisea subsericea densius vestitum. Antennae capite thoraceque paulo breviores, minus tenues, articulo secundo tertioque subaequalibus, apicem versus leniter subincrassatae, articulo ultimo ovato, apice subacuminato, totae rufo-testaceae. Palpi testacei. Caput crassiusculum, deflexum, confertim subtilissime punctatum, piceum. Thorax coleopteris perparum angustior, latitudine dimidio brevior, antrorsum vix angustatus, basi leviter, lateribus fortius rotundatus, apice submarginatus, angulis omnibus obtusis, subrotundatis, leviter transversim convexus, confertim subtilissime punctatus, basi obsolete impressus, rufo-testaceus, disco picescente, margine dilutiore. Abdomen apicem versus parum angustatum, confertissime subtilissimeque pubescens, nigrum, ano segmentorumque marginibus latis rufo-testaceis. Pectus medio piceum. Pedes testacei.

Mas abdominis segmento ventrali sexto apice obtuse triangulariter producto.

- ✕ \* 4. *O. opaca*: *Brevior, nigra, fusco-pubescent, antennarum basi pedibusque rufis, elytris brunneis, thorace convexo, aequali.* — Long.  $1\frac{1}{2}$  lin.

*Mannerh. Brachelytr.* 70. 4. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 538. 1. — *Er. Col. March. I.* 344. 2.

*Aleochar. opaca* Grav. *Micr.* 89. 31. *Mon.* 163. 41. — *Latr. Hist. nat. Crust. et Ins.* IX. 386. 31. — *Gyll. Ins. Suec. II.* 423. 45. — *Sahlb. Ins. Fenn. I.* 390. 79. — *Zett. Faun. Lapp. I.* 103. 20. *Ins. Lapp.* 79. 29.

*Aleochar. pulla* Grav. *Micr.* 96. 43. *Mon.* 172. 65. — *Latr. Hist. nat. Crust. et Ins.* IX. 389. 43.

Habitat in Europa.

Habitu fere Tachypori, nigra, parum nitida, pube subtili, densa, fusco-sericea oblecta. Antennae capite thoraceque longiores, validiusculae, articulo secundo tertio paulo brevior, ultimo oblongo-ovato, praecedentibus ambobus longiore, nigrae, basi rufo-piceae. Caput confertissime subtilius punctatum. Thorax coleopterorum latitudine, longitudine dimidio lator, antrorsum angustatus, angulis anterioribus subrectis, deflexis, subtiliter confertissime punctatus, limbo laterali rufo-piceo. Elytra thorace paulo longiora, confertissime punctata, undulatum subrugosa, brunnea, interdum piceo-testacea, basi nigricantia. Abdomen breviusculum, apicem versus attenuatum, segmentorum marginibus supra rufo-piceis, subtus testaceis. Pedes testacei.

Mas abdominis segmento ventrali sexto apice triangulariter producto distinctus.

- \* 5. *O. umbrata*: *Elongata, nigra, subnitida, sericeo-pubescent, antennarum basi pedibusque rufo-testaceis, elytris obscure brunneis, thorace convexo, antice subcompresso, lateribus piceo.* — Long.  $1\frac{1}{2}$  lin.

*Mannerh. Brachelytr.* 70 5. — *Er. Col. March. I.* 344. 3.

*Aleochar. umbrata* Gyll. *Ins. Suec. II.* 424. 46. — *Sahlb. Ins. Fenn. I.* 390. 80.

Habitat in Germania, Suecia.

Praecedente parum brevior, at multo angustior, nigra, subnitida, pube tenuissima griseo-sericea vestita, omnium subtilissime confertissimeque punctata. Antennae capitis thoracisque longitudine, articulo secundo tertio subaequali, apicem versus subincrassatae, articulis penultimis tribus transversis, ultimo subovato, apice acuminato, piceae, articulis duobus primis rufo-testaceis. Palpi testacei. Caput thorace angustius, deflexum, rotundatum, fronte convexiuscula. Thorax latitudine paulo brevior, basi coleopterorum latitudine, antrorsum leviter angustatus, basi et lateribus modice rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, fortiter transversim convexus, lateribus praecipue antice deflexis, limbo laterali piceo, interdum rufescente. Elytra thorace paulo longiora, fusco-picea, opaca. Abdomen elongatum, apice attenuatum, segmentis ventralibus ferrugineo-marginatis. Pedes rufo-testacei.

- ✕ \* 6. *O. longiuscula*: *Elongata, nigra, subnitida, tenuiter sericeo-pubescent, antennarum basi pedibusque rufo-testaceis, thorace leviter convexo, basi foveolato.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I.* 345. 4.

*Aleoch. longiuscula* Grav. *Micr.* 80. 20. *Mon.* 155. 21. — *Latr. Hist. nat. Crust. et Ins.* IX. 383. 20.

*Oxypoda procerula* Mannerh. *Brachelytr.* 71. 9.

*Aleoch. procerula* Sahlb. *Ins. Fenn.* I. 392. 82.

Habitat in Germania; in Anglia, Dom. Rudd.

Praecedente minus convexa, nigra, parum nitida, confertissime subtiliter punctata, pube tenuissima griseo-sericea vestita. Antennae capite thoraceque longiores, apicem versus incrassatae, articulo tertio secundo paulo brevior, 4—10 omnibus crassiusculis, subcylindricis, crassitie haud brevioribus, ultimo maris magno, praecedentibus ambobus longiore, oblongo, apice subacuminato, feminae maiusculo, ovato, nigrae, articulis duobus primis piceo-testaceis. Palpi piceo-testacei. Caput thorace angustius, deflexum, rotundatum. Thorax latitudine paulo brevior, basi coleopterorum prope latitudine, antrorsum angustatus, apice duplo fere angustior, basi leviter rotundatus, utrinque subsinuatus, lateribus modice rotundatus, apice truncatus, angulis anterioribus rotundatis, deflexis, posterioribus subrectis, modice transversim convexus, basi foveola transversa impressus. Elytra thorace paulo longiora, picea, interdum cum thorace rufo-picea. Abdomen apice subangustatum, ano late segmentorumque ventralium marginibus rufo-piceis. Pedes rufo-testacei.

\* 7. *O. alternans*: *Elongata, testacea, capite, pectore, elytrorum angulo exteriori abdominisque cingulo postico nigricantibus.* — Long.  $1\frac{1}{2}$  lin.

Mannerh. *Brachelytr.* 71. 8. — Boisd. et Lacord. *Faun. Ent. Paris.* I. 539. 3. — Er. *Col. March.* I. 345. 5.

*Aleoch. alternans* Grav. *Micr.* 85. 26. *Mon.* 163. 38. — *Latr. Hist. nat. Crust. et Ins.* IX. 385. 26. — Gyll. *Ins. Suec.* II. 419. 41. — Sahlb. *Ins. Fenn.* I. 385. 71. — Zetterst. *Faun. Lappon.* I. 103. 20. *Ins. Lappon.* 78. 28.

Habitat in Germaniae, Sueciae boletis.

Statura angusta praecedentium, dilute testacea, confertim subtilissime punctata, pube tenuissima griseo-sericea vestita. Antennae capitis thoracisque longitudine, apicem versus leviter incrassatae, articulo secundo tertioque aequalibus, 5—10 transversis, ultimo praecedentibus tribus coniunctis vix brevior, subovato, apice acuminato, rufae vel rufo-piceae vel nigricantes, basi testaceae. Caput oblongum, deflexum, subtiliter punctulatum, nigricans aut piceum, ore testaceo. Thorax coleopteris paulo angustior, latitudine baseos dimidio brevior, antrorsum angustatus, apice dimidio angustior, basi et lateribus modice rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, transversim leviter convexus, confertim subtiliter punctatus, basi foveola parva impressus. Elytra thorace sesqui longiora, confertim subtilius punctata, fusco-testacea, circa scutellum et in angulo apicis exteriore fusciscentia vel nigricantia. Abdomen apicem versus sensim attenuatum, confertissime subtiliter punctatum, segmento quarto quintoque basi nigricantibus. Pectus nigricans.

Variat thoracis disco anteriore infuscato.

\* 8. *O. togata*: *Elongata, nigra, capite piceo, thorace elytrisque rufis, antennarum basi pedibusque testaceis.* — Long.  $1\frac{2}{3}$  lin.

Er. *Col. March.* I. 346. 8.

Habitat Berolini.



Statura omnino *O. umbratae*. Corpus pube sericea subtilissima et minus quam in affinis obvia. Antennae capitis thoracisque longitudine, apicem versus leniter incrassatae, articulo secundo tertio paulo maiore, articulis penultimis vix transversis, ultimo oblongo-ovato, praecedentibus ambobus paulo longiore, apice acuminato, piceae, basi testaceae. Palpi testacei. Caput deflexum, piceum, nitidum, dense punctatum. Thorax latitudine paulo brevior, basi coleopterorum latitudine, antrorsum angustatus, basi leviter, lateribus fortius rotundatus, apice truncatus, angulis anterioribus obtusis, subrotundatis, deflexis, posterioribus rotundatis, transversim convexus, subtilissime confertim punctatus, rufus. Elytra thorace paululum longiora, confertissime subtilius punctata, rufa. Abdomen elongatum, fere parallelum, apice angustatum, confertissime subtilius punctatum, nigrum, nitidum, segmentis piceo-marginatis. Pectus nigrum. Pedes rufo-testacei.

\* 9. *O. sagulata*: *Elongata, nigra, capite piceo, thorace basi foveolato elytris rufis, antennis pedibusque testaceis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Pensylvania, Dom. Zimmermann.

Praecedenti proxima. Corpus pube brevissima, vix observanda, fulvescente vestitum. Antennae capitis thoracisque longitudine, apicem versus leniter incrassatae, articulo secundo tertio paulo maiore, sequentibus sensim paulo crassioribus, 7—10 leviter transversis, ultimo maiusculo, ovato, apice obtuso, testaceae. Caput deflexum, subtilissime punctulatum, nigro-piceum, ore rufo-piceo. Thorax latitudine dimidio brevior, basi coleopteris latitudine aequalis, antrorsum angustatus, basi perparum, lateribus modice rotundatus, apice truncatus, angulis omnibus obtusis, subrotundatis, anterioribus deflexis, transversim leviter convexus, subtilissime punctulatus, basi plerumque obsolete foveolatus, rufus. Elytra thorace paulo longiora, confertissime distinctius punctata, thoraci concolora. Abdomen elongatum, subparallelum, apice attenuatum, supra creberrime minus subtiliter punctatum, subnitidum, nigrum, segmentorum marginibus apiceque obscure rufis. Pedes rufo-testacei.

Obs. Praecedente paulo minor, antennarum articulo ultimo apice obtuso, thorace brevior, basi foveolato sat distincta.

\* 10. *O. abdominalis*: *Elongata, pallide rufa, griseo-sericea, thorace convexo, coleopteris latiore, abdomine nigro, segmentorum marginibus apiceque rufis.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 346. 7.*

*Sphenoma abdominale* Mannerh. *Brachelytr.* 69. 1.

*Aleochara abdominalis* Sahlb. *Ins. Fenn. I. 93. 83.*

Habitat in Germania, Fennia.

Dilute rufa, subnitida, pube griseo-sericea distinctiore vestita. Antennae capitis thoracisque longitudine, apicem versus leniter incrassatae, articulo secundo tertio subaequali, articulis 7—10 leniter transversis, ultimo praecedentibus ambobus aequali, subovato, apice obtusiusculo, rufo-testaceae, basi dilutiores. Palpi testacei. Caput subtilissime punctulatum, deflexum, subrotundatum, rufo-piceum. Thorax basi latitudine coleoptera excedens,

longitudine sesqui lator, antrosum angustatus, basi perparum, lateribus leniter rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, transversim convexus, subtilissime punctulatus. Elytra thorace vix longiora, confertissime subtiliter punctata. Abdomen apicem versus sensim attenuatum, confertissime subtiliter punctatum, nigrum, segmentorum marginibus apiceque rufis. Pectus rufum. Pedes testacei.

- \* 11. *O. testacea*: *Elongata, rufo-testacea, subtiliter sericeo-pubescent, thorace brevi, leviter convexo, abdomine distinctius punctato, segmentis intermediis basi fusciscentibus.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I.* 347. 8.

Habitat in Germania septentrionali.

Praecedentibus aequalis, at paulo magis linearis, dilute rufo-testacea, pube tenui griseo-sericea vestita. Antennae capite thoraceque breviores, crassiusculae, articulo secundo tertio paulo maiore, 4—10 brevibus, fortiter transversis, penultimis longitudine plus duplo crassioribus, ultimo maiusculo, subgloboso, apice obtuso. Caput subtilissime punctatum. Thorax latitudine coleopteris aequalis eademque vix dimidio brevior, antrosum angustatus, basi subtruncatus, lateribus leviter rotundatus, apice truncatus, angulis anterioribus obtusis, deflexis, posterioribus subrectis, transversim convexus, subtilissime punctatus. Elytra thorace vix longiora, confertissime subtilius punctata. Abdomen elongatum, parallelum, apice subangustatum, confertissime subtilius punctatum, obscure rufo-testaceum, segmentis intermediis basi nigricantibus, apice pallidiore.

- \* 12. *O. helvola*: *Elongata, linearis, testacea, sericeo-pubescent, thorace leviter convexo, elytris thorace subbrevioribus, abdomine segmento quinto basi nigricante.* — Long. 1 lin.

*Er. Col. March. I.* 347. 9.

Habitat in Germania.

Statura angusta praecedentium, at multo minor, laete rufo-testacea, subtilissime confertim punctata, pube tenui flavescenti-sericea densius vestita. Antennae capite thoraceque breviores, crassiusculae, articulo secundo tertio paulo longiore, 4—10 transversis, ultimo subgloboso, obtusiusculo. Thorax coleopterorum latitudine, longitudine sesqui lator, antrosum angustatus, lateribus et basi leviter rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, transversim convexus. Elytra thorace paulo breviora, minus subtiliter punctata. Abdomen elongatum, lineare, apice angustatum, segmento quinto basi nigricante.

- \* 13. *O. ferruginea*: *Elongata, linearis, ferruginea, tenuiter pubescent, thorace leviter convexo, elytris thorace subbrevioribus, depressis, abdomine nigro, apice testaceo.* — Long. 1 lin.

Habitat in Germania.

Praecedenti affinis, at paulo angustior et magis depressa, pube subtili depressa griseo-pubescent. Antennae capite thoraceque paulo breviores, apicem versus leviter sensim incrassatae, articulo secundo tertio duplo maiore,

4 — 10 longitudine subaequalibus, sensim paulo crassioribus, transversis, ultimo praecedentibus ambobus fere aequali, oblongo-ovato, apice acuminato, pubescentes, ferrugineae, basi testaceae. Palpi testacei. Caput thorace paulo angustius, oblongum, deflexum, confertim subtiliter punctulatum, obscure ferrugineum. Thorax coleopteris paulo latior, basi leviter, lateribus fortius rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus subrotundatis, deflexis, transversim leviter convexus, confertim subtiliter punctatus, basi foveola minuta obsolete impressus, ferrugineus, subnitidus. Elytra thorace paulo breviora, fortiter depressa, crebre minus subtiliter punctata, ferruginea. Abdomen elongatum, lineare, apice attenuatum, confertim subtiliter punctatum, nigrum, subnitidum, segmentis singulis piceo-marginatis, quinto apice late sextoque toto rufo-testaceis. Pectus piceum. Pedes testacei, coxis omnibus concoloribus.

Obs. A praecedente statura tenuiore, antennis paulo longioribus, minus incrassatis, elytris depressis et colore obscuriore distincta videtur.

\* 14. *O. promiscua*: *Elongata, sublinearis, picea, capite abdomineque nigris, ano, antennarum basi pedibusque testaceis, thorace leviter convexo, aequali, elytris thorace paulo longioribus.* — Long. 1 lin.

Habitat in Germania; Berolini a Dom. Schüppel lecta.

Statura et summa affinitas proxime praecedentium, elytris autem longioribus satis distincta. Parva, linearis, nitida, pube minus subtili depressa grisea densius vestita. Antennae capite thoraceque paulo breviores, apicem versus sensim incrassatae, articulo tertio secundo dimidio minore, 4 — 10 longitudine aequalibus, brevibus, sensim crassioribus et fortius transversis, ultimo praecedentibus ambobus longitudine aequali, crassiore, acuminato, fuscae, basi testaceae. Caput thorace paulo angustius, oblongum, deflexum, subtilissime punctatum, nigrum, nitidum, ore testaceo. Thorax basi coleopterorum prope latitudine, apicem versus leviter angustatus, latitudine dimidio brevior, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus rotundatis, fortiter deflexis, posterioribus obtusis, transversim convexus, aequalis, densius subtilissime punctatus, fusco-piceus, nitidus. Elytra thorace paulo longiora, minus subtiliter confertim punctata, dilutius fusco-picea. Abdomen elongatum, apicem versus leviter angustatum, confertim subtiliter punctatum, nigrum, nitidum, segmentis singulis piceo-marginatis, quinto apice late sextoque toto lacte rufo-testaceis. Pectus piceum. Pedes testacei.

\* 15. *O. praecox*: *Testacea, immaculata, thorace transversim convexo, aequali, elytris thorace paulo longioribus, abdomine apicem versus sensim fortius attenuato, nigro-pilosello.* — Long. 1 lin.

Habitat in Austria, Dom. Schüppel.

Habitu omnino *O. abdominalis*, at multo minor, *O. helvolae* aequalis, tota dilute testacea, subnitida, pube subtili tenuique fulvescente subsericans. Antennae capite thoraceque paulo breviores, apicem versus sensim fortius incrassatae, articulo tertio secundo duplo brevior et paulo tenuiore, 4 — 10 breviusculis, longitudine aequalibus, at sensim paulo crassioribus et fortius transversis, ultimo praecedente sesqui longiore, crassiusculo, subcom-



presso, apice truncato. Caput thorace plus duplo angustius, oblongum, deflexum, confertim subtiliter punctatum, oculis nigris. Thorax basi coleopterorum latitudine, antrorsum leviter angustatus, apice dimidio quam basi angustior, latitudine baseos dimidio brevior, basi et lateribus modice rotundatus, apice truncatus, angulis anterioribus obtusis, subrotundatis, deflexis, posterioribus subrectis, transversim convexus, aequalis, confertim subtilius punctatus. Elytra thorace paulo longiora, confertim minus subtiliter punctata. Abdomen longiusculum, apicem versus sensim sat distincte attenuatum, confertim subtiliter punctatum, lateribus praecipue versus apicem breviter nigropilosellum. Pedes flavo-testacei.

Obs. *O. helvolae* magnitudine et colore similis, distincta autem corpore non lineari sed fusiformi, elytris longioribus et abdomine pilosello.

- \* 16. *O. exoleta*: *Elongata, testacea, sericeo-pubescent, thorace leviter convexo, elytris thorace longioribus, abdomine ante apicem nigricante.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Germania.

Statura elongata *O. helvolae*, at paulo maior et minus linearis, sordide testacea, parum nitida, pube brevissima densa griseo-fulvescente sericans. Antennae capite thoraceque paulo breviores, apicem versus leviter sensim incrassatae, articulo tertio secundo dimidio minore, 4—10 breviusculis, longitudine inter se aequalibus, at sensim paulo crassioribus et magis transversis, ultimo praecedentibus ambobus longitudine subaequali, apice rotundato, obtuso, fusco-testaceae, articulo primo flavo-testaceo. Caput thorace paulo angustius, oblongum, deflexum, confertim subtiliter punctatum, fusco-testaceum, ore flavescens. Thorax basi coleopterorum latitudine, antrorsum angustatus, apice basi tertia parte angustiore, latitudine baseos plus dimidio brevior, basi leviter, lateribus vix rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, transversim convexus, aequalis, confertim subtiliter punctatus, rufescenti-testaceus. Elytra thorace paulo longiora, confertim subtilius punctata, fusciscenti-testacea. Abdomen elongatum, apicem versus sensim leniter attenuatum, confertim subtiliter punctatum, rufotestaceum, segmento quarto toto, quinto basi nigricantibus, quinto apice late sextoque toto laete rufo-testaceis. Pectus rufo-testaceum. Pedes flavotestacei.

- \* 17. *O. cuniculina*: *Brevior, nigra, sericeo-pubescent, thorace breviusculo, leviter convexo elytris piceis, abdomine apicem versus attenuato, segmentis piceo-marginatis.* — Long.  $1\frac{1}{4}$  lin.

*Er. Col. March. I.* 348. 10.

Habitat in Germania: Berolini, Dom. Schüppel, in Bavaria, Dom. Waltl, in Austria, Dom. Ullrich.

*O. umbratae* affinis, at multo brevior, nigra, subtilissime confertissimeque punctata, pube tenui griseo-sericea obducta. Antennae capite thoraceque paulo breviores, apicem versus sensim modice incrassatae, articulo secundo subcylindrico, tertio obconico, secundo paulo brevior, 3—10 arcte contiguis, 6—10 transversis, ultimo maiusculo, ovato, obtuso, piceae, basi dilutiones. Thorax latitudine duplo brevior, basi coleopterorum latitudine antrorsum angustatus, basi et lateribus rotundatus, apice truncatus, angulis omnibus ob-

tusis, anterioribus deflexis, transversim convexus, supra scutellum obsolete impressus, piceus, disco nigricans. Elytra thorace paulo longiora, rufo-picea. Abdomen apicem versus attenuatum, nigrum, segmentis margine rufo-piceis. Pedes testacei.

- \* 18. *O. exigua*: *Nigra, nitidula, thorace leviter convexo elytris piceis, antennarum basi pedibusque testaceis, abdomine distinctius punctulato.*  
— Long. 1 lin.

*Er. Col. March. I.* 348. 11.

Habitat Berolini.

Habitu omnino praecedentis, at minuta, minus convexa, nitidula, nigra, thorace elytris pube tenuissima sericeo-micantibus. Antennae fere capitis thoracisque longitudine, articulo secundo tertio paulo longiore, apicem versus subincrassatae, articulis 7—10 transversis, ultimo maiusculo, breviter ovato, obtuso, nigrae, basi flavescentes. Palpi piceo-testacei. Caput deflexum, rotundatum, subtilissime punctatum. Thorax latitudine plus dimidio brevior, coleopterorum latitudine, antrorsum parum angustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis posterioribus subrectis, anterioribus rotundatis, leviter transversim convexus, subtiliter punctulatus, nigro-piceus. Elytra thoracis longitudine, picea, subtiliter dense punctata. Abdomen elongatum, apice angustatum, distinctius vage punctatum, pilis longis depressis parce vestitum, apice pilosellum. Pedes rufo-testacei.

- \* 19. *O. cunctans*: *Brevis, nigra, opaca, sericeo-pubescent, antennarum basi pedibusque rufo-testaceis, thorace brevi, leviter convexo, postice leviter bisinuato, abdomine supra confertissime subtiliter punctato.* — Long.  $1\frac{1}{3}$  lin

*Er. Col. March. I.* 349. 12.

Habitat Berolini.

Praecedentibus brevior, latior et supra minus convexa, nigra, opaca, pube tenui grisescenti-sericea obducta. Antennae capitis thoracisque longitudine, apicem versus leviter incrassatae, articulo secundo tertio paulo crassiore, 4—10 cylindricis, crassiusculis, crassitie autem vix brevioribus, ultimo oblongo, praecedentibus ambobus paulo longiore, apice obtuse acuminato, piceae, basi testaceae. Palpi testacei. Caput subtilissime confertim punctatum, fronte convexa. Thorax coleopterorum latitudine, antrorsum parum angustatus, longitudine duplo latior, lateribus rotundatus, basi utrinque leviter sinuatus, apice truncatus, angulis omnibus obtusis, subrotundatis, parum convexus, subtilissime dense punctatus, supra scutellum leviter impressus. Elytra thorace sesqui longiora, subtilissime dense punctata. Abdomen apicem versus leviter angustatum, confertim subtiliter punctatum, subtus apice et marginibus segmentorum ventralium singulorum late rufo-piceis. Pedes testacei.

- \* 20. *O. lentula*: *Brevior, nigra, opaca, sericeo-pubescent, antennarum pedibusque piceis, thorace transverso, parum convexo, basi foveola transversa impresso.* — Long.  $1\frac{1}{4}$  lin.

*Er. Col. March. I.* 349. 13.

Habitat Berolini.

Praecedente paulo minor, pube tenui griseo-sericea vestita, confertissime subtilissimeque punctata, opaca, vel tota nigra, vel thorace elytrisque piceis. Antennae capitis thoracisque longitudine, articulo secundo tertioque subaequalibus, sequentibus leviter transversis, sensim paulo crassioribus, ultimo praecedentibus ambobus subaequali, subovato, apice obtusiusculo, apicem versus parum incrassatae, nigro-piceae, vel piceae, basi dilutiores. Caput deflexum, fronte convexa, oris partibus inferioribus testaceis, palpis picescentibus. Thorax coleopterorum latitudine, antrorsum parum angustatus, latitudine duplo fere brevior, basi leviter, lateribus fortiter rotundatus, apice truncatus, angulis anterioribus rotundatis, posterioribus obtusis, subrotundatis, transversim convexus, basi foveola transversa impressus. Elytra thorace parum longiora. Abdomen breviusculum, apicem versus parum angustatum, apice piceum. Pedes picei.

Obs. A praecedente praecipue elytris brevioribus abdomineque subtilius punctatis distincta.

\* 21. *O. curtula*: Brevis, nigra, subnitida, distinctius punctata, subtilissime parce pubescens, antennis pedibusque ferrugineis, thorace brevi, transverso, basi leviter bisinuato. — Long. 1 lin.

Er. Col. March. I. 350. 14.

Habitat Berolini.

Aleocharae morioni similis, at huius generis videtur. Brevis, nigra nitidula, dense ac minus subtiliter punctata, pube subtilissima vix observanda obducta. Antennae capitis thoracisque longitudine, articulo secundo tertio paulo maiore, praecipue crassiore, apicem versus sensim incrassatae, articulis penultimis transversis, ultimo maiusculo, subgloboso, rufo-piceae, basi testaceae. Caput breviusculum, subtilius punctatum. Thorax basi coleoptera latitudine aliquantulum superans, longitudine duplo latior, antrorsum angustatus, lateribus rotundatus, basi utrinque leviter sinuatus, apice truncatus, angulis anterioribus obtusis, subrotundatis, posterioribus rectis, retrorsum leviter prominulis, supra parum convexus, confertissime punctatus, basi medio obsolete impressus, margine laterali haud deflexo. Elytra thorace fere duplo longiora et paulo fortius punctata, apice medio conjunctim emarginata. Abdomen apicem versus leviter angustatum, confertissime punctatum. Pedes testacei, femoribus nigricantibus.

Obs. Specimen unicum coleopteris utrinque prope scutellum tuberculo minuto prominulo notatis insigne: num proprio speciei an' sexui?

Species mihi invisae, forte huius generis et sectionis.

22. *O. melanaria*: Nigra, nitida, subpubescens, subtilissime punctulata, elytris piceo-testaceis, segmentorum ventralium marginibus, ano pedibusque rufo-testaceis, antennis filiformibus, thorace foveola postica obsolete impresso. — Long.  $1\frac{1}{4}$  lin.

Mannerh. Brachelytr. 70. 3.

*Aleochara melanaria* Sahlb. Ins. Fenn. I. 389. 78.

Habitat in Fennia.

Magnitudo Homalotae lividipennis, sed convexior, nitidior, levius punctata, antennae longiores et thorax antice minus angustatus. Caput thorace



multo minus, rotundatum, convexum, nigrum, nitidum, parce pubescens, subtilius adhuc quam in *Hom. lividipenni* et vix visibiliter punctulatum, ore et palpis piceo-rufescentibus. Antennae thorace multo longiores, filiformes, nigrae, articulo primo elongato, obconico, secundo brevior et ceteris fere angustiore, tertio iterum elongato, obconico, quarto insequentibus paulo brevior, 5—10 longitudine et latitudine inter se aequalibus, ultimo elongato, acuminato. Thorax antice paulo angustior, truncatus, lateribus et postice rotundatus, supra convexus, niger, nitidus, subtilissime et parcus punctulatus, tenuiter pubescens, foveola ante scutellum obsoletissime impressus. Scutellum breve, triangulare, nigrum, punctulatum. Elytra thorace vix latiora, sed paulo longiora, apice intra angulum anteriorem emarginata, supra depressa, piceo-testacea, subtilissime et crebre punctulata, subpubescentia. Abdomen pone elytra reliquo corpore longius, nigrum, subnitidum, subtilissime punctulatum, subpubescens. Corpus subtus fusco-nigrum, subnitidum, subtilissime punctulatum, densius pubescens, marginibus segmentorum ventralium anoque rufo-testaceis. Pedes elongati, tenues, rufo-testacei, articulo tarsorum primo insequente paulo longiore. (Sahlb.)

23. *O. pellucida*: Nigro-fusca, pubescens, antennarum basi, thorace, ano, marginibus segmentorum pedibusque piceo-ferrugineis, thoracis depressiusculi lateribus testaceis, angulo elytrorum exteriori infusato. — Long.  $1\frac{1}{4}$  lin.

*Mannerh. Brachelytr.* 70. 6. — *Runde Brachelytr. Hal.* 29. 1.

*Aleochar. pellucida* Sahlb. *Ins. Fenn. I.* 391. 81.

Habitat in fungis Fenniae.

*O. opacae* affinis, sed brevior, latior, caput magis prominulum, pedes postici minus elongati, abdomen parum attenuatum, colorque alius. Caput thorace multo minus, rotundatum, convexiusculum, fusco-nigrum, nitidum, subtilissime punctulatum, palpis testaceis. Antennae longitudine thoracis, piceo-ferrugineae, extrorsum fusciscentes, articulo primo elongato, incrassato, secundo et tertio itidem elongatis, obconicis et inter se aequalibus, 4—10 brevibus, extrorsum sensim crassioribus, ultimo magno, conico-ovato, subacuto. Thorax transversus, longitudine fere duplo latior, antice angustior, truncatus, angulis compressis, lateribus et postice rotundatus, angulis tamen parum prominulis, supra subdepressus, obsolete canaliculatus, piceo-ferrugineus, subtilissime punctulatus, pubescens, lateribus pellucido-testaceis. Scutellum minutum, triangulare, fusciscentes. Elytra thorace paulo latiora et longiora, apice subtruncata, intra angulum anteriorem vix sinuata, supra depressa, piceo-ferruginea vel ferrugineo-testacea, angulis apicis externis, interdum etiam sutura infuscatis, subtilissime punctulata, pubescentia. Abdomen pone elytra reliquo corpore brevius corpusque subtus nigro-fusca, pubescentia, marginibus segmentorum, imprimis subtus, anoque ferrugineis. Pectus distinctius punctatum. Pedes mediocres, piceo-ferruginei, interdum testacei; articulus tarsorum primus insequente paulo longior. (Sahlb.)

24. *O. lateralis*: Nigra, punctulata, pubescens, antennis thorace longioribus basi, thorace lateribus, humeris, marginibus segmentorum ventralium pedibusque rufo-testaceis. — Long.  $1\frac{1}{4}$  lin.

*Mannerh. Brachelytr.* 70. 7.

*Aleochar. lateralis* Sahlb. *Ins. Fenn. I.* 391. 82.

Habitat in fungis Fenniae.

Praecedenti simillima, differt autem statura angustiore, antennis thorace longioribus, pubescentia densiore, elytris obscurioribus, ad angulum exterioriorem evidentius sinuatis et ano nunquam rufo. *O. opacae* etiam affinis, sed caput maius, prominulum, pedes postici minus elongati, abdomen parum tantum attenuatum. Caput ut in praecedente. Antennae thorace paulo longiores, extrorsum incrassatae, nigrae, articulis tribus baseos elongatis, rufo-testaceis, ultimo magno, conico-ovato, subacuto. Thorax nigro-fuscus, lateribus rufo-testaceis, foveola ante scutellum obsolete impressus, structura ut in praecedente. Scutellum triangulare, nigrum, punctulatum, pubescens. Elytra thorace paulo longiora et latiora, apice intra angulum exterioriorem nonnihil sinuata, supra depressa, fusca, subtilissime punctulata, pubescentia, humeris indeterminate rufo-testaceis. Abdomen pone elytra reliquo corpore brevius corpusque subtus nigra, subtilissime punctulata, pubescentia, marginibus segmentorum ventralium rufo-testaceis. Pedes mediocres, rufo-testacei, articulo tarsorum primo insequente paulo longiore. (Sahlb.)

25. *O. sericata*: Fusca, opaca, subtiliter punctulata, sericeo-pubescent, antennarum basi, pedibus anoque pallidioribus.

Mannerh. *Brachelytr.* 71. 10.

Habitat Petropoli.

*O. umbratae* quodammodo similis, magnitudine duplo minore, capite vix occulto, antennis longioribus, punctura subtiliore, pubescentia densiore ut et colore diversa. (Mannerh.)

26. *O. cingulata*: Livido-testacea, subtilissime punctata, griseo-pubescent, capite abdominisque cingulo nigri, thorace basi foveolato.

Mannerh. *Brachelytr.* 71. 11.

Habitat Petropoli.

Praecedenti iterum statura et magnitudine fere aequalis, antennae vero breviores, articulo ultimo magis globoso, thorax antice angustior, postice foveolatus, et color diversus. (Mannerh.)

## \*\*\* Lineares.

- \*27. *O. corticina*: Piceo-rufa, abdomine ante apicem capiteque piceis, ano, antennis pedibusque ferrugineis, thorace transverso, subdepresso, angulis posterioribus rectis. — Long.  $1\frac{1}{3}$  lin.

Er. Col. March. I. 351. 15.

Habitat in Germania sub cortice Pinus.

Piceo-rufa, parum nitida, pube tenuissima fusciscenti densius oblecta. Antennae capite thoraceque vix breviores, apicem versus subincrassatae, articulis 4—10 transversis, ultimo ovato, subacuminato, ferrugineae. Caput piceum, subtilissime punctatum, fronte convexa, ore ferrugineo. Thorax coleoptera latitudine fere excedens, longitudine plus sesqui latior, antrorsum vix angustatus, basi medio rotundatus, utrinque subsinuatus, lateribus modice rotundatus, apice late emarginatus, angulis anterioribus acutiusculis, leviter deflexis, posterioribus subrectis, leviter transversim convexus, confertissime

subtilius punctatus. Elytra rufa, fortius punctata. Abdomen parallelum, rufo-piceum, ante apicem nigricans, apice ferrugineum. Pectus fortius punctatum, piceum. Pedes ferruginei.

28. *O. amicta*: *Nigra, opaca, antennis, elytris, ano pedibusque laete rufis, thorace apicem versus subangustato. — Long. 1 lin.*

Habitat in Sardinia, Prof. Gené, Mus. Reg. Taurin.

Praecedenti affinis, at gracilior, nigra, opaca, tota sat fortiter confertissime punctata, pube subtili depressa grisea vestita. Antennae capitis thoracisque longitudine, validiusculae, articulo secundo tertio vix brevior, apicem versus leviter incrassatae, articulis penultinis vix transversis, ultimo oblongo, praecedentibus ambobus paulo longiore, apice obtusiusculo, totae rufo-ferrugineae. Palpi rufo-ferruginei. Caput thorace multo minus, deflexum. Thorax latitudine vix dimidio brevior, coleopterorum latitudine, basi medio parum rotundatus, utrinque subsinuatus, lateribus ultra medium rectis et parallelis, dein apicem versus rotundato-angustatis, apice truncatus, angulis anterioribus obtusis, deflexis, posterioribus rectis, parum convexus, margine laterali anguste rufo. Elytra thorace longiora, laete rufa. Abdomen apicem versus sensim subangustatum, apice segmentorumque singulorum margine tenui apicali rufo-testaceis. Pedes graciles, rufo-testacei.

- \* 29. *O. analis*: *Rufo-picea, abdomine ante apicem nigricante, apice, antennis pedibusque ferrugineis, thorace transverso, subdepresso, angulis posterioribus obtusis. — Long. 1½ lin.*

*Aleochar. analis* Gyll. II. 388. 11. — *Sahlb. Ins. Fenn. I. 354. 16.*

Habitat in Suecia, Dom. Schüppel.

Corpus lineare, rufo-piceum vel piceo-rufum, opacum, pube subtili depressa flavescenti-grisea parcius vestitum. Antennae capitis thoracisque vix longitudine, articulo secundo tertio subaequali, apicem versus sensim incrassatae, articulis 5 — 10 transversis, ultimo brevior, apice acuminato, totae rufo-ferrugineae. Caput thorace parum angustius, sublaeve, fusco-piceum, ore attenuato, testaceo. Thorax latitudine dimidio brevior, coleopterorum latitudine, lateribus ante medium modice rotundatus, basin versus subangustatus, basi leviter rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, posterioribus obtusis, parum convexus, confertim subtiliter punctulatus, basi leviter impressus. Elytra thorace sesqui longiora, confertim minus subtiliter punctata. Abdomen lineare, crebre minus subtiliter punctulatum, ante apicem nigro-piceum, apice ipso ferrugineo. Pedes ferruginei.

Obs. Intermedia quasi inter *O. corticinam* et *prolixam*, ab illa statura magis elongata, thorace minus lato, angulis posterioribus obtusis, ab hac statura magis depressa et punctura multo subtiliore distincta.

- \* 30. *O. gracilis*: *Elongata, rufo-testacea, abdomine ante apicem, capite elytrisque picescentibus, thorace basin versus angustato, canaliculato, margine subreflexo. — Long. 1½ lin.*

*Er. Col. March. I. 351. 16.*

Habitat Berolini, Dom. Weber.



Statura admodum gracili, antennis pedibusque elongatis insignis, rufo-testacea, parum nitida, pube subtili, brevi, flavescente leviter vestita. Antennae capite thoraceque multo longiores, apicem versus perparum incrassatae, articulo secundo tertio paulo brevior, 4 — 10 subcylindricis, sensim paulo brevioribus crassioribusque, decimo ipso crassitie paulo longiore, ultimo oblongo, praecedentibus ambobus paulo longiore, apice acuminato-pallidoque. Caput dilute piceum, confertim subtiliter punctatum, fronte supra antennarum basin obsolete impressa, ore testaceo. Thorax coleopteris prope angustior, latitudine fere dimidio brevior, lateribus rotundatus, praesertim ante medium, basin versus angustatus, basi leviter rotundatus, apice late emarginatus, angulis posterioribus obtusis, anterioribus acutis, prominulis, supra subdepressus, confertim subtiliter punctatus, obsolete canaliculatus, margine laterali acuto, subreflexo. Elytra thorace sesqui longiora, subpicescentia, fortius punctata. Abdomen elongatum, parallelum, confertim subtiliter punctatum, ante apicem supra picescens. Pedes elongati, tenues.

Obs. Habitu fere Tachysuae, at huius generis videtur, et in hac sectione collocanda.

- \* 31. *O. proluxa*: *Elongata, rufo-ferruginea, nitida, fortius punctata, ab domine ante apicem capiteque piceis, ano, antennis pedibusque ferrugineis, thorace transversim quadrato, convexiusculo, basi foveolato. — Long. 1½ lin.*

*Er. Col. March. I. 352. 17.*

*Aleoch. proluxa Grav. Micr. 71. 5. Mon. 165. 45. — Latr. Hist. nat. Crust. et Ins. IX. 379. 5.*

Habitat in Germania, Gallia, Suecia.

*O. corticinae* coloribus similis, at nitidior, magis elongata, fortius punctata, pube brevi, parum depressa, flavescente tenuiter vestita. Antennae capite thoraceque breviores, apicem versus modice incrassatae, articulo secundo tertioque subaequalibus, 7 — 10 transversis, ultimo praecedentibus ambobus longitudine aequali, ovato, apice subacuminato, ferrugineae. Caput thorace parum angustius, deflexum, vage punctatum, piceum, ore attenuato, ferrugineo. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus modice rotundatus, basin versus subangustatus, basi leviter rotundatus, apice leviter emarginatus, angulis posterioribus obtusis, anterioribus deflexis, subrectis, modice convexus, confertim minus subtiliter punctatus, supra scutellum transversim impressus. Elytra thorace sesqui longiora, dense fortiter punctata, rufa, circa scutellum et lateribus infuscata. Abdomen parallelum, crebre punctatum, rufum, ante apicem nigricans, apice ferrugineum. Pedes ferruginei.

Obs. Specimen e Suecia meridionali, a Dom. Schönherr nom. *Al. obfuscatae* communicatum dedit Dom. Schöppel.

- \* 32. *O. ruficollis*: *Elongata, nigro-picea, nitida, thorace transverso, basin versus subangustato, rufo, ano, pedibus antennisque ferrugineis, his medio infuscat. — Long. 1½ lin.*

Habitat Brunsvigae, Knoch, in Hercynia, sub cortice Abietis, Dom. Saxesen, Lutetiae, Mus. Dom. Aubé, in Anglia, Dom. Rudd.

Præcedenti affinis, at dimidio minor. Nigro-picea, nitida, parce pubescens. Antennae capite thoraceque breviores, apicem versus incrassatae, articulo secundo tertio paulo maiore, 7—10 sensim magis transversis crassioribusque, ultimo subgloboso, ferrugineae, ante apicem fuscae. Caput thoracis fere latitudine, parce subtiliter punctatum, ore attenuato, testaceo. Thorax coleopteris haud angustior, latitudine dimidio brevior, lateribus ante medium leviter rotundatis, basin versus subangustatus, basi leviter rotundatus, apice vix emarginatus, angulis anterioribus rectis, deflexis, posterioribus subrotundatis, supra leviter convexus, aequalis, parce subtilissime punctatus, rufus. Elytra thoracis longitudine, nigra, nitida, minus dense fortiter punctata, medio subimpressa. Abdomen elongatum, parallelum. vel apicem versus subincrassatum, laeve, ano ferrugineo. Pedes ferruginei.

\* 33. *O. fumida*: Nigra, subnitida, antennarum basi, ore pedibusque testaceis, ano rufo, thorace basin versus subangustato, basi foveolato. — Long.  $1\frac{1}{3}$  lin.

Habitat Lutetiae, Dom. Aubé, in Thuringia, Dom. Kellner.

Statura omnino praecedentis, *O. corticinae* aequalis, nigra, minus nitida, pube grisea canescens. Antennae capitis thoracisque prope longitudine, apicem versus leniter incrassatae, articulo secundo tertio paulo maiore, 5—10 sensim paulo crassioribus, penultimis transversis, ultimo breviter ovato, subacuminato, fuscae, articulis tribus primis laete testaceis. Caput thorace paulo angustius, subtiliter punctulatum, ore angustato, testaceo. Thorax latitudine dimidio brevior, coleopteris paulo angustior, basin versus subangustatus, lateribus ante medium modice rotundatus, basi leviter rotundatus, apice late emarginatus, angulis anterioribus acutis, deflexis, posterioribus subrectis, leviter convexus, confertim subtilius punctatus, basi supra scutellum obsolete impressus. Elytra thorace sesqui longiora, sat crebre fortiterque punctata. Abdomen parallelum, supra crebrius subtiliter punctatum, ano rufo-ferrugineo. Pedes laete testacei.

Mas abdominis segmento dorsali penultimo tuberculo minuto oblongo, satis prominulo, insignis.

Obs. *O. anali* proxime affinis, colore autem thoraceque coleopteris angustiore distincta.

\* 34. *O. maura*: Nigra, nitida, ore testaceo, antennis pedibusque piceis, thorace quadrato, convexo, coleopteris angustiore, elytris fortius punctulatis. — Long.  $1\frac{1}{3}$  lin.

Er. Col. March. I. 352. 18.

Habitat in Germania, Berolini, Dom. Schüppel et ipse.

Statura fere *O. prolixae*, at minus elongata, nigra, nitida, tenuiter griseo-pubescens. Antennae capitis thoracisque longitudine, articulo secundo tertio duplo maiore, apicem versus incrassatae, articulis 4—10 sensim crassioribus, penultimis transversis, ultimo maiusculo, ovato, subacuminato, pubescentes, piceae, basi rufescentes. Caput dense subtilius punctatum, ore attenuato, testaceo. Thorax coleopteris paulo angustior, latitudine haud brevior, basi leviter rotundatus, lateribus subrectis, apice truncatus, angulis omnibus subrectis, anterioribus deflexis, supra convexus, aequalis, confertim

minus subtiliter punctatus. Elytra thorace paulo longiora, leviter convexa, dense fortius punctata. Abdomen parallelum, subtilius parcusque punctatum, segmentis tribus primis basi fortiter transversim depressis. Pedes picei, geniculis tarsisque rufescentibus.

Species mihi invisae, forte huius generis et sectionis.

35. *O. suturalis*: *Nigra, nitida, tenuiter pubescens, ore, antennis, elytrorum sutura anoque rufis, antennarum articulo ultimo oblongo-ovato, elytris intra apicem anteriorem profunde sinuatis.* — Long. 1 lin.

*Bolitochara suturalis* Mannerh. *Brachelytr.* 82. 47.

*Aleochara suturalis* Sahlb. *Ins. Fenn.* 1. 376. 55.

Habitat in Fennia australi, rarissime.

Caput thorace angustius, rotundatum, basi truncatum, nigrum, nitidum. tenuiter pubescens, ore palpisque rufis. Antennae longitudine thoracis, versus apicem vix incrassatae, articulo ultimo oblongo-ovato, rufo, in medio nonnihil obscuriores. Thorax transversus, longitudine duplo fere latior, antice truncatus, angulis compressis, lateribus rotundatus, postice subtruncatus, angulis nonnihil prominulis, supra valde convexus, niger, nitidus, pubescens. Scutellum breve, triangulare, nigrum. Elytra thorace vix latiora, sed eodem longiora, apice intra angulum anteriorem profundius sinuato-emarginata, supra subdepressa, nigra, nitida, subtilissime punctulata, pubescentia, sutura margineque apicis rufis. Abdomen pone elytra vix reliqui corporis longitudine corpusque subtus nigra, nitida, marginibus segmentorum ventralium anoque rufis. Pedes mediocres, toti rufi. (Sahlb.)

36. *O. pulchella*: *Nigro-picea, opaca, crebre punctulata, subpubescens, antennis extrorsum crassioribus, ore, elytris, ano pedibusque rufis, articulo antennarum ultimo subgloboso, elytris apice emarginatis.* — Long. 1 lin.

*Bolitochara pulchella* Mannerh. *Brachelytr.* 83. 48.

*Aleochara pulchella* Sahlb. *Ins. Fenn.* 1. 377. 56.

Habitat in Fennia australi, rarissime, semel capta.

A praecedente differt colore, thorace paulo longiore et angustiore, antennis versus apicem incrassatis et articulo harum ultimo subgloboso. Caput omnino ut in praecedente. Antennae thorace fere longiores, versus apicem incrassatae, articulis tribus baseos, ut in ceteris, elongatis, 4 — 8 sensim paulo crassioribus, ultimo maximo, subgloboso, rufae, versus apicem lutescentes. Thorax subtransversus, longitudine dimidio latior, antice truncatus, angulis compressis, lateribus rotundatus, postice subtruncatus, angulis nonnihil prominulis, supra convexus, nigro-piceus, opacus, subtilissime punctulatus, pubescens, foveola ante scutellum obsolete impressa. Scutellum subtriangulare, nigro-piceum, subtilissime punctulatum. Elytra thoracis latitudine, illoque vix longiora, apice intra angulum anteriorem sinuato-emarginata, supra depressa, obscura rufa, crebre punctulata. Abdomen longitudine fere reliqui corporis, nigro-piceum, punctulatum, pubescens, segmentis ventralibus primis fere totis, marginibus insequentium et ano rufescentibus. Pedes mediocres, pallide rufi. (Sahlb.)



37. *O. elegantula*: *Lineari-elongata, rufa, nitida, capite, thoracis disco abdominisque cingulo nigro-piceis, thorace elytris apice emarginatis angustiore, basi foveolato, antennarum articulo ultimo ovato.* — Long  $1\frac{1}{6}$  lin.

*Bolitochara elegantula* Mannerh. *Brachelytr.* 377. 57.

*Aleochara elegantula* Sahlb. *Ins. Fenn. I.* 377. 57. — *Zetterst. Ins. Lapp.* 78. 23.

Habitat in Fennia, Suecia, Lapponia sub cortice Pinus.

A praecedente colore, thorace elytris angustiore et minus convexo, nec non forma articuli antennarum ultimi distincta. Caput ut in proxime praecedentibus. Antennae longitudine thoracis, versus apicem sensim crassiores, rufescentes, extrorsum fusciscentes, articulo ultimo ovato. Thorax subtransversus, longitudine vix dimidio latior, antice truncatus, angulis compressis, lateribus rotundatus, postice subtruncatus, angulis parum prominulis, supra convexus, nigro-piceus, limbo omni anguste rufescente, subnitidus, pubescens. Scutellum breve, triangulare, piceo-rufescens. Elytra thorace latiora et longiora, apice intra angulum externum sinuato-emarginata, supra depressa, rufescentia, disco posteriore fusco-picea, crebre punctulata, pubescentia. Abdomen pone elytra reliquo corpore fere longius corpusque subtus rufescentia, nitida, pubescentia, cingulo abdominis piceo-nigro, ano pallide rufo. Pedes toti rufi. (Sahlb.)

### **Aleochara Grav.**

*Mannerh. Brachelytr. p.* 66. — *Er. Col. March. I. p.* 353.

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula brevis, bifida; paraglossae brevissimae.*

*Palpi labiales 4-articulati, articulis sensim minoribus, ultimo parvo, subulato.*

*Tarsi omnes 5-articulati, postici articulo primo elongato.*

Corpus oblongum, crassiusculum, alatum. Caput exsertum, deflexum, basi haud constrictum, thorace plerumque angustius, oculis mediocribus, oblongis, parum prominulis. Labrum transversum, apice truncatum. Mandibulae breviusculae, muticae. Maxillae mala exteriore cornea, basi apiceque membranea, portione membranea apicali pubescente, mala interiore exteriore paulo breviori, membranea, margine exteriore corneo, apice introrsum oblique truncata, sinu spinulis brevibus ciliato. Palpi maxillares leviter elongati, articulo secundo tertioque longitudine subaequalibus, hoc parum incrassato, quarto minore, subulato. Labium (t. III. f. 9.) mento brevi, apice late emarginato, ligula brevi, palporum labialium articuli primi apicem vix attingens, latiuscula, bifida, paraglossis brevissimis, latis, subacuminatis, interne haud ciliatis. Palpi labiales 4-articulati, articulo primo maiore, sequentibus decrescentibus, quarto minimo, at distincto. Antennae plerumque breviores, articulis tribus primis magis elongatis, secundo tertioque obconicis, primo crassiusculo. Thorax transversus, basi coleopterorum latitudine, antrorsum angustatus, apice truncatus, basi et lateribus rotundatus. Elytra breviter truncata. Abdomen plerumque parallelum, rarius apicem versus angustatum, supra planum, subtus convexum, in vivis reclinatum. Pedes breviores, intermedii basi leviter distantes, tarsi omnibus 5-articulatis, posticis articulo primo elongato.

## Observationes.

1. Maxillae omnino ut in Homalota (t. III. f. 22). Ligula brevissima biloba, paraglossis latis brevissimis vix longiore, et palpis labialibus 4-articulatis hoc genus insigne. Unicum hucusque coleopterorum genus, in quo mihi ultra tres palporum labialium articuli veri observati sunt, palporum stipites enim saepius articuli basalis specimen prae se ferunt, ut in Cicindelae plurimisque Carabis.

2. Facies huius generis quodammodo peculiaris pube haud sericante, corpore crassiusculo, elytris, antennis pedibusque brevioribus, abdomine saepius revoluta. A Myrmedonia distinguitur praeter palporum structuram tarsis anticis 5-articulatis.

3. Victus in fimo, cadaveribus, saepius etiam in quisquiliis.

4. Mas distinguitur abdominis segmento inferiore sexto apice utrinque sinuato, medio leviter triangulariter producto.

\* 1. *A. fuscipes*: Nigra, nitida, antennis brevibus, fusiformibus basi pedibusque rufis, elytris thorace brevioribus, rufis, lateribus fuscis, abdomine supra parcius punctato. — Long.  $2\frac{1}{2}$  — 3 lin.

Grav. Micr. 92. 36. Mon. 170. 56. — Latr. Hist. nat. Crust. et Ins. IX. 388. 36. — Gyll. Ins. Suec. II. 428. 50. — Germ. Faun. Ins. Europ. XIII. 8. — Boisd. et Lacord. Faun. Ent. Paris. I. 528. 1. — Mannerh. Brachelytr. 66. 1. — Runde Brachelytr. Hal. 28. 1. — Er. Col. March. I. 354. 1.

Staph. fuscipes Fab. Syst. Ent. 266. 12. Spec. Ins. I. 336. 15. Mant. Ins. I. 221. 22. Ent. Syst. I. II. 527. 38. Syst. El. II. 598. 47. — Payk. Faun. Suec. III. 397. 39. — Schrank Faun. Boic. I. 645. 845. — Walk. Faun. Ent. Paris. I. 271. 13. — Cederh. Faun. Ingr. 119. 364. — Marsh. Ent. Britt. 525. 77.

Staph. brachypterus Fourc. Ent. Par. I. 167. 12.

Staph. à étuis très-courts Geoffr. Hist. Ins. I. 365. 12.

Var. *Elytris concoloribus, antennarum basi pedibusque piceis.*

*Aleochara lata* Grav. Micr. 186. 2. Mon. 170. 57.

Habitat in Europa et in America septentrionali.

Magnitudine variabilis, speciminibus minutis A. tristis individuis maioribus aequalibus, plerumque duplo vel triplo maior, lata, nigra, nitida, capite thoraceque nonnunquam subaeneis, parcius flavescenti-villosa. Antennae capite sesqui longiores, crassae, articulo secundo tertio dimidio brevior, quarto sequentibus angustiore, vix transverso, 5 — 10 fortiter transversis, crassitie duplo brevioribus, inter se aequalibus, ultimo praecedentibus duobus coniunctis longitudine aequali, subconico, nigrae fuscae, basi rufae vel rufo-testaceae, apice etiam nonnunquam rufescentes, Palpi rufi. Caput thorace multo angustius, deflexum, parce punctatum. Thorax latitudine dimidio brevior, basi coleopterorum prope latitudine, basi et lateribus rotundatus, antrum angustatus, angulis omnibus obtusis, anterioribus deflexis, modice convexus, sat crebre distincteque punctatus. Elytra thorace paulo breviora, angulo exteriori rotundato, creberrime sat fortiter punctata, punctis oblique impressis, rufa, lateribus fusca. Abdomen parallelum, utrinque sat fortiter, supra minus confertim punctatum, segmentis dorsalibus ultimis duobus, ventralibus omnibus saepius margine apicali rufo-piceo, rarius concolore. Pedes rufi, femoribus posticis plerumque magis minusve piceis.

Variat elytris totis nigris, antennis basi pedibusque piceis, his geniculis tarsisque rufis.

Obs. 1. Specimina Americana cum Europaeis omnino congruunt. Varietas elytris concoloribus rarior, individua Pensylvanica supersunt e Mus. Knochiano et Hoffmannseggiano, exemplum Austriacum addidit Dom. Schüppel.

2. *Staph. fuscipes Fabricii* certe huic speciei adscribitur, ipse enim insectum in auctoris Museo vidi. Reliqua synonyma indubia puto. Marshamianum locum non sine haesitatione allegavi, temere enim auctor synonyma contraria (Linnei, Fabricii et Panzeri) composuit et descriptione Linneana illustravit. *Staph. fuscipes* Panz. apud Tachinos citandus, *Staph. fuscipes* Linnei species dubia quidem est, at certe distincta a nostra, „thorace, capite et maxillis insignibus,” forte ad *Oxytelum* vel potius *Platysthetum* quendam spectans.

- \* 2. *A. chrysorrhoea*: *Nigra, nitida, antennis brevibus fusiformibus basi, ano pedibusque anterioribus testaceis, elytris thorace subbrevioribus, abdomine supra parcius punctato. — Long. 3 lin.*

Habitat in Brasilia, Dom. Virmond.

Statura praecedentis, nigra, nitida, parcius griseo-pilosa. Antennae capite paulo longiores, articulo secundo tertio dimidio brevior, sequentibus crassis, 5—8 fortiter transversis, crassitie plus duplo brevioribus, praecedentibus duobus his paulo angustioribus, licet transversis, at longitudine vix duplo crassioribus, ultimo iterum paulo angustiore, oblongo-ovato, medio leviter constricto, fuscae, articulis tribus primis testaceis, ultimo apice subferrugineo. Palpi picei, articulo terminali pallide testaceo: Caput thorace multo minus, deflexum, fronte planiuscula, crebre fortiterque punctata, linea media longitudinali laevi. Thorax latitudine dimidio brevior, basi coleopteris paululum angustior, basi leviter, lateribus fortius dilatatis, apicem versus modice angustatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, leviter convexus, sat crebre fortiterque punctatus. Elytra thorace perparum breviora, angulo apicali exteriore rotundato, dense fortiter punctata. Abdomen apicem versus subangustatum, utrinque parcius punctatum, segmento penultimo apice ultimoque totis fulvis, segmentis ventralibus piceo-marginatis. Pedes testacei.

- \* 3. *A. dilatata*: *Nigra, nitida, antennis brevibus, fusiformibus basi, ano pedibusque rufo-testaceis, elytris thorace brevioribus, abdomine supra parcius punctato. — Long. 2 lin.*

Habitat in Madagascar, Dom. Goudot.

Magnitudine *A. bipunctatae*, at brevior, nigra, nitida, subtiliter puberula. Antennae capite parum longiores, crassae, articulo secundo tertio dimidio brevior, quarto leviter transverso, 5—10 brevissimis, longitudine triplo crassioribus, ultimo praecedentibus duobus coniunctis sublongiore, ovato-conico, fuscae, articulis 3 primis testaceis. Palpi testacei. Caput parvum, deflexum, fronte depressa, parce punctata. Thorax latitudine plus dimidio brevior, basi coleopterorum prope latitudine, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus rotundatis, deflexis, posterioribus obtusissimis, transversim leviter convexus, crebrius punctatus. Elytra thorace dimidio prope breviora, angulo apicali exteriore rotundato, densius profunde punctata. Abdomen breviusculum, apicem versus subangustatum, supra parcius profundiusque punctatum, segmento penultimo margine apicali ultimoque toto testaceis, subtus crebrius punctatum, densius griseo-pilosum, segmentis singulis margine apicali tenui ferrugineo. Pedes testacei.



- \* 4. *A. Cayennensis*: *Rufo-testacea, capite abdominisque segmentis quatuor intermediis nigris, antennis nigris, basi testaceis, elytris thorace brevioribus, abdomine supra parcius punctato. — Long. 2 lin.*

Laporte *Etud. Ent.* p. 136.

Habitat in Cayenna, Dom. Buquet.

Statura *A. rufipennis*, rufo-testacea, subnitida, fulvescenti-puberula. Antennae capite sesqui longiores, articulo secundo tertio paulo brevior, quarto turbinato, crassitie haud brevior, 5 — 10 transversis, longitudine sesqui fere crassioribus, ultimo praecedentibus crassitie aequali, oblongo-ovato, praecedentibus tribus coniunctis vix brevior, fusco-nigrae, articulis 4 primis testaceis. Palpi picei. Caput deflexum, nigrum, parce obsoletius punctatum. Thorax latitudine dimidio brevior, basi coleopterorum prope latitudine, apicem versus angustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis omnibus obtusissimis, subrotundatis, anterioribus deflexis, transversim leviter convexus, dorso subdepressus, subtilius dense punctatus, rufo-testaceus, medio lineis duabus approximatis, utrinque abbreviatis, obscuris. Elytra thorace paulo breviora, confertim punctata, angulo apicali exteriori rotundato, subfusco. Abdomen parallelum, supra parcius profundiusque punctatum, basi obscure rufo-testaceum, segmentis 2 — 5 nigris, subtus margine apicali ferrugineo, ultimis duobus dilute testaceis. Pedes dilute testacei.

- \* 5. *A. lateralis*: *Nigra, antennis medio subincrassatis basi pedibusque rufo-testaceis, ano elytrisque rufis, his thorace brevioribus, macula laterali nigra, abdomine supra crebrius punctato. — Long. 2½ lin.*

Habitat in Columbia, Dom. Moritz.

*A. fuscipede* angustior, nigra, nitida, griseo-puberula. Antennae capite sesqui longiores, ab articulo quarto inde incrassatae, medio paulo crassiores, apicem versus parum attenuatae, articulo quarto cyathiformi, crassitie paulo brevior, 5—10 transversis, crassitie duplo brevioribus, ultimo ovato-subconico, praecedentibus duobus coniunctis longitudine aequali, fuscae, articulis tribus primis testaceo-rufis. Labrum rufum. Palpi testaceo-rufi. Caput deflexum, disperse sat profunde punctatum. Thorax latitudine plus dimidio brevior, basi coleopterorum prope longitudine, apicem versus subangustatus, basi leviter, lateribus fortius rotundatus, angulis omnibus obtusis, leviter transversim convexus, confertim punctatus. Elytra thorace paulo breviora, angulo apicali exteriori rotundato, confertissime punctata, rubra, macula marginali, humerum haud attingente, nigra. Abdomen apice parum angustatum, supra fortius crebriusque punctatum, longius pilosum, segmento penultimo apice ultimoque toto saturate rubris, subtus dense punctatum, segmentis singulis margine apicali rufo-piceo. Pedes testaceo-rufi.

6. *A. breviuscula*: *Piceo-nigra, nitida, antennarum basi pedibusque testaceis, thoracis lateribus elytrisque rufo-piceis, his thorace brevioribus, lateribus nigris, abdomine supra parce punctato, antennis crassis, subfusiformibus. — Long. 1½ lin.*

Habitat in Columbia, Dom. Moritz.

Habitu fere praecedentium, at minor, *A. pullae* similis et eadem pa-

rum longior, at crassula et antennis crassis dissimilis, piceo-nigra, nitida, tenniter griseo-puberula. Antennae capite sesqui longiores, crassae, medio subincrassatae, articulis 4—10 transversis, fuscae, basi rufo-testaceae. Caput parce obsoletius punctatum, ore testaceo. Thorax coleopteris vix angustior, latitudine dimidio brevior, modice transversim convexus, crebrius punctatus, lateribus rufo-piceus. Elytra thorace paulo breviora, crebrius punctata, rufo-picea, lateribus nigra. Abdomen breviusculum, apicem versus parum angustatum, supra parce punctatum, ano segmentisque singulis margine apicali rufo-piceis. Pedes flavescenti-testacei.

- ✕ \* 7. *A. rufipennis*: Nigra, nitida, elytris thorace subbrevioribus rufis, lateribus nigris, pedibus piceis, abdomine supra parce punctato. — Long. 2 lin.

*Aleo. laevigata* Boisd. et Lacord. Faun. Ent. Paris. I. 530. 6.

Habitat in Austria, Gallia, Italia.

Habitu omnino *A. tristis*, nigra, nitida, densius subtiliter griseo-puberula. Antennae capite duplo longiores, crassiusculae, apicem versus incrassatae, articulo secundo tertio paulo brevior, 5—10 transversis, ultimo his crassitie aequali, ovato, praecedentibus duobus coniunctis fere longiore, basi obscure piceae. Palpi maxillares picei, articulo terminali pallide testaceo. Caput deflexum, parce obsoletius punctatum. Thorax latitudine dimidio brevior, coleopterorum prope latitudine, antrorsum vix angustatus, basi leviter, lateribus fortius rotundatus, apice truncatus, angulis omnibus obtusissimis, subrotundatis, anterioribus deflexis, transversim leviter convexus, crebre subtilius punctatus. Elytra thorace paululum breviora, angulo apicali exteriorē rotundato, dense punctata, laete rufa, latere nigro. Abdomen apicem versus subangustatum, supra sub-glabrum, parcius profundiusque punctatum, subtus crebrius punctatum, longius pilosum, segmentis singulis margine apicali obscure piceo. Pedes picei, geniculis tarsisque rufis.

- ✕ \* 8. *A. tristis*: Nigra, nitida, elytris thorace brevioribus, macula apicali rufo-testacea, pedibus piceis, thorace aequaliter punctato, abdomine parallelo, supra confertissime punctato. — Long.  $1\frac{1}{2}$ — $2\frac{1}{4}$  lin.

*Grav. Mon.* 170. 59. — *Gyll. Ins. Suec.* II. 430. 59. — *Sahlb. Ins. Fenn.* I. 395. 86. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 529. 2. — *Mannerh. Brachelytr.* 66. 2. — *Runde Brachelytr. Hal.* 28. 2. — *Er. Col. March.* I. 355. 2.

*Staph. geometricus* Schrank *Faun. Boica* I. 642. 869.

*Var. Elytris rufo-testaceis.*

*Aleo. crassiuscula* Sahlb. *Ins. Fenn.* I. 396. 89.

Habitat in Europa et in Oriente.

Statura fere *A. fuscipedis*, at plerumque duplo minor, magnitudine magnopere varians, nigra, subaeneo-nitidula, breviter dense cinereo-puberula. Antennae capite paulo longiores, articulo secundo tertio subaequali, apicem versus sensim leviter incrassatae, articulis penulimīs transversis, ultimo praecedentibus duobus coniunctis longitudine subaequali, subconico, nigrae, articulo primo interdum piceo. Palpi picei. Caput deflexum, parcius punctatum. Thorax latitudine plus dimidio brevior, coleopteris latitu-

dine aequalis, apicem versus vix angustatus, basi leviter, lateribus fortius punctatus, apice truncatus, angulis omnibus obtusissimis, subrotundatis, anterioribus deflexis, transversim modice convexus, confertim punctatus, disco interdum punctis quatuor, quadratim positis, vel sex evidentioribus. Elytra thorace paulo breviora, angulo apicali exteriori rotundato, confertissime punctata, subrugulosa, prope suturam macula magna apicali rufo-testacea notata. Abdomen parallelum, crassiusculum, supra infraque confertissime punctatum. Pedes picei, tarsi rufescentibus.

Variat interdum elytris rufo-testaceis, saepius basi et circa scutellum, maculaque laterali fusciscentibus.

Obs. Occurrunt nonnunquam individua minuta, long.  $1\frac{1}{2}$  —  $1\frac{2}{3}$  lin., elytris totis rufo-testaceis, thorace parcius obsoletiusque punctato, ab hac vix specie distinguenda. Huc forte referenda *Al. crassicornis* Boisd. et Lacord.

✕ \* 9. *A. bipunctata*: Nigra, nitida, elytris thorace dimidio brevioribus, apice sanguineis, pedibus testaceis, thorace convexo, creberrime punctato, abdomine apicem versus attenuato, supra crebre punctato. — Long.  $2\frac{1}{4}$  lin.

*Grav. Micr.* 93. 37. *Mon.* 171. 61. — *Latr. Hist. nat. Crust. et Ins.* IX. 338. 37. *Gen. Crust. et Ins.* I. 301. 1. — *Gyll. Ins. Suec.* II. 430. 52. — *Zetterst. Faun. Lappon.* I. 104. 23. *Ins. Lapp.* 79. 33. — *Sahlb. Ins. Fenn.* I. 95. 87. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 529. 3. — *Er. Col. March.* I. 355. 3.

*Staph. bipunctatus* Oliv. *Ent.* III. 42. 31. 43. t. 5. f. 44. (forte)

*Al. intricata* Mannerh. *Brachelytr.* 66. 4. — *Runde Brachelytr.* Hal 28. 4. — *Sahlb. Ins. Fenn.* I. 396. 88.

Habitat in Europā.

Præcedentibus brevior et magis convexa, nigra, nitidula, pube brevi subtili depressa fusca vestita. Antennae capite paulo longiores, tenuiores, articulo secundo tertio longitudine aequali, apicem versus leniter incrassatae, articulis penultimis subtransversis, ultimo præcedentibus ambobus vix longiore, oblongo-ovato, nigrae, basi obscure piceae. Palpi maxillares nigri, articulo terminali testaceo. Caput deflexum, dense punctatum. Thorax latitudine dimidio brevior, basi coleopterorum latitudine, antrorsum angustatus, basi lateribusque rotundatus, apice truncatus, angulis anterioribus subrectis, deflexis, posterioribus obtusis, sat convexus, subtiliter confertim punctatus. Elytra thorace dimidio breviora, angulo apicali exteriori rotundato, confertissime vel ruguloso-punctata, opaca, apice sanguinea. Abdomen apicem versus sensim attenuatum, supra dense, subtus fortius punctatum, segmentis ventralibus singulis margine posteriore rufo-piceo. Pedes rufo-testacei.

\* 10. *A. brevipennis*: Nigra, nitida, antennarum basi pedibusque testaceis, elytris thoracis longitudine, creberrime punctatis, immaculatis, abdomine apicem versus attenuato, supra parcius punctato. — Long. 2 lin.

*Grav. Mon.* 169. 53. — *Er. Col. March.* I. 356. 4.

*Al. carnivora* Gyll. *Ins. Suec.* II. 431. 53. — *Sahlb. Ins. Fenn.* I. 397. 90. — *Zetterst. Faun. Lapp.* I. 104. 24. *Ins. Lapp.* 79. 34. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 350. 4. — *Mannerh. Brachelytr.* 66. 5.

Habitat in Europā.

Statura *A. bipunctatae*, nigra, subaeneo-nitida, lanugine depressa fla-



vescenti-grisea, densius vestita. Antennae capite duplo longiores, articulo secundo tertio paulo brevior, apicem versus sensim leniter incrassatae, articulis penultimis longitudine haud crassioribus, ultimo praecedentibus ambobus longitudine aequali, oblongo-ovato, apice compresso, nigrae, articulo primo rufo. Palpi rufi. Caput deflexum, fronte parce punctata. Thorax latitudine plus dimidio brevior, basi coleopterorum latitudine, apicem versus fortiter angustatus, basi leviter, lateribus fortius rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, sat fortiter convexus, parcius subtiliter punctatus. Elytra thoracis longitudine, angulo apicali exteriore rotundato, profunde confertissime punctata, subrugulosa, nigro-subaenea, opaca, apice concolore. Abdomen apicem versus sensim leviterque angustatum, supra parcius, subtus crebrius fortiter punctatum, segmentis ventralibus apice piceo-marginatis. Pedes testacei, femoribus interdum fusciscentibus.

- \* 11. *A. verberans*: *Nigra, nitida, antennarum basi pedibusque rufis, femoribus posticis nigris, elytris thoracis longitudine crebre, fortiterque punctatis, apice rufescentibus, abdomine parallelo, supra parce punctato. — Long. 2 lin.*

Habitat in Brasilia, Columbia.

Statura et summa similitudo *A. lanuginosae*. Corpus nigrum, nitidum, pube tenuiore depressa fulvescente parce vestitum. Antennae capite sesqui longiores, articulo secundo tertioque aequalibus, apicem versus sensim incrassatae, articulis 7—10 transversis, ultimo subovato, apice fortiter compresso, praecedentibus ambobus longitudine aequali, nigrae, articulo primo secundoque rufis. Palpi picei. Caput deflexum, fronte parce obsolete, vertice crebre distinctius punctatum. Thorax latitudine dimidio brevior, coleopterorum prope latitudine, antrosum vix angustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus obtusis, deflexis, posterioribus obtusissimis, subrotundatis, transversim leviter convexus, basi distincte, apicem versus sensim obsolete punctatus, basi obsolete biimpressus, ut in *A. lanuginosa*. Elytra thoracis longitudine, angulo apicali exteriore rotundato, creberrime profundius punctata, interstitiis subrugosis, sub-aeneo-nigra, apice magis minusve rufescentia. Abdomen parallelum, supra parce punctatum, subtus segmentis singulis margine apicali ferrugineo. Pedes rufi, femoribus posticis nigris, apice rufis.

Obs. Specimina huius speciei duo, prope Caracas Columbiae lecta communicavit D. Moritz, alterum Brasiliensibus magnitudine aequale, alterum duplo maius.

- \* 12. *A. vulnerata*: *Nigra, nitida, antennarum basi, pedibus elytrisque rufis, his thorace brevioribus, sutura latereque late nigris, abdomine apicem versus attenuato, supra profunde punctato. — Long. 2½ lin.*

Habitat in Madagascar, Dom. Goudot.

Statura fere praecedentis, nigra, nitida, subtiliter griseo-lanuginosa. Antennae capite sesqui longiores, articulo secundo tertioque aequalibus, apicem versus leniter incrassatae, articulis penultimis longitudine paulo crassioribus, ultimo his crassitie et penultimis duobus coniunctis longitudine aequali, oblongo-ovato, nigrae, basi rufae. Thorax latitudine dimidio bre-

vior, antrorsum angustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus obtusis, deflexis, posterioribus subrotundatis, transversim convexus, confertim punctatus. Elytra thorace paulo breviora, angulo apicali exteriore subrotundato, confertissime punctata, rufa, sutura basi dilatata limboque lato laterali, antice abbreviato, nigris. Abdomen apicem versus sensim attenuatum, supra parce pilosum, profundius minus dense punctatum, longitudinaliter subrugosum, subtus segmentis singulis ferrugineo-marginatis. Pedes rufi.

- \* 13. *A. puberula*: *Nigra, fusco-pubescent, antennarum basi, thoracis limbo, elytrorum disco pedibusque rufis, elytris thoracis longitudine, abdomine dense punctato. — Long. 2 lin.*

*Klug Coleopt. Madagasc. 51. 44.*

Habitat in Madagascar, Dom. Goudot.

Statura intermedia fere inter *A. tristem* et *brevipennem*. *Nigra*, subnitida, pube depressa fusca densius vestita. Antennae capitis thoracisque longitudine, articulo secundo tertio paulo longiore, apicem versus incrassatae, articulis penultimis transversis, ultimo praecedentibus ambobus longitudine aequali, conico, acuminato, nigrae, basi rufo-testaceae, apice ferrugineo. Caput subglobosum, deflexum, confertim subtiliter punctatum. Thorax latitudine dimidio brevior, basi coleopterorum latitudine, lateribus et basi leviter rotundatus, antrorsum fortius angustatus, apice truncatus, angulis anterioribus rectis, fortiter deflexis, posterioribus obtusis, leviter transversim convexus, confertim subtilius punctatus, limbo omni obscure rufo. Elytra thoracis longitudine, angulo apicali exteriore rotundato, fortius confertissime punctata, rufa, basi, sutura maculaque magna laterali fuscis. Abdomen apicem versus sensim attenuatum, fortius confertim punctatum, nigrum, segmentorum marginibus rufo-piceis. Pedes testacei.

- \* 14. *A. taeniata*: *Nigra, subopaca, antennarum basi, elytrorum vitta lata arcuata pedibusque testaceis, elytris thorace dimidio brevioribus, abdomine supra densius punctato. — Long. 2 lin.*

Habitat in Americae ins. St. Johannis, Dom. Moritz.

Statura fere *A. tristis*, at paulo magis depressa. *Nigra*, vix nitida, densius griseo-pubescent. Antennae capite sesqui longiores, articulo secundo tertio brevior, apicem versus incrassatae, articulis penultimis transversis, ultimo praecedentibus ambobus vix longiore, ovato, nigrae, basi testaceae. Palpi maxillares picei, articulo terminali testaceo. Caput deflexum, subtiliter punctatum. Thorax latitudine dimidio brevior, coleopterorum latitudine, basi lateribusque leviter rotundatus, antrorsum subangustatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, leviter transversim convexus, subtiliter densius punctatus. Elytra thorace dimidio breviora, angulo apicali exteriore rotundato, creberrime minus fortiter punctata, testacea, sutura infuscata, macula magna laterali nigra. Abdomen apice attenuatum, supra densius punctatum, nigrum, segmentis anterioribus apice obscure testaceis, subtus segmentorum singulorum margine lato posteriore anoque ferrugineis. Pedes testacei, coxis picei.

- \* 15. *A. fumata*: Nigra, subnitida, antennis basi pedibusque testaceis, thoracis lateribus elytrisque piceis, his thoracis longitudine, creberrime punctatis, abdomine apicem versus sensim subangustato, supra parcius profunde punctato. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 357. 6.

*Al. fumata* var. 2 Grav. Micr. 96. 42. Mon. 172. 64. — Latr. Hist. nat. Crust. et Ins. IX. 389. 42.

*Al. curta* Sahlb. Ins. Fenn. I. 400. 97.

Habitat in Europa.

Statura *A. brevipennis*, at triplo minor, pube depressa grisea densius vestita. Antennae capitis thoracisque fere longitudine, articulo secundo tertio paulo brevior, apicem versus incrassatae, articulis 6—10 leviter transversis, ultimo praecedentibus ambobus aequali, ovato, subacuminato, rufae, basi testaceae. Palpi rufi. Caput deflexum, fronte subtiliter parceque punctata. Thorax basi coleopterorum latitudine, hac duplo prope brevior, basi leviter, lateribus fortius rotundatus, antrorsum angustatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, sat convexus, confertim subtiliter punctatus, lateribus obscure piceis. Elytra thorace vix breviora, angulo apicali exteriore rotundato, confertim profundius punctata, nigro-vel fusco-picea, minus nitida. Abdomen apicem versus sensim subangustatum, supra parce profundeque punctatum, subtus segmentis singulis late piceo-marginatis. Pedes testacei.

Obs. 1. Proxima *A. brevipenni*, differt autem statura multo minore, thorace lateribus et elytris piceis, his paulo brevioribus, minus dense punctatis.

2. Ab *A. moerente*, quacum olim mihi confusa, discedit statura brevior et elytrorum angulo apicali exteriore rotundato.

- \* 16. *A. bisignata*: Nigra, nitida, pedibus piceis, thorace parce subtiliterque aequaliter punctato, elytris thoracis longitudine, crebre punctatis, macula apicali rufa, abdomine parallelo, supra parcius profunde punctato. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 357. 7.

Habitat in Europa et in Oriente.

Statura omnino *A. nitidae*, nigra, nitida, parcius griseo-lanuginosa. Antennae capite paulo longiores, crassiusculae, articulo secundo tertioque aequalibus, 5—10 transversis, longitudine plus duplo crassioribus, inter se aequalibus, ultimo praecedentibus ambobus vix longitudine aequali, subovato, acummatum, totae nigrae. Palpi maxillares nigri, articulo terminali testaceo. Caput deflexum, laeve. Thorax latitudine duplo prope brevior, coleopterorum latitudine, basi leviter, lateribus modice rotundatus, antrorsum subangustatus, apice truncatus, angulis omnibus obtusis, anterioribus subdeflexis, leviter transversim convexus, subtiliter parceque punctatus. Elytra thoracis longitudine, angulo apicali exteriore rotundato, profunde fortiterque crebre punctata, nigro-picea, macula magna apicali rufa prope suturam signata. Abdomen parallelum, supra minus crebre fortius punctatum, nitidum. Pedes picei, geniculis tarsisque rufescentibus.



- \* 17. *A. bimaculata*: *Nigra, nitida, antennarum basi pedibusque rufo-piceis, elytris thorace subbrevioribus, macula magna testacea, thorace lateribus crebre, dorso biserialim, abdomine supra confertim punctato.* — Long 2 lin.

Grav. Micr. 187. 3. Mon. 170. 58. — Say Act. Acad. nat. Scienc. Philadelph. III. p. 157.

Habitat in Pensylvania.

Statura *A. tristis*. Nigro-subaenea, nitida, pube parca depressa fulvescente vestita. Antennae capite parum longiores, articulo secundo tertio vix brevior, apicem versus paulum incrassatae, articulis 6 — 10 transversis, ultimo praecedentibus ambobus vix brevior, subconico, basi obscure rufae. Palpi rufi. Caput deflexum, fronte utrinque parce sparsim punctata, vertice laevissimo. Thorax coleopterorum latitudine, hac dimidio brevior, basi leviter, lateribus modice rotundatus, antrorsum subangustatus, apice truncatus, angulis anterioribus obtusis, subrotundatis, deflexis, posterioribus rotundatis, transversim convexus, lateribus crebre obsoletius punctatus, lineis duabus longitudinalibus impressis, irregulariter fortius punctatis, spatio intermedio laevi. Elytra thorace paululum breviora, angulo apicali exteriori rotundato, dense fortiter punctata, macula magna apicali rufo-testacea. Abdomen parallelum, crassiusculum, supra crebre punctatum, subtus segmentis singulis margine apicali obscure piceo. Pedes picei, tarsis rufis, interdum rufi, femoribus posticis solis piceis.

Variat statura duplo maiore.

18. *A. duplicata*: *Nigra, nitida, antennarum basi pedibusque rufo-piceis, elytris thorace brevioribus, macula apicali testacea, thorace lateribus parce subtiliterque, dorso biserialim, abdomine supra confertim punctato* — Long.  $1\frac{3}{4}$  lin.

Habitat prope Caracas Columbiae, Dom. Moritz.

Praecedentibus affinis, *A. bimaculata* duplo fere minor, *A. nitidae* aequalis, nigro-subaenea, nitida, elytris abdomineque subtilissime puberulis. Antennae capite sesqui longiores, apicem versus sensim incrassatae, nigrae, basi rufescenti-piceae. Caput deflexum, disperse obsoletius punctatum. Thorax lateribus disperse subtiliusque punctatus, medio laevis, lineis duabus longitudinalibus impressis, inaequaliter punctatis. Elytra thorace paulo breviora, dense punctata, macula magna apicali triangulari rufo-testacea. Abdomen parallelum, creberrime punctatum. Pedes rufi, femoribus piceis.

- \* 19. *A. notula*: *Nigra, nitida, antennarum basi, pedibus elytrorumque macula apicali rufo-testaceis, his thorace brevioribus, thorace biserialim, elytris abdomineque parcius punctatis.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Americae merid. ins. St. Thomae, Dom. Moritz.

*A. nitidae* affinis, at paulo minor, nigro-subaenea, nitida, elytris abdomineque subtiliter puberulis. Antennae capite paulo longiores, articulo secundo tertioque aequalibus, apicem versus sensim leniter incrassatae, articulis 5 — 10 transversis, ultimo maiusculo, praecedentibus ambobus longiore, subconico, piceae, basi testaceae. Palpi testacei. Caput deflexum, utrinque punctis raris impressum. Thorax latitudine plus dimidio brevior, coleopte-

ris basi latitudine subaequalis, antrorsum leviter angustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis omnibus obtusissimis, anterioribus deflexis, transversim convexus, dorso obsolete bilineatim impressus, in lineis obsolete irregulariter punctatus, punctis quatuor (binis) maioribus, quadratim positis, in medio dorsi notatus, lateribus parce vageque subtilissime punctulatus, punctoque uno alterove maiore impressus. Elytra thorace paulo breviora, angulo apicali exteriore rotundato, sat fortiter profundeque minus dense punctata, macula magna apicali rufo-testacea. Abdomen parallelum, parcius punctatum, subtus apice et singulorum segmentorum margine apicali lato ferrugineis. Pedes testaceo-rufi.

\*20. *A. nitida*: *Nigra, nitida, thorace medio biserialim punctato, elytris thoracis longitudine, profunde punctatis, macula apicali rufa, abdomine supra parcius punctato.* — Long.  $1\frac{1}{4}$  —  $1\frac{3}{4}$  lin.

Grav. Micr. 97. 44. Mon. 172. 63. — Latr. Hist. nat. Crust. et Ins. IX. 390. 44. — Gyll. Ins. Suec. II. 435. 57. — Sahlb. Ins. Fenn. I. 401. 98. — Boisd. et Lacord. Faun. Ent. Paris. I. 531. 8. — Mannerh. Brachelytr. 63. 14. — Er. Col. March. I. 358. 8.

*Var. Elytris immaculatis.*

*Al. bilineata* Gyll. Ins. Suec. II. 436. 58. — Sahlb. Ins. Fenn. I. 401. 98. — Boisd. et Lacord. Faun. Ent. Paris. I. 532. 9. — Mannerh. Brachelytr. 68. 15.

Habitat in Europa et in Oriente.

Nigra, nitida, parce griseo-puberula. Antennae capite sesqui longiores, articulo secundo tertioque subaequalibus, apicem versus subincrassatae, articulis 5—10 transversis, ultimo praecedentibus ambobus magnitudine aequali, subovato, nigrae. Palpi maxillares nigro-picei, articulo terminali pallide testaceo. Caput deflexum, fronte utrinque parce punctata. Thorax coleopterorum latitudine, hac duplo brevior, basi leviter, lateribus modice rotundatus, antrorsum subangustatus, apice truncatus, angulis omnibus obtusissimis, subrotundatis, transversim convexus, antrorsum subcompressus, medio biserialim, lateribus vage punctatus. Elytra thoracis longitudine, angulo apicali exteriore subsinuato, profunde fortiterque minus dense punctata, interstitiis laevibus, apice macula magna rufa notata. Abdomen parallelum, parcius punctatum. Pedes picei, geniculis tarsisque dilutioribus.

Elytra interdum fere tota testacea, saepius etiam tota nigra.

Obs. In Gravenhorstii Mon. allegatus apud hanc speciem Staph. bipunctatus Oliv. et Staph. bipustulatus Lin. Ille, nisi ad *Al. bipunctatam* Grav., potius ad *Al. tristem* pertinere mihi videtur, Staph. bipustulatus Lin. (Faun. Suec. 232. 847. Syst. Nat. I. II. 685. 16) autem ob „magnitudinem minoris pediculi, antennis clavatas et elytra breviora” forte apud hanc speciem respiciendus esset, nisi punctum ferrugineum in elytri singuli medio notatum esset, quare hoc citatum nimis dubium existimo.

\*21. *A. lanuginosa*: *Nigra, nitida, pube erecta vestita, antennis totis nigris, elytris thoracis longitudine, crebre fortiter punctatis, apice rufescentibus, abdomine parallelo, supra parcius punctato.* — Long. 2 lin.

Grav. Micr. 94. 48. Mon. 171. 62. — Latr. Hist. nat. Crust. et Ins. IX. 388. 8. — Gyll. Ins. Suec. II. 432. 54. — Sahlb. Ins. Fenn. I. 398. 93. — Germ. Faun. Ins. Europ. XIII. 9. — Boisd. et Lacord. Faun. Ent. Paris. I. 530. 5. — Mannerh. Brachelytr. 67. 8. — Runde Brachelytr. Hal. 28. 6. — Er. Col. March. I. 356. 5.

Habitat in Europa.

Nigra, capite, thorace elytrisque subaeneis, nitida, pube erecta grisea

vestita. Antennae capite duplo longiores, articulo secundo tertio subaequali, apicem versus leviter incrassatae, articulis penultimis longitudine vix crassioribus, ultimo praecedentibus duobus coniunctis longitudine prope aequali, ovato-subconico, nigrae. Palpi picei. Caput deflexum, fronte parcius punctata. Thorax latitudine plus dimidio brevior, vix coleopterorum latitudine, basi et lateribus rotundatus, antrorsum leviter angustatus, apice truncatus, angulis anterioribus obtusis, deflexis, posterioribus rotundatis, leviter transversim convexus, crebre punctatus, basi foveis duabus approximatis longitudinalibus obsoletis vel obsoletissimis, at densius fortiusque punctatis. Elytra thoracis longitudine, vel eodem sublongiora, apice ad angulum anteriorem subsinuata, fortiter profunde denseque punctata, interstitiis planis, margine apicali rufescente. Abdomen parallelum, supra segmentis singulis basi densius, apice parcius punctatis, subtus fortius punctatum, segmentis singulis margine posteriore piceo. Pedes nigri, tibiis piceis, tarsis rufescentibus.

Variat 1, antennis elytrisque rufo-piceis, pedibus rufis.

2, antennis rufo-piceis, elytris, ano pedibusque rufo-testaceis.

Obs. Pube praecipue thoracis erecta ab affinibus distinguenda.

\* 22. *A. moerens*: Nigra, nitida, antennarum basi, pedibus anoque rufo-testaceis, elytris thoracis longitudine, dense punctatis, castaneis, abdomine apicem versus subangustato, parcius profundiusque punctato. — Long. 2 lin.

Gyll. Ins. Suec. IV. 493. 53—54. — Sahlb. Ins. Fenn. I. 398. 91. — Mannerh. Brachel. 66. 6. Aleoch. fumata var. 1 Grav. Micr. 96. 42.

Habitat in Europa.

Nigra, nitida, tenuiter griseo-puberula. Antennae capite thoraceque paulo breviores, articulo secundo tertioque aequalibus, apicem versus sensim incrassatae, articulis 5 — 10 sensim fortius transversis, ultimo praecedentibus ambobus longitudine aequali, acuminato, piceae, basi rufo-testaceae. Palpi rufo-testacei. Caput deflexum, parce obsoleteque punctatum, ore piceo. Thorax latitudine plus dimidio brevior, basi coleopterorum latitudine, antrorsum angustatus, basi et lateribus modice rotundatus, apice truncatus, angulis posterioribus obtusis, anterioribus subrotundatis, deflexis, modice transversim convexus, crebrius punctatus, lateribus interdum piceus. Elytra thoracis longitudine, apice ad angulum anteriorem leviter sinuata, dense punctata, punctis obliquis, rufo-picea castaneave. Abdomen apicem versus sensim subangustatum, subtus densius, supra parcius longius pilosum, subtus crebre, supra segmentis anterioribus crebrius, posterioribus parcius punctatis, ano rufo-testaceo, segmentis ventralibus piceo-marginatis. Pedes rufo-testacei.

Mas abdominis segmento ventrali sexto apice medio triangulariter producto distinctus.

Obs. 1. Statura intermedia quasi inter *A. fumatam* et *moestam*, ab illa elytris apice subsinuatis, ab hac thorace abdomineque apicem versus subangustatis, hoc crebrius punctato distinguenda. Ab *Al. lanuginosa* pube depressiuscula et elytris dense minusque fortiter punctatis discedit.

2. Specimina Suecica, a Gyllenhalio communicata dedit Dom. Schüppel.



- \* 23. *A. moesta*: *Elongata, subdepressa, nigra, nitida, parce subtiliter punctata, antennarum basi pedibusque piceis, elytris thoracis longitudine, fortius crebre punctatis, abdomine supra sublaevigato.* — Long.  $1\frac{1}{2}$  —  $2\frac{1}{4}$  lin.

Grav. Micr. 96. 41. Mon. 170. 55. — Latr. Hist. nat. Crust. et Ins. I. 389. 41. — Er. Col. March. I. 358. 9.

*Al fumata* Gyll. Ins. Suec II. 434. 56. — Sahlb. Ins. Fenn. I. 400. 96. — Zetterst. Faun. Lappon. I. 105. 25. Ins. Lapp. 79. 36. — Mannerh. Brachelytr. 67. 10. — Runde Brachelytr. Hal. 29. 8.

*Al. haemorrhoidalis* Mannerh. Brachelytr. 67. 7. — Sahlb. Ins. Fenn. I. 398. 92.

Habitat in Europa.

Praecedentibus paulo magis elongata depiesaque, nigra, nitida, parcius griseo-puberula. Antennae capite thoraceque paulo breviores, articulo secundo tertioque subaequalibus, apicem versus sensim subincrassatae, articulis 5 — 10 leviter transversis, ultimo praecedentibus ambobus longitudine aequali, ovato-subconico, nigrae aut piceae, basi piceae aut rufo-piceae, articulo primo dilutiore, rufo. Palpi rufi. Caput parce punctatum. Thorax coleopteris paulo angustior, latitudine dimidio brevior, basi et lateribus leviter rotundatus, antrorsum subangustatus, apice truncatus, angulis omnibus obtusis, anterioribus leviter deflexis, parum convexus, parcius punctatus. Elytra thoracis longitudine, apice ad angulum anteriorem subsinuata, crebre profundius punctata, aut nigro-, aut rufo-picea. Abdomen longiusculum, parallelum, supra vix punctatum. Pedes picei aut rufo-picei.

Mas abdominis segmento ventrali sexto apice medio triangulariter producto distinctus.

Obs. Proxime affinis *A. lanuginosae*, at statura minore, graciliore, et abdomine supra nonnisi raris punctis impresso distinguenda.

- \* 24. *A. ruficornis*: *Ferruginea, capite, thoracis disco abdominisque segmentorum intermediorum basi piceis, antennis elongatis, elytris thorace longioribus.* — Long.  $2\frac{2}{3}$  lin.

Grav. Micr. 91. 34. Mon. 169. 51. — Latr. Hist. nat. Crust. et Ins. IX. 387. 34.

Habitat in Germania media.

Statura magna *A. fuscipedis*, at gracilior, et antennis multo longioribus discedens. Rufo-testacea vel rufo-ferruginea, pube flavescente parcius vestita. Antennae capite thoraceque paulo longiores, graciles, articulo secundo tertio paululum brevior, apicem versus sensim subincrassatae, articulis 6 — 10 longitudine perparum crassioribus, ultimo praecedentibus ambobus longitudine aequali, sensim acuminato, ferrugineae, articulo ultimo testaceo. Caput deflexum, parcius punctatum, nigrum, ore rufo. Thorax latitudine dimidio prope brevior, coleopterorum fere latitudine, antrorsum vix angustatus, lateribus et basi leviter rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, minus convexus, crebre punctatus, disco picescente. Elytra thorace sublongiora, apice ad angulum anteriorem subsinuata, creberrime punctata. Abdomen parallelum, supra parcius punctatum, segmentis intermediis basi picescentibus. Pedes dilute testacei.

Mas abdominis segmento dorsali secundo apice dente erecto, apice truncato, tertio medio tuberculo minuto acutiusculo, sexto ante apicem me-

dio transversim elevato, sexto ventrali apice bisinuato, medio leviter triangulariter producto insignis.

\* 25. *A. erythroptera*: *Nigra, antennarum basi, elytris, ano pedibusque rufis, antennis elongatis, elytris thorace longioribus. — Long. 2 lin.*

*Grav. Mon.* 158. 26.

*Al. moesta* var. *b* *Er. Col. March.* I. 359. 10.

Habitat in Germania.

Habitu omnino praecedentis, colore, antennis apicem versus magis incrassatis etc. distincta, nigra, minus nitida, parcius flavescenti-puberula. Antennae capite thoraceque longiores, articulo secundo tertio paulo breviori, apicem versus sensim incrassatae, articulis 4—10 subobconicis, penultimis etiam crassitie vix brevioribus, ultimo ovato, fuscae vel rufo-piceae, articulis tribus primis rufis. Palpi rufi. Caput deflexum parce obsoleteque punctatum, fronte convexa, inter antennis leviter impressa. Thorax latitudine dimidio brevior, basi coleopterorum latitudine, antrorsum angustatus, basi et lateribus modice rotundatus, apice truncatus, angulis posterioribus obtusis, anterioribus rotundatis, deflexis, transversim convexus, minus crebre punctulatus, lateribus interdum obscure piceus. Scutellum confertissime punctatum, nigrum. Elytra thorace paulo longiora, apice ad angulum anteriorem subsinuata, dense fortius punctata, laete rufa, parum nitida. Abdomen apicem versus sensim subangustatum, subtus parcius, supra parce punctatum, nigrum, nitidum, ano et subtus segmentorum singulorum margine posteriore lato rufis. Pedes toti cum coxis rufo-testacei.

Ma s abdomine supra segmento secundo et sexto ante apicem medio tuberculo minuto munitis, subtus segmentis secundo tertioque basi transversim fortiter impressis, in impressione fortius villosis, sexto apice medio triangulariter producto distinctus. Antennae articulo ultimo maris praecedente duplo, feminae sesqui longiore.

Obs. 1. *Oxyporus erythropterus* Panz., a Gravenhorstio ad hanc speciem allegatus, in *Tachyporo* genere quaerendus.

2. Cum hanc speciem satis distinctam *A. moestae* subiungerem, feminam solum cognovi.

\* 26. *A. amplicolis*: *Nigra, nitida, parce subtilissime pubescens, antennis, pedibus anoque piceis, elytris thoracis longitudine, apice ad angulum anteriorem profundius sinuata, abdomine supra parcius punctato. — Long 2 lin.*

Habitat in Madagascar, Dom. Goudot.

Nigra, nitida, pube subtili depressa grisea parce vestita. Antennae capitis thoracisque fere longitudine, apicem versus leniter incrassatae, articulo tertio secundo paulo longiore, 4—10 sensim brevioribus et crassioribus, penultimis duobus crassitie paulo brevioribus, ultimo his ambobus longitudine aequali, oblongo, apice subacuminato, compresso, piceae, basi dilutiores. Caput parce subtiliterque punctatum, ore rufo-piceo. Thorax latitudine dimidio brevior, basi coleopterorum latitudine, antrorsum angustatus, lateribus et basi rotundatus, apice emarginatus, angulis omnibus obtusis.

anterioribus leviter deflexis, medio laevisimus, versus latera subtilissime punctulatus. Elytra thoracis longitudine, crebre punctata, basi secundum suturam elevata, apice ad angulum exteriorem profundius sinuata. Abdomen apicem versus subattenuatum, supra parcius subtiliusque punctatum, segmento quinto medio subtilissime longitudinaliter biseriatim tuberculato, subtus minus subtiliter pubescens, ano piceo. Pedes picei.

Obs. Specimen unicum, mutilatum, habitu fere *Gymnusae brevicollis*, at huius generis videtur.

- \* 27. *A. peltata*: *Nigra, nitida, antennarum basi, pedibus elytrisque rufo-piceis, his thorace sublongioribus, angulo posteriore exteriore nigricante, abdomine supra laevigato.* — Long. 2 lin.

Habitat in Mexico.

Nigra, nitida, pube depressiuscula, minus densa grisea vestita. Antennae capite duplo longiores, articulo secundo tertio paulo brevior, apicem versus incrassatae, articulis 6 — 10 transversis, crassitie duplo brevioribus, ultimo praecedentibus ambobus longitudine aequali, subovato, apice valde acuminato, nigrae, articulis duobus primis testaceis, ultimo apice flavescente. Palpi maxillares testacei, articulo tertio piceo. Mandibulae rufae. Caput deflexum, vage punctatum. Thorax latitudine plus dimidio brevior, coleopteris paululum angustior, basi, angulis posterioribus lateribusque rotundatis, antrorsum subangustatus, apice submarginatus, angulis anterioribus obtusis, leviter deflexis, parum convexus, sat crebre fortiter profundeque punctatus, margine basali elevato. Elytra thorace paululum longiora, apice ad angulum exteriorem subsinuata, minus dense fortiter punctata, brunneo-rufa, angulo posteriore exteriore late fuliginoso. Abdomen apice subangustatum, dorso laevigatum glabratumque, subtus parcius punctatum, longius pubescens, segmentis singulis utrinque margine posteriore rufo-piceo. Pedes rufo-picei.

- \* 28. *A. vaga*: *Fusca, antennarum basi apiceque, thoracis limbo, elytrorum vitta obsoleta, abdominis segmentorum marginibus pedibusque rufis, abdomine supra densius punctato.* — Long. 1½ lin.

Habitat ad Promont. bon. Spei, Dom. Bergius.

A. pulla paulo maior et multo latior, fusca, creberrime punctata, pube tenui grisea densius vestita. Antennae capite sesqui longiores, articulo secundo tertio paululum brevior, apicem versus sensim subincrassatae, articulis 5 — 10 transversis, longitudine dimidio crassioribus, ultimo oblongo, praecedentibus ambobus longiore, acuminato, basi rufae, articulo ultimo acuminato, ferrugineo. Caput subporrectum, parcius punctulatum, nigrum, ore rufo. Thorax latitudine dimidio brevior, basi coleopterorum latitudine, antrorsum angustatus, basi et lateribus modice rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, transversim convexus, creberrime subtiliter punctatus, fuscus, limbo omni rufo. Elytra thoracis longitudine, apice ad angulum exteriorem leviter sinuata, creberrime punctata, vitta ab humero incipiente, arcuatim deflexa, obsoletiore, rufa. Abdomen breviusculum, apice subangustatum, supra profundius densiusque punctatum, nitidum, nigrum, segmentis margine rufis. Pedes rufi.



- \* 29. *A. deserta*: Nigra, antennarum basi apiceque pedibusque testaceis, thorace, elytris anoque dilute rufo-piceis, abdomine supra crebre punctato. — Long.  $1\frac{1}{2}$  lin.

Habitat in Arabia deserta, Dom. Prof. Ehrenberg.

Sequente vix longior, at paulo latior. Antennae capitis thoracisque longitudine, parum incrassatae, articulo secundo tertioque aequalibus, 5—10 transversis, longitudine sesqui fere crassioribus, ultimo praecedentibus ambobus paulo longiore, oblongo-ovato, apice acuminato, nigrae, articulis primis et ultimo testaceis. Caput subdeflexum, nigrum, subtiliter punctatum, ore testaceo. Thorax latitudine dimidio brevior, basi coleopterorum latitudine, apicem versus angustatus, basi et lateribus leviter rotundatus, angulis omnibus obtusis, anterioribus deflexis, transversim convexus, subtiliter confertim punctatus, rufo-piceus, limbo dilutiore. Elytra thoracis longitudine, apice ad angulum exteriorem subsinuata, confertim subtiliter punctata, dilute rufo-picea. Abdomen apicem versus subangustatum, supra fortius crebre punctatum, segmentis singulis piceo-marginatis, ano dilute rufo-piceo. Corpus totum pube tenui depressa fulvescente vestitum. Pedes testacei.

- \* 30. *A. pulla*: Nigra, subnitida, confertissime punctata, dense grisescenti-pubescentis, elytris thorace longioribus, brunneis, antennis nigris, basi apiceque pedibusque testaceis. — Long.  $1\frac{1}{2}$  lin.

Gyll. Ins. Suec. IV. 494. 56—57. — Mannerh. Brachelytr. 68. 13. — Boisd. et Lacord. Faun. Ent. Paris. I. 533. 10. — Er. Col. March. I. 359. 10.

Habitat in Germania, Suecia.

Nigra, parum nitida, confertissime punctata, pube brevi flavescenti-grisea nitida densius vestita. Antennae capite thoraceque parum breviores, crassiusculae, articulo secundo tertio paulo longiore, 4—10 fortiter transversis, longitudine duplo crassioribus, ultimo ovato, subacuminato, nigrae, articulis primis ultimoque rufo-testaceis. Palpi rufo-testacei. Caput deflexum, confertim fortiter profundeque punctatum, linea media longitudinali tenui laevi. Thorax brevis, latitudine duplo fere brevior, basi coleopteris vix angustior, apicem versus modice angustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, leviter transversim convexus, minus subtiliter dense aequaliterque punctatus. Elytra thorace paulo longiora, apice ad angulum exteriorem leviter sinuata, eodem modo punctata, aut nigro-fusca, aut rufo brunnea, thorace sesqui longiora. Abdomen apice subangustatum, utrinque dense et minus subtiliter punctatum, segmentis singulis margine apicali piceo. Pedes rufo-testacei.

31. *A. spissicornis*: Nigra, nitida, antennis crassis basi, pedibus, elytris anoque rufis, abdomine supra parce punctato. — Long. vix ultra 1 lin.

Habitat Lutetiae, coll. Dom. Aubé.

Minuta, antennarum structura praecipue insignis, nigra, nitida, pube depressa grisea vestita. Antennae capite thoraceque paululum breviores, crassae, apicem versus sensim crassiores, articulo secundo tertio sesqui longiore, hoc obconico, 5—10 fortiter transversis, longitudine triplo crassio-

ribus, perfoliatis, ultimo magno, subacuminato, rufo-piceae, basi testaceo-rufae. Caput deflexum, parce obsoleteque punctulatum, ore rufo-testaceo. Thorax latitudine duplo fere brevior, basi coleopterorum prope latitudine, apicem versus subangustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis posterioribus obtusis, anterioribus rotundatis, deflexis, modice transversim convexus, subtiliter punctulatus, niger, nitidus, lateribus piceus. Elytra thoracis longitudine, apice ad angulum anteriorem subsinuata, fortius oblique punctata, lateritia, nitida. Abdomen parallelum, supra parce fortius punctatum, ano rufo, subtus segmentis singulis margine apicali rufo-piceo. Pedes testaceo-rufi.

- \* 32. *A. laevigata*: Nigra, nitida, parce subtiliterque punctata, elytris thoracis longitudine, crebre punctatis pedibusque rufis, abdomine parallelum, supra parcius profundiusque punctato. — Long.  $1\frac{2}{3}$  lin.

Gyll. Ins. Suec. II. 433. 55. — Sahlb. Ins. Fenn. I. 399. 95. — Mannerh. Brachel. 68. 11.

Habitat in Suecia, Dom. Schüppel.

Statura omnino *A. nitidae*, nigra, nitida, parce subtiliterque pubescens. Antennae capite paulo longiores, articulo secundo tertioque aequalibus, apicem versus sensim leviter incrassatae, articulis 5 — 10 transversis, ultimo maiusculo, praecedentibus ambobus paulo longiore, subacuminato, nigrae, basi piceae. Palpi picei. Caput deflexum, utrinque subtilissime obsoleteque punctulatum. Thorax latitudine duplo brevior, coleopterorum latitudine, basi leviter, lateribus modice rotundatus, antrosum vix angustatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, transversim convexus, subtilissime obsoleteque parce punctatus. Elytra thoracis longitudine, apice ad angulum anteriorem leviter sinuata, dense fortius punctata, punctis obliquis, densius pubescentia, rufa, subnitida. Abdomen parallelum, minus confertim profundius punctatum, subtus segmentorum margine posteriore piceo. Pedes rufi.

- \* 33. *A. restricta*: Nigra, nitidula, griseo-pubescens, antennis, tibiis tarsisque rufis, elytris thorace sesqui longioribus, fortius punctatis, abdomine apicem versus attenuato, supra confertim punctato. — Long.  $1\frac{1}{3}$  lin.

Habitat in Brasilia, Dom. Prof. Germar.

Habitus et summa affinitas *A. morionis*, at paulo maior, nigra, nitidula, pube longiore depressa fusciscenti-grisea densius vestita. Antennae capite sesqui fere longiores, articulo secundo tertio duplo prope minore, apicem versus sensim incrassatae, articulis 4 — 10 transversis, sensim paulo crassioribus, ultimo praecedentibus ambobus prope longiore, apice abrupte acuminato, dilute rufae, basi testaceae. Palpi pallide testacei. Caput deflexum, subtilissime punctulatum. Thorax basi coleopterorum prope latitudine, antrosum leviter angustatus, latitudine baseos duplo fere angustior, basi medio rotundatus, utrinque subsinuatus, lateribus rotundatus, apice truncatus, angulis anterioribus deflexis, rotundatis, posterioribus subrectis, sat fortiter convexus, crebrius subtiliter punctatus. Elytra thorace sesqui longiora, apice ad angulum anteriorem sinuata, confertim fortius punctata, nigro-picea, humeris et margine summo apicali picescentibus. Abdomen apicem versus sensim attenuatum, subtilius crebre punctatum, segmentis singulis margine po-

steriore rufescentibus, apice summo rufo. Pedes picei, tibilis tarsisque dilute rufis.

\* 34. *A. morion*: *Nigra, subnitida, antennarum basi pedibusque rufis, elytris thorace longioribus, fortius punctatis, abdomine apicem versus attenuato, supra crebre punctato, apice rufo-piceo.* — Long.  $\frac{3}{4}$ — $1\frac{1}{4}$  lin.

Grav. Micr. 97. 45. Mon. 175. 69. — Latr. Hist. nat. Crust. et Ins. IX. 390. 45. — Sahlb. Ins. Fenn. I. 402. 100. — Mannrh. Brachel. 68. 16. — Er. Col. March. I. 361. 13.

*Aleocho. exigua* Mannrh. Brachelytr. 68. 17. — Sahlb. Ins. Fenn. I. 402. 101.

Habitat in Europa et in America meridionali.

Habitu omnino *A. bipunctatae*, at multo minor, nigra, subnitida, pube rara subtilique vestita. Antennae capitis thoracisque longitudine, articulo secundo tertioque aequalibus, apicem versus incrassatae, articulis 4 — 10 transversis, ultimo maiusculo, ovato, nigrae, basi rufae. Palpi rufo-picei. Caput deflexum, parce subtilius punctatum. Thorax latitudine duplo brevior, basi coleopterorum latitudine, modice rotundatus, utrinque subsinuatus, lateribus leviter rotundatus, antrorsum angustatus, apice truncatus, angulis omnibus obtusis, anterioribus deflexis, transversim convexus, confertim subtilius punctatus. Elytra thorace paulo longiora, apice ad angulum anteriorem sinuata, fortiter dense punctata, reliquo corpore densius pubescentia. Abdomen apicem versus attenuatum, supra sat crebre licet segmentis posterioribus minus dense et subtilius punctatum, apice rufo-piceum. Pedes rufi.

Obs. 1. *Al. exigua* antennis longioribus praecipue ab *Al. morione* discedens nisi sexu nullo modo ab hac differre videtur. Magnitudine haec species, uti plures huius generis valdopere variat, quod e longitudinis mensura indicata patet.

2. Specimina Americana cum Europaeis exactissime conveniunt; exemplum in Brasilia a Dom. Beske lectum dedit Dom. Thorey, alterum e Columbia a Dom. Moritz allatum. Forte etiam Americam septentrionalem habitat.

\* 35. *A. angulata*: *Rufo-brunnea, antennarum basi apiceque anoque ferrugineis, thorace depresso, obsolete canaliculato, angulis posticis acutis.* — Long.  $1\frac{1}{2}$  lin. Thy. 204

Er. Col. March. I. 360. 11.

Habitat Berolini; Lutetiae etiam in societate Formicae rufae lecta a Dom. Aubé.

Statura fere *A. pullae*, at supra magis plana, pube tenuissima depressa grisea parce vestita, rufo-brunnea. Antennae capite sesqui longiores, articulo secundo tertioque subaequalibus, apicem versus sensim leniterque incrassatae, articulis arcte contiguis, penultimis leviter transversis, ultimo praecedentibus tribus longitudine vix inferiore, oblongo, apice subacuminato, rufo-brunneae, articulis primis ultimoque ferrugineis. Caput parvum, subdeflexum, obsolete punctulatum, opacum, nigrum, ore rufo. Thorax coleopterorum latitudine, eaque duplo brevior, antrorsum angustatus, lateribus parum, basi praecipue medio fortius rotundatus, basi utrinque sinuatus, apice truncatus, angulis anterioribus obtusis, deflexis, posterioribus acutis, supra subdepressus, obsolete canaliculatus, basi distincte transversim impressus, subtiliter obsoleteque punctatus, opacus. Elytra thorace vix longiora, apice prope angulum anteriorem sinuata, dense punctata, opaca. Abdomen apice



angustatum, infra longius pubescens, supra parcius punctatum, nitidulum, ante apicem obscurum, apice late rufo-testaceum. Pedes rufo-testacei.

- \* 36. *A. praetexta*: Nigra, nitida, antennis, elytrorum limbo omni, abdo-  
minis apice segmentorumque marginibus pedibusque rufis, thorace brevi,  
angulis posticis acutiusculis. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 361. 12.

Habitat in Germania, Gallia.

Linearis, nigra, nitida, pube depressa flavescente parcius vestita. Antennae capite thoraceque breviores, articulo secundo tertio paulo maiore, 4 — 10 transversis, arcte contiguis, ultimo ovato, apice compresso, totae rufae. Caput porrectum, maiusculum, obsolete punctatum, ore rufo. Thorax coleopterorum latitudine, eaque duplo brevior, antrorsum vix angustatus, basi modice, lateribus leviter rotundatus, basi utrinque leviter sinuatus, angulis posterioribus acutiusculis, apice truncatus, angulis anterioribus obtusis deflexis, transversim parum convexus, parcius subtiliter punctatus. Elytra thorace sesqui longiora, apice ad angulum anteriorem profundius sinuata, subtiliter punctata, limbo suturali, apicali et lateribus rufis. Abdomen parallelum, parcius punctatum, segmentis 5 primis rufo-marginatis, ultimis duobus totis rufo-testaceis. Pedes rufi.

- \* 37. *A. obscurella*: Atra, opaca, pube brevi crassiuscula grisea densius vestita, antennis pedibusque rufo-piceis, elytris thorace sesqui longioribus. — Long.  $1\frac{2}{3}$  lin.

Grav. Mon. 159. 27. — Gyll. Ins. Suec. II. 403. 25.

*Oxyroda sericea* Boisd. et Lacord. Faun. Ent. Paris. I. 533. 2.

Habitat in Germania septentrionali, Suecia, Gallia.

Elongata, subdepressa, atra, opaca, pube brevi, parum depressa, grisea, micante densius vestita. Antennae capite parum longiores, rufo-piceae, articulo secundo tertio longitudine aequali, licet paulo crassiore, apicem versus sensim subincrassatae, articulis 5 — 10 transversis, ultimo ovato-subconico. Palpi rufo-picei. Caput thorace parum angustius, porrectum, rotundatum, fortius punctatum, linea media subelevata laevi. Thorax coleopterorum latitudine, eaque plus dimidio brevior, antrorsum vix angustatus, lateribus et basi leviter rotundatus, apice truncatus, angulis anterioribus obtusis, deflexis, posterioribus subrotundatis, transversim parum convexus, parce subtiliterque punctatus. Elytra thorace sesqui longiora, apice truncata, crebre punctata, picescentia. Abdomen parallelum, crebrius punctatum, subtilius pubescens, supra nitidulum. Pedes picei, geniculis tarsis rufescentibus.

#### Species mihi invisae.

38. *A. haemorrhoidalis*: Nigro-coerulea, nitida, antennis fusiformibus, capite, thorace elytrisque crebre punctatis, abdomine nigro, apice ferrugineo, pedibus atris.

Guérin Voy. d. l. Coquille, Ins. t. 1. f. 24. — Boisd. Faun. Ent. Ocean. 57. 1.

Habitat in Nova Hollandia. (Port Jackson.)

Obs. A chrysorrhoeae affinis videtur.

39. *A. villosa*: *Sublinearis, nigra, opaca, creberrime punctata, pube brevi fusca undique tecta, antennis thorace longioribus basi pedibusque piceis.* — Long.  $1\frac{3}{4}$  lin.

Mannerh. *Brachelytr.* 67. 9. — Sahlb. *Ins. Fenn. I.* 399. 94. — Zetterst. *Ins. Lappon.* 79. 35. — Runde *Brachelytr. Hal.* 29. 7.

Habitat in Fennia, Lapponia.

Ab *A. lanuginosa* differre videtur corpore angustiore, magis lineari, antennis thorace longioribus, basi piceis, thorace elytris fere angustiore, margine elytrorum nunquam rufo, et pubescentia brevior, fusciscente. De cetero eidem adeo similis est, ut ulteriori non opus sit descriptione. (Sahlb.)

Obs. Mihi omnino dubia.

40. *A. crassicornis*: *Nigra, nitida, antennis crassis, fuscis, pedibus elytrisque rufo-testaceis, his margine laterali infuscato.* — Long.  $1\frac{1}{4}$ — $1\frac{1}{3}$  lin.

Boisd. et Lacord. *Faun. Ent. Paris. I.* 531. 7.

Habitat Lutetiae.

*A. rufipenni* quodammodo affinis, at multo minor. Antennae breves, crassae, fuscae. Caput rotundatum, nigro-subaeneum, nitidum, subtilissime punctatum et pubescens, palpis maxillaribus fuscis, labialibus testaceis. Thorax ut in praecedentibus, subtilissime punctatus et pubescens, nigro-subaeneus, nitidus, margine laterali subferrugineo. Scutellum triangulare, laeve, piceum. Elytra dilute testacea, margine laterali magis minusve anguste fuliginoso, apice truncata, depressa, thorace et capite fortius punctata, tenuiter pubescentia. Abdomen totum nigro-subaeneum, nitidissimum, elongatum, punctatum et pubescens ut in praecedentibus (*A. rufipenni*, *lanuginosa* etc.) Pedes rufo-testacei. (Boisd. et Lacord.)

Obs. An *A. tristis* var. minor, capitis thoracisque punctura obsoleta?

## Phytosus Rudd.

Curt. Britt. *Ent. XV. pl.* 718.

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula subintegra; paraglossae angustae, acuminatae.*

*Palpi labiales 3-articulati, articulo secundo et tertio longitudine aequalibus.*

*Tarsi antici 4-, posteriores 5-articulati, postici articulis 4 primis aequalibus.*

*Tibiae anteriores spinulosae.*

Corpus elongatum, lineare, subdepressum. Caput exsertum, porrectum, basi haud constrictum, thoracis latitudine. Labrum apice truncatum, mandibularum apices haud obtegens. Mandibulae muticae. Maxillae mala exteriori cornea, basi apiceque membranacea, portione membranacea apicali pubescente, mala interiore exteriori vix brevior, membranacea, margine exteriori corneo, apice oblique introrsum fortius emarginata, sinu spinulis brevibus ciliato. Palpi maxillares elongati, articulo tertio secundo paulo longiore, apicem versus sensim leniter incrassato, quarto minuto, subulato. Labium mento transversum, antrorsum angustatum, apice late emarginato, ligula brevi, apice vix emarginata, paraglossis sat elongatis, angustis, acuminatis, interne ciliatis. Palpi labiales triarticulati, articulo primo secundoque minus perspicue distinctis, hoc paulo brevior, tertio secundo longitudine subaequali,

basi tenuiore, apice incrassatulo. Antennae breviores, articulis duobus primis elongatis. Thorax coleopteris paulo angustior, basin versus subangustatus. Scutellum triangulare. Elytra truncata. Abdomen elongatum, lineare, segmento septimo distincte conspicuo. Pedes breves, intermediū basi approximati, tibiis anterioribus externe seriatim sat fortiter spinulosi, posticis muticis, tarsis brevibus, anticis 4-, posticis 5-articulatis, articulis primis brevibus, aequalibus, ultimo elongato, illis coniunctis fere aequali.

#### O b s e r v a t i o n e s.

1. Maxillae fere ut in *Homalota* (t. III. f. 22.), at mala interior apice fortius sinuata, et palpi maxillares articulo tertio magis elongato, haud tumidulo. Palpi labiales ut in *Falagria* (t. III. f. 3.), at articulo secundo tertio haud brevior, paraglossae ut in *Autalia* (ibid. f. 2.), mentum ut in *Bolitochara* (ibid. f. 4); ligula mihi forte haud satis accurate observata, at brevis et latiuscula mihi visa est, ut in *Aleochara*, at apice haud biloba, immo vix emarginata.

2. Habitu fere *Pronomaeae*, at corpus dense sericeo-pubescent. Proxime affinis videtur *Aleocharae*, (v. c. *Al. obscurellae*), at palporum labialium et tarsorum structura discedit. Tarsorum structura cum *Homalota* congruit, differt autem paraglossis distinctis. Tibiis anterioribus fortiter spinulosi hoc genus in hac tribu valde distinctum, tibiis anticis fossoriis et abdominis segmento septimo conspicuo pulchre *Oxytelinorum* tribum in animum revocans, nuperrime ab auctore ipso benevole mihi communicatum hoc loco inserui, cum non modo ligulae structura intermedium inter *Aleocharam* et sequentia genera, sed summa etiam affinitatis specie *Aleocharae obscurellae* iunctum videatur.

3. Sexus differentia infra indicata.

\* 1. *Ph. spinifer*: *Niger*, dense cinereo-pubescent, antennis pedibusque rufis. — Long.  $1\frac{1}{3}$  lin.

Curtis Britt. Ent. XV t. 718.

Habitat in Anglia, Dom. Rudd.

Cinerascenti-niger, vix nitidus, pube minus subtili, depressa, cinerea, sericea dense vestitus. Antennae capite sesqui longiores, apicem versus parum incrassatae, articulis duobus primis maioribus, longitudine fere aequalibus, primo crassiusculo, secundo elongato-obconico, tertio obconico, secundo duplo brevior, sequentibus breviusculis, sensim paulo crassioribus, penultimis 4 transversis, ultimo praecedentibus ambobus longitudine aequali, ovato, apice obtuse acuminato, dilute rufae, articulo ultimo fusco. Caput thoracis prope latitudine, rotundatum, confertim subtiliter punctatum, fronte leviter convexa, ore subacuminato, rufescente. Thorax coleopteris tertia parte fere angustior, latitudine haud brevior, basin versus subangustatus, lateribus anterieus rotundatis, posterius subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus obtusis, posterioribus subrectis, leviter convexus, confertim subtiliter punctatus, obsolete canaliculatus, basi leviter foveolatus. Elytra subdepressa, confertim subtiliter punctata. Abdomen confertim subtiliter punctatum, segmento quinto margine apicali subtestaceo. Pedes pallide rufi, femoribus nigricantibus.

Mas Alatus, elytris thoracis longitudine.

Femina Aptera, elytris thorace dimidio brevioribus.

Variat minus maturus testaceus, abdomine ante apicem capiteque nigricantibus.



## Oligota Mannerh.

Mannerh. *Brachelytr.* p. 72. — *Er. Col. March. I.* p. 362.

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula brevis bifida; paraglossae nullae.*

*Palpi labiales 3-articulati, articulis duobus primis crassiusculis, subaequalibus, tertio tenui, acuminato.*

*Antennae 10-articulatae, apice fortiter incrassatae.*

*Tarsi omnes 4-articulati, postici articulo primo elongato.*

Corpus minimum, vel elongatum et sublineare, vel subovatum, alatum. Caput thorace angustius, deflexum, basi haud constrictum, oculis mediocribus, parum prominulis. Labrum transversum, breve, apice truncatum, mandibularum apices haud obtegens. Mandibulae muticae. Maxillae malis breviusculis, membraneis, exteriori apice pubescente, interiori huic subaequali, apice introrsum oblique subemarginata, sinu spinulis brevibus ciliato. Palpi maxillares modice elongati, articulo secundo tertioque longitudine subaequalibus, hoc sat tumidulo, quarto minuto, subulato. Labrum (t. III. f. 10.) mento transversum, apice leviter emarginato, ligula minuta, bifida, paraglossis nullis. Palpi labiales triarticulati, articulo primo secundoque grossiusculis, inter se aequalibus, tertio minuto, subulato. Antennae breviusculae, 10-articulatae, articulis duobus primis crassiusculis, reliquis longioribus, sequentibus minutis, ultimis 3 vel 4 clavum oblongam formantibus. Thorax perbrevis, basi coleopteris latitudine subaequalis, leviter bisinnotatus, modice convexus. Coleoptera apice coniunctim late emarginata. Abdomen vel lineare, vel apicem versus sensim angustatum. Pedes breviusculi, intermedii basi distantes, tarsi omnes 4-articulati, postici articulo primo subelongato.

### Observationes.

1. Maxillae omnino ut in *Homalota* (t. III. f. 22), palpi maxillares autem articulo ultimo longiore aciculari differunt. Labii structura pari modo cum *Homalotis* satis convenit, palpi labiales vero itidem aliter constructi. Ceterum hoc genus admodum insigne antennis clavatis 10-articulatis et tarsis omnibus 4-articulatis.

2. Habitu hoc genus, speciebus minimae magnitudinis compositum, duas continet formas. Alteram constituunt *O. pusillima*, *atomaria* et *subtilis*, corpore elongato, lineari, abdomine apicem versus haud angustato, alteram *O. granaria*, *flavicornis* et *apicata*, corpore crassiusculo breviusculoque, utrinque attenuato, forma *Tachyporina*, ceterum autem omnibus characteribus cum illis congruentia, ut speciebus tam paucis vix sectione distinguantur. Antennarum tarsorumque structura *Oligotae* insigniter cum *Hypocrypto*, *Tachyporinorum* genere conveniunt, cuius habitum *O. granaria*, *flavicornis* et *apicata* concinne simulant.

3. Sexus differentia latet.

4. Victus aliis sub arborum cortice, aliis in pratis.

5. Genus *Microcera* Mannerh. (*Brachelytr.* p. 72.) *Oligotae* affine videtur et dubito an revera sit distinctum. Convenit palporum maxillarium structura et antennarum articulis ultimis crassioribus, differt praecipue corpore anterieus attenuato, postice latiore, ano truncato.

\* 1. *O. pusillima*: *Linearis, nigra, nitida, antennis, pedibus anoque rufis, antennarum articulis ultimis tribus abrupte crassioribus.* — Long.  $\frac{1}{2}$  lin.

Mannerh. *Brachelytr.* 72. I. — *Er. Col. March. I.* 363. I.

*Aleoeh. pusillima* Grav Mon. 175. 71. — *Gyll. Ins. Suec. IV.* 491. 38 — 39. — *Sahlb. Ins. Fenn. I.* 384. 70.

## Habitat in Europa.

Corpus angustum, nigrum, nitidum, parce subtiliter pubescens. Antennae capite sesqui longiores, articulis duobus primis maiusculis crassiusculisque, 3—7 minutis, 7to praecedentibus subaequali, ultimis tribus multo maioribus, 8to et 9to aequalibus, fortiter transversis, ultimo his ambobus parum minore, subgloboso, ferrugineae, clava fuscescente. Caput thorace paulo angustius, deflexum, subtilissime punctulatum, ore ferrugineo. Thorax coleopteris haud angustior, latitudine duplo brevior, basin versus haud angustatus, basi medio leviter rotundatus, utrinque subsinuatus, lateribus leviter rotundatus, apice truncatus, angulis omnibus obtusis, subrotundatis, anterioribus fortius deflexis, convexus, subtilissime punctulatus. Elytra thorace sesqui longiora, apice coniuncta late emarginata confertim fortius punctata. Abdomen parallelum, confertim subtiliter punctatum, apice rufescens. Pedes rufi.

Obs. 1. Al. pusillimam Gyll olim O. subtili allegavi, specimina autem Suecica, quae Dom. Schüppel a Dom. Schönherr sub hoc nomine accepit, ad hanc speciem pertinent, neque descriptio Gyllenhaliana huic contraria.

2. Statura magis lineari praecipue a proxime sequentibus distincta.

\* 2. *O. atomaria*: *Linearis, nigra, pedibus antennisque fusco-piceis, his articulis ultimis tribus abrupte crassioribus.* — Long.  $\frac{1}{2}$  lin.

Er. Col. March. I. 363. 2.

Habitat in Germania; specimen singulum Berolini legi.

Praecedenti affinis, at paulo latior, nigra, nitidula, parce subtiliter pubescens. Antennae capite sesqui longiores, articulis duobus primis maiusculis crassiusculisque, 3—7 minutis, 6 et 7 praecedentibus vix crassioribus, 8—10 clavam validam oblongam formantibus, 8 et 9 crassitie brevioribus, 10 his ambobus paulo minore, breviter ovato, apice obtusiusculo, piceae, basi testaceae. Palpi fusco-picei. Caput thorace duplo prope angustius, deflexum, subtiliter punctatum, ore piceo. Thorax coleopterorum prope latitudine, antrorsum modice angustatus, latitudine baseos duplo, apicis dimidio brevior, basi medio late rotundatus, utrinque subsinuatus, lateribus rotundatus, apice truncatus, angulis omnibus obtusis, anterioribus leviter deflexis, leviter transversim convexus, confertim subtiliter punctatus. Elytra thorace duplo longiora, confertim minus subtiliter punctata. Abdomen apicem versus subangustatum, confertim subtilius punctatum, apice piceum. Pedes piceo-testacei.

\* 3. *O. subtilis*: *Linearis, picea, elytris anoque rufo-piceis, pedibus antennisque rufis, his articulis ultimis quatuor sensim crassioribus.* — Long.  $\frac{1}{2}$  —  $\frac{1}{3}$  lin.

Er. Col. March. I. 364. 3.

Habitat in Germania.

Statura praecedentium, at saepius paulo maior, nigro-picea, pube subtili depressa densius vestita, elytris, ano, interdum etiam thorace rufo-piceis. Antennae capite sesqui longiores, articulis duobus primis maiusculis, 3—6 minutis, licet sexto praecedentibus paulo crassiore, septimo intermedio inter praecedentes et sequentes, illis distincte maiore, his distincte minore, ultimis

tribus reliquis crassioribus, penultimis duobus transversis, ultimo his ambobus magnitudine aequali, subconico, rufae, basi pallide testaceae. Caput thorace plus dimidio angustius, deflexum, subtilissime punctulatum, ore testaceo. Thorax elytris paulo angustior, antrorsum subangustatus, latitudine baseos duplo, apicis plus dimidio brevior, basi medio leviter rotundatus, utrinque subsinuatus, lateribus leviter rotundatus, apice truncatus, angulis anterioribus fortiter deflexis, subrotundatis, posterioribus rectis, transversim convexus, subtilissime punctatus. Elytra thorace duplo fere longiora, confertim subtilius punctata. Abdomen lineare, confertim subtiliter punctatum, ano late piceo. Pedes rufo-testacei.

Obs. *O. pusillimae* valde affinis et aegre distinguenda, praecipue corpore paulo latiore, thorace antrorsum leviter angustato, minus convexo, angulis posterioribus rectis, elytris longioribus et antennarum structura differe videtur.

\* 4. *O. granaria*: *Ovata, nigra, pedibus antennisque ferrugineis, his articulis ultimis quatuor sensim crassioribus.* — Long.  $\frac{1}{2}$  lin.

Er. Col. March. I. 364. 4.

Habitat in Germania; Berolini specimina duo legi.

Statura omnino sequentis, at sesqui maior, nigra, nitidula, subtiliter parce pubescens, confertim subtiliter punctulata. Antennae breves, crassiusculae, articulo secundo primo multo maiore, obconico, 4—6 minutis sensim paulo crassioribus, 7—9 magnis et crassis, itidem sensim paulo crassioribus, transversis, ultimo maiore, praecedentibus ambobus longitudine aequali, at paulo crassiore, subovato, apice obtuse acuminato, obscure ferrugineae. Caput thorace multo angustius, deflexum, ore piceo-testaceo. Thorax basi coleopterorum prope longitudine, antrorsum fortiter angustatus, latitudine baseos duplo, apicis vix dimidio brevior, apice truncatus, lateribus et angulis anterioribus rotundatis, basi medio leviter rotundatus, angulis posterioribus subrectis, leviter convexus. Coleoptera thorace duplo prope longiora, planiuscula, lateribus subrotundata, thorace distinctius punctata. Abdomen apicem versus attenuatum, apice piceum. Pedes ferruginei.

Obs. Sequenti affinis, at structura et colore antennarum satis distincta.

\* 5. *O. flavicornis*: *Ovata, nigra, pedibus ferrugineis, antennis flavis, articulis ultimis tribus abrupte crassioribus.* — Long. vix  $\frac{1}{2}$  lin.

Er. Col. March. I. 364. 5.

*Hypocyphus flavicornis* Boisd. et Lacord. Faun. Ent. Paris. I. 521. 4.

Habitat in Germania, Gallia.

*O. pusillima* longitudine fere aequalis, at plus sesqui latior, utrinque attenuata, statura fere Hypocypti, nigra, nitida, parce subtiliter pubescens. Antennae capite sesqui longiores, articulis duobus primis maiusculis, subaequalibus, 3—7 minutis, aequalibus, 8—10 abrupte multo crassioribus, magnis, 8 et 9 aequalibus, transversis, decimo magno, globoso, totae laete flavae. Palpi pallide flavi. Caput thorace multo angustius, deflexum, confertim subtiliter punctatum, ore flavo. Thorax basi coleopterorum prope longitudine, antrorsum angustatus, latitudine baseos duplo prope, apicis dimidio angustior, apice truncatus, lateribus parum rotundatis, angulis anterioribus subrectis, deflexis, basi medio leviter rotundatus, utrinque subsinuatus,



angulis posterioribus obtusis, subrotundatis, leviter convexus, confertim subtiliter punctatus. Coleoptera thorace sesqui longiora, lateribus subrotundata, medio thorace latiora, confertim minus subtiliter punctata. Abdomen breve, apicem versus attenuatum. Pedes ferruginei, posteriores fusciscentes.

Obs. *Hypocyphus flavicornis* Faunae Parisiensis teste Dom. Aubé certe ad hoc insertum revocandus.

\* 6. *O. apicata*: *Ovata, nigra, thorace elytrisque castaneis, ano late pedibusque fulvis.* — Long.  $\frac{1}{2}$  lin.

*Er. Col. March. I.* 365. 6.

Habitat in Germania: Berolini singulum specimen legi.

Statura praecedentis. Antennae mutilatae, articulis basalibus fulvis, secundo primo paulo longiore. Caput thorace angustius, deflexum, nigrum, nitidum, subtilissime punctatum, ore testaceo. Thorax basi coleopterorum prope latitudine, antrorsum angustatus, latitudine baseos duplo, apicis dimidio brevior, apice utrinque subsinuatus, lateribus et angulis anterioribus subrotundatis, basi medio leviter rotundatus, utrinque subsinuatus, angulis posterioribus subrectis, subtilissime punctatus, castaneus. Coleoptera thorace duplo longiora, lateribus subrotundata, medio thorace latiora, confertim punctata, subtiliter pubescentia, castanea. Abdomen apicem versus attenuatum, nigrum, segmentis duobus ultimis laete fulvis. Pedes fulvi.

### ***Gyrophæna Mannerh.***

*Mannerh. Brachelytr. p.* 74. — *Er. Col. March. I.* p. 365.

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula brevis, integra, rotundata; paraglossae nullae.*

*Palpi labiales 2-articulati, articulo primo cylindrico, secundo tenui, acuminato.*

*Tarsi antici 4-, posteriores 5-articulati, postici articulo primo elongato.*

Corpus parvum, latiusculum, subtus convexum, alatum. Caput exsertum, deflexum, basi haud constrictum, thorace paulo angustius, oculis plerumque sat magnis, modice prominulis. Labrum latiusculum, lateribus rotundatum, apice truncatum, mandibulas obtegens. Mandibulae muticae. Maxillae malis corneis, exteriore basi apiceque membranæ, membrana apicali malam interiorem superante, hac interne membranæ, apice oblique introrsum subemarginata, sinu spinulis confertis ciliato. Palpi maxillares parum elongati, articulo secundo tertioque subaequalibus, hoc leviter incrassato, quarto parvo, subalato. Labium mento perbreve, apice leviter emarginato, ligula brevi, rotundata, integra, paraglossis nullis. Palpi labiales biarticulati, articulo primo cylindrico, secundo hoc parum brevior, tenui, subacuminato. Antennae ab articulo quarto inde incrassatae, articulis tribus primis leviter elongatis, longitudine subaequalibus, tertio graciliore. Thorax coleopteris angustior, transversus, basi rotundatus, tenuiter marginatus. Scutellum transversim triangulare. Elytra truncata. Abdomen lateribus magis minusve rotundatum, supra planum, margine latiore, medio elevato, subtus convexum, in vivis reclinatum. Pedes breviusculi, tenues, intermediis basi modice distantes, tarsis anticis 4-, posterioribus 5-articulatis, posticis articulo primo parum elongato.

## O b s e r v a t i o n e s .

1. Maxillæ omnino ut in *Homalota* (t. III. f. 22). Labium ab eodem omnium præcedentium generum ligula integra et palpis labialibus biarticulatis discedit.

2. Habitus huius generis peculiaris corpore minuto, breviusculo, glabro, thorace coleopteris angustiore, transverso, subtiliter marginato, capite deflexo, fronte planiuscula, oculis prominulis, abdomine rotundato, lateribus late marginato, supra plano, subtus convexo, in vivis reclinato, antennis crassiusculis, pedibus intermediiis basi distantibus.

3. Genus *Encephalus* Kirby antennis brevioribus, apicem versus sensim incrassatis et thorace antorsum angustato haud satis distinctum et, reliquis omnibus notis habitualibus et pedum et oris structura pariter congruentibus, huic generi subiungendum mihi videtur.

4. Differentia sexualis abdomine tuberculis notato saepius indicata.

5. Victus in fungis.

\* 1. *G. complicans*: *Brevis, nigra, nitida, thorace basi subtiliter marginato, laevissimo, elytris medio brunneis, antennis pedibusque ferrugineis.* — Long. 1 lin.

*Er. Col. March. I.* 366. 1.

*Encephalus complicans* Westwood Guérin *Mug. Zool. III. cl. IX. t. 69.* — *Steph. Illustr. V. t. 26. f. 4.*

Habitat in Germania, Anglia.

Lata, crassiuscula, nigra, nitida. Antennae capitis longitudine, apicem versus sensim incrassatae, articulo tertio secundo plus duplo minore et tenuiore, 4 — 7 minutis, tenuioribus, at sensim paulo brevioribus et crassioribus, 8 — 10 inter se aequalibus, transversis, ultimo crassitie vix longiore, apice acuminato, ferrugineae. Caput laeve, ore ferugineo. Thorax basi fere coleopterorum latitudine, antorsum angustatus, latitudine baseos plus duplo brevior, basi leviter rotundatus, apice profundius sinuatus, lateribus parum rotundatis, cum angulis anterioribus productis acutis sat fortiter deflexis, angulis posterioribus obtusis, transversim convexus, laevis, basi medio supra scutellum leviter marginatus, margine summo piceo. Scutellum magnum, triangulare, laeve. Elytra brevissima, thorace vix longiora, laevia, medio simul depressa, brunnea, margine exteriori tumidulo, nigro. Abdomen piceum, laeve, subtus convexum, supra planum, margine laterali elevato, lato, rotundato, dentato. Pedes testacei, femoribus posticis nigricantibus.

Obs. Incedit hoc insectum abdomine reclinato, reliquum corpus obtegente, globuli instar, Agathidii cuiusdam speciem simulans. (Conf. Kirby et Spence *Introduct. II. p. 232.*)

\* 2. *G. nitidula*: *Brevis, nigra, nitida, antennarum basi, elytrorum disco pedibusque testaceis, thorace brevissimo, parce punctato, basi late marginato.* — Long. 1½ lin.

*Mannerh. Brachelytr. 74. 1.*

*Aleoch. nitidula* Gyll. *Ins. Suec. II.* 413. 35.

Habitat in Suecia, Dom. Schüppel.

Sequenti affinis, at duplo maior, nigra, nitida. Antennae capitis thoracis prope longitudine, articulo tertio secundo vix brevior, at multo tenuiore, quarto minuto, 5 — 10 inter se aequalibus, crassitie vix brevioribus, ultimo præcedente sesqui longiore, acuminato, densius pubescentes et

pilosellae, fusco-nigrae, articulis tribus primis pallide testaceis. Caput planiusculum, fronte utrinque longitudinaliter subimpressa et parce punctata, medio laevi, ore ferrugineo. Thorax latitudine plus duplo brevior, coleopteris angustior, basi et lateribus fortius rotundatus, apice truncatus, angulis anterioribus deflexis, posterioribus obtusis, subrotundatis, basi praesertim medio late marginatus, supra convexus, ad margines punctis paucis sparsis, medio punctis obsoletioribus crebris biserialis impressus, parce griseo-pubescentis. Elytra thorace sesqui longiora, subdepressa, humeris elevatis, crebre rugoso-punctata, tenuiter pubescentia, saturate rufo testacea, regione scutelli triangulariter fusca, angulo apicis exteriori late nigro. Abdomen ovatum, crassum, nigro-piceum, nitidum, segmentorum marginibus anoque subtus laete, supra obsoletius rufo-testaceis. Pedes rufo-testacei.

\* 3. *G. nana*: *Picea*, *antennis*, *pedibus*, *thoracis limbo*, *elytris abdominisque basi pallide testaceis*, *thorace marginato*, *parce punctato*, *elytris disco creberrime rugoso-punctatis*. — Long. 1 lin.

*Mannerh. Brachelytr.* 74. 2. — *Runde Brachel. Hal.* 20. 1. — *Er. Col. March.* I. 367. 2. *Staphyl. nanus* Payk. *Faun. Suec.* III. 408. 52.

*Aleochar. nana* Grav. *Micr.* 98. 47. *Mon.* 176. 72. — *Latr. Hist. nat. Crust. et Ins.* IX. 390. 46. — *Gyll. Ins. Suec.* II. 414. 36. — *Sahlb. Ins. Fenn.* I. 382. 66. — *Zetterst. Faun. Lappon.* I. 102. 19. *Ins. Lapp.* 78. 27.

*Gyrophæna nitidula* Boisd. et Lacord. *Faun. Ent. Paris.* I. 536. 2.

*Staphyl. fasciatus* Marsh. *Ent. Britt.* 514. 47.

Habitat in Europæ fungis.

Antennae capite thoraceque multo breviores, articulo tertio secundo multo minore, tenui, quarto breviter obconico, 5 — 10 aequalibus, transversis, ultimo praecedentibus ambobus longitudine subaequali, acuminato, pubescentes, totae flavo-testaceae. Caput nigrum, nitidum, fronte plana, utrinque leviter impressa et parce punctata, ore testaceo. Thorax coleopteris plus sesqui angustior, latitudine duplo prope brevior, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus subrectis, deflexis, posterioribus obtusis, utrisque subrotundatis, basi tenuiter marginatus, parum convexus, punctis paucis, dorso biserialis, terminalibus distinctis, reliquis obsoletis impressus, niger, nitidus, margine omni testaceo. Elytra thorace sesqui longiora, dense intricato-punctata, testacea, angulo apicis exteriori nigro. Abdomen basi testaceum, apice nigrum, ano iterum testaceo. Pectus nigrum. Pedes flavi.

Abdominis segmentum dorsale penultimum in utroque sexu pliculis sex elevatis ornatum, ultimum maris tridentatum, feminae simplex.

Variat 1, colore aut pallidiore, corpore testaceo, capite abdominisque cingulo nigris, aut saturatiore, corpore nigro, thoracis limbo et abdominis basi obscure ferrugineis, elytris nigris, macula magna humerali testacea.

2, punctura elytrorum subtiliore, obsolescente. Specimina nonnulla, ceterum omnino cum genuinis congruentia, Austriaca misit Dom. Ullrich.

\* 4. *G. affinis*: *Testacea*, *capite abdominisque cingulo piceis*, *thorace marginato*, *seriatim punctato*, *elytris parce punctatis*. — Long. 1 lin.

*Mannerh. Brachelytr.* 74. 3. — *Er. Col. March.* I. 368. 3.

*Aleochar. affinis* Sahlb. *Ins. Fenn.* I. 383. 67.



*Gyrophæna. amabilis* Botsd. et Lacord. Faun. Ent. Paris. I. 535. 1.

*Gyrophæna. nana* Boisd. et Lacord. Faun. Ent. Paris. I. 537. 3.

Habitat in Europae fungis.

Praecedenti simillima. Antennae capitis thoracisque longitudine, articulo secundo elongato, parum incrassato, tertio hoc triplo minore, minuto, quarto obconico, 5—10 aequalibus, subcylindricis, longitudine haud crassioribus, ultimo praecedente sesqui longiore, apice acuminato, pubescentes, parce pilosellae, testaceae, basi dilutiores. Caput depressum, fronte utrinque subimpressa et parce punctata, piceum, ore testaceo. Thorax coleopteris plus dimidio angustior, latitudine dimidio brevior, lateribus ante medium et basi leviter rotundatus, apice truncatus, angulis anterioribus subrectis, leviter deflexis, posterioribus obtusis, utrisque subrotundatis, subdepressus, basi tenuissime marginatus, testaceus, punctis paucis, dorso biseriatis impressus. Elytra thorace paulo longiora, depressa, omnium subtilissime alutacea, parce punctata, aut flava, angulo apicis exteriore fusco nigrove, aut fusco-testacea, humeris luteis. Abdomen rufo-testaceum, segmentis duobus penultimis piceis. Pedes flavi.

Abdominis segmentum dorsale penultimum feminae aequale, maris tuberculo minuto ante marginis posterioris medium munito, ultimum feminae simplex, maris bispinum, spinis subhamatis.

Variat abdomine toto rufo-testaceo, supra ante apicem macula minuta nigricante notato.

\* 5. *G. gentilis*: Testacea, capite nigro, elytris apice, abdominis macula dorsali pectoreque piceis, thorace marginato, biseriatim punctato, elytris dense subtiliter punctatis. — Long.  $1\frac{1}{4}$  lin.

Habitat in Carinthia, Dom. Schmidt.

Antennae debiles, capite sesqui longiores, articulo tertio secundo vix brevior, at multo tenuiore, quarto paulo crassiore, obconico, 5—10 subaequalibus, leviter transversis, ultimo praecedente duplo fere longiore, apice acuminato, pilosellae, fusco-testaceae, basi flavae. Caput thorace paululum angustius, fronte plana, utrinque subimpressa et crebrius fortiusque punctata, nigrum, nitidum, ore testaceo. Thorax coleopteris multo angustior, latitudine dimidio fere brevior, basi modice, lateribus ante medium leviter rotundatus, apice truncatus, angulis anterioribus subrectis, leviter deflexis, posterioribus obtusissimis, utrisque subrotundatis, basi tenuiter marginatus, parum convexus, dorso bilineatum leviter impressus, lineis punctis obsoletioribus at sat crebris notatis, praeterea punctis raris dispersis impressus, testaceus, disco piceo. Scutellum testaceum. Elytra thorace sesqui longiora, depressa, sat crebre subtiliter punctata, tenuiter pubescentia, testacea, apice praecipue extrorsum picea. Abdomen lateribus minus rotundatum, testaceum, dorso segmento antepenultimo nigro-piceo. Pectus piceum. Pedes flavi.

Sexus differentia latet. Abdominis segmentum dorsale penultimum aequale.

Obs. *G. nana* duplo fere maior, latior, elytris subtilius et minus intricatim punctatis, abdominisque segmento penultimo laevissimo, pliculis elevatis nullis, satis distincta. A *G. affini* statura elytrorumque punctura differt.

- \* 6. *G. vinula*: *Testacea, nitida, capite, elytrorum angulis exterioribus abdominisque macula nigris, thorace marginato, subtiliter punctato, elytris parce punctatis.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Pensylvania, Dom. Zimmermann.

*G. nana* paulo maior et magis oblonga depressaque, rufo-testacea, nitida. Antennae capite thoraceque paulo breviores, minus validae, articulo secundo tertioque elongatis, hoc tenuiore et brevior, quarto sub-obconico, 5 — 10 crassitie aequalibus, subcylindricis, sensim paulo brevioribus, ultimo praecedentibus ambobus longitudine prope aequali, subcylindrico, apice acuminato, pubescente, pilosellae, dilute testaceae, basi flavae. Caput nigerrimum, fronte utrinque subseriatim punctata, ore testaceo. Oculi magni. Thorax coleopteris paulo angustior, latitudine duplo brevior, basi modice, lateribus parum rotundatus, apice truncatus, angulis anterioribus subrectis, leviter deflexis, posterioribus obtusis, utrisque subrotundatis, basi tenuiter marginatus, subdepressus, parce subtiliter punctatus, punctis medio irregulariter biseriatis. Elytra thorace sesqui longiora, depressa, subtiliter parce griseo-pubescentia, parce subtiliterque punctata, angulo apicis exteriori nigerrimo. Abdomen supra ante apicem macula magna nigra.

- \* 7. *G. dissimilis*: *Picea, capite nigro, thorace, elytris, abdominis basi, antennis pedibusque testaceis, thorace marginato, parce punctato, elytris aut parce punctatis aut subtiliter granulatis.* — Long. 1 lin.

Habitat in Carolina, Dom. Zimmermann.

*G. nanae* aequalis. Antennae capitis thoracisque longitudine, articulo secundo crassiusculo, tertio hoc paulo brevior et multo tenuiore, quarto tertio duplo brevior et crassior, 5 — 10 subaequalibus, longitudine paulo crassioribus, ultimo praecedentibus ambobus aequali, oblongo-ovato, apice obtuse acuminato, pilosellae, testaceae, basi flavescentes. Caput nigerrimum, nitidum, fronte punctis nonnullis iuxta utrumque oculum impressa, ore testaceo. Oculi magni. Thorax coleopteris paulo angustior, latitudine duplo brevior, basi leviter, lateribus perparum marginatus, apice truncatus, angulis anterioribus subrectis, minus deflexis, posterioribus obtusissimis, vix ullis, parum convexus vel subdepressus, dorso punctis parvis biseriatis, terminalibus in utraque serie binis maioribus, reliquis sat subtilibus, ceterum laevissimus, testaceus, nitidus. Elytra thorace sesqui longiora, subdepressa, aut vage subtilius punctata, aut in altero sexu densius subtiliterque granulata, parce tenuiterque grisea-pubescentia, testacea, apice praecipue extrorsum fusca. Abdomen nigrum, basi supra late, infra anguste testaceum. Pectus nigrum. Pedes testacei.

Sexus differentia praeter elytrorum puncturam mihi non observata. Specimina elytris granulatis mascula esse mihi videntur.

Obs. Caput maiusculum, fronte minus depressa, oculis magnis convexis omnino ut in *G. vinula*.

- \* 8. *G. dispar*: *Testacea, pectore nigro, capite, elytrorum apice abdominisque cingulo piceis, thorace marginato, parce punctato, elytris aut granulatis aut laevigatis.* — Long. 1 lin.

Habitat in Brasilia.

Statura omnino *G. nanae*. Antennae capitis thoracisque prope longitudine, articulo secundo crassiusculo, tertio hoc duplo minore, quarto tertio brevior at paulo crassior, 5—10 subaequalibus, longitudine haud crassioribus, ultimo praecedente duplo longiore, apice acuminato, totae testaceae. Caput nigro-piceum, nitidum, fronte depressa, punctis raris dispersis impressa, ore testaceo. Thorax coleopteris angustior, latitudine duplo latior, basi modice, lateribus ante medium leviter rotundatus, apice truncatus, angulis anterioribus subrectis, leviter deflexis, posterioribus obtusis, utrisque subrotundatis, basi tenuiter marginatus, dorso biseriatim punctatus, punctis serierum parvis, subtilibus, obsoletis, terminali posteriore solo sat conspicuo, ceterum laevissimus, nitidus, testaceus, medio piceus. Elytra thorace plus sesqui longiora, depressa, alterius sexus dense granulata, alterius laevigata, testacea, apice praecipue exterius picea. Abdomen testaceum, pone medium macula supra minore, infra maiore picea. Pectus nigrum. Pedes testacei.

Mas praeter elytra granulata antennis longioribus distinctus.

- \* 9. *G. congrua*: Testacea, capite abdominisque cingulo piceis, thorace fusco-testaceo, dorso biseriatim punctato, elytris subtilissime punctulatis. — Long.  $\frac{3}{4}$ —1 lin.

Er. Col. March. I. 368. 4.

Habitat in Germania.

*G. nana* plerumque paulo minor et brevior. Antennae capite thoraceque paulo longiores, articulo secundo crassiusculo tertioque hoc plus duplo brevior obconicis, quarto fere transverso, sequentibus crassiusculis, 5—10 subaequalibus, crassitie paulo brevioribus, ultimo praecedente duplo longiore, subovato, apice subacuminato, pilosellae, fuscae, basi testaceae. Caput thorace vix angustius, fronte planiuscula, utrinque subtilissime et parce punctata, piceum, parum nitidum, ore testaceo. Thorax coleopteris dimidio angustior, latitudine plus dimidio brevior, basi modice, lateribus leviter rotundatus, apice truncatus, angulis anterioribus rectis posterioribusque obtusis subrotundatis, leviter convexus, basi tenuiter marginatus, omnium subtilissime alutaceus, dorso punctis parvis, biseriatis, obsoletis, terminali posteriore solo maiore, lateribus puncto uno alterove subtili impressus, fusco-testaceus, nitidus. Elytra thorace sesqui longiora, subdepressa, subtilissime alutacea, subtilissimeque punctulata, luteo-testacea, angulo exteriori apicis nigro. Abdomen rufo-testaceum, segmentis duobus penultimis nigricantibus. Pedes testacei.

Sexus differentia latet.

- \* 10. *G. lucidula*: Nigra, nitida, antennarum basi, pedibus elytrisque palide testaceis, his angulo apicali exteriori nigro, thorace dorso biseriatim punctato, elytris laevissimis. — Long.  $\frac{2}{3}$  lin.

Er. Col. March. I. 369. 5.

Habitat Berolini.

Statura praecedentium, at minor, nigra, nitida. Antennae capitis thoracisque longitudine, articulo secundo sat magno, tertio hoc plus duplo minore, praecipue brevior, quartoque tertio brevior et crassior obconicis, sequentibus crassiusculis, 5—10 aequalibus, leviter transversis, ultimo prae-



cedente duplo longiore, acuminato, pilosellae, fuscae, basi testaceae. Caput thorace paulo angustius, fronte depressiuscula, utrinque leviter longitudinaliter subimpressa, at impunctata, laevissimum, nitidum, ore testaceo. Thorax coleopteris dimidio prope angustior, latitudine duplo brevior, lateribus et basi leviter rotundatus, apice subemarginatus, angulis anterioribus subrectis, leviter deflexis posterioribusque obtusis subrotundatis, leviter convexus, basi distincte marginatus, dorso punctis paucis biseriatis obsoletissimis, excepto pari singulo pone medium sat forti impressus, ceterum laevissimus. Elytra thorace plus sesqui longiora, laevigata, testacea, angulo apicis exteriore late nigro. Abdomen totum nigrum. Pedes flavi.

Sexus differentia latet.

- \* 11. *G. flaveola*: Testacea, antennis fuscis, basi pallidis, thorace marginato, convexo, elytris parce punctatis. — Long.  $\frac{3}{4}$  lin.

Habitat in Brasilia.

Statura *G. affinis* et minimis eius individuis aequalis, testacea, nitida, laevis. Antennae capitis thoracisque prope longitudine, articulo secundo crassiusculo, tertio hoc multo minore et angustiore, quarto breviusculo, 5 — 10 aequalibus, longitudine vix crassioribus, ultimo apice acuminato, pilosae, nigro-fuscae, articulis 3 primis flavis. Caput thorace paulo angustius, fronte prope utrumque oculum punctis paucis impressa. Thorax coleopteris paulo angustior, latitudine duplo fere brevior, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus subrectis, leviter deflexis, posterioribus obtusissimis, subrotundatis, leviter convexus, basi tenuiter marginatus, laevissimus, dorso punctis parvis biseriatis notatus, terminalibus distinctioribus, reliquis subtilibus, vix conspicuis. Elytra thorace sesqui prope longiora, punctis subtilibus raris sparsa, interstitiis laevissimis, parce pubescentia, magis lutea. Abdomen laeve, tenuissime pubescens. Pedes lutei.

Mas abdominis segmento dorsali quinto pliculis duabus longitudinalibus notato, sexto apice bispinoso, femina iisdem simplicibus distinguuntur.

- \* 12. *G. splendidula*: Nigra, nitida, antennarum basi pedibusque testaceis, thorace marginato, disco 4-punctato, elytris aut laevibus aut parce granulatis, callo humeralo testaceo. — Long.  $\frac{3}{4}$  —  $1\frac{1}{4}$  lin.

Habitat in Americae merid. insula St. Thomae, Dom. Moritz, in Brasilia, Dom. Prof. Germar.

Habitu *G. politae*, magnitudine instabilis, individuis maximis *G. nanam* prope superantibus, nigra, nitida. Antennae capite thoraceque paulo longiores, articulo secundo haud incrassato, tertio hoc parum tenuiore et paulo longiore, quarto obconico, praecedente paulo brevior et crassiore, 5 — 10 subaequalibus, longitudine haud crassioribus, ultimo paulo longiore, apice acuminato, fuscae, articulis 4 primis flavo-testaceis. Caput thorace multo angustius, fronte plana, aequali, utrinque sparsim parce punctata, ceterum laevissimum, ore magis minusve rufo-piceo. Thorax basi coleopteris parum angustior, antrorsum subangustatus, latitudine duplo fere latior, basi et lateribus fortius rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, posterioribus obtusis, transversim sat fortiter convexus, tenuiter mar-

ginatus, disco punctis 4 sat profundis, quadratim dispositis impressus, inter medium et marginem puncto uno alterove obsolete notatus, ceterum laevissimus. Elytra thorace paulo longiora, laevigata, in altero sexu laevissima, in altero tuberculis vel punctis elevatis raris, in series duas oblique introrsum vergentes digestis obsita, glaberrima, nitida, nigra, callo humerali testaceo. Abdomen tenuiter pubescens, subtus nigrum, basi apiceque piceum, supra obscure testaceum, segmentis intermediis basi piceis. Pedes flavo-testacei.

Mas elytris tuberculatis et abdominis segmento dorsali ultimo medio producto, apice truncato, utrinque spinis duabus porrectis, lobum intermedium prope superantibus armato insignis.

\* 13. *G. corruscula*: *Nigra, nitida, antennis pedibusque testaceis, elytris abdominisque basi piceis, thorace obsolete subtilissime biseriatim punctato, elytris parce obsoleteque punctatis.* — Long.  $\frac{1}{2}$  —  $\frac{2}{3}$  lin.

Habitat in Carolina meridionali, Dom. Zimmermann.

Habitu omnino *G. affinis*, at minor, nigra, nitida. Antennae capite thoraceque paulo longiores, validiusculae, articulo secundo sat magno, primo subaequali, tertio hoc dimidio brevior et multo tenuiore, quarto iterum dimidio brevior, at crassior, obconicis, 5—10 aequalibus, longitudine crassioribus, ultimo praecedente plus dimidio longior, apice acuminato, laete testaceae, basi flavae. Caput thorace paulum angustius, laevigatum, glaberrimum, fronte aequali, utrinque parce punctata, ore testaceo. Thorax coleopteris paulo angustior, latitudine plus duplo brevior, basi et lateribus leviter rotundatus, apice submarginatus, angulis anterioribus subrectis, leviter deflexis, posterioribus obtusis, leviter transversim convexus, basi tenuiter marginatus, dorso subtilissime obsoleteque parce biseriatim punctatus, ceterum laevissimus. Elytra thorace sesqui longiora, subdepressa, parce obsoleteque subtiliter punctata, interstitiis laevissimis, pilis raris brevissimis subtilissimis obsita, picea, nitida, lateribus nigricantia, callo humerali dilutior. Abdomen crassiusculum, laeve, nigrum, basi piceum. Pedes testacei.

Mas abdominis segmento dorsali secundo tuberculo minuto acutiusculo in medio margine posteriore, quinto tuberculis duobus leviter distantibus ante apicem, sexto tuberculis duobus obsoletioribus approximatis in medio munitis distinctus.

\* 14. *G. socia*: *Nigra, nitida, antennis pedibusque testaceis, elytris abdominisque basi piceis, thorace utrinque vage subtiliter punctato, elytris crebrius punctatis.* — Long.  $\frac{1}{2}$  lin.

Habitat in Carolina meridionali, Dom. Zimmermann.

Praecedenti affinis et minoribus eius speciminibus aequalis, nigra, nitida. Antennae capite thoraceque fere longiores, articulo secundo crassiusculo, primo paulo brevior, tertio hoc duplo minore, praecipue tenuiore, quartoque hoc brevior et vix crassior obconicis, 5—10 aequalibus, crassis, longitudine duplo fere crassioribus, ultimo crassitie laud longior, obtuse acuminato, testaceae, basi flavae. Caput thorace vix angustius, fronte aequali, parce disperse punctata, ore testaceo. Thorax coleopteris parum angustior, latitudine plus duplo brevior, basi modice, lateribus leviter ro-

tundatus, apice truncatus, angulis anterioribus rectis, parum deflexis, posterioribus obtusis, parum convexus, basi minus subtiliter marginatus, lateribus parce vageque subtiliter punctatus, medio longitudinaliter laevis pilis longioribus albidis parce adspersus. Elytra thorace plus sesqui longiora, subdepressa, crebrius subtiliusque punctata, parcius longius albedo-pubescentia, picea, humeris dilutioribus, subtestaceis. Abdomen supra subglabrum, subtus parcius longiusque albedo-pubescentia, nigrum, basi apiceque piceum. Pedes flavo-testacei.

Obs. Cum praecedente occurrere, nihilominus specie distincta videtur, thoracis elytrorumque punctura pubescentiaque differentibus, neque pro altero eius sexu haberi potest, cum uterque illius sexus, mas abdomine tuberculato, femina abdomine simplice, obvius sit. Omnia, quaecunque vidi, huius speciei individua abdominis segmento apicali apice bifido instructa sunt, omnia forte mascula.

\* 15. *G. manca*: *Nigra, nitida, antennis pedibusque flavo-testaceis, thorace crebrius, elytris crebre punctatis.* — Long.  $\frac{2}{3}$  lin.

Habitat in Germania: Berolini, ipse.

Statura omnino praecedentis et parum maior, nigra, nitida, elytris concoloribus. Antennae capitis thoracisque longitudine, articulo secundo crassiusculo, obconico, tertio hoc dimidio brevior et praecipue tenuiore, quarto hoc brevior et paulo crassior, 5—10 crassis, inter se aequalibus, longitudine dimidio crassioribus, ultimo praecedente duplo prope crassior, acuminato, flavo-testaceae. Caput thorace angustius, fronte aequali, utrinque punctis paucis obsoletioribus impressa, ore testaceo. Thorax coleopteris paulo angustior, latitudine duplo brevior, lateribus et basi leviter rotundatus, apice truncatus, angulis anterioribus subrectis, deflexis, posterioribus obtusis, leviter transversim convexus, basi subtiliter marginatus, crebrius, lateribus disperse, dorso subseriatim subtilius punctatus, interstitiis laevissimis, glaber. Elytra thorace sesqui longiora, sat crebre minus subtiliter punctata, interstitiis laevissimis, tenuiter albedo-pubescentia. Abdomen subtilissime punctulatum, tenuissime subtilissimeque albedo-pubescentia, ano lutescente. Pedes flavo-testacei.

Sexus differentia latet.

Obs. Olim cum *G. polita* mihi confusa, a qua thoracis elytrorumque sculptura, nec non corpore latiusculo differt. Propius accedit ad praecedentem, a qua thorace elytrisque crebrius punctatis, thorace basi subtilius marginato, et elytris concoloribus distinguenda videtur.

\* 16. *G. polita*: *Oblonga, picea, antennis pedibusque pallidis, thorace disco obsolete biseriatim punctato, elytris parce obsoletius punctatis, interstitiis subtilissime alutaceis.* — Long.  $\frac{2}{3}$  lin.

*Mannerh. Brachelytr.* 74. 4. — *Er. Col. March.* I. 369. 6.

*Al. polita Grav. Micr.* 99. 48. *Mon.* 176. 73. — *Latr. Hist. nat. Crust. et Ins.* IX. 391. 47. — *Gyll. Ins. Suec.* II. 415. 37. — *Sahlb. Ins. Fenn.* I. 483. 68.

Habitat in Europa.

Nigro-picea, nitidula. Antennae capite sesqui longiores, articulo secundo crassiusculo, obconico, tertio hoc parum brevior, at tenuiore, quarto minuto, transverso, 5—10 crassiusculis, transversis, ultimo praecedente vix duplo longiore, crassiusculo, subgloboso, apice acuminato, testaceae.



Caput minutum, fronte aequali, utrinque parce obsoleteque punctata, ore testaceo. Thorax coleopteris dimidio angustior, latitudine dimidio brevior, antrorsum subangustatus, basi leviter rotundatus, lateribus subrectis, apice truncatus, angulis anterioribus rectis, fortiter deflexis, posterioribus obtusis, parum convexus, basi subtiliter marginatus, seriebus duabus dorsalibus punctorum subtilium obsoletorumque. Elytra thorace sesqui longiora, subtilissime alutacea, vage obsolete punctata, picea. Abdomen latiusculum, apice testaceum. Pedes testacei.

- \* 17. *G. strictula*: *Oblonga, nigra, nitidula, abdomine piceo, ante apicem nigro, antennis pedibusque testaceis, thorace laevi, elytris subtilissime alutaceis, obsoletissime punctulatis.* — Long.  $\frac{2}{3}$  lin.

Habitat in Hungaria, Dom. Frivaldsky, in Gallia, Mus. Dom. Aubé.

Statura omnino praecedentis, nigra, nitidula. Antennae capite thoraceque breviores, apicem versus sensim subincrassatae, articulo secundo crassiusculo, obconico, tertio hoc dimidio brevior et multo tenuiore, quarto minuto, transverso, 5—10 crassiusculis, transversis, sensim paulo crassioribus, ultimo maiusculo, breviter ovato, apice subacuminato, laete flavo-testaceae. Caput thorace paulo angustius, fronte aequali, planiuscula, sparsim obsolete punctata. Thorax coleopteris plus dimidio angustior, latitudine dimidio brevior, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus subrectis, leviter deflexis, posterioribus obtusissimis, leviter convexus, basi minus subtiliter marginatus, laevis vel obsolete biserialim impressus, pone medium punctorum maiorum pari singulo notatus. Elytra thorace plus sesqui longiora, omnium subtilissime alutacea, parce subtiliter et valde obsolete punctata, nigro-picea. Abdomen nigrum, apice, interdum etiam basi piceum. Pedes flavi.

Abdominis segmentum dorsale penultimum in altero sexu pliculis brevissimis longitudinalibus instructum, in altero laeve.

Obs. Differt a praecedente statura paulo maiore, antennis apicem versus sensim crassioribus, elytris subtiliter punctatis, thorace apicem versus nullo modo angustato, basi minus subtiliter marginato.

- \* 18. *G. Boleti*: *Elongata, nigro-fusca, nitida, subtilissime punctulata, ore, antennis pedibusque flavis, elytris fusco-testaceis, thorace marginato, subtilissime crebre punctato.* — Long.  $\frac{1}{2}$  lin.

*Staphyl. Boleti* Linn. Faun. Suec. n. 859. Syst. Nat. I. II. 686. 26. — De Geer Ins. IV. 28. 13. t. 1. f. 15—17. — Müll. Prodr. 97. 1093. — Goeze Ent. Beitr. I. 722. 26. — De Vill. Ent. I. 419. 26.

*Aleochar. Boleti* Gyll. Ins. Suec. II. 416. 38. — Sahlb. Ins. Fenn. I. 384. 69.

*Bolitochara Boleti* Mannerh. Brachelytr. 82. 46.

Habitat in Suecia, Germania.

Minima, picea, subtiliter pubescens. Antennae captis thoracisque fere longitudine, crassiusculae, extrorsum sensim crassiores, articulo secundo crassiusculo tertioque hoc duplo minore obconicis, quarto minuto, transverso, 5—10 transversis, crassis, sensim paulo crassioribus, ultimo subgloboso, flavae. Caput laeve, nigrum, nitidum, ore testaceo. Thorax coleopteris vix angustior, latitudine plus dimidio brevior, basi et lateribus leviter

rotundatus, apice truncatus, angulis anterioribus rectis, leviter deflexis, posterioribus obtusis, leviter transversim convexus, lateribus et basi rotundatus, basi anguste marginatus, subtiliter crebre punctulatus, tenuiter albido-pubescent. Elytra thorace sesqui longiora, crebre subtiliter punctata, tenuiter albido-pubescentia. Abdomen nigro-piceum, subtilissime punctulatum, tenuiter subtiliterque albido-pubescent, ano obscure testaceo. Corpus subtus pallidius. Pedes flavi.

Variat corpore fusco-testaceo, capite abdominisque medio nigro-fuscis, elytris pallide testaceis.

\* 19. *B. minima*: *Rufo-testacea, nitida, thorace medio obsolete punctato, elytris parce punctatis.* — Long.  $\frac{1}{2}$  lin.

*Er. Col. March. 1. 370. 7.*

Habitat Berolini, Dom. Kirstein.

Praecedenti affinis. Corpus dilute ferrugineum vel rufo-testaceum, nitidum. Antennae capitis thoracisque longitudine, articulo secundo crassiusculo tertioque hoc multo minore, praecipue angustiore obconicis, quarto minuto, breviusculo, tertio autem paulo crassiore, sequentibus crassis, 5—10 aequalibus, transversis, ultimo praecedente duplo fere longiore, subacuminato. Caput thorace vix angustius, laeve, vertice piceo. Thorax coleopteris angustior, latitudine duplo brevior, lateribus et basi leviter rotundatus, apice truncatus, angulis anterioribus rectis, posterioribus obtusis, subdepressus, basi tenuiter marginatus, nitidus, laevis, medio longitudinaliter vage subtiliterque punctatus. Elytra thorace sesqui longiora, subdepressa, parce punctata, subtilissime pubescentia, angulo apicis exteriori piceo. Abdomen ante apicem interdum piceum.

## Pelusia Er.

*Maxillae mala interiore mutica, intus spinulis ciliata.*

*Ligula brevis, integra; paraglossae nullae.*

*Palpi labiales 2-articulati, articulis subaequalibus, secundo sublunato.*

*Tarsi antici 4-, posteriores 5-articulati, postici articulo primo elongato.*

Corpus oblongum, subtus leviter convexum, alatum. Caput exsertum, deflexum, basi haud constrictum, oculis mediocribus, modice prominulis, ore producto, clypei spatio membranaceo. Labrum maiusculum, lateribus leviter rotundatum, apice truncatum, mandibulas obtegens. Mandibulae muticae. Maxillae malis corneis, exteriori basi apiceque membranacea, portione membranacea apicali malam interiori superante, hac interne membranacea, apice oblique introrsum submarginata, sinu spinulis confertis ciliato. Palpi maxillares modice elongati, articulo secundo tertioque longitudine subaequalibus, hoc parum incrassato, quarto parvo, subulato. Labium (t. III. f. 12.) mento apice profunde emarginato, angulis anterioribus spiniformibus, porrectis, ligula brevi, rotundata, integra, paraglossis nullis. Palpi labiales biarticulati, articulis subaequalibus, primo cylindrico, secundo sublunato. Antennae breves crassiusculaeque, subclavatae, articulo primo crassiusculo, secundo tertioque clavatis, elongatis, sequentibus sensim paulo crassioribus, ultimo apice abrupte acuminato. Thorax coleopterorum prope latitudine, transversus,

basi rotundatus, leviter convexus. Scutellum minutum, triangulare. Elytra truncata, ad angulum anteriorem subsinuata. Abdomen parallelum, supra planum, subtus convexum, reclinandum. Pedes breviores, intermedii basi approximati, tarsi anticis 4-, posterioribus 5-articulatis, posticis articulo primo leviter elongato.

#### Observationes.

1. Maxillae omnino ut in *Homalota* (t. III. f. 22.), ligula omnino ut in *Gyrophæna*, a qua ceterum mento longiore, antrosum angustato, profunde emarginato, angulis anterioribus spiniformibus et palporum labialium articulo secundo differt.

2. Habitus intermedius quasi inter *Myrmedoniam* et *Aleocharam*, ab illa coxis intermediis approximatis, ab hac tarsis anticis 4-articulatis discedens. Caput deflexum, fronte planiuscula, et antennae breves subclavatae affinitatem cum *Gyrophæna* produnt, a qua autem praeter notas supra allatas coxis intermediis basi approximatis facile distinguitur. Corpus parce pilosellum, ut in *Aleochara*, neque pube sericea obductum.

3. Sexus differentia antennarum structura et abdominis segmentis dorsalibus duobus in mare tuberculatis in specie unica huius generis constituitur.

4. De victu nihil compertum.

\* 1. *P. labiata*: *Nigro-picea, ore albido, antennis pedibusque rufis.* — Long 2 lin.

T. 1. f. 1.

Habitat in Madagascar, Dom. Goudot.

*Nigro-picea*, subnitida, subtiliter pubescens. Antennae capite parum longiores, maris articulis tribus primis sequentibus coniunctis prope longioribus, primo secundo tertioque coniunctis vix brevioribus, crassiusculo, his fortiter clavatis, tertio paululum longiore, 4 — 10 sensim paulo brevioribus et crassioribus, penultimis brevissimis, perfoliatis, ultimo praecedentibus ambo paulo longiore, apice abrupte acuminato, feminae articulis 3 primis sequentibus coniunctis paulo brevioribus, primo secundo sesqui longiore, crassiusculo, tertio hoc paulo longiore, at multo tenuiore, elongato-obconico, sequentibus apicem versus sensim leviter incrassatis, 4 — 10 sensim paululum crassioribus, non autem brevioribus, ultimo praecedentibus ambobus paulo longiore, apice abrupte magis quam in mare acuminato, rufo-testaceae vel rufae. Caput fronte plana, vage punctata, clypeo labioque testaceis, palpis rufis. Thorax coleopteris vix angustior, latitudine plus dimidio brevior, basi modice, lateribus fortius rotundatus, apice truncatus, angulis anterioribus subrectis, leviter deflexis, posterioribus obtusis, omnibus subrotundatis, leviter convexus, crebrius minus subtiliter punctatus. Elytra thorace paulo longiora, apice ad angulum anteriorem subsinuata, crebrius fortiusque punctata. Abdomen supra subconvacum, parcius et minus subtiliter punctatum, parce pubescens, concolor, subtus convexum, minus dense punctatum, longius pubescens, segmentis 4 primis margine apicali piceo, quinto sextoque totis rufo-testaceis. Caput et thorax subtus rufo-testacei. Pedes rufo-testacei.

Mas abdominis segmento dorsali secundo medio, quinto ante apicem tuberculo minuto munitis, femina segmentis iisdem muticis.



**Placusa Er.***Er. Col. March. I. p. 370.**Maxillae mala interiore mutica, intus spinulis ciliata.**Ligula brevis, integra; paraglossae breves.**Palpi labiales breves, 2-articulati, articulo primo crasso, secundo tenui.**Tarsi antici 4-, posteriores 5-articulati, postici articulo primo subelongato.*

Corpus oblongum, depressum, alatum. Caput thorace minus, exsertum, porrectum, basi haud constrictum, os versus triangulariter attenuatum, oculis mediocribus, modice prominulis. Labrum transversum, truncatum, mandibularum apicem haud obtegens. Mandibulae muticae. Maxillae mala exterioriore anteriorem paululum superante, cornea, basi apiceque membranacea, portione membranacea apicali subtiliter pubescente, mala interiori membranacea, margine exteriori corneo, apice oblique introrsum subemarginata, sinu spinulis confertis ciliato. Palpi maxillares minus elongati, articulo secundo tertioque longitudine subaequalibus, hoc modice incrassato, quarto minuto, subulato. Labium (t. III. f. 13.) mento transversum, apice leviter emarginato, ligula latiuscula, integra, paraglossis distinctis, at minutis, ligula multo brevioribus, intra marginem anteriorem subtilissime ciliatis. Palpi labiales biarticulati, articulo primo crasso, secundo hoc parum brevior, at multo tenuiore. Antennae apicem versus sensum leviter incrassatae, articulis secundo tertioque obconicis, hoc praecedente paulo minore, articulo ultimo rotundato. Thorax transversus, plerumque basi utrinque sinuatus. Scutellum triangulare. Elytra apice truncata. Abdomen parallelum, vel apicem versus leviter angustatum. Pedes breves, intermedii basi approximati, tarsi antici 4-, posteriores 5-articulati, postici articulo primo leviter elongato.

**Observationes.**

1. Maxillae omnino ut in *Homalota* (t. III. f. 22.). Labium distinctum ligula suborbiculari et paraglossis distinctis. Paraglossae olim me fugerant.

2. *A. Gyrophaena* coxis intermediis approximatis et reliquis generibus, quibuscum ligula integra et palpis labialibus biarticulatis convenit, corpore valde depresso discedit. Habitu intermedia quasi inter *Homalota* et *Oxypoda*, hac autem magis depressa, et tarsis anticis 4-articulatis, ab illa tarsorum posticorum articulo primo leviter elongato distincta.

3. Victus, quantum constat, in fungis.

4. Mas abdominis segmento dorsali sexto utrinque spina armato, apice tuberculato aut denticulato, femina segmento eodem simplice distinguuntur.

\* 1. *P. complanata*: *Depressa, nigra, opaca, antennarum basi pedibusque testaceis, elytris fuscis, thorace coleopterorum latitudine, depresso, basi bisinuato, angulis posterioribus obtusis.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Germania, Dom. Schüppel.

Corpus latiusculum, complanatum, nigrum, opacum, pube brevissima fusca densius vestitum. Antennae capite thoraceque paulo breviores, tenuiores, apicem versus vix incrassatae, articulis tribus primis longioribus, secundo tertioque aequalibus, primo paulo brevioribus, 4 — 10 aequalibus, crassitie paulo brevioribus, ultimo breviter ovato, apice acutiusculo, fuscae, basi testaceae. Palpi testacei. Caput thorace duplo prope angustius, subtriangulare, confertissime subtiliter punctatum, fronte depressa, ore rufo-

piceo. Thorax coleopterorum latitudine, latitudine duplo brevior, lateribus rotundatus, antrorsum subangustatus, apice submarginatus, angulis anterioribus obtusis, leviter deflexis, basi medio late sat fortiter rotundatus, utrinque sinuatus, angulis posterioribus acutiusculis, retrorsum prominulis, fortiter depressus, confertissime subtiliter punctatus. Elytra thorace sesqui longiora, apice ad angulum anteriorem leviter sinuata, depressa, omnium confertissime subtiliter punctata, testaceo-fusca, ad suturam et angulum apicis anteriorem nigricantia. Abdomen depressum, apicem versus sensim subangustatum, supra segmentis tribus primis confertim subtiliter, quarto quintoque parce subtilissime punctatis, parce tenuiterque pubescens, nitidulum, apice summo testaceo, subtus subtilius parciusque aequaliter punctatum, nigrum, nitidum, segmentis singulis piceo-marginatis. Pedes testacei, femoribus fusciscentibus.

Mas abdominis segmento dorsali sexto apice tuberculo prominulo terminato, utrinque spina longiore tenui, introrsum vergente, armato cognoscitur.

\* 2. *P. pumilio*: Depressa, nigra, opaca, antennarum basi pedibusque testaceis, elytris fuscis, thorace coleopterorum latitudine, subdepresso, basi bisinuato, angulis posterioribus subrectis. — Long. 1 lin.

*Aleo ch. pumilio* Grav. Micr. 98. 46. Mon. 175. 70. — Latr. Hist. nat. Crust. et Ins. IX. 390. 46. — Gyll. Ins. Suec. II. 411. 33. — Sahlb. Ins. Fenn. I. 382. 64. — Zetterst. Faun. Lapp. I. 102. 18. Ins. Lapp. 78. 26.

*Bolitochara pumilio* Mannerh. Brachelytr. 82. 44. — Boisd. et Lacord. Faun. Ent. Paris. I. 552. 20.

Habitat in Germania, Suecia, Gallia.

Praecedente paulo minor, nigra, opaca, pube brevissima grisea densius vestita. Antennae capitis thoracisque prope longitudine, apicem versus subincrassatae, articulo secundo crassiusculo, tertio hoc paulo minore, obconico, 4 - 10 transversis, crassitie duplo brevioribus, ultimo maiusculo, subgloboso, apice obtusiusculo, pilosellae, fusco-nigrae, basi fusco-testaceae. Palpi fusco-testacei. Caput thorace dimidio angustius, triangulare, basi rotundatum, confertissime subtiliter punctatum, opacum. Thorax coleopterorum latitudine, latitudine duplo brevior, antrorsum vix angustatus, lateribus rotundatus, apice submarginatus, angulis anterioribus obtusis, subrotundatis, leviter deflexis, basi medio late rotundatus, utrinque sinuatus, angulis posterioribus rectis, minutis, prominulis, subdepressus seu perparum convexus, confertissime subtiliter punctatus. Elytra thorace sesqui longiora, apice truncata, depressa, confertissime subtiliter punctata, testaceo-fusca seu fusco-testacea. Abdomen depressum, apicem versus sensim subangustatum, supra segmentis tribus primis confertim, quarto quintoque parcius subtiliter, subtus aequaliter parcius subtiliter punctatum, densius pubescens, subnitidum, nigrum, ano testaceo. Pedes testacei.

Mas abdominis segmento dorsali sexto tuberculo prominulo terminato, utrinque spina longiore tenui armato distinctus.

Obs. Affinis praecedenti, satis distincta autem antennarum articulis 4 - 10 fortius transversis, thorace minus depresso, antrorsum vix angustato, basi medio minus fortiter rotundato, utrinque minus fortiter sinuato, angulis posterioribus minoribus, minus acutis, elytris apice externe non sinuatis, statura minore, non adeo plana.

- \* 3. *P. humilis*: *Depressa, nigra, opaca, antennarum basi pedibusque testaceis, thorace coleopterorum prope longitudine, subdepresso, basi rotundato, angulis posterioribus obtusis.* — Long. 1 lin.

*Placusa pumilio* Er. Col. March. I. 371. 1.

Habitat in Germania.

Statura depressa *P. complanatae*, at paulo minor et angustior, nigra, opaca, pube brevissima fusca densius vestita. Antennae capite thoraceque paulo breviores, tenuiores, articulo secundo crassiusculo tertioque obconico longitudine aequalibus, 4 — 10 subaequalibus, crassitie paulo brevioribus, ultimo ovato, apice obtusiusculo, fusco-nigrae, basi testaceae. Caput thorace duplo fere angustius, triangulare, confertissime subtiliter punctatum, fronte plana, ore rufo-testaceo. Thorax basi coleopteris haud angustior, latitudine duplo brevior, antrorsum subangustatus, apice truncatus, angulis anterioribus rotundatis, leviter deflexis, basi rotundatus, ad angulos posteriores vix sinuatus, his obtusis, fortiter depressus, confertissime subtilissimeque punctatus. Elytra thorace sesqui longiora, apice truncata, confertissime subtilissimeque punctata, fusco-nigra. Abdomen depressum, apicem versus sensim subangustatum, confertim subtiliter punctatum, supra parcius pubescens, nitidulum, subtile opacum, nigrum, segmentis anterioribus late piceo-marginatis, ano testaceo. Pedes testacei, femoribus saepe fusciscentibus.

Mas abdominis segmento dorsali sexto tuberculo prominulo terminato, utrinque spina longiore tenui, introrsum vergente, armato distinctus.

Variat thorace elytrisque fuscis, abdomine rufo-piceo, ante apicem nigro.

Obs. *P. complanatae* magis quam praecedenti affinis corpore omnino complanato et antennis tenuioribus, ab utraque autem thoracis forma satis distincta. Hae tres species a sequentibus abdominis in masculis structura discedunt, inter se autem conveniunt.

- \* 4. *P. infima*: *Nigra, antennarum basi pedibusque testaceis, elytris fuscis, thorace coleopterorum prope longitudine, transversim convexo, basi leviter bisinuato, angulis posterioribus rectis.* — Long. 1 — 1½ lin.

Habitat in variis Germaniae regionibus; in montibus Uralensibus, Dom. Prof. Eversmann.

Praecedente paulo minus depressa, nigra, subnitida, pube subtili brevique grisea tenuius vestita. Antennae capitis thoracisque prope longitudine, graciliores, apicem versus vix incrassatae, articulo secundo tertioque subaequalibus, quarto sequentibus minore, 5 — 10 transversis, subaequalibus, ultimo breviter ovato, apice obtusiusculo, parce pilosellae, fuscae, basi testaceae. Palpi testacei. Caput thorace duplo fere angustius, densius subtiliter punctatum, fronte minus depressa. Thorax basi coleopteris vix angustior, latitudine baseos duplo prope brevior, antrorsum leviter angustatus, lateribus rotundatus, apice truncatus, angulis anterioribus obtusis, subrotundatis, leviter deflexis, basi medio late rotundatus, prope angulos posteriores leviter sinuatus, his rectis, at parum prominulis, leviter transversim convexus, confertim subtiliter punctatus. Elytra thorace sesqui longiora, apice truncata, depressa, confertim subtiliter punctata, fusca, seu fusco-testacea. Abdomen apice sensim attenuatum, confertissime subtilissimeque punctatum,



supra brevissime, subtus longius pubescens, nitidulum, nigrum, apice summo testaceum. Pedes testacei.

Mas abdominis segmento dorsali sexto supra granulis nonnullis instructo, margine apicali subtiliter denticulato, utrinque spina laterali brevior armato distinguendus.

Variat aut elytris nigricantibus, aut dilutior, thorace fusco, elytris fusco-testaceis, abdomine fusco, ante apicem nigro.

Obs. Habitus huius speciei, minus quam praecedentes depressae, intermedius quasi inter Homalotam et Oxypodam.

\* 5. *P. despecta*: Nigra, pedibus testaceis, elytris fuscis, thorace coleopterorum latitudine, transversim convexo, angulis posterioribus obtusis. — Long.  $\frac{3}{4}$  lin.

Habitat in Carolina meridionali, Dom. Zimmermann.

Habitu fere Homalotae aterrimae, at paulo magis depressa, nigra, parum nitida, pube subtili grisea vestita. Antennae capite thoraceque paulo breviores, apicem versus sensim subincrassatae, articulo secundo crassiusculo, tertio hoc paulo brevior et duplo fere tenuiore, obconico, 4 — 10 sensim paulo crassioribus, crassitie paulo brevioribus, ultimo maiusculo, praecedentibus ambobus maiore, ovato, apice obtuso, pilosellae, nigrae, basi piceae. Caput thorace multo angustius, confertim subtilissime punctatum, fronte minus depressa, ore piceo. Thorax basi coleopterorum latitudine, latitudine dimidio brevior, antrorsum leviter angustatus, lateribus rotundatus, apice truncatus, angulis anterioribus obtusis, subrotundatis, deflexis, basi late modice rotundatus, ad angulos posteriores subsinuatus, his obtusis, haud prominulis, leviter transversim convexus, confertim subtilissime punctatus. Elytra thorace paulo longiora, apice ad angulum anteriorem subsinuata, depressa, confertissime subtiliter punctata, fusca seu fusco-testacea, circa scutellum infuscata. Abdomen apicem versus sensim angustatum, minus confertim subtiliter punctatum, piceum, medio nigrum. Pedes testacei.

Mas abdominis segmento dorsali sexto supra granulis nonnullis notato, margine apicali acute denticulato, utrinque spina laterali brevior armato distinctus.

Variat thorace rufo-piceo, elytris testaceis, abdomine rufo-piceo, ante apicem nigro.

Obs. Cum praecedente thorace transversim convexo convenit, differt autem statura minore brevioraque, thorace minus brevi, angulis posterioribus obtusis, abdomine minus elongato. Abdominis segmentum sextum maris omnino ut in illa.

\* 6. *P. adscita*: Linearis, subdepressa, nigra, antennarum basi pedibusque testaceis, elytris fuscis, thorace coleopteris angustiore, subdepresso. — Long. 1 lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura elongata P. humilis, at paulo minor et magis depressa, nigra, subnitida, tenuiter subtilissime griseo-pubescens. Antennae capite thoraceque paulo breviores, apicem versus fortius incrassatae, articulo secundo crassiusculo tertioque paulo tenuiore longitudine aequalibus, obconicis, quarto minuto, prope transversum, 5 — 10 aequalibus, crassiusculis, transversis, cras-

sitie duplo brevioribus, ultimo breviter ovato vel subgloboso, apice obtusiusculo, pilosellae, pubescentes, fuscae, basi fusco-testaceae. Palpi testacei. Caput thorace paulo angustius, basi leviter constrictum, confertissime subtiliter punctatum, fronte parum convexa, antice transversim depressa. Thorax coleopteris dimidio prope angustior, latitudine dimidio brevior, antrosum non angustatus, basi et lateribus rotundatus, apice truncatus, angulis omnibus obtusis, subrotundatis, anterioribus leviter deflexis, parum convexus seu subdepressus, confertim subtiliter punctatus. Elytra thorace sesqui longiora, apice truncata, depressa, crebre minus subtiliter punctata, fusca. Abdomen lineare, depressum, confertim subtiliter punctatum, nitidulum. Pedes testacei.

Mas. abdominis segmento dorsali sexto apice spina media acuta, basi bidenticulata, utrinque spina tenui laterali, intermediae longitudine aequali, introrsum curvata, armato insignis.

Obs. Corporis habitus et thoracis forma omnino Homalotae, at certe huius generis, etiam secundum oris structuram.

Species mihi invisae, forte huius generis.

7. *P. (?) atrata*: Brevis, lata, depressa, nigra, subnitida, antennis longitudine thoracis pedibusque fusciscentibus, tibiis tarsisque dilutioribus, thorace brevissimo, ante scutellum obsolete foveolato. — Long.  $\frac{2}{3}$  lin.

*Bolitochara atrata* Mannerh. *Brachelytr.* 82. 45.

Var. b: *Ore elytrisque fusco-testaceis, pedibus totis pallidioribus, thorace haud foveolato.*

*Aleochara atrata* Sahlb. *Ins. Fenn.* I. 375. 54. — *Zetterst. Ins. Lappon.* 77. 17.

Habitat in Fennia australi, rarius. Var. b in Lapponia legit F. Sahlberg.

*P. pumihoni* statura satis similis, sed differt thorace depresso, postice vix sinuato. Caput thorace multo angustius, rotundatum, nigrum, nitidum, subtilissime punctulatum, ore palpisque fusco-piceis. Antennae thoracis longitudine, fusciscentes, apice parum incrassatae, articulo ultimo ovato, obtuso. Thorax transversus, longitudine duplo latior, antice truncatus, angulis depressis, lateribus et postice rotundatus, supra depressus, ater, opacus, densius pubescens, subtilissime punctulatus, foveola ante scutellum obsolete impressa. Scutellum triangulare, nigrum, subtilissime punctulatum. Elytra thorace parum latiora et dimidio longiora, apice truncata, supra depressa, subtiliter punctulata, pubescentia, nigra, subopaca. Abdomen pone elytra reliquo corpore brevius, latum, depressum, nigrum, subnitidum, vix punctulatum, pubescens. Corpus subtus nigrum, nitidum, subtiliter punctulatum, pubescens. Pedes mediocres, femoribus fuscis, tibiis tarsisque pallidioribus.

Var. b colore tantum pallidiore et thorace paulo densius pubescente atque haud foveolato differt. Vix diversa species. — Specimen var. b et colore et statura simillimum, sed duplo maius, e Lapponia acceptum posideo, quod huc pertinere credo. (Sahlberg.)

Obs. De genere vix mihi dubium est. Statura plana et thoracis angulis posterioribus haud prominentibus affinis forte *P. humili*, differre autem videtur statura minore, colore nitidiore, magis nigro et thorace basi foveolato. Var. b forte a *P. humili* non distincta, praesertim specimen allatum maius

**Euryusa Er.***Er. Col. March. I. p. 371**Maxillae mala interiore mutica, intus spinulis ciliata.**Ligula elongata, integra; paraglossae nullae.**Palpi labiales 2-articulati, articulo primo paulo longiore.**Tarsi antici 4-, posteriores 5-articulati, postici articulo primo elongato.*

Corpus oblongum, latiusculum, supra depressum, subtus leviter convexum, alatum. Caput nāiusculum, deflexum, basi rotundatum, apice leviter angustatum, oculis parvis, rotundatis, leviter prominulis. Labrum transversum, apice subrotundatum, mandibularum apicem haud obtegens. Mandibulae muticae. Maxillae (t. III. f. 23.) mala exteriore cornea, basi apiceque membranacea, portione membranacea apicali pubescente, malam interiorem parum superante, hac cornea, margine exteriore incrassato, interiore membranacea, apice introrsum oblique subemarginato, sinu spinulis confertis ciliato. Palpi maxillares modice elongati, articulo secundo tertio paulo brevior, hoc incrassato, quarto parvo, subulato. Labium (t. III. f. 14.) mento transverso, apice truncato, ligula angusta, lineari, palporum labialium articulo primo fere aequali, integra, paraglossis nullis. Palpi labiales biarticulati, articulis crassitie subaequalibus, primo paulo longiore. Antennae breviusculae crassiusculaeque, apicem versus leviter incrassatae, articulis tribus primis leviter elongatis, secundo tertioque leviter obconicis, articulis 6 — 10 transversis, ultimo maiore, subacuminato. Thorax coleoptera latitudine excedens, antrorsum parum angustatus, basi bisinuatus, apice truncatus. Scutellum transversim triangulare. Elytra apice truncata, ad angulum exteriorem emarginata. Abdomen apicem versus sensim leviter angustatum, supra planum, subtus convexiusculum. Pedes breviusculi, intermedii basi approximati, tarsi antici 4-, posteriores 5-articulati, postici articulo primo modice elongato.

**Observationes.**

1. Habitus corporis omnino Dinardae, ut nisi antennis paulo longioribus, apicem versus sensim incrassatis vix distinguatur, at tanto magis oris partibus ab illa discedit. Sunt enim huic generi maxillae mala interiore ut in Homalota (t. III. f. 22.) illi apice unco terminata, huic mentum apice truncatum, illi profundius emarginatum, ligula huic integra, illi bifida, paraglossae huic nullae, illi distinctae, palpi labiales huic 2-, illi 3-articulati, palporum maxillarium articulus ultimus huic tenuissimus, subulatus, illi conicus, acuminatus. Alii cuidam generi Euryusa neque habitu, neque characteribus accedit. Ligula elongata et lineari reliquis generibus, quibus palpi labiales 2-articulati sunt, discrepat.

2. Palpos labiales olim triarticulatos descripsi, nuper autem, oris examine reiterato, duos tantummodo articulos observo, quorum primum maiorem solidum nec duobus compositum, ut olim putavi, cognosco.

3. Sexus differentiam non observavi.

4. Victus in formicetis.

\* 1. *E. sinuata*: Testacea, subtiliter punctulata, thorace lato, brevi, basi leviter bisinuato. — Long.  $1\frac{1}{4}$  lin.

Tab. I. f. 2.

*Er. Col. March. I. 372. 1.*

Habitat in Germania: Berolini, Weber, in Austria Dom. Ullrich; in Gallia, Dom. Aubé.



Corpus confertim subtiliter punctulatum, pube tenuissima depressa flavescenti vestitum, fusco-testaceum, abdomine ante apicem nigro. Antennae thorace sesqui longiores, apicem versus sensim leviter incrassatae, articulo tertio secundo paulo longiore, quarto tertio paulo crassiore at dimidio brevior, 5 — 10 longitudine aequalibus, transversis, sensim paulo crassioribus, subcylindricis, ultimo maiusculo, praecedentibus tribus coniunctis longitudine prope aequali et paulo crassiore, subovato, apice subacuminato, rufae, articulo ultimo ferrugineo. Caput fronte leviter convexa. Thorax coleopteris paulo latior, latitudine duplo brevior, antrorsum leviter angustatus, lateribus et angulis anterioribus rotundatus, apice leviter emarginatus, basi leviter bismuatus, angulis posterioribus retrorsum leviter prominulis, rotundatis, supra parum convexus, margine laterali late planato. Elytra thorace paulo longiora, subdepressa. Abdomen longius pubescens et praecipue versus apicem parcius profundiusque punctatum.

### **Dinarda Leach.**

*Mannerh. Brachelytr. p. 65. — Er. Col. March. I. p. 373.*

*Lomechusa Curtis Britt. Ent. IX. t. 410.*

*Maxillae mala interiore unco simplice terminata.*

*Ligula angusta, bifida; paraglossae latae, breves, rotundatae.*

*Palpi maxillares 3-articulati, articulo secundo paulo brevior.*

*Tarsi antici 4-, posteriores 5-articulati, postici articulo primo vix elongato.*

Corpus latiusculum, alatum. Caput parvum, rotundatum, insertum, oculis parvis rotundatis, vix prominulis. Labrum breve, apice truncatum, lateribus rotundatum, mandibulas haud obtegens. Mandibulae muticae. Maxillae (t. III. f. 24.) mala exteriore interiorem haud superante, cornea, apice membranacea, dense pubescente, mala interiore cornea, margine interiore membraneo, longius dense introrsum barbato, margine exteriore incrassato, apice unco simplice introrsum spectante malam terminante. Palpi maxillares breviores, articulo secundo tertio paulo brevior, hoc parum incrassato, quarto tenui, subulato. Labium (t. III. f. 15.) mento brevi, apice emarginato, ligula angusta, palporum labialium articulum primum paulum excedente, bifida, paraglossis ligula duplo brevioribus, triangulariter acuminatis. Palpi labiales triarticulati, articulo primo crassiusculo, maiore, secundo tertio paulo brevior. Antennae breves, crassae, medium versus incrassatae, articulo secundo tertio duplo fere brevior, hoc leviter elongato, obconico, articulis 5—10 subtransversis, arcte coactis, ultimo longiore, subacuminato. Thorax coleopterorum latitudine, brevis, antrorsum angustatus, apice profunde emarginatus, basi bisinuatus, margine laterali dilatato, plano, angulis posterioribus elongatis, acuminatis. Scutellum triangulare. Elytra apice truncata, ad angulum exteriorem profunde emarginata, hoc elongato, acuminato. Abdomen apicem versus sensim angustatum, supra planiusculum, subtus leviter convexum, segmento septimo conspicuo, licet minuto. Pedes breves, intermedii approximati, tarsi antici 4-, posteriores 5-articulati, postici articulis 4 primis sensim subdecrecentibus.

### **O b s e r v a t i o n e s.**

1. Genus satis distinctum, cuius affinitates partim cum *Lomechusa*, partim

cum Aleochara esse videntur, illis thoracis elytrorumque forma et maxillarum mala interiore apice uncinata, his corporis habitu et labii structura propius accedit.

2. Differt ab Aleochara imprimis ligula graciliore, palpis labialibus triarticulatis, maxillarum mala interiore apice uncinata, palpis maxillaribus articulo conico-acuminato terminatis, thorace lateribus dilatato, tarsis anticis 4-articulatis.

3. Differt a Lomechusa praecipue antennis brevibus, fusiformibus, ligula bifida, paraglossis distinctis, maxillarum mala interiore unco simplice terminata, thorace apice emarginato, abdomine non revoluta, segmento sexto dorsali integro, pedibus brevibus, intermediis basi approximatis.

4. Differentias ab Euryusa in huius generis illustratione notavi.

5. Dom. Curtis Dinardam dentatum pro generis Lomechusae typo depinxit. Oris partes dissectae ab eadem desumtae, at uncus corneus, quo maxillarum mala interior terminatur, et paraglossae auctorem fugerunt, ceterum illae sat exacte depictae.

6. Sexus differentia abdominis segmento inferiore penultimo in maribus emarginato constare videtur.

7. Victus in formicetis.

\* 1. *D. dentata*: Nigra, thoracis lateribus elytrisque brunneis, capite thorace elytrisque crebre punctatis. — Long.  $1\frac{1}{2}$ —2 lin.

Mannerh. *Brachelytr.* 65. 1. — Boisd. et Lacord. *Faun. Ent. Paris.* I. 524. 1. —

— Runde *Brachelytr. Hal.* 27. 1. — Er. Col. *March. J.* 374. 1.

*Lomechusa dentata* Grav. *Mon.* 181. 4. — Gyll. *Ins. Suec.* II. 441 4. — Zetterst. *Ins. Lappon.* 74. 3. — Curtis *Britt. Ent.* IX. t. 410.

*Staph. strumosus* Payk. *Fn. Suec.* III. 402. 45.

Habitat in Europa.

Nigro-picea, sericeo-micans. Antennae capite sesqui longiores, crassiusculae, subfusiformes, articulo quarto praecedente dimidio brevior, at paulo crassiore, 5—10 transversis, sensim paulo crassioribus, ultimo praecedentibus ambobus longiore, at minus crasso, apicem versus sensim attenuato, nigrae, basi apiceque ferrugineae. Caput dense punctatum, fronte subimpressa, antice retusa, ore rufescente. Thorax coleopteris paulo latior, latitudine baseos plus duplo brevior, antorsum angustatus, lateribus fortiter ampliato-rotundatus, apice profunde emarginatus, basi profunde bisinuatus, angulis posterioribus retrorsum valde prominulis, anterioribus obtusis acuminatis, supra lateribus late explanatis, medio parum convexus, obsolete canaliculatus, confertim punctatus, breviter pubescens, niger, lateribus late obscure ferrugineis, seu saturate rufo-ferrugineus, disco nigro. Elytra thoracis longitudine, confertim punctata, breviter pubescentia, ferruginea, angulo apicis exteriore retrorsum valde prominulo, acuminato. Abdomen segmentis tribus anterioribus parcius, posterioribus tribus parce punctatis, supra parcius, subtus densius longiusque fulvescenti-pubescent, segmentorum marginibus anoque ferrugineis. Pedes ferruginei.

Obs. Insectum rarum, plerumque in Formicae rufae nidis obvium. Dom. Aubé eam etiam in Formicae cuniculariae societatibus observavit, et exempla prope Lutetiam in silva Boloniensi ex utroque formiceto cum formicis benevole mihi communicavit. Specimina cum *F. rufa* degentia omnino congruunt cum iis, quae e Germania, Hungaria et Suecia hic exstant, individua e nidis *F. cuniculariae* recepta illis quidem sunt duplo minora, ceterum vero nullo modo distincta. Cave igitur, ne in variis formicetis varias hospitem species gigni credas. Forte eiusdem speciei magnitudinis varietas e vario insecti loco natali derivanda, cum apud formicam maiorem individua maiora, apud minorem minora nascentur.

**Lomechusa Grav.**

*Grav. Mon. p. 178. — Mannerh. Brachelytr. p. 65. — Er. Col. March. I. p. 375.*

*Maxillae mala interiore unco duplice terminatae.*

*Ligula brevis, integra; paraglossae nullae.*

*Palpi labiales 3-articulati, articulo primo paulo maiore.*

*Tarsi antici 4-, posteriores 5-articulati, postici articulo primo elongato.*

Corpus latiusculum, crassiusculum, alatum. Caput minutum, sub-protectum, fronte planiuscula, oculis parvis, rotundatis, modice prominulis. Labrum transversum, apice truncatum, mandibulas haud obtegens. Mandibulae muticae. Maxillae (t. III. f. 25.) mala exterior cornea, apice membranea pubescenteque, malam interiorē paulo superante, hac cornea, margine interiore membraneo, pubescente, exteriorē incrassato, apice unco bispino, introrsum spectante, malam terminante. Palpi maxillares breviores, articulo secundo apicem versus leviter incrassato, tertio hoc paulo crassiore, haud incrassato, quarto parvo, subulato. Labium (t. III. f. 16.) mento transverso, apice emarginato, ligula lata, rotundata, integra, paraglossis nullis. Palpi labiales 3-articulati, articulo primo crassiusculo, maiore, secundo hoc duplo minore, tertio secundo tenuiore et parum longiore. Antennae elongatae, fortes, apicem versus subattenuatae, articulo primo incrassato, ultimo leviter elongato, acuminato. Thorax coleopterorum latitudine, apice truncatus, basi bisinuatus, margine laterali dilatato, incrassato, angulis anterioribus rotundatis, posterioribus acuminatis, disco plano. Scutellum triangulare. Elytra apice truncata. Abdomen parallelum, subtile convexum, supra planum, revolutum, segmentis tribus primis margine barbato, sexto apice late profundeque emarginato, lobis acuminatis, septimo superiore in praecedentis sinu sat conspicuo, inferiore occulto. Pedes elongati, crassi, intermedii basi late distantes, tibiae apice subattenuatae, tarsi tenues, haud elongati, antici 4-, posteriores 5-articulati, postici articulo primo modice elongato.

**Observationes.**

1. Habitus huius generis valde singularis, corpore in hac tribu maiusculo, crasso, abdomine fortiter revolutum, capite minuto, antennis validioribus, filiformibus, elongatis, acuminatis, pedibus elongatis, sat robustis, thorace margine dilatato, basi profunde bisinuato, abdominis segmento dorsali sexto emarginato. Maxillarum structura eadem fere ut in *Dinarda*, at mala interior unco duplice sive fisso terminata. Labii forma magis cum *Gyrophaena* convenit, at ligula latior et palpi labiales triarticulati. Habitu *Myrmedoniae* nunnulæ quodammodo *Lomechusae* speciem prae se ferre videntur antennis fortioribus rigidis, abdomine revolutum et pedibus intermediis basi late distantibus.

2. Gravenhorst genus *Lomechusam* praecipue secundum thoracis elytrorumque formam condidit, quare etiam *L. dentata*, reliquis omnibus notis adeo distinctam, huic generi subiunxit. Mannerheim, *Lomechusam* et *Dinardam* palporum maxillarum articulo ultimo conico-acuto a reliquis Aleocharinis distinxit, in *Lomechusa* autem palpi maxillares articulo terminali eodem modo quam in reliquis huius tribus generibus constructo mihi videntur.

3. *Atemeles Dillwyn* seu *Goniodes Kirby*, (Stephens Illustr. V. p. 107.) certe idem hoc genus, cui *L. strumosa* adscribitur, thoracis angulis posterioribus obtusis a *Lomechusa* thoracis angulis posterioribus acuminatis spinosis distinctum. *Lomechusae* typum Dom. Stephens *L. emarginata* affert, dubito autem, an eadem sit ac nostra, cum ei antennae medio magis incrassatae et fortius fusiformes quam *Lomechusae* et *Dinardae* tribuantur (p. 109.); neque suspicandum est, hanc meram esse *Dinardam* dentatam, cum abdominis segmentum penultimum emarginatum, processibus duobus styliformibus instructum et pedes elongati descripta sint.



4. Hospitatur hoc genus in formicetis. Villis in abdominis basi forte quaedam analogia est cum illis, quos Müller (Germ. Mag. III. p. 79. et 95.) in Clavigero foveolato observavit. Notatu dignum videtur, quod reliqua huius tribus insecta, quaecunque in formicetis degunt, his villis destituta sunt.

\* 1. *L. strumosa*: Ferruginea, thorace subtiliter granulato, canaliculato, antrorsum angustato. — Long. 3 lin.

Grav. Mon. 179. 1. — Gyll. Ins. Suec. II. 437. 1. — Sahlb. Ins. Fenn. I. 403. 1. — Zetterst. Ins. Lappon. 64. 1. — Mannerh. Brachelytr. 65. 1. — Runde Brachelytr. Hal. 27. 1.

*Staph. strumosus* Fab. Ent. Syst. I. II. 525. 28. Syst. El. II. 597. 36.

*Aleoch. strumosa* Grav. Micr. 91. 35. — Latr. Hist. nat. Crust. et Ins. IX. 387. 35.

Habitat in Germania, Suecia.

Obscure rufo-ferruginea, elytris pedibusque paulo dilutioribus. Antennae articulo primo breviusculo, crasso, tertio secundo duplo prope longiore, clavato seu apice subito incrassato, 4 — 10 cyathiformibus, sensim paululum longioribus, ultimo praecedente duplo prope longiore, acuminato, pube brevissima subtilissimaque depressa albida obducta. Caput laeve, pilis parvis depressis griseis adpersum, antice subretusum, fronte late impressa. Thorax basi coleopteris parum angustior, antrorsum angustatus, lateribus et angulis anterioribus rotundatus, apice truncatus, basi profunde bisinuatus, angulis posterioribus retrorsum vix prominulis, subrectis, apice summo rotundatis, lobo intermedio fortiter producto, rotundato, supra medio subdepressus, subtiliter canaliculatus, margine laterali late elevato, crassiusculo, parce subtiliter granulatus, pilis subtilissimis depressis griseis adpersus. Elytra thorace paululum longiora, apice truncata, subtiliter punctata, pube subtilissima depressa grisea adpersa. Abdomen medio fuscum, segmentis basalibus margine laterali et apicali fulvo-villosis, segmento dorsali sexto minus profunde emarginato, lobis lateralibus triangularibus. Pedes fulvo-tomentosi, basi glabri.

Differt sexu abdominis segmenti dorsalis quinti angulo apicali aut simpliciter aut lateraliter elevato prominuloque.

Obs. Prof. Sahlberg hanc speciem in societate cum *Formica*, (*Myrmica*) rubra lectam esse memorat.

\* 2. *L. paradoxa*: Ferruginea, thorace laevi, utrinque leviter impresso, lateribus rotundato-ampliato. — Long. 2 — 2½ lin.

Grav. Mon. 180. 3. — Gyll. Ins. Suec. II. 438. 2. — Germ. Faun. Ins. Europ. V. 2. — Mannerh. Brachelytr. 65. 2. — Guér. Iconogr. Regn. Amin. Ins. t. 10 f. 6. — Boisd. et Lacord. Faun. Ent. Paris. I. 526. 1. t. 2. f. 2. — Runde Brachelytr. Hal. 27. 2. — Er. Col. March. I. 376. 1.

*Staph. emarginatus* var. Payk. Mon. Staph. p. 57. not. Faun. Suec. III. p. 402. not. — Oliv. Ent. III. 42. t. 2. f. 12. b.

Habitat in Europa.

Sequenti valde affinis, at plerumque paulo maior, saturate ferruginea vel brunneo-testacea. Antennae articulo primo crasso, tertio elongato, secundo duplo prope longiore, apicem versus sensim leviter incrassato, 4 — 10 crassitie aequalibus, sensim paulo longioribus, prioribus transversis, ultimis crassitie haud longioribus, ultimo praecedentibus ambobus longiore, sensim attenuato et acuminato, pube subtilissima longiore albida vestita,

certo situ quasi velatae. Caput inter antennas leviter impressum, glabrum, nigricans, ore ferrugineo. Thorax latitudine dimidio prope brevior, coleopteris vix angustior, antrorsum haud angustatus, lateribus ampliatus, pone medium subsinuatus, ante medium cum angulis anterioribus rotundatus, apice truncatus, basi profundius bisinuatus, angulis posterioribus oblique productis, acutis, lobo intermedio subtruncato, laevis, glaber, obsolete canaliculatus, utrinque leviter impressus, margine laterali elevato, incrassato, Elytra thorace paulo longiora, apice sinuata, confertissime subtilissimeque punctata, subtilissime fulvo-pubescentia. Abdomen parce subtilissime punctulatum, segmentis basalibus margine laterali fulvo-villoso, segmentis quarto quintoque nigris, segmento dorsali sexto profunde emarginato, lobis acuminatis, spiniformibus, apice villosis. Pedes glabri.

Sexus differentia latet.

Obs. 1. In *Myrmicae rubrae* nidis D. Aubé hanc speciem cepit, „in societate cum *Formica parva nigra*” eam repertam esse refert Gyllenhal (Ins. Suec. IV. p. 495.); ipse eam in gramine prope *Formicae rufae* nidos legi. Conf. quae apud Dinardam dentatam observavi.

2. Specima illa, quae ipse prope oppidum Freyenwalde legi cum exemplis Suecis a Dom. Schüppel communicatis congruunt statura maiore et colore saturatione. Exempla collectionis Hoffmannseggianae, a Dahl in Germania transrhenana lecta his sunt minora et pallidiora, statura et colore L. emarginatae aequalia. Specimina Lutetiae collecta, a Dom. Aubé mihi communicata, his intermedia sunt.

3. *Staphylinus emarginatus* Oliv. a Gravenhorstio ad hanc, a Gyllenhalio ad sequentem speciem allegatur, mihi ambae ab Oliviero sub nomine illo confusae videntur; fig. a. L. emarginatam, fig. b. L. paradoxam, praecipue secundum thoracis formam bene exhibet.

\* 3. *L. emarginata*: *Ferruginea*, *thorace subtilissime punctulato*, *antrorsum angustato*. — Long. 2 lin.

Grav. Mon. 179. 2. — Gyll. Ins. Suec. II. 440. 3. — Sahlb. Ins. Fenn. I. 404. 2. — Mannerh. Brachel. 65. 3. — Boisd. et Lacord. Faun. Ent. Paris. I. 526. 2. — Runde Brachelytr. Hal. 27. 3. — Er. Col. March. I. 376. 2.

*Staphyl. emarginatus* Payk. Mon. Staph. 56. 40. Faun. Suec. III. 401. 44. — Fab. Ent. Syst. I. II. 528. 46. Syst. El. II. 600. 57. — Oliv. Ent. III. 42. 31. 44. t. 2. f. 12. a.

Habitat in Europa.

Ferruginea, vel rufo-testacea, abdomine ante apicem pectoreque piceis nigrisve, saepius quoque capite, nonnunquam etiam thoracis disco vel thorace fere toto nigricantibus. Antennae articulo primo crasso, secundo tertioque obconicis, hoc paululum longiore, 4 — 10 cylindricis, sensim paulo longioribus, prioribus crassitie vix brevioribus, ultimis eadem paulo longioribus, apicali elongato, praecedentibus tribus coniunctis longitudine aequali, sensim attenuato et acuminato, dense subtilissime longius albido-pubescentes ut in praecedente. Caput subtilissime punctulatum, inter antennas obsolete impressum. Thorax latitudine baseos dimidio brevior, basi coleopterorum prope latitudine, antrorsum aequaliter angustatus, ante medium cum angulis anterioribus rotundatus, apice truncatus, basi profunde bisinuatus, angulis posterioribus oblique productis, valde acuminatis, lobo intermedio leviter rotundato, supra subtilissime punctulatus, obsolete canaliculatus, utrinque leviter impressus, margine laterali elevato, haud incrassato. Elytra thorace

paulo longiora, apice exterius leviter sinuata, confertissime subtiliter punctata, pube subtilissima depressa fulva densius vestita. Abdomen parce subtilius punctatum, segmentis duobus primis margine laterali fulvo-villoso, dorsali penultimo profunde emarginato, lobis acuminatis, spiniformibus, apice villosis. Pedes glabri.

Sexus differentia latet.

Obs. 1. Praecedenti simillima et aegre distinguenda, differt, autem thorace antrorsum sensim angustato, angulis posterioribus paulo acutioribus, dorso punctulato et antennis paulo longioribus, articulo ultimo magis, articulo tertio autem non elongato.

2. Habitat secundum Gyllenhal semper in Formicae rufae nidis, Dom. Aubé autem specimen in Formicae cuniculariae societate legit. Forte, ut affines, in variis formicetis obvia.

3. In Museo Regio exemplum inveni patria America septentrionali signatum, certe ad hanc speciem pertinet, an etiam patria eius indubia?

### Species mihi invisae.

4. *L. inflata*: Parum punctata, fusco-rufescens, antennis subfiliformibus, thoracis lateribus incrassatis, elytris pedibusque saturate rufo-testaceis, thoracis elytrorumque angulis posticis prominulis.

Zetterst. Faun. Lappon. I. 95. 1. Ins. Lapp. 74. 2.

Habitat in Lapponia rarissime, sub cortice Betulae albae emortuae et trunco perforato unicum specimen cum Formica rufa ad Wittargi Lapponiae Tornensis, initio mensis Junii inveni.

*L. strumosae* similis et affinis, sed differt antennis dilutioribus, thoracis lateribus magis incrassatis, disco obscuriori, et quod duplo minor et ultra, *L. paradoxae* aequalis. Caput parvum, nigrum, laeve, fronte foveata. Os rufescens. Antennae elongatae, rufo-ferrugineae, articulo primo crassissimo, 2 brevissimo, 3 longiore, obconicis, 4—10 nodiformibus, remotis, subaequalibus. ultimo elongato, angustiore, conico, acuto. Thorax fere ut in *L. strumosa* formatus, scilicet brevis, transversus, postice in medio productus, utrinque sinuatus, angulis subrectis, minime prominulis, supra discum depressus, inaequalis, vix canaliculatus, margine antico et postico parum, lateribus valde elevatis, et his etiam incrassatis, fusco-piceus, nitidus, lateribus ferrugineo-testaceis, opacis, totus parce et obsoletissime punctulatus, subpilosus. Elytra, abdomen et pedes omnino ut in *L. strumosa*, etiam quoad pubescentiam. (Zett.)

Obs. Pedibus tomentosis propius ad *L. strumosam*, quam ad *L. paradoxam* accedere videtur.

### *Silusa* Er.

Er. Col. March. I. p. 377.

*Maxillae malis inaequalibus, interiore elongata, cornea, apice uncinata, intus subserata.*

*Ligula elongata, angusta, integra.*

*Palpi labiales exarticulati, setacei.*

*Tarsi antici 4-, postici 5-articulati, omnes articulis primis aequalibus.*

Corpus elongatum, lineare, supra depressum, subtus sat convexum, ala-



tum. Caput exsertum, deflexum, basi haud constrictum, subglobosum, ore attenuato, leviter elongato, oculis mediocribus, minus prominulis. Labrum minutum, seniorbiculare. Mandibulae muticae. Maxillae (t. III. f. 26.) mala interiore valde elongata, cornea, cultriformi, apice acuto introrsum adunco, margine interiore ante apicem spinulis minutis brevissimis serrulata, dein tenuiter ciliata, mala exteriora haud elongata, interiore duplo breviora, medio cornea, basi apiceque membranacea, portione membranacea apicali pubescente. Palpi maxillares malam interiorem parum excedentes, articulo secundo tertioque subaequalibus, illo elongato-subobconico, hoc tumidulo, quarto minuto, subuliformi. Labrum (t. III. f. 17.) mento sat magno, apice sat fortiter emarginato, ligula tenui, filiformi, integra, paraglossis nullis. Palpi labiales setacei, valde elongati, prominuli, subarticulati, articulo primo secundoque subaequalibus, tertio minimo. Antennae crassiusculae, articulo secundo tertioque obconicis, subaequalibus, 5 — 10 subtransversis, ultimo oblongo-ovato. Thorax transversus, basi coleopterorum prope latitudine, apicem versus parum angustatus. Scutellum triangulare. Elytra apice truncata, ad angulum anteriorem profunde excisa. Abdomen parallelum. Pedes breviusculi, intermedii basi approximati, tarsi antici 4-, posteriores 5-articulati, articulis 4 primis aequalibus.

#### O b s e r v a t i o n e s.

1. Genus oris fabrica satis distinctum, mandibularum mala altera solita structura, altera elongata, ambiguum quasi inter praecedentia et sequentia.

2. Habitu Aleocharae generis quibusdam speciebus accedit, a quibus tarsorum structura differt. Tarsis anticis 4-articulatis, posterioribus articulis 4 primis inter se aequalibus cum Homalotis convenit, a quibus praeter caput basi haud constrictum et os attenuatum nisi ore accuratius inspecto aegre distinguatur.

3. Abdominis segmentum inferius sextum maris apice bisinuatum, medio leviter productum, feminae rotundatum.

4. De victu nihil mihi compertum.

\* 1. *S. rubiginosa*: *Confertim punctata, nigra, antennis, pedibus elytrisque rufis, his circa scutellum anguloque exteriori apicis fuscis, thorace basi foveolato.* — Long.  $1\frac{3}{4}$  lin.

*Er. Col. March. I.* 378. 1.

Habitat Berolini, Dom. Schüppel et Dom. Weber, in Austria, Dom. Schüppel, Lutetiae, Mus. Dom. Aubé.

Corpus totum griseo-puberulum. Antennae vix capitis thoracisque longitudine, rufae. Caput laeve, subnitidum, nigrum, ore rufescente. Thorax coleopteris paulo angustior, latitudine duplo fere brevior, lateribus leviter rotundatus, antrorsum vix angustatus, angulis anterioribus rotundatis, posterioribus rectis, parum convexus, subtilissime punctatus, basi foveola transversa impressus, niger, margine laterali obscure rufescente. Elytra thorace paulo longiora, confertim fortiter punctata, sutura elevata, opaca, rufa, macula communi scutellari anguloque posteriore exteriori fuscis, seu fusca, vitta lata oblique ab humero ad angulum interiorem descendente rufa. Abdomen densius fortiter punctatum, minus subtiliter pubescens, nigrum, segmentis quatuor primis margine piceis. Pedes rufi.

Mas abdominis segmento dorsali quinto acute carinato distinctus.

Obs. 1. *Al. foveolatae* Dej. nomine hoc insectum in collectione D. Aubé signatum vidi.

2 Labrum maiusculum, mandibulas omnino obtegens.

\* 2. *S. rubra: Punctatissima, rufa, abdomine supra parce punctato, ante apicem capiteque piceis, thorace basi fovea lata impresso. — Long.  $1\frac{3}{4}$  lin.*

Habitat in Germania meridionali, Dom. Schüppel.

Statura omnino praecedentis, rubra seu rubro-testacea, punctatissima, subtiliter pubescens. Antennae capitis thoracisque longitudine, apicem versus subincrassatae. Caput confertim sat fortiter punctatum, infra antennas laeve, nigro-piceum, oris partibus rufo-testaceis. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus leviter rotundatus, angulis omnibus subrectis, anterioribus deflexis, leviter convexus, confertim sat profunde punctatus, basi fovea transversa, antrorsum explanata, media et utrinque puncto impressus. Elytra thorace paulo longiora, fortiter punctata, subrugosa. Abdomen elongatum, obscure rufum, ante apicem nigricans, supra parce punctatum, segmentis posterioribus sublaevibus. Pectus piceum. Pedes flavo-testacei.

Mas abdominis segmento dorsali quinto bicarinato insignis. Femina mihi invisae.

Variatur colore minus saturo, rufo-testaceo, capite pectoreque concoloribus.

Obs. Labrum minus quam in praecedente, et mandibulae sub eodem prominentes. Nihilominus cum praecedente, quantum examinatione exteriore videre licet, oris reliqui et praecipue maxillarum structura haud differt.

### **Pronomaea Er.**

*Er. Col. March. I. p. 378.*

*Maxillae malis aequalibus, elongatis, interiore cornea, apice uncinata, intus serrata.*

*Ligula bifida, minuta, sub mento occulta; paraglossae nullae.*

*Palpi labiales exarticulati, setacei.*

*Tarsi antici 4-, posteriores 5 articulati, postici articulo primo elongato.*

Corpus elongatum, lineare, supra subdepressum, infra modice convexum, alatum. Caput exsertum, cernuum, basi haud constrictum, ore attenuato, valde elongato, oculis mediocribus, parum prominulis. Labrum dependens, fere oblongum, apice rotundatum, mandibulas obtegens. Mandibulae muticae, membranula fimbriatae, apice libero, acuto, introrsum adunco. Maxillae (t. III. f. 27.) mala utraque elongata, lineari, tenui, interiore cornea, apice acuto, introrsum adunco, margine interiore apicem versus spinulis minutis deorsum versis serrulato, inferius ciliato, mala exteriore membranacea, margine tenuissimo interiore apiceque membranacea, apice pubescente, malae interiori aequali. Palpi maxillares elongati, articulis secundo tertioque aequalibus, parum incrassatis, quarto minimo, subulato. Labium (t. III. f. 18.) mento sat magno, apice profunde exciso, ligula brevissima, biloba, sub mento occulta, paraglossis nullis. Palpi labiales valde elongati, setacei, exarticulati. Antennae apicem versus sensim subincrassatae, articulo secundo

tertioque subaequalibus, ultimo ovato. Thorax coleopterorum latitudine, apicem versus haud angustatus. Scutellum triangulare. Coleoptera apice coniunctim leviter emarginata, ad angulum apicalem anteriorem profundius excisa. Abdomen lineare. Pedes breviusculi, intermediis brevi spatio distantes, tarsi antici 4-, posteriores 5-articulati, postici articulo primo subelongato.

#### Observationes.

1. Mentum magnum, ligulam omnino occultans, quae nonnisi mento destructo conspicitur.

2. Habitus corporis linearis Oxypodae Sect. II, a qua autem non modo tarsis anticis 4-articulatis distinguitur, sed capite etiam tumido, ore producto, attenuato, fere rostrato, palpis maxillaribus elongatis et labialibus quam plurimum exsertis, setaceis discedit.

3. Sexus differentiam non cognovi.

4. Victus in silvis in musco et sub foliis deciduis.

\* 1. *P. rostrata*: *Nigra, nitida, thorace basi foveolato, elytris pedibusque piceis.* — *Long.* 1½ lin.

*T.* 1. *f.* 3.

*Er. Col. March.* I. 379. 1.

Habitat in Germania, Gallia. In Sardinia etiam a Dom. Prof. Gené lecta.

Linearis, nigra, nitida, subtiliter punctata, tenuiter pubescens. Antennae capite thoraceque breviores, crassiusculae, obscure piceae, basi dilutiores, apice summo ferrugineo. Caput sat magnum, subtilissime punctulatum, fronte leviter convexa, clypeo apice membranaceo, luteo, labro piceo. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus subrectis, basin versus subangustatus, angulis anterioribus rotundatis, deflexis, posterioribus obtusis, supra leviter convexus, subtiliter punctulatus, basi foveola impressus. Elytra thorace paulo longiora, crebre fortius punctata, longius pubescentia, picea. Abdomen supra parce punctatum, fere laevigatum, parce pubescens, ano piceo. Pedes rufo-picei.

### **Diglossa Haliday.**

*Entomol. Mag.* IV. p. 252.

*Maxillae malis aequalibus, elongatis, exteriore cornea, apice uncinata, intus serrata.*

*Ligula sub mento occulta.*

*Palpi labiales exarticulati, setacei.*

*Tarsi omnes 4-articulati.*

Corpus elongatum, apterum. Caput exsertum, porrectum, thorace haud angustius, basi haud constrictum, ore attenuato, producto, oculis minutis. Labrum transversum, apice leviter sinuatum. Mandibulae productae, prominulae, angustae, acutae, apice leviter introrsum incurvae, medio obsolete dentatae. Maxillae malis elongatis, longitudine aequalibus, exteriore setacea, interiore lineari, apice introrsum curvata, cornea, margine interiore subtiliter denticulato. Palpi maxillares elongati, articulo primo minuto, secundo tertioque longis, longitudine subaequalibus, hoc vix clavato, quarto obsoleto. Mentum transversum, antrorsum angustatum, apice trun-



catum. Ligula sub mento occulta. Palpi labiales exarticulati, setacei, porrecti. Antennae graciliores, articulo primo cylindrico, secundo paulo longiore, clavato, tertio brevior, obconico, sequentibus brevibus, subglobosis, extremis sensim paulo crassioribus, fere transversis, ultimo iterum longiore, subgloboso-ovato. Thorax coleopteris angustior, basin versus leviter angustatus. Elytra truncata, ad angulum anteriorem vix sinuata. Abdomen elongatum, lineare. Pedes breviores, tibiis anticis angulo apicali posteriore profunde emarginato, sinu dentibus prominulis acutis terminato, tarsis omnibus 4-articulatis, anterioribus brevissimis, posterioribus brevibus, articulo primo paulo longiore, quarto (ungiculari) primo brevior, crassior, unguiculis brevibus, fortiter curvatis, validis.

1. *D. mersa*: *Nigra, subtiliter pubescens punctataque, palpis pedibusque ferrugineis.* — Long. 1 lin.

*Haliday Ent. Mag. IV. p. 252.*

Habitat in arenis maritimis Hiberniae, aestu alterno opertus et reiectus.

*Nigra*, leviter pubescens, capite, thorace elytrisque subtiliter punctulatis. Antennae capite thoraceque paulo longiores, apicem versus perparum incrassatae, obscure ferrugineae. Palpi saturate ferruginei. Mandibulae dilutius ferrugineae. Thorax capite elytrisque angustior, latitudine longior, lateribus ante medium rotundatus, basin versus angustatus, basi apiceque truncatus. Elytra thorace haud longiora. Pedes saturate ferruginei, tibiis pubescentibus medio femoribusque obscurioribus, tarsis dilutis.

Obs. Insectum mihi invisum, ab auctore citato l. c. accuratius descriptum et icone ligno inciso illustratum, corporis habitu et oris structura summam cum *Pro-nomaea rostrata* nostra similitudinem prae se fert. Tarsi 4-articulati, articulo ultimo brevi incrassato, unguiculis brevibus, fortibus, fortiter curvatis instructo et tibiae anticae apice emarginatae huic generi peculiares. Victus Blediorum quorundam. Analogia cum *Oxytelis* tibiatarum anticarum structura probatur. Nomen generis ex errore, quo auctor palpos labiales pro ligulae lacinias habuit, ortum, non sine dubio immutatum recepi.

## **Myllaena Er.**

*Er. Col. March. I. p. 382.*

*Centroglossa Matthews Ent. Mag. V. p. 194.*

*Maxillae malis aequalibus, elongatis, anteriore cornea, apice uncinata, intus serrata.*

*Ligula brevis, integra.*

*Palpi labiales exarticulati, setacei.*

*Tarsi antici 4-, posteriores 5-articulati, antici articulo primo brevissimo, posteriori subelongato.*

Corpus oblongum, antice obtusum, anum versus sensim attenuatum, acuminatum, convexum, alatum. Caput basi haud constricta a thorace receptum, deflexum, subglobosum, ore attenuato, modice elongato, fronte convexa, oculis lateralibus, mediocribus, parum prominulis. Labrum coriaceum, semi-orbiculare, mandibulas obtegens. Mandibulae muticae. Maxillae (t. III. f. 29.) mala utraque valde elongata, anteriore cornea, lineari-cultriformi, apice acuto introrsum adunco, margine anteriore apicem versus

spinulis brevissimis subtiliter serrulato, exteriore membranea, lineari. interiori aequali, apice pilosulo. Palpi maxillares malas parum superantes, articulo secundo satis elongato, tertio hoc paulo brevior, tumidulo, quarto minuto, subulato. Labium (t. III. f. 20.) mento transverso, apice leviter emarginato, utrinque spina porrecta, lineari, valde elongata terminata, ligula brevissima, rotundata, integra, paraglossis nullis. Palpi maxillares valde elongati, prominuli, setacei, exarticulati. Antennae tenuiusculae, articulo secundo tertio sesqui longiore, ultimo oblongo-ovato, apice acuminato. Thorax basi coleopterorum latitudine, eaque amplectens, lateribus rotundatus, apicem versus fortiter angustatus, convexus. Scutellum thoracis margine obtectum. Coleoptera apice coniunctim emarginata, utrinque ad angulum exteriorem excisa. Abdomen apicem versus sensim attenuatum, acuminatum, stylis analibus exsertis. Pedes breviusculi, intermedii basi approximati, tibiis glabris, tarsis anticis 4-, posterioribus 5-articulatis, anticis articulo primo brevissimo, posticis elongato.

#### Observationes.

1. Corpus totum subtilissime punctatum, pube densa subtilissima brevissimaque sericeum. Thorace amplo et abdomine apicem versus attenuato habitus omnino Tachypororum, Oxypodis sectionis prioris quoque non dissimiles minutae hae species, quibuscum insuper elytris apice ad angulum exteriorem excisis congruunt, distinctae tamen ore attenuato, subrostrato. Palpi labiales setacei plerumque prominent.

2. Ingeniose Com. Mannerheim Myllaenam sequenti generi adscribit, attamen accuratius examinata oris et tarsorum structura, tibiisque haud spinulosis satis distincta.

3. Abdominis segmentum inferius sextum in altero sexu apice truncatum, in altero acute rotundatum.

4. Victus in silvis in musco et sub foliis deciduis.

\* 1. *M. dubia*: Nigra, opaca, cinereo-sericea, thoracis angulis posticis prominulis. — Long.  $1\frac{1}{4}$  lin.

Er. Col. March. I. 383. 1.

Aleochar. dubia Grav. Mon. 173. 67. — Gyll. Ins. Suec. II. 426. 48. — Sahlb. Ins. Fenn. I. 395. 85. — Zetterst. Ins. Lappon. 79. 32.

Gymnusa dubia Mannerh. Brachelytr. 66. 2.

Centroglossa conuroides Matthews Ent. Mag. V. p. 195. f. 1.

Habitat in Europa.

Antennae fuscae, articulo primo flavo. Palpi flavi, articulo tertio fusco. Thorax antrorsum angustatus, lateribus angulisque anterioribus rotundatis, angulis posterioribus retrorsum prominulis, basi profundius bisinuatus. Abdomen apice segmentorumque ventralium marginibus rufo-piceis. Pedes flavi, femoribus fuscis.

\* 2. *M. intermedia*: Nigra, opaca, cinereo-sericea, thoracis angulis posticis subrectis. — Long.  $\frac{3}{4}$  lin.

Er. Col. March. I. 383. 2.

Habitat in Germania.

Praecedenti affinis, at plus duplo minor. Antennae fuscae, articulo primo flavo. Os flavum. Thorax antrorsum angustatus, lateribus minus

rotundatus, basi leviter bisinuatus, angulis posterioribus parum prominulis, subrectis. Abdomen apice testaceum. Pedes testacei, postici fuscescentes.

Variat immatura colore rufo-testaceo, abdominis basi pectoreque nigris.

\* 3. *M. minuta*: Nigra, opaca, cinereo-sericea, thoracis angulis posticis obtusis. — Long.  $\frac{1}{2}$  lin.

Er. Col. March. I. 384. 3.

*Aleoeh. minuta* Grav. Mon. 174. 68. — Gyll. Ins. Suec. II. 427. 29.

Habitat in Germania, Suecia.

Praecedenti simillima, at paulo angustior et duplo minor. Antennae articulo primo concolore. Os fusco-testaceum, palpis maxillaribus fuscis. Thorax antrorsum vix angustatus, lateribus parum rotundatus, basi levissime bisinuatus, angulis posterioribus obtusis. Pedes fusco-testacei.

### **Gymnusa Karsten.**

Grav. Mon. p. 173. — Mannerh. Brachelytr. p. 66. — Er. Col. March. I p. 380.

*Maxillae malis aequalibus, elongatis, interiore cornea, apice uncinata, intus inaequaliter serrata.*

*Ligula bifida, laciniis porrectis, elongatis, setaceis, palpis aequalibus; paraglossae nullae.*

*Palpi labiales setacei, exarticulati.*

*Tarsi omnes 5-articulati, postici articulo primo elongato.*

Corpus oblongum, utrinque attenuatum, convexum, alatum. Caput basi haud constricta a thorace receptum, deflexum, subglobosum, ore attenuato, elongato, fronte convexa, oculis facialibus, mediocribus, parum prominulis, ovatis. Labrum maiusculum, suborbiculare, dependens, mandibularum apicem tegens. Mandibulae elongatae, lineares, apice acute bidentatae, dente altero apicali, altero ante apicem sito, margine interno infra dentes membranula interne acute denticulata fimbriato. Maxillae (t. III. f. 28.) mala utraque cornea, valde elongata, interiore cultriformi, apice acuto, adunco, margine interiore ante apicem inaequaliter acute serrato, exteriori lineari, ad basin sinuato-excisa, apice setulis duabus barbulaque introrsum vergente terminata. Palpi maxillares itidem valde elongati, tennes, articulo secundo longissimo, tertio hoc paulo brevior, apicem versus parum incrassato, quarto minimo, subulato. Labium (t. III. f. 19.) mento sat magno, apice profunde emarginato, ligula elongata, bifida, laciniis coriaceis, linearibus vel subsetaceis, palpis subaequalibus, paraglossis nullis. Palpi labiales elongati, setacei, apice obsolete articulati, ita ut, si articulatos respicias, articulus primus sit longissimus, secundus minimus, tertius hoc paululum longior. Antennae tennes, articulis 2—10 subobconico-teretibus, 11mo ovato. Thorax basi coleopterorum latitudine, eaque amplexans, basi utrinque subsinuatus, antrorsum angustatus, lateribus rotundatus, supra convexus. Scutellum thoracis margine obtectum. Coleoptera apice coniunctim emarginata, utrinque ad angulum exteriorem excisa. Abdomen apicem versus sensim attenuatum, stylis analibus distinctis. Mesosternum subtiliter carinatum. Pedes brevius-



sculi, intermedii basi approximati, tibiis spinulosis, tarsis omnibus 5-articulatis, postici articulo primo valde elongato.

#### O b s e r v a t i o n e s.

1. Genus insigni modo distinctum, neque cum aliquo confundendum. Mandibulae acute dentatae, oculi in capitis facie anteriore siti, tibiae spinulosae et styli anales exserti characteres sunt singulares in hac tribu, in cuius finem collocavi hoc genus, pariter ad Tachyporinos atque ad Staphylininos vergens.

2. Abdomen segmento sexto inferiore maris apice leviter emarginato, septimo fortiter inciso, feminae segmento sexto apice utrinque leviter sinuato, medio acute rotundato, septimum occultante. Qua sexus differentia a reliquis Aleocharinis hoc genus quodammodo discedit, magis autem Staphylininis quam Tachyporinis appropinquatur.

3. Victus sub musco et ad arborum radices.

\* 1. *G. brevicollis*: *Nigra, antennarum articulo primo tarsisque testaceis.*  
— *Long.*  $2\frac{1}{2}$  lin.

*Mannerh. Brachelytr.* 66. 1. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 534. 1. — *Er. Col. March. I.* 381. 1.

*Staphyl. brevicollis* Payk. *Faun. Suec. III* 398. 40.

*Aleochar. brevicollis* Gyll. *Ins. Suec. II.* 425. 47. — *Sahlb. Ins. Fenn. I.* 394. 84. — *Zetterst. Faun. Lappon. I.* 103. 22. *Ins. Lappon.* 79. 31.

*Aleochar. carnivora* Grav. *Mon.* 171. 60.

*Aleochar. excusa* Grav. *Mon.* 172. 66.

Habitat in Germania, Suecia.

Nigerrima, subnitida, subtilissime tenuissimeque cinereo-pubescens. Antennae articulo primo rufo. Caput laeve. Thorax basi coleopterorum latitudine, latitudine paulo brevior, basi subsinuatus, angulis posticis obtusiusculis, convexus, subtiliter punctatus. Elytra thoracis longitudine, confertim subtiliter punctata, opaca. Abdomen minus subtiliter punctatum. Pedes nigri, tarsis rufo-testaceis.

2. *G. laticollis*: *Atra, tenuiter sericeo-pubescens, palpis tarsisque luteofuscis.* — *Long.*  $1\frac{3}{4}$  lin.

Habitat in Germania, Gallia.

Habitu omnino praecedentis, at plus duplo minor, tota confertissime subtilissimeque punctulata, atra, pube tenuissima cinereo-sericea dense vestita. Antennae filiformes, basi obscure fuscescentes. Labrum transversim orbiculatum palpique flavescenti-fusca. Thorax coleopteris sublatior, brevis, longitudine duplo latior, basi bisinuatus, angulis posterioribus subrectis, retrorsum prominulis, supra leviter convexus. Elytra thorace longiora. Pedes nigri, geniculis tarsisque fusco-luteis.

Obs. Statura triplo fere minore, pube densa, sericante, labro transverso, thorace brevior et latior a praecedente facile distinguitur.

## **Tribus secunda.**

### **Tachyporini.**

*Stigmata prothoracica conspicua.*

*Antennae sub frontis margine laterali insertae.*

---

Labrum integrum, corneum, fimbria et laciniiis membraneis nullis.

Mandibulae haud prominentes, muticae, trigonae aut subfalcatae, limbo membraneo, interne ciliato, margini interiori affixo intus fimbriatae, apice libero acuto in plerisque;

valde elongatae, tenues, falcatae, acutae in Tanygnatho.

Maxillae cardine stipiteque corneis, malis coriaceis membranaceisve, exteriore apice, interiore margine interiore barbatis in plerisque;

malis, elongatis, linearibus, interiore apice unicata in Trichophyo.

Palpi maxillares 4-articulati, articulo primo minuto, secundo tertioque elongatis, quarto

minuto, subulato in Hypocypto, Conuro, Tachyporo, Mycetoporo; elongato, apice valde acuminato in Habrocero.

praecedenti aequali, apice leviter acuminato in Tachino, Trichophyo, Boletobio, Tanygnatho.

Mentum corneum, transversum, antrorsum angustatum, apice truncatum aut leviter emarginatum.

Labium membranaceum magis minusve exserendum.

Ligula membranea,

lata biloba, paraglossis ei connatis, linea ciliata indicatis in Conuro, Tachyporo, Habrocero, Tachino;

angustiore, paraglossis nullis in Boletobio et Mycetoporo;

rotundata, medio producta, paraglossis nullis in Hypocypto;

bifida, paraglossis eadem brevioribus in Trichophyo.

Palpi labiales triarticulati plerisque;

filiformes Conuro, Tachyporo, Habrocero, Tachino, Boletobio; articulo ultimo subulato aut fere subulato Hypocypto, Mycetoporo, Tanygnatho;

biarticulati (?) Trichophyo.

Antennae infra oculos sub capitis margine laterali insertae,

11-articulatae, filiformes plerisque;

tenuis, articulis singulis verticillato-pilosis, primis duobus crassiusculis in *Habrocero* et *Trichophyo*;

10-articulatae, clavatae in *Hypocyppto*.

Oculi parvi rotundati aut ovals, haud prominentes.

Caput thoraci insertum, basi haud constrictum, deflexum vel inflexum.

Thorax amplus, convexus, postice coleopterorum basin, antice caput amplectens.

Prosternum triangulare, spatio pone coxas anteriores membraneo.

Pectus metathoracis epimeris magnis, retrorsum prominentibus.

Scutellum triangulare.

Elytra pectore distincte longiora in *Hypocyppto*, *Conuro*, *Tachyporo*, *Habrocero*, *Tachino*;

aut pectus longitudine vix superantia in *Trichophyo*, *Boletobio*, *Mycetoporo*, *Tanygnatho*.

Abdomen apicem versus attenuatum, segmentis sex primis distinctis, septimo saepius occulto, stylis analibus plerumque haud distinctis;

marginatum in plerisque;

immarginatum in *Conuro*.

Pedes coxis anticis exsertis, conicis, posticis parallelopipedis, trochanteribus posticis fulcrantibus, tibiis aut omnibus aut posterioribus plerumque, nonnunquam etiam femorum apice spinulosis, tarsis

5-articulatis in plerisque,

4-articulatis in *Hypocyppto* et *Tanygnatho*,

anticis saepe dilatatis;

unguiculis tenuibus, falcatis, simplicibus.

Sexus differentia saepius abdominis segmentorum posteriorum figura et tarsis anticis maris fortius dilatatis cognoscitur, in generibus et speciebus variis vario modo se habens, et in iisdem describendis accuratius illustranda.

#### O b s e r v a t i o n e s .

1. Corpus minutum, agile, habitu quodam peculiari, convexus, ante et postice attenuatum, licet antice obtusiusculum, abdomine in vivis modice elongato.

2. Huius tribus minus numerosae plerasque sibi cognitae species Fabricius immerito *Oxyporo* subiunxit, Gravenhorst genera *Tachyporum* et *Tachinum* ex iis constituit, illud primo (*Micr. p. 124.*) species palporum maxillarium articulo quarto subulato, dein (*Mon. p. 1.*), methodo magis naturali, species elytris longioribus continens, hoc primo (*Micr. p. 134.*) species palpis maxillaribus filiformibus, dein (*Mon. p. 22.*) species elytris pectore haud longioribus amplectens. Genera hodierna pleraque iam ab auctore celebri familiarum nomine in utroque genere suo distincta sunt.

### **Hypocypthus Schüpp.**

*Er. Col. March. I. p. 387.*

*Hypocypthus Mannerh. Brachelytr. p. 58. — Boisd. et Lacord. Faun. Ent. Paris. I. p. 519.*

*Cypha Stephens Illustr. V. p. 187.*

*Antennae 10-articulatae, articulis ultimis tribus crassioribus.*

*Palpi maxillares articulo quarto subulato.*



*Ligula rotundata, integra.*  
*Elytra pectore longiora.*  
*Tarsi 4-articulati.*

Corpus parvum, globatum, alatum. Caput latum, inflexum, oculis rotundatis, parum prominulis. Labrum transversum, integrum. Mandibulae minutae, muticae. Maxillae (t. IV. f. 1. b.) malis membraneis, interiore apice oblique truncata, ibique barbata, exteriore extus profunde sinuata, apice introrsum barbata, interiorem paulum superante. Palpi maxillares articulo primo minimo, secundo elongato, subelavato, tertio ovato, tumidulo, quarto minimo, subulato. Labium (t. IV. f. 1. a.) mento transverso, apice truncato, ligula membranea, semiorbiculari, medio processu minuto umbonata, paraglossis nullis. Palpi labiales articulis duobus primis crassiusculis, secundo brevior, tertio sat elongato, tenui, subuliformi. Antennae 10-articulatae, articulis primis duobus crassiusculis, ultimis tribus leviter incrassatis, clavam oblongam formantibus. Thorax transversus, basi coleopteris latitudine subaequalis, antrorsum angustatus, rotundatus, basi utrinque sinuatus, leviter convexus. Scutellum obtectum. Elytra convexuscula, apice singula rotundata. Abdomen minus elongatum, apicem versus attenuatum. Pedes breves, debiles, intermedii basi satis distantes, interstitio aequali, coxae posticae haud amplae, parallelipedae, tibiae omnes laeves vel subtilissime setulosae; tarsi omnes 4-articulati, articulo primo leviter elongato, tarsorum anteriorum incrassato, sequentibus duobus minimis, ultimo primo paulo brevior.

#### Observationes.

1. Corpus laeve, parce pubescens, capite thoraceque fortiter inflectendis globatile, globatum Agathidiorum more seminulum mentiens.

2. Affinitates huius generis cum Aleocharinis admodum obviae videntur, ligulaeque forma affirmantur. Antennarum insertio autem huius Tribus. Cum genere Oligota arctissime coniungitur, non solum corporis figura, sed etiam articulorum antennarum tarsorumque numero comminuto.

3. Victus in sylvis, vel sub lignorum cortice recentiore, nondum exsiccato latitantes vel in gramine herbisque vagantes reperias. Reliqua vivendi ratio latet. Praeter indigenas paucas nulla mihi innotuit species.

4. Genus hocce sub Scaphidio apud Payk<sup>1</sup> et Gyllenhal, sub Tachyporo apud Gravenhorst militans, primus cognovit et proposuit Dom. Schüppel. Vulgo scribunt Hypocyptus contra regulas etymologicas, scribendum esset aut Hypocyphtus aut Hypocyptus. Hoc ultimum nomen ab inventore erudito pro suo agnoscitur.

\* 1. *H. longicornis*: Niger, nitidus, parce pubescens, antennis apicem versus sensim incrassatis pedibusque testaceis, thoracis angulis posterioribus subrectis. — Long.  $\frac{2}{3}$  —  $\frac{3}{4}$  lin.

Mannerh. Brachelytr. 58. 1. — Boisd. et Lacord. Faun. Ent. Paris. I. 521. 3. — Er. Col. March. I. 388. 1.

Scaphidium longicorne Payk. Faun. Suec. III. 340. 3. — Gyll. Ins. Suec. I. 191. 6 IV. p. 294. — Sahlb. Ins. Fenn. I. 63. 4.

Scaphidium acuminatum Marsh. Ent. Britt. 234. 3.

Tachyporus granulum Grav. Mon. 3. 1.

Hypocyptus granulum Boisd. et Lacord. Faun. Ent. Paris. I. 520. 1.

Hypocyptus globulus Boisd. et Lacord. Faun. Ent. Paris. I. 521. 2.

Habitat in Europa, sub arborum cortice.

Niger, nitidus, pube subtili depressa cinerea parce adpersus. Antennae testaceae, articulis duobus primis sat magnis, primo subgloboso, secundo cylindrico, 3 — 5 tenuibus, inter se aequalibus, crassitie vix longioribus, 6 et 7 paulo crassioribus et sensim paulo longioribus, subobconicis, sequentibus tribus clavam valde elongatam, parum incrassatam, fusiformem formantibus, penultimis duobus aequalibus, obconicis, ultimo his coniunctis longitudine aequali, at non crassiore, apicem versus sensim attenuato. Palpi testacei. Caput fronte planiuscula, laevi. Thorax basi coleopterorum prope latitudine, antrorsum angustatus, latitudine baseos plus duplo brevior, lateribus cum angulis anterioribus fortiter rotundatus, apice truncatus, basi leviter bisinuatus, angulis posterioribus subrectis, fortiter convexus, margine laterali dilatato, pallido, pellucido. Elytra thorace sesqui longiora, apice rotundata, convexa, parcius subtilissime punctata, nigro-picea. Abdomen parce subtilissime punctatum, apice plerumque piceum. Pedes testacei.

Mas abdominis segmento sexto inferiore apice profunde triangulariter exciso, femina eodem integro rotundato. Tarsi antici in utroque sexu simplices.

Obs. H. granulum Boisd. et Lacord. thorace elytris piceis, H. globulus eorundem auctorum nonnisi statura paulo minore a genuinis discedunt.

\* 2. *H. discoideus*: Niger, pubescens, antennis articulis ultimis tribus abrupte maioribus pedibusque testaceis, elytris disco rufis, thorace angulis posterioribus rotundatis. — Long.  $\frac{1}{2}$  lin.

Er. Col. March. I. 388. 2.

Habitat Berolini, Dom. Schüppel et ipse.

Praecedente dimidio vel duplo minor, nigro-piceus, pube longiore, depressa, fulvescenti-grisea densius vestitus. Antennae flavo-testaceae, pilosae, articulis 3—7 tenuioribus, 8 et 9 his duplo crassioribus, subobconicis, ultimo his coniunctis longitudine aequali, elongato, apice acuminato, his ultimis tribus clavam valde elongatam at parum incrassatam formantibus. Caput parcius subtiliter punctatum, nigrum, nitidum, ore testaceo. Thorax basi coleopterorum latitudine, antrorsum angustatus, latitudine baseos plus duplo brevior, basi subbisinuatus, angulis posterioribus rotundatis, transversim convexus, parcius subtilissime punctatus, piceus, margine basali tenui et laterali subdilatato testaceo-pallidis. Elytra thorace sesqui longiora, apice oblique truncata, leviter convexa, crebre subtiliter punctata, testaceo-rufa, basi et lateribus fusca. Abdomen parce subtiliter punctatum, piceum, supra segmentis anterioribus quatuor margine posteriore rufescente, ultimis duobus totis rufo-testaceis, subtus rufo-testaceum, basi piceum. Pedes testacei.

Mas abdominis segmento sexto inferiore apice emarginato, tarsis anticis articulo primo dilatato.

Femina abdominis segmento sexto inferiore apice rotundato, tarsis anticis simplicibus.

✕ \* 3. *H. laeviusculus*: Niger, antennis articulis ultimis tribus abrupte maioribus pedibusque nigricantibus, thorace angulis posterioribus subrectis. — Long.  $\frac{1}{2}$  lin.

Mannerh. Brachelytr. 58. 2. — Er. Col. March. I. 389. 3.

*Scaphid. longicorne var. b* Gyll. Ins. Suec. II. 191. 6. — Sahlb. Ins. Fenn. I. 63. 4.

Habitat in Europae pratis.

*H. longicorni* duplo minor, niger, nitidus, parce subtiliterque cinereo-pubescent. Antennae articulis duobus primis crassiusculis, 3 — 7 minutis, subaequalibus, 8 et 9 maioribus, inter se aequalibus, ultimo his coniunctis longitudine aequali et paulo crassiore, apice acuminato, cum iis clavam elongatam minus incrassatam formante, albido-pilosellae, nigro- vel fusco-piceae, basi nonnunquam dilutiores. Palpi nigro-picei. Caput parce subtilissime punctatum. Thorax basi coleopterorum latitudine, antrorsum angustatus, latitudine baseos plus triplo fere brevior, basi subbisinuatus, angulis posterioribus subrectis, acumine summo rotundatis, transversim convexus, parce subtilissime punctatus, margine laterali anguste obsoleteque obscure testaceo-pellucente. Coleoptera thorace sesqui longiora, apice oblique truncata, convexa, crebrius subtiliter punctata. Abdomen crebrius punctulatum, longius pubescens, segmentis singulis piceo-marginatis. Pedes vel nigro-vel fusco-picei, tarsis subtestaceis.

Mas abdominis segmento sexto inferiore apice emarginato, tarsis anticis articulo primo dilatato.

Femina abdominis segmento sexto inferiore apice rotundato, tarsis anticis simplicibus.

Variat immaturus colore corporis ferrugineo.

\* 4. *G. seminulum*: Niger, nitidus, antennis ultimis tribus abrupte maioribus pedibusque testaceis, thorace angulis posterioribus subrectis. — Long.  $\frac{1}{2}$  lin.

Er. Col. Morch. I. 389. 4.

Habitat in Germania: Berolini a me lectus.

Praecedente plerumque paulo minor, niger, nitidus, tenuiter subtiliterque cinereo-pubescent. Antennae articulis duobus primis crassiusculis, 3 — 5 minutis, subaequalibus, 6 et 7 vix crassioribus, 8 et 9 distincte maioribus, ultimo his coniunctis longitudine subaequali et paulo crassiore, apice acuminato, longius pilosellae, totae rufo-testaceae. Palpi rufi. Caput parce subtilissime punctatum. Thorax basi coleopterorum latitudine, antrorsum angustatus, latitudine baseos triplo fere brevior, basi leviter bisinuatus, angulis posterioribus subrectis, convexus, laevis, margine laterali tenui pallido-pellucido. Elytra thorace tertia parte longiora, apice oblique truncata, crebrius subtiliter punctata. Abdomen parcius punctulatum, subtus apice piceo-rufescens. Pedes rufo-testacei, femoribus posticis nigro-piceis.

Mas latet.

Femina abdominis segmento sexto inferiore apice rotundato, tarsis anticis simplicibus.

\* 5. *H. pulicarius*: Niger, nitidus, thorace elytrisque piceis, antennis articulis ultimis tribus abrupte maioribus pedibusque testaceis, thorace angulis posterioribus subrotundatis. — Long.  $\frac{1}{2}$  lin.

Habitat in Germania: Berolini a Dom. Schüppel lectus.



Praecedenti affinis, at paulo maior, niger, nitidus, pube longiore depressa grisea parce vestitus. Antennae longiores, articulis duobus primis maiusculis, 3 — 5 minutis, aequalibus, 6 et 7 his paulo longioribus, at vix crassioribus, 8 et 9 distincte maioribus, subobconicis, inter se aequalibus, ultimo elongato, his coniunctis paulo longiore et paulo crassiore, fusiformi, apice fortiter acuminato, parce pilosellae, totae laete rufo-testaceae. Caput laeve, ore cum palpis rufo-testaceo. Thorax basi coleopterorum latitudine, apicem versus angustatus, latitudine baseos plus duplo brevior, basi vix sinuatus, angulis posterioribus subrotundatis, convexus, laevis, piceus, margine anteriore et posteriore dilutiore, rufescente, laterali latiore, testaceo-pellucido. Elytra thorace tertia parte longiora, oblique truncata, convexa, parce subtilissime punctata, picea. Abdomen parce subtilius punctatum, apice rufo-piceum. Pedes dilute testacei.

Masculatet.

Femina abdominis segmento sexto inferiore apice rotundato, tarsis anticis simplicibus.

Obs. Praecedenti proxime affinis, differe autem videtur statura maiore, antennis longioribus et pedibus dilutioribus, thoracis angulis posterioribus potius rotundatis, elytris parce punctatis.

### **Conurus Steph.**

*Steph. Illustr. V. p. 188.*

*Tachyporus \* Er. Col. March. I. p. 390.*

*Antennae 11-articulatae, filiformes.*

*Palpi maxillares articulo quarto subulato.*

*Ligula biloba, lobis emarginatis.*

*Elytra pectore longiora.*

*Abdomen immarginatum.*

*Tarsi 5-articulati.*

Corpus oblongum, antice obtuse rotundatum, postice sensim attenuatum, convexum, alatum. Caput thoraci insertum, deflexum, convexum, ore obtusiusculo, oculis mediocribus rotundatis, parum prominentibus. Labrum perbreve, integrum. Mandibulae breves, trigonae, intus membrana angusta fimbriatae. Maxillae malis membranaceis, apice barbatis. exteriore interiore vix superante. Palpi maxillares articulo primo minuto, secundo tertioque longitudine subaequalibus, hoc leviter incrassato, quarto parvo subulato. Labium (t. IV. f. 2.) mento transversum, ligula membranea, incisione profundiore biloba, lobis latis, fortiter emarginatis, apice introrsum spinulis nonnullis validioribus munitis, paraglossis ligulae aequalibus eique omnino connatis, nonnisi ciliarum serie indicatis. Palpi labiales articulo primo crassiusculo, secundo hoc duplo breviori, tertio primo longitudine aequali, tenuiore, acuminato. Antennae filiformes, apicem versus saepius subincrassatae. Thorax convexus, basi elytris arcte applicatus, antrorsum magis minusve angustatus, apice truncatus, angulis anterioribus tamen leviter prominentibus, acutiusculis. Elytra pectore paulo longiora, lateribus marginata, angulo apicali exteriore recto. Abdomen immarginatum, apicem versus sensim fortiter attenuatum, vivis sat elongatum. Mesosternum carinatum. Pedes elongati, coxis posterioribus parallalelopipedis, ad femorum basin

scutiformi-elevatis, antici femoribus tibiisque muticis, posteriores femoribus apice tibiisque totis magis minusve spinulosis, spinis tibiarum terminalibus omnibus subtilibus, tarsis omnibus articulis 4 primis decrescentibus, posterioribus valde elongatis, undique glabris, articulo ultimo leviter elongato.

#### O b s e r v a t i o n e s.

1. Corpus totum pube subtili depressa sericante vestitum, abdomine lateribus et apice magis minusve pilosellum.

2. Genus summa affinitate cum *Tachyporo* coniunctum, at notis pluribus, praeter habitum peculiarem distinctum videtur. Differt imprimis ligulae laciniis emarginatis, mesosterno carinato, abdomine immarginato, femoribus posterioribus apice spinulosis, tibiis autem minus spinulosis, tarsis subtus haud pubescentibus, articulo quarto non adeo minuto et corpore toto pube sericante obducto.

3. Differentia sexualis fere ut in *Tachyporo* :

Mas abdominis segmento sexto superiore integro, vel apice subacuminato, inferiore apice exciso, septimo conspicuo.

Femina abdominis segmento sexto superiore apice quadrifido, inferiore apice subacuminato, setis rigidis dense ciliato.

4. Victus sub arborum cortice, in truncis putridis et sub foliis deciduis.

\* 1. *C. littoreus*: *Fuscus, sericeo-pubescent, antennis basi apiceque, thoracis angulis posterioribus, elytrorum macula basali pedibusque rufo-testaceis.* — Long. 2 lin.

*Staphyl. littoreus* Linn. Faun. Suec. n. 852. Syst. Nat. I. II. 685. 18. — Goeze Ent. Beitr. I. 720. 18. — De Vill. Ent. I. 416. 18.

*Tachyporus littoreus* Er. Col. March. I. 390. 1.

*Oxyporus cellaris* Fab. Ent. Syst. I. II. 532. 5. Syst. El. II. 605. 5.

*Staphyl. cellaris* Payk. Faun. Suec. III. 421. 67.

*Tachypor. cellaris* Grav. Micr. 131. 10. Mon. 5. 7. — Latr. Hist. nat. Crust. et Ins. X. 8. 10. — Gyll. Ins. Suec. II. 244. 9. — Sahlb. Ins. Fenn. I. 292. 9. — Zetterst. Ins. Lappon. 55. 8. — Mannerh. Brachelytr. 60. 11. — Boisd. et Lacord. Faun. Ent. Paris. I. 518. 9.

*Staph. pubescens* Payk. Mon. Carab. App. 138. 53—54.

*Staph. tomentosus* Rossi Faun. Etr. Mant. 95. 219. (ed. Hellw. 443. 218.)

*Oxyporus 2-maculatus* Fab. Ent. Syst. I. II. 533. 8. Syst. El. II. 606. 8.

*Staph. maculatus* Fourc. Ent. Par. I. 169. 20.

*le Staph. noir à taches jaunes* Geoffr. Hist. Ins. I. 369. 20.

Habitat in Europa.

Leviter convexus, fuscus, opacus, dense subtiliterque sericeo-pubescent. Antennae capite thoraceque paulo longiores, apicem versus vix incrassatae, articulis 5 — 10 subobconicis, sensim paulo brevioribus et crassioribus, ultimo ovato, abrupte acuminato, testaceae, medio fuscae. Caput fronte leviter convexa, parce subtilissime punctatum, nigrum, ore flavo-testaceo. Thorax coleopteris perparum latior, latitudine paululum brevior, antrorsum angustatus, apice plus dimidio quam basi angustiore, lateribus praecipue pone medium rotundatis, basi utrinque subsinuatus, angulis posterioribus postice prominulis, acutis, modice convexus, dense subtilissime punctatus, angulis posterioribus macula magna rufo-testacea notatis. Elytra thorace sesqui prope longiora, apice oblique truncata, parum convexa, confertissime subtilissimeque punctata, macula magna statim pone basin, aequali spatio a sutura et a margine exteriori distante, rhomboidea rufo-testacea. Abdo-

men parcius subtilissime punctulatum, longius pubescens, minus pilosellum, segmentis singulis margine apicali rufescentibus. Mesosternum modice carinatum. Pedes testacei, coxis anterioribus concoloribus, tibiis parce spinulosis, tarsis posticis valde elongatis.

Mas abdominis segmento sexto subtus apice sat fortiter triangulariter exciso, tarsis anticis articulis tribus primis leviter dilatatis.

Femina abdominis segmento sexto inferiore apice medio triangulariter producto, setis rigidis dense ciliato, superiore quadrifido, laciniis intermediis profunde divisis, tarsis anticis simplicibus.

Variat elytrorum macula effusa, discum totum occupante, margine tenui, angulo scutellari latius, apice exteriori late fuscis. Thoracis latera ante maculam testaceam saepe fusco-rufa.

Obs. 1. *Oxyporum himaculatum* Fabr. nonnisi secundum eius descriptionem huic speciei allegavi, specimen enim descriptum in collectione citata desideratur.

2. Exemplum nomine *Tachypori trinotati* a Dahlio missum, in Carinthia lectum, communicavit Dom. Schüppel.

\* 2. *C. grossus*: *Fusco-niger, sericeo-pubescens, elytris thorace paulo longioribus, pedibus rubris, antennis longis, articulis duobus primis rubris.* — Long. 3 lin.

Habitat in Madagascar, Dom. Goudot.

Reliquis multo maior, sat convexus, fusco-niger, subnitidus, pube fusco-nigra vestitus. Antennae elongatae, coleopterorum apicem fere attingentes, apicem versus haud incrassatae, articulis a tertio inde omnibus elongatis, ultimo apice sensim acuminato, nigro-fuscae, articulis duobus primis rubris. Palpi rubri. Caput parcius subtiliter punctatum, fronte leviter convexa, nigrum, infra oculos rufo-piceum. Thorax coleopteris paulo latior, latitudine baseos paulo brevior, antrorsum fortiter angustatus, apice basi triplo fere brevior, lateribus aequaliter rotundatus, margine basali recto, angulis posterioribus fortiter prominentibus, subrotundatis, fortiter convexus, sat dense subtiliter punctatus. Elytra thorace paululum longiora, apice oblique truncata, transversim convexa, confertissime subtiliter punctata. Abdomen minus confertim subtiliter punctatum, parcius pubescens, fortiter nigro-pilosellum, summo apice nonnunquam rufescente. Pectus fortiter carinatum. Pedes rubri, coxis anticis concoloribus, femoribus posterioribus apice fortius, tibiis interne parcius breviterque spinulosis, tarsis posticis sat elongatis.

Mas abdominis segmento quinto subtus apice leviter emarginato, sexto inferiore profunde inciso, superiore sat acute acuminato, tarsis anticis articulis tribus primis fortiter dilatatis, antennis longioribus, articulis ultimis magis elongatis.

Femina abdominis segmento sexto inferiore apice rotundato, setis fulvis dense ciliato, superiore quadrifido, laciniis intermediis sat profunde divisis, tarsis anticis articulis tribus primis leviter dilatatis, antennis articulis ultimis minus fortiter elongatis.

Variat corpore subtus toto rufo-brunneo vel castaneo.



- \* 3. *C. australis*: *Niger, fusco-pubescentis, antennis basi apiceque testaceis, pedibus rubris, elytris thorace tertia parte longioribus, apice oblique truncatis.* — Long. 2 lin.

Habitat in Terra Van-Diemenii, Dom. Schayer.

Habitu omnino *C. pubescentis*, at paulo maior, praecipue latior, *C. littoreo* aequalis, hoc autem multo convexior, niger, subnitidus, pube brevi, depressa, fusca, subsericante vestitus. Antennae capitis thoracisque prope longitudine, articulis 3 et 4 elongatis, teretibus, 5 — 10 subobconicis, sensim paulo brevioribus et crassioribus, ultimo ovato, apice acuminato, articulis duobus primis pallide testaceis, tertio testaceo, apice fusco, 5 — 10 fuscis, ultimo testaceo. Caput convexiusculum, laevigatum, parcius pubescens, nigrum, nitidum, ore rufo-testaceo. Thorax coleopteris paulo latior, latitudine paulo brevior, antrorsum angustatus, apice duplo quam basi angustior, lateribus rotundatus, margine basali recto, angulis posterioribus leviter retrorsum prominentibus, subrectis, convexus, confertim subtilissime punctatus, margine basali piceo. Elytra thorace tertia parte longiora, apice oblique truncata, convexa, confertissime subtiliter punctata. Abdomen confertim subtiliter punctulatum, segmentis singulis piceo-marginatis, lateribus et apice fortius pilosellum. Mesosternum acute carinatum. Pedes rubri seu ferruginei vel testacei, coxis anticis concoloribus, tarsis posticis sat elongatis.

Mas abdominis segmento sexto inferiore apice triangulariter exciso, superiore apice rotundato, tarsis anticis articulis tribus primis modice dilatatis.

Femina abdominis segmento sexto inferiore apice obtuse acuminato, setis rigidis ferrugineis dense ciliato, superiore quadrifido, laciniis intermediis ad segmenti medium usque divisis.

Obs. *C. pubescenti* proximus, differt autem statura paulo robustiore, antennis brevioribus et debilioribus magis fuscis, thoracis angulis posterioribus magis prominentibus, elytris apice oblique truncatis, abdomine fortius pilosello.

- \* 4. *C. pubescens*: *Fusco-niger, griseo-pubescentis, elytris thorace longioribus, antennis pedibusque rufo-testaceis.* — Long. 2 lin.

*Tachyporus pubescens* Grav. *Micr.* 130. 9. *Mon.* 5. 8. — *Latr. Hist. nat. Crust. et Ins.* X. 8. 9. — *Gyll. Ins. Suec.* II. 243. 8. — *Sahlb. Ins. Fenn.* I. 292. 8. — *Mannerh. Brachelytr.* 60. 10. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 518. 10. — *Er. Col. March.* I. 390. 2.

*Staph. pubescens* var.  $\beta$  Payk. *Mon. Carab. App.* 138. 53 — 54.

*Staph. cellaris* var.  $\beta$  Payk. *Faun. Suec.* III. 421. 67.

*Staph. tomentosus* var.  $\beta$  Rossi *Faun. Etr. Mant.* 97. 218. (*Ed. Hellw. I.* 443. 218.)

*Var. a: Fuscus, elytris brunneis.*

*Tachyporus sericeus* Boisd. et Lacord. *Faun. Ent. Paris.* I. 519. 11.

*Var. b: Totus pallide testaceus, immaturus.*

*Oxyporus testaceus* Fab. *Ent. Syst.* I. II. 535. 18. *Syst. El.* II. 607. 18. — *Panz. Ent. Germ.* 361. 14.

Habitat in Europae silvis, in truncis putridis. In China meridionali etiam lectus a Prof. Meyen.

Pube brevi depressa griseo-sericea dense vestitus. Antennae capitis

thoracisque longitudine, apicem versus sensim subincrassatae, articulis 3—5 elongatis, teretibus, 6—10 subobconicis, sensim paulo crassioribus brevioribusque, ultimo apice abrupte acuminato, testaceae, medio fuscae. Os cum palpis testaceum. Caput modice convexum, laevigatum, parcius subtiliter pubescens, nigrum, nitidum. Thorax coleopteris latior, latitudine parum brevior, antrorsum fortiter angustatus, apice duplo prope angustior quam basi, lateribus aequaliter rotundatus, basi utrinque subsinuatus, angulis posterioribus postice vix prominulis, subrectis, fortiter convexus, omnium subtilissime confertim punctatus, piceo-niger, pube derasa nitidus, margine summo basali rufescente. Elytra thorace tertia fere parte longiora, convexa, confertissime subtiliter punctata, apice truncata. Abdomen fuscum, apice pilosello segmentorumque singulorum margine apicali late castaneis. Pectus nigrum, nitidum, mesosterno acute carinato. Pedes testacei, coxis anticis concoloribus, posteriores femoribus apice tibiisque parce spinulosis, tarsis posticis sat elongatis.

Mas abdominis segmento quinto subtus apice leviter emarginato, sexto inferiore leviter triangulariter exciso, superiore apice rotundato, tarsis anticis articulis tribus primis modice dilatatis, antennis paulo longioribus.

Femina abdominis segmento sexto inferiore apice leviter triangulariter producto, setis rigidis ferrugineis dense ciliato, superiore quadrifido, lacinis intermediis ad medium segmenti usque divisis, tarsis anticis articulis tribus primis leviter dilatatis.

Variat colore corporis aut nigro, elytris aut concoloribus aut castaneis, aut fusco, elytris dilutius rufo-brunneis, aut toto ferrugineo, aut minus maturus testaceo.

Obs. Specimina nigra elytris concoloribus nomine *Tachypori alpestris* ex albis Carinthiae, et *T. tristis* e Styria, specimen immaturum, corpore testaceo nomine *T. rufescentis* ex Austria a Dahlio sibi missa, dedit Dom. Schüppel. *Oxyporus testaceus* Fab. secundum ipsum specimen ab auctore descriptum hoc loco allegatus.

- \* 5. *C. crassus*: *Fusco-niger, tenuiter griseo-pubescens, antennis testaceis, thoracis margine, elytrorum basi, ano pedibusque rufescentibus, elytris thorace parum longioribus.* — — Long.  $2\frac{1}{2}$  lin.

*Tachyporus crassus* Grav. Micr. 190. 1. Mon. 6. 9.

Habitat in America septentrionali.

Praecedente triplo maior, multo latior, fortiter convexus, fusco-niger, nitidus. Antennae capitis thoracisque prope longitudine, apicem versus parum incrassatae, testaceae, articulis 3 et 4 elongatis, teretibus, 5—10 subobconicis, sensim paulo brevioribus et crassioribus, basi fusciscentibus, ultimo oblongo-ovato, toto flavo. Caput parum convexum, omnium subtilissime punctatum, parcius pubescens, nigrum, ore cum palpis rufo-testaceo. Thorax coleopterorum basi parum latior, latitudine paulo brevior, antrorsum angustatus, apice plus duplo quam basi angustior, lateribus fortius aequaliter rotundatis, basi utrinque subsinuatus, angulis posterioribus retrorsum parum prominulis, rotundatis, valde convexus, omnium subtilissime punctulatus, pube depressa griseo-fusca tenuiter vestitus, margine laterali basaque sanguineo. Elytra thorace paululum longiora, apice oblique truncata,

convexa, confertim minus subtiliter punctata, nigro-fusca, tenuiter pubescentia, margine basali sanguineo, densius pubescente. Abdomen breviusculum, subtilissime punctulatum, pube depressa griseo-fulva sericante densius vestitum, lateribus et apice fortius pilosellum, segmentorum singulorum margine apicali rufo-piceo, ano rufo-testaceo. Mesosternum fortiter carinatum. Pedes rufo-testacei, coxis anticis concoloribus, postici femoribus apice fortiter, tibiis interne parce breviterque spinulosis, tarsis posticis modice elongatis.

Mas mihi invisus.

Femina abdominis segmento sexto inferiore apice acuminato, setis rigidis fulvis dense ciliato, superiore quadrifido, laciniis intermediis ultra segmenti medium divisis, tarsis anticis articulis tribus primis leviter dilatatis.

6. *C. tumidus*: *Fuscus, griseo-pubescentis, thoracis margine summo elytrorumque basi rufescentibus, pedibus rufo-testaceis, antennis nigris, articulis 4 primis ultimoque testaceis.* — Long. 2 lin.

Habitat prope Puerto Cabello Columbiae, Dom. Moritz.

Statura omnino praecedentis, at paulo minor, T. littoreo paulo latior et multo convexior, fuscus, nitidus, subtiliter punctulatus, pube minus subtili depressa fulvo-grisea densius vestitus. Antennae capitis thoracisque prope longitudine, apicem versus sensim leviter incrassatae, articulis 4 primis testaceis, tertio quartoque elongatis, teretibus, 5—10 obconicis, sensim paulo brevioribus et crassioribus, nigris, ultimo ovato, apice acuminato, flavo, basi fusciscente. Palpi dilute testacei. Caput leviter convexum, parce subtilissime punctulatum, ore et subtus totum rufo-testaceum. Thorax coleopteris paulo latior, antrorsum angustatus, apice quam basi plus duplo angustior, latitudine baseos paulo brevior, lateribus aequaliter rotundatus, basi truncatus, angulis posterioribus retrorsum haud prominulis, obtusis, subrotundatis, fortiter convexus, parce subtilissime punctatus, margine laterali et basali rufescente. Elytra thorace paulo longiora, convexa, apice vix oblique truncata, parcius subtilissime punctata, tascia basali, parum distincta, externe abbreviata rufescente. Abdomen crebrius subtiliter punctatum, densius pubescens, segmentis singulis piceo-marginatis, apice testaceum et pilosellum. Mesosternum subtiliter at sat acute carinatum. Pedes rufo-testacei, posteriores femoribus apice tibiisque fortius spinulosis, tarsis posticis fortiter elongatis.

Mas abdominis segmento sexto inferiore apice fortiter triangulariter exciso, tarsis anticis articulis tribus primis valde dilatatis.

Femina latet.

- \* 7. *C. bipustulatus*: *Niger, subtiliter griseo-pubescentis, antennis basi apiceque pedibusque testaceis, elytris thorace paulo longioribus, macula basali rhombea rufa.* — Long. 2 lin.

*Tachyporus bipustulatus* Grav. Micr. 132. 11. — Latr. Hist. nat. Crust. et Ins. X. 9. 11. — Gyll. Ins. Suec. II. 245. 10.

*Tachyporus bimaculatus* Grav. Mon. 5. 6.

Habitat in Hungaria, Dahl, in Styria et in Suecia, Dom. Schüppel.

Statura omnino *C. pubescentis*, at paulo maior, convexus, niger, ni-



nitidulus, pube depressa subtili fulvescenti-grisea tenuiter vestitus. Antennae capite thoraceque parum breviores, apicem versus leviter sensim incrassatae, testaceae, articulis 3 et 4 elongatis, teretibus, 5 — 10 subobconicis, sensim paulo brevioribus et crassioribus, fuscis, ultimo oblongo-ovato, flavo. Caput modice convexum, parce omnium subtilissime punctatum, ore rufo-testaceo. Thorax coleopteris paulo latior, antrorsum angustatus, apice duplo quam basi angustior, latitudine baseos paulo brevior, lateribus sat rotundatus, margine basali medio late recto, angulis posterioribus retrorsum modice prominulis, subrectis, convexus, minus confertim subtilissime punctatus. Elytra thorace paulo longiora, apice parum oblique truncata, convexa, confertim subtiliter punctata, macula rhombea subtransversa testaceo-rufa ad basin, basin ipsam haud attingente, a margine exteriori paulo magis quam a sutura remota, sat distincta et terminata. Abdomen sat dense subtiliter punctatum, segmentis ultimis rufo-marginatis, apice pilosellum. Mesosternum subtiliter carinatum. Pedes rufi, tibiis intermediis parce, posticis vix spinulosis, tarsis posticis tibia parum longioribus.

Mas abdominis segmento sexto inferiore apice fortiter triangulariter exciso, superiore apice rotundato, tarsis anticis articulis tribus primis modice dilatatis.

Femina abdominis segmento sexto inferiore apice rotundato, setis rigidis fulvis dense ciliato, superiore quadrifido, laciniis intermediis profunde divisis, tarsis anticis articulis tribus primis leviter dilatatis.

Obs. Nomine Tachypori interrupti a Dahlio et Zieglero missus.

\* 8. *C. binotatus*: *Fuscus, fulvescenti-pubescent, antennis pedibusque rufo-testaceis, elytris thorace sesqui longioribus, macula magna basali rufa.* — Long.  $1\frac{2}{3}$  lin.

*Tachyporus binotatus* Grav. *Micr.* 132. 12. *Mon.* 5. 5. — *Latr. Hist. nat. Crust. et Ins.* X. 9. 12. — *Er. Col. March.* I. 391. 3.

Habitat in Germania.

C. pubescente paulo minor, fuscus, nitidulus, pube subtiliore depressa fulvescente vestitus. Antennae capite thoraceque paulo breviores, apicem versus leviter incrassatae, articulis 5—10 sensim paulo brevioribus et crassioribus, obconicis, ultimo ovato, apice abrupte acuminato, pilosellae, rufo-testaceae, articulo ultimo flavo. Caput leviter convexum, parce subtilissime punctatum, nigrum, nitidum, ore testaceo. Thorax coleopteris parum latior, antrorsum angustatus, apice vix duplo quam basi angustior, latitudine baseos dimidio prope brevior, lateribus sat rotundatus, basi utrinque subsinuatus, angulis posterioribus retrorsum sat prominulis, rotundatis, modice convexus, densius subtiliter punctatus, margine basali rufescente. Elytra thorace sesqui longiora, apice truncata, leviter convexa, confertim subtiliter punctata, macula magna rotundata rufo-testacea, a basi ad elytri medium usque descendente, margini exteriori et suturae aequaliter approximata, minus distincte terminata, margine apicali summo rufescente. Abdomen densius subtiliter punctatum, segmentis singulis margine rufis, apice testaceum et parcius pilosellum. Mesosternum subtiliter carinatum. Pedes rufo-testacei, tibiis intermediis longius, posticis vix spinulosis, tarsis posticis tibia paulo longioribus.

Mas abdominis segmento sexto inferiore apice sat fortiter triangulariter exciso, superiore apice obtuse acuminato, tarsis anticis articulis tribus primis leviter dilatatis.

Femina abdominis segmento sexto inferiore apice rotundato, pilis brevissimis tenuiter ciliato, superiore quadrifido, laciniis intermediis sat profunde divis, tarsis anticis levissime dilatatis.

- \* 9. *C. basalis*: *Fuscus, nitidus, griseo-pubescens, elytris macula basali transversali summoque margine apicali rufescentibus, antennis elongatis pedibusque testaceis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Carolina meridionali, Dom. Zimmermann.

*C. binotato* paulo minor, convexus, fuscus, subnitidus, pube depressa fulvescenti-grisea sericante densius vestitus. Antennae sat elongatae, graciles, apice leviter incrassatae, articulis 3 — 5 elongatis, teretibus, 6 — 10 elongato-obconicis, sensim paulo crassioribus at vix brevioribus, ultimo oblongo-ovato, apice abrupte acuminato, obscure testaceae, basi flavescens, articulo ultimo dilute flavo. Caput convexus, vix punctulatum, nigrum, nitidum, ore late testaceo. Thorax coleopteris lator, antrorsum angustatus, apice duplo fere quam basi angustior, latitudine baseos paulo brevior, lateribus aequaliter rotundatus, margine basali medio late recto, angulis posterioribus modice retrorsum prominentibus, acutiusculis, fortiter convexus, confertissime subtiliter punctatus, nigro-fuscus, margine basali et apicali rufescente. Elytra thorace longiora, apice truncata, convexa, confertissime subtiliter punctata, fusca, macula magna in ipsa basi, a margine exteriori magis quam a sutura distante, rufa, margine summo apicali rufescente. Abdomen subtilissime punctulatum, segmentis singulis apice piceis, posterioribus subtestaceis, densius pubescens, lateribus et apice fortiter nigro-pilosum. Mesosternum acute carinatum. Pedes testacei, coxis anticis concoloribus, femoribus posterioribus apice parce, tibiis vix spinulosis, tarsis posticis sat elongatis, anticis articulis tribus primis in utroque sexu leviter dilatatis.

Mas abdominis segmento sexto inferiore apice triangulariter exciso, antennis capite thoraceque longioribus.

Femina abdominis segmento sexto superiore quadrifido, laciniis intermediis profunde divis.

Variat corpore dilutiore, fusco-rufescente.

10. *C. interruptus*: *Fuscus, griseo-pubescens, coleopteris fascia basali, medio interrupta pedibusque rufis, antennis testaceis, medio fuscis.* — Long.  $1\frac{1}{2}$  lin.

Hacitat in valle Araguensi Columbiae, Dom. Moritz.

Praecedenti affinis, at paulo angustior, convexus, fuscus, nitidulus, subtiliter punctulatus, pube depressa griseo-fulvescente subsericante tenuiter vestitus. Antennae capitis thoracisque longitudine, apice sensim incrassatae, articulis 4 primis testaceis, tertio quartoque teretibus, modice elongatis, 5 — 10 subobconicis, sensim crassioribus at vix brevioribus, ultimo crassiusculo, ovato, apice acuminato, 5 — 9 fuscis, ultimis duobus laete testaceo-flavis. Caput modice convexus, vix punctulatum, ore cum palpis testaceo.

Thorax coleopteris parum latior, antrorsum modice angustatus, apice dimidio fere quam basi angustior, latitudine baseos parum brevior, lateribus leviter rotundatus, basi truncatus, utrinque subsinuatus, angulis posterioribus retrorsum vix prominulis, rectis, valde convexus, parcius omnium subtilissime punctulatus, margine basali et laterali obsolete rufescentibus. Elytra thorace parum longiora, apice oblique truncata, leviter convexa, densius subtiliter punctata, fascia basali ad suturam abbreviata rufo-testacea, margine summo apicali rufescente. Abdomen subtilissime punctulatum, apice testaceum, densius pubescens, lateribus et apice fortiter nigro-pilosum. Mesosternum leviter carinatum. Pedes testacei, coxis anticis concoloribus, posteriores femoribus apice tibiisque parce spinulosis, tarsis posticis modice elongatis.

Mas segmento abdominis sexto inferiore apice leviter triangulariter exciso, tarsis anticis articulis tribus primis fortius dilatatis.

Femina latet.

Obs. Affinis praecedenti, statura autem paulo angustiore, antennis articulis 5 — 9 distincte fuscis, coleopterorum fascia ad marginem anteriorem continuata etc. satis distinctus.

\* 11. *C. cinctulus*: *Niger, subnitidus, tenuiter pubescens, antennis basi apiceque pedibusque testaceis, coleopteris fascia basali integra rufa, sutura abdomineque densius pubescentibus.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Carolina meridionali, Dom. Zimmermann.

Statura et summa affinitas *C. basalis*. Antennae capitis thoracisque longitudine, apicem versus sensim leviter incrassatae, articulis 2 — 4 subelongatis, teretibus, 5 — 10 subobconicis, sensim paulo brevioribus et crassioribus, penultimis transversis, ultimo breviter ovato, apice subacuminato, articulis tribus primis flavis, sequentibus fuscis, ultimo testaceo. Caput modice convexum, parce subtiliter punctulatum, parce griseo-pubescens, nigrum, nitidum, ore testaceo. Thorax coleopteris paulo latior, antrorsum leviter angustatus, apice dimidio fere quam basi angustior, latitudine baseos distincte brevior, lateribus rotundatus, margine basali medio late recto, angulis posterioribus retrorsum leviter prominentibus, acutiusculis, fortiter convexus, dense subtiliter punctatus, fusco-niger, subnitidus, margine basali et laterali tenui rufescente, pube depressa flavescenti-grisea tenuiter vestitus. Scutellum fuscum. Elytra thorace paulo longiora, apice oblique truncata, leviter convexa, confertissime subtiliter punctata, nigro-fusca, subnitida, fascia integra in ipsa basi rufa, tenuiter fusco-pubescentia, fascia basali et sutura pube densa depressa griseo-fulvescente sericante obductis. Abdomen confertissime subtiliter punctatum, nigrum, segmentis singulis margine apicali piceo, quinto testaceo, pube depressa griseo-fulvescente dense vestitum, apice nigro-pilosum. Pectus nigrum. Pedes rufo-testacei, coxis anticis concoloribus, posteriores femoribus apice tibiisque parcius spinulosis, tarsis posticis minus elongatis, tibiae longitudine aequalibus.

Mas abdominis segmento sexto inferiore apice triangulariter exciso, tarsis anticis articulis tribus primis leviter dilatatis.

Femina latet.



Obs. Differt a praecedentibus coleopterorum fascia integra, antennarum colore, abdomine lateribus haud piloso etc.

- \* 12. *C. varius*: *Niger, sericeo-pubescent, thorace pedibusque testaceis, elytris nigro testaceoque variegatis.* — Long.  $1\frac{1}{2}$  lin.

Habitat prope Caracas Columbiae, Dom. Moritz.

C. fusculo paulo angustior, totus pube minus subtili, depressa, griseo-fulvescente minus dense vestitus. Antennae capite thoraceque parum breviores, apicem versus leviter incrassatae, articulis 2 — 4 elongatis, subteretibus, 5 — 10 brevioribus, subobconicis, sensim paulo crassioribus, ultimo maiusculo, subovato, apice acuminato, rufo-testaceae, basi flavescentes. Caput parum convexum, vix punctulatum, nigro-piceum, nitidum, ore testaceo. Thorax basi coleopteris paulo latior, antrorsum angustatus, apice quam basi duplo angustior, latitudine baseos distincte brevior, lateribus parum rotundatus, basi utrinque subsinuatus, angulis posterioribus retrorsum haud prominulis, obtusiusculis, modice convexus, parce subtilissime punctulatus, testaceus, nitidus. Scutellum nigrum. Elytra thorace tertia parte longiora, apice truncata, convexa, subtiliter sat crebre punctata, nigra, maculis oblongis confluentibus testaceis variegata. Abdomen piceum, lateribus et apice fortiter nigro-pilosum. Pectus nigrum. Pedes testacei.

Mas latet.

Femina abdominis segmento sexto inferiore apice leviter producto, setis rigidis densius ciliato, superiore apice quadrifido, tarsis anticis leviter dilatatis.

13. *C. venustulus*: *Tenuiter pubescens, piceus, capite, thorace pectoreque rufis, elytris nigris, his macula magna basali pedibusque flavis.* — Long.  $1\frac{1}{4}$  lin.

Habitat prope Puerto Cabello Columbiae, Dom. Moritz.

Statura omnino C. fusculi. Corpus totum pube depressa fulvescenti-grisea parce vestitum. Antennae capite thoraceque paulo breviores, apicem versus sensim leviter incrassatae, articulis 2 — 4 elongatis, teretibus, 5 — 10 sensim paulo brevioribus et crassioribus, 5 — 7 subobconicis, 8 — 10 transversis, ultimo breviter ovato, apice obtuse acuminato, testaceae, articulis 7 — 10 fuscis. Palpi testacei. Caput parum convexum, parce disperse subtiliter punctatum, nitidum, testaceo-rufum. Thorax coleopteris perparum latior, antrorsum angustatus, apice dimidio fere quam basi angustior, latitudine baseos parum brevior, lateribus leviter rotundatus, basi truncatus, angulis posterioribus retrorsum paululum prominulis, acutiusculis, modice convexus, parcius subtiliter punctatus, testaceo-rufus, nitidus. Scutellum testaceum. Elytra thorace longiora, apice vix oblique truncata, modice convexa, dense minus subtiliter punctata, nigerrima, nitida, macula magna densius pubescente in ipsa basi flavo-testacea. Abdomen piceum, densius pubescens, lateribus et apice fortiter nigro-pilosum. Pectus rufum. Pedes flavo-testacei, coxis anticis concoloribus, postici femoribus apice tibusque parce longius spinulosi, tarsis posticis satis elongatis.

Mas latet.

Femina abdominis segmento sexto inferiore apice producto et setis rigidis parciorebus ciliato, superiore quadrifido, laciniis intermediis profunde divis, tarsis anticis articulis tribus primis parum dilatatis.

- \* 14. *C. obsoletus*: *Fuscus, flavescenti-pubescens, thorace elytrisque rufo-testaceis, his thorace sesqui longioribus, antennarum basi pedibusque testaceis.* — Long.  $1\frac{1}{4}$  lin.

Habitat prope Caracas Columbiae, Dom. Moritz.

Statura omnino *T. fusculi*. Corpus totum pube tenui depressa griseo-flavescente subsericante minus dense vestitum. Antennae capite thoraceque paulo breviores, apicem versus vix incrassatae, articulis 2—4 modice elongatis, teretibus, 5—10 brevioribus, subobconicis, sensim paulo brevioribus at vix crassioribus, ultimo praecedentibus ambobus paulo longiore, apice acuminato, fuscae, articulis 5 primis pallide testaceis. Palpi flavi. Caput parum convexum, laeve, testaceo-rufum vel rufo brunneum, nitidum, ore rufescente. Thorax basi coleopterorum latitudine, antrorsum angustatus, apice duplo prope quam basi angustior, latitudine baseos brevior, lateribus parum rotundatus, apice truncatus, angulis posterioribus retrorsum haud prominulis, subrectis, modice transversim convexus, vix punctulatus, rufo-testaceus, nitidus. Scutellum fuscum. Elytra thorace sesqui longiora, leviter convexa, subtilissime punctulata, rufo-testacea, lateribus medio infuscat. Abdomen fuscum, lateribus et apice fortiter nigro-pilosum. Pectus fuscum. Pedes flavo-testacei, postici femoribus apice et tibiis parce spinulosis, tarsis posticis sat elongatis.

Sexus differentia in specime descripto haud perspicua.

- \* 15. *C. fumatus*: *Fuscus, nitidus, fulvescenti-pubescens, antennarum basi pedibusque testaceis, thorace limbo rufescente, angulis posterioribus haud prominulis, rotundatis, elytris thorace longioribus.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Terra Van-Diemenii, Dom. Schayer.

Statura intermedius quasi inter *C. pubescentem* et *fuscum*, utrique affinis, ab utroque autem pube minus subtili et thoracis angulis posterioribus haud prominulis rotundatis satis distinctus. Nigro-fuscus, nitidus, pube depressa minus subtili fulvescente sericante leviter vestitus. Antennae longiores, articulis 4—10 subobconicis, sensim paulo brevioribus at vix crassioribus, fuscae, articulis 4 primis testaceis; (ultimus specimini unico deest). Caput fortiter convexum, laevissimum, ore testaceo. Thorax coleopteris vix latior, antrorsum sat angustatus, apice duplo quam basi angustior, latitudine baseos dimidio brevior, lateribus fortiter rotundatus, basi truncatus, utrinque subsinuatus, angulis posterioribus retrorsum haud prominulis, rotundatis, modice transversim convexus, parce subtiliter punctatus, limbo omnipiceo-rufo, disco indeterminate fusco-piceo. Elytra thorace tertia parte longiora, apice oblique truncata, leviter convexa, densius subtiliter punctata, fusca, margine summo apicali rufescente. Abdomen parce subtiliterque punctatum, parcius pubescens, lateribus et apice longius nigro-pilosum, apice et segmentorum singulorum margine posteriore rufo-piceo. Mesosternum subtiliter carinatum. Pedes rufo-testacei, tibiis posterioribus vix spinulosis, tarsis posticis minus elongatis, tibia haud longioribus.

Mas abdominis segmento sexto inferiore apice sat fortiter triangulariter exciso, superiore apice rotundato; tarsis anticis articulis tribus primis leviter dilatatis.

\* 16. *C. fuscus*: *Fusco-niger, griseo-pubescentis, antennis basi testaceis, apice flavis, pedibus rufis, elytris thorace longioribus.* — Long.  $1\frac{1}{4}$  lin.

*Tachyporus fuscus* Er. Col. March. I. 591. 4.

*Tachyporus pedicularius* Boisd. et Lacord. Faun. Ent. Paris. I. 519. 12.

Habitat, in Germania.

Habitus et summa similitudo *C. pubescentis*, at triplo minor, convexus, fusco-niger, pube subtili depressa grisea sericante vestitus. Antennae capitis thoracisque fere longitudine, articulis 3 et 4 elongatis, teretibus, 5—10 subobconicis, sensim brevioribus et paulo crassioribus, primis eorum sat elongatis, ultimo praecedente parum maiore, fuscae, articulis tribus primis testaceis, ultimis duobus flavis. Caput leviter convexum, laevigatum, nigrum, nitidum, ore testaceo. Thorax coleopteris paulo latior, antrorsum angustatus, apice duplo quam basi angustior, latitudine baseos paulo brevior, lateribus modice rotundatus, basi utrinque subsinuatus, angulis posterioribus retrorsum paululum prominulis, acutis, convexus, confertim subtilissime punctatus, margine basali summo rufescente. Elytra thorace paululum longiora, apice truncata, leviter convexa, confertissime subtiliter punctata, margine apicali rufescente. Abdomen confertim subtiliter punctatum, segmentis singulis margine postico rufescente, apice pilosellum. Mesosternum acute carinatum. Pedes rufo-ferruginei, coxis anticis concoloribus, tibiis intermediis parce, posticis vix spinulosis, tarsis posticis modice elongatis, tibia parum longioribus.

Mas abdominis segmento sexto inferiore apice sat fortiter triangulariter exciso, superiore subacuminato, tarsis anticis articulis tribus primis modice dilatatis.

Femina abdominis segmento sexto inferiore apice rotundato, setis rigidis fulvis dense ciliato, superiore apice quadrifido, laciniis intermediis brevissimis divisis, tarsis anticis articulis tribus primis leviter dilatatis.

Variat minus maturus corpore ferrugineo vel rufo-testaceo.

Obs. *C. pediculario* similis, statura autem sesqui maiore, antennis medio fuscis, thoracis angulis posterioribus acutis et elytris longioribus facile distinguendus.

\* 17. *C. lividus*: *Rufo-testaceus, fulvescenti-pubescentis, pectore piceo, elytris thorace paulo longioribus.* — Long. 1 lin.

Habitat Lutetiae, Dom. Dr. Aubé, in Sardinia, Dom. Prof. Gené.

Statura et summa affinitas *C. pedicularii*, rufo-testaceus, pube depressa minus subtili fulvescente densius vestitus. Antennae longiores, apice sensim subincrassatae, articulis teretibus, penultimis tribus quatuorve subobconicis, ultimo praecedente sesqui maiore, apice abrupte acuminato, flavo-testaceae. Caput leviter convexum, laevissimum. Thorax coleopteris paululum latior, antrorsum angustatus, tertia parte quam basi angustior, latitudine baseos paulo brevior, lateribus rotundatus, margine basali medio recto, angulis posterioribus leviter retrorsum prominentibus, acutiusculis, convexus, omnium subtilissime punctatus. Elytra thorace paulo longiora, apice oblique truncata,



convexa, confertim subtilius punctata. Abdomen subtilissime punctatum, segmentis duobus primis summa basi nigricantibus, apice nigro-pilosellum. Pectus piceum. Mesosternum acute carinatum. Pedes testacei, tibus posterioribus haud spinulosis, tarsis posticis tibia paulo brevioribus.

Mas abdominis segmento sexto inferiore apice fortiter triangulariter exciso, superiore apice obtuse rotundato, tarsis anticis leviter dilatatis, antennis capite thoraceque longioribus.

Femina abdominis segmento sexto inferiore apice leviter triangulariter producto et dense fulvo-ciliato, superiore apice quadrifido, tarsis anticis simplicibus, antennis capitis thoracisque longitudine.

Obs. A C. pediculario differt antennis elytrisque longioribus, thoracis angulis posterioribus acutioribus et mesosterni carina fortiore, acuta etc., a C. fusculo statura minore elytris brevioribus, ab utraque pube minus subtili. Ceterum speciminibus eorum immaturis similis, at color testaceus huic speciei normalis.

\* 18. *C. pedicularius*: Nigro-piceus, nitidus, subtiliter griseo-pubescent, antennis pedibusque testaceis, elytris thorace subbrevioribus. — Long. 1 lin.

*Tachyporus pedicularius* Grav. Micr. 133. 14. Mon. 4. 2. — Latr. Hist. nat. Crust. et Ins. X. 10. 14. — Gyll. Ins. Suec. II. 246. 11. — Sahlb. Ins. Fenn. I. 293. 11. — Mannerh. Bruchelytr. 60. 13. — Runde Brachel. Hal. 25. 7. — Er. Col. March. I. 392. 5.

Var.: Totus testaceus, immaturus.

*Tachypor. truncatellus* Grav. Mon. 5. 3.

Habitat in Europa.

Minutus, convexus, nigro-piceus, nitidulus, pube subtili depressa grisea sericante densius vestitus. Antennae capitis thoracisque prope longitudine, apicem versus subincrassatae, articulis 5 — 10 sensim paulo brevioribus et crassioribus, subobconicis, ultimo praecedente sesqui maiore, apice abrupte acuminato, totae testaceae. Caput convexum, laeve, nigrum, ore testaceo. Thorax coleopteris latior, antrorsum angustatus, apice duplo fere quam basi angustior, lateribus sat rotundatus, margine basali medio recto, angulis posterioribus retrorsum fortiter prominulis, rotundatis, valde convexus, densius omnium subtilissime punctatus, margine basali summo rufescente. Elytra thoracis longitudine vel paulo breviora, apice oblique truncata, convexa, confertim subtiliter punctata, margine apicali rufescente. Abdomen dense subtilissime punctulatum, segmentis singulis margine postico rufescente, apice parcius nigro-pilosum. Mesosternum leviter carinatum. Pedes rufo-testacei, tibus posterioribus muticis, tarsis posticis tibia paulo brevioribus.

Mas abdominis segmento sexto inferiore apice fortiter triangulariter exciso, superiore apice obtuse rotundato.

Femina abdominis segmento sexto inferiore apice leviter triangulariter producto et subtilius fulvo-ciliato, superiore apice quadrifido, tarsis anticis haud dilatatis.

Variat minus maturus corpore ferrugineo vel testaceo.

\* 19. *C. bipunctatus*: Niger, nitidus, tenuiter sericeo-pubescent, elytrorum macula basali arcuata pedibusque rubris, antennis testaceis, medio fuscis. — Long. 1 lin.

*Tachyporus bipunctatus* Grav. *Micr.* 133. 13. *Mon.* 5. 4. — *Latr. Hist. nat. Crust. et Ins.* X. 9. 13. — *Gyll. Ins. Suec.* III. 702. 10 — 11. — *Sahlb. Ins. Fenn.* I. 293. 10. — *Mannerh. Brachelytr.* 60. 12. — *Er. Col. March.* I. 392. 6.

Habitat in Germania, sub cortice Salicis.

C. pediculario statura et magnitudine aequalis, niger, nitidus, pube subdepressa grisea tenuiter vestitus. Antennae capite thoraceque breviores, apicem versus sensim leviterque incrassatae, articulis 3 et 4 leviter elongatis, subobconicis, 5—10 transversis, sensim paulo crassioribus, ultimo praecedente paulo longiore, apice subacuminato, testaceae, medio fuscae. Caput parce subtiliter punctatum, nigrum, nitidum, ore testaceo. Thorax coleopteris paulo latior, antrorsum angustatus, apice duplo fere quam basi angustior, latitudine baseos paulo brevior, lateribus rotundatus, basi utrinque subsinuatus, angulis posterioribus retrorsum sat prominulis, rotundatis, convexus, parce subtilissime punctatus. Elytra thorace sesqui fere longiora, apice oblique truncata, convexa, densius subtiliusque punctata, margine laterali setoso, macula magna testaceo-rufa, ante basin transversali, dein secundum suturam elongata, ideoque arcuatim ab humero ad angulum suturalem apicalem pertingente. Abdomen parce subtilissime punctatum, segmentis singulis margine postico rufo-piceo, apice rufum, dorso parce, lateribus et apice crebre nigro-pilosum. Mesosternum subtiliter carinatum. Pedes rufi, tarsis testaceis, tibiis posterioribus vix spinulosis, tarsis posticis tibia haud longioribus.

Mas abdominis segmento sexto inferiore apice triangulariter exciso, superiore apice obtuso, tarsis anticis leviter dilatatis.

Femina abdominis segmento sexto inferiore apice rotundato ciliatoque, superiore quadrifido, laciniis intermediis profundius divisis, tarsis anticis fix dilatatis.

Variat elytrorum macula obliqua basali et litura longitudinali subsuturali disiunctis.

### **Tachyporus Grav.**

*Tachyporus* \*\* *Er. Col. March.* I. p. 392.

*Antennae* 11-articulatae, filiformes.

*Palpi maxillares articulo quarto subulato.*

*Ligula biloba, lobis rotundatis.*

*Elytra pectore longiora.*

*Abdomen marginatum.*

*Tarsi 5-articulati.*

Corpus oblongum, antice obtusiusculum, postice sensim attenuatum, convexum, alatum. Caput thoraci insertum, fronte convexiuscula, antice angustata, oculis mediocribus, rotundatis, convexis. Labrum transversum, integrum. Mandibulae breves, trigonae, intus membrana angusta interne ciliata fimbriatae. Maxillae (t. III. f. 3. b.) malis corneis, apice barbatis, exteriori interiorem parum superante, hac interne membranea. Palpi maxillares articulo primo minuto, secundo tertioque modice elongatis, longitudine subaequalibus, hoc incrassato, quarto parvo, subulato. Labium (t. IV. f. 3. a.) mento transversum, ligula lata, membranea, biloba, lobis integris, rotundatis, divergentibus, paraglossis ligulae connatis, nonnisi ciliarum serie cognoscen-

dis. Palpi labiales articulis duobus primis crassiusculis, subaequalibus, tertio haud brevior at tenuiore, acuminato. Antennae filiformes, apicem versus plerumque subincrassatae. Thorax amplus, transversim convexus, transversus, basi coleopteris arcu applicatus, antrorsum angustatus, apice truncatus, licet angulis anterioribus sat prominentibus, acutis. Scutellum triangulare. Elytra pectore longiora, tenuiter marginata, angulo apicali exteriori truncato. Abdomen distincte marginatum, apicem versus sensim at sat fortiter angustatum, vivis modice elongatum. Mesosternum simplex. Pedes modice elongati, coxis posticis parallelopipedis, ad femorum basin scutiformi-elevatis, femoribus omnibus laevibus, tibiis anticis muticis, posterioribus sat fortiter spinulosis, tarsis posterioribus elongatis, omnibus articulis 3 primis decrescentibus, subtus pubescentibus, quarto minimo, quinto leviter elongato.

#### O b s e r v a t i o n e s.

1. Caput et thorax laevissima, glaberrima, elytra subtiliter punctata, iam glabra, iam tenuiter pubescentia, margine laterali pilosello. Abdomen saepius subtilissime pubescens, lateribus et apice pilosellum.

2. Mas abdominis segmento sexto superiore integro, inferiore apice exciso, septimo conspicuo.

Femina abdominis segmento sexto superiore apice quadrifido, laciniis aequalibus, acutis, inferiore integro, apice pilis rigidis dense ciliato.

3. Victus in quisquiliis, sub foliis deciduis, musco etc.

\* 1. *T. obtusus*: *Testaceo-rufus, nitidus, pectore, elytrorum basi abdominisque apice nigris*. — Long.  $1\frac{2}{3}$  lin.

*Mannerh. Brachelytr.* 59. 6. — *Er. Col. March.* I. 392. 7.

*Staph. obtusus* Linn. *Syst. Nat.* I. II. 684. 9. — *Fab. Syst. Ent.* 266. 10. *Spec. Ins.* I. 336. 11. *Mant. Ins.* I. 221. 17. — *Schränk Enum. Ins. Aust.* 237. 451. — *Goeze Ent. Beitr.* I. 718. 9. — *De Vill. Ent.* I. 414. 9. — *Rossi Faun. Etr.* I. 251. 621. (*Ed. Hellw.* 311. 621.) — *Marsh. Ent. Britt.* 520. 61.

*Orypor. analis* Fab. *Mant. Ins.* I. 222. 3. *Ent. Syst.* I. II. 533. 10. *Syst. El.* II. 606. 10. — *Panz. Faun. Germ.* 22. 16. *Ent. Germ.* 360. 6.

*Tachypor. analis* Grav. *Micr.* 129. 8. *Mon.* 6. 10. — *Latr. Hist. nat. Crust. et Ins.* X. 8. 8. — *Gyll. Ins. Succ.* II. 239. 4. — *Zetterst. Faun. Lappon.* I. 58. 4. *Ins. Lapp.* 55. 5. — *Sahlb. Ins. Fenn.* I. 290. 4. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 516. 4. — *Runde Brachelytr. Hal.* 24. 3.

*Staph. vernalis* Müll. *Prod.* 98. 1103. — *Goeze Ent. Beitr.* I. 727. 27.

*Staph. dispar* var.  $\gamma$  Payk. *Mon. Staph.* 78. 54. *Faun. Suec.* III. 423. 69.

*Staph. chrysomelinus* var. 2 Walk. *Faun. Paris.* I. 273. 19.

*Staphyl. melanurus* Marsh. *Ent. Britt.* 525. 80.

Habitat in Europa.

Antennae capite thoraceque paulo longiores, apicem versus parum incrassatae, testaceae, articulis 4 primis flavis. Caput laevissimum, rufo-testaceum, oculis nigris. Thorax coleopteris paulo latior, latitudine dimidio brevior, angulis posterioribus obtusis, externe rotundatis, basi utrinque subsinuatus, transversim convexus, laevissimus, totus rufo-testaceus. Scutellum nigrum. Elytra thorace sesqui longiora, subtiliter punctata, rufo-testacea, basi lata nigra. Abdomen rufo-testaceum, segmentis ultimis duobus nigris, omnibus basi punctatis. Pectus nigrum. Pedes testacei.

Mas tarsis anticis leviter, femina vix dilatatis.



- \* 2. *T. abdominalis*: *Rufo-testaceus, nitidus, pectore abdomineque fusciscentibus, elytris thorace parum longioribus.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 393. 8.*

*Tachypor. abdominalis var. a* Gyll. *Ins. Suec. II. 238. 3.* — *Zetterst. Faun. Lappon. I. 58. 3. Ins. Lappon. 55. 4.* — *Sahlb. Ins. Fenn. I. 290. 3.* — *Mannerh. Brachelytr. 59. 4.*

Habitat in Europa.

Habitu omnino praecedentis, at paulo minor, rufo-testaceus, nitidus, pectore abdomineque toto fusciscentibus, hoc segmentis singulis margine late testaceis. Antennae capitis thoracisque longitudine, apice sensim subincrassatae fusciscentesque. Caput laevissimum. Thorax latitudine plus dimidio brevior, basi utrinque subsinuatus, angulis posterioribus retrorsum parum prominulis, rotundatis, transversim convexus, laevissimus. Elytra thorace parum longiora, parcius subtilissime punctata, tenuissime pubescentia, immaculata. Abdomen parcius subtiliter punctulatum.

Mas tarsis anticis leviter dilatatis, femina simplicibus.

Obs. Synonymia huius speciei minus perspicua, et dubito an loci a Gyllenhalio citati omnes huc referri possint. *Tachyp. abdominalis* Grav. certe ad *T. brunneum* pertinet. *Oxyporus abdominalis* Fab. (*Spec. Ins. Suppl. 501. 6. Mant Ins. I. 222. 7. Ent. Syst. I. II. 535. 17. Syst. El. II. 507. 17.*) mihi omnino dubius, in Museo etiam a Fabricio citato invisus, minime ad speciem nostram revocandus videtur, huic enim minime est abdomen atrum, nitidum. — *T. abdominalis var. b* apud Gyllenhal aegre ad hanc speciem alleganda, specimina ab auctore ipso sub hoc nomine missa, quae Dom. Schüppel mihi communicavit, potius specimina omnino immatura alius cuiusdam speciei vel aliarum quarundam specierum esse puto.

- \* 3. *T. rufus*: *Rufo-testaceus, nitidus, pectore abdomineque nigris, elytris thorace sesqui longioribus, basi prope scutellum nigricantibus.* — Long.  $1\frac{1}{2}$  lin.

*Tachypor. abdominalis* Boisd. et Lacord. *Faun. Ent. Paris. I. 515. 5.*

Habitat Lutetiae, Dôm. Aubé.

Statura omnino *T. chrysomelini*. Antennae capite thoraceque paulo longiores, tenuiores, apicem versus leviter sensim incrassatae, articulo ultimo apice abrupte acuminato, flavae, articulis penultimis tribus fuscis, ultimo testaceo. Caput laevissimum, dilute rufo-testaceum, nitidum. Thorax coleopteris paulo latior, latitudine baseos dimidio brevior, basi utrinque subsinuatus, angulis posterioribus retrorsum leviter prominulis, externe rotundatis, transversim convexus, laevissimus, dilute rufo-testaceus. Scutellum piceum. Elytra thorace sesqui longiora, subtiliter crebre punctata, basi prope scutellum nigricantia. Abdomen crebrius punctatum, nigrum, nitidum, segmentis 4 primis margine apicali obscure testaceis, quinto fere toto testaceo, sexto nigro. Pectus mesothoracis rufo-testaceum, metathoracis nigrum. Pedes dilute testacei, coxis omnibus concoloribus, posticis basi nigris.

Mas tarsis anticis fortius, femina leviter dilatatis.

Obs. Differt a praecedente praecipue statura sesqui maiore, abdomine magis nigro, elytris longioribus et basi nigricantibus. A *T. chrysomelino* facile distinguitur capite rufo-testaceo.

- \* 4. *T. saginatus*: *Niger, nitidus, thorace, elytris pedibusque rufis, antennis brevioribus, latiusculis, compressis, ferrugineis.* — Long 2 lin.

Grav. Mon. 6. 11. — Mannerh. Brachelytr. 59. 1. — Er. Col. March. I. 393. 9.

Habitat in Germania.

*T. chrysomelino* duplo maior, praecipue latior. Antennae capite thoraceque breviores, crassiusculae, apicem versus sensim incrassatae, ab articulo quarto inde sensim magis compressae, ferrugineae. Caput convexum, laevissimum, nigro-piceum, nitidum, subtus oreque rufis. Thorax coleopterorum latitudine, latitudine baseos dimidio brevior, basi truncatus, angulis posterioribus postice haud prominulis, rotundatis, transversim convexus, laevissimus, testaceo-rufus, nitidus. Scutellum piceum. Elytra thorace paulo longiora, subtiliter crebrius punctulata, testaceo-rufa, basi prope scutellum fusciscentia. Abdomen subtiliter punctulatum, nigrum, nitidum, segmentis singulis margine summo postico obscure testaceo. Pectus nigrum. Pedes rufo-testacei, coxis anterioribus concoloribus.

Mas tarsis anterioribus levissime dilatatis, femina simplicibus.

- \* 5. *T. erythropterus*: *Niger, nitidus, elytris pedibusque rufis, antennis brevioribus, crassis, ferrugineis.* — Long.  $1\frac{3}{4}$  lin.

*Oxyporus erythropterus* Panz. Faun. Germ. 27. 21.

Habitat in Germania meridionali, Dom. Dr. Waltl.

Habitu omnino *T. saginati*, at minor, *T. hypnorum* perparum maior. Antennae capite thoraceque breviores, crassiusculae, apicem versus incrassatae, ab articulo quarto inde sensim fortius compressae, ferrugineae. Caput laevissimum, nigrum, nitidum, ore testaceo. Thorax coleoptera latitudine parum excedens, latitudine baseos paulo brevior, antrorsum minus fortiter angustatus, lateribus minus rotundatis, basi utrinque leviter sinuatus, angulis posterioribus retrorsum modice prominentibus, late rotundatis, fortiter convexus, laevissimus, niger, nitidus, margine summo basali lateralique rufescente. Scutellum nigrum. Elytra thorace parum longiora, subtiliter punctata, saturate rufa, nitida, basi et circa scutellum tenui spatio nigricantia. Abdomen nigrum, nitidum, parcius subtiliter punctatum, segmentis singulis margine summo postico rufescente. Pectus nigrum. Pedes rufi, coxis anticeis piceis, brevisculi fortiusculique, tarsis brevioribus, subcompressis.

Mas tarsis anticeis vix dilatatis. Femina latet.

Obs. *T. saginato* antennarum structura affinis, differt autem thorace scutelloque nigris, thorace angulis posterioribus prominentibus, elytris distinctius punctulatis. *A. T. hypnorum* antennarum structura satis distinctus.

- X \* 6. *T. hypnorum*: *Niger, thoracis margine testaceo, pedibus elytrisque rufis, his macula marginali nigra.* — Long.  $1\frac{2}{3}$  lin.

Er. Col. March. I. 394. 11.

*Staph. hypnorum* Fab. Syst. Ent. 266. 13. Spec. Ins. I. 336. 16. Mant. Ins. I. 221. 23. — Goeze Ent. Beitr. I. 724. 7. — Marsh. Ent. Britt. 525. 79.

*Oxyporus hypnorum* Fab. Ent. Syst. I. II. 535. 22. Syst. El. II. 607. 22. — Panz. Ent. Germ. 361. 17.

- Staphyl. nitidulus* Fab. Spec. Ins. I. 337. 17. Mant. Ins. I. 221. 24. — Oliv. Ent. III. 42. 34. 48. t. 3. f. 28. — Marsh. Ent. Britt. 520. 64.
- Oxyporus minutus* Fab. Ent. Syst. I. II. 535. 20. Syst. El. II. 607. 20.
- Oxyporus marginatus* Punz. Ent. Germ. 359. 4. Faun. Germ. 27. 17.
- Tachyporus marginatus* Grav. Micr. 127. 5. Mon. 7. 13. — Latr. Hist. nat. Crust. et Ins. X. 6. 5 — Gyll. Ins. Suec. II. 237. 2. — Zetterst. Faun. Lappon. I. 57. 2. Ins. Lappon. 55. 3. — Sahlb. Ins. Fenn. I. 290. 2. — Mannerh. Brachelytr. 59. 3. — Guér. Iconogr. t. 10. f. 9. — Boisd. et Lacord. Faun. Ent. Paris. I. 514. 1. — Ru de Brachelytr. Hal. 21. 2.
- Staph. Blattinus* Schrank Enum. Ins. Aust. 236. 447. Faun. Boic. I. 446. 878.
- Staph. conicus* De Vill. Ent. I. 425. 72.
- Staph. martialis* Schrank Enum. Ins. Aust. 237. 450.
- Staph. dispar* var.  $\beta$  Payk Mon. Staph. 78. 54. Faun. Suec. III. 421. 69.
- Staph. chrysomelinus* var. 1 Walk. Faun. Paris. I. 272. 19.

Habitat in Europa.

Statura omnino *T. chrysomelini*. Antennae capitis thoracisque longitudine vel paulo longiores, graciliores, apicem versus parum crassiores, testaceae, apice subinfuscaetae. Caput laevissimum, nigrum, ore testaceo. Thorax coleopteris paulo latior, latitudine baseos dimidio brevior, basi truncatus, angulis posterioribus postice haud prominentibus, obtusis, externe rotundatis, fortiter convexus, niger, nitidus, margine basali angusto lateralique basin versus sensim latiore testaceis. Scutellum nigrum. Elytra thorace paulo longiora, subtiliter punctata, rufa, macula parva triangulari circa scutellum maculaque oblonga laterali nigris, tenuissime pubescentia. Abdomen crebrius subtiliter punctatum, nigrum, segmentis singulis margine apicali ferrugineo. Pectus nigrum. Pedes testacei, coxis piceis.

Mas tarsis anticis leviter, femina vix dilatatis.

- \* 7. *T. anticus*: Niger, nitidus, thorace elytris pedibusque rufo-testaceis, thorace disco nigro. — Long.  $1\frac{1}{2}$  lin.

Habitat in Aegypto, Dom. Prof. Ehrenberg.

*T. hypnorum* paulo minor et gracilior. Antennae capitis thoracisque prope longitudine, apicem versus haud incrassatae, testaceae, apice fuscescentes. Caput convexus, laevissimum, nigrum, nitidum, palpis testaceis. Thorax coleopterorum latitudine, latitudine baseos dimidio brevior, basi truncatus, angulis posterioribus haud prominulis, rotundatis, leviter transversim convexus, laevissimus, nitidus, testaceus, disco anteriore late nigro, seu niger, margine laterali et basali lato, anteriore angusto testaceo. Scutellum piceum. Elytra thorace sesqui prope longiora, subtiliter punctata, rufo-testacea, immaculata, nitida. Abdomen subtiliter punctatum, nigrum, segmentis singulis margine postico, quinti latiore, ferrugineo. Pectus nigrum. Pedes testacei, coxis anticis concoloribus.

Obs. *T. hypnorum* simillimus, distinctus autem statura minore, minus convexus, thorace coleopteris haud latiore, elytris constanter immaculatis, coxis anticis testaceis etc. A. *T. brunneo* differt statura duplo maiore, antennis apicem versus haud incrassatis etc.

- \* 8. *T. chrysomelinus*: Niger, nitidus, antennarum basi, pedibus, thorace elytrisque testaceo-rufis, his thorace paulo longioribus, crebre subtiliter punctatis, circa scutellum et margine laterali nigris. — Long.  $1\frac{2}{3}$  lin.



- Grav. Micr.* 128. 7. *Mon.* 7. 12. — *Latr. Hist. nat. Crust. et Ins.* X. 7. 7. t. 79. f. 9. *Gen. Crust. et Ins.* I. 300. 1. — *Gyll. Ins. Suec.* II. 236. 1. — *Sahlb. Ins. Fenn.* I. 289. 1. — *Zetterst. Faun. Lappon.* I. 57. 1. *Ins. Lappon.* 54. 1. — *Manne. h. Brachelytr.* 59. 2. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 515. 2. — *Runde Brachelytr. Hal.* 24. 1. — *Er. Col.-March.* I. 393. 10.
- Staph. chrysomelinus* Linn. *Faun. Suec. n.* 855. *Syst. Nat.* I. II. 685. 21. — *Fab. Syst. Ent.* 267. 17. *Spec. Ins.* 337. 21. *Mant. Ins.* I. 221. 28. — *Schrank Enum. Ins. Austr.* 236. 448. *Faun. Boic.* I. 646. 789. — *Ström. Act. Nidros.* III. 405. 28. — *Müll. Prod.* 98. 1105. — *Goeze Ent. Beitr.* I. 720. 21. — *De Vill. Ent.* I. 417. 21. — *Walk. Faun. Paris.* 1. 272. 19. — *Foucr. Ent. Paris.* 169. 18. — *Rossi Faun. Etr.* I. 250. 616. (*Ed. Helhv.* 309. 616. var. 1. a.) — *Oliv. Ent.* III. 42. 33. 47. t. 3. f. 22. — *Marsh. Ent. Britt.* 519. 42.
- Oxyporus chrysomelinus* Fab. *Ent. Syst.* I. II. 534. 15. *Syst. El.* II. 606. 14. — *Panz. Faun. Germ.* 9. 14. *Ent. Germ.* 360. 12. — *Cederh. Faun. Ingr.* 120. 371.
- Oxyporus melanocephalus* Fab. *Ent. Syst.* I. II. 534. 16. *Syst. El.* II. 607. 16.
- Staph. merdarius* Marsh. *Ent. Britt.* 521. 68.
- Oxyporus merdarius* Panz. *Ent. Germ.* 360. 8. *Faun. Germ.* 27. 18.
- Staph. chrysomelinus* var.  $\alpha$  Payk. *Mon. Staph.* 57. 74.
- Staph. dispar* var.  $\alpha$  Payk. *Faun. Suec.* III. 423. 69.
- Le Staph. couleur de paille* Geoffr. *Hist. Ins.* I. 368. 18.

Habitat in Europa.

Statura et magnitudo T. obtusi. Antennae capitis thoracisque longitudine, graciles, apice vix incrassatae, articulis 5—10 sensim paulo brevioribus at parum crassioribus, subobconicis, ultimo his paulo crassiore, ovato, apice abrupte acuminato, articulis tribus primis dilute testaceis, reliquis rufo-fuscescentibus. Caput mediocre, leviter convexum, laevisimum, nitidum, nigrum, ore testaceo. Thorax coleopteris paulo latior, latitudine baseos tertia fere parte brevior, basi truncatus, angulis posterioribus retrorsum haud prominulis, obtusis, externe fortius rotundatis, transversim convexus, laevissimus, rufo-testaceus, nitidus. Scutellum nigrum. Elytra thorace tertia fere parte longiora, leviter convexa, crebrius subtiliter punctata, glabra, nitida, rufo-testacea, spatio parvo communi triangulari circa scutellum, et margine laterali, postice abbreviato, nigris. Abdomen crebrius subtiliter punctatum, glabrum, lateribus et apice pilosellum, nigrum, segmentis singulis margine posteriore late rufo-testaceo, ultimo piceo-testaceo. Pectus nigrum. Pedes testacei, coxis anticis concoloribus.

Mas tarsis anticis modice, femina perparum dilatatis.

- X \* 9. *T. solutus*: Niger, nitidus, ore, antennis, pedibus, thorace elytrisque rufo-testaceis, his thorace paulo longioribus, circa scutellum infuscatitis. — Long.  $1\frac{2}{3}$  lin.

*Tachyporus saginatus* Boisd. et Lacord. *Faun. Ent. Paris.* I. 515. 13.

Habitat in Europa.

Statura et summā affinitas T. chrysomelini, et maximis eius individuis magnitudine aequalis. Antennae capite thoraceque paulo longiores, apicem versus distinctius incrassatae, articulis ultimis sex evidenter crassioribus, crassitie autem haud brevioribus, leviter obconicis, excepto ultimo oblongo-ovato, apice acuminato. Caput mediocre, leviter convexum, laevisimum, nitidum, nigrum, ore magis minusve testaceo. Thorax coleopteris paulo latior, lat-

tudine baseos tertia fere parte brevior, basi truncatus, angulis posterioribus obtusis, externe subrotundatis, sat fortiter transversim convexus, laevis, nitidus, rufo-testaceus. Scutellum nigrum. Elytra thorace tertia fere parte longiora, leviter convexa, sat crebre subtiliter punctata, glabra, rufo-testacea, nitida, basi spatio communi triangulari nigricante. Abdomen sat crebre subtiliter punctatum, glabrum, lateribus et apice nigro-pilosellum, nigrum, nitidum, segmentis 4 primis margine tenui posteriore testaceo. Pectus nigrum. Pedes testacei, coxis anticis concoloribus.

Mas tarsis anticis modice, femina leviter dilatatis.

Obs. 1. Differt a *T. chrysomelino* antennis extrorsum crassioribus, totis testaceis, elytris margine laterali concolore, ano nigro, a *T. saginato* statura minore, minus lata, et antennarum structura.

2. *Tachypori chrysomelini* var. b Gyll. Sahlb. Zett. vix ad hanc speciem referenda, quae speciei genuinae magnitudine inferior fertur. *Oxyporus merdarius* Panz. pariter potius *T. chrysomelinum* quam hanc speciem exhibere videtur, hac enim minor est, et antennae apicem versus fuscescunt.

\* 10. *T. tersus*: *Niger, nitidus, antennarum basi, pedibus thorace elytrisque rufo-testaceis, his thorace sesqui longioribus, parce obsoleteque punctatis, margine laterali et circa scutellum nigris.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Germania, Kirstein.

*T. chrysomelino* similis, at minor, angustior et minus convexus, statura omnino *T. pusilli*. Antennae capitis thoracisque longitudine, apicem versus perparum incrassatae, articulis 5 — 10 sensim paululum brevioribus et crassioribus, subobconicis, ultimo praecedente paulo crassiore, ovato, apice abrupte acuminato, fuscae, articulis 4 primis testaceis. Caput medio cre, leviter convexus, laevis, politus, nigrum, ore concolore, oris partibus inferioribus testaceis. Thorax coleopterorum latitudine, latitudine baseos dimidio fere brevior, basi utrinque subsinuatus, angulis posterioribus rotundatis, transversim convexus, laevis, nitidus, rufo-testaceus. Elytra thorace sesqui longiora, leviter convexa, parce subtiliter obsoleteque punctata, glabra, nitida, rufo-testacea, spatio parvo communi triangulari circa scutellum, et margine laterali tenuiore, summo apice abbreviato, nigris. Abdomen parcius subtiliter punctatum, glabrum, lateribus et apice parce nigro-pilosellum, nigrum, nitidum, segmentis singulis margine apicali obscure testaceis. Pectus nigrum. Pedes testacei, coxis anticis concoloribus.

Mas tarsis anticis leviter dilatatis, femina simplicibus.

Obs. Affinis certe *T. chrysomelino*, at distinctus statura minore, angustiore, minus convexus, thorace brevior, coleopteris haud brevior, his parce obsoleteque punctatis, fere laevigatis, et ore testaceo.

\* 11. *T. arduus*: *Niger, nitidus, ore, antennarum basi, pedibus, thorace elytrisque rufo-testaceis, his thorace paulo longioribus, parcius punctatis, circa scutellum et margine laterali nigris.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Pennsylvania, Dom. Zimmermann.

*T. chrysomelino* minor et angustior, statura omnino *T. scituli*. Antennae capite thoraceque paulo longiores, apicem versus sensim leviter incrassatae, articulis 5 — 10 sensim paulo brevioribus et crassioribus, ultimo praecedente paulo crassiore, subovato, apice abrupte acuminato, rufo-fuscae,

articulis 4 primis testaceis. Caput mediocre, convexum, laevissimum, nitidum, nigrum, ore testaceo. Thorax coleopteris vix latior, latitudine baseos tertia fere parte brevior, basi truncatus, angulis posterioribus retrorsum haud prominulis, rotundatis, transversim convexus, laevissimus, nitidus, rufo-testaceus. Scutellum nigrum. Elytra thorace tertia fere parte longiora, leviter convexa, parcius et minus subtiliter punctata, subtiliter tenuiterque pubescentia, rufo-testacea, nitida, spatio parvo communi triangulari circa scutellum, et vitta marginali abbreviata, nonnunquam obsoleta, nigricantibus. Abdomen crebrius minus subtiliter punctatum, subtilissime tenuissimeque pubescens, lateribus et apice longius nigro-pilosellum, nigrum, nitidum, segmentis singulis margine posteriore, anterioribus 4 angustiore, quinto latiore rufo-testaceis, ultimo apice saepius piceo. Pectus nigrum. Pedes testacei, coxis anticis concoloribus.

Mas tarsis anticis leviter dilatatis, femina simplicibus.

Variat elytris spatio inter maculam scutellarem et marginalem magis minusve infuscato.

Obs. Ab Europaeis omnino distinctus, differt a sequentibus ore testaceo, a praecedentibus statura minore, elytris pubescentibus, minus subtiliter punctatis.

\* 12. *T. humerosus*: *Niger, nitidus, antennarum basi, pedibus, thorace elytrisque rufo-testaceis, his thorace longioribus, crebrius subtiliter punctatis, pubescentibus, lateribus et macula scutellari nigris.* — *Long.*  
 $1\frac{1}{3}$  —  $1\frac{1}{2}$  lin.

*Tachyporus ruficollis* var. (*humerosus* Knoch) Grav. Mon. p. 8.

*Tachyporus lateralis* Grav. Micr. 127. 4. Mon. 7. 14. — Latr. Hist. nat. Crust. et Ins. X. 6. 4. (*forte*)

*Tachyporus ruficollis* Runde Brachelytr. Hal. 25. 4.

Habitat in Germania.

*T. chrysomelino* dimidio vel duplo minor, brevior et convexior. Antennae capitis thoracisque longitudine, tenuiores, apicem versus vix incrassatae, articulis 4 — 10 sensim paulo brevioribus at vix crassioribus, subobconicis, ultimo praecedente duplo prope longiore, subovato, apice abrupte acuminato, fuscae, basi testaceae. Palpi testacei. Caput magnum, convexum, laevissimum, nitidum, nigrum, ore concolore. Thorax coleopteris vix latior, latitudine baseos tertia fere parte brevior, antrorsum leviter angustatus, basi truncatus, angulis posterioribus rotundatis, sat fortiter transversim convexus, laevissimus, rufo-testaceus, nitidus. Scutellum nigrum. Elytra thorace tertia parte longiora, sat convexa, crebrius subtiliter punctata, densius subtiliter pubescentia, nitida, rufa, spatio minuto communi triangulari circa scutellum, margineque basali et laterali, hoc apice abbreviato, nigris, spatio interiora saepius magis minusve infuscato, limbo apicali magis rufo-testaceo. Abdomen crebrius subtiliusque punctatum, subtilissime tenuiter pubescens, lateribus et apice nigro-pilosellum, nigrum, nitidum, segmentis 4 primis margine posteriore testaceo. Pectus nigrum. Pedes testacei, coxis anterioribus concoloribus.

Mas tarsis anticis leviter dilatatis, femina simplicibus.

Obs. *T. chrysomelino* similis, distinguitur autem corpore minore, convexiore, elytris pubescentibus et ore concolore, *T. ruficollis* magis affinis, differt autem an-



tennis longioribus, praeter basin fuscis, thorace brevior, antrorsum evidenter angustato, elytris longioribus, et tarsis anticis maris leviter dilatatis.

\* 13. *T. ruficollis*: *Niger, nitidus, thorace pedibusque rufis, elytris thoracis longitudine, piceis, apice rufescentibus.* — Long.  $1\frac{1}{2}$  lin.

*Grav. Micr.* 128. 6. *Mon.* 7. 16. — *Latr. Hist. nat. Crust. et Ins.* X. 7. 6.

*Staph. chrysomelinus* var. *1. a. β Rossi Faun. Etr. Ed. Helhv.* I. 309. 616.

Habitat in Germania.

Antennae capitis thoracisque prope longitudine, tenues, apicem versus vix incrassatae, articulis 5—10 sensim paululum brevioribus et crassioribus, subobconicis, ultimo praecedente perparum maiore, subobconico, apice abrupte acuminato, totae testaceae. Palpi testacei. Caput magnum, convexum, laevissimum, nigrum, nitidum, ore concolore. Thorax coleopteris paulo latior, latitudine baseos paulo brevior, antrorsum parum angustatus, basi truncatus, angulis posterioribus retrorsum haud prominentibus, rotundatis, fortiter convexus, laevissimus, testaceo-rufus, nitidus. Scutellum piceum, Elytra thorace vix longiora, convexa, subtiliter punctata, subtilissime tenuiter pubescentia, iam nigro-, iam fusco-picea, nitida, margine apicali rufescente. Abdomen crebre subtiliter punctatum, subtiliter tenuiterque pubescens, lateribus et apice parce nigro-pilosellum, nigrum, nitidum, segmentis singulis margine summo postico testaceo. Pectus nigrum. Pedes rufo-testacei, coxis anterioribus concoloribus.

Mas tarsis anticis fortiter dilatatis, femina simplicibus.

Variat elytris rufo-piceis vel omnino rufis.

Obs. Omnino distinctus statura brevi, convexa, capite magno, antennis brevibus tenuibusque, totis testaceis, thorace longiore et antrorsum perparum angustato. Var. ad calcem memorata *T. humeroso* similis, notis autem modo laudatis distinguenda. *T. ruficollis* Gyll. ad *T. transversalem* referendus.

\* 14. *T. pusillus*: *Niger, nitidus, thoracis margine elytrisque fuscis, his thorace sesqui longioribus, macula marginali nigra.* — Long.  $1\frac{1}{4}$  lin.

*Grav. Mon.* 9. 18. — *Gyll. Ins. Suec.* II. 241. 6. — *Zetterst. Faun. Lappon.* I. 59. 6. *Ins. Lappon.* 55. 7. — *Sahlb. Ins. Fenn.* I. 291. 6. — *Mannerh. Brachel.* 60. 8. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 516. 6. — *Runde Brachelytr. Nat.* 25. 5. — *Er. Col. March.* I. 394. 12.

Habitat in Europa.

Antennae capitis thoracisque longitudine, apicem versus parum crassiores, fuscae, basi testaceae. Palpi picei, articulis duobus primis flavo-testaceis. Caput mediocre, leviter convexum, laevissimum, nigrum, nitidum, ore concolore. Thorax coleopterorum latitudine, latitudine baseos dimidio brevior, basi truncatus, angulis posterioribus haud prominentibus, obtusis, externe rotundatis, modice convexus, laevissimus, niger, nitidus, margine basali angustiore lateraliq. lato fusco-testaceo. Scutellum nigrum. Elytra thorace sesqui longiora, crebre subtiliter punctata, fusca, margine laterali nigro. Pectus et abdomen nigra, nitida, confertim subtiliter punctata. Pedes testacei, coxis anticis piceis.

Mas tarsis anticis leviter dilatatis, femina simplicibus.

Obs. Tachyporo hypnorum colore, licet obscuriore, affinis, differt autem statura minore, minus convexa, thoraceque coleopteris haud latiore.

- \* 15. *T. scitulus*: *Niger, nitidus, thoracis margine laterali elytrorumque apicali rufo-piceis, antennarum basi pedibusque rufis, elytris thorace paulo longioribus.* — Long.  $1\frac{1}{4}$  lin.

*Er. Col. March. I. 395. 13.*

*Tachyporus pusillus* var. *Grav. Mon. 9. 18.*

*Tachypor. pusillus* var. *b* *Gyll. Ins. Suec. II. 241. 6. — Zetterst. Faun. Lappon. I. 59. 6. Ins. Lappon. 55. 7. — Sahlb. Ins. Fenn. I. 291. 6. — Mannerh. Brachelytr. 60. 8.*

Habitat in Europa.

Statura *T. pusilli*, at paulo convexior, nitidus. Antennae capitis thoracisque fere longitudine, apicem versus parum incrassatae, fuscae, basi testaceae. Caput mediocre, convexum, laevissimum, nigrum, ore concolore, oris partibus piceo-testaceis. Thorax coleopteris parum latior, latitudine baseos dimidio brevior, basi truncatus, angulis posterioribus haud prominulis, obtusis, externe rotundatis, modice transversim convexus, margine laterali latiore summoque basali piceo-rufescentibus. Elytra thorace paulo longiora, subtilissime punctulata, tenuiter subtiliterque pubescentia, margine apicali piceo-rufescente. Abdomen crebre subtiliter punctatum, segmentis 2 — 5 saepius tenuiter rufo-marginatis. Pedes rufo-testacei, coxis anticis picescentibus.

Mas tarsis anticis leviter dilatatis, femina simplicibus.

Variat thoracis margine laterali et basali elytrorumque apicali laete rufo-testaceis.

Obs. *T. pusillo* similis, differt corpore paulo convexiore, elytrorum colore, et punctura parciore et subtiliore.

- \* 16. *T. nanus*: *Niger, nitidus, ore, antennarum basi, elytrorum apice pedibusque rufis, thoracis margine piceo, elytris thorace longioribus, distinctius punctatis.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Statura et summa affinitas praecedentis. Niger, nitidus. Antennae capitis thoracisque prope longitudine, graciles, apice paulo crassiores, piceo-rufae, basi testaceae. Caput mediocre, convexum, laevissimum, ore testaceo. Thorax coleopteris paululum latior, latitudine baseos dimidio brevior, basi truncatus, angulis posterioribus haud prominentibus, rotundatis, modice convexus, laevissimus, margine basali laterali que tenuiore piceo-rufescente. Elytra thorace tertia parte longiora, crebre minus subtiliter punctata, parce pubescentia, apice (tertia elytri parte) laete rufa. Abdomen parce subtiliter punctatum, segmentis singulis margine postico rufescente. Pedes rufo-testacei, coxis anticis femoribusque picescentibus.

Mas tarsis anticis leviter dilatatis. Femina latet.

Obs. *A. T. scitulo* distinctus antennis brevioribus, dilutioribus, ore testaceo, elytris distinctius punctatis, apice latius rufis, etc.

- \* 17. *T. transversalis*: *Niger, nitidus, thorace elytrisque rufis, antice dimidiato-nigris, pedibus testaceis.* — Long.  $1\frac{1}{4}$  lin.

*Grav. Mon. 8. 17. — Er. Col. March. I. 395. 14.*

*T. ruficollis* Gyll. *Ins. Suec. II.* 240. 5. — *Zetterst. Faun. Lappon. I.* 58. 5. *Ins. Lappon.* 55. 6. — *Sahlb. Ins. Fenn. I.* 291. 5. — *Mannerh. Brachelytr.* 59. 7.  
*T. dispar* var. ♂ *Payk. Mon. Staph.* 78. 54. *Faun. Suec. III.* 424. 69.

Habitat in Germania.

Statura *T. pusilli*. Antennae capitis thoracisque longitudine, apicem versus parum crassiores, ferrugineae, medio fuscescentes. Palpi testacei, articulo tertio fusco. Caput minusculum, leviter convexum, laevissimum, nigrum, nitidum, ore testaceo. Thorax coleopteris paulo latior, latitudine baseos dimidio brevior, basi truncatus, modice convexus, angulis posterioribus haud prominentibus, rotundatis, satis convexus, laevissimus, testaceo-rufus, antice plerumque dimidiato-nigricans, nitidus. Scutellum nigrum. Elytra thorace paulo longiora, crebre subtiliter punctata, parce subtiliterque pubescentia, nitida, antice nigra, pone medium testaceo-rufa. Abdomen crebre subtiliter punctatum, nigrum, segmentis singulis margine postico late rufo. Pectus nigrum. Pedes rufo-testacei, coxis anticis concoloribus.

Mas tarsis anticis vix dilatatis, femina simplicibus.

\* 18. *T. brunneus*: Rufo-testaceus, capite, thoracis disco pectoreque piceis, elytris thorace sesqui longioribus. — *Long.*  $1\frac{1}{4}$  lin.

*Er. Col. March. I.* 395. 15.

*Oxyporus brunneus* *Fab. Ent. Syst. I. II.* 535. 19. *Syst. El. II.* 607. 19. — *Panz. Ent. Germ.* 361. 15.

*Staphyl. nitidulus* *Oliv. Ent. III.* 42. 34. 48 t. 3. f. 28.

*Tachyporus nitidulus* *Grav. Micr.* 126. 2. *Mon.* 11. 20. — *Latr. Hist. nat. Crust. et Ins. X.* 6. 2. — *Gyll. Ins. Suec. II.* 242. 7. — *Sahlb. Ins. Fenn. I.* 292. 7. — *Mannerh. Brachelytr.* 60. 9. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 517. 7. — *Runde Brachelytr. Hal.* 25. 6.

*Var.: Totus testaceus.*

*Tachyporus abdominalis* *Grav. Micr.* 127. 3. *Mon.* 7. 15. — *Latr. Hist. nat. Crust. et Ins. X.* 6. 3.

*Tachyporus scutellaris* *Boisd. et Lacord. Faun. Ent. Paris. I.* 517. 8.

Habitat in Europa; in Aegypto, Dom. Prof. Ehrenberg; in America septentrionali, Dom. Zimmermann.

Praecedentibus minor, angustior et minus convexus. Antennae capitis thoracisque longitudine, apice vix crassiores, totae testaceae. Caput minusculum, leviter convexum, laevissimum, piceum, nitidum, ore testaceo. Thorax coleopterorum latitudine, latitudine baseos dimidio brevior, basi truncatus, angulis posterioribus haud prominulis, obtusis, externe rotundatis, leviter convexus, laevissimus, testaceus, nitidus, disco piceo. Scutellum piceo-testaceum. Elytra thorace sesqui longiora, crebre subtilius punctata, tenuiter subtiliterque pubescentia, testacea, nitida. Abdomen crebre minus subtiliter punctatum, fusco-testaceum, segmentis singulis apice dilutioribus. Pectus piceum. Pedes flavo-testacei.

Mas tarsis anticis leviter dilatatis, femina simplicibus.

Variat capite, thoracis disco pectoreque testaceis concoloribus.



## Species mihi invisae.

19. *T. nigricornis*: *Nigro-piceus, nitidus, antennarum articulo ultimo, thoracis lateribus pedibusque posticis rufo-testaceis, elytris rufis, regione scutelli nigris.*

*Gyll. Ins. Suec. IV. 469. 1 — 2.*

Habitat in Vestrogothia.

*T. chrysomelino* et affinis magnitudine, statura et partium formatione simillimus, colore antennarum et pedum anteriorum a reliquis praecipue distinctus. Caput nigrum, nitidissimum, glabrum, laeve, ore palpisque etiam nigris, pubescentibus. Antennae paulo breviores et validiores quam in affinis, extrorsum crassiores, nigrae, articulo ultimo solo testaceo. Thorax amplus, nigro-brunneus, lateribus rufescens. Scutellum nigrum. Elytra rufo-testacea, circa scutellum nigricantia, et iuxta humerum litura fusca notata. Abdomen nigro-piceum, segmentorum marginibus dilutioribus. Corpus sub-nigro-piceum. Pedes quatuor anteriores nigro-picei, postici et tarsi anteriores testacei. (*Gyll.*)

20. *T. obscurellus*: *Convexus, nitidus, laevis, glaber, niger, antennarum basi, thorace pedibusque anterioribus rufo-testaceis, elytris fuscis. — Long. 1¼ lin.*

*Zetterst. Ins. Lappon. 54. 2.*

Habitat in Lapponia Umensi rarius.

Similis *T. chrysomelino*, marginato, etc., colore ab illis tantum differt. Caput nigrum. Antennae fuscae, articulo primo testaceo. Thorax rufus. Elytra obscura s. fusco-rufa, margine summo apicali pallido. Abdomen nigrum. Pedes obscure rufo-testacei, femoribus posticis nigricantibus s. fuscis, tibiis eiusdem parvis testaceis, tarsis pallidis. (*Zetterst.*)

Obs. Colore corporis cum *T. ruficollis* Gr. convenire videtur, at antennarum et pedum posticorum colore differt.

21. *T. nigriceps*: *Latus, subconvexus, pallidus, capite nigro, ano infusato.*

*Mannerh. Brachelytr. 59. 5.*

Habitat in Fennia: ad Wilnäs semel captus.

*T.* abdominali duplo minor, capite nigro, postice transversim profundius impresso, et thorace elytris parum latiore diversus. (*Mannerh.*)

Obs. -Species perquam dubia, forte castiganda.

**Habrocerus Er.**

*Er. Col. March. I. p. 400.*

*Antennae 11-articulatae, capillares.*

*Palpi maxillares articulo quarto acuminato.*

*Ligula biloba, lobis truncatis.*

*Elytra pectore longiora.*

*Tarsi 5-articulati.*

Corpus oblongum, leviter convexum, alatum. Caput thoraci insertum, rotundatum, convexum, deflexum, oculis mediocribus, rotundatis, modice pro-

minutis. Labrum subquadratum, apice leviter sinuatum. Mandibulae minutae. Maxillae (t. IV. f. 4. b.) mala inferiore rhomboidea, externe coriacea, interne membranacea, apice breviter barbata, mala superiore interne coriacea, externe et apice dilatato-rotundato barbata, membranacea. Palpi maxillares elongati, articulo primo minuto, secundo tertioque longitudine subaequalibus, hoc apicem versus sensim incrassato, quarto huic fere aequali, elongato-conico, acuminato. Labium (t. IV. f. 4. a.) mento transversum, ligula lata, truncata, medio incisa, membranacea, paraglossis ligulae connatis, eique aequalibus, non nisi linea ciliata indicatis. Palpi labiales ligulam vix superantes, articulo primo sat magno, cylindrico, secundo tertioque minutis, subaequalibus, hoc apice truncato. Antennae (t. IV. f. 4. c.) 11-articulatae, articulis duobus primis crassiusculis, reliquis tenuibus, verticillato-pilosellis. Thorax transversus, basi coleopterorum latitudine, antrorsum leviter angustatus, transversim convexus, basi truncatus, apice pro receptione capitis emarginatus, lateribus leviter rotundatis, angulis posterioribus obtusis, subrotundatis. Scutellum minutum, triangulare. Elytra thorace paulo longiora, angulo apicali posteriore oblique truncato. Abdomen marginatum, apicem versus sensim angustatum, apice depressum, segmentis sexto septimoque occultis. Mesosternum antice carinatum. Pedes breviusculi, coxis intermediis approximatis, posticis triangularibus, trochanteribus posticis maiusculis, tibiis omnibus subtilissime spinulosis, tarsis 5-articulatis, tenuibus, posterioribus articulo primo elongato, sequentibus tribus sensim longitudine decrescentibus, anticis in utroque sexu simplicibus.

#### Observationes.

1. Habitus corporis omnino Tachypori, at minus convexus. Caput et thorax laevissima glaberrimaque, hic margine parce pilosello, elytra punctulata, glabra, margine parce pilosello, abdomen subtiliter tenuiterque pubescens, lateribus et apice longius pilosellum.

2. Quodammodo intermedium videtur hoc genus inter Tachyporum et Tachinum, palporum maxillarum articulus tertius apice acutus ut in Tachyporo, at elongatus, et basi tertio crassitie aequalis, ut in Tachino; ab utroque praeterea notis pluribus discedit, praecipue antennarum abdominisque structura nec non coxis posticis triangularibus. Antennarum figura insolita cum Trichophyo convenit.

3. Mas abdominis segmento quinto inferiore apice leviter emarginato, sexto paululum eminulo, femina segmento quinto apice truncato, sexto omnino occulto distinguuntur.

\* 1. *H. capillaricornis*: Nigro-piceus, nitidus, antennis pedibusque testaceis. — Long.  $1\frac{1}{4}$  lin.

Er. Col. March. I. 401. 1.

*Tachyporus capillaricornis* Grav. Mon. 10. 19.

Habitat in Germania, in silvis, sub foliis deciduis, in Gallia, Mus. Dom. Dr. Aubé, in Sardinia, Dom. Prof. Gené.\*

Piceus vel nigro-piceus, nitidus, ore, antennis pedibusque testaceis, capite, thorace scutelloque glaberrimis, thorace apice punctis quatuor transversim positus, sat distinctis notato. Elytra obsolete parceque punctata, glabra. Abdomen confertim subtiliter punctatum, subtiliter pubescens, pilosellum.

**Tachinus Grav.***Tachinus Fam. I. Grav. Micr. I. p. 135.**Antennae 11-articulatae, filiformes.**Palpi maxillares filiformes.**Ligula biloba.**Elytra pectore longiora.**Tarsi 5-articulati.*

Corpus oblongum, magis minusve convexum, plerumque alatum. Caput thoraci insertum, fronte convexiuscula, antice angustata, oculis medicoribus, rotundatis, leviter prominulis. Labrum transversum, apice leviter sinuatum. Mandibulae trigonae, obtusiusculae, intus membrana tenui, subtiliter ciliata fimbriatae. Maxillae (t. IV. f. 5. b.) malis basi subcorneis, apice membranaceis, interiore apice oblique truncato, exteriore rotundato barbatis, hac interiori aut aequali aut eam leviter superante. Palpi maxillares modice elongati, filiformes, articulo primo minuto, brevi, secundo tertio paulo longiore, quarto tertio vel paulo longiore, vel subaequali, apice acuminato. Labium (t. IV. f. 5. a.) mento transversum, ligula lata, medio incisa, lobis oblique introrsum truncatis, membranacea, paraglossis ligulae omnino connatis eamque haud excedentibus, interne subtiliter ciliatis. Palpi labiales filiformes, articulo primo secundo paulo longiore, tertio praecedentibus haud tenuiore, primo longitudine aequali, apice subacuminato. Antennae filiformes, apice saepius leniter sensimque subincrassatae, articulo ultimo ovato. Thorax transversus, basi coleopterorum latitudine, vel iis paulo latior, antrorsum angustatus, apice pro receptione capitis emarginatus, lateribus rotundatus, basi vel truncata vel utrinque leviter sinuata, coleopteris arcu applicata, angulis posterioribus saepe postice prominentibus, anterioribus acutiusculis. Scutellum rotundatum. Elytra sat magna. Abdomen marginatum, apicem versus sensim attenuatum. Pedes breviusculi, coxis posticis paralletopipedis, ad femorum basin scutiformi-elevatis, tibiis omnibus spinulosis, tarsis 5-articulis, minus elongatis, articulis 4 primis longitudine sensim decrescentibus.

**Fam. I.***Mesosternum carinatum.**Tarsi antici in utroque sexu simplices.*

Statura Tachypori, corpus saepius convexius, posterius attenuatum, antice obtuse rotundatum. Palpi maxillares articulo ultimo magis acuminato, nonnunquam conico, penultimo saepius haud longiore. Femora valida, compressa.

Mas abdominis segmento sexto superiore apice quadrifido, laciniis spiniformibus, intermediis longioribus, inferiore fisso, quinto inferiore plerisque integro.

Femina segmento sexto superiore et inferiore apice quadrifidis, laciniis spiniformibus, aequalibus.

**Observationes.**

1. Corpus minutum, convexum, posterius valde attenuatum, capite, thorace elytrisque glaberrimis, abdomine saepius tenuissime subtilissimeque pubescente, la-



teribus et apice parce pilosello. Scutellum magnum. Elytra convexa, lateribus subtiliter marginata, secundum marginem saepius longitudinaliter impressa, angulo apicali exteriori obtuso,

2. Habitus omnino Tachypori, notas autem distinctivas praebeant abdominis segmentum sextum aliter constructum, mesosternum carinatum, tarsi antici in utroque sexu simplices, posteriores haud elongati, tibia breviores.

3. A Fam. altera haecce notis supra laudatis discedit, reliqui characteres satis exacte congruunt, oris etiam structura, qua nonnisi palpis maxillaribus paululum magis acuminatis differt. Generi igitur peculiari condendo characteres aegre sufficient.

\* 1. *T. silphoides*: Niger, nitidus, thoracis limbo, elytrorum vitta submarginali, sutura apiceque pedibusque flavis. — Long.  $1\frac{1}{3}$  lin.

Gyll. Ins. Suec. II. 267. 14. — Sahlb. Ins. Fenn. I. 303. 13. — Mannerh. Brachelytr. 62. 13. — Boisd. et Lacord. Faun. Ent. Paris. I. 513. 11. — Runde Brachelytr. Hal. 26. 6. — Er. Col. March. I. 396. 1.

Staphyl. silphoides Linn. Syst. Nat. I. II. 684. 11. — Goeze Ent. Beitr. I. 719. 10. — De Vill. Ent. I. 414. 11. — Marsh. Ent. Britt. 521. 66.

Oxyporus suturalis Panz. Faun. Germ. 18. 20. Ent. Germ. 360. 7.

Tachinus suturalis Grav. Micr. 144. 15. — Latr. Hist. nat. Crust. et Ins. IX. 397. 15.

Tachyporus suturalis Grav. Mon. 13. 22.

Tachinus marginalis Grav. Micr. 192. 4.

Tachyporus marginalis Grav. Mon. 13. 23.

Tachyporus limbatus Grav. Mon. 12. 21. (forte)

Staph. dispar var.  $\zeta$  Payk. Mon. Staph. 79. 54.

Staph. dispar var.  $\varepsilon$  Payk. Faun. Suec. III. 424. 69.

Habitat in Europa et in America septentrionali.

Statura omnino Tachypori ruficollis, niger, nitidus. Antennae capite thoraceque paulo longiores, apice haud crassiores, articulis 4 — 10 subobconicis, crassitie longioribus, ultimo oblongo, apice acuminato, ferrugineae, basi flavo-testaceae. Palpi testacei. Caput magnum, convexum, laevissimum. Thorax coleopterorum latitudine, latitudine dimidio fere brevior, antrosum parum angustatus, lateribus leviter rotundatus, basi utrinque leviter sinuatus, angulis posterioribus retrorsum leviter prominentibus, subrectis, externe rotundatis, convexus, laevissimus, limbo laterali lato margineque tenui basali et apicali flavo-testaceis. Elytra thorace paulo longiora, modice convexa, angulo apicali exteriori oblique truncato, dense subtilissime punctulata, marginē suturali et apicali vittaque submarginali, iam ante flavedinem abrupta, iam cum illa confluenta, flavo-testaceis. Abdomen dense subtiliter punctulatum, densius subtilissime pubescens, lateribus et apice nigro-pilosellum, subtus apice testaceum. Mesosternum subtiliter carinatum. Pedes testacei.

Obs. 1. Elytra signaturis flavis iam angustioribus iam latioribus, ita ut nonnunquam evadant flavo-testacea, macula oblonga submarginali alteraque discoidali nigris, utraque cum nigridine baseos cohaerente.

2. *T. marginalis* Americanus cum *T. silphoide* Europaeo exacte convenit. Tachyp. limbatus Grav., quem solis palpis acuminatis differre affert Grav., neque Tachyporum, neque speciem peculiarem esse mihi persuasum, quamvis a Dejean ipso idem Latreillianae collectionis specimen sub genere illo enumeratur.

3. Mas abdominis segmento inferiore quinto apice emarginato a reliquis huius sectionis speciebus discedit.

- \* 2. *T. pictus*: *Niger, nitidus, pedibus, ano, thoracis limbo elytrisque flavis, his macula rotundata media nigra.* — Long.  $1\frac{1}{3}$  lin

Habitat in Madagascar, Dom. Goudot.

Statura et summa affinitas praecedentis. Antennae capite thorace paulo longiores, apice vix crassiores, articulis 3 — 10 subobconicis, crassitie longioribus, ultimo elongato, cylindrico, apice obtuse acuminato, nigrae, articulis duobus primis flavis. Palpi flavi, apice picei. Caput sat magnum, convexum, laevissimum, nigrum, nitidum. Thorax coleopterorum latitudine, latitudine dimidio fere brevior, antrorsum parum angustatus, lateribus modice rotundatus, basi utrinque subsinuatus, angulis posterioribus retrorsum parum prominentibus, rotundatis, modice transversim convexus, laevissimus, nitidissimus, niger, margine anteriore et posteriore tenui limboque laterali lato flavis. Scutellum laeve, nigrum. Elytra thorace paulo longiora, modice convexa, angulo apicali exteriore oblique truncato, parcius subtilissime punctulata, nitida, flava, macula magna ovali media nigra, margine laterali medio infuscato. Abdomen crebre subtiliter punctatum, densius subtilissime pubescens, lateribus et apice parce nigro-pilosellum, nigrum, apice lato testaceo. Pectus nigrum, mesosterno subtiliter carinato. Pedes flavi.

Obs. Differt constanter a praecedente praecipue elytrorum macula media antrorsum haud elongata, basi nigredine nulla tincta, margine laterali medio solummodo infuscato, ano superne et inferne testaceo, elytris parcius punctulatis, thoracis angulis posterioribus paulo magis rotundatis. Sexus differentia omnino ut in illo.

- \* 3. *T. piceus*: *Leviter convexus, piceus, nitidus, capite, thorace elytrisque laevissimis, antennarum basi pedibusque testaceis, thoracis margine elytrorumque limbo apicali rufescentibus, his thorace paulo longioribus.* — Long. 1 lin.

Habitat in ins. Puerto Rico, Dom. Moritz.

Sequenti affinis, leviter convexus, piceus, nitidus. Antennae capitis thoracisque fere longitudine, apice crassiores, articulis 2 et 3 elongatis, longitudine subaequalibus, 5 — 10 praecedentibus distincte crassioribus, aequalibus, transversis, ultimo subgloboso, nigro-piceae, articulis 4 primis dilute testaceis. Caput parum convexus, laevissimum, oris partibus testaceis. Thorax basi coleopterorum basi haud latior, latitudine baseos duplo brevior, basi utrinque late leviterque sinuatus, angulis posterioribus retrorsum parum prominulis, acutiusculis, externe rotundatis, modice convexus, laevissimus, piceus, margine basali laterali dilutiore. Elytra thorace vix sesqui longiora, apicem versus sensim angustata, angulo apicali exteriore oblique truncato, secundum marginem lateralem profundius impressa, laevissima, margine laterali apicali dilutiore. Abdomen subtiliter punctulatum, tenuiter subtilissime pubescens, apice dilutius. Mesosternum subtiliter carinatum. Pedes rufo-testacei.

- \* 4. *T. cinctellus*: *Convexiusculus, nigro-piceus, nitidus, capite, thorace elytrisque laevissimis, antennarum basi apiceque pedibusque testaceis, thoracis elytrorumque margine laterali apicali piceis, his thorace plus sesqui longioribus.* — Long. 1 lin.

Habitat in Brasilia, Dom. Thorey.

Corpus parvum, convexiusculum, piceo-nigrum, nitidum. Antennae

capitis thoracisque longitudine, apicem versus sensim subincrassatae, articulis 2 et 3 leviter elongatis, sequentibus sensim paulo brevioribus et paululum crassioribus, subobconicis, ultimo breviter ovato, apice abrupte acuminato, fusco-nigrae, articulis 4 primis ultimoque testaceis. Caput laevissimum, ore rufo-piceo. Thorax basi coleopterorum basi haud latior, latitudine baseos duplo brevior, antrorsum fortius angustatus, lateribus rotundatis, basi bisinuatus, angulis posterioribus retrorsum leviter prominentibus, latis, rotundatis, convexus, laevissimus, margine laterali basaliq[ue] indeterminate piceo-rufescentibus. Elytra thorace plus sesqui longiora, apicem versus sensim angustata, angulo apicali exteriore oblique truncato, secundum marginem lateralem leviter impressa, laevissima, apice (ad tertiam elytri partem usque) margineque tenui laterali indeterminate piceo-rufis. Abdomen densius subtilissime pubescens, subtus apice piceum. Mesosternum subtiliter carinatum. Pedes rufo-testacei.

- \* 5. *T. pulchellus*: *Leviter convexus, nigro-piceus, nitidus, capite thorace elytrisq[ue] laevissimis, pedibus, thoracis limbo, elytris postice margineq[ue] laterali piceo-rufis, antennis basi testaceis.* — Long. 1 lin.

*Var.*: *Thorace elytrisq[ue] testaceis, illis disco his basi piceis.*

Habitat in Puerto Rico et in Columbia, Dom. Moritz.

Praecedentibus simillimus, modice convexus, nigro-piceus, nitidus. Antennae capitis thoracisque prope longitudine, apice paulo crassiores, articulis 2 et 3 elongatis, aequalibus, sequentibus sensim paulo brevioribus et crassioribus, ultimo ovato, apice subacuminato, nigro-piceae, basi testaceae. Palpi rufo-testacei. Caput leviter convexus, laevissimum, ore dilutiore. Thorax basi coleopterorum basi haud latior, latitudine baseos duplo brevior, antrorsum fortiter angustatus, lateribus rotundatus, basi leviter bisinuatus, angulis posterioribus retrorsum parum prominentibus, rotundatis, modice convexus, limbo basali lateraliq[ue] iam tenuiore iam latiore piceo-rufo. Elytra thorace sesqui longiora, apicem versus sensim angustata, angulo apicali exteriore oblique truncato, secundum marginem lateralem vix impressa, laevissima, apice lato (ad dimidiam fere elytri partem usque) margineq[ue] laterali testaceo-rufis. Abdomen confertim subtiliter punctulatum, dense subtiliter pubescens, apice piceum. Mesosternum subtiliter at acute carinatum. Pedes testacei.

Variat thorace elytrisq[ue] testaceis, illo disco, his basi piceis, vel interdum totus testaceus, capite, thoracis disco elytrorumq[ue] basi paulo saturatioribus.

- \* 6. *T. nitidulus*: *Leviter convexus, nigro-piceus, nitidus, capite, thorace elytrisq[ue] laevissimis, capite, thorace, pedibus antennisque testaceis, elytris abdomineq[ue] apice rufescentibus.* — Long.  $\frac{3}{4}$  lin.

Habitat in Puerto-Rico, Dom. Moritz.

Habitu omnino praecedentium, at paulo minor. Corpus leviter convexus. Antennae capitis thoracisque prope longitudine, apice paulo crassiores, articulis 2 et 3 elongatis, aequalibus, sequentibus sensim paulo brevioribus et crassioribus, penultimis crassitie sua brevioribus, ultimo breviter acuminato, testaceae, apice fuscae. Caput parum convexus, rufo-testa-



cum, laevissimum. Thorax basi coleopterorum basi haud latior, latitudine baseos duplo brevior, antrorsum fortiter angustatus, lateribus rotundatus, basi vix sinuatus, angulis posterioribus vix prominentibus rotundatis, leviter convexus, laevissimus, rufo-testaceus, nitidus. Scutellum nigro-piceum. Elytra thorace plus sesqui longiora, apicem versus sensim leviter angustata, angulo apicali exteriori obtuso, secundum marginem lateralem leviter impressa, laevissima, nitida, obscure fusco-picea, basi fere nigra, margine laterali apicalique indeterminate testaceo-rufescentibus. Abdomen subtiliter punctulatum, parce subtilissime pubescens, nigro-piceum, apice late testaceo-rufum. Pectus nigrum, mesosterno subtiliter carinato. Pedes rufo-testacei.

- \* 7. *T. convexus*: *Convexus, piceo-niger, nitidus, capite thoraceque laevissimis, elytris parce subtilissime punctatis, antennarum basi pedibusque rufis.* — Long. 2 lin.

Habitat in Brasilia, Dom. Melly.

Corpus convexum, nigro-piceum, nitidum. Antennae capite thoraceque paulo breviores, apice parum crassiores, articulis 2 et 3 leviter elongatis, sequentibus sensim paululum brevioribus et crassioribus, subobconicis, ultimo ovato, fuscae, articulis tribus primis rufo-testaceis. Caput laevissimum fronte convexiuscula, ore rufo, palpis testaceis. Thorax basi coleopterorum basi latitudine aequalis, latitudine baseos duplo brevior, basi utrinque sinuatus, antrorsum valde angustatus, lateribus fortiter rotundatus, angulis posterioribus postice leviter prominentibus, acutiusculis, acumine rotundato, valde convexus, laevissimus, margine basali laterali summo piceo-rufescente. Elytra thorace sesqui prope longiora, apicem versus sensim leviter angustata, valde convexa, angulo apicali exteriori obtusiusculo, secundum marginem lateralem haud impressa, convexa, minus dense subtilissime punctata. Abdomen subtiliter punctulatum, parce subtilissime pubescens, piceum, apice segmentorumque singulorum margine postico rufescente. Mesosternum fortiter acuminatum, valde prominens. Pedes rufi.

- \* 8. *T. elatus*: *Convexus, piceo-niger, nitidus, capite thoraceque laevissimis, hoc lateribus castaneo, elytris parce subtilissime punctatis, ore, antennarum basi pedibusque rufo-testaceis.* — Long. 2 lin.

Habitat Novae Valenciae in Columbia, Dom. Moritz.

Statura et summa similitudo praecedentis, fortius convexus, piceo-niger, nitidus. Antennae capite thoraceque paulo breviores, tenuiores, apice vix incrassatae, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus et paululum crassioribus, subobconicis, ultimo subovato, apice abrupte acuminato, nigrae, articulis tribus primis testaceis. Caput laevissimum, fronte parum convexa, ore testaceo. Thorax basi coleopterorum basi haud, latitudine baseos duplo fere brevior, antrorsum valde angustatus, lateribus fortiter rotundatus, basi utrinque levissime sinuatus, angulis posterioribus retrorsum leviter prominentibus, acutis, sat convexus, laevissimus, nitidus, dorso piceo-niger, margine laterali indeterminate castaneo. Elytra thorace tertia fere parte longiora, apicem versus sensim leviter angustata, angulo apicali exteriori rotundato, secundum marginem lateralem haud impressa, valde convexa, parcius subtiliter punctata. Abdomen crebre

subtilius punctulatum, parce subtilissime pubescens, segmentis singulis rufo-marginatis. Mesosternum acute carinatum. Pedes rufi.

Obs. T convexo simillimus, at non adeo convexus et antennis in utroque sexu longioribus tenuioribusque, mesosternoque minus prominenter carinato distinctus.

- \* 9. *T. hepaticus*: *Convexus, piceo-castaneus, nitidus, capite thoraceque laevissimis, elytris vage subtilissime punctatis, antennis basi rufis, apice ferrugineis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

T. convexo dimidio minor et paulo minus convexus, piceo-castaneus, nitidus. Antennae capitis thoracisque fere longitudine, apice leviter crassiores, articulo tertio secundo paululum longiore et tenuiore, 4 — 10 sensim paulo brevioribus et crassioribus, subobconicis, ultimo oblongo, praecedentibus ambobus prope longiore, subcylindrico, apice abrupte acuminato, fuscae, articulis tribus primis rufo-testaceis, ultimo ferrugineo. Palpi rufo-testacei. Caput vix convexum, laevissimum, nigrum, ore rufo-piceo. Thorax basi coleopterorum basi vix latior, latitudine duplo brevior, antrorsum fortiter angustatus, lateribus modice rotundatus, basi leviter bisinuatus, angulis posterioribus retrorsum prominentibus, acutiusculis, acumine extrorsum rotundato, modice convexus, laevissimus. Elytra thorace vix tertia parte longiora, apicem versus sensim leviter angustata, angulo apicali exteriori obtuso, subrotundato, modice convexa, minus dense subtilissime punctata. Abdomen subtiliter punctulatum, omnium subtilissime parce pubescens. Mesosternum fortiter acuteque carinatum, modice prominens. Pedes rufo-testacei.

- \* 10. *T. brunneus*: *Convexus, piceus, nitidus, capite thoraceque laevissimis, elytris vage subtilissime punctatis, antennarum basi pedibusque rufis.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Columbia, Dom. Moritz.

Praecedenti affinis, at dimidio minor et paulo magis convexus. Antennae capite thoraceque breviores, tenuiores, apice leviter crassiores, articulo tertio secundo tenuiore at vix brevior, sequentibus sensim paulo brevioribus et crassioribus, ultimo subovato, apice abrupte acuminato, fuscae, articulis tribus primis rufo-testaceis. Palpi rufo-testacei. Caput leviter convexum, laevissimum, nigro-piceum, ore rufo-piceo. Thorax basi coleopterorum basi paulo latior, latitudine vix duplo brevior, antrorsum fortiter angustatus, lateribus fortius rotundatis, basi utrinque leviter sinuatus, posterioribus retrorsum leviter prominentibus, acutis, externe rotundatis, sat convexus, laevissimus, nitidus, saturate piceo-castaneus, dorso indeterminate piceo-nigro. Scutellum nigro-piceum, nitidum. Elytra thorace tertia fere parte longiora, apicem versus sensim leviter angustata, angulo apicali exteriori obtuso, secundum marginem lateralem vix impressa, convexa, minus confertim subtilissime punctata, saturate fusco-picea, nitida, margine laterali apicalique summo castaneo. Abdomen subtiliter punctulatum, vix perspicue pubescens, piceum, nitidum, subtus apice late rufo-piceum. Mesosternum subtiliter carinatum. Pedes testaceo-rufi.

Obs. Differt a praecedente statura minore, magis convexa, thorace paulo longiore, convexiore, antennis tenuioribus, apice haud ferrugineis, mesosterno subtiliter carinato, haud prominente.

- \* 11. *T. terminalis*: *Convexus, nigro-piceus, nitidus, capite thoraceque laevissimis, elytris crebre subtiliter punctatis, pedibus antennisque rufis, his apice ferrugineis.* — Long.  $1\frac{1}{3}$  lin.

Habitat in ins. Puerto Rico, Dom. Moritz.

Statura omnino praecedentis. Antennae capitis thoracisque prope longitudine, apice leviter crassiores, articulo tertio secundo paulo longiore et tenuiore, sequentibus sensim paulo brevioribus at vix crassioribus, ultimo praecedentibus ambobus longitudine subaequali, oblongo-ovato, apice abrupte acuminato, piceo-rufae, articulis tribus primis rufo-testaceis, ultimo flavo-testaceo. Palpi rufo-testacei. Caput leviter convexum, laevissimum, piceo-nigrum, nitidum, ore piceo-subrufo. Thorax basi coleopterorum basi paulo latior, latitudine duplo fere brevior, antrorsum fortiter angustatus, lateribus modice rotundatus, basi utrinque sinuatus, angulis posterioribus retrorsum leviter prominentibus, acutiusculis, externe rotundatis, convexus, laevissimus, nigro-piceus, marginibus dilutioribus. Scutellum nigro-piceum. Elytra thorace tertia fere parte longiora, apicem versus sensim leviter angustata, angulo apicali exteriori obtuso, secundum marginem lateralem obsolete impressa, convexa, densius subtiliter punctata, nigro-picea, nitida, marginibus dilutioribus, parce subtilissime brevissimeque pubescens. Abdomen parcius subtiliter punctulatum, piceum, subtus apicem versus rufescens. Mesosternum subtiliter carinatum. Pedes testaceo-rufi.

Obs. Praecedentibus admodum affinis, elytris autem crebrius minusque subtiliter punctatis distinctus.

- \* 12. *T. apicalis*: *Convexiusculus, nigro-piceus, nitidus, capite thoraceque laevissimis, elytris subtilissime punctatis, pedibus antennarumque basi testaceis, his articulo apicali flavo.* — Long. 1 lin.

Habitat in ins. Puerto Rico, Dom. Moritz.

Statura omnino *T. picei* et vix paululum maior, praecedente plus dimidio minor, modice convexus, piceo-niger, nitidus. Antennae capite thoraceque paulo breviores, apicem versus sensim leviter incrassatae, articulo tertio secundo tenuiore et paulo longiore, sequentibus sensim brevioribus et paulo crassioribus, penultimus transversis, ultimo breviter ovato, apice acuminato, nigrae, articulis 4 primis testaceis, ultimo flavo. Caput minusculum, fronte parum convexa, ore dilute testaceo. Thorax basi coleopteris vix latior, latitudine triplo prope brevior, antrorsum fortius angustatus, lateribus leviter rotundatis, basi truncatus, angulis posterioribus haud prominulis, acumine summo rotundatis, laevissimus, margine summo laterali brunneo. Elytra thorace duplo prope longiora, apicem versus sensim angustata, angulis posterioribus obtusiusculis, secundum marginem lateralem haud impressa, omnium subtilissime punctulata. Abdomen parce omnium subtilissime punctulatum, vix pubescens, apice rufescens. Mesosternum subtiliter carinatum. Pedes dilute testacei.

Obs. Differt a praecedente statura multo minore, elytris omnium subtilissime, vix perspicue punctulatis etc. a *T. piceo* capite maiore, antennarum articulo ultimo flavo, elytris subtilissime punctulatis.



- \* 13. *T. humilis*: *Parum convexus, nigro-piceus, nitidus, capite thoraceque laevissimis, elytris subtilissime punctulatis, ore testaceo, antennarum basi pedibusque rufis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Brasilia, Dom. Prof. Germar.

Corpus leviter convexum, nigro-piceum, nitidum, Antennae capitis thoracisque longitudine, crassiusculae, apice sensim crassiores, articulo tertio secundo evidenter longiore, sequentibus sensim paulo brevioribus at vix crassioribus, ultimo oblongo, praecedentibus ambobus longitudine aequali, apice sat fortiter acuminato, fuscae, articulis tribus primis rufis. Caput parum convexum, laevissimum, ore testaceo. Thorax basi coleopteris parum lator, latitudine plus duplo brevior, antrorsum fortiter angustatus, lateribus modice rotundatus, basi utrinque vix sinuatus, angulis posterioribus retrorsum haud prominulis, acumine summo rotundatis, leviter transversim convexus, laevissimus, nigro-piceus, marginibus dilutioribus. Elytra thorace duplo prope longiora, apicem versus sensim leviter angustata, angulo apicali exteriore obtusiusculo, secundum marginem lateralem canaliculata, subtilissime punctulata. Abdomen subtilissime punctulatum, vage tenuissime subtilissimeque pubescens. Pedes testaceo-rufi.

Obs. Distinctus a praecedentibus statura minus convexa thoraceque brevior, basi vix sinuato, angulis posterioribus postice haud prominentibus.

14. *T. brevicollis*: *Subdepressus, nigro-piceus, nitidus, capite thoraceque laevissimis, elytris omnium subtilissime punctulatis, abdomine evidenter punctato, antennarum basi pedibusque testaceis.* — Long.  $1\frac{1}{2}$  lin.

Habitat Novae Valentiae in Columbia, Dom. Moritz.

Statura et summa affinitas praecedentis, perparum convexus, nigro-piceus, nitidus. Antennae capitis thoracisque prope longitudine, apicem versus leviter incrassatae, articulo tertio secundo paululum longiore, 4 — 10 sensim paulo brevioribus at vix crassioribus, ultimo ovato, apice abrupte acuminato, nigrae, articulis tribus primis testaceis. Oris partes testaceae. Caput laevissimum, fronte parum convexa. Thorax basi coleopteris parum lator, latitudine baseos triplo fere brevior, antrorsum fortiter angustatus, lateribus leviter rotundatus, basi utrinque subsinuatus, angulis posterioribus retrorsum prominentibus, externe rotundatis, leviter transversim convexus, laevissimus, nigro-piceus, marginibus rufescentibus. Elytra thorace plus sesqui longiora, apicem versus sensim leviter attenuata, angulo apicali exteriore obtusiusculo, secundum marginem lateralem canaliculata, subtilissime parcius punctulata. Abdomen crebre distincteque licet sat subtiliter punctulatum, tenuissime subtilissimeque pubescens, apice piceo-testaceum. Mesosternum subtiliter carinatum. Pedes rufo-testacei.

Obs. *T. humili* proximus, abdomine autem crebre evidenter punctato satis differt.

- \* 15. *T. depressus*: *Subdepressus, niger, nitidus, capite, thorace elytrisque confertim subtiliter punctatis, his piceo-castaneis, antennarum basi pedibusque testaceis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Corpus parum convexum, nigrum, nitidum. Antennae capite thoraceque paulo longiores, apice vix crassiores, articulo secundo tertioque aequalibus,

sequentibus sensim paulo brevioribus, at vix crassioribus, ultimo oblongo-ovato, apice abrupte acuminato, fusco-nigrae, articulo primo toto, secundo tertioque basi testaceis. Palpi testacei. Caput parum convexum, subtilissime punctulatum. Thorax basi coleopterorum basi vix latior, latitudine plus duplo brevior, antrorsum fortiter angustatus, lateribus leviter rotundatus, basi utrinque leviter sinuatus, angulis posterioribus retrorsum modice prominentibus, acutiusculis, externe rotundatis, parum convexus vel subdepressus, confertim subtiliter punctulatus, margine postico piceo-pellucente. Elytra thorace sesqui longiora, apicem versus sensim subangustata, angulo apicali exteriore oblique truncato, subdepressa, secundum marginem lateralem longitudinaliter leviter impressa, confertim subtiliter punctulata, piceo-castanea, nitida, lateribus nigra. Abdomen crebre subtiliter punctatum, tenuissime subtilissimeque pubescens, segmentis ultimis duobus apice piceo-castaneis. Mesosternum subtiliter carinatum. Pedes rufo-testacei.

- \* 16. *T. melanarius*: *Parum convexus, niger, nitidus, antennarum basi pedibusque rufo-piceis, capite thoraceque subtilissime punctulatis, elytris confertim subtiliter punctatis.* — Long. 1 lin.

Habitat in Bengala, Dom. Hope.

*T. gibbulus* minor et multo minus convexus, fere subdepressus, niger, nitidus. Antennae capitis thoracisque prope longitudine, apice vix crassiores, articulo secundo tertioque aequalibus, obconicis, sequentibus inter se subaequalibus, leviter compressis, ultimo paulo maiore, subovato, compresso, apice abrupte acuminato, nigricantes, basi piceae. Caput parum convexum, subtilissime punctulatum, oris partibus piceis. Thorax basi coleopterorum basi latitudine aequalis, apice duplo angustior, latitudine baseos duplo brevior, lateribus sat fortiter rotundatus, basi leviter bisinuatus, angulis posterioribus retrorsum leviter prominentibus, acutis, leviter transversim convexus, subtilissime punctulatus, margine baseos summo rufescente. Scutellum subtiliter punctulatum. Elytra thorace plus sesqui longiora, apicem versus sensim subangustata, angulo apicali exteriore oblique truncato, parum convexa, nigra, nitida, confertim subtiliter punctata, secundum marginem lateralem canaliculata, margine apicali summo piceo-rufescente. Abdomen crebre subtiliter punctulatum, segmentis anterioribus apice piceo-marginatis, subtus apice rufo-piceum. Pedes rufo-picei.

- \* 17. *T. gibbulus*: *Convexus, piceo-niger, nitidus, capite thoraceque laevissimis, elytris confertim subtiliter punctatis, ore antennarum pedibusque rufis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in America septentrionali, Dom. Zimmermann. Nominis *Tachypori gibbuli* etiam a Com. Dejean communicatus.

Corpus convexus, nigrum, nitidum. Antennae capitis thoracisque prope longitudine, apice sensim crassiores, articulo tertio secundo tenuiore et prope brevior, 3—5 obconicis, sensim brevioribus et crassioribus, 6—10 sensim magis transversis, ultimo breviter ovato, apice acuminato, totae rufae. Caput leviter convexus, laevissimum, ore cum palpis rufo-testaceo. Thorax basi coleopterorum basi parum latior, latitudine duplo brevior, antrorsum fortiter angustatus, lateribus fortiter rotundatus, basi utrinque leviter sinuatus,

angulis posterioribus retrorsum modice prominentibus, acutiusculis, externe rotundatis, fortius convexus, laevisimus, margine postico lateraliq[ue] piceo-pellucente. Elytra thorace sesqui longiora, apicem versus sensim subangustata, angulo apicali exteriore subrecto, secundum marginem lateralem longitudinaliter modice impressa, crebrius subtiliter punctata, margine laterali et apicali indeterminate rufo-piceis. Abdomen subtilissime punctulatum, vix pubescens, segmentis singulis margine apicali piceo. Pedes testaceo-rufi.

\* 18. *T. rutilus*: *Leviter convexus, rufus, nitidus, capite thoraceque laevissimis, elytris parcius subtiliter punctatis.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Americae ins. Puerto Rico et St. Thomae, in Columbia, Dom. Moritz.

Statura omnino *T. gibbuli*, totus testaceo-rufus, nitidus. Antennae capitulis thoracisque prope longitudine, apicem versus leniter crassiores, articulo tertio secundo tenuiore et paulo brevior, 4—10 sensim paulo brevioribus et crassioribus, subobconicis, ultimo crassiore, subovato, apice abrupte acuminato, testaceo, medio fuscescentes. Caput laevissimum. Thorax basi coleopteris paulo latior, latitudine duplo brevior, antrorsum fortiter angustatus, lateribus modice rotundatus, utrinque leviter sinuatus, angulis posterioribus retrorsum leviter prominentibus, rotundatis, satis convexus, laevisimus. Elytra thorace sesqui longiora, apicem versus sensim leviter angustata, angulo apicali exteriore obtusiusculo, secundum marginem lateralem medio impressa, parcius subtiliter obsoleteq[ue] punctata. Abdomen subtiliter punctulatum, tenuissime subtilissimeq[ue] pubescens. Mesosternum subtiliter carinatum. Pedes rufo-testacei.

## Fam. II.

*Mesosternum simplex.*

*Tarsi antici maris dilatati.*

Corpus plerumque maius, magis oblongum et minus convexum. Palpi maxillares articulo ultimo subcylindrico, apice acuminato, saepius penultimo duplo longiore. Mesosternum simplex. Tarsi posteriores articulis duobus primis subaequalibus.

Mas tarsis anticis articulis 3 primis dilatatis, abdominis segmento superiore sexto apice 4-dentato, inferiore incisione profunda bifido, lobis acuminatis, saepius compressis, septimo conspicuo, quinto inferiore apice magis minusve emarginato.

Femina tarsis anticis simplicibus, abdominis segmento sexto superiore vel trifido vel quadrifido, laciniis lateralibus profundius divis, inferiore apice 6-fido, laciniis intermediis duabus obtusioribus, apice setulis rigidis fulvis ciliatis, reliquis acuminatis, lateralibus brevioribus, nisi segmento fortiter exserto observandis, intermediis 4 longitudine subaequalibus.

## Observationes.

1. Caput, thorax et elytra magis minusve subtiliter, saepius subtilissime punctata, interstitiis nonnunquam omnium subtilissime alutaceis, glaberrima, abdomen omnium subtilissime tenuissimeq[ue] pubescens, pube omnium brevissima, apice vix unquam pilosellum. Scutellum medio cre. Elytra lateribus subtiliter marginata, angulo apicali exteriore subrotundato.



2. Dom. Prof. Gravenhorst primo secundum palporum structuram Tachinum a Tachyporo distinxit, dein corporis habitum magis respiciens, Boletobium et Mycetoporum sub hoc nomine coniunxit, quod auctores recentiores generi nostri restituerunt. Habitu generali, coxarum posticarum et ligulae figura magis ad Tachyporum accedit, palporum autem et tarsorum structura facile distinguitur.

3. Victus plerisque in fimo. Europae et Americae septentrionalis regionibus temperatis et frigidis huius sectiones species propriae videntur, intra tropos etiam in Europa australi vix obviae.

✕ \* 19. *T. rufipes*: Convexus, piceo-niger, nitidus, antennarum basi pedibusque rufis, elytris thorace sesqui longioribus, puncto humerali margineque apicali picescentibus. — Long. 3 lin.

*Er. Col. March* I. 397. 2.

*Staphyl rufipes* Degeer *Ins.* IV. 24. 11. t. 1. f. 14. — *Fab. Syst. Ent.* 267. 17. *Spec. Ins.* I. 337. 22. *Mant. Ins.* 222. 31. — *Payk. Mon. Staph.* 75. 53. *Faun. Suec.* III. 418. 65. — *Walk. Faun. Paris.* I. 273. 20.

*Staph. rufipes* Schrank *Fauna Boica* I. 644. 874. (forte)

*Oxyporus rufipes* *Fab. Ent. Syst.* I. II. 235. 21. *Syst. El.* II. 607. 21. — *Cederh. Faun. Ingr.* 121. 372. — *Panz. Ent. Germ.* 361. 16.

*Oxyporus rufipes* *Panz. Faun. Germ.* 27. 20. (male)

*Tachinus pullus* *Grav. Micr.* 140. 6. — *Latr. Hist. nat. Crust. et Ins.* IX. 395. 6. — *Gyll. Ins. Suec.* II. 262. 10. IV. 472. 10. — *Zetterst. Faun. Lappon.* I. 62. 4. *Ins. Lappon.* 57. 4. — *Sahlb. Ins. Fenn.* I. 299. 8. — *Mannerh. Brachelytr.* 61. 8.

*Tachyporus pullus* *Grav. Mon.* 16. 30.

*Tachinus signatus* *Grav. Micr.* 139. 5. — *Latr. Hist. nat. Crust. et Ins.* IX. 395. 5. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 508. 3.

*Tachyporus signatus* *Grav. Mon.* 17. 32.

*Var. a*: Thoracis limbo laterali elytrisque castaneo-testaceis, his dorso subinfuscat. (Minus maturus.)

*Tachinus laticollis* *Gyll. Ins. Suec.* IV. 471. 5—6. — *Sahlb. Ins. Fenn.* 297. 4. — *Mannerh. Brachelytr.* 61. 5. (forte.)

*Var. b*: Rufo-testaceus, capite thoracisque disco fusco-piceis. (Immaturus.)

*Tachinus pallens* *Gyll. Ins. Suec.* II. 259. 7. — *Sahlb. Ins. Fenn.* I. 298. 6.

Habitat in Europa.

Piceo-niger, nitidus. Antennae capite thoraceque paulo longiores, graciles, apice vix crassiores, articulo tertio secundo paululum longiore, 5—10 elongato-obconicis, subaequalibus, ultimo oblongo-ovato, subacuminato, apice compresso, nigri-picei, articulo primo rufo-testaceo, sequentibus tribus rufo-piceis. Palpi rufo-picei. Caput convexum, omnium subtilissime punctulatum, ore concolore. Thorax coleopteris lator, latitudine baseos quarta parte brevior, antrorsum modice angustatus, apice profundius emarginatus, lateribus fortiter rotundatus, basi truncatus, angulis anterioribus modice prominentibus posterioribusque obtusis, subrotundatis, convexus, omnium subtilissime punctulatus, niger, nitidus, margine summo laterali saepe piceo. Scutellum parce punctatum, nigrum. Elytra thorace vix sesqui longiora, confertissime subtilissimeque alutaceo-strigosa, subtiliter vage punctulata, nigro-picea, puncto humerali margineque summo apicali dilutioribus. Abdomen distinctius haec subtiliter crebrius punctatum, pube brevissima subtilissimaque

adpersum, nigrum, segmentis singulis margine summo apicali rufo-piceo. Pedes rufi.

Mas abdominis segmento superiore sexto quadridentato, dentibus acutis, fere spiniformibus, intermediis magis prominentibus, segmentis inferioribus 3—5 subimpressis, quinto apice leviter triangulariter emarginato, margine intra sinum spongioso, sexto bifido, laciniis vix elongatis, compressis, leviter curvatis, acuminatis.

Femina abdominis segmento superiore sexto trifido, laciniis intermediis longioribus, spiniformibus, intermedio lato apice lanceolato-acuminato.

Obs. *T. pallens* Gyllenh. secundum exemplum a Dom. Schüppel communicatum, ab autore ipso missum, ad *T. humeralem* quidem pertineat, licet specimenibus genuinis minor, descriptio Gyllenhali autem potius huic speciei adaptanda mihi videtur, praecipue ob caput et thoracem multo subtilius punctata.

\* 20. *T. flavipes*: *Piceo-niger, pedibus elytrisque brunneo-ferrugineis, his thorace sesqui longioribus, dorso plerumque infuscat, antennis basi nigricantibus.* — Long. 3 lin.

*Er. Col. March. I. 398. 3.*

*Oxyporus flavipes* Fab. *Syst. Ent.* 268. 3. *Spec. Ins. I.* 339. 5. *Mant. Ins. I.* 222. 6. *Ent. Syst. I. II.* 534. 14. *Syst. El. II.* 606. 13. — *Panz. Ent. Germ.* 360. 11. *Staphyl rufipes* Marsh. *Ent. Brit.* 518. 60.

*Tachinus rufipes* Grav. *Micr.* 137. 4. — *Latr. Hist. nat. Crust. et Ins. IX.* 294. 4. *Gen. Crust. et Ins. I.* 299. 1. — *Gyll. Ins. Suec. II.* 260. 9. — *Zetterst. Faun. Lappon. I.* 62. 3. *Ins. Lappon.* 57. 3. — *Sahlb. Ins. Fenn. I.* 298. 7. — *Mannerh. Brachelytr.* 61. 7. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 507. 2. — *Runde Brachelytr. Hal.* 26. 3.

*Tachyporus rufipes* Grav. *Mon.* 17. 15.

*Var. Thoracis limbo laterali latiore castaneo-testaceo.*

*Tachinus castaneus* Grav. *Micr.* 140. 7. — *Latr. Hist. nat. Crust. et Ins. IX.* 395. 7.

*Tachyporus fimetarius* var. Grav. *Mon. p.* 15.

*Tachyporus rufipes?* Grav. *Mon. p.* 18.

*Tachinus dubius* Gyll. *Ins. Suec. II.* 258. 6. — *Sahlb. Ins. Fenn. I.* 298. 5. — *Mannerh. Brachelytr.* 61. 6. — *Runde Brachelytr. Hal.* 25. 2.

Habitat in Europa.

Statura fere *T. rufipedis*, at minus convexus. Antennae capitis thoracisque longitudine, apice paulo crassiores, articulo tertio secundo paulo longiore, 5—10 subobconicis, sensim paululum brevioribus, penultimis crassitie subbrevioribus, ultimo breviter ovato, apice compresso, piceae, basi nigricantes. Palpi nigro-picei. Caput confertim subtiliter punctulatum, nigrum, ore concolore. Thorax coleopteris vix latior, latitudine baseos tertia parte brevior, antrorsum leviter angustatus, apice late et parum profunde emarginatus, basi truncatus, angulis anterioribus leviter prominentibus posterioribusque rotundatis, parum convexus, confertim subtiliter punctatus, nitidus, niger, limbo tenui basali lateraliq. testaceo. Scutellum piceum, subtiliter punctulatum. Elytra thorace plus sesqui longiora, confertim subtiliter punctata, castaneo-testacea, dorso magis minusve infusato, praecipue in femina. Abdomen minus confertim subtiliter punctatum, pube brevissima subtilissimaque adpersum, nigrum, nitidum, segmentis singulis margine summo apicali

piceo-testaceo. Pectus nigrum. Pedes rufo-testacei, femoribus basi nonnunquam picescentibus.

Mas abdominis segmento superiore sexto quadridentato, dentibus acutiusculis, intermediis multo magis prominentibus; segmento inferiore quinto apice leviter emarginato, margine intra sinum spongioso, sexto profunde inciso, laciniis triangularibus, acuminatis.

Femina abdominis segmento superiore sexto trifido, laciniis longitudine aequalibus, lateralibus spiniformibus, intermedia latiore, sensim angustata, acute acuminata.

Variat thoracis limbo laterali latius testaceo, elytris dorso haud aut parum infuscatis. Feminae semper obscuriores.

Obs. 1. Differt a *T. rufipede* antennis basi nigricantibus, statura minus convexa, capite, thorace elytrisque distinctius punctatis.

2. *Tachinus intricatus* Sahlb. Ins. Fenn. I. 300. 9. absque dubio ad hanc speciem pertinet: elytrorum sculptura „punctis in lineolas confuse intricatas aliquatenus ordinatis” certe fortuita.

\* 21. *T. frigidus*: *Niger, nitidus, antennarum basi, palpis, pedibus thoracisque limbo laterali rufo-ferrugineis, elytris thorace sesqui longioribus, piceis, puncto humerali margineque apicali ferrugineis.* — Long.  $2\frac{1}{2}$  lin.

Habitat in ins. Unalaschka, Dom. v. Chamisso.

*Niger, nitidus.* Antennae capitis thoracisque longitudine, apicem versus sensim leviter incrassatae, articulo tertio secundo paulo longiore, 5—10 subobconicis, sensim paulo brevioribus et crassioribus, ultimo ovali, acuminato, apice compresso, fuscae, basi rufae, articulo primo dilutiore. Palpi toti rufo-testacei. Caput subtiliter punctulatum. Thorax coleopterorum latitudine, latitudine baseos dimidio brevior, antrorsum leviter angustatus, apice profundius emarginatus, lateribus rotundatus, basi truncatus, angulis anterioribus leviter prominentibus posterioribusque subrotundatis, modice convexus, confertim subtiliter punctatus, margine postico tenui laterali que latiore testaceo. Scutellum subtilissime punctulatum, nigrum. Elytra thorace sesqui longiora, dense subtilius punctata, vel nigro-vel rufo-picea, puncto humerali margineque apicali ferrugineis. Abdomen distinctius punctatum, pube parca subtilissima brevissimaque adspersum, ano subtus rufo-piceo. Pedes testacei.

Mas abdominis segmento superiore sexto quadridentato, dentibus acutiusculis, intermediis fortiter prominentibus; segmentis inferioribus 3—5 subimpressis, quinto apice leviter emarginato, margine intra sinum spongioso, sexto bifido, laciniis minus elongatis, compressis, modice curvatis, acuminatis.

Femina latet.

Obs. Affinis *T. pallipedi*, *rufipedi* et *flavipedi*, ab hoc antennis basi rufis, ab illis statura minus convexa et capite thoraceque crebre evidentius punctatis distinctus.

X \* 22. *T. humeralis*: *Niger, nitidus, antennarum basi, pedibus thoracisque limbo rufo-ferrugineis, elytris thorace plus sesqui longioribus, rufo-ferrugineis, dorso magis minusve fusciscentibus.* — Long.  $3\frac{1}{3}$  lin.

Grav. Micr. 130. 3. var. 2. 3. — Latr. Hist. nat. Crust. et Ins. IX. 394. 3.



— Zetterst. Faun. Lappon. I. 62. 2. Ins. Lappon. 56. 2. — Sahlb. Ins. Fenn. I. 296. 3. — Mannerh. Brachelytr. 61. 4. — Boisd. et Lacord. Faun. Ent. Paris. I. 507. 1. — Runde Brachelytr. Hal. 25. 1.

*Tachyporus humeralis* Grav. Mon. 19. 36.

*Tachinus humeralis* var. *b* Gyll. Ins. Suec. II. 257. 5.

*Staphyl. cinctus* Marsh. Ent. Britt. 519. 61.

Habitat in Europa.

*T. rufipede* maior, latior et minus convexus. Antennae capite thorace paulo longiores, apice parum crassiores, articulo tertio secundo paulo longiore, 5—10 aequalibus, obconicis, ultimo fusiformi, nigro-fuscae, basi rufae. Palpi rufi. Caput crebre subtiliter punctatum, nigrum, nitidum. Thorax coleopteris vix latior, latitudine baseos dimidio brevior, antrorsum modice angustatus, apice minus profunde emarginatus, lateribus rotundatus, basi utrinque subsinuatus, angulis anterioribus leviter prominentibus posterioribusque subrotundatis, leviter convexus, crebre subtiliter punctatus, nitidus, disco nigro, limbo antico posticoque angustioribus lateralique latiore rufo-ferrugineis. Elytra thorace duplo prope longiora, confertim subtiliter punctata, nitidula, ferruginea, dorso magis minusve infusata. Abdomen crebre distinctius punctatum, pube subtilissima brevissimaque parcius adpersum, nigrum, nitidum, segmentis singulis margine postico piceo. Pectus nigrum, nitidum. Pedes dilute rufi.

Mas abdominis segmento superiore sexto quadridentato, dentibus omnibus obtusis, subrotundatis, intermediis magis prominentibus; segmento inferiore quinto medio late impresso, apice emarginato, margine intra sinum retuso et spongioso, sexto bifido, laciniis modice elongatis, compressis, acuminatis, parum curvatis.

Femina abdominis segmento superiore sexto trifido, laciniis lateralibus paulo longioribus, spiniformibus, intermedia lata, apice triangulariter acuminata.

\* 23 *T. picipes*: *Piceo-niger*, nitidus, ore antennarumque basi rufis, pedibus piceis, elytris thorace sesqui longioribus. — Long.  $3\frac{1}{2}$  lin.

Habitat in America septentrionali, Dom. Zimmermann.

Habitu omnino *T. rufipedis*, at duplo maior, *T. fimbriati* minoribus individuis aequalis, piceo-niger, nitidus, capite, thorace, scutello elytrisque omnium confertissime subtilissimeque alutaceo-strigosis. Antennae capite thorace paulo breviores, crassiusculae, articulo tertio secundo duplo prope longiore, 5—10 crassioribus, crassitie aequalibus, sensim paulo brevioribus, penultimis fere transversis, ultimo ovato, apice compresso, nigrae, articulis 4 primis rufis, ultimo ferrugineo. Caput subtilissime punctulatum, ore rufo. Thorax coleopterorum latitudine, latitudine baseos tertia parte brevior, antrorsum modice angustatus, apice minus profunde emarginatus, lateribus modice rotundatus, basi utrinque leviter sinuatus, angulis anterioribus leviter prominentibus, subrotundatis, posterioribus subrectis, modice convexus, minus confertim subtilissime punctulatus, margines versus nonnunquam picescens. Scutellum piceum, laeve. Elytra thorace plus sesqui longiora, vage subtiliter punctata. Abdomen crebre subtiliter punctatum. Pedes picei, antici rufescentes, posteriores nigricantes.

Mas abdominis segmento superiore sexto quadridentato, dentibus lateralibus obtusiusculis, intermediis parum magis prominentibus, acutis; segmento inferiore quinto late at parum profunde triangulariter impresso, apice emarginato et utrinque obtuse bidentato, sexto bifido, laciniis modice elongatis, gracilioribus, compressis, leviter curvatis, acuminatis.

Femina abdominis segmento superiore sexto quadrifido, laciniis longitudine aequalibus, acuminatis, intermediis vix ad segmenti medium usque divisis.

Variat immaturus corpore testaceo, capite obscuriore.

- \* 24. *T. memnonius*: *Nigro-piceus, opacus, ore, antennarum basi pedibusque rufis, thoracis limbo elytrorumque puncto humerali rufo-piceis, his thorace plus sesqui longioribus.* — Long. 3 lin.

*Tachinus memnonius* Grav. Micr. 192. 3. (Mas)

*Tachyporus memnonius* Grav. Mon. 17. 31.

*Tachinus batychrous* Grav. Micr. 191. 1. (Fem.)

*Tachyporus batychrous* Grav. Mon. 20. 37.

Habitat in America septentrionali.

Statura omnino *T. flavipedis*, nigro- aut fusco-piceus, opacus, capite, thorace, scutello elytrisque omnium confertissime subtilissimeque alutaceo-strigosis. Antennae capitis thoracisque prope longitudine, extrorsum paululum crassiores, articulo tertio secundo paulo longiore, 5—10 longitudine subaequalibus, at sensim paululum crassioribus, obconicis, ultimo subovato, apice acuminato, fuscae, articulis 4 primis rufis. Caput impunctatum, nigrum, ore late rufo. Thorax coleopteris vix latior, latitudine baseos dimidio brevior, antrorsum modice angustatus, apice leviter emarginatus, lateribus rotundatus, basi truncatus, angulis anterioribus leviter prominentibus, rotundatis, posterioribus obtusis, subrotundatis, parum convexus, parce obsoleteque subtilissime punctatus, margine omni piceo-rufescente. Elytra thorace duplo prope longiora, subtilissime vage punctata, opaca, macula maiore minoreve humerali, saepe obsoleta, margineque summo apicali rufo-piceis. Abdomen distinctius crebre punctatum, parce subtilissime brevissimeque pubescens, segmentis singulis margine postico piceo-rufescente. Pedes testaceo-rufi.

Mas abdominis segmento superiore sexto acute quadridentato, dentibus intermediis longioribus et magis prominentibus; segmento inferiore quinto late triangulariter impresso, impressione spongiosa, apice late profundius emarginato, intra sinum utrinque setis rigidis densis vel contiguis, deorsum curvatis, ciliato, sexto bifido, laciniis elongatis, validis, compressis, curvatis, acuminatis.

Femina abdominis segmento superiore sexto trifido, laciniis lateralibus paulo longioribus, linearibus, intermedia latiore, canaliculata, apice bifida, laciniis valde acuminatis.

- \* 25. *T. fimbriatus*: *Niger, nitidus, pedibus piceis, ore antennarumque basi rufis, elytris substriato-punctatis, castaneis, apice nigro-marginatis.* — Long. 4 lin.

Grav. Micr. 191. 2. — Mannerh. Brachelytr. 60. 1.

*Tachyporus fimbriatus* Grav. Mon. 17. 34.

Habitat in America septentrionali.

Habitu omnino *T. humeralis*, at duplo maior, niger, nitidus. Antennae capitis thoracisque prope longitudine, articulo tertio secundo plus sesqui longiore, 5 — 10 aequalibus, subobconicis, crassitie haud brevioribus, ultimo subovato, apice compresso, nigricantes, articulis 4 primis rufis, ultimo apice ferrugineo. Caput subtiliter punctulatum, interstitiis confertissime subtiliter alutaceis, ore rufo, mandibulis piceis. Thorax coleoptera latitudine vix superans, latitudine baseos tertia parte brevior, antrorsum leviter angustatus, basi leviter emarginatus, lateribus modice rotundatus, basi utrinque leviter sinuatus, angulis anterioribus parum antrorsum prominentibus posterioribusque obtusis, modice convexus, confertissime subtilissimeque alutaceus, vage subtilissime obsoleteque punctulatus. Elytra thorace sesqui longiora, subtilissime confertissimeque alutacea, distincte subseriatim punctata, dilute castanea, nitida, margine apicali nigricante. Abdomen crebre distinctius punctatum. Pedes rufo-picei, femoribus obscurioribus.

Mas abdominis segmento superiore sexto quadridentato, dentibus omnibus acutis, intermediis magis prominentibus segmento inferiore quinto apice profundius emarginato, sexto bifido, laciniis longioribus, validis, compressis, parum curvatis, acuminatis.

Femina abdominis segmento superiore sexto trifido, laciniis laterali-bus spiniformibus, intermedia his brevior, apice acute emarginata, bicuspi.

Variat minus maturus thoracis margine laterali rufescente, elytris apice concoloribus, vel totus dilute castaneus, fronte nigricante.

\*26. *T. subterraneus*: *Niger, nitidus, antennarum basi elytrorumque macula humerali oblonga rufo-testaceis, pedibus piceis.* — *Long.* 3 lin.

*Grav. Micr.* 135. 1. — *Latr. Hist. nat. Crust. et Ins.* IX. 393. 1. — *Gyll. Ins. Suec.* II. 252. 2. — *Sahlb. Ins. Fenn.* I. 295. 1. — *Mannerh. Brachelytr.* 60. 2. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 510. 6.

*Staph. subterraneus* *Linn. Faun. Suec. n.* 849. *Syst. Nat.* I II. 684. 12. — *Müll. Prodr.* 98. 1111. — *Goeze Ent. Beitr.* I. 719. 12. — *De Vill. Ent.* I. 414. 12. — *Payk. Mon. Staph.* 73. 52. *Faun. Suec.* III. 420. 66. — *Marsh. Ent. Britt.* 518. 59.

*Oxyporus subterraneus* *Fab. Ent. Syst.* I. II. 532. 4. *Syst. El.* II. 605. 4.

*Tachyporus subterraneus* *Grav. Mon.* 21. 40.

*Var. Thoracis lateribus elytrisque rufo-testaceis, his sutura margineque apicali nigris.*

*Oxyporus marginatus* *Fab. Ent. Syst.* I. II. 532. 6. *Syst. El.* II. 605. 6.

*Tachinus marginatus* *Gyll. Ins. Suec.* II. 255. 4.

*Tachyporus bicolor* *Grav. Mon.* 17. 33.

*Tachinus biplagiatus* *Boisd. et Lacord. Faun. Ent. Paris.* I 511. 7.

*Staphyl. latus* *Marsh. Ent. Britt.* 524. 76.

Habitat in Europa.

Corpus parum convexum, nigrum, nitidum. Antennae capitis thoracisque longitudine, articulo secundo tertioque subaequalibus, 5 — 10 subobconicis, inter se aequalibus, ultimo oblongo-ovato, cinereo-pubescentes, nigricantes, articulis 4 primis rufo-testaceis, glabris. Palpi rufo-picei. Caput crebre subtiliter punctulatum, ore concolore. Thorax basi coleopterorum latitudine, latitudine baseos dimidio brevior, antrorsum leviter angustatus,



apice profundius emarginatus, angulis anterioribus subrotundatis, lateribus rotundatus, basi recte truncatus, angulis posterioribus obtusiusculis, leviter convexus, subtilissime alutaceus, crebre subtiliter punctulatus, margine laterali saepe rufescente. Elytra thorace duplo longiora, subtilissime alutacea, subtiliter punctulata, macula magna oblonga, prope marginem anteriorem ab humero ultra medium descendente, satis determinata, rufo-testacea. Abdomen crebre profundius punctulatum, segmentis singulis margine apicali rufo-piceis. Pedes iam rufo-, iam fusco-picei.

Mas abdominis segmento superiore sexto apice quadridentato, dentibus omnibus obtusis, intermediis magis prominentibus; segmento inferiore quinto apice emarginato, utrinque spina valida, apice bifida, lacinia anteriore muricata, armato, intra sinum late impresso, sexto bifido, laciniiis validis, compressis, deorsum curvatis, acuminatis.

Femina abdominis segmento superiore sexto apice quadrifido, laciniiis lateralibus validis, linearibus, intermediis his multo brevioribus et tenuioribus, spiniformibus.

Variat elytrorum macula dilatata: nonnunquam elytra tota rufo-testacea, sutura margineque laterali exceptis.

\* 27. *T. bipustulatus*: Nigro-piceus, antennis, pedibus, thoracis limbo maculaque transversali baseos elytrorum rufis. — Long. 3 lin.

Grav. Micr. 135. 2. — Latr. Hist. nat. Crust. et Ins. IX. 394. 2. — Gyll. Ins. Suec. II. 254. 3. — Mannerh. Brachelytr. 61. 3.

*Oxyporus bipustulatus* Fab. Ent. Syst. I. II. 533. 9 Syst. El. II. 606. 9. — Panz. Faun. Germ. 16. 21. Ent. Germ. 369. 5. — Cederh. Faun. Ingr. 120. 370.

*Tachyporus bipustulatus* Grav. Mon. 21. 39.

*Staph. subterraneus* var. Payk. Faun. Suec. III. 420. 66. not.

Habitat in Europa.

Corpus modice convexus, nigro-piceum, nitidum. Antennae capite thoraceque breviores, extrorsum crassiores, articulo tertio secundo paulo longiore, articulis 5 — 10 sensim paululum brevioribus, ultimo ovato, compresso, totae rufae. Palpi rufi. Caput subtilissime punctulatum, fronte utrinque leviter impressa, nigrum, ore concolore. Thorax basi coleoptera latitudine vix superans, latitudine baseos paulo brevior, apice emarginatus, lateribus rotundatus, basi utrinque subsinuatus, angulis anterioribus antrorsum leviter prominentibus posterioribusque rotundatis, modice convexus, omnium subtilissime confertissimeque alutaceus, crebre subtilissime punctulatus, nigro-piceus, limbo omni indeterminate rufescente. Elytra thorace plus sesqui longiora, confertissime subtilissimeque alutacea, crebre subtiliter punctata, nigro-picea, macula magna basali transversa indeterminata rufa. Abdomen distinctius crebreque punctatum, pube subtili brevissima cinerea adpersum, segmentis omnibus margine postico rufo-piceis. Pedes rufi.

Mas abdominis segmento superiore sexto quadridentato, dentibus acutiusculis, intermediis multo magis prominentibus; segmento inferiore quinto apice late profundius emarginato, sexto bifido, laciniiis breviusculis, validis, compressis, leviter curvatis, acutiusculis.

Femina abdominis segmento superiore sexto quadrifido, laciniiis late-

ralibus validioribus, linearibus, intermediis his parum brevioribus, tenuibus, spiniformibus, acutis, inter se ad segmenti medium usque divisus.

Obs. *Oxyporum bipustulatum* Fab. potius ad *Conurum bipustulatum* quam ad hanc speciem pertinere autumat Gyllenhal. (Ins. Suec. II. p. 225.) Equidem autem insectum modo descriptum sub hoc nomine in collectione Fabriciana vidi, quare hoc synonymum indubie attuli.

\* 28. *T. pallipes*: *Niger, nitidus, antennarum basi, pedibus thoracisque limbo laterali rufo-ferrugineis, elytris thorace paulo longioribus, nigropiceis, puncto humerali margineque apicali rufescentibus.* — Long. 3 lin.

*Boisd. et Lacord. Faun. Ent. Paris. I. 508. 4.*

*Tachyporus pallipes* Grav. Mon. 20. 38.

*Tachinus humeralis* var. 1 Grav. Micr. 130. 3.

*Tachinus humeralis* var. a Gyll. Ins. Suec. II. 256. 5.

Habitat Lutetiae et in Hercyniae, Austriae Uralensibusque montibus.

Statura omnino *T. rufipedis*, niger, nitidus. Antennae capite thoraceque paulo longiores, apice paulo crassiores, articulo tertio secundo paulo longiore, 5 — 10 subobconicis, sensim paulo brevioribus et crassioribus, ultimo subovato, articulo primo rufo-testaceo, reliquis piceis. Palpi picei, basi testacei. Caput omnium subtilissime punctulatum, ore concolore. Thorax coleopteris vix latior, latitudine baseos tertia parte brevior, antrorsum leviter angustatus, apice late emarginatus, lateribus rotundatus, basi truncatus, angulis anterioribus leviter prominentibus posterioribusque rotundatis, convexus, omnium subtilissime punctatus, margine anteriore posterioreque tenui obscurius, laterali latiore distinctoque testaceis. Scutellum parvum, laeve, nigro-piceum. Elytra thorace sesqui longiora, subtiliter punctulata, nigro-picea, nitida, puncto humerali margineque apicali dilutioribus, rufo-piceis. Abdomen distinctius licet subtiliter punctatum, pube brevissima subtilissimaque adpersum. Pedes rufo-testacei.

Mas abdominis segmento superiore sexto acute quadridentato, dentibus intermediis multo magis prominentibus; segmentis inferioribus 3 — 5 medio leviter impressis, quinto apice late at minus profunde emarginato, sexto bifido, laciniis minus elongatis, parum curvatis, apice compressis, acuminatis.

Femina abdominis segmento superiore sexto quadridentato, laciniis acuminatis, spiniformibus, intermediis acutissimis, ad segmenti medium usque divisus.

Variat elytrorum macula basali transversali indeterminata testacea, pedibus pallidioribus, et minus maturus elytris ferrugineis, dorso posteriori infuscato.

Obs. 1. Ab affinis facile distinguitur femina segmento abdominis subquadridentato.

2. A Gyllenhalio cum *T. humeralis* mare confusus, quod ex eius descriptione patet: exemplum etiam femineum *T. pallipedis* pro *T. humeralis* mare a Schönherro missum in collectione Dom. Prof. Germar exstat.

\* 29. *T. axillaris*: *Piceo-niger, nitidus, antennarum basi, pedibus thoracisque limbo rufo-testaceis, elytris thorace sesqui longioribus, macula humerali summoque margine apicali rufis.* — Long.  $2\frac{2}{3}$  lin.

Habitat in America septentrionali, Dom. Zimmermann.

Statura omnino *T. flavipedis*, at paulo minor, piceo-niger, nitidus. Antennae capite thoraceque fere longiores, apice paulo crassiores, articulo tertio secundo sesqui longiore, 5 — 10 crassioribus, subobconicis, inter se subaequalibus, ultimo oblongo-ovato, compresso, nigrae, articulis duobus primis rufis. Palpi nigro-picei. Caput omnium subtilissime punctulatum, ore concolore. Thorax coleopterorum latitudine, latitudine baseos dimidio brevior, antrorsum leviter angustatus, apice leviter emarginatus, lateribus rotundatus, basi truncatus, angulis anterioribus parum prominentibus posterioribusque rotundatis, leviter convexus, omnium subtilissime alutaceus, crebre subtilissime punctatus, margine basali apicalique tenui lateraliq[ue] latiore castaneo-testaceo. Scutellum piceum. Elytra thorace sesqui longiora, omnium subtilissime alutacea, crebre subtiliter punctata, nigro-vel brunneo-picea, macula humerali indeterminata, saepius per totam basin extensa, margineque summo apicali rufis. Abdomen crebre subtilius punctatum, segmentis singulis margine postico rufo. Pedes rufi.

Mas abdominis segmento superiore sexto quadridentato, dentibus lateralibus acutiusculis, intermediis valde prominentibus, acutis, prope spiniformibus; segmento inferiore quarto leviter triangulariter impresso, quinto apice late profundius emarginato, secundum totam longitudinem late et parum profunde impresso, sexto bifido, laciniis longioribus validioribusque, leviter curvatis, acuminatis.

Femina abdominis segmento superiore sexto quadrifido, laciniis omnibus aequalibus, spiniformibus.

\* 30. *T. basalis*: *Niger, nitidus, antennarum basi pedibusque rufis, elytris thorace paulo longioribus, basi rufis.* — Long.  $2\frac{1}{2}$  lin.

Habitat in montibus Uralensibus, Dom. Prof. Eversmann.

Statura fere *T. marginelli* et maioribus eius individuis aequalis, niger, nitidus. Antennae capitis thoracisque longitudine, extrorsum crassiores, articulis 2 et 3 aequalibus, 5—10 leviter obconicis, sensim paulo brevioribus et crassioribus, ultimo crassiusculo, ovato, piceae, articulis 4 primis rufo-testaceis. Palpi rufo-picei. Caput laeve, parce omnium subtilissime punctulatum, ore concolore. Thorax coleopteris paulo latior, latitudine dimidio brevior, apice leviter emarginatus, lateribus rotundatus, basi truncatus, angulis anterioribus leviter prominentibus, obtusiusculis, posterioribus rotundatis, leviter convexus, parcius obsolete subtiliterque punctatus. Scutellum piceum. Elytra thorace longiora, fortius crebre punctata, nitida, fusco-picea, basi indeterminate castaneo-rufa. Abdomen parce subtiliter punctatum, nigrum, nitidum, segmentis singulis margine apicali rufo-piceo. Pedes dilute rufi.

Mas abdominis segmento superiore sexto quadridentato, dentibus omnibus acutiusculis, intermediis multo magis productis, approximatis; segmento inferiore quinto late leviter emarginato, sexto profunde inciso, laciniis acuminatis, externe unidentatis.

Femina latet.



- \* 31. *T. marginellus*: *Nigro-piceus, nitidus, antennarum basi, pedibus, thoracisque limbo castaneo-testaceis, elytris thorace sesqui longioribus, nigro-piceis, vitta intramarginali margineque apicali dilutionibus.* — Long.  $1\frac{1}{2}$  — 2 lin.

*Grav. Micr.* 143. 14. — *Latr. Hist. nat. Crust. et Ins.* IX. 397. 14. — *Gyll. Ins. Suec.* II. 265. 12. — *Zetterst. Faun. Lappon.* I. 63. 6. *Ins. Lappon.* 57. 6. — *Sahlb. Ins. Fenn.* I. 301. 11. — *Mannevh. Brachelytr.* 62. 11. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 512. 9. — *Runde Brachelytr. Hal.* 26. 5. — *Er. Col. March.* I. 399. 5.

*Staphyl. marginellus* *Fab. Spec. Ins.* I. 337. 20. *Mant. Ins.* I. 221. 27.

*Oxyporus marginellus* *Fab. Ent. Syst.* I. II. 536. 23. *Syst. El.* II. 607. 23. — *Panz. Faun. Germ.* 9. 13. *Ent. Germ.* 361. 18.

*Tachyporus marginellus* *Grav. Mon.* 13. 27.

*Tachinus laticollis* *Grav. Micr.* 141. 10. — *Latr. Hist. nat. Crust. et Ins.* IX. 396. 10. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 509. 5.

*Tachyporus laticollis* *Grav. Mon.* 15. 29.

*Staphylin. laevigatus* *Marsh. Ent. Britt.* 519. 62.

*Tachinus intermedius* *Mannerh. Brachelytr.* 62. 9.

*Staphyl. rufipes* var.  $\beta$  *Payk. Mon. Staph.* 76. 53.

*Staphyl. rufipes* var. 2 *Payk. Faun. Suec.* III. 419. 65.

*Staphyl. rufipes* var. 1 *Walk. Faun. Paris.* I. 273. 20.

*Staphyl. marginatus* *Fourcr. Ent. Par.* I. 169. 18.

*Le Staph. à étuis bordés de jaune* *Geoffr. Hist. Ins.* I. 168. 19:

*Var. Testaceus, capite thoracisque disco fuscis (immaturus).*

*Tachinus immaturus* *Grav. Micr.* 142. 12. — *Latr. Hist. nat. Crust. et Ins.* IX. 396. 12.

*Tachyporus immaturus* *Grav.* 13. 25.

Habitat in Europa.

Habitu omnino rufipedis, at triplo fere minor. Antennae capite thoraceque paulo longiores, apice parum crassiores, articulo tertio secundo perparum longiore, 5 — 10 subobconicis, sensim paulo brevioribus, ultimo oblongo, subovato, nigro-piceae, basi rufae, articulo primo rufo-testaceo. Palpi picei. Caput subtilissime punctulatum, nigrum, nitidum, ore concolore. Thorax coleopteris paulo latior, latitudine baseos tertia parte brevior, antrosum modice angustatus, apice profundius emarginatus, lateribus fortius rotundatus, basi truncatus, angulis anterioribus modice prominentibus posterioribusque obtusis, convexus, dense subtiliter punctatus, niger, nitidus, limbo laterali latiore posticoque tenui testaceis. Scutellum nigrum, subtilissime punctulatum. Elytra thorace sesqui longiora, dense minus subtiliter punctata, obscure brunneo-picea, vitta intramarginali, interdum obsoleta, margineque apicali testaceis. Abdomen minus dense subtiliter punctatum, pubes subtilissima grisea adpersum, piceo-nigrum, nitidum, segmentis singulis margine postico rufo-piceo. Pedes rufo-testacei.

Mas abdominis segmento superiore sexto quadridentato, dentibus omnibus valde acutis, intermediis paulo magis prominentibus; segmento inferiore quinto apice profunde sinuato, margine intra sinum late spongioso, sexto profundo inciso, laciniis triangularibus, apice fortiter acuminatis seu mucronatis.

Femina abdominis segmento superiore sexto quadridentato, laciniis omnibus spiniformibus, acuminatis, intermediis paulo brevioribus tenuioribusque.

Obs. *Oxyp. marginellus* Panz., a Gravenhorstio ad *T. humeralem* relatus, ob magnitudinem minorem potius ad hanc speciem pertinere videtur, et quidem ad individua obscuriora, elytrorum vitta intramarginali obsoleta. *T. laticollis* nonnisi statura paulo maiore, thorace lateribus magis rotundato eiusque limbo laterali minus determinato a *T. marginello* genuino differt.

- \* 32. *T. fimetarius: Niger, nitidus, pedibus, thoracis limbo elytrisque castaneo-testaceis, his dorso plerumque infuscatis, antennis nigris.* — Long. 2 lin.

*Grav. Micr.* 141. 9. — *Latr. Hist. nat. Crust. et Ins.* IX. 396. 9. — *Gyll. Ins. Suec.* II. 263. 11. — *Zetterst. Faun. Lappon.* I. 63. 5. *Ins. Lappon.* 57. 5. — *Sahlb. Ins. Fenn.* I. 300. 10. — *Mannerh. Brachelytr.* 62. 10. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 511. 8. — *Runde Brachelytr. Hal.* 26. 4. — *Er. Col. March.* I. 398. 4.

*Tachyporus fimetarius* Grav. Mon. 14. 28.

*Tachinus sordidus* Grav. Micr. 141. 8. — *Latr. Hist. nat. Crust. et Ins.* IX. 395. 8.

*Staphyl. fuscipes* Panz. Ent. Germ. 356. 28. Faun. Germ. 27. 12.

*Spec. intermed. inter T. laticollem et fimetarium:* Grav. Mon. p. 15.

Habitat in Europa.

Habitu omnino *T. flavipedis*, at duplo minor. Antennae capitis thoracisque longitudine, apice vix crassiores, articulo tertio secundo paulo longiore, 5—10 elongato-obconicis, subaequalibus, ultimo elongato, fusiformi, acuminato, totae nigrae. Palpi nigro-picei. Caput laeve, nigrum, nitidum, ore concolore. Thorax coleopterorum latitudine, latitudine baseos quarta fere parte brevior, antrorsum modice angustatus, apice leviter emarginatus, lateribus rotundatus, basi truncatus, angulis anterioribus parum prominentibus posterioribusque rotundatis, leviter transversim convexus, laevis, niger, nitidus, limbo antico posticoque tenuibus lateralique latiore castaneo-testaceis. Scutellum piceum, laeve. Elytra thorace plus sesqui longiora, parce subtiliter punctulata, castaneo-testacea, nitidula, dorso magis minusve infuscata. Abdomen nigrum, nitidum, parcius subtiliter punctulatum, pube parca brevissima subtilissimaque adpersum. Pectus nigrum, nitidum. Pedes testacei.

Mas abdominis segmento superiore sexto quadridentato, dentibus longitudine subaequalibus, acutis, lateralibus fere spiniformibus; segmento inferiore quinto apice medio profunde sinuato, sinu utrinque denticulo acuto terminato, sexto profunde inciso, laciniis triangularibus, fortiter acuminatis.

Femina abdominis segmento superiore sexto apice quadrifido, laciniis subaequalibus, fortiter acuminatis.

Obs. *T. flavipedi* habitu et colore similis, differt autem statura duplo minore, capite thoraceque impunctatis, abdominisque in utroque sexu structura. — *T. castaneus* Grav. secundum specimina typica ad *T. flavipidem* pertinet.

- \* 33. *T. collaris: Dense punctatus, nigro-piceus, antennis, pedibus thoraceque rufis, hoc disco plerumque infuscato.* — Long.  $1\frac{1}{2}$  —  $1\frac{2}{3}$  lin.

*Grav. Micr.* 133. 13. — *Latr. Hist. nat. Crust. et Ins.* IX. 397. 13. — *Gyll. Ins. Suec.* II. 266. 13. — *Sahlb. Ins. Fenn.* I. 302. 12. — *Mannerh. Brachelytr.* 62. 12. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 512. 10. — *Er. Col. March.* I. 400. 6.

*Tachyporus collaris* Grav. Mon. 13. 24.

*Var. Pallidus, immaturus.*

*Tachinus corticinus* Grav. Micr. 131. 11. — *Latr. Hist. nat. Crust. et Ins.* IX. 396. 11.

*Tachyporus corticinus* Grav. Mon. 13. 16.

*Tachinus flavellus* Zetterst. Faun. Lappon. I. 64. 7. Ins. Lappon. I. 57. 7. (forte)

Habitat in Europa.

Statura fere *T. marginelli*. Antennae capite thoraceque paulo longiores, apice paulo crassiores, articulo tertio secundo paulo longiore, 5 — 10 obconicis, longitudine subaequalibus, at sensim paulo crassioribus, ultimo oblongo, apice acuminato et subcompresso, rufae, apicem versus obscuriores. Palpi picei. Caput crebre subtiliter punctatum, nigrum, nitidum, ore concolore. Thorax coleopteris paulo latior, latitudine baseos dimidio brevior, antrorsum modice angustatus, apice profundius emarginatus, lateribus rotundatus, basi utrinque subsinuatus, angulis anterioribus modice prominentibus, rotundatis, posterioribus subrectis, externe rotundatis, convexus, dense subtilius punctatus, piceo-rufus, disco saepius infuscato. Scutellum piceum. Elytra thorace sesqui longiora, dense minus subtiliter punctata, iam fusco-iam rufo-picea, nitida. Abdomen parcius subtiliter punctatum, pube subtilissima brevissimaque adpersum, nigrum, nitidum, segmentis singulis margine postico piceo. Pectus nigrum. Pedes rufo-testacei.

Mas abdominis segmento superiore sexto quadridentato, dentibus breviusculis, intermediis multo magis prominentibus; segmento inferiore quinto apice fortiter triangulariter exciso, sexto inciso, laciniis triangularibus, apice acuminatis, fere mucronatis.

Femina abdominis segmento superiore sexto apice quadrifido, laciniis subaequalibus, acuminatis.

\* 34. *T. elongatus*: *Elongatus, nigro-piceus, nitidus, antennis pedibusque rufis, elytris subtiliter substriato-punctatis, margine apicali rufescente.* — Long.  $3\frac{1}{2}$  lin.

Gyll. Ins. Suec. II. 251. 1. — Zetterst. Faun. Lappon. I. 61. 1. Ins. Lappon. 56. 1.

Habitat in Suecia, Dom. Schüppel, in Helvetia, Dom. Guyot, in Unalashka, Dom. Chamisso.

Statura elongata, linearis, fere subcylindrica a congeneribus discedens, piceo-niger, subnitidus. Antennae capitis thoracisque longitudine, crassiusculae, articulo tertio secundo sesqui longiore, 5 — 10 sensim paulo brevioribus at vix crassioribus, ultimo subovato, apice abrupte compresso, totae rufae. Palpi rufi. Caput omnium subtilissime parce punctulatum. Thorax coleopteris vix latior, latitudine baseos quarta parte brevior, antrorsum parum angustatus, apice profundius emarginatus, lateribus leviter rotundatus, basi truncatus, angulis anterioribus modice prominentibus, acutiusculis, acumine rotundato, posterioribus rotundatis, transversim convexus, omnium subtilissime parce punctulatus. Elytra thorace vix sesqui longiora, subseriatim subtilius punctata, nigro-picea, margine apicali rufescente. Abdomen parallelum, subtiliter punctatum, versus latera tenuiter griseo-pubescentis, segmentis piceo-marginatis. Pedes rufi.

Mas abdominis segmento superiore sexto quadridentato, dentibus minutis, intermediis paulo magis prominentibus; segmento inferiore quinto apice late profundeque emarginato, sexto bifido, laciniis validis, elongatis, compressis, fortius deorsum curvatis, apice acuminatis.



Femina abdominis segmento superiore sexto apice quadrifido, lacinii subaequalibus, acuminatis.

Variat thorace elytrisque rufo-piceis.

- \*35. *T. discoideus*: *Nigro-piceus, nitidus, antennis pedibusque rufis, elytris fortius punctatis, rufis, margine omni piceo.* — Long.  $1\frac{2}{3}$  lin.

Habitat in Austria.

Antennae capitis thoracisque fere longitudine, validiusculae, apicem versus leviter incrassatae, articulo 2 et 3 subaequalibus, 5—10 longitudine subaequalibus at sensim paulo crassioribus, ultimo oblongo-ovato, apice acuminato, totae rufo-testaceae. Caput parvum, parcius subtiliter punctatum, nigro-piceum, nitidum, ore rufo-testaceo. Thorax basi coleopterorum latitudine, latitudinē baseos quarta fere parte brevior, antrorsum leviter angustatus, apice profundius emarginatus, lateribus leviter rotundatus, basi truncatus, angulis anterioribus modice prominentibus, leviter deflexis posterioribusque obtusis subrotundatis, parum convexus, parce vageque subtiliter punctatus, iam nigro- iam rufo-piceus, nitidus. Scutellum laeve, piceum. Elytra thorace sesqui prope longiora, crebrius fortiusque punctata, punctis hinc inde subseriatis, nitida, disco laete rufo, limbo omni piceo. Abdomen minus subtiliter punctulatum, nigro-piceum, segmentis singulis margine apicali rufo-piceo. Pectus nigrum, nitidum. Pedes rufo-testacei.

Mas abdominis segmento superiore sexto quadridentato, dentibus minutis, intermediis fortius prominentibus; segmento inferiore quinto apice leviter triangulariter emarginato, sexto profunde inciso, lobis triangularibus, apice acuminatis.

Femina abdominis segmento superiore sexto apice quadrifido, lacinii subaequalibus, acuminatis.

#### Species mihi invisae:

36. *T. rufipennis*: *Niger, nitidus, glaber, antennis basi rufo-piceis, elytris profunde punctatis, rufis, apice nigro-marginatis, tarsis ferrugineis.*

*Gyll. Ins. Suec. II. 259. 8.*

Habitat in Sueciae stercoratis rarius.

Statura omnino *T. humeralis*, sed adhuc paulo maior, et vere distinctus. Caput rotundatum, nigrum, nitidissimum, subtiliter punctulatum, linea tenui angulata inter antennis impressum; os et palpi nigro-picei. Antennae thorace paulo longiores, nigro-fuscae, basi rufo-piceae. Thorax latus, transversus, fere ut in *T. humerali*, antice tamen profundius emarginatus, supra valde convexus, subtilissime punctulatus, totus niger, nitidissimus. Scutellum triangulare, rufo-piceum, subtiliter punctulatum. Elytra thorace multo angustiora, sed fere duplo longiora, apice truncata, angulis ut in prioribus, supra convexa, crebre et sat profunde vage punctata, saturate rufa, nitida, margine postico late nigro, nigredine iuxta marginem lateralem altius adscendente, sutura etiam anguste nigra. Abdomen pone elytra illarum longitudine, convexus, totum nigrum, nitidum, crebre punctatum; anus secundum sexum ut in *T. rufipede* armatus. Corpus totum subtus nigrum, nitidum,

profunde punctatum. Pedes elongati, nigri, nitidi, spinulis tibiæ tarsisque ferrugineis. (Gyll.)

### **Trichophyus Er.**

*Er. Col. March. I. p. 402.*

*Trichophya Mannerh. Brachelytr. p. 73.*

*Antennae 11-articulatae, capillares.*

*Palpi maxillares filiformes.*

*Ligula apice emarginata, paraglossis distinctioribus, ligula duplo brevioribus.*

*Elytra pectori longitudine aequalia.*

*Tarsi 5-articulati.*

Corpus oblongum, subdepressum, alatum. Caput deflexum, thorace minus et angustius, subtriangulare, oculis parvis, rotundatis, prominulis. Labrum transversum, apice rotundatum. Mandibulae muticae, margine interno limbo membranaceo fimbriato. Maxillae (t. IV. f. 8. b.) mala utraque leviter elongata, lineari, membranacea, longitudine altera alteri subaequali, exteriori apice barbata, interiori apice leviter uncinata, dein spinulis tribus minutis armata, margine interiore medio barbato. Palpi maxillares sat elongati, articulo secundo elongato, tertio quartoque singulis hoc paulo brevioribus, inter se subaequalibus, quarto apice acuminato. Labium (t. IV. f. 8. a.) mento transversum, ligula angustata, prominula, apice emarginata, paraglossis ligulae connatis, et eadem brevioribus. Palpi labiales biarticulati, articulo secundo parvo, acuminato. Antennae longitudine dimidii corporis, articulo primo secundoque crassiusculis, primo clavato, secundo subgloboso, sequentibus valde tenuibus, sensim angustioribus, pilis longis verticillatis undique adpersis. \*) Thorax coleopterorum prope latitudine, antrorsum haud angustatus, parum convexus. Scutellum triangulare. Elytra pectus haud excedentia. Abdomen apicem versus subangustatum. Pedes breves, intermedii basi adproximati, coxae posticae triangulares, tibiae pubescentes, tarsi omnes 5-articulati, antici articulis 4 primis leviter dilatatis, posteriores articulo primo leviter elongato.

### **Observationes.**

1. Genus admodum distinctum, statura autem minus convexa, thorace minus amplo, capite magis exserto, huic tribui quodammodo alienum videtur. Aleocharis adscribi nequit, antennae enim infra oculos sub capitis margine laterali insertae, palporum maxillarum articulus ultimus haud subulatus, et tarsi antici dilatati. Quibus notis cum Tachyporinis congruit: oris autem structura in Trichophyo singularis, Aleocharorum genera postrema in animum revocans. Palpi labiales perbreves biarticulati mihi videntur.

2. Antennarum structura singulari cum Habrocero sat quidem convenit, reliquis autem characteribus omnibus discedit.

3. Sexus differentia latet.

4. Victus sub arborum emortuarum cortice, sec. Gyll. Specimen nostrum in foro lignario volitans captum est.

5. Nomen generis, huic tribui adscripti, aliquantum mutavi, quoniam omnia Tachyporinorum nomina masculina sunt.

\*) Antennae in nostro specimine mutilatae, quare earum descriptionem secundum Gyllenhalium supplavi.

1. *T. pilicornis*: *Niger, punctatus, pubescens, elytris fuscis, pedibus fusco-testaceis.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I.* 403. 1.

*Trichophya pilicornis* *Mannerh. Brachelytr.* 73. 1.

*Aleochara pilicornis* *Gyll. Ins. Suec. II.* 417. 39.

Habitat in Germania, Suecia.

Caput crebrius punctatum, nigrum, nitidum, ore fusco, oris partibus antennarumque articulis primis rufo-testaceis. Thorax latitudine plus sesqui brevior, lateribus rotundatus, basin versus non minus quam antrorsum angustior, basi leviter rotundatus, angulis posterioribus obtusis, apice leviter emarginatus, supra parum convexus, lateribus basique planiusculis, confertim punctatus, tenuiter griseo-pubescens, fusco-niger, nitidus. Elytra thorace sesqui longiora, apice oblique introrsum truncata, dense punctata, tenuius pubescentia, fusca, opaca. Abdomen pectusque dense punctata, tenuiter pubescentia, nigra, ano fusco. Pedes testacei.

### **Boletobius Leach.**

*Mannerh. Brachelytr. p.* 64.

*Megacronus Steph. Illustr. V. p.* 165 et *Boletobius* *ibid. p.* 171.

*Bryocharis* *Boisd. et Lacord. Faun. Ent. Paris. I. p.* 502 et *Boletobius* *ibid. p.* 497.

*Antennae 11-articulatae, filiformes.*

*Palpi maxillares filiformes.*

*Ligula apice emarginata.*

*Elytra pectori subaequalia.*

*Tarsi 5-articulati.*

Corpus elongatum, utrinque attenuatum et deorsum curvatum, alatum. Caput thoraci immissum, deflexum, aliis breviusculum, aliis oblongum vel elongatum, oculis mediocribus, ovalibus, vix prominulis. Labrum aliis transversum, aliis (capite elongato) quadratum, apice truncatum. Mandibulae minores, apice falcatae, acutae, interne basi membrana fimbriatae. Maxillae (t. IV. f. 6. b.) malis coriaceis, interiore interne, exteriori apice ciliatis. Palpi maxillares leviter elongati, articulo primo minuto obconico, secundo tertioque vel subaequalibus, vel secundo paulo longiore, apicem versus leviter incrassatis, quarto praecedenti aequali vel eodem paulo longiore, plurimis cylindrico, apice obtuse acuminato. Labium (t. IV. f. 6. a.) mento transversum, parte membranacea ante palporum basin elongata, ligula rotundata, medio incisa, membranacea, paraglossis ligulae multo brevioribus, eique connatis et nonnisi linea ciliata distinguendis. Palpi labiales articulis duobus primis subaequalibus, tertio paulo longiore, apice obtuse acuminato. Antennae utrinque infra oculos insertae, filiformes, apicem versus plerumque leniter incrassatae. Thorax basi fere coleopterorum latitudine, apicem versus angustatus, latitudine parum brevior, lateribus rotundatus, basi truncatus, apice pro receptione capitis emarginatus, lateribus, praecipue anterieus, deflexis. Scutellum triangulare. Elytra apice truncata. Abdomen apicem versus sensim attenuatum, deorsum curvatum. Mesosternum haud carinatum. Pedes graciliores, coxis intermediis valde approximatis, posticis amplis, parallelipipedis, scutatis, femoribus saepius apice obsolete setosis,



tibiis omnibus spinulosis, spinis terminalibus validioribus, tarsis 5-articulatis, elongatis, gracilioribus, posterioribus articulo primo elongato, sequentibus sensim decrescentibus, ultimo iterum longiore.

#### Observationes.

1. Corpore graciliore, incurvo, elytris coxas posticas vix superantibus, his parallelopidedis, amplis, ut in *Mordellis* scutatis, pedibus etiam ut in hac familia constructis a praecedentibus discedens, summa affinitate cum genere sequente iungitur.

2. Caput laevissimum; thorax laevissimus, ad margines punctis paucis, interdum etiam dorso punctorum pari impressus; scutellum laeve; elytra plerisque punctorum seriebus tribus ordinariis notata, intima striae leviter exaratae suturali, extima striae tenui marginali impositis, intermedia discoidali; abdomine punctato.

3. *Megacronus* Steph. nonnisi capite minus elongato et antennis minus gracilibus, *Bryocharis* Boisd. et Lacord. nonnisi tarsis anticis in mare dilatatis distinguuntur. Distinctio illa omnino arbitraria, haec temeraria; tarsi enim dilatati simplicesve vix unquam in tota Staphylinorum familia characterem genericum praebent, speciebus inter se maxime affinibus saepius hac nota discrepantibus.

4. Sexus differentia varia, apud singulas species notanda.

5. Victus in silvis, in boletis et sub muscis et foliis deciduis.

\* 1. *B. analis*: *Niger, nitidus, pedibus cum coxis anticis, antennis articulis 4 primis ultimoque testaceis, ano elytrisque rufis.* — Long.  $2\frac{1}{2}$  — 3 lin.

*Mannerh. Brachelytr.* 64. 3. — *Runde Brachelytr. Hal.* 26. 1. — *Er. Col. March.* I. 403. 1.

*Staphyl. analis* Payk. *Mon. Staph.* 47. 34. *Faun. Suec. III.* 394. 34. — *Walk. Faun. Paris.* I. 272. 17. — *Oliv. Ent. III.* 42. 28. 38. t. 3. f. 24. — *Marsh. Ent. Britt.* 522. 70.

*Tachinus analis* Grav. *Mon.* 148. 20. — *Latr. Hist. nat. Crust. et Ins. IX.* 399. 20. — *Sahlb. Ins. Fenn. I.* 303. 14. — *Zetterst. Ins. Lappon.* 58. 8.

*Tachinus analis* var. 2 Grav. *Mon.* 31. 16.

*Tachinus analis* var. b Gyll. *Ins. Suec. IV.* 472. 13.

*Staph. analis* Schrank *Faun. Boica. I.* 649. 890. (*forte*)

*Var. Thorace elytris anoque testaceo-rufis.*

*Staph. analis* var.  $\beta$  Payk. *Mon. Staph.* 47. 34. *Faun. Suec. III.* 395. 34.

*Tachinus merdarius* Gyll. *Ins. Suec. II.* 270. 16. — *Sahlb. Ins. Fenn. I.* 304. 15.

Habitat in Europa.

*Niger, nitidus.* Antennae capite thoraceque paulo longiores, apicem versus sensim leniterque subincrassatae, articulo tertio secundo sesqui longiore, 5 — 10 sensim paulo brevioribus et crassioribus, fuscae, articulis 4 primis ultimoque testaceis, hoc maris oblongo, cylindrico, apice obtuse rotundato, praecedente plus duplo maiore, feminae ovato, praecedente sesqui maiore. Caput parvum, laevissimum, ore late testaceo. Thorax coleopterorum prope latitudine, latitudine baseos parum brevior, lateribus rotundatus, antice paulo magis quam postice angustatus, basi leviter rotundatus, angulis posterioribus obtusissimis, fere rotundatis, satis convexus, laevissimus. Scutellum laevissimum. Elytra thorace tertia parte longiora, rufa; nitida, seriebus tribus ordinariis punctorum remotorum. Abdomen parce punctatum, parceque longius pubescens, segmento ultimo penultimoque rufis, hoc basi nigro. Pedes testacei, antici coxis concoloribus.

Mas tarsi anticis leviter dilatatis, abdominis segmento inferiore sexto apice medio leviter triangulariter producto, antennarum articulo ultimo praecedentibus tribus coniunctis subaequali.

Femina tarsis anticis simplicibus, abdominis segmento inferiore sexto apice rotundato, antennarum articulo ultimo praecedentibus ambobus vix aequali.

Obs. Tach. merdarius Gyll. nonnisi thorace rufo differt, quare vix pro specie peculiari rite habendus: differentiae a Gyllenhalio notatae ad sequentem speciem spectant. Specimina Suecica dedit Dom. Schüppel.

\* 2. *B. cingulatus*: Niger, nitidus, antennis articulis 3 primis ultimisque duobus pedibusque testaceis, his coxis anticis nigris, ano elytrisque rufis. — Long. 3 — 3½ lin.

Mannerh. Brachelytr. 64. 2. — Er. Col. March. I. 404. 2.

*Tachinus analis* var. 1 et 3 Grav. Mon. 31. 16.

*Tachinus analis* Gyll. Ins. Suec. II. 269. 15.

*Bryocharis analis* Boisd. et Lacord. Faun. Ent. Paris. I. 502. 1.

*Oxyporus bicolor* Rossi Faun. Fr. I. 253. 625. (Ed. Helw. I. 314. 626.) forte.

Habitat in Europa.

Statura et summa affinitas praecedentis, at plerumque paulo maior et vere distinctus. Corpus nigrum, nitidum. Antennae capite thoraceque paulo longiores, apicem versus vix incrassatae, articulo tertio secundo duplo prope longiore, 5—10 sensim paulo brevioribus at perparum crassioribus, articulis duobus primis testaceis, tertio rufo-testaceo, vel ferrugineo, 4—9 nigris, feminae penultimo ferrugineo, ultimo hoc paulo maiore, ovato, subacuminato, testaceo, maris ultimis duobus sulphureis, ultimo elongato, praecedentibus duobus longiore. Caput minutum, laevissimum, nigrum, ore late testaceo. Thorax basi coleopterorum latitudine, ibique longitudine paulo latior, apice basi duplo fere angustior, lateribus et basi leviter rotundatus, angulis posterioribus obtusis, leniter convexus, laevissimus. Elytra thorace tertia parte longiora, rufa, nitida, punctorum seriebus tribus ut in praecedente omnino. Abdomen parce punctatum, parceque longius pubescens, ano piloso, segmento sexto toto quintoque apice lato rufis, septimo, quando conspicitur, nigro-piceo. Pedes rufo-testacei, coxis omnibus nigris.

Mas tarsi anticis fortius dilatatis, abdominis segmento inferiore sexto canaliculato, apice leviter producto, antennis articulo ultimo praecedentibus ambobus paulo longiore.

Femina tarsis anticis simplicibus, abdominis segmento inferiore sexto apice rotundato, antennis articulo ultimo praecedentibus ambobus brevioribus.

Variat 1, antennis articulis 3 et 10 piceis, ultimo solo flavo. Specimen Austriacum dedit Dom. Schüppel.

2, antennis articulis ultimis tribus flavis. Specimen in Germania septentrionali lectum communicavit Dom. Schüppel.

Obs. 1. Distinguendus a praecedente praeter antennarum colorem articulo earum ultimo apice acuminato neque obtuso, thorace minus convexo, postice multo latiore, angulis posterioribus minus obtusis, coxis anticis nigris, tarsis anticis maris fortius dilatatis.

2. Oxy. bicolor Rossi ob magnitudinem ultra 3 lin. potius ad hanc quam ad praecedentem speciem pertinere videtur.

- \* 3. *B. inclinans*: *Rufo-testaceus, nitidus, capite, pectore abdomineque nigris, hoc segmentis singulis piceo-marginatis, elytris crebre punctatis.* — Long.  $3\frac{1}{2}$  lin.

*Er. Col. March. I.* 405. 3.

*Tachinus inclinans* Grav. *Mon.* 33. 20.

Habitat in Germania.

Antennae capite thoraceque longiores, apice haud incrassatae, articulo tertio secundo plus sesqui longiore, sequentibus sensim paulo brevioribus, ultimo praecedente paulo longiore, acuminato, testaceae, medio fusciscentes. Caput laevissimum, nigrum, nitidum, ore late testaceo. Thorax posterius coleopteris paululum latior, latitudine summa paulo brevior, apice duplo quam basi angustior, lateribus leviter rotundatis, basi subsinuatus, angulis posterioribus obtusis, leviter convexus, dorso ante medium punctorum pari impressus, rufo-testaceus, nitidus. Scutellum laeve, rufo-testaceum. Elytra thorace sesqui longiora crebre punctata, parcius ferrugineo-pubescentia, rufo-testacea, nitida. Abdomen parce punctatum, parceque pubescens, nigrum, nitidum, segmentis margine apicali piceis, ultimo penultimoque dimidio piceis. Pectus nigrum, nitidum. Pedes testacei, coxis anterioribus concoloribus.

Mas tarsis anticis articulis duobus primis fortiter dilatatis, tibiis intermediis spinis apicalibus fortioribus, tarsis intermediis articulo primo compresso-dilatato, abdominis segmento inferiore quinto apice leviter inciso.

Femina tarsis anterioribus simplicibus, abdominis segmento inferiore quinto integro.

- \* 4. *B. formosus*: *Rufo-testaceus, nitidus, capite abdomineque nigris, hoc ano rufo, antennis fuscis, basi testaceis, elytris vage punctatis.* — Long.  $2\frac{1}{2}$  lin.

*Mannerh. Brachelytr.* 64. 1.

*Tachinus formosus* Grav. *Mon.* 32. 19.

*Megacrönus formosus* Steph. *Illustr. Bt. E. V. t.* 26. f. 5.

*Tachinus merdarius* Grav. *Micr.* 149. 21. — *Latr. Hist. nat. Crust. et Ins.* IX. 399. 21.

Habitat in Germania.

Corpus gracilius, rufo-testaceum, nitidum. Antennae capite thoraceque multo longiores, apice haud crassiores, articulo tertio secundo plus sesqui longiore, sequentibus sensim paulo brevioribus, ultimo praecedente paulo longiore, acuminato, fuscae, articulis duobus primis ultimoque testaceis. Caput laevissimum, nigrum, nitidum, ore testaceo. Thorax posterius coleopteris paulo latior, latitudine baseos paulo brevior, apice duplo prope angustior, lateribus leviter rotundatus, basi subtrisinuatus, angulis posterioribus subrectis, convexus, dorso ante medium punctorum pari impressus. Scutellum laeve. Elytra thorace haud longiora, parcius fortiusque vage punctata, puncto quoque pilo brevi reclinato instructo. Abdomen parce sat fortiter punctatum, parce longius pubescens, nigrum, nitidum, segmento quinto dimidio sextoque toto testaceo-rufis. Pectus cum pedibus testaceum.

Obs. *Tachin. merdarius* Grav., secundum specimen typicum coll. Hoffmanns-



eggianae a Staph. merdario Fab. et Oliv omnino distinctus, cum T. formoso postea descripto omnino convenit.

5. *B. obscurus*: *Nigro-piceus, nitidus, antennis testaceis, pedibus anoque rufo-piceis, elytris vage punctatis*. — Long. 2 lin.

Habitat in ins. Puerto-Rico, Dom. Moritz.

Statura angusta B. formosi, nigro-piceus, nitidus. Antennae capite thoraceque multo breviores, apicem versus sensim modice incrassatae, articulo tertio secundo plus sesqui longiore, sequentibus sensim paulo brevioribus et parum crassioribus, penultimis crassitie brevioribus, ultimo breviter subovato, compresso, totae testaceae. Palpi flavi, maxillares articulo ultimo conico, penultimo haud longiore. Caput breve, laevissimum, ore testaceo. Thorax basi coleopterorum prope latitudine, latitudine baseos vix brevior, antrorsum angustatus, apice plus dimidio quam basi angustior, basi et lateribus leviter rotundatus, angulis posterioribus subrectis, convexus, dorso ante medium punctorum pari impressus. Scutellum laeve. Elytra thorace sesqui longiora, crebre fortius vage punctata, stria suturali impressa, tenuiter pubescentia. Abdomen parcius punctatum, longius pubescens, apice segmentorumque singulorum marginibus rufo-piceis. Pedes rufo-testacei.

6. *B. seriatatus*: *Nigro-piceus, nitidus, antennarum basi pedibusque testaceis, elytris crebre seriatim punctatis*. — Long. 2 lin.

Habitat Novae Valentiae in Columbia, Dom. Moritz.

Statura omnino B. striati, at minor, nigro-piceus, nitidus. Antennae capite thoraceque paulo breviores, apicem versus sensim sat fortiter incrassatae, articulo tertio secundo plus sesqui longiore, sequentibus sensim paulo brevioribus et crassioribus, penultimis transversis, ultimo subovato, apice acuminato, nigrae, articulis duobus primis testaceis. Caput breviusculum, laevissimum, ore flavo-testaceo. Thorax basi coleopterorum prope latitudine, latitudine baseos paulo brevior, antrorsum angustatus, apice dimidio quam basi angustior, basi et lateribus leviter rotundatus, angulis posterioribus subrectis, convexus, marginibus summis brunneis. Elytra thorace sesqui longiora, dorso punctorum sat fortium seriebus pluribus (sex) striaque suturali impressa, parce pubescentia. Abdomen parcius punctatum, longius pubescens, segmentis singulis castaneo-marginatis. Pedes testacei.

- \* 7. *B. cernuus*: *Niger, nitidus, antennarum basi pedibusque testaceis, elytris rufis, punctorum seriebus obsoletioribus, interstitiis obsoletissime punctulatis*. — Long. 2 —  $2\frac{1}{2}$  lin.

Mannerh. *Brachelytr.* 64. 4. — Boisd. et Lacord. *Faun. Ent. Paris.* I. 500. 5. — Er. *Col. March.* I. 406. 4.

*Tachinus cernuus* Grav. *Mon.* 31. 18. — Gyll. *Ins. Suec.* II. 271. 17. — Sahlb. *Ins. Fenn.* I. 304. 16. — Zetterst. *Faun. Lappon.* I. 64. 8. *Ins. Lappon.* 58. 2.

*Var. Thorace elytrisque rufis.*

*Staph. merdarius* Oliv. *Ent.* III. 42. 29. 39. t. 5. f. 45.

*Bolitobius merdarius* Boisd. et Lacord. *Faun. Ent. Paris.* I. 501. 6.

Habitat in Europa.

Statura omnino B. striati. Antennae capitis thoracisque prope longi-

tudine, tenuiores, apice parum incrassatae, articulo tertio secundo plus sesqui longiore, sequentibus sensim paulo brevioribus et crassioribus, ultimo ovato, praecedente sesqui longiore, apice subcompresso, acuminato, pilosellae, fuscae, articulis tribus primis testaceis, ultimo interdum obscure testaceo. Palpi maxillares articulo quarto praecedente paulo brevior, conico. Caput breviusculum, nigrum, nitidum, ore testaceo. Thorax basi coleopterorum latitudine, latitudine baseos paulo brevior, apice plus dimidio quam basi angustior, lateribus praecipue antice deflexis, leviter rotundatis, basi subtruncatus, angulis posterioribus obtusis, laevissimus, margine punctis singulis, dorso ante apicem punctorum pari impressus, nitidus, iam nigerrimus, iam piceo-rufescens, iam testaceo-rufus. Scutellum piceum, nitidum, laevissimum. Elytra thorace tertia parte longiora, rufa vel testaceo-rufa, nitida, punctorum sat remotorum subtilium seriebus tribus, prima ad suturam, secunda in medio dorso, tertia ad marginem lateralem, interstitiis punctis distantibus seriatis subtilibus obsoletisque impressis. Abdomen vage fortius punctulatum, longius griseo-pubescens, nigrum, nitidum, segmentis 4 primis margine postico, quinto-apice, sexto toto rufis. Pectus nigrum, nitidum. Pedes rufo-testacei, cōxis anterioribus iam concoloribus, iam nigro-piceis, tarsis anticis in utroque sexu simplicibus.

- \* 8. *B. rufus*: *Rufo-testaceus, nitidus, abdominis segmentis anterioribus basi fuscescentibus, antennis fuscis, basi testaceis, elytris punctorum seriebus obsoletioribus, interstitiis subtiliter subseriatim punctatis.* — Long.  $2\frac{1}{2}$  lin.

*Er. Col. March. I.* 401. 5.

*Var. Capite pectoreque nigro-piceis.*

Habitat in Germania.

Statura et summa affinitas praecedentis. Antennae capitis thoracisque prope longitudine, apicem versus fortius incrassatae, articulo tertio secundo sesqui longiore, 4—7 sensim brevioribus et crassioribus, 7—10 subaequalibus, longitudine paulo crassioribus, ultimo praecedente sesqui longiore, breviter subovato, apice compresso, obtuse rotundato, fuscae, articulis quatuor primis rufo-testaceis. Palpi maxillares articulo tertio modice incrassato, quarto hoc dimidio prope brevior et duplo tenuiore, valde acuminato. Caput minutum, triangulare, saturate rufo-testaceum, nitidum, ore dilutiore, vertice nonnunquam fuscescente. Thorax basi coleopterorum prope latitudine, apice duplo angustior, latitudine baseos tertia parte prope brevior, basi et lateribus leviter rotundatis, angulis posterioribus subrotundatis, laevissimus vel interdum subtiliter obsoletissimeque vage punctulatus, apice prope marginem anteriorem punctorum pari impressus, saturate rufo-testaceus, nitidus. Scutellum concolor, laevissimum, nitidum. Elytra thorace sesqui longiora, punctorum maiorum seriebus tribus, interstitiis parcius subtiliter obsoleteque subseriatim punctatis, rufo-testacea, nitida. Abdomen supra vage profundius punctatum, subtus crebre fortiter profundeque punctatum, parce fulvo-pubescens, piceo-testaceum seu castaneum, nitidum, segmentis singulis margine apicali rufescente. Pectus saturate rufo-testaceum. Pedes testacei.

*Variat nonnunquam capite pectoreque nigro-piceis.*

Obs. Differt a praecedente praeter colorem antennis apicem versus magis incrassatis, thorace lateribus minus rotundato, punctorum pari apici propiore, elytris in interstitiis distinctius punctulatis. Palporum maxillarium articulo quarto brevior conicoque anceps quasi inter Boletobium et Mycetoporum.

\* 9. *B. bicolor*: *Niger, nitidus, capite, thorace, scutello, pedibus elytrisque rufis, his angulo exteriori apicis nigro.* — Long.  $2\frac{1}{2}$  —  $3\frac{1}{2}$  lin.

*Tachinus bicolor* Grav. Mon. 28. 9.

Habitat in Hercynia, Borussia.

B. anali robustior et saepe duplo maior. Antennae vix capitis thoracisque longitudine, apice paulo crassiores, articulo tertio secundo plus sesqui longiore, 5 — 10 modice compressis, subtriangularibus, latitudine apicis vix brevioribus, inter se aequalibus, ultimo oblongo, praecedente plus duplo longiore, compresso, apice rotundato, fusco-nigrae, articulis 4 primis testaceis, ultimo apice summo ferrugineo. Caput oblongum, testaceo-rufum, ore testaceo. Thorax basi coleopterorum prope latitudine, latitudine summa non brevior, apice paulo angustior, basi et lateribus leviter rotundatus, angulis posterioribus obtusis subrotundatis, leniter convexus, laevissimus, testaceo-rufus, nitidus. Scutellum laevissimum, testaceo-rufum, nitidum. Elytra thorace paulo longiora, punctorum seriebus tribus ordinariis, punctis sat distinctis, serie dorsali quaque striae leviter exaratae imposita, testaceo-rufa, nitida, angulo apicali exteriori a medio margine laterali usque ad suturae apicem nigerrimo. Abdomen crebrius punctatum, parcius nigro-pubescent, nigrum, nitidum. Pectus nigrum, nitidum, mesosterno rufo-testaceo. Pedes testacei, coxis anterioribus concoloribus, femoribus apice fortius spinulosis, tarsis anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore quinto ante apicem carinula parva longitudinali dentiformi instructo.

Femina abdominis segmento inferiore quinto mutico.

\* 10. *B. axillaris*: *Piceus, nitidus, antennarum basi apiceque pedibusque anticis testaceis, capite elytrisque nigris, his vitta longitudinali margineque apicali testaceis.* — Long. 3 lin.

*Tachinus axillaris* Grav. Mon. 29. 11.

Habitat in America septentrionali.

Statura omnino praecedentis. Antennae capitis thoracisque longitudine, apice parum crassiores, articulo tertio secundo sesqui longiore, 5 — 10 crassioribus, modice compressis, subtriangularibus, 6 — 10 latitudine apicis paulo brevioribus, aequalibus, ultimo praecedente duplo longiore et paulo angustiore, oblongo, compresso, nigrae, articulis duobus primis ultimoque testaceis. Caput oblongum, nigrum, nitidum, labro palpisque testaceis. Thorax basi coleopterorum prope latitudine, apice dimidio angustior, latitudine baseos paulo brevior, basi et lateribus leviter rotundatus, angulis posterioribus obtusis, laevissimus, nigro-piceus, nitidus. Scutellum nigrum, nitidum. Elytra thorace sesqui fere longiora, serie dorsali punctis fere sex satis distantibus composita, nigra, nitida, margine apicali vitta longitudinali, ab angulo humerali apicis medium petente, marginem apicalem haud attingente,



testaceis. Abdomen supra parcius, subtus crebrius fortiusque punctatum, piceum, nitidum, segmentis singulis margine apicali dilutiore. Pedes antici cum coxis testacei, posteriores picei, femoribus piceo-testaceis.

Mas abdominis segmento inferiore quinto ante apicem carinula longitudinali dentiformi instructo.

Femina latet.

- \* 11. *B. niger*: *Niger, nitidus, antennarum articulo primo testaceo, abdominis segmento quinto apice rufo, elytris punctorum seriebus obsoletioribus.* — Long.  $4\frac{1}{2}$  — 6 lin.

*Tachinus niger* Grav. Micr. 193. 5. Mon. 30. 12.

*Var. Abdominis segmento quinto apice concolore.*

Habitat in America septentrionali.

Magnus, niger, nitidus. Antennae capite thoraceque paulo breviores, articulis 3 primis teretibus, 2 et 3 subobconicis, hoc praecedente sesqui longiore, sequentibus latioribus, modice compressis, triangularibus, 5 — 10 latitudine apicis paulo brevioribus, aequalibus, ultimo paulo angustiore, oblongo-ovato, nigrae, articulo primo testaceo, secundo piceo, ultimo apice ferrugineo. Caput oblongum, clypeo pallido, labro palpisque piceis. Thorax basi coleopterorum prope latitudine, latitudine baseos haud brevior, apice dimidio angustior, basi et lateribus leviter rotundatus, angulis posterioribus obtusis, laevissimus. Scutellum laevissimum. Elytra thorace parum longiora, punctorum seriebus tribus ordinariis, punctis earum obsoletis, inter se sat remotis. Abdomen segmentis anterioribus parcius, posterioribus crebrius punctatum. Pedes nigro-picei, tarsi antici in utroque sexu simplicibus.

Mas abdomine apice concolore, segmento inferiore sexto tricarinato, carina intermedia apice, lateralibus basi abbreviatis, apice inter carinas laterales emarginato.

Femina abdominis segmento quinto apice late sanguineo, sexto simplice.

Obs. Utriusque sexus nonnisi unicum specimen vidi, nescio igitur an differentia coloris in abdominis apice semper cum differentia sexus conveniat.

- \* 12. *B. striatus*: *Nigro-piceus, nitidus, pedibus segmentorumque abdominis marginibus rufis, elytris rufis, angulo apicali exteriori nigro.* — Long. 2 —  $2\frac{2}{3}$  lin.

*Mannerh. Brachelytr.* 64. 6. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 500. 4. — *Runde Staphelytr. Hal.* 26. 3.

*Staphyl. striatus* Oliv. Ent. III. 42. 28. 37. t. 5. f. 47.

*Tachinus striatus* Grav. Micr. 146. 17. Mon. 27. 6. — *Latr. Hist. nat. Crust. et Ins.* IX. 398. 17. — *Gyll. Ins. Suec. II.* 272. 18. — *Sahlb. Ins. Fenn. I.* 304. 17.

*Staphyl. angularis* Payk. Faun. Suec. III. 395. 35.

*Staphyl. analis* var.  $\gamma$  Payk. Mon. Staph. 48. 34.

Habitat in Europa.

Antennae capite thoraceque breviores, fortius incrassatae, articulo secundo tertioque aequalibus, 5 — 10 crassis, transversis, ultimo praecedente sesqui longiore, compresso, ovato, fuscae, articulis 4 primis rufo-testaceis, ultimo testaceo. Caput oblongum, nigro-piceum, nitidum, ore rufo-testaceo.

Thorax basi coleopterorum prope latitudine, apice dimidio angustior, latitudine baseos paulo brevior, basi et lateribus leviter rotundatus, angulis posterioribus omnino rotundatis, laevissimus, nigro-piceus, nitidus, margine summo piceo-rufo. Scutellum laevissimum, rufo-piceum, nitidum. Elytra thorace paulo longiora, punctorum seriebus tribus ordinariis, punctis sat remotis, rufa, nitida, macula magna transversa apicali, exterius plerumque latiore nigro-picea, margine apicali ipso summo rufo. Abdomen fortius punctatum, longius pubescens, nigro-piceum, nitidum, segmentis 4 primis margine postico, quinto apice, sexto toto rufis. Pectus nigro-piceum, nitidum. Pedes testacei, coxis anticis concoloribus, tarsis anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore sexto apice sub-bisinuato.

Femina abdominis segmento inferiore sexto apice rotundato.

\* 13. *B. dimidiatus*: *Niger, nitidus, elytrorum basi pedibusque rufis.* — *Long.*  $2\frac{1}{2}$  lin.

Habitat in America septentrionali.

Statura omnino praecedentis, niger, nitidus. Antennae capite thoraceque multo breviores, apicem versus fortiter incrassatae, articulo tertio secundo paulo longiore, 5—10 crassis, sensim paulo crassioribus, transversis, ultimo angustiore, breviter subovato, apice subacuminato, nigrae, articulis tribus primis rufo-testaceis, ultimo ferrugineo. Caput oblongum, laevissimum, ore testaceo. Thorax basi coleopterorum latitudine, apice duplo prope angustior, latitudine baseos parum brevior, basi, lateribus angulisque posterioribus rotundatis, laevissimus. Elytra thoracis longitudine, serie dorsali punctis sat distinctis sex composita, nitida, nigra, basi prope ad medium usque rufa. Abdomen crebrius fortiusque punctatum, griseo-pubescens, segmentis singulis margine apicali rufo-piceo. Pedes rufi, coxis anticis picescentibus.

Mas abdominis segmento inferiore sexto apice sub-bisinuato.

Femina abdominis segmento inferiore sexto apice rotundato.

X \* 14. *B. atricapillus*: *Rufus, nitidus, capite, pectore, ano elytrisque nigris, his lunula basali margineque apicali albidis.* — *Long.*  $2\frac{1}{2}$ —3 lin.

*Mannerh. Brachelytr.* 64. 7. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 497. 1. — *Runde Brachelytr. Hal.* 27. 5. — *Er. Col. March.* I. 408. 6.

*Staph. atricapillus* *Fab. Syst. Ent.* 267. 15. *Spec. Ins.* I. 337. 19. *Mant. Ins.* I. 221. 26. *Ent. Syst.* I. II. 527. 40. *Syst. El.* II. 599. 49. — *Goeze Ent. Beitr.* I. 724. 9. — *Payk. Mon. Staph.* 49. 35. *Faun. Suec.* III. 414. 60. — *Oliv. Ent.* III. 42. 29. 40. t. 4. f. 39. — *Panz. Ent. Germ.* 456. 30. — *Schränk Fauna Boica* I. 645. 877. — *Marsh. Ent. Britt.* 522. 71.

*Tachinus atricapillus* *Grav. Micr.* 148. 19. *Mon.* 30. 14. — *Latr. Hist. nat. Crust. et Ins.* IX. 399. 19. — *Gyll. Ins. Suec.* II. 273. 19. — *Sahlb. Ins. Fenn.* I. 305. 18. — *Guér. Iconogr. Ins.* t. 10. f. 8.

*Oxyporus lunulatus* *Fab. Syst. Ent.* 268. 2. *Spec. Ins.* I. 338. 2. *Mant. Ins.* I. 222. 2. *Ent. Syst.* I. II. 532. 3. *Syst. El.* II. 605. 3. (*forte*) — *Panz. Ent. Germ.* 359. 3. *Faun. Germ.* 22. 15.

*Tachinus lunulatus* *Zetterst. Faun. Lappon.* I. 65. 9. *Ins. Lappon.* 58. 10.

Habitat in Europae boletis.

Antennae capitis thoracisque prope longitudine, apice crassiores, articulo tertio secundo paulo longiore, 5—10 sensim paululum brevioribus et crassioribus, subobconicis, ultimo oblongo-ovato, praecedentibus ambobus longitudine aequali, fuscae, articulis 4 primis ultimoque testaceis. Caput valde elongatum, angustatum, nigrum, nitidum, ore flavo. Thorax basi coleopterorum prope latitudine, latitudine baseos parum brevior, apice plus dimidio quam basi angustior, basi et lateribus leviter rotundatus, angulis posterioribus obtusis, laevissimus, apice punctorum pari notatus, testaceo-rufus, nitidus. Scutellum nigrum, nitidum. Elytra thorace paulo longiora, punctorum remotorum seriebus tribus ordinariis, nigra, nitidissima, macula lunata humerali margineque apicali albidis. Abdomen parce punctatum, rufum, nitidum, segmentis ultimis duobus nigris, penultimo apice albidum. Pectus nigrum. Pedes flavi, coxis anticis concoloribus, tarsis anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore sexto apice truncato.

Femina abdominis segmento inferiore sexto apice rotundato.

Obs. 1. Ox. lunulatus Fabricii ob „abdomen apice nigrum, annulo albo” ad hanc speciem referendus videtur: in hac enim annulus albus in segmenti penultimi margine apicali situs conspicuus, in sequente specie tenuissimus neque nisi oculo armato observandus. Staph. atricapilli nomine signata in Museo Fabriciano Boletochara lunulata exstat.

2. Dom. Prof. Zetterstedt speciem hanc pro vero Staph. lunulato Lin. habet, praecipue quod frequentius obvius sit, descriptio Linneana autem aegre ab hoc insecto desumta.

\* 15. *B. speciosus*: *Rufus, nitidus, capite, pectore, ano elytrisq.ue nigris, his basi margineque apicali inaequaliter testaceis, stria dorsali multipunctata.* — Long. 4 lin.

Habitat in Hungariae montibus, Dom. Schüppel.

Habitu omnino praecedentis, at plus duplo maior. Antennae capitis thoracisque longitudine, apice parum crassiores, subcompressae, articulo tertio secundo sesqui prope longiore, 5—10 subobconicis, sensim paululum brevioribus at vix crassioribus, ultimo praecedente parum longiore, ovato, compresso, nigrae, articulis 4 primis ultimoque testaceis. Caput fortiter elongatum, angustum, nigerrimum, nitidum, ore testaceo. Thorax coleopterorum basi paululum angustior, apice duplo prope quam basi angustior, latitudine baseos vix brevior, basi et lateribus leviter rotundatus, angulis posterioribus obtusis, laevissimus, testaceo-rufus, nitidus. Scutellum testaceum. Elytra thorace tertia fere parte longiora, punctorum seriebus tribus ordinariis, punctis seriei intermediae numerosis (circiter 10), nitidissima, nigra, fascia basali, postice profunde sinuata, margineque inaequali apicali testaceis. Abdomen segmentis 4 primis parce punctatis, nigro-pilosellis, rufis, nitidis, reliquis crebre punctatis, nigerrimis, nitidis, quinto margine apicali tenuissimo albo. Pectus nigerrimum. Pedes pallidi, coxis anticis concoloribus, posterioribus nigris, posticis margine postico albidum.

Mas abdominis segmento inferiore sexto apice leviter emarginato.

Femina latet.



- \* 16. *B. lunulatus*: *Rufus, nitidus, capite, pectore, ano, elytrisque nigris, his basi tota margineque apicali testaceis, serie dorsali parumpunctata.* — Long.  $2\frac{1}{2}$  lin.

*Mannerh. Brachelytr.* 64. 6. — *Runde Brachelytr. Hal.* 27. 4. — *Er. Col. March.* I. 408. 7.

*Staphyl. lunulatus* Linn. *Faun. Suec.* n. 845. *Syst. Nat.* I. II. 684. 7. — *Müll. Faun. Frid.* 23. 225. *Prodr.* 98. 1104. — *Goeze Ent. Beitr.* I. 718. 7. — *De Vill. Ent.* I. 413. 7. — *Payk. Mon. Staph.* 58. 41. *Faun. Suec.* III. 415. 61. — *Marsh. Ent. Britt.* 523. 72.

*Oxyporus lunulatus* Cederh. *Faun. Ingr.* 120. 369.

*Tachinus lunulatus* Grav. *Micr.* 147. 18. *Mon.* 30. 13. — *Latr. Hist. nat. Crust. et Ins.* IX. 398. 18. — *Gyll. Ins. Suec.* II. 274. 20. — *Sahlb. Ins. Fenn.* I. 305. 19.

*Tachinus atricapillus* Zetterst. *Faun. Lappon.* I. 65. 10. *Ins. Lappon.* 58. 11.

Habitat in Europa.

Statura et summa affinitas praecedentis. Antennae capitis thoracisque longitudine, apice leviter crassiores, articulo tertio secundo sesqui longiore, 6 — 10 sensim paululum brevioribus et crassioribus, subobconicis, ultimo ovato, praecedente sesqui longiore, compresso, apice abrupte acuminato, fuscae, articulis 4 primis ultimoque testaceis. Caput modice elongatum, nigrum, nitidum, ore flavo. Thorax basi coleopterorum prope latitudine, apice dimidio quam basi angustior, latitudine baseos vix brevior, basi et lateribus leviter rotundatus, angulis posterioribus obtusis, laevissimus, testaceo-rufus, nitidus. Scutellum testaceum, nitidum. Elytra thorace paulo longiora, punctorum seriebus tribus ordinariis, punctis seriei intermediae paucis, distantibus, nigra, nitidissima, fascia basali integra, elytri quadrantem fere occupante, margineque apicali testaceis. Abdomen parce punctatum, rufum, nitidum, segmentis ultimis duobus nigris, penultimo margine apicali tenuissimo albo. Pectus nigrum. Pedes flavi, coxis anterioribus concoloribus, tarsis anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore sexto apice truncato.

Femina abdominis segmento inferiore sexto apice subacuminato.

- \* 17. *B. cinctus*: *Rufus, nitidus, capite, pectore, ano elytrisque nigris, his basi, sutura margineque apicali testaceis.* — Long.  $2\frac{2}{3}$  lin.

*Tachinus cinctus* Grav. *Micr.* 193. 6. *Mon.* 30. 15.

*Tachinus atricaudatus* Say *Journ. of the Acad. of natur. scienc. of Philadelph.* III. p. 158.

Habitat in America septentrionali.

Statura et summa affinitas praecedentium. Antennae capitis thoracisque longitudine, apice leniter crassiores, subcompressae, articulo tertio secundo sesqui longiore, 5 — 10 subobconicis, sensim paulo crassioribus et paululum brevioribus, ultimo praecedente sesqui longiore, ovato, apice acuminato, fuscae, articulis 4 primis ultimoque testaceis. Caput elongatum, angustum, nigrum, nitidum, ore testaceo. Thorax basi coleopterorum latitudine, antice plus dimidio quam basi angustior, latitudine baseos paulo brevior, basi et lateribus leviter rotundatus, angulis posterioribus obtusis, laevissimus, testaceo-rufus, nitidus. Scutellum rufo-testaceum, nitidum. Elytra thorace paulo longiora, punctorum seriebus ordinariis tribus, punctis seriei interme-

diae paucis, distantibus, nitidissima, nigra, basi tota, sutura basi leviter dilatata, margineque apicali testaceis. Abdomen parce punctatum, nitidum, segmentis quatuor primis rufis, ultimis duobus nigris, penultimo apice testaceo. Pectus nigrum. Pedes flavi, coxis anterioribus concoloribus, posticis nigris, margine postico testaceo.

- X \* 18. *B. trimaculatus*: Niger, nitidus, thorace elytrisque testaceis, illo disco, his angulo exteriori late nigris, antennis pedibusque testaceis, elytrorum serie dorsali multipunctata. — Long.  $2\frac{1}{2}$  lin.

Mannerh. Brachelytr. 65. 9. — Boisd. et Lacord. Faun. Ent. Paris. I. 498. 2.

Oxyporus trimaculatus Fab. Ent. Syst. I. II. 533. 7. Syst. El. II. 605. 7.

Staphyl. trimaculatus Payk. Faun. Suec. III. 422. 68.

Tachinus trimaculatus Gyll. Ins. Suec. II. 275. 21. — Sahlb. Ins. Fenn. I. 306.

20. — Zetterst. Ins. Lappon. 58. 12.

Staphyl. littoreus Payk. Mon. Staph. 80. 55.

Habitat in Suecia, Fennia, Dom. Schüppel.

Sequentibus affinis, at maior, B. cernuo aequalis, medio tamen latior, et apicem utrumque versus magis angustatus. Antennae capitis thoracisque prope longitudine, apice parum crassiores, articulo tertio secundo paulo longiore, 5—10 leviter obconicis, subaequalibus, ultimo elongato-subovato, apice acuminato, testaceae, medio magis minusve fuscrescentes. Caput breviusculum, laevissimum, nigrum, nitidum, ore testaceo. Thorax basi coleopteris distincte angustior, apicem versus leviter angustatus, apice dimidio quam basi angustior, latitudine baseos parum brevior, basi et lateribus parum rotundatus, angulis posterioribus obtusis subrotundatis, laevissimus, nitidus, rufo-testaceus, disco longitudinali basin versus angustato nigro. Scutellum laevissimum, nigrum. Elytra thorace sesqui longiora, luteo-testacea, nitida, angulo posteriore exteriori a suturae apice inde usque ad lateris medium margineque toto inflexo nigerrimo, stria dorsali ordinaria circiter 12-punctata. Abdomen breviusculum, sparsim fortius punctatum, pube longiore depresso grisea parcius vestitum, nigrum, nitidum, segmentis singulis margine apicali piceo-testaceo. Pectus nigerrimum. Pedes testacei, coxis anterioribus concoloribus, posticis nigris, margine posteriore albido, tarsis anticis in utroque sexu simplicibus.

Obs. Differt a B. trinotato et pygmaeo elytrorum stria dorsali multipunctata, a B. exoleto statura multo maiore et duplo latiore ut et colore.

- X \* 19. *B. trinotatus*: Testaceus, nitidus, capite, elytrorum macula scutellari aliaque postica nigris, antennis fuscis, basi testaceis, elytris serie dorsali multipunctata. — Long. 2 lin.

Er. Col. March. I. 309. 8.

Staph. bimaculatus Schrank Faun. Boic. I. 644. 872. (forte)

Habitat in Germania.

Antennae capitis thoracisque longitudine, apice leviter crassiores, articulis 2 et 3 longitudine aequalibus, 4—10 subobconicis, sensim paululum brevioribus et crassioribus, ultimo ovato, apice acuminato, pilosae, fuscae, articulis 4 primis dilute testaceis. Caput oblongum, nigrum, nitidum, labro, palpis reliquisque oris partibus dilute testaceis. Thorax basi coleopteris

angustior, apice basi vix dimidio angustior, latitudine baseos parum brevior, basi et lateribus medio leviter rotundatus, angulis posterioribus obtusis, laevis, rufo-testaceus, nitidus. Scutellum testaceum. Elytra thorace plus sesqui longiora, flavo-testacea, nitida, macula communi circa scutellum aliaque sat magna intra angulum apicalem anteriorem rotundatis nigris, saepius magis minusve obsoletis, serie dorsali ordinaria punctis circiter 8—9 composita. Abdomen rufo-testaceum, segmentis singulis rarius basi piceis, nitidum, supra punctis raris impressum, subtus crebrius fortiter punctatum. Pectus nigrum, nitidum. Pedes flavo-testacei, coxis anterioribus concoloribus, posticis nigris, margine postico testaceo, tarsis anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore sexto longitudinaliter obsolete impresso.

Femina abdominis segmento inferiore sexto simplice.

- \* 20. *B. exoletus*: Testaceus, nitidus, capite nigro, elytris macula apicali obsoleta fusca, antennis ante apicem fuscis, elytris serie dorsali multipunctata. — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I.* 409. 9.

Habitat in Germania, ipse, in Gallia, Dom. Dr. Aubé, in Sardinia, Dom. Prof. Gené.

Statura sequentis et maioribus eius individuis aequalis, at paulo angustior. Antennae capite thoraceque paulo longiores, apice leviter crassiores, articulo tertio secundo paulo longiore, 5—10 leviter obconicis, aequalibus, ultimo ovato, apice acuminato, testaceae, ante apicem paulo obscuriores. Caput breviusculum, nigrum, nitidum, oris partibus testaceis. Thorax basi coleopterorum latitudine, apice dimidio fere angustior, latitudine baseos vix brevior, basi parum, lateribus leviter rotundatus, angulis posterioribus obtusis, testaceus, nitidus. Scutellum testaceum. Elytra thorace plus sesqui longiora, thoracis basi haud latiora, testacea, nitida, apice macula transversa fusca obsoletiore, in angulo exteriori sita, notata, margine apicali summo plerumque testaceo, serie dorsali ordinaria punctis sat confertis, numero circiter 12—14 composita. Abdomen utrinque parce punctatum, testaceum, nitidum, segmentis singulis basi obscurioribus. Pectus piceum. Pedes flavi, coxis anterioribus concoloribus, posticis piceis, margine postico dilutiore, tarsis anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore sexto apice summo subemarginato.

Femina abdominis segmento inferiore sexto apice rotundato.

Variat thoracis disco piceo, elytrorum macula nigerrima, maiore, ultra elytri tertiam partem dilatata.

Obs. Praecedenti et sequenti admodum affinis, ab illo autem statura minore, angustiore, capite brevior etc., ab hoc elytrorum serie dorsali multipunctata satis distinctus.

- \* 21. *B. pygmaeus*: Piceo-testaceus, nitidus, capite elytrorumque macula angulari nigris, antennis fuscis, basi testaceis, elytrorum serie dorsali parumpunctata. — Long.  $1\frac{1}{3}$  — 2 lin.

*Mannerh. Brachelytr.* 65. 10. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 499. 3. — *Er. Col. March. I.* 410. 10.



- Oxyporus pygmaeus* Fab. Spec. Ins. I. 339. 4. Mant. Ins. I. 222. 5. Ent. Syst. I. II. 534. 13. Syst. El. II. 606. 12. — Panz. Ent. Germ. 360. 10. Faun. Germ. 27. 19.
- Tachinus pygmaeus* Gyll. Ins. Suec. II. 277. 22. — Zetterst. Faun. Lappon. I. 66. 11. Ins. Lappon. 58. 13. — Sahlb. Ins. Fenn. I. 306. 21.
- Oxyporus thoracicus* Fab. Spec. Ins. I. 338. 3. Mant. Ins. I. 222. 4. Ent. Syst. I. II. 534. 12. Syst. El. II. 606. 12. Panz. Ent. Germ. 360. 9.
- Tachinus melanocephalus* Grav. Micr. 144. 16. Mon. 27. 7. — Latr. Hist. nat. Crust. et Ins. IX. 398. 16.
- Staphyl. melanocephalus* Marsh. Ent. Brit. 523. 73. (forte)
- Staphyl. trimaculatus* var.  $\beta$ ,  $\gamma$  Payk. Faun. Suec. III. 422. 68.
- Boletobius merdarius* Runde Brachelytr. Hal. 26. 2.
- Staph. pallidus* Rossi Faun. Etr. Mant. 99. 221. (Ed. Hellw. I. 444. 221.) forte.

Habitat in Europa.

Magnitudo variat, specimina maxima B. trinotato aequalia, pleraque autem duplo minora et ultra. Antennae capite thoraceque breviores, apice crassiores, articulis 2 et 3 aequalibus, 5 — 10 subobconicis, sensim paulo crassioribus, ultimo compresso, breviter ovato, apice obtuso, fuscae, articulis 5 primis testaceis. Caput breviusculum, nigrum, nitidum, ore testaceo. Thorax basi coleopteris paulo angustior, apice duplo prope quam basi angustior, latitudine baseos paulo brevior, basi et lateribus leviter rotundatus, angulis posterioribus obtusis, laevissimus, testaceus, nitidus, disco magis minusve picescente. Scutellum piceo-testaceum. Elytra pallide testacea, nitida, macula magna rotundata intra angulum apicalem, rarius obsoleta, picea, nonnunquam etiam circa scutellum picescentia, serie dorsali ordinaria punctis sat distantibus, circiter 5 — 6 composita. Abdomen supra parcius, subtus crebrius sat fortiter punctatum, piceum, nitidum, apice segmentorumque singulorum margine postico rufo-testaceis, nonnunquam omnino rufo-testaceum. Pectus nigro-piceum. Pedes pallide testacei, coxis anterioribus plerumque concoloribus, rarius picescentibus, posticis nigro-piceis, margine postico testaceo, tarsis anticis in utroque sexu simplicibus.

Mas abdominis segmento sexto inferiore medio obsolete tuberculato, pone tuberculum leviter longitudinaliter impresso.

Femina abdominis segmento inferiore sexto simplice.

Variat rarius thorace abdomineque omnino nigro-piceis, frequentius omnino testaceus, capite solo nigro.

### Species mihi invisae.

22. *B. pulchellus*: Rufo-testaceus, nitidus, capite, antennarum medio, pedore, elytrorum fascia postica anòque nigris.

Mannerh. Brachelytr. 64. 8.

Habitat in Fennia, prope Aboam.

B. atricapillo similis, paulo tamen angustior et aliter coloratus. B. lunulato etiam affinis, sed quadruplo minor, nitidior et respectu magnitudinis angustior, antennarum articulo ultimo longiore toto pallido. (Mannerh.)

### **Mycetoporus** Mannerh.

Mannerh. Brachelytr. p. 62.

*Ischnosoma* Steph. Illustr. V. p. 168.

Antennae 11-articulatae, filiformes.

*Palpi maxillares articulo ultimo subulato.*

*Ligula integra, apice rotundata.*

*Elytra pectori longitudine aequalia.*

*Tarsi 5-articulati.*

Corpus elongatum, aut lineare aut utrinque attenuatum, utrinque deorsum curvatum, alatum. Caput thoraci immissum, deflexum, oculis rotundatis, parum prominulis. Labrum subquadratum, apice leviter emarginatum. Mandibulae parvae, apice incurvae, acutae, interne basi membrana ciliatae. Maxillae (t. IV. f. 7. b.) malis coriaceis, interiore interne, exteriori apice barbatis. Palpi maxillares leviter elongati, articulo primo minuto, secundo tertioque inter se longitudine subaequalibus, subobconicis, tertio reliquis crassiore, quarto minuto, tenui, subulato. Labium (t. IV. f. 7. a.) mento transversum, parte membranacea ante palporum basin sat elongata, ligula rotundata, membranacea, paraglossis ligulae connatis, nonnisi linea ciliata distinguendis, eamque haud superantibus. Palpi labiales breviusculi, crassiusculi, articulo primo cylindrico, secundo brevior, subgloboso, tertio minuto, subulato. Antennae ante mandibularum basin insertae, filiformes, apicem versus magis minusve incrassatae. Thorax coleopterorum fere latitudine, antrorsum angustatus, latitudine parum brevior, lateribus rotundatus, basi truncatus, antice pro receptione capitis leviter emarginatus, transversim convexus. Scutellum triangulare. Elytra apice truncata, angulo apicali exteriori oblique truncato. Abdomen aut lineare aut apicem versus sensim attenuatum, deorsum curvatum. Pedes breviusculi, coxis intermediis valde approximatis, posticis amplis, parallelipedis, scutatis, tibiis omnibus spinulosis, tarsis 5-articulatis, posterioribus elongatis, gracilibus, articulo primo elongato, anticis crassiusculis, articulo primo reliquis paulo longiore, in utroque sexu simplicibus.

#### Observationes.

1. Genus praecedenti proximum, at ligula integra et palporum maxillarium articulo ultimo subulato sat distinctum.

2. Caput laevissimum, thorax laevissimus, margine punctis singulis, dorso saepius punctis nonnullis inter medium et latera sitis, apice semper punctis quatuor transversim positis, margine anteriori magis minusve approximatis impressus, scutellum laevissimum, elytra punctorum seriebus tribus ordinariis, una suturali, secunda dorsali, tertia marginali; abdomen parcius fortiusque punctatum.

3. Mas plerumque antennis apice paulo crassioribus et abdominis segmento inferiore sexto minus rotundato, septimoque igitur conspicuo distinguendus.

4. Victus in silvis, sub muscis et foliis deciduis.

\* 1. *M. splendens*: *Oblongus, niger, antennis brevioribus, subclavatis, thorace disco laevissimo, pedibus elytrisque rubris, his punctorum serie dorsali simplice, — Long. 2 lin.*

*Er. Col. March. I. 411. 1.*

*Staphyl. splendens Marsh. Ent. Britt. 524. 75.*

Habitat in Germania.

Habitu fere *Boletobii* cernui, at paulo minor, niger, nitidus. Antennae capite thoraceque paulo breviores, apicem versus fortius incrassatae, apice subcompressae, articulo tertio secundo paulo longiore, 4 — 6 obconicis, sensim crassioribus et paulo brevioribus, 7 — 10 transversis, ultimo breviter

ovato, pilosae, fuscae, articulis duobus primis testaceis. Palpi testacei. Caput breve, subrotundatum. Thorax basi coleopterorum prope latitudine, apice dimidio angustior, latitudine baseos parum brevior, lateribus modice, basi leviter rotundatus, angulis posterioribus rotundatis, modice convexus, ad apicem punctis quatuor transversim positis impressus. Elytra thorace paulo longiora, punctorum seriebus tribus ordinariis, interstitiis laevissimis, saturate rufa, nitida, margine laterali nigro. Abdomen apicem versus leviter attenuatum, pilis brevibus depressis parce pubescens, segmentis singulis fortius punctatis, apice laevioribus, margine postico rufo-piceo. Pedes anteriores rufi, postici rufo-picei, femoribus nigricantibus.

- \* 2. *M. punctus*: *Oblongus, niger, pedibus antennisque testaceis, his ante apicem fusciscentibus, thorace rufo, disco utrinque punctis duobus, elytris rufis, lateribus nigris, intra seriem dorsalem punctis nonnullis impressis.* — Long. 2 lin.

Mannerh. *Brachelytr.* 63. 5. — *Er. Col. March.* I. 412. 2.

*Tachyporus punctus* Gyll. *Ins. Suec.* II. 248. 13. — *Zetterst. Ins. Lapp.* 56. 11.

*Tachinus punctus* Grav. *Mon.* 27. 5. (*forte*)

Habitat in Germania.

Statura omnino praecedentis, punctura vero *M. lepidi*. Antennae capite thoraceque parum breviores, apicem versus leviter incrassatae, articulo tertio secundo paulo longiore, 5 — 10 subcylindricis, sensim paululum brevioribus et crassioribus, penultimis crassitie parum brevioribus, ultimo ovato, apice subacuminato, testaceae, ante apicem flavo-testaceum fusciscentes. Caput breviusculum, nigro-piceum, ore testaceo. Thorax basi coleopterorum fere latitudine, apice duplo angustior, latitudine baseos paulo brevior, basi et lateribus leviter rotundatus, angulis posterioribus subrotundatis, leviter convexus, dorso utrinque punctis duobus, a margine laterali et a medio aequaliter remotis, apice punctis quatuor, transversim positis, notatus, rufus, nitidus. Scutellum piceum, nitidum. Elytra thorace sesqui prope longiora, punctorum seriebus tribus ordinariis, punctis sat distantibus, et praetera interstitio interiore punctis nonnullis, saepius seriem dorsalem accessoriam constituentibus impressa, rufa, nitida, margine laterali nigro. Abdomen apicem versus sensim attenuatum, supra parce fortiter punctatum, parce griseo-pubescens, nigrum, segmentis singulis margine postico late rufo-piceo. Pectus nigrum. Pedes rufo-testacei.

Variat minus maturus testaceus, capite solo nigro-piceo.

- \* 3. *M. longulus*: *Elongatus, niger, nitidus, antennis filiformibus, thorace disco utrinque puncto uno alterove impresso, pedibus elytrisque rubris, his punctorum serie dorsali simplice.* — Long. 2 — 2½ lin.

Mannerh. *Brachelytr.* 63. 4. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 503. 1. — *Er. Col. March.* I. 413. 3.

Habitat in Germania.

*M. splendenti* affinis, at magis elongatus, niger, nitidus. Antennae capitis thoracisque longitudine, apice parum incrassatae, articulo tertio secundo sesqui longiore, 5 — 10 sensim paulo brevioribus at parum crassioribus, omnibus, penultimis etiam, crassitie longioribus, ultimo oblongo-ovato, apice acuminato, fuscae, articulis duobus primis testaceis. Caput breve, apicem



versus angustatum, ore testaceo. Thorax basi coleopterorum latitudine, apice duplo prope angustior, latitudine baseos paulo brevior, basi, lateribus et angulis posterioribus rotundatis, leviter convexus, medio utrinque punctis duobus tribusve, apice punctis quatuor, transversim positis, impressus. Elytra thorace sesqui prope longiora, punctorum seriebus tribus ordinariis, punctis sat distantibus, interstitiis laevissimis, rufa, nitida, margine laterali summoque basali nigro. Abdomen elongatum, apicem versus sensim attenuatum, densius griseo-pubescens, segmentis singulis crebre fortiter punctatis, margine postico rufo-piceo. Pedes anteriores rufi, postici rufo-picei, femoribus nigricantibus.

\* 4. *M. lepidus*: *Elongatus, linearis, piceus, pedibus testaceis, thorace disco utrinque punctis duobus, elytris prope seriem dorsalem punctis duobus impressis.* — Long.  $1\frac{1}{2}$  — 2 lin.

*Mannerh. Brachelytr.* 63. 1. — *Er. Col. March.* I. 413. 4.

*Tachinus lepidus* Grav. *Mon.* 26. 4.

*Tachyporus lepidus* Gyll. *Ins. Suec.* II. 247. 12. — *Zetterst. Faun. Lappon.* I. 59. 7. *Ins. Lappon.* 55. 9. — *Sahlb. Ins. Fenn.* I. 294. 12.

*Staphyl. punctato-striatus* var.  $\alpha$  —  $\gamma$  Payk. *Faun. Suec.* III. 417. 64.

*Staphyl. merdarius* Schrank *Faun. Boic.* I. 647. 884.

*Staph. merdarius* Fab. *Syst. Ent.* 266. 14. *Spec. Ins.* I. 337. 18. *Mant. Ins.* I. 221. 25. *Syst. El.* II. 598. 46. — *Goeze Ent. Beitr.* I. 724. 8. (*forte*)

*Oxyporus merdarius* Fab. *Ent. Syst.* I. II. 533. 11. (*forte*)

*Var. a: Nigro-piceus, elytris rufis.*

*Mycetoporus lepidus* Boisd. et Lacord. *Faun. Ent. Paris.* I. 504. 2.

*Var. b: Nigro-piceus, elytris rufis, macula postica nigricante.*

*Mycetoporus bimaculatus* Boisd. et Lacord. *Faun. Ent. Paris.* I. 505. 3.

*Var. c: Minor, nigro-piceus, antennarum basi pedibusque picescentibus.*

*Tachinus tristis* Grav. *Mon.* 29. 10.

Habitat in Europa.

*Elongatus, angustus, linearis, plerumque iam obscure piceus, iam castaneus, vel interdum testaceus, nitidus. Antennae capitis thoracisque longitudine, apicem versus parum incrassatae, articulo tertio secundo sesqui prope longiore, 5—10 sensim paululum-crassioribus at vix brevioribus, penultimis longitudine paulo crassioribus, ultimo breviter ovato, subacuminato, piceae, basi testacei. Caput breve, rotundatum, nigro-piceum, ore testaceo. Thorax basi coleopteris paulo latior, apice duplo fere angustior, latitudine baseos vix brevior, basi et lateribus modice rotundatus, angulis posterioribus obtusis, subrotundatis, modice transversim convexus, dorso utrinque punctis duobus tribusve, apice punctis quatuor, exterioribus in ipso margine, impressus. Elytra thorace paulo longiora, punctorum seriebus tribus ordinariis, punctis modice distantibus, et prope seriem intermediam interne punctis duobus impressa, plerumque corpori concolora, rarius rufa, interdum macula nigra ante apicem notata. Abdomen elongatum, parallelum, fortius punctatum, parce griseo-pubescens. Pedes testacei vel rufo-testacei.*

Obs. 1. Magnitudine et colore instabilis, statura lineari, elongata facilius cognoscitur.

2. *Staph. merdarius* Fab. secundum specimen in collectione auctoris conservatum, ad hanc speciem pertinet, cum autem ab eo Museum Banksianum citatum sit, non sine dubio descriptionem mancā allegavi.

3. *Staph. brunneus* Marsh. Ent. Britt. 524 74, secundum specimen auctoris ipsius a Dom. Stephens in hoc genere enumeratus, forte ad hanc speciem pertinet. Marsham descriptionem a Paykulo mutuavit, qui hoc nomine *Omalium* significavit, locus Marshamianus igitur neque apud hoc neque apud illud insectum citandus.

\* 5. *M. Americanus: Oblongus, niger, nitidus, antennarum basi, thorace, elytris pedibusque rufo-testaceis, thorace dorso utrinque punctis duobus impresso, elytris punctorum serie dorsali simplice.* — Long.  $1\frac{1}{2}$  lin.

Habitat in America septentrionali, Dom. Zimmermann.

Habitu omnino *M. proni*, at sesqui fere maior. Antennae apicem versus incrassatae, articulo tertio secundo vix aequali, obconico, sequentibus sensim crassioribus, nigrae, basi rufo-testacei (articuli ultimi desunt). Caput parvum, breve, nigro-piceum, nitidum, ore rufo. Thorax basi coleopterorum fere latitudine, apice duplo angustior, latitudine baseos distincte brevior, basi et lateribus leviter rotundatus, angulis posterioribus obtusis, modice transversim convexus, dorso utrinque punctis duobus, a margine laterali et a medio aequaliter distantibus, et paulo ante apicem punctis 4, exterioribus margini magis approximatis, impressus, rufo-testaceus, nitidus. Scutellum demersum, piceum. Elytra thorace paulo longiora, serie dorsali sat regulari, punctis crebrioribus, interstitiis punctis impressis nullis, rufo-testacea, nitida, circa scutellum picea, margine laterali nigro. Abdomen apicem versus sensim attenuatum, crebrius fortiter punctatum, fulvo-pubescent, nigro-piceum, segmentis singulis margine postico rufo-piceo. Pectus nigro-piceum. Pedes rufo-testacei, coxis anticis concoloribus.

Obs. *M. pronus* affinis at paulo maior et thoracis punctis discoidalibus lateralibus distinctus.

\* 6. *M. pronus: Elongatus, rufo-testaceus, nitidus, pectore abdomineque nigricantibus, thorace disco laevissimo, elytris punctorum serie dorsali simplice.* — Long.  $1\frac{1}{2}$  —  $1\frac{3}{4}$  lin.

Er. Col. March. I. 414. 5.

Habitat in Germania.

*M. lepidus* minor minusque linearis. Antennae capite sesqui longiores, apicem versus fortiter incrassatae, articulo tertio secundo tenuiore nec longiore, obconico, sequentibus sensim crassioribus, 6–10 fortiter transversis, ultimo subgloboso, apice obtuso, rufo-testaceae, apice fusciscentes, vel fuscae, basi testaceae. Caput breviusculum, rufo-testaceum, vertice interdum piceo. Thorax basi coleopterorum prope latitudine, apice duplo fere angustior, latitudine baseos paulo brevior, lateribus et basi leviter rotundatus, angulis posterioribus obtusiusculis, modice convexus, ante apicem punctis quatuor, transversim positis, sat distinctis, impressus, disco laevissimus, rufo-testaceus, nitidus. Elytra thorace parum longiora, punctorum seriebus tribus ordinariis, interstitiis laevissimis, rufo-testacea, nitida. Abdomen apicem versus leviter attenuatum, parce fortiter punctatum, parce griseo-pubescent, nigro-piceum, nitidum, apice segmentorumque singulorum margine postico rufis. Pectus nigro-piceum. Pedes testacei.

Mas abdominis segmento inferiore sexto apice truncato, antennisque crassioribus cognoscitur.

Variat 1, Elytris circa scutellum et margine laterali nigricantibus. Genuinis plerumque paulo maior, colore saturatiore, et antennis minus fortiter incrassatis, praecipue in feminis.

2, Capite, ore excepto, piceo. Elytra saepius ut in var. 1, nonnunquam pone medium macula discoidali fuscescente notata. Genuinis paulo minor.

Obs. Distinguitur corporis statura fere fusiformi, antennis crassis vel crassiusculis, elytrorum serie punctorum dorsali simplice et thoracis disco punctis lateralibus nullis, punctis ad apicem impressis autem spatio quodam a margine anteriore remotis.

\* 7. *M. nanus*: Oblongus, niger, nitidus, antennarum basi pedibusque piceis, thorace disco laevissimo, elytris serie punctorum dorsali sesquialtera. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 415. 6.

*Tachinus nanus* Grav. Mon. 28. 8.

Habitat in Germania, Polonia.

Habitu praecedentis omnino, at paulo minor, niger, nitidus. Antennae capite thoraceque breviores, apicem versus leviter incrassatae, articulo tertio secundo vix longiore, sequentibus sensim paululum crassioribus, penultimis vix transversis, ultimo breviter ovato, apice obtusiusculo, compresso, nigrae, basi piceae. Caput oblongum, ore cum palpis piceo. Thorax basi coleopterorum latitudine, apice duplo angustior, latitudine baseos hand brevior, basi et lateribus leviter rotundatus, angulis posterioribus obtusis, modice convexus, apice punctis quatuor transversim positis impressus, disco laevissimo. Elytra thorace paulo longiora, punctorum seriebus tribus ordinariis, serie intermedia punctis crebrioribus composita, interne punctorum serie accessoria dimidia approximata quasi duplicata, margine apicali piceo. Abdomen apicem versus sensim angustatum, parce subtiliter punctatum, parce griseo-pubescentis, segmentis singulis margine postico piceis. Pedes picei.

Variat. Thorace elytris piceis, rufo-piceis, vel testaceis: his caput piceum, ore testaceo, antennae testaceae, basi dilutiores, pedes testacei.

Obs. 1. Specimen thorace elytris rufo-testaceis pro Tachini nani typo benevole communicavit Dom. Prof. Gravenhorst, hoc synonymum igitur indubium, quamvis descriptio anceps videtur: thorax enim punctis 2 dorsalibus impressus potius ad *M. lepidum* spectare videtur. Color etiam typi mihi communicati aegre cum descriptione convenit.

2. A praecedente differt statura paulo minore, antennis parum incrassatis, thoracis punctis ad apicem impressis margini anteriori approximatis, elytrorum seriebus crebre punctatis, intermedia quasi duplicata.

\* 8. *M. lucidus*: Oblongus, piceus, nitidus, thorace disco medio laevissimo, elytris dorso vage punctatis. — Long. 2 lin.

Er. Col. March. I. 415. 7.

Habitat in Germania.

*M. lepidus* longitudine fere aequalis, at in elytris duplo latior, utrinque attenuatus, fusiformis. Antennae capite duplo longiores, apicem versus le-



viter incrassatae, articulo tertio secundo paulo longiore, sequentibus sensim paululum brevioribus et crassioribus, penultimis etiam autem vix transversis, ultimo breviter ovato, apice compresso, rufo-piceae, basi testaceae. Caput breviusculum, nigro-piceum, nitidum, ore testaceo. Thorax basi coleopterorum fere latitudine, apice duplo angustior, latitudine baseos paulo brevior, lateribus modice, basi leviter rotundatus, angulis posterioribus obtusis, leviter convexus, longe ante apicem punctis quatuor, transversim positus, impressus, disco laevissimus, rufo-piceus, nitidus, disco nonnunquam nigricante. Scutellum piceum. Elytra thorace sesqui longiora, striis tribus ordinariis subtiliter regulariterque punctatis, interstitio interiore vage punctato, picea, lucida. Abdomen apicem versus sensim attenuatum, nigro-piceum, nitidum, segmentis singulis margine postico rufo-piceo. Pectus nigro-piceum. Pedes rufo-testacei.

Variat minus maturus colore corporis rufo-testaceo.

Obs. Elytris dorso fere 4-seriatim punctatis et thoracis punctis 4 transversalibus medio inter medium et apicem sitis valde distinctus.

\* 9. *M. splendidus*: *Elongatus, testaceus, vertice, elytrorum, basi, pectore abdomineque piceis, elytris serie punctorum dorsali simplice.* — Long.  $1\frac{1}{2}$  — 2 lin.

Mannerh. *Brachelytr.* 63. 2. — Boisd. et Lacord. *Faun. Ent. Paris. I.* 505. 4. — Er. Col. *March. I.* 416. 8.

*Tachinus splendidus* Grav. *Mon.* 24. 3.

*Tachyporus splendidus* Gyll. *Ins. Suec. II.* 249. 13. — Zetterst. *Faun. Lappon. I.* 60. 8. *Ins. Lappon. I.* 55. 10. — Sahlb. *Ins. Fenn. I.* 295. 13.

Habitat in Europa.

Antennae elongatae, capitis thoracisque longitudine, apice leviter incrassatae, articulo tertio secundo sesqui longiore, 4 — 10 subobconicis, longitudine subaequalibus et sensim perparum crassioribus, ultimo ovato, testaceae, articulo secundo sequentibusque nonnullis fuscis. Caput elongatum, rufo-testaceum, basi plerumque piceum. Thorax basi coleopterorum fere latitudine, apice dimidio angustior, latitudine baseos paulo brevior, lateribus modice, basi vix rotundatus, angulis posterioribus obtusis, vix rotundatis, leviter convexus, apice punctis quatuor transversim positus notatus, disco laevissimus, testaceus, nitidus. Scutellum piceum. Elytra thorace sesqui prope longiora, punctorum seriebus tribus ordinariis, serie intermedia punctis sat distantibus, testacea, basi plerumque picea. Abdomen apicem versus leviter attenuatum, parce punctatum, undique pilosellum, fuscum, apice segmentorumque singulorum margine lato testaceis. Pectus piceum. Pedes elongati testacei.

Mas abdominis segmento inferiore sexto profunde inciso, quinto margine dense piloso.

Femina abdominis segmento inferiore sexto integro, apice leviter rotundato, quinto simplice.

Variat 1, capite nigro-piceo, ore solo testaceo, elytris basi ultra medium piceis.

2, Corpore rufo-testaceo. Hic plerumque genuinis sesqui maior, ceterum autem nullo modo distinctus videtur.

Obs. *M. pallidulus* Mannerh. Brachelytr. 63. 3, „*M. splendido minor*, angustior, elytrorum striis vix conspicuis et articulo antennarum ultimo brevior” vix satis ab hoc differe videtur, forte nonnisi sexu distinctus.

## **Tanygnathus Er.**

*Er. Col. March. I. p. 417.*

*Antennae 11-articulatae.*

*Palpi maxillares elongati, filiformes.*

*Elytra pectori longitudine aequalia.*

*Tarsi 4-articulati.*

Corpus oblongum, utrinque attenuatum, alatum. Corpus thoraci immisum, deflexum, oblongum, oculis mediocribus, ovalibus, parum prominulis. Labrum amplum, apice leviter emarginatum, membranaceum. Mandibulae longae, tenues, falcatae, decussatae. Palpi maxillares valde elongati, tenues, articulo primo minuto, reliquis tribus inter se longitudine aequalibus, ultimo acuminato. Palpi labiales articulo primo secundo paulo brevior, hoc cylindrico, tertio hoc paulo brevior, tenui, acuminato. Antennae capite thoraceque longiores, subrefractae, filiformes, apicem versus haud incrassatae, articulis cylindricis, primo leviter elongato, 2—10 subaequalibus, ultimo iterum paulo longior, apice acuminato. Thorax transversus, basi coleoptera amplexans, apicem versus angustatus. Scutellum sat magnum, triangulare. Coleoptera apice coniunctim leviter late emarginata, ad angulum anteriorem profundius excisa. Abdomen apicem versus angustatum, pilosellum. Pedes breviusculi, validiusculi, intermediis basi approximatis, coxis posticis parvis, angustis, interne scuti instar productis, femorum insertionem obtegentibus, tibiis parce spinulosis, tarsis 4-articulatis, omnibus simplicibus, articulo primo modice elongato, quarto haud brevior, sequentibus duobus sensim minoribus, quarto tenui.

### O b s e r v a t i o n e s .

1. Oris structuram accuratius examinare nondum mihi contigit, genus autem satis distinctum, mandibulis elongatis, falciformibus, decussatis a reliquis huius tribus discedens, habitu intermedium inter Tachyporum et Heterothopem, elytris pectore non longioribus, apice ad angulum anteriorem profunde sinuatis Oxypodam quodammodo referens.

2. Caput et thorax laevissima, glaberrima, polita, hoc punctorum pari uno alterove notato; scutellum, elytra et abdomen punctata et pubescentia.

3. Sexus differentia latet.

4. De victu nihil constat.

\* 1. *T. terminalis*: Niger, nitidus, thorace piceo, palpis, pedibus elytrorumque margine postico rufis, antennis elongatis, nigris, articulo primo testaceo. — Long.  $1\frac{3}{4}$  lin.

Tab. 1. f. 4.

*Er. Col. March. I. 418. 1.*

Habitat Berolini, rarissimus.

Antennae capite thoraceque longiores, nigrae, apice fusciscentes, articulo primo testaceo. Oris partes testaceae. Caput laevissimum, nigrum, nitidum. Thorax basi coleopterorum latitudine, antrorsum angustatus, basi et lateribus leviter rotundatus, angulis posterioribus obtusis, latitudine baseos

paulo brevior, transversim convexus, dorso medio utrinque puncto singulo impressus, ceterum laevissimus, nitidus, disco nigro-piceo, margine laterali basaliue late rufo-piceis. Scutellum confertim subtiliter punctatum, tenuiter pubescens, nigrum. Elytra thorace paulo longiora, confertim subtiliter punctata, pube tenui depressa grisea densius vestita, nigra, parum nitida, margine apicali tenui rufo. Abdomen minus subtiliter minusque dense punctatum, minus subtiliter griseo-pubescens, pilis raris rigidis nigris sparsum, nigrum, subnitidum, segmentis singulis rufo-marginatis. Pectus nigrum. Pedes rufo-testacei, coxis anticis concoloribus.

- \* 2. *T. collaris*: *Niger, nitidus, thorace, scutello palpis pedibusque rufo-testaceis, antennis testaceis, medio fuscis.* — Long.  $1\frac{2}{3}$  lin.

Habitat in Guyana Venezuelensi, Dom. Moritz.

Statura omnino praecedentis, at paulo minor. Antennae capitis thoracisque longitudine, articulo primo testaceo, sequentibus fuscis, ultimis 3 vel 4 ferrugineis. Oris partes pallide testaceae. Caput nigrum, nitidum. Thorax basi coleopterorum latitudine, apicem versus leviter angustatus, basi et lateribus leviter rotundatus, angulis posterioribus obtusis, latitudine parum brevior, transversim convexus, dorso medio et apice utrinque puncto singulo impressus, ceterum laevissimus, rufo-testaceus, nitidus. Scutellum confertim subtiliter punctatum, rufo-testaceum. Elytra thoracis longitudine, confertim subtiliter punctata, pube subtili depressa cinerea vestita, nigra, parum nitida, margine apicali rufescente. Abdomen creberrime subtilius punctatum, fortius densiusque griseo-pubescens, nigro-piceum, subnitidum, segmentorum marginibus rufo-piceis. Pectus piceum. Pedes testacei, coxis anterioribus concoloribus.

Variat thorace piceo, marginibus piceo-rufis.

3. *T. laticollis*: *Rufo-piceus, nitidus, capite, thoracis disco elytrisque nigro-piceis, thorace coleopteris latiore.* — Long. 2 lin.

Habitat in Amer. mer. ins. St. Vincentii, Mus. Dom. Hope.

Antennae capite thoraceque longiores, testaceae, apice flavae. Oris partes pallide testaceae. Caput laevissimum, fusco-piceum, nitidum. Thorax basi coleopteris multo latior, apicem versus fortiter angustatus, basi modice, lateribus fortiter rotundatus, angulis posterioribus rotundatis, latitudine baseos plus dimidio brevior, leviter transversim convexus, dorso ante medium et prope basin utrinque puncto singulo minuto subtilique impressus, punctis posterioribus multo magis distantibus, ceterum laevissimus, piceus, nitidus, margine omni rufescente. Scutellum parce subtilissime punctatum, piceum. Elytra thorace paulo breviora, parcius subtiliter obsoleteque punctata, tenuiter subtiliterque pubescentia, margine laterali nigro-pilosello, picea, nitida, margine summo apicali rufescente. Abdomen crebrius subtiliter punctatum, longius griseo-pubescens, piceo-rufum, nitidulum. Pectus piceo-rufum. Pedes testacei, femoribus anticis basi lineola longitudinali nigra.



## Tribus *tertia*.

### **S t a p h y l i n i n i .**

*Stigmata prothoracica conspicua.*

*Antennae in frontis margine anteriore insertae,*

---

Labrum plerisque bilobum, saepius limbo exteriore membranaceo coriaceove, piloso.

Mandibulae magis minusve prominentes, falcatae, acutae, medio saepius dentatae, basi lacinia membranacea libera, interne ciliata instructae.

Maxillae cardine stipiteque corneis, malis membranaceis coriaceisve, interiore interne, exteriore apice fortius barbatis.

Palpi maxillares 4-articulati, articulo primo minuto, secundo tertioque magis minusve elongatis, quarto his subaequali vel securiformi vel subulato.

Mentum breve, corneum.

Labium membranaceum, exsertum.

Ligula membranacea membranaceave, minuta, iam biloba, iam integra, paraglossis magnis multo brevior plerisque;

sat magna, biloba, paraglossis nullis *Platyprosopo*, *Oxypro*.

Palpi labiales omnibus triarticulati.

Antennae in capitis margine anteriore intra mandibularum basin insertae, 11-articulatae, articulis 3 primis reliquis paulo magis elongatis, filiformes, vel apicem versus sensim paulo crassiores.

Oculi laterales, parum prominentes.

Caput exsertum, saepius porrectum, basi magis minusve constrictum.

Thorax basi lateribusque marginatus, apice truncatus, margine posteriore rotundato coleopterorum basi applicatus.

Prosternum triangulare, spatio pone coxas membranaceo.

Scutellum triangulare.

Elytra pectoris longitudine.

Abdomen marginatum, segmentis sex distinctis, septimo saepius conspicuo, stylis analibus plerumque exsertis.

Pedes plerumque breviusculi, coxis anticis exsertis, conicis, posticis subconicis, trochanteribus posticis simplicibus, tibiis vel omnibus vel posterioribus saepius spinulosis, spinis apicalibus validioribus, tarsis 5-articulatis, anticis saepe dilatatis.

Sexus differentia plerumque abdominis segmento inferiore sexto maris apice emarginato vel inciso, feminae rotundato integroque obvia; tarsi antici etiam maris saepe magis quam feminae dilatati.

#### O b s e r v a t i o.

Remotis Lathrobiis, quae stigmátibus prothoracis occultis et antennarum insertionem differunt, haec tribus cum sectione Brachelytrorum fissilabrium Latreillei et Mannerheimii convenit, at characterem a labro derivato relinquendum putavi, in nonnullis generibus et speciebus (e. g. Quedio molochino) omnino obsoletum. Cum genere Staphylinio Gravenhorstii autem exactissime congruit tota haec tribus, antennarum insertionem et stigmatibus prothoracis conspicuis a reliquis satis determinata.

### Subtribus prima.

## X a n t h o l o n i n i.

*Stigmata prothoracica detecta.*

*Antennae basi inter se non magis quam ab oculis distantes.*

Labrum transversum, bilobum, lateribus coriaceum membranaceumve plerisque;

totum corneum, bilobum Sterculiae;

totum corneum, sexdentatum Scytalino;

integrum, apice membrana elongata porrecta biloba auctum Platyprosopo.

Mandibulae plerumque haud prominentes.

Maxillae mala exteriore coriacea, oblonga, apice haud dilatata Platyprosopo, Othio, Sterculiae;

apice extrorsum leviter dilatata et praeter barbam introrsum vergentem setosa Holiso, Diocho, Xantholino, Leptacino;

segmento inferiore corneo apice extrorsum fortiter dilatato Scytalino.

Palpi maxillares filiformes Platyprosopo, Othio, Sterculiae, Scytalino, Xantholino;

articulo ultimo subulato Holiso, Diocho, Leptacino.

Ligula minuta, integra, rotundata Othio, Holiso, Sterculiae, Scytalino, Xantholino;

minuta, rotundata, medio incisa Diocho, Leptacino;

magna, biloba Platyprosopo.

Paraglossae ligula multo longiores, ciliatae plerisque;

nullae Platyprosopo.

Palpi labiales filiformes Platyprosopo, Othio, Xantholino;

articulo ultimo securiformi Sterculiae, Scytalino;

articulo ultimo subulato Holiso, Diocho, Leptacino.

Antennae rectae Platyprosopo, Othio, Holiso, Diocho;

fractae Sterculiae, Scytalino, Xantholino, Leptacino.

Caput collo quasi tenui brevique thoraci affixum plerisque;  
 basi modice constrictum, thoraci insertum Othio;  
 basi haud aut leviter constrictum, thoraci insertum *Platyprosopo*.

Thorax oblongus, apice truncatus plerisque;  
 oblongus vel elongatus, apice attenuatus *Sterculiae*;  
 trapezoideus *Holiso*.

Elytra sutura integerrima *Platyprosopo*, *Othio*, *Holiso*, *Diocho*;  
 sutura imbricata *Sterculiae*, *Scytalino*, *Xantholino*, *Leptacino*.

Abdomen elongatum, plerisque lineare.

Pedes breviusculi plerisque;

elongati *Sterculiae*;

intermedii basi approximati *Platyprosopo*, *Othio*, *Holiso*, *Diocho*,  
*Leptacino* \* et \*\*\*\*;

basi distantes *Sterculiae*, *Scytalino*, *Xantholino*, *Leptacino* \*\* et \*\*\*,  
 tibiis omnibus spinulosis *Othio*, *Holiso*, *Diocho*;

anticis externe incrassatis et muricato-spinulosis, *Scytalino*, *Xantholino*, *Leptacino*;

anticis inermibus *Platyprosopo*;

omnibus inermibus vel subtilissime spinulosis *Sterculiae*;

tarsis breviusculis,

anticis simplicibus *Holiso*, *Sterculiae*, *Scytalino*, *Xantholino*,  
*Leptacino* \* et \*\*;

anticis dilatatis *Platyprosopo*, *Othio*, *Diocho*, *Leptacino* \*\*\* et \*\*\*\*.

#### Observationes.

1. Antennarum structura et elytrorum sutura in duas sectiones haec subtribus dividi potest, altera antennis rectis et elytrorum sutura integerrima *Platyprosopum*, *Othium*, *Holisum* et *Diochum*, altera antennis fractis et elytrorum sutura imbricata *Sterculiam*, *Scytalinum*, *Xantholinum* et *Leptacinum* continente. Exceptis paucis *Leptacini* speciebus genera illa coxis intermediis approximatis, haec coxis intermediis distantibus conveniunt. *Platyprosopus* labri ligulaeque structura magis forte discrepat.

2. Corpus plerumque magis quam in *Staphylininis* genuinis elongatum, a quibus haec subtribus praecipue antennarum insertionem distinguitur.

\* Antennae rectae. Elytrorum sutura integerrima.

#### ***Platyprosopus* Mannerh.**

*Mannerh. Brachelytr. p. 36.*

*Antennae rectae.*

*Ligula biloba; paraglossae nullae.*

*Palpi omnes filiformes.*

Corpus elongatum, alatum. Caput sat magnum, thoracis fere latitudine, porrectum, basi haud constrictum, oculis parvis, haud prominulis. Labrum (t. IV. f. 25. c.) transversum, membrana biloba, antrorsum producta auctum, lobis acuminatis. Mandibulae sat validae, medio bidentatae. Maxillae (t. IV. f. 25. b.) mala interiore membranacea, interne barbata, exteriori elongata, portione basali cornea, apicali membranacea, apice barbata. Palpi maxillares articulo primo brevior, reliquis longitudine subaequalibus. Labium (t. IV. f. 25. a.) mento transversum, ligula sat elongata, membranacea, biloba,



lobis margine interiore ciliatis, paraglossis nullis. Palpi labiales articulo primo secundoque leviter obconicis, hoc paulo maiore, tertio iterum paulo maiore, apice truncato. Antennae basi ab oculis magis quam inter se distantes, rectae, apicem versus sensim attenuatae, articulo primo leviter elongato, secundo obconico, sequentibus sensim leniter decrescentibus, ultimo apice oblique truncato submarginatoque. Thorax coleopteris latitudine aequalis vel amplior, basi lateribusque rotundatus. Scutellum sat magnum, triangulare. Elytra apice oblique introrsum truncata, sutura integerrima. Abdomen elongatum, apice attenuatum. Pedes breviusculi, validiusculi, tibiis posterioribus spinulosis, anticis dense pubescentibus, tarsis anticis dilatatis, posterioribus articulo primo elongato, sequentibus tribus triangularibus, sensim paululum minoribus.

#### Observationes.

1. Capite basi haud constricto et thorace ampliore ad Tachyporinos habitu, oris structura quodammodo ad Oxytelinos et Omalinos accedit, stigmatibus autem prothoracis detectis conspicuisque et antennarum insertione ab iis distinguitur. Corporis habitu Xantholininos cum Tachyporinis coniungit.

2. In hac subtribu hoc genus valde insigne labro haud sinuato, processu membraneo valde elongato bilobo aucto, et ligula biloba, paraglossis omnino nullis, quoniam cum Oxyporo, generi altero hanc tribum terminante, congruit.

3. Mas abdominis segmento inferiore sexto apice emarginato, femina rotundato distinguuntur.

\* 1. *P. beduinus*: Niger, fusco-pubescent, capite thoraceque glabris, pedibus rufis. — Long.  $6\frac{1}{2}$  — 9 lin.

Nordm. Symb. 133. 1.

Habitat in Nubia, Dom. Prof. Ehrenberg.

Corpus elongatum, subdepressum, nigrum, nitidum, capite thoraceque nitidis. Antennae capite thoraceque paulo breviores, articulo secundo tertio dimidio brevior, rufo-piceae, apice dilutiores, articulo secundo nigricante. Oris partes rufae. Caput magnum, coleopterorum latitudine, subovatum, supra parum convexum, versus latera dense subtiliter, medio parcius subtilissime punctatum, utrinque punctis paucis maioribus sparsum et puncto singulo grossiusculo notatum, glabrum, lateribus et infra confertissime punctatum, fusco-pubescent. Thorax capite et coleopteris latior, latitudine tertia parte brevior, basin versus subangustatus, lateribus leviter rotundatis, basin versus subsinuatis, basi leviter rotundatus, apice truncatus, angulis posterioribus obtusis, anterioribus rectis, leviter deflexis, leviter transversim convexus, confertim subtilissime punctatus, medio utrinque inter medium et latera puncto singulo grossiusculo notatus, glaberrimus. Elytra thorace tertia parte breviora, subdepressa et scutellum confertissime subtiliter punctata, pube depressa fusco-grisea densius vestita. Abdomen coleopteris paulo angustius, apicem versus sensim attenuatum, confertissime subtiliter punctatum, punctis raris maioribus sparsum, pube depressa griseo-fusca dense obductum, parce pilosum, segmento quinto margine apicali, sexto supra apice, infra toto, septimoque obscure rufis. Pedes rufi, coxis anticis concoloribus.

- \* 2 *P. elongatus*: Niger, griseo-pubescent, thorace glabro coleopterorum latitudine, his antennis pedibusque rufis. — Long. 6 lin.

Mannerh. *Brachelytr.* 36. 1. — Nordm. *Symb.* 154. 2.

Habitat Grusia, Dom. Marschall a Bieberstein, in Chersoneso Taurica, Dom Schüppel.

*Elongatus*, subdepressus, niger. Antennae capite thoraceque paulo breviores, articulo tertio secundo sesqui longiore, dilute rufi. Oris partes rufae. Caput thoracis latitudine, oblongum, lateribus subrectis, supra leviter convexum, creberrime subtilius punctatum, linea tenui media longitudinali laeviore, pube depressa grisea densius vestitum. Thorax coleopteris vix latior, latitudine sesqui prope longior, lateribus vix rotundatis, pone medium subsinuatis, basi modice rotundatus, apice truncatus, angulis posterioribus subrotundatis, anterioribus rectis, parum convexus, parce subtilissime punctatus, punctorum maiorum seriebus quatuor longitudinalibus notatus, seriebus intermediis 5—6-, lateralibus 3-punctatis, glaberrimus, limbo laterali confertissime subtiliter punctato, griseo-pubescente. Elytra thorace paulo longiora et scutellum testaceo-rufa, dense subtiliter punctata, pube depressa fulvescentigrisea obducta. Abdomen elongatum, dense subtiliter punctatum, segmentis singulis margine apicali, ultimo apice latiore rufo-piceis. Pectus rufo-piceum. Pedes cum coxis anterioribus rufi.

### **Othius Leach.**

*Steph. Illustr.* V. p. 253. — *Er. Col. March.* I. p. 420.

*Cafius* Boisd. et Lacord. *Faun. Ent. Paris.* I p. 411.

*Antennae rectae.*

*Ligula integra; paraglossae lineares.*

*Palpi omnes filiformes.*

Corpus mediocre, elongatum, lineare, alatum. Caput subovale, basi modice constrictum, collo brevi angusto thoraci insectum, oculis parvis, rotundatis, haud prominulis. Labrum angustum, transversum, medio incisum. Mandibulae breves, medio obtuse dentatae. Maxillae (t. IV. f. 24. b.) mala interiori ampliore membranacea, interne barbata, exteriori tenuiore, basi cornea, subcylindrica, apice coriacea, barbata. Palpi maxillares articulo primo minuto, secundo tertioque longitudine subaequalibus, quarto minore, subacuminato. Labium (t. IV. f. 24. a.) mento transversum, ligula rotundata, integra, membranacea, paraglossis divergentibus, membranaceis, interne ciliatis, ligulam superantibus. Palpi labiales articulis sensim paulo longioribus, ultimo apice subacuminato. Antennae basi ab oculis totidem atque inter se distantes, breviusculae, filiformes, rectae, articulo primo modice elongato, secundo tertioque obconicis, ultimo ovato-subacuminato. Thorax oblongus, basi rotundatus, apice truncatus. Scutellum triangulare. Elytra apice truncata, sutura integerrima. Abdomen lineare. Pedes breves, coxis intermediis contiguis, tibiis spinulosus, tarsis posterioribus articulo primo leviter elongato, anticis articulis 4 primis dilatatis, subtus spongioso-tomentosis.

### **Observationes.**

1. A *Xantholinis*, quibuscum a plerisque auctoribus coniungitur, hoc genus distinctum antennis rectis, elytrorum sutura integerrima, coxis intermediis approximatis et tarsis anticis dilatatis.

2. Dom. Lacordaire recte quidem hoc genus a Xantholino distinxit, nomine autem Cafii, ab Anglis auctoribus aliis huius tribus speciebus tributo, errore quodam inductus, usus est. Com. Dejean in recentissimo collectionis suae Catalogo idem hoc genus nomine Sauriodes distinxit, nomen autem Anglis prius usitatum praevalere debet.

3. Mas abdominis segmento sexto subtus apice leviter emarginato, femina rotundato distinguuntur.

4. Victus in silvis sub musco, foliis deciduis lapidibusque et sub cortice truncorum emortuorum.

\* 1. *O. fulvipennis*: *Niger, nitidus, antennis, pedibus, ano elytrisque rufis, his punctulatis.* — Long.  $4\frac{1}{2}$  —  $5\frac{1}{2}$  lin.

*Er. Col. March. I. 420. 1.*

*Paederus fulvipennis* Fab. *Ent. Syst. I. II. 537. 4. Syst. El. II. 609. 4.* — *Panz. Ent. Germ. 362. 5. Faun. Germ. 27. 3.*

*Staph. fulgidus* Payk. *Mon. Staph. 22. 14. Faun. Suec. III. 377. 14.*

*Staph. fulminans* Grav. *Micr. 47. 70. Mon. 105. 107.* — *Latr. Hist. nat. Crust. et Ins. IX 332. 89.* — *Gyll. Ins. Suec. II. 357. 72.* — *Sahlb. Ins. Fenn. I. 334. 71.*

*Gyrophyn. fulminans* Mannerh. *Brachelytr. 33. 9.* — *Runde Brachelytr. Hal. 11. 7.* — *Nordm. Symb. 119. 14.*

*Cafius fulminans* Boisd. et Lacord. *Faun. Ent. Paris. I. 411.*

*Staph. fulvopterus* Fourc. *Ent. Par. I. 167. 11. (forte)*

*Le Staph. à étuis marons pointillés* Geoffr. *Hist. Ins. I. 365. 11. (forte)*

*Staph. ustulatus* Grav. *Micr. 46. 69.* — *Latr. Hist. nat. Crust. et Ins. IX. 332. 88.*

Habitat in Europa.

Elongatus, linearis, niger, nitidus. Antennae capite paulo longiores, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus et crassioribus, pilosellae, rufae. Palpi rufi. Caput thoracis latitudine, oblongum, antrosum subangustatum, angulis posterioribus fortiter rotundatis, supra parum convexum, utrinque supra oculos parce vageque punctatum, sulculis duobus anticis inter antennas leviter convergentibus, lateralibusque obliquis ante oculos brevioribus obsoletioribusque notatum, inter oculos medio obsolete transversim impressum. Thorax coleopteris perparum angustior, latitudine sesqui longior, basin versus haud angustatus, lateribus pone medium subsinuatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, dorso punctis utrinque tribus, duobus versus apicem, tertio pone medium impressus, versus margines plerumque rufescens. Scutellum laeve. Elytra vix thoracis longitudine, parcius punctata, parce subtiliterque cinereo-pubescentia, rufa, nitida. Abdomen minus crebre subtiliter punctulatum, tenuiter griseo-pubescentis, segmentis singulis margine apicali, et subtus segmento ultimo toto rufis. Pedes cum coxis rufi.

Variat thorace etiam rufo, interdum medio picescente, capite, pectore, abdominisque segmentorum singulorum basi rufo-piceis.

Obs. *Staph. ustulatus* Grav. specimen est coleopteris disco nigricantibus (conf. Grav. *Mon. p. 106.*)

\* 2. *O. melanocephalus*: *Piceus, capite nigro, thorace antennis pedibusque testaceis, elytris punctatis, thorace paulo brevioribus.* — Long.  $2\frac{1}{2}$  lin.

*Er. Col. March. I. 421. 2.*

*Staph. melanocephalus* Grav. *Mon. 107. 111.* — *Gyll. Ins. Suec. II. 360. 74.* — *Sahlb. Ins. Fenn. I. 336. 73.*



*Gyrophyn. melanocephalus* Mannerh. *Brachelytr.* 34. 15. — Nordm. *Symb.* 120. 21.

Habitat in Europa.

Antennae capite paulo longiores, articulo tertio secundo sesqui longiore, 4—10 sensim paulo brevioribus et crassioribus, testaceae. Os rufo-testaceum. Caput thorace paulo angustius, oblongo-ovatum, supra leviter convexum, utrinque supra oculos parce vage punctulatum, sulculis duobus anticis inter antennis profundioribus, subparallelis, lateralibus obliquis ante oculos omnino fere obsoletis, fronte punctis 4 quadratim positis impressa, nigrum, nitidum. Thorax coleopteris perparum angustior, latitudine sesqui longior, basin versus haud angustatus, lateribus pone medium vix sinuatis, apice truncatus, angulis anterioribus deflexis, leviter rotundatis, dorso utrinque punctis duobus, altero ante, altero pone medium, testaceus, nitidus. Scutellum nigrum, laeve. Elytra thorace paulo breviora, fusco-picea, nitida, parcius profundius punctata. Abdomen subtilissime punctulatum, subtiliter parceque pubescens, piceum, nitidum, subtus segmentorum marginibus apiceque testaceis. Pectus piceum. Pedes cum coxis testacei.

\* 3. *O. punctipennis*: Niger, nitidus, pedibus testaceis, elytris punctatis, thorace paulo longioribus. — Long.  $2\frac{1}{2}$  lin.

*Staph. punctipennis* Boisd. et Lacord. *Faun. Ent. Paris. I.* 409. 44.

Habitat in Austria, Helvetia, Italia, Gallia, Anglia.

Statura et summa similitudo praecedentis, niger, nitidus. Antennae capite sesqui longiores, apicem versus haud incrassatae, articulo secundo tertioque aequalibus, 4—10 sensim paulo brevioribus at vix crassioribus, penultimis leviter transversis, obscure ferrugineae, articulis tribus primis nigris. Os piceum. Caput vix thoracis latitudine, subovatum, supra parum convexum, supra oculos utrinque parce punctatum, fronte punctis quatuor quadratim positis notata, sulculis duobus anticis inter antennis parallelis, minus profundis licet sat distinctis, lateralibus ante oculos omnino nullis. Thorax ut in praecedente omnino, at nigerrimus, et puncta in dorso utrinque impressa paulo magis distantia. Scutellum laeve. Elytra thorace paulo longiora, profundius punctata, parce griseo-pubescentia, nigro-picea, margine apicali piceo-testaceo. Abdomen subtilissime parce punctulatum, tenuiter griseo-pubescens. Pedes testacei, coxis posterioribus nigris.

Obs. 1. Praecedenti admodum affinis, praeter colorem autem antennarum articulis 2 et 3 aequalibus, capite minus oblongo, et elytris longioribus sat distinctus.

2. Nomine Othii laeviusculi Kirby (Steph. *Illustr. V.* 255. 8.) ex Anglia a Dom. Rudd benevole communicatus.

\* 4. *O. pilicornis*: Niger, nitidus, antennis pedibusque piceis, elytris laevibus, stria suturali impressa. — Long. 3 lin.

*Er. Col. March. I.* 421. 3.

*Staph. pilicornis* Payk. *Mon. Car. App.* 335. 14—15. *Faun. Suec. III.* 379. 16. — Schrank *Faun. Boic. I.* 642. 868. — Gyll. *Ins. Suec. II.* 359. 73. — Sahlb. *Ins. Fenn. I.* 335. 72.

*Xantholinus pilicornis* Zetterst. *Faun. Lappon. I.* 81. 3. *Ins. Lappon.* 66. 5.

*Gyrophyn. pilicornis* Mannerh. *Brachelytr.* 34. 10. — Nordm. *Symb.* 119. 16.

*Var. b: Rufus, capite, elytris abdominisque segmentis duobus penultimis nigro-piceis.*

*Staph. pilicornis var. b Payk. Faun. Suec. III. 379. 16. — Gyll. Ins. Suec. II. 359. 73. — Sahlb. Ins. Fenn. I. 335. 72.*

*Gyrophypn. pilicornis var. b Mannerh. Brachel. 34. 10.*

*Staph. alternans Grav. Micr. 48. 72. Mon. 107. 109. — Latr. Hist. nat. Crust. et Ins. IX. 333. 91.*

*Gyrophypn. alternans Mannerh. Brachelytr. 31. 12. — Runde Brachelytr. Hal. 12. 8. — Nordm. Symb. 119. 16.*

*Gyrophypn. nigriceps Mannerh. Brachelytr. 34. 11. — Nordm. Symb. 120. 17.*

*Var. c: Fusco-rufescens, subtilus dilutior, capite picescente.*

*Staph. pilicornis var. Payk. Faun. Suec. III. 380. 16. — Gyll. Ins. Suec. II. 359. 73. — Sahlb. Ins. Fenn. I. 335. 72.*

*Xanth. pilicornis var. b Zetterst. Ins. Lappon. 66. 5.*

*Staph. affinis Payk. Mon. Staph. 24. 16.*

Habitat in Suecia, Germania, sub cortice Pinuum.

Linearis, subdepressus. Antennae capite sesqui longiores, apicem versus leviter incrassatae, articulo tertio secundo vix longiore, 4 — 10 sensum crassioribus, 6 — 10 crassis, transversis, ultimo conico, rufo-piceae vel rufae, pilosellae. Palpi rufi. Mandibulae prominentes, rufo-piceae. Caput thorace latius, subquadratum, angulis posterioribus fortiter rotundatis, convexum, laeve, punctis subtilibus 4 inter oculos nonnullisque aliis in fronte impressum, sulcis duobus anticis inter antennis minus profundis, parallelis, apice obsoletis, lateralibus ante oculos omnino nullis, nigro-piceum, nitidum. Thorax coleopteris paulo angustior, latitudine sesqui longior, lateribus subrectis, angulis anterioribus leviter rotundatis, leviter convexus, puncto unico utrinque in medio dorsi, unicoque maiore angulos anteriores versus notatus. Elytra thoracis longitudine, laevia vel parce subtiliterque obsoletissime punctulata, stria suturali impressa. Abdomen laeve. Pedes rufo-testacei.

Variat colore corporis vel nigro-piceo, vel rufo-testaceo, capite elytrisque nigro-piceis et abdominis segmento quarto quintoque fuscis, vel interdum corpore toto testaceo, capite solo magis minusve picescente. Var. secunda apud nos frequentior occurrit.

Obs. *Gyrophypnus nigriceps* Mannerheim statura paulo maiore et capite latiore specie haud satis distinctus.

### Species mihi invisae.

5. *O. macrocephalus: Piceus, nitidus, palpis, antennis, capite postico segmentorumque marginibus rufescentibus, elytrorum dimidia parte anteriore pedibusque rufo-testaceis, thorace utrinque 3-punctato. — Long. 2 lin.*

*Gyrophypnus macrocephalus Nordm. Symb. 124. 26.*

Habitat in America boreali occidentali, ad littora ins. Sithae a cel. Eschscholtz lectus.

Antennae capite longiores, extrorsum sat distincte crassiores, rufae, longius pilosae. Palpi rufescentes. Caput thorace latius, quadratum, antice vix angustatum, supra convexum, nigrum, basi, praesertim lateribus, piceo-

rufescens, nitidum, angulis verticis et regione ophthalmica punctis nonnullis impressum, in antico frontis inter antennis foveola, certo situ conspicua, infra rufescens, glabrum (?), tenue pubescens (?). Clypeus valde angustus, pallide flavescens, labro rufo-ferrugineo. Collum nigro-piceum, nitidum, laeve. Thorax antice subrotundatus, paulo latior quam basi, angulis modice deflexis, lateribus subobliquis, basi attenuatus, postice fere truncatus, supra convexus, nigro-piceus, nitidus, laevis, antice ad latus prope marginem utrinque punctum unum maiusculum, pauloque ab eo in disco puncta duo remota, quorum ultimum obsoletum. Scutellum parvum, apice subacuminatum, nigro-piceum, impunctatum. Elytra thorace paulo latiora, sed vix longiora, supra minus convexa, nitida, antice laete flavo-testacea, dimidia parte posteriore nigra, superficie laevi vel vix conspicue punctulata. Abdomen elytris triplo longius, nigrum, nitidum, vix pubescens, segmentorum marginibus, praesertim infra et ano rufo-ferrugineis. Pedes pallide flavescens. — *O. pilicorni* similis, differt thorace et elytris paulo angustioribus, colore etc. (Nordm.)

### Holus Er.

#### *Antennae rectae.*

*Ligula rotundata, integra, brevissima; paraglossae amplae.*

*Palpi omnes articulo ultimo subulato.*

Corpus elongatum, lineare, valde depressum, alatum. Caput sat magnum, quadratum, angulis rotundatis, basi profunde constrictum, collo quasi brevi tenui thoraci affixum, exsertum, porrectum, oculis parvis, rotundatis, haud prominulis. Labrum transversum, bilobum. Mandibulae apice adunca, medio obtuse dentatae. Maxillae (t. IV. f. 23. b.) mala interiore membranacea, introrsum barbata, exteriore segmento primo triangulari, corneo, altero membraneo, barbato. Palpi maxillares leviter elongati, articulo primo minuto, brevi, secundo tertioque longitudine subaequalibus, hoc apicem versus sensim leviter incrassato, quarto tenui, subulato. Labium (t. IV. f. 23. a.) mento brevi, transversum, ligula brevissima, ultra palporum insertionem haud prominente, paraglossis sat prominulis, acuminatis, membranaceis, pubescentibus. Palpi labiales articulo primo cylindrico, modice elongato, secundo hoc paulo brevior, ultimo praecedenti longitudine aequali, at tenui acutoque. Antennae aequae ab oculis atque inter se distantes, filiformes, rectae, articulo primo reliquis maiore, obconico, incurvo, apicem versus sensim subincrassatae, articulo ultimo ovato. Thorax quadrangulus, angulis rotundatis, basin versus angustatus, basi truncatus, apice utrinque oblique truncatus. Scutellum triangulare. Elytra apice truncata, sutura integerrima. Abdomen lineare, subtilius marginatum. Pedes breves, coxis intermediis contiguis, tibiis omnibus subtiliter spinulosis, tarsis brevibus, articulis 4 primis subaequalibus, crassiusculis, quinto brevi, tenui, unguiculis tenuibus, elongatis, tarsis anticis simplicibus.

### O b s e r v a t i o n e s.

1. Corpore valde depresso, plano Achenio assimilis, at huius tribus, oris structura Xantholino propinquus, at sutura elytrorum integerrima, ligula brevissima, paraglossae ampliores, et palporum utrorumque articulus ultimus subulatus: a *Leptacino* ligula integra distinctus: *Othio* antennis rectis propior, at para-



glossis amplis, ligula brevissima, maxillarum mala exteriore apice dilatata, palporum articulo ultimo subulato et tarsis anticis simplicibus diversus.

2. Sexus differentia latet.

3. Species omnes ex America meridionali allatae. De victu earum nihil compertum.

\* 1. *H. humilis*: *Niger, nitidus, antennis pedibusque piceis, capite thoraceque vage punctatis, linea media longitudinali laevi.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Brasilia.

Piceo-niger, nitidus. Antennae capite paulo longiores, articulo secundo tertioque subaequalibus, 5 — 10 sensim paulo crassioribus, ultimo ovato, apice abrupte acuminato, nigro-picei, basi dilutiores. Palpi rufo-testacei. Caput thorace multo maius, coleopterorum latitudine, oblongo-subquadratum, angulis posterioribus obtusis, subrotundatis, punctis profundioribus aequaliter sparsum. Thorax coleopteris paulo angustior, latitudine antica vix brevior, basin versus angustatus, lateribus leniter rotundatus, ante basin subsinuatus, basi leviter rotundatus, apice utrinque oblique truncatus, angulis anterioribus rotundatis, posterioribus obtusis, planus, supra punctis parvis sparsisque subtilioribus impressus, linea media longitudinali laevi. Elytra thorace tertia parte longiora, dense subtilius punctata, tenuiter griseo-pubescentia. Abdomen parcius minus subtiliter punctatum, tenuiter griseo-pubescent, segmento ultimo apice piceo. Pedes picei.

\* 2. *H. analis*: *Niger, nitidus, ano rufo, femoribus piceis, capite thoraceque utrinque parce punctatis.* — Long.  $3\frac{1}{2}$  lin.

T. 1. f. 5.

Habitat prope Caracas Columbiae, Dom. Moritz.

Habitu omnino praecedentis, at multo maior, niger, nitidus. Antennae capite plus sesqui longiores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus et crassioribus, ultimo ovato, apice abrupte acuminato, pilosellae, nigrae. Palpi rufo-picei. Caput thoracis latitudine, subquadratum, angulis posterioribus subrotundatis, supra punctis rarioribus vagis profundis impressum, secundum marginem lateralem punctorum serie sat regulari notatum. Thorax latitudine anteriore paulo brevior, basin versus angustatus, basi truncatus, apice utrinque oblique truncatus, lateribus leviter rotundatus, angulis omnibus obtusis, planus, dorso utrinque punctis rarioribus vagis profundis impressus. Elytra thorace sesqui longiora, plana, dense subtiliter punctata, tenuiter griseo-pubescentia. Abdomen sat crebre minus subtiliter punctatum, tenuiter pubescens, lateribus parce pilosellum, segmento ultimo apice dimidiato-testaceo. Pedes picei, tibiis nigricantibus.

3. *H. debilis*: *Piceus, nitidus, antennarum basi, pedibus, elytris anoque testaceis, thorace utrinque capiteque nigro crebrius punctatis.* — Long. 2 lin.

Habitat in Americae meridionalis ins. St. Johannis, Dom. Moritz.

Statura omnino *H. humilis*, at paululum maior. Antennae capite paulo longiores, rufo-piceae, articulis duobus primis testaceis. Os rufo-piceum, palpis testaceis. Caput thorace paulo latius, subquadratum, angulis posterioribus leviter rotundatis, planum, crebrius sat profunde vage punctatum, fronte

antice sulcis duobus subtilibus obsoletioribus obliquis notata, nigrum. Thorax latitudine antica haud brevior, basin versus angustatus, lateribus leviter rotundatus, basi truncatus, apice utrinque oblique truncatus, angulis anterioribus rotundatis, planus, dorso utrinque vage punctatus, linea lata longitudinali media laevi, piceus, nitidus. Elytra dense subtiliter punctata, tenuiter griseo-pubescentia, luteo-testacea, nitida. Abdomen minus dense punctatum, tenuiter pubescens, nigro-piceum, nitidulum, segmento ultimo toto, penultimo apice, reliquis margine postico testaceis. Pectus et pedes flavo-testacea.

4. *H. Guildingii*: *Niger, nitidus, antennis pedibusque rufo-piceis, elytris pallidis, basi fuscis, thorace capiteque utrinque parce punctatis, hoc antice bistriato.* — Long.  $2\frac{1}{2}$  lin.

Habitat in Americae meridionalis ins. St. Vincentii, Mus. Dom. Hope.

Habitu *H. humilis*, at duplo maior, niger, nitidus. Antennae capite paulo longiores, articulo secundo tertioque subaequalibus, 4 — 10 sensim paulo brevioribus et crassioribus, ultimo apice abrupte acuminato, rufo-piceae, basi dilutiores. Palpi rufo-testacei. Caput thorace paulo latius, subquadratum, angulis posterioribus leviter rotundatis, utrinque parcius punctatum, punctis oblongis et sat profundis, antice inter antennas sulcis duobus parallelis, sat profundis, in fronte desinentibus, impressum. Thorax coleopteris angustior, latitudine paulo longior, basin versus subangustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus rotundatis, depressus, utrinque parce subtilius, dorso sub-biseriatim punctatus. Elytra thorace paulo longiora, crebrius subtiliter punctata, tenuiter pubescentia, pallide flavo-testacea, basi fusca. Abdomen parce obsoleteque punctulatum, segmentis singulis margine tenui apicali piceo. Pedes rufo-picei.

### **Diochus Er.**

*Antennae rectae.*

*Ligula emarginata; paraglossae lineares.*

*Palpi omnes articulo ultimo subulato.*

Corpus pusillum, leviter elongatum, lineare, minus depressum, alatum. Caput oblongum, antrorsum leviter angustatum, basi truncatum, collo quasi brevissimo tenui thoraci affixum, exsertum, liberum, oculis parvis, rotundatis, ante capitis medium sitis. Labrum angustum, transversum, medio sinuatum. Mandibulae breves, haud prominentes. Maxillae mala interiore membranacea, interne barbata, exteriori cornea, apice membranacea barbataque. Palpi maxillares articulo primo parvo, sequentibus duobus longitudine subaequalibus, tertio crassiore, quarto minuto, subulato. Labium mento transversum, ligula brevi, rotundata, medio emarginata, membranacea, paraglossis ligulam superantibus, membranaceis. Palpi labiales articulo secundo primo paulo longiore, ultimo minuto, subulato. Antennae inter se aequae atque oculis approximatae, breves, crassiusculae, rectae, articulo primo reliquis paulo maiore, secundo tertioque subaequalibus, obconicis, sequentibus brevibus, transversis, ultimo subacuminato. Thorax oblongo-subquadratus, basi angulisque posterioribus rotundatus. Scutellum triangulare. Elytra apice truncata, sutura integerrima. Abdomen lineare. Pedes breves, coxis inter-

mediis approximatis, tibiis subtiliter spinulosis, tarsis anticis leviter dilatatis, posterioribus articulo primo elongato.

O b s e r v a t i o n e s.

1. Habitus huius insecti fere minimi Philonthi, at antennae basi minus distantes et caput a thorace magis remotum. Oris structura omnino cum Leptacino (conf. t. IV. f. 22.) convenit, discedit autem ab eo antennis rectis et sutura elytrorum integerrima. A genere praecedente paraglossis linearibus, palporum articulo ultimo multo brevior et tenuior et tarsis anticis dilatatis, ab Othio palporum articulo ultimo subulato, ab utroque ligula biloba et corporis habitu distinguitur.

2. Sexus differentia latet.

3. De victu nihil constat.

\* 1. *D. nanus*: *Nigro-piceus, nitidus, antennis pedibusque testaceis, capite thoraceque oblongis, politis, hoc 4-seriatim parce punctato, elytris parce obsoleteque punctatis.* — Long.  $1\frac{1}{3}$  lin.

T. 1. f. 6.

Habitat in variis Americae meridionalis regionibus, e Columbia a Dom. Moritz allatus.

Minutus, nigro-piceus, capite, thorace elytrisque nitidissimis. Antennae capite parum longiores, crassiusculae, articulo tertio secundo paulo minore, 4 — 10 transversis, subaequalibus, ultimo subovato, apice abrupte acuminato, testaceae, basi dilutiores. Os testaceum. Caput oblongum, thorace paulo angustius, basi truncatum, angulis posterioribus subrotundatis, antrorsum leviter angustatum, postice laevissimum, anterius parce subseriatim punctatum, sulculis duobus apicalibus brevibus obsoleteque, lateralibus ante oculos omnino nullis. Thorax coleopteris parum angustior, latitudine paulo longior, basin versus haud angustatus, lateribus subrectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, leviter transversim convexus, seriebus quatuor punctorum subtilium, duobus dorsalibus, approximatis, 4-punctatis, reliquis utraque laterali, 4-punctata, ceterum laevissimus, dilutius piceus. Elytra thorace haud longiora, parce obsoleteque seriatim punctata. Abdomen breviusculum, omnium subtilissime punctulatum, segmentis singulis margine apicali piceo. Pedes cum coxis testacei.

Variat thorace rufo-testaceo, capite elytrisque fusco-piceis, his apice indeterminate rufo-testaceis, abdominis singulis segmentis margine apicali lato rufo-piceis.

\*\* Antennae fractae. Elytrorum sutura imbricata.

**Sterculia Laporte.**

*Laporte Etud. Ent. p. 118.*

*Agrodes Nordm. Symb. 161. 29.*

*Aracocnemis Nordm. Symb. 163. 30.*

*Antennae fractae.*

*Ligula integra.*

*Labrum totum corneum, bilobum.*

Corpus maiusculum, elongatum, alatum. Caput sat magnum, exsertum, pedunculo angusto thoraci affixum, porrectum, oculis minutis, lateralibus,



inter medium et apicem sitis. Labrum angustum, transversum, medio sinuatum, totum corneum. Mandibulae longiores brevioresve, basi denticulatae, et ante denticulum membrana ciliata instructae. Maxillae (t. IV. f. 19. b.) malis brevibus, coriaceis, introrsum barbatis. Palpi maxillares minus elongati, crassiusculi, articulo primo minuto, angusto, secundo leviter elongato, obconico, quarto subovato. Labium (t. IV. f. 19. a.) mento transversum, ligula minuta, rotundata, integra, coriacea, paraglossis sat amplis, ligulam longe superantibus, membranaceis, intus ciliatis. Palpi labiales articulis duobus primis subaequalibus, tertio crassiore, subovato. Antennae intra mandibularum basin insertae, inter se aequae atque oculis approximatae, fractae, articulo primo elongato, secundo brevi, obconico, tertio iterum elongato, primo autem duplo fere brevior, reliquis inter se arcte coactis, 4—10 transversis, ultimo obtuse acuminato, interdum apicem versus leniter incrassatae. Thorax oblongus, coleopteris angustior, apice angustatus. Scutellum triangulare. Elytra truncata, sutura imbricata. Abdomen mediocriter elongatum. Pedes elongati, graciles, coxis intermediis late distantibus, tibiis pubescentibus, inter pubem subtilissime spinulosis, anticis haud incrassatis, tarsis omnibus simplicibus, posterioribus articulis duobus primis subelongatis.

#### O b s e r v a t i o n e s.

1. Genus valde insigne, habitu thorace antrorsum attenuato et pedibus elongatis a *Xantholino* discedens, a quo ceterum praecipue labro lateribus haud membranaceo, ligula brevior, palporum structura et tibiis haud distincte spinulosis, anticis haud incrassatis differt.

2. Prof. Nordmann genera duo *Agrodem* palpis labialibus truncatis, et *Araeocnemum* palpis filiformibus distinxit. Palporum structura autem omnes fere species magis minusve inter se discedunt, quam ob rem hoc genus omnino naturale dissolvi nequit. In Catalogo Dejeaniano nomine *Plochionoceri* signatum est.

3. Sexus differentia latet.

4. Victus latet. Species hucusque cognitae in America meridionali obviae.

\* 1. *St. coelestina*: *Coerulea, nitida, thorace elytris brevior, capite lato, subquadrato, mandibulis capitis longitudine*. — Long. 11 lin.

Habitat in Brasilia, Dom. Virmond.

*St. fulgenti* duplo maior, laete viridi-coerulea, nitida, tenuiter nigropubescent. Antennae apicem versus vix incrassatae, articulo tertio elongato, secundo plus duplo longior, 4—10 leviter transversis, inter se aequalibus, ultimo acuminato, nigrae, articulis tribus primis cyaneis. Palpi maxillares articulo ultimo penultimo paulo brevior, labiales articulo tertio fortiter securiformi. Mandibulae capitis longitudine. Caput thorace latius, basi leviter, angulis posterioribus fortius rotundatum, lateribus subrectis, neque retrorsum, neque antrorsum angustatum, latitudine vix angustius, posterius levissime convexum, confertissime punctatum, longitudinaliter subrugulosum. Thorax coleopterorum basi dimidio angustior, latitudine baseos plus sesqui longior, antrorsum angustatus, lateribus ante medium leviter rotundatis, modice transversim convexus, utrinque subtilissime disperse punctatus, spatio medio longitudinali laevissimo, postice utrinque oblique impressus, margine angulari posteriore late leviter elevato. Scutellum punctulis nonnullis impressum. Elytra thorace paulo longiora, apicem versus sensim paulo latiora, subtiliter parce punctata. Abdomen breviusculum, apicem versus sen-

sim angustatum, depressiusculum, late marginatum, subtiliter parce punctulatum, lateribus et subtus breviter nigro-pilosellum. Alae nigro-violaceae.

\* 2. *St. fulgens*: *Cyanea, nitida, thorace elytris brevior, capite oblongo, mandibulis capite brevioribus.* — Long. 7 lin.

D'Orbigny It. Amer. mer. Ins. t. 5. f. 6.

*Staph. fulgens* Fab. Ent. Syst. I. II. 522. 13. Syst. El. II. 592. 15. — Grav. Mon. 91. 87.

*Araeocnemis fulgens* Nordm. Symb. 164. 1. t. 2. f. 8.

*Staph. violaceus* Oliv. Ent. III. 42. 8. 3. t. 1. f. 8.

Habitat Jalapae in Mexico, Dom. Deppe, Cayennae et Surinami sec. Fab. et Oliv.

Corpus violaceo-cyaneum, nitidum, tenuiter nigro-puberulum, subtus laetius cyaneum, abdomine certo sito viridi-nitente. Antennae apicem versus vix incrassatae, articulo tertio secundo duplo prope longiore, 4 — 10 sensim paululum crassioribus, ultimo apice abrupte acuminato, nigrae, articulis tribus primis cyaneis. Palpi maxillares articulo quarto tertio aequali, labiales articulo tertio securiformi. Mandibulae capite breviores. Caput thorace paulo latius, latitudine paulo longius, basi cum angulis posterioribus rotundatum, lateribus subrectum, neque antrorsum neque basin versus angustatum, subdepressum, creberrime fortius rugoso-punctatum, opacum. Thorax coleopterorum basi dimidio angustior, latitudine sesqui fere longior, antrorsum subangustatus, lateribus leviter, angulis anterioribus deflexis fortiter rotundatis, modice transversim convexus, subtilissime disperse punctatus, spatio longitudinali medio et plaga utrinque laterali laevissimis, postice utrinque oblique impressus, margine angulari posteriore elevato. Scutellum punctulis nonnullis impressum. Elytra thorace longiora, subtiliter parce punctata. Abdomen apicem versus parum angustatum, depressiusculum, margine laterali sat elevato, subtiliter parce punctatum. Alae nigro-violaceae.

\* 3. *St. elegans*: *Cyanea, nitida, thorace angusto, elytrorum longitudine, capiteque elongatis, hoc basi attenuato, mandibulis capite triplo brevioribus.* — Long. 8 lin.

*Agrodes elegans* Nordm. Symb. 162. 1. t. 2. f. 9.

Habitat in Para interiore Brasiliae, Sieber.

Magnitudo praecedentis. Corpus nigro-pilosellum, coerulescenti-cyaneum, subtus viridi-coeruleum. Antennae apicem versus haud incrassatae, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus, ultimo apice acuminato, nigrae, articulis tribus primis cyaneis. Palpi maxillares articulo quarto tertio longiore, labiales articulo tertio cylindrico, apice acuminato. Mandibulae capite multo-breviores. Caput thorace paulo latius, elongato-subovatum, latitudine duplo longius, basi attenuatum, supra perparum transversim convexus, creberrime punctatum, utrinque longitudinaliter subrugosum, apice ante oculos laeviusculum. Thorax angustus, elongatus, coleopterorum basi duplo fere angustior, latitudine triplo fere longior, antrorsum sensim attenuatus, modice transversim convexus, subtilissime disperse punctatus, linea media longitudinali laevi, pone medium utrinque oblique leniter impressus. Elytra thoracis longitudine, subtilissime pun-

ctulata. Abdomen crassiusculum, densius subtiliter punctatum. Alae nigro-violaceae.

- \* 4. *St. Leprieuri*: *Coerulea, nitida, pubescens, thorace elytrorum longitudine, capite oblongo-ovato, mandibulis capite triplo brevioribus.* — Long. 6 lin.

*Laporte Etud. ent. I. 118. 1. (perperam descripta.)*

Habitat in Cayenna, Dom. Buquet.

Praecedentibus minor, laete coerulea, nitida, elytris cyaneis, subtiliter nigro-puberula, elytrorum basi margineque apicali abdominisque segmentorum marginibus longius cinereo-pubescentibus. Antennae apicem versus leviter incrassatae, articulo tertio secundo duplo prope longiore, 4 — 10 sensim paululum crassioribus, transversis, ultimo crassiusculo, subovato, apice acuminato, nigrae, articulis tribus primis coeruleis. Palpi maxillares articulo tertio quarto brevior, labiales articulo tertio subsecuriformi, s. apicem versus subincrassato, apice truncato. Mandibulae capite multo breviores. Caput thorace haud latius, latitudine duplo prope longius, basin versus magis quam antrorsum angustatum, oblongo-obovatum, subdepressum, confertissime punctatum, opacum. Thorax basi coleopteris duplo prope angustior, latitudine baseos plus duplo longior, lateribus pone medium rectis, parallelis, a medio usque ad apicem sensim attenuatus, leviter transversim convexus, utrinque subtilissime densius punctatus, spatio longitudinali medio angusto laevi, pone medium utrinque oblique impressus. Elytra thoraci longitudine aequalia abdomenque apicem versus subangustatum densius subtiliter punctata.

- \* 5. *St. formicaria*: *Nigro-subaenea, opaca, griseo-pubescens, capite oblongo-ovali.* — Long. 6 — 7 lin.

*Laporte Etud. ent. I. 119. 2. t. 3. f. 3.*

*Araeocnemis flagellicornis Nordm. Symb. 165. 2.*

*Var. Densius pubescens.*

*Araeocnemis pubescens Nordm. Symb. 167. 3.*

Habitat in Brasilia, Dom. Sellow, Surinami, Dom. Cordua, Cayennae sec. Laporte, in Columbia, Dom. Moritz.

Corpus nigro-subaeneum, opacum, in illaesis speciminibus pube griseo-lutescente obductum. Antennae apicem versus sensim incrassatae, articulo tertio secundo vix sesqui longiore, 4 — 10 transversis, sensim paululum crassioribus, ultimo apice obtusius acuminato, nigrae, articulo secundo basi rufo. Mandibulae capite triplo breviores, validae. Caput thorace haud latius, latitudine plus sesqui longius, basi cum angulis posterioribus fortiter, lateribus leviter rotundatum, subtilius confertissime punctatum. Thorax basi coleopteris duplo fere angustior, latitudine baseos duplo fere longior, lateribus paulo ante medium subangulato-rotundatis, pone medium subsinuatatis, antice angustatus, leviter convexus, omnium confertissime punctatus, linea media longitudinali laevi elevata, pone medium tenui. Elytra thoracis longitudine, subtilissime dense punctata. Abdomen subparallelum, crassiusculum, subtiliter crebre punctatum, nitidulum. Alae fuscae.



## Scytalinus Er.

*Antennae fractae.*

*Labium totum corneum, sexdentatum.*

*Palpi maxillares filiformes, labiales articulo ultimo securiformi.*

*Ligula integra.*

Corpus valde elongatum, lineare, subcylindricum, alatum. Caput subquadratum, angulis rotundatis, supra leviter convexum, basi profunde constrictum, collo quasi brevi cylindrico thoraci affixum, exsertum, porrectum, oculis parvis, rotundatis, vix prominulis. Labrum angustum, transversum, totum corneum, 6-dentatum, margine fortiter setosum. Mandibulae sat fortes, subfalcatae, medio fortiter dentatae. Maxillae (t. IV. f. 20. b.) mala inferiore brevi, coriacea, introrsum barbata, exteriore cornea, depressa, triangulari, angulo exteriore producto, setoso, interiore lobo coriaceo barbato instructo. Palpi maxillares filiformes, articulo primo minuto, reliquis tribus subaequalibus. Labium (t. IV. f. 20. a.) mento transversum, ligula brevissima, integra, membranea, paraglossis eam longe superantibus, coriaceis. Palpi labiales articulo primo parvo, secundo leviter obconico, tertio modice securiformi. Antennae basi inter se magis quam oculis approximatae, geniculatae, articulo primo modice elongato, leviter incurvo, subclavato, secundo tertioque leviter obconicis, hoc praecedente sesqui longiore, articulis reliquis brevibus, pubescentibus, ultimo breviter ovato, subacuminato. Thorax coleopteris paulo angustior, latitudine longior, basin versus leviter angustatus, basi rotundatus, apice truncatus, angulis posterioribus rotundatis, anticis subrectis, subdeflexis. Scutellum triangulare. Elytra apice truncata, sutura imbricata. Abdomen lineare. Pedes breves, postici anterioribus paulo debiliores, coxis intermediis distantibus, tibiis spinulosis, anticis leviter incrassatis, extus spinulis densioribus muricatis, tarsis omnibus simplicibus, articulis 1 — 4 subobconicis, sensim decrescentibus.

### Observationes.

1. Habitu omnino Xantholini, et huic generi nimis fere affinis, distinguendus autem videtur: 1, labro lateribus haud membranaceo, medio haud profundius inciso, apice 4-, lateribus utrinque 1-dentato; 2, maxillarum malae segmento inferiore corneo apice extrorsum fortiter producto; 3, palpis labialibus articulo ultimo securiformi.

2. Sexus differentia latet.

\* 1. *Sc. serpentinus*: Niger, nitidus, abdomine apice summo rufo, capite subquadrato, fronte inaequaliter punctata. — Long. 8 lin.

T. 1. f. 7.

Var. *Elytris rufis.*

Habitat in Brasilia, Dom. Prof. Germar.

Corpus elongatum, subcylindricum, nigrum, nitidum, elytris aut concoloribus aut rufis, abdomine segmenti quinti margine sextique apice rufis. Antennae capite paulo longiores, articulo tertio modice elongato, secundo sesqui longiore, 4—10 sensim paulo crassioribus, et paulo magis transversis, piceae, articulis tribus primis nigris. Caput thorace latius et brevius, subquadratum, parum depressum, lateribus laevissimum, fronte sparsim punctata, plaga media inaequali laeviore, ante utrinque longitudinaliter subim-

pressa. Thorax coleopteris angustior, latitudine plus sesqui longior, basin versus modice angustatus, lateribus subrectis, apice utrinque oblique truncatus, angulis anterioribus rectis, transversim modice convexus, laevis, apice haud procul ab angulo anteriore utrinque puncto profundo, basi medio lineola brevissima subtili impressus. Scutellum apice 4-punctatum. Elytra thorace paulo longiora, dorso fortius subseriatim punctata, ad latera laevissima, lateribus inflexis profundius punctatis. Abdomen supra nitidum, subtus opacum, utrinque punctis raris subtilibus sparsum, parce griseo-pubescent, apice rufo longius fulvo-pilosum. Tarsi picei.

### **Xantholinus Dahl.**

*Encycl. Meth. X. p. 475. — Zetterst. Faun. Lappon. I. p. 80. Ins. Lappon. p. 66.*

*— Boisd. et Lacord. Faun. Ent. Paris. I. p. 411. — Er. Col. March. I. p. 422.*

*Gyrohypnus Kirby Mannerh. Brachelytr. p. 33. — Steph. Illustr. V. p. 258. — Nordm. Symb. p. 113.*

*Eulissus Mannerh. Brachelytr. p. 35.*

*Antennae fractae.*

*Labium bilobum, lateribus membranaceum.*

*Palpi omnes filiformes.*

*Ligula integra.*

Corpus elongatum, lineare, plerumque subdepressum, alatum. Caput plerisque oblongum, basi profunde constrictum, collo quasi brevi cylindrico thoraci affixum, exsertum, porrectum, oculis parvis, rotundatis, apicem versus positus, vix prominulis. Labrum angustum, transversum, medio profunde sinuatum, apice setosum, corneum, limbo laterali membranaceo. Mandibulae breviores, medio dentatae, basi lacinia triangulari membranacea, interne ciliata. Maxillae (t. IV. f. 21. b.) mala interiore brevior, coriacea, interne barbata, exterior cornea, obconica, apice barbata. Palpi maxillares filiformes, articulo primo parvo, sequentibus tribus subaequalibus vel ultimo praecedentibus parum brevior, apice subacuminato. Labium (t. IV. f. 21. a.) mento transversum, ligula brevi, rotundata, integra, membranacea, paraglossis ligulam superantibus, membranaceis. Palpi labiales filiformes, articulis sensim longioribus, ultimo apice subacuminato. Antennae basi inter se non magis quam ab oculis distantes, breviusculae, geniculatae, articulo primo modice elongato, secundo tertioque leviter obconicis, hoc plerumque longior, reliquis brevibus, pubescentibus, ultimo ovato, apice subacuminato. Thorax plerumque oblongus, basin versus subangustatus, basi rotundatus, apice truncatus, angulis anterioribus saepius lateraliter prominentibus. Scutellum triangulare. Elytra apice truncata, sutura imbricata. Abdomen lineare. Pedes breves, postici anterioribus debiliores, coxis intermediis late distantibus, tibiis spinulosis, anticis subincrassatis, extus spinuloso-muricatis, tarsis omnibus simplicibus, articulo primo secundoque subaequalibus, intermediis paulo longioribus.

### **Observationes.**

1. Caput antice sulculis quatuor abbreviatis impressum, duobus interioribus, inter antennarum basin adscendentibus, frontem versus plerumque convergentibus, saepius distinctioribus, alteris duobus exterioribus obliquis ante oculos sitis, plerumque obsoletioribus. Thorax saepe punctorum seriebus quatuor notatus, seriebus dorsalibus rectis, lateralibus lituis s. sinuatis, antice circumflexis, Elytra dorso sae-

pius subseriatim punctata, secundum marginem lateralem laevia, margine laterali inflexo iterum punctato.

2. Nomen Xantholinus praetuli, prius a Com. Dejean vulgatum, et ab auctoribus Encyclopediae methodicae et a Latreillio (Regn. An. IV. p. 435.) receptum; praeterea nomen Gyrohypnus perperam compositum. Eulissus Mannerheim mihi nullo modo ab hoc genere distinguendus videtur: idem iam monuit Dom. Brullé. (Hist. nat. Ins. VI. p. 72.)

3. Mas saepius abdominis segmento inferiore sexto apice truncato, septimo paulum conspicuo, femina segmento inferiore sexto apice rotundato, septimo occulto, distinguuntur.

4. Victus in silvis et campis sub lapidibus, foliis deciduis muscisve et in fimo. Corpus in quiete spirae instar convolutum.

5. Larvae habitu et moribus omnino imaginum, quibuscum occurrunt. Xantholini punctulati larvam Dom. Bouché observavit et hoc modo descripsit: Corpus lineare, posterius attenuatum, pilosellum, albedo-flavescens. Caput oblongum, subquadrangulum, angulis rotundatis, dilute fuscum. Antennae subulatae, 4-articulatae (articulis sensim decrescentibus). Labrum semiorbiculatum, medio emarginatum. Mandibulae validae, falcatae. Maxillae oblongae, (mala singula), palpis validioribus, subulatis, 4-articulatis. Prothorax antrorsum angustatus, testaceus. Pedes subulati. Abdomen incisuris profundis, segmentis singulis rotundatis, convexis, segmento anali caudis duabus triarticulatis, subulatis, articulis earum cylindricis, secundo primo dimidio brevior, tertio setiformi. Anus tubulatus. Habitat vere in fimo equino; viciat rapina. (Naturgesch. d. Ins. I. 181. 3. t. VIII. f. 9—13.)

# 1) Thorax punctorum seriebus dorsalibus lateralibusque nullis.

\* 1. *X. canaliculatus*: *Depressus, nigerrimus, nitidus, ano rufo-testaceo, elytris laevigatis, capite oblongo, canaliculato, utrinque sulcato, angulis posterioribus rectis.* — Long. 6 lin.

Habitat Bogatae in Columbia, a Dom. Buquet nom. Xanth. canaliculati Dej. communicatus; in Mexico, collect. Dom. Chevrolat.

Statura fere Belonuchi haemorrhoidalis, totus nigerrimus, politus, nitidus, abdominis segmento quinto apice sextoque toto rufo-testaceis. Antennae capite paulo longiores, articulo tertio elongato, secundo duplo longiore, 4—11 rufo-piceis, cinerascens. Palpi picei. Caput thorace paulo longius et sesqui latius, basi truncatum, lateribus angulisque posterioribus rectis, his prominulis, antrorsum vix angustatum, planum, supra laevissimum, sulculis duabus obliquis inter antennas in media fronte fere conniventibus, dein usque ad basin canaliculatum, utrinque prope oculos sulco profundo, postice abbreviato, longitudinali, impressum, vertice punctorum serie transversali notatum, lateribus crebre punctatum, subtus medio et prope latera longitudinaliter canaliculatum. Thorax latitudine longior, antice coleopteris latitudine aequalis, apice utrinque oblique truncatus, basin versus sensim modiceque angustatus, lateribus haud sinuatus, angulis anterioribus obtusis, modice prominentibus, depressus, laevissimus. Scutellum laeve. Elytra thorace paulo longiora, laevissima, lateribus pilosella. Abdomen parce punctulatum parceque pilosellum, coleopteris angustius, apice attenuatum, subtus longius pilosum. Pedes nigri, spinulis tibiarum tarsisque piceis.

2. *X. flavipennis*: *Niger, nitidus, elytris rufo-testaceis, laevigatis, capite oblongo, canaliculato, utrinque subsulcato, angulis posterioribus obtusis.* — Long. 5½ lin.



Habitat in Mexico, collect. Dom. Chevrolat.

Præcedenti affinis, licet paulo angustior et minus depressus, nigerimus, pernitidus. Antennae capite vix longiores, articulo tertio minus elongato, secundo vix sesqui longiore, piceae, articulo primo nigro. Caput thorace paulo latius, oblongum, antrorsum subangustatum, basi utrinque oblique truncatum, angulis posterioribus obtusis, depressum, laeve, fronte canalicula tenui media, in summa basi abbreviata, antice cum sulculis duobus obliquis ordinariis ad antennarum basin, in fronte coniunctis, confluyente, sulcoque secundum capitis marginem lateralem, brevior, latiusculo, obsoletere exarata, lateribus supra oculos dense subtiliter punctatum, subtus medio et secundum latera canaliculatum. Thorax coleopteris paulo angustior, latitudine sesqui fere longior, basin versus leviter angustatus, apice utrinque oblique truncatus, lateribus haud sinuatus, subdepressus, laevissimus, versus angulos anteriores utrinque puncto singulo obsoletere impressus. Scutellum laeve. Elytra thoracis longitudine, laevigata, testaceo-rufa, nitida. Abdomen laeve, nigerrimum, ano concolore, subtus nitidum, griseo-pilosellum. Pedes nigro-picei, geniculis tarsisque rufescentibus.

3. *X. divisus*: *Niger, nitidus, elytris laevigatis, pectore, ano pedibusque rufo-testaceis, capite oblongo, canaliculato, utrinque sulcato, angulis posterioribus obtusis.* — Long. 6 lin.

Habitat Vera-Cruce in Mexico, collect. Dom. Chevrolat.

Statura præcedentium. Antennae capite vix longiores, articulo tertio modice elongato, secundo duplo prope longiore, piceae. Palpi rufo-picei. Caput thorace latius, oblongum, antrorsum vix angustatum, basi utrinque oblique truncatum, lateribus rectum, angulis posterioribus obtusis, subprominentibus, depressum, nigerrimum, nitidum, fronte canalicula media, postice subabbreviata, antice cum sulculis duobus ordinariis ad antennarum basin, in fronte connexis, confluyente, sulcoque secundum marginem lateralem, postice abbreviato, exarata, ceterum supra laevissimum, lateribus creberrime punctatum, subtus medio longitudinaliter impressum et subtiliter canaliculatum, secundum marginem lateralem sulcatum. Thorax apice coleopterorum latitudine, basin versus sensim angustatus, latitudine apicis sesqui fere longior, apice utrinque oblique truncatus, angulis anterioribus subrectis, modice prominentibus, lateribus rectis, nullo modo sinuatis, subdepressus, laevissimus, puncto utrinque singulo prope angulum anteriorem impressus, nigerrimus, nitidus. Scutellum testaceo-rufum, laeve. Elytra thoracis longitudine, in medio disci punctis tribus, longitudinaliter dispositis, impressa, ceterum laevigata, rufo-testacea, nitida. Abdomen parce punctulatum parceque pilosellum, subtus longius griseo-pilosum, nitidum, nigrum, segmento quinto apice sextoque toto testaceo-rufis. Pectus rufo-testaceum. Pedes cum coxis omnibus testaceo-rufi, tibiis anticis piceis.

- \* 4. *X. laticeps*: *Depressus, niger, nitidus, pedibus piceis, elytris subseriatim punctatis, capite lato, suborbiculato, subtiliter punctulato.* — Long. 5 lin.

Habitat in Java, Dom. Com. Hoffmannsegg.

Depressus, niger, nitidus. Antennae capite paulo longiores, articulo tertio modice elongato, secundo duplo prope longiore, piceae, articulis tri-

bus primis nigricantibus. Palpi picei. Caput magnum, suborbiculatum, depressum, fronte medio transversim subtiliter punctulata, apice sulculis duobus intermediis sat profundis, inter oculos foveolis 5 minutis, rotundatis, transversim positis, supra oculos punctis grossis utrinque tribus oblique positis, vertice pluribus irregulariter transversim sitis impressum, lateribus dense punctatum, subtus canaliculatum, parce subtiliterque sparsim punctatum. Thorax coleopteris angustior, latitudine sesqui longior, basin versus angustatus, lateribus subsinuatus, apice utrinque oblique truncatus, angulis anterioribus obtusiusculis, modice prominulis, depressus, laevis, in angulo utroque anteriore punctis tribus subtilioribus impressus. Scutellum leviter impressum, quadripunctatum. Elytra thoracis longitudine, dorso subseriatim punctata, lateribus inflexis subtiliter vage punctatis. Abdomen parce punctulatum, parcus nigro-pilosum, apice piceum, subtus totum piceo-rufescens. Pedes picei, tarsi rufescentibus.

- \* 5. *X. lynceus*: *Nigerrimus, nitidus, abdomine subcoerulescente, elytris punctatis, obscure coeruleis, capite orbiculato, limbo punctato.* — Long. 6 — 7 lin.

Habitat Bogotae in Columbia interiore, a Dom. Buquet sub hoc nomine communicatus.

Subdepressus, nigerrimus. Antennae capite sesqui longiores, articulo tertio elongato, secundo plus duplo longiore, nigrae, cinereo-pruinosa, articulis tribus primis glaberrimis. Palpi apice picei. Caput maris thorace duplo, feminae vix sesqui latius, orbiculatum, depressum, utrinque subtilius punctulatum, et, ut in vertice, punctis nonnullis grossis impressum, fronte media late laevis, nitidissima. Thorax latitudine sesqui longior, coleopteris dimidio prope angustior, basin versus vix angustatus, lateribus subsinuatus, apice utrinque oblique truncatus, angulis anterioribus obtusis, subrotundatis, supra parum convexus, in angulo utroque anteriore punctis tribus impressus, ceterum laevis, nitidissimus. Scutellum punctis nonnullis impressum. Elytra thoracis longitudine, dorso parcus subseriatim, lateribus crebrius subtiliusque vage punctata, obscure coerulea, nitida. Abdomen apice angustatum, parce subtiliterque punctulatum, nigro-subcoerulescens, nitidum, nigro-pilosum. Pedes graciliores, nigri, tarsi concoloribus.

- \* 6. *X. chalybeus*: *Coeruleus, nitidus, pedibus rufis, tibis tarsisque nigris, capite triangulari, laevissimo.* — Long. 8 — 9 lin.

Brullé Hist. nat. Ins. VI. 72. 1.

*Eulissus chalybaeus* Mannerh. *Brachelytr.* 35. 1. — Nordm. *Symb.* 129. 1.

*Staphyl. saphyrinus* Perty *Delect. An. art. p.* 31, *t.* 7, *f.* 5.

Habitat in Brasilia.

Magnus, robustus, laete coeruleus, pernitidus, abdomine subvirescente. Antennae vix capitis longitudine, articulo tertio elongato, secundo duplo longiore, nigro-piceae, articulo ultimo apice ferrugineo. Palpi picei. Caput basi thoracis apice vix latius, basi truncatum, angulis posterioribus rotundatis, lateribus subrectis, antrorsum angustatum, triangulare, supra modice convexum, laevissimum, fronte sulculis inter antennis obsoletis, inter oculos punctis duobus maioribus distantibus impressa. Thorax apice coleopteris

vix angustior, ibique longitudini latitudine subaequalis, basin versus angustatus, lateribus subrectis, apice truncatus, angulis anterioribus acutiusculis, leviter deflexis, leviter convexus, laevissimus, puncto singulo utrinque haud procul ab angulo anteriore impressus. Scutellum punctulatum. Elytra apice et lateribus vage subtiliter punctulata, dorso biserialim obsolete punctata, serie altera suturali, altera in medio dorsi. Abdomen breviusculum, apicem versus attenuatum, parce subtiliter punctulatum, longius nigro-pilosum. Pedes tibiis tarsisque nigris, femoribus omnibus coxisque anticis totis, intermediis infra rufis. Alae fuscae.

- \* 7. *X. Mannerheimii*: *Laete viridi-aeneus, pedibus flavis, elytris, pectoris lateribus abdominisque margine laterali luteis, capite triangulari.*  
— Long. 7 lin.

*Eulissus Mannerheimii* Laporte *Etud. Ent. I* 118. 2. t. 3. f. 5.

Habitat Cayennae.

Statura omnino praecedentis, at dimidio minor. Antennae vix capitis longitudine, articulo tertio leviter elongato, secundo sesqui longiore, 4—10 sensim paulo minoribus, piceo-testaceae, apice dilutiores, articulo ultimo flavo. Palpi et labrum rufo-picea. Caput thoracis longitudine eoque haud latius, basi truncatum, angulis posterioribus obtusis, antrorsum sensim angustatum, supra sat convexus, laete viridi-aeneum, nitidum, laevissimum, sulculis inter antennis distinctis, puncto maiusculo utrinque ad oculi marginem interiorem, alioque ad marginem eius superiorem impressum, margine verticali punctato. Thorax coleopteris parum angustior, longitudine paulo longior, basin versus leviter angustatus, lateribus rectis, nullo modo sinuatis, apice utrinque oblique truncatis, angulis anterioribus subrectis, leviter deflexis, supra leviter convexus, laevissimus, puncto singulo prope angulum utrumque anteriorem notatus, laete lucideque viridi-aeneus, margine angulari antico flavo. Scutellum parce punctatum, flavum, aeneo-nitidulum. Elytra thoracis longitudine, lutea, dorso biserialim punctata, serie altera suturali, altera in medio dorsi, lateribus inflexis subtilius punctatis. Abdomen densius griseo-pilosum, subtiliter punctulatum, apice attenuatum, lucide laeteque viridi-aeneum, margine laterali luteo. Pectus viridi-aeneum, flavo-limbatum. Pedes dilute flavi, coxis anterioribus disco viridi-aeneo.

- \* 8. *X. limbatus*: *Niger, nitidus, elytris margine laterali luteo, capite subtriangulari, confertim punctulato.* — Long. 7 lin.

*Staphyl. limbatus* Klug *Col. Madag.* 50. 42. t. 1. f. 13.

Habitat in Madagascar, Dom. Goudot.

Statura robustior *X. chalybei*, at minor, niger, nitidus, elytrorum margine laterali inflexo solo luteo. Antennae capite vix longiores, articulo tertio secundo sesqui longiore, nigrae, articulis 4—10 cinerascensibus, ultimo ferrugineo. Palpi rufo-picei. Caput thorace parum latius, eique longitudine aequale, basi truncatum, angulis posterioribus rotundatis, denticulo minutissimo prominulo notatis, antrorsum angustatum, lateribus vix rotundatum, supra infraque confertim punctulatum. Thorax apice coleopteris haud angustior, latitudine antica parum longior, basin versus angustatus, lateribus medio subsinuatis, apice utrinque oblique truncatus, angulis anterioribus acutiusculis, leviter deflexis,



leviter convexus, laevis, puncto utrinque singulo haud procul ab angulo anteriore impressus. Scutellum punctulatum. Elytra thoracis latitudine, dorso subtiliter rugulosa, subtilissime punctulata, biseriatim fortius punctata, serie altera suturali, altera dorsali, lateribus vage subtiliter punctatis, luteis. Abdomen sparsim punctatum, parce pilosum. Pedes nigri, tarsi piceis.

\* 9. *X. cyanopterus*: *Nigerrimus, nitidus, elytris subtiliter triseriatim punctatis, chalybeis, capite oblongo, laevi*. — Long.  $5\frac{1}{2}$  lin.

Habitat in Terra Van-Diemenii, Dom. Schayer.

Habitu fere praecedentis at paulo minor, totus nigerrimus, nitidissimus, elytris cyaneis. Antennae capite parum longiores, articulo tertio secundo duplo longiore, ultimo apice ferrugineo. Palpi apice picei. Caput thoracis latitudine, latitudine paulo longius, basi et lateribus leviter, angulis posterioribus fortiter rotundatis, antrorsum paululum angustatum, supra convexum, subtilissime punctulatum, punctis utrinque tribus in capitis margine postico, duobus ad oculum utrumque impressum, sulculis duobus anticis intermediis parallelis. Thorax coleopteris parum angustior, latitudine paulo longior, basin versus subangustatus, apice truncatus, angulis anterioribus lateribusque subrectis, leviter convexus, puncto utrinque singulo intra angulum anteriorem sat profundo impressus. Scutellum obsolete 4-punctatum. Elytra thorace paulo longiora, triseriatim punctata, serie prima suturali, secunda dorsali, tertia marginali, omnibus sat regularibus, margine apicali et lateribus inflexis vage punctatis. Abdomen parce punctatum, parce nigro-pilosum. Pedes nigri, tarsi piceis. Alae fusco-hyalinae.

\* 10. *X. chloropterus*: *Niger, capite triangulari laevi thoraceque opacis, elytris viridibus, abdomine nigro-subaeneo, nitidulis*. — Long. 9 lin.

Habitat in Terra Van-Diemenii, Dom. Schayer.

*X. chalybeo* longitudine aequalis, at gracilior, corporis habitu potius *X. glabrati*. Antennae capite vix longiores, articulo tertio elongato, secundo duplo longiore, 4—10 crassitie aequalibus, at sensim paulo brevioribus, nigrae, ultimo apice subferrugineo. Palpi nigri, apice rufo-picei. Caput thoracis longitudine et basi eius apici latitudine aequale, antrorsum angustatum, oblongo-subtriangulare, basi et lateribus perparum, angulis posterioribus fortiter rotundatum, supra leviter convexum, puncto maiusculo singulo ad interiorem, alteroque itidem singulo ad superiorem oculi marginem impresso, ceterum praeter sulculos anticos ordinarios laeve, nigrum, subopacum. Thorax apice coleopterorum latitudine, basin versus modice angustatus, latitudine antica paulo longior, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus subrotundatis, modice prominentibus, leviter convexus, longe intra apicem puncto singulo impressus, ceterum laevis, niger, subopacus. Scutellum medio punctatum, nigrum, parum nitidum. Elytra thorace vix longiora, triseriatim punctata, serie prima suturali, secunda dorsali, tertia marginali, margine inflexo summoque apice vage punctatis, viridia, nitidula. Abdomen elongatum, parallelum, parce subtiliter punctatum, parce tenuiterque nigro-pilosum, nigro-subaeneum, nitidulum. Corpus subtus nigro-subaeneum, nitidum. Pedes nigri. Alae fuscae.

11. *X. chalcopterus*: *Niger, nitidus, elytris triseriatim punctatis, cyaneo-aeneis, pedibus rufo-fulvis, capite oblongo, laevi.* — Long. 5 lin.

Habitat in Nova Hollandia, ad Cygnorum flumen, Mus. Dom. Hope.

Habitu omnino *X. fulgidi*, at duplo maior, niger, pernitidus. Antennae capite paulo longiores, articulo tertio secundo plus sesqui longiore, nigrae, articulo secundo tertioque basi rufo-piceis, ultimo apice ferrugineo. Palpi picei. Caput thorace paulo latius, latitudine longius, basi cum angulis posterioribus rotundatus, lateribus subrectis, antrorsum haud angustatum, subdepressum, omnium subtilissime obsoletissimeque punctulatum, puncto maiusculo singulo ad interiorem, duobus approximatis ad superiorem utriusque oculi marginem, pluribusque irregulariter transversim positis verticis impressum, sulculis duobus anticis intermediis prope parallelis. Thorax latitudine paulo longior, basin versus angustatus, lateribus pone medium subsinuatis, basi coleopteris angustior, parum convexus, puncto singulo ad angulum utrumque anteriorem impressus. Scutellum parumpunctatum. Elytra thorace paululum longiora, subtilius triseriatim punctata, serie prima suturali, secunda in medio dorsi, tertia marginali, omnibus sat regularibus, cyaneo-aenea, nitida. Abdomen parce vageque punctulatum, tenuiter pilosum. Pedes omnes cum coxis anterioribus rufo-fulvi. Alae fuscae.

- \* 12. *X. speciosus*: *Cyaneus, nitidus, elytris anoque rufis, capite subquadrato, punctulato, angulis posterioribus subdentatis.* — Long. 9 lin.

Habitat in Brasilia, Dom. Virmond.

Magnus. Corpus subtus cum pedibus nigrum, supra cyaneum, nitidum, elytris rufis, abdominis segmento quinto rufo-testaceo, basi cyaneo, sexto toto testaceo. Antennae capite vix longiores, articulo tertio secundo sesqui longiore, nigrae, articulo ultimo apice testaceo. Palpi picei. Mandibulae elongatae. Caput magnum, subquadratum, thorace plus sesqui latius, basi utrinque oblique truncatum, lateribus rectum, angulis posterioribus obtusis, inferne denticulo armatis, antrorsum subangustatum, depressum, totum crebrius punctatum punctisque maioribus utrinque nonnullis impressum. Thorax coleopteris angustior, latitudine plus sesqui longior, basin versus vix angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, leviter convexus, omnium subtilissime punctulatus, dorso medio punctis quatuor, quadratim positis, obsoletis impressus. Elytra thorace paulo longiora, subtilissime punctulata, dorso punctorum maiorum distantium serie instructa, rufo-testacea, nitida, coeruleo-subnitentia. Abdomen parallelum, sublaeve, punctis subtilibus raris impressum, parce nigro-pilosum. Pedes leviter elongati, graciliores, nigri, tarsis piceis.

- \* 13. *X. insignis*: *Nigro-subaeneus, nitidus, elytris anoque rufo-testaceis, capite subtriangulari, angulis posterioribus rotundatis.* — Long. 8 lin.

Habitat in Brasilia, Dom. Prof. Germar.

Affinis praecedenti, obscure nigro-aeneus, pernitidus, abdominis segmento quinto apice sextoque toto elytris rufo-testaceis. Antennae capitis longitudine, articulo tertio secundo paulo longiore, nigro-piceae, articulis 4—11 dense griseo-tomentosis. Palpi picei. Caput magnum, basi thorace sesqui latius et utrinque oblique truncatum, angulis posterioribus

rotundatis, lateribus subrectis, antrorsum fortius angustatum, depressum, vage subtiliter punctatum, fronte utrinque punctis grossis subvariolosa. Thorax latitudine sesqui longior, basin versus angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus subrectis, modice prominentibus, leviter convexus, laevissimus, punctis tribus in angulo utroque anteriore impressus. Scutellum punctis nonnullis fortibus notatum. Elytra thorace paulo longiora, dorso parce subseriatim punctata, rufo-testacea, metallico-nitida. Abdomen apice attenuatum, parce nigro-pilosum, laevigatum, punctis raris subtilibus prope segmentorum margines sparsum, apice rufo-testaceum. Corpus subtus nigro-subaeneum. Pedes nigri, tarsis piceis.

\* 14. *X. festivus*: *Cyaneus, nitidus, elytris rufo-testaceis, capite oblongo, punctulato*. — Long. 8 lin.

Habitat in Brasilia, Dom. Prof. Germar.

Corpus cyaneum, splendidum, elytris solis rufo-testaceis. Antennae capite paulo longiores, articulo tertio secundo paulo longiore, nigrae, articulo secundo piceo, articulis 4—11 dense griseo-tomentosis. Palpi picei. Caput thoracis longitudine eoque parum latius, basi rotundatum, lateribus subrectis, angulis posterioribus obtusis, denticulo minutissimo obtusiusculo notatis, antrorsum subangustatum, supra subdepressum, sparsim subtilius punctatum, fronte utrinque punctis nonnullis maioribus notata, sulculis intermediis in fronte antica conniventibus, fortiter impressis. Thorax latitudine duplo prope longior, basin versus vix angustatus, lateribus rectis, apice truncatus, angulis anterioribus obtusiusculis, transversim convexus, laevissimus, intra angulum utrumque anteriorem punctis nonnullis notatus. Scutellum punctis paucis impressum. Elytra thorace sesqui latiora et paulo longiora, subtiliter punctata, punctorum maiorum seriebus tribus, prima suturali, secunda dorsali media, tertia marginali, punctis earum autem parum profundis et sat distantibus. Abdomen elongatum, parallelum, laevigatum, vel punctis subtilissimis raris impressum, glabrum, solo apice nigro-piloso. Pedes graciliores, tarsis piceis.

\* 15. *X. lucidus*: *Niger, nitidissimus, elytris anoque rufo-testaceis, thorace dorso bipunctato, capite oblongo, vage punctulato*. — Long. 5½ lin.

Habitat in Brasilia, Dom. Prof. Germar.

Corpus nigrum, nitidissimum, abdominis segmento penultimo apice, ultimo toto elytris testaceo-rufis. Antennae capite paulo longiores, articulo tertio secundo parum longiore, rufo-piceae, articulo primo nigro. Palpi rufi. Caput thoracis fere longitudine, eoque parum latius, antrorsum subangustatum, basi utrinque oblique truncatum, angulis posterioribus obtusis, subrotundatis, depressiusculum, vage parcius punctatum, fronte utrinque punctis nonnullis grossiusculis notata, sulcis ordinariis in antica fronte inter antennis sat longis profundeque impressis. Thorax coleopteris paulo angustior, latitudine paulo longior, basin versus subangustatus, lateribus pone medium subsinuatis, apice utrinque oblique truncatus, angulis anterioribus obtusis, leviter convexus, omnium subtilissime punctulatus, punctis duobus in medio dorsi et tribus intra angulum utrumque anteriorem impressus. Scutellum bipunctatum. Elytra thorace longiora, subtiliter punctulata, pun-



ctorum maiorum serie in medio dorsi, rufo-testacea, nitida, sub-pellucida. Abdomen parallelum, segmentis singulis utrinque parcius subtiliter punctatis, parce pilosum, apice rufo-testaceum. Pedes graciles, nigri, tarsi rufo-piceis.

16. *X. phoenicopterus*: *Niger, nitidus, tarsi piceis, elytris rubris, purpureo-nitentibus, triseriatim punctatis, capite oblongo, laevi.* — Long. 5 lin.

Habitat in Nova Hollandia, Mus. Dom. Hope.

Statura omnino *X. analis*, nigerrimus, pernitidus. Antennae capite parum longiores, articulo tertio elongato, secundo duplo longiore, nigrae. Palpi picei. Caput thoracis latitudine, latitudine paulo longius, antrosum subangustum, lateribus rectis, angulis posterioribus rotundatis, punctis utrinque singulo maiusculo ad interiorem, binis approximatis, oblique positis, ad superiorem oculi marginem, paucisque irregulariter transversim positis verticis impressum, ceterum praeter sulculos anticos ordinarios laevissimum. Thorax coleopteris paulo angustior, longitudine paulo longior, basin versus modice angustatus, lateribus medio vix sinuatis, apice utrinque oblique truncatus, angulis anterioribus obtusis, modice prominentibus, parum convexus, intra angulum anteriorem utrinque puncto magno profundo et in margine anteriore punctis utrinque nonnullis impressus. Scutellum punctulatum. Elytra thorace paululum longiora, punctorum seriebus tribus sat regularibus notata, prima suturali, secunda in medio dorsi, tertia marginali, rubra, nitore purpureo resplendentia. Abdomen parallelum, parcius subtiliter punctulatum, nigrum, pernitidum, immaculatum. Pedes breves, nigri, tarsi piceis. Alae fusco-hyalinae.

- \* 17. *X. analis*: *Niger, nitidus, elytris anoque rufis, thorace dorso quadripunctato, capite oblongo, utrinque parce punctato.* — Long.  $4\frac{1}{2}$ — $5\frac{1}{2}$  lin.

*Gyrophypnus analis* Nordm. Symb. 123. 25.

Habitat in Brasilia.

Corpus nigrum, pernitidum, abdominis segmento penultimo apice, ultimoque toto elytris rufis. Antennae capite parum longiores, articulo tertio secundo paulo longiore, 4—10 transversis, sensim paululum crassioribus, rufo-piceae, articulo primo nigro, ultimo ferrugineo. Palpi rufi. Caput thoracis longitudine eoque paulo latius, basi et angulis posterioribus rotundatum, lateribus subrectum, antrosum vix angustatum, supra leviter convexum, fronte utrinque punctis grossiusculis profundisque parvis sparsa, ceterum praeter sulculos anticos ordinarios sat profundos laevissimum. Thorax coleopteris angustior, latitudine sesqui fere longior, basin versus leviter angustatus, lateribus subrectis, apice utrinque oblique truncatus, angulis anterioribus subrectis, modice prominentibus, leviter convexus, dorso medio punctis subtilioribus quatuor, quadratim positis, pone medium puncto uno alterove, lateribus utrinque ante medium punctis profundioribus quatuor impressus. Scutellum parce punctatum. Elytra thoracis longitudine, dorso et lateribus parce subtiliterque subseriatim punctata, seriebus tribus punctorum maiorum, prima suturali, secunda dorsali, tertia marginali. Abdomen punctis raris subtilioribus impressum, parce nigro-pilosellum, apice laete

rufo, densius nigro-rufoque piloso. Pedes graciliores, tarsis piceis. Alae fusco-hyalinae.

- \* 18. *X. illucens*: *Niger, nitidus, elytris testaceis, scutello punctulato, capite oblongo, utrinque parum-punctato, subtilissime strigoso.* — Long. 5 lin.

Habitat in Columbia, Dom. Moritz.

Corpus nigrum, pernitidum, aeneo-relucens, elytris testaceis. Antennae capite paulo longiores, articulo tertio secundo vix longiore, 4—10 sensim paulo brevioribus, penultimis transversis, rufo-piceae, articulo basali nigro, apicali ferrugineo. Palpi rufi. Caput thoracis longitudine eoque vix latius, basi et lateribus parum, angulis posterioribus fortiter rotundatis, antrorsum parum angustatum, supra modice convexum, subtilissime minus dense longitudinaliter strigosum, fronte utrinque punctis nonnullis (circiter 5) grossiusculis impressa. Thorax coleopteris angustior, latitudine paulo longior, basin versus leviter angustatus, lateribus subrectis, apice truncatus, angulis anterioribus subrotundatis, supra modice convexus, laevis, lateribus subtilissime strigosis, utrinque ante medium punctis tribus impressus, angulis anterioribus summis rufis. Scutellum punctulatum, longe parceque fulvopilosum. Elytra thoracis longitudine, crebrius punctata, punctis dorso subseriatis. Abdomen parcius subtiliter punctatum, segmentis ventralibus tenuiter testaceo-marginatis, subtiliter tenuiterque griseo-pilosum. Pedes nigropicei, geniculis tarsisque rufo-piceis, trochanteribus omnibus testaceis, coxis anticis testaceis, disco anteriore et posteriore nigro. Alae fuscae.

- \* 19. *X. castigatus*: *Niger, nitidus, antennarum basi, pedibus, elytris, thorace subtilis, pectore anoque rufis, capite suborbiculato, utrinque fortiter punctulato.* — Long. 4½ lin.

Habitat in Columbia, Dom. Moritz.

Subdepressus. Antennae capite paulo longiores, articulo tertio secundo tertia parte longiore, 4—10 sensim paulo brevioribus et crassioribus, penultimis fortiter transversis, rufo-piceae, articulo primo rufo, ultimo ferrugineo. Palpi rufi. Mandibulae piceae. Caput thorace latius, suborbiculatum, depressum, fronte utrinque punctis grossiusculis oblongis vage fortiterque impressa, nigrum, nitidum. Thorax coleopteris angustior, latitudine apicis paulo longior, basin versus angustatus, lateribus rectis, apice truncatus, angulis anterioribus subrectis, modice prominentibus, subdepressus, supra utrinque punctis 5 sat profundis impressis, nigro-piceus, nitidus, marginibus rufescentibus, subtus rufus. Scutellum rufum, parce punctulatum. Elytra thoracis longitudine, dorso parce subseriatim, lateribus crebrius vage subtiliterque punctata, testaceo-rufa, nitida. Abdomen parallelum, parce subtiliterque punctulatum, supra parce tenuiterque pilosum, subtus longius griseo-pilosum, nigrum, segmento quinto apice, sexto toto rufis. Pectus cum pedibus rufum. Alae fusco-hyalinae.

20. *X. vagus*: *Niger, nitidus, pedibus posterioribus rufo-brunneis, elytris punctatis, nigro-subaeneis, capite oblongo, lateribus parce punctato,* — Long. 4¾ lin.

Habitat prope Caracas Columbiae, Dom. Moritz.

Habitu fere *X. linearis*, at triplo fere maior, niger, pernitidus. Antennae capite paulo longiores, articulo tertio secundo parum longiore, 4—10 sensim paululum crassioribus, nigrae. Palpi rufo-picei. Caput thoracis latitudine, oblongo-ovatum, fronte leviter convexa, utrinque parce punctata, sulculis anticis intermediis obsoletioribus, parallelis. Thorax coleopteris paulo angustior, latitudine plus sesqui longior, basin versus haud angustatus, lateribus pone medium subsinuatis, angulis anterioribus rotundatis, posterioribus obtusis, leviter convexus, utrinque lateribus antice punctis paucis maioribus, postice punctis nonnullis sparsis subtilioribus impressus. Scutellum punctis quatuor impressum. Elytra thorace paulo longiora, dorso et lateribus crebre punctata, nigro-subaenea, nitida. Abdomen parallelum, parce subtiliter punctatum, tenuiter pilosum, nigro-subaeneum, nitidum, apice summo piceo-testaceo, densius piloso. Pedes antici nigro-picei, posteriores rufo-brunnei. Alae fuscae.

- \* 21. *X. gaganus*: *Nigerrimus, nitidus, elytris angulo humerali testaceo, capite oblongo-subquadrato, crebre fortiterque punctato. — Long. prope 5 lin.*

Habitat in Madagascar, Dom. Goudot.

Habitu corporis *X. fulgidi*, at duplo prope maior, nigerrimus, nitidus. Antennae capite paulo longiores, crassiusculae, apicem versus sensim leniter incrassatae, articulo tertio secundo paulo longiore, 4—10 transversis, sensim crassioribus, piceae, articulo primo nigro. Caput thoracis latitudine eodemque paulo latius, basi truncatum, angulis posterioribus leviter rotundatis, lateribus subrectis, antrorsum vix angustatus, supra modice convexus, punctis magnis, profundis, oblongis, crebrius variolosum, sulcis anticis intermediis sat profundis, posterius leviter conniventibus, interstitio sulcis duobus approximatis obsoletioribus canaliculato, minus nitidum. Thorax coleopteris paulo angustior, latitudine apicis plus sesqui longior, basin versus sensim leniter angustatus, lateribus subrectis, apice truncatus, angulis anterioribus subrectis, modice prominentibus, sat convexus, nitidissimus, subtilissime parce punctatus, antice utrinque punctis maiusculis tribus, pone medium ad marginem lateralem singulo notatus. Scutellum impressum. Elytra thorace paulo longiora, dorso punctorum seriebus tribus, prima suturali, secunda discoidali, tertia marginali, notatis, interstitiis parce, lateribus inflexis crebrius subtiliter punctatis, angulo summo humerali testaceo. Abdomen parce subtilius punctatum, tenuiter griseo-pubescent. Tarsi picei.

22. *X. anachoreta*: *Niger, nitidus, antennis pedibusque nigro-piceis, elytris triseriatim punctatis, apice lutescentibus, capite subtriangulari, lateribus crebre punctato. — Long. 6 lin.*

Habitat in Bengala, Mus. Dom. Hope.

Statura omnino *X. glabrati*, nigerrimus, nitidus. Antennae capite paulo longiores, articulo tertio secundo sesqui longiore, nigro-piceae. Palpi picei. Caput thorace vix latius, latitudine baseos vix longior, subtriangulare, angulis posterioribus fortiter rotundatis, supra leviter convexum, laevissimum, prope oculos punctis nonnullis grossis rugiformibus profundis impressum, lateribus crebre subtilius punctatum, medio laevissimum, sulcis anticis inter-



mediis sat profundis, ad mediam frontem usque adscendentibus, parum conniventibus. Thorax apice coleopterorum latitudine, basin versus sensim leviter angustatus, latitudine apicis paulo longior, lateribus pone medium subsinuatis, ante medium leviter rotundatis, angulis anterioribus obtusis, parum convexus, laevis, puncto utrinque singulo intra angulum anteriorem impressus. Scutellum parce punctatum. Elytra thorace haud longiora, triseriatim punctata, serie prima suturali secundaque in medio dorsi distinctioribus, tertia marginali valde obsoleta, nigra, apice indeterminate lutescentia. Abdomen crebrius punctulatum, segmentis dorsalibus medio longitudinaliter laevibus, tenuiter griseo-pubescentis. Pedes nigro-picei, geniculis tarsisque rufescentibus, tibiis fortius spinulosus.

23. *X. dispilus*: Niger, nitidus, antennis pedibusque rufo-piceis, elytris margine inflexo maculaque discoidali testaceis, capite triangulari, fronte punctata, lateribus sulcata. — Long.  $3\frac{2}{3}$  lin.

Habitat in Bengala, Mus. Dom. Hope.

Subdepressus, niger, nitidus. Antennae capite parum longiores, articulo secundo tertioque longitudine aequalibus, apicem versus subattenuatae, rufo-piceae. Palpi piceo-rufi. Caput thorace latius, latitudine haud longius, triangulare, basi truncatum, angulis posterioribus rotundatis, subdepressum, spatio transverso frontali dense subtiliter punctato, lateribus prope oculos sulculo profundo, apice sulcis quatuor longitudinalibus impressum, intermedii eorum ordinariis ad mediam frontem usque adscendentibus, parum conniventibus, vertice punctis nonnullis maioribus profundis, irregulariter transversim positis, notato. Thorax apice coleopteris haud angustior, basin versus sensim angustatus, latitudine apicis parum longior, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusiusculis, subdepressus, laevis, puncto utrinque singulo prope angulum anteriorem impressus. Scutellum subtiliter punctulatum. Elytra thorace haud longiora, triseriatim punctata, serie prima suturali secundaque in medio dorsi sat distinctis, tertia marginali obsoletiore, nigro-picea, macula magna discoidali testacea. Abdomen apicem versus sensim attenuatum, subtiliter punctulatum, dorso linea lata media longitudinali laevi, parce griseo-pilosellum, segmentis singulis marginibus piceis. Pedes piceo-rufi vel piceo-testacei.

- \* 14. *X. volvulus*: Subcylindricus, niger, nitidus, antennis pedibusque piceis, ano rufo, elytris laevibus, capite oblongo, punctato. — Long. 4 lin.

Habitat in Brasilia, Dom. Virmond.

Statura tenuis fere *X. linearis*, at magis cylindricus, niger, pernitidus, abdominis segmento quinto apice, sexto toto rufo-testaceis. Antennae capite paulo longiores, articulo tertio secundo paulo brevior, 4—10 sensim paulo crassioribus, rufo-piceae, articulo primo nigro. Palpi rufi. Os piceum. Caput thorace aliquantum latius, at paulo brevius, basi truncatum, lateribus subrectis, angulis posterioribus rotundatis, antrorsum haud angustatum, supra leviter convexum, crebrius profunde punctatum, utrinque punctis nonnullis maioribus impressum. Thorax coleopteris angustior, latitudine duplo longior, lateribus posterius subsinuatus, apice truncatus, angulis anterioribus subrectis, subcylindricus, ante medium utrinque punctis duobus

oblique positus impressus, ceterum laevissimus. Scutellum laeve. Elytra thorace paulo longiora, obsolete subtilissime punctata. Abdomen parce punctulatum parceque griseo-pilosellum, apice rufo-testaceum. Pedes picei.

Species mihi invisae:

25. *X. rutilus*: *Aureo-viridis, nitidus, pectore, pedibus elytrisque brunneo-flavis, his aureo-micantibus.* — Long. 5 lin.

*Staph. rutilus* Perty *Delect. an. art. p. 30. t. 7. f. 2.*

Habitat in prov. Minarum Brasiliae.

Statura omnino *X. chalybei* videtur, at minor. Caput viridi-aureum, nitidissimum, laevigatum, striola et punctum utrinque inter oculos, punctis duobus utrinque pone oculos, et punctorum serie irregulari ad marginem posticum impressis. Mandibulae nigrae. Palpi brunnei. Thorax viridi-aureus, nitidissimus, punctis duobus utrinque ad angulos anteriores, et singulo ad angulos posteriores valde rotundatos impressus. Scutellum magnum, dilute brunneum, vage punctulatum. Elytra flavo-brunnea, aureo-micantia, punctulata, ad apicem ciliata. Abdomen viride, nitidum, apice flavicante. Antennae capite thoraceque breviores, ad basin brunneae, apice grisescenti-pilosae, articulo ultimo flavicante. Pedes brunneo-flavi.

26. *X. ater*: *Niger, nitidus, elytris biseriatim punctatis.* — Long. 10 lin.  
*Eulissus ater* Laporte *Etud. Ent. I. 117. 1. t. 3. f. 4.*

Habitat in Senegalia.

Statura omnino *X. lateralis*, at maior, totus ater, immaculatus, nitidus. Caput thorace latius, magnum, antice sulculis duobus impressum. Thorax coleopteris parum angustior, latitudine antica vix longior, basin versus angustatus, laevissimus, puncto utrinque singulo prope angulum anteriorem impressus. Elytra thoracis longitudine, seriebus duabus punctorum instructa. Abdomen punctulatum.

27. *X. vagepunctus*: *Nitidus, nigricans, laevis, coleopterorum disco vage punctato.* — Long. 5 lin.

*Staph. vagepunctus* Grav. *Mon. 88. 83.* — Latr. *Hist. nat. Crust. et Ins. IX. 301. 14.*

Habitat?

Caput orbiculatum, thoracis magnitudine, nitidum, laeve, punctis tantum paucis in baseos angulis impressum, antennae capitis longitudine. Thorax convexus, quadratus, angulis obtusis, coleopteris brevior et angustior, nitidus, laevis, puncto uno laterali distincto impressus. Coleoptera quadrata, nitida, depilia, excepto margine apicali setis rigidis fimbriato, laevia, margine laterali confertim punctato, disco vage punctato, nigricantia, margine summo laterali rufescente. Abdomen nigrum, ano fusciscente. Pedes fusciscentes.

Variat coleopteris fusciscentibus, margine laterali pallidiore, pedibus segmentorumque marginibus fuscis. (Grav.)

Obs. Elytra secundum descriptionem Latreillianam dorso seriatim punctata. Forte recte ad hoc genus allatus.

2) Thorax punctorum seriebus dorsalibus nullis, lateralibus distinctis.

\* 28. *X. fulgidus*: Niger, nitidus, elytris rufis, capite profunde punctato, thorace lateribus unisulcato. — Long. 4 lin.

Er. Col. March. I. 423. 1.

*Staph. fulgidus* Fab. Mant. Ins. I. 220. 14.

*Paederus fulgidus* Fab. Syst. El. II. 609. 6. Ent. Syst. I. II. 537. 6.

*Staph. pyropterus* Grav. Mon. 102. 103. — Latr. Hist. nat. Ins. Crust. et Ins.

IX. 334. 95. — Gyll. Ins. Suec. II. 356 71. — Sahlb. Ins. Fenn. I. 334. 70.

*Xanthol. pyropterus* Boisd. et Lacord. Faun. Ent. Paris. I. 413. 2.

*Gyrohypn. pyropterus* Mannerh. Brachelytr. 33. 8. — Nordm. Symb. 119. 15.

Habitat in Europa, minus frequens.

Nigerrimus, pernitidus, elytris rufis. Antennae capite paulo longiores, articulo tertio elongato, secundo duplo prope longiore, 4—10 sensim paulo brevioribus et crassioribus, piceae, articulo primo nigro. Palpi rufo-picei. Caput thoracis longitudine eoque paulo latius, basi truncatus, angulis posterioribus fortiter rotundatis, lateribus rectis, antrorsum haud angustatum, punctis magnis, profundis, oblongis, lateribus in series continuas sulciformes digestis, in media fronte rarioribus, impressum, fronte utrinque sulco longitudinali abbreviato notata. Thorax coleopteris paulo angustior, latitudine plus sesqui longior, basin versus angustatus, lateribus pone medium subsinuatus, angulis anterioribus obtusiusculis, modice prominentibus, parum convexus, dorso laevis, lateribus sulco profundo, leviter flexuoso, puncta 8—9 continente, et praeterea punctis binis approximatis utrinque ad marginem anteriorem impressus. Scutellum punctis nonnullis notatum. Elytra thoracis longitudine, biseriatim punctata, serie altera suturali, altera discoïdali, irregularibus, spatio inter series punctis raris sparsis impressum, spatio ante marginem lateralem laevi, margine laterali inflexo subtilius punctato. Abdomen lineare, subtiliter parce punctatum, parce nigro-pilosum. Pedes picei, tarsis rufescentibus.

3) Thorax punctorum seriebus dorsalibus rectis lateralibusque lituis impressus.

\* 29. *X. glabratus*: Niger, nitidus, antennis pedibusque piceis, elytris rufis, capite lateribus disperse punctato thoraceque parce subtilissime punctulatis, hoc seriebus dorsalibus 5—7 punctatis. — Long.  $5\frac{1}{2}$ — $6\frac{1}{2}$  lin.

Er. Col. March. I. 424. 2.

*Staph. glabratus* Grav. Micr. 178. 38 Mon. 101. 100.

*Gyrohypn. glabratus* Nordm. Symb. 113. 1.

*Staph. fulgidus* Grav. Micr. 48. 71. Mon. 106. 103. — Oliv. Ent. III. 42. 18. 19.

t. 4. f. 34. a. d. — Latr. Hist. nat. Crust. et Ins. IX. 832. 90. Gen. Crust. et Ins. I. 288. 4.

*Staph. nitidus*. Panz. Faun. Germ. 27. 8.

*Staph. cruentatus* Marsh. Ent. Britt. 516. 56.

*Xanthol. fulgidus* Boisd. et Lacord. Faun. Ent. Paris. I. 412. 1. — Brullé Hist. nat. Ins. VI. 73. 2.

*Gyrohypn. relucens* Nordm. Symb. 115. 5.

Var. a: *Pedibus rufis*.

*Gyrohypn. merdarius* Nordm. Symb. 116. 6.



*Var. b: Elytris pallidius testaceis, antennis pedibusque rufis.*

Oliv. Ent. III. 42. t. 4. f. 34. b.

Staph. relucens Grav. Mon. 101. 102.

Xanthol. cadaverinus Boisdu. et Lacord. Faun. Ent. Paris. I. 414. 4.

Gyrohypn. ochropterus Nordm. Symb. 118. 13.

Habitat in Europa temperata et meridionali, hic frequentior.

Nigerrimus, nitidus, elytris rufis. Antennae capite paulo longiores, articulo secundo tertioque aequalibus, quarto globoso, 5 — 10 subaequalibus, rufo-piceae. Palpi rufi. Caput thoracis longitudine eodemque basi parum latius, basi truncatum, angulis posterioribus rotundatis, lateribus subrectis, antrorsum leniter angustatum, supra leniter convexum, subtilissime parce punctulatum, utrinque punctis magnis sparsis impressum. Thorax coleopteris paulo angustior, latitudine sesqui prope longior, basin versus sensim angustatus, lateribus pone medium subsinuatus, angulis anterioribus obtusis, leviter convexus, subtilissime parce punctatus, seriebus dorsalibus 5—7-punctatis, serie laterali litua 6—7-punctata. Scutellum laeve. Elytra thoracis longitudine, parcius irregulariter punctata. Abdomen parce subtiliter punctatum, subtiliter tenuiterque griseo-pubescentis, parce nigro-pilosum. Pedes picei, tarsis rufescentibus.

Mas abdominis segmento ventrali sexto truncato, septimo apice inciso, femina segmento sexto apice rotundato, septimo integro, distinguuntur.

Obs. Capite thoraceque subtilissime interpunctatis bene distinctus. Var. a tantummodo pedibus potius rufis quam piceis, var. b pedibus rufo-testaceis et elytris pallide testaceis, statura saepius paulo minore magis a genuinis discedit, at haec forte minus matura minime speciem peculiarem constituere videtur.

30. *X. erythropterus: Subdepressus, niger, nitidus, elytris, ano pedibusque rufis, capite subquadrato, utrinque parce punctato, thorace seriebus dorsalibus parumpunctatis. — Long. 6 lin.*

Habitat in Nova Hollandia, Mus. Hope.

X. glabratum longitudine haud superans, at paulo latior et magis depressus, niger, nitidus. Antennae capite parum longiores, articulo tertio elongato, secundo duplo prope longiore, 4 — 10 cylindricis, leviter transversis, aequalibus, piceae, articulo primo rufo-piceo, ultimo apice ferrugineo. Palpi et labrum piceo-rufa. Caput thorace latius, subquadratum, basi truncatum, angulis posterioribus rotundatis, lateribus rectis, antrorsum vix angustatus, supra subdepressum, utrinque punctis nonnullis sparsis impressum, sulculo laterali ante oculos brevissimo, obsoleto, interne foveola latiore parum profunda terminato. Thorax coleopteris paulo angustior, basin versus vix angustatus, latitudine paulo longior, angulis anterioribus subrectis, modice prominentibus, subdepressus, seriebus dorsalibus punctis 4 — 5, 3 vel 4 ante medium, et singulo a reliquis distante pone medium compositis, serie laterali 5- vel 6-punctata, punctis omnibus subtilioribus. Scutellum punctulatum. Elytra thorace paulo longiora, vage subtiliter punctata, margine laterali laevi, rufa, nitida. Abdomen parce subtiliter punctatum parceque pilosum, segmento quinto apice sextoque toto rufo-testaceis. Pedes piceo-rufi.

- \* 31. *X. lubricus*: *Niger, nitidus, elytris abdominisque segmentis duobus ultimis rufis, capite oblongo, utrinque parce punctato, thorace seriebus dorsalibus parum-punctatis.* — Long. 4 lin.

Habitat prope San João del Rey Brasiliae, Dom. Sellow.

Statura omnino *X. fulgidi*, niger, pernitidus, elytris abdominisque segmentis duobus apicalibus rufis. Antennae capite paulo longiores, crassiusculae, articulo tertio modice elongato, secundo duplo prope longiore, 4—10 sensim paulo crassioribus, 5—10 transversis, rufo-piceae, articulo primo tertioque nigris, secundo rufo, ultimo apice ferrugineo. Palpi rufi. Caput thoracis longitudine eoque paulo latius, basi truncatum, angulis posterioribus rotundatis, lateribus subrectis, antrorsum leviter angustatum, supra parum convexum, utrinque parce sparsim punctatum. Thorax coleopteris angustior, latitudine sesqui longior, basin versus angustatus, lateribus pone medium subsinuatus, angulis anterioribus obtusiusculis, subrotundatis, leviter convexus, seriebus dorsalibus parum-punctatis, serie laterali flexuosa circiter 11-punctata. Scutellum punctis nonnullis obsoletis impressum. Elytra thoracis longitudine, dorso parcius subseriatim, lateribus subtilius crebriusque vage punctata, rufa, nitida. Abdomen parce punctulatum, parce griseo pilosum, apice late rufum. Pedes nigri, tarsi piceis.

Obs. Speciem unicum, thoracis medio acu laesum, quare punctorum in serie dorsali numerus dubius est.

- \* 32. *X. calidus*: *Niger, nitidus, elytris rufis, abdomine summo apice rufo-piceo, capite utrinque fortiter punctato, thorace seriebus dorsalibus 6—7-punctatis.* — Long. 5 lin.

Habitat in Columbia, Dom. Moritz.

Statura *X. glabrati* et minoribus eius individuis aequalis, niger, pernitidus, elytris rufis. Antennae capite paulo longiores, articulo tertio modice elongato, secundo sesqui longiore, 4—10 sensim paulo brevioribus et crassioribus, 5—10 sensim paulo magis transversis, rufo-piceae, articulis tribus primis nigris, ultimo apice ferrugineo. Palpi rufi. Caput thoracis prope longitudine eoque parum latius, basi et lateribus subrectis, angulis autem posterioribus late fortiterque rotundatis, antrorsum leviter angustatum, supra leviter convexum, utrinque parcius fortiusque vage punctatum. Thorax coleopteris angustior, latitudine tertia parte longior, basin versus subangustatus, lateribus pone medium subsinuatus, angulis anterioribus obtusis, parum convexus, seriebus dorsalibus 6- vel 7-punctatis, postice duplicatis, serie laterali flexuosa circiter 11-punctata. Scutellum punctis paucis obsoletis impressum. Elytra thoracis longitudine, dorso subseriatim, lateribus crebre punctata, rufa, nitida. Abdomen parce subtiliterque punctulatum, teniter griseo-pilosum, segmento penultimo margine summo apicali, sexto apice rufo-piceis. Pedes picei, tarsi dilutioribus. Alae fusco-hyalinae.

- \* 33. *X. castanopterus*: *Niger, nitidus, antennis pedibusque rufo-piceis, elytris rufis, capite oblongo, crebre fortiterque punctato, thorace seriebus dorsalibus 10-punctatis.* — Long. 4 lin.

Habitat in Brasilia, Dom. Prof. Germar.

Statura omnino *X. fulgidi*, niger, pernitidus, elytris saturate rufis. An-

tennae capite paulo longiores, crassiusculae, articulo tertio secundo parum longiore, 4—10 transversis, subaequalibus, rufo-piceae, articulo primo rufo-piceo, secundo rufo, ultimo apice ferrugineo. Palpi rufi. Mandibulae et labrum rufo-picea. Caput thorace parum brevius eique latitudine aequale, basi cum angulis posterioribus rotundatum, lateribus subrectis, antrorsum leviter angustatum, supra leviter convexum, fortiter profundeque sat crebre punctatum, spatio medio longitudinali laeviore, sulcis anticis intermediis sat profundis, ad mediam frontem usque adscendentibus, vix conviventibus; lateralibus valde obsoletis. Thorax coleopteris paulo angustior, latitudine plus sesqui longior, basin versus vix angustatus, lateribus pone medium subsinuatis, angulis anterioribus rotundatis, supra leviter convexus, seriebus dorsalibus minus regularibus, punctis circiter 10 compositis, serie laterali flexuosa fere 12-punctata, praetera punctis binis pone medium prope seriem dorsalem et pluribus intra utrumque angulum anteriorem impressus, punctis omnibus profundioribus. Scutellum impressum, impunctatum. Elytra thoracis longitudine, dorso parcius subtiliusque subseriatim, lateribus crebrius subtiliter punctata, saturate rufa, nitida, parce griseo-pubescentia. Abdomen parce subtilius punctatum, supra parce, subtus densius griseo-pubescentia, segmento quinto sextoque margine postico rufo-piceo. Pedes nigro-picei, tibiis tarsisque rufis.

\* 34. *X. pectoralis*: *Niger, nitidus, elytris, ano, pectore pedibusque rufis, capite utrinque parce punctato, thorace seriebus dorsalibus 6-punctatis.* — Long.  $4\frac{1}{2}$  — 5 lin.

Habitat in Columbia, Dom. Moritz.

Statura praecedentium, niger, nitidus, elytris, abdominis segmento quinto margine apicali sextoque toto, pedibus pectoreque rufis, hoc medio nigricante. Antennae capite parum longiores, articulo tertio secundo sesqui longiore, 4—10 sensim paulo brevioribus crassioribusque et magis transversis, rufo-piceae, articulo ultimo apice ferrugineo. Palpi rufi. Caput thoracis longitudine eoque parum latius, basi et apice parum, angulis posterioribus late fortiterque rotundatis, antrorsum modice angustatum, supra leviter convexum, utrinque parce punctatum. Thorax coleopteris paululum angustior, latitudine sesqui longior, basin versus subangustatus, lateribus pone medium subsinuatis, angulis anterioribus rotundatis, leviter convexus, seriebus dorsalibus 6-punctatis, serie laterali irregulari, flexuosa, circiter 12-punctata. Scutellum punctis nonnullis obsoletis impressum. Elytra thoracis longitudine, dorso parcius subseriatim, lateribus crebrius subtiliusque vage punctata, laete rufa, nitida, parce tenuiterque griseo-pubescentia. Abdomen parce subtiliterque punctatum, tenuiter griseo-pilosum, apice rufum. Pectus rufum, medio nigrum. Pedes cum coxis omnibus laete rufi.

Obs. Hanc speciem nomine *X. spreti* in Catalogo collectionis suae signatum esse ill. Com. Dejean litteris certiores nos fecit.

\* 35. *X. rufipennis*: *Niger, nitidus, elytris pedibusque rufis, capite subovato, utrinque parce punctato, thorace seriebus dorsalibus 10—11-punctatis.* — Long.  $3\frac{1}{2}$  lin.

Habitat in Sicilia, Dom. Dr. Schultz, in Syria, Dom. Dr. Aubé.



*X. glabrato triplo minor, niger, nitidus, elytris pedibusque rufis.* Antennae capite parum longiores, articulo secundo tertioque obconicis, aequalibus, 4 — 10 subcylindricis, crassitie vix brevioribus, itidem inter se subaequalibus, rufo-piceae. Palpi rufi, articulo ultimo pallido. Labrum rufum. Caput thorace paulo brevius et parum latius, basi et lateribus leviter, angulis posterioribus fortiter rotundatis, antrorsum modice angustatum; supra leviter convexum, utrinque minus subtilius at sat profunde disperse punctatum, fronte media laevi, sulculis anticis intermediis parallelis, sulculis obliquis ante oculos obsoletioribus, interne puncto maiusculo terminatis. Thorax coleopteris paulo angustior, latitudine sesqui longior, basin versus leviter angustatus, lateribus pone medium vix sinuatis, angulis anterioribus rotundatis, leviter convexus, seriebus dorsalibus minus regularibus, 10 — 11-punctatis, serie litua laterali 11 — 12-punctata, spatio inter seriem dorsalem et lateralem punctis nonnullis dispersis notato. Scutellum obsoletius tripunctatum. Elytra thorace parum longiora, dorso minus subtiliter subseriatim punctata, secundum marginem lateralem laevia, margine laterali inflexo subtilius punctato, laete rufa, nitida, fulvo-pubescentia. Abdomen parce subtiliterque punctatum, subtus segmentis duobus ultimis piceis, tenuiter parceque griseo-pubescentis, subtiliter parceque nigro-pilosum. Pedes cum coxis anticis laete rufi.

\* 36. *X. elegans: Rufo-testaceus, nitidus, capite abdomineque nigris, hoc segmentorum marginibus testaceis, capite oblongo, utrinque subtiliter parce punctato, thorace seriebus dorsalibus subtiliter 8-punctatis.* —

*Staph. elegans Oliv. Ent. III. 42. 19. 20. t. 5. f. 50. — Latr. Hist. nat. Crust. et Ins. IX. 331. 87.*

*Xanthol. meridionalis Boisd. et Lacord Faun. Ent. Paris. I. 413. 3.*

*Gyrophypnus meridionalis Nordm. Symb. 117. 12.*

Habitat in Gallia meridionali, Hispania.

Statura omnino *X. linearis*, at sesqui maior. Antennae capite paulo longiores, articulo secundo tertioque subaequalibus. 4 — 10 sensim paulo brevioribus et crassioribus, penultimis leviter transversis, rufo-piceae, articulis tribus primis rufis, ultimo apice ferrugineo. Palpi rufi. Os piceum. Caput thoracis fere longitudine eoque parum latius, basi et lateribus parum, angulis posterioribus fortiter rotundatis, antrorsum leviter angustatum, supra leviter convexum, nigrum, pernitidum, utrinque subtiliter parce punctatum, spatio medio longitudinali laevi, sulculis intermediis inter antennis brevibus, parallelis, parum profundis, sulculis obliquis ante oculos longioribus, sat distinctis. Thorax coleopteris paulo angustior, latitudine sesqui longior, basin versus angustatus, lateribus subrectis, apice utrinque oblique truncatus, angulis anterioribus lateraliter sat prominentibus, subrotundatis, leviter convexus, rufo-testaceus, pernitidus, seriebus dorsalibus 8-punctatis, serie litua laterali circiter 9-punctata, punctis omnibus sat subtilibus. Scutellum piceum, obsolete bipunctatum. Elytra thorace prope breviora, testacea, nitida, dorso subseriatim punctata, secundum marginem lateralem laevia, margine laterali inflexo subtilius punctato. Abdomen subtiliter parceque punctatum, nitidulum, supra nigro-piceum, segmentis ultimis margine testaceis, subtus rufo-testaceum, segmentis singulis basi picescentibus. Pectus piceo-testaceum. Pedes breviusculi, graciliores, testacei.

- \* 37. *X. collaris*: *Niger, nitidus, ano pedibusque rufis, capite oblongo, utrinque parce punctato thoraceque sanguineo subtilissime parcius punctulatis, hoc seriebus dorsalibus 6 — 7-punctatis. — Long. 3½ — 4 lin.*

*Er. Col. March. I. 424. 3.*

Habitat prope Berolinum, Dom. Schüppel, in Germania meridionali, Dom. Godet, in Sardinia, Dom. Prof. Gené.

Statura omnino *X. lenti*, at paulo maior. Antennae capite parum longiores, crassiusculae, articulo tertio secundo paulo longiore, 4 — 10 transversis, sensim paulo brevioribus et crassioribus, rufo-piceae, articulo ultimo apice ferrugineo. Palpi rufi. Caput thoracis fere longitudine eoque paulo latius, basi truncatum, lateribus rectis, angulis posterioribus subrotundatis, antrorsum vix angustatum, subdepressum, nigrum, nitidum, subtilissime disperse punctulatum, utrinque punctis grossiusculis oblongis sparsum. Thorax coleopteris paulo angustior, latitudine plus sesqui longior, basin versus subangustatus, lateribus subrectis, apice utrinque oblique truncatus, angulis anterioribus rectis, modice prominentibus, parum convexus, rufo-sanguineus, nitidus, subtilissime parce punctulatus, seriebus dorsalibus 6 — 7-punctatis, serie litua laterali 6-punctata, punctis sat subtilibus. Scutellum nigrum. Elytra thoracis longitudine, dorso parcius subseriatim punctata, secundum marginem lateralem laevia, lateribus inflexis subtilius punctatis, nigro-picea, margine apicali pallido. Abdomen parce subtilissime punctatum, nigrum, nitidum, segmento quinto supra apice, subtus fere toto, sextoque toto rufis. Pectus nigrum. Pedes cum coxis anterioribus rufo-testacei.

- \* 38. *X. decorus*: *Rufus, nitidus, capite, elytris ano pectoreque nigris, capite oblongo, parce subtilius punctato, thorace seriebus dorsalibus 7 — 8-punctatis. — Long. 3½ lin.*

Habitat in Austria, Dom. Schüppel, in Hungaria montana, Dom. Waltl.

*X. glabro* statura similis, at gracilior. Antennae capite sesqui longiores, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo crassioribus et magis transversis, totae rufae. Palpi rufo-brunnei. Caput thorace paulo latius, oblongo-ovatum, supra leviter convexum, parce sparsim subtilius punctatum, sulcis anticis intermediis brevibus et parum profundis, sulculis obliquis ante oculos distinctis, nigrum, pernitidum, mandibulis rufo-piceis. Thorax coleopteris vix angustior, latitudine plus sesqui longior, lateribus parallelis rectis, angulis omnibus rotundatis, convexus, seriebus dorsalibus 7 — 8-, lateralibus lituis fere 12-punctatis, punctis omnibus subtilibus, laete rufus, pernitidus. Scutellum subimpressum, nigrum. Elytra thoracis longitudine, parcius subseriatim punctata, parceque griseo-pubescentia, nigerima, pernitida. Abdomen parce punctatum, parceque griseo-pilosellum, nitidum, segmentis anterioribus 4 rufis, reliquis nigris. Pectus nigrum. Pedes rufo-testacei.

Obs. Corporis colore *Paederum* simulat.

- \* 39. *X. emmesus*: *Testaceus, nitidus, capite oblongo, utrinque parce punctato, nigro, thorace rufo-testaceo, seriebus dorsalibus 4-punctatis. — Long. 2½ lin.*

*Staph. emmesus Grav. Micr. 176. 35. Mon. 98. 97.*

Habitat in America septentrionali.

Statura et summa affinitas *X. glabri*. Antennae capite paulo longiores, articulo secundo tertioque breviusculis, aequalibus, 4—10 transversis, rufo-piceae, articulis 3 primis rufis, ultimo apice ferrugineo. Os rufo-piceum. Caput thoracis fere longitudine eoque parum latius, basi truncatum, angulis posterioribus fortiter rotundatis, lateribus rectis, antrorsum haud angustatum, supra leviter convexum, utrinque parcius fortiter punctatum, medio longitudinaliter laeve, nigrum, nitidum. Thorax coleopteris perparum angustior, latitudine tertia parte longior, basin versus subangustatus, lateribus subrectis, angulis anterioribus obtusis subrotundatis, modice convexus, rufo-testaceus, nitidus, seriebus dorsalibus 4-punctatis, serie laterali litua 5—6-punctata. Scutellum testaceum, laeve. Elytra thoracis longitudine, luteo-testacea, nitida, triseriatim punctata, serie prima suturali irregulari, secunda in medio dorsi, tertia marginali, his sat regularibus, lateribus inflexis laevibus. Abdomen parce punctulatum, subtus testaceum, supra segmentis anterioribus 5 basi fusciscentibus. Pectus et pedes testacei.

\* 40. *X. glaber*: *Niger, nitidus, elytris pedibusque brunneo-testaceis, capite suborbiculato, utrinque parce subtiliter punctato, thorace seriebus dorsalibus 5—9-punctatis.* — Long.  $3\frac{1}{2}$  lin.

*Er. Col. March. I.* 425. 4.

*Gyrophyn. glaber Nordm. Symb.* 114. 4.

*Staph. glaber var. 2 Grav. Mon.* 100. 99.

*Xantholin. lentus var. b Zetterst. Faun. Lappon. I.* 81. 2.

*Staph. diaphanus Marsh. Ent. Britt.* 514. 46. (*forte*)

Habitat in Germania.

*Niger, nitidus.* Antennae capite parum longiores, articulo secundo tertioque brevioribus, subaequalibus, 4—10 brevibus, transversis, sensim paulo crassioribus, ultimo ovato, apice compresso, rufo-piceae, articulo primo medio nigricante. Palpi rufo-picei. Caput thorace vix latius et paulo brevius, subovatum, antrorsum leviter angustatum, supra leviter convexum, utrinque punctis parvis sparsis rotundis subtilioribus impressum, interstitiis laevissimis, sulculis anticis intermediis sat profundis, parum conniventibus, lateralibus ad oculum utrumque obsoletioribus, licet distinctis. Thorax coleopteris parum angustior, latitudine sesqui prope longior, basin versus subangustatus, lateribus pone medium vix sinuatis, apice truncatus, angulis anterioribus subrotundatis, leviter convexus, laevissimus, seriebus dorsalibus 8—9-punctatis, serie laterali litua 7—8-punctata. Scutellum laeve. Elytra thorace parum longiora, dorso seriebus tribus parce punctatis, prima suturali, secunda discoidali, tertia marginali, interstitiis serierum praeterea punctis raris subtilibus impressis; lateribus parcius subseriatim punctatis, brunneo-testacea, nitida. Abdomen parce subtiliter obsoleteque punctulatum, griseo-pilosellum, segmentis singulis margine tenui, segmento ultimo toto piceo-testaceo. Pedes brunneo-testacei, femoribus piceis, tarsis dilutioribus.

\* 41. *X. lentus*: *Subdepressus, niger, nitidus, elytris pedibusque brunneo-testaceis, capite subquadrato, utrinque crebrius fortiter punctato thoraceque subtilissime parce punctulatis, hoc seriebus dorsalibus 7—8-punctatis.* — Long.  $3\frac{1}{2}$  lin.



*Zetterst. Faun. Lappon. I.* 80. 2. *Ins. Lappon. 66.* 3. — *Er. Col. March. I.* 426. 5.  
*Staph. lentus Grav. Mon.* 101. 101. — *Gyll. Ins. Suec. II.* 354. 69. — *Sahlb. Ins. Fenn. I.* 333. 68.

*Gyrohypn. lentus Mannerh. Brachelytr.* 33. 6. — *Nordm. Symb.* 118. 10.

*Staph. glaber var. 1 Grav. Mon.* 100. 99.

*Staph. tricolor var. b Payk. Faun. Suec. III.* 378. 15.

Habitat in Germania, Suecia.

Statura et magnitudo omnino *X. punctulati*, subdepressus, niger, nitidus. Antennae capite paulo longiores, articulo secundo tertioque longitudine subaequalibus, 4—10 sensim paululum crassioribus, penultimis leviter transversis, rufae. Palpi rufi. Mandibulae piceae. Caput thoracis prope latitudine eodemque basi latius, basi leviter, angulis posterioribus fortius rotundatis, lateribus subrectis, antrorsum subangustatum, supra parum convexum, utrinque punctis sat magnis oblongis crebrius sparsum, undique subtilissime parce punctulatum, sulculis duobus anticis intermediis brevioribus, sat latis, at parum profundis, lateralibus ad oculi utriusque marginem inferiorem obsoletis. Thorax coleopteris paulo angustior, latitudine sesqui longior, basin versus leniter angustatus, lateribus pone medium subsinuatis, apice truncatus, angulis anterioribus subsinuatis, subdepressus, parce subtilissime punctulatus, seriebus dorsalibus 7—8-punctatis, serie laterali litua 4—5-punctata. Scutellum laeve. Elytra thorace parum longiora, crebrius vage punctata, secundum marginem lateralem laevia, brunneo-testacea, nitida. Abdomen subtilissime parce punctatum, tenuiter griseo-pubesceus, subtus segmento penultimo margine summo, ultimo apice picescentibus. Pedes cum coxis omnibus rufo-testacei.

\* 42. *X. consentaneus*: Subdepressus, niger, nitidus, elytris pedibusque brunneo-testaceis, capite oblongo, utrinque parce punctato thoraceque subtilissime disperse punctulatis, hoc seriebus dorsalibus 6—7-punctatis. — Long. 3 — 3½ lin.

*Gyrohypn. consentaneus Nordm. Symb.* 122. 24.

*Gyrohypn. flavipennis Nordm. Symb.* 121. 23.

◊ Habitat in America septentrionali.

Statura subdepressa et summa affinitas praecedentis, at magis elongatus, niger, nitidus. Antennae capite vix longiores, articulo secundo tertioque aequalibus, 4—10 transversis, subaequalibus, rufae. Palpi testacei. Caput oblongum, thoracis fere longitudine, eoque parum latius, basi truncatum, angulis posterioribus rotundatis, lateribus subrectis, antrorsum vix angustatum, supra parum convexum, utrinque parcius subseriatim punctatum, undique subtilissime parce punctulatum, sulculis duobus anticis intermediis sat profundis, leviter conniventibus, lateralibus obliquis ad oculorum marginem inferiorem obsoletioribus, licet distinctis. Thorax coleopteris paulo angustior, latitudine plus sesqui longior, basin versus subangustatus, lateribus pone medium subsinuatus, apice truncatus, angulis anterioribus subrectis, parum convexus, parce subtilissime punctatus, seriebus dorsalibus 6—7-punctatis, punctis posterioribus magis distantibus, seriebus lateralibus lituis 6-punctatis, punctis laeve. Elytra thoracis longitudine, dorso parcius subseriatim, lateribus subtilius crebriusque vage punctata, testacea, nitida. Abdomen lineare, parce

subtiliterque punctulatum, tenuiter griseo-pilosellum, subtus piceum, segmentis singulis apice testaceis, seu piceo-testaceum, segmentis singulis basi lateribusque nigricantibus. Pedes toti testacei.

Obs. Specimina sub nomine *Gyrophypni flavipennis* a Dom. Nordmann descripta nonnisi statura paulo minore et colore elytrorum pedumque dilutius differunt.

- \* 43. *X. brunneipennis*: *Niger, nitidus, elytris pedibusque brunneo-testaceis, capite utrinque disperse, medio biserialim punctato, thorace seriebus dorsalibus 7 — 9-punctatis.* — Long.  $3\frac{2}{3}$  lin.

Habitat in Columbia, Dom. Moritz.

Statura, magnitudo et summa affinitas *X. lenti*. *Niger*, pernitidus. Antennae capite paulo longiores, articulo secundo tertioque aequalibus, 4 — 10 transversis, sensim paulo crassioribus, rufo-piceae, articulo primo tertioque obscurioribus, ultimo apice ferrugineo. Palpi rufi. Labrum rufo-piceum. Caput thoracis fere longitudine eoque parum latius, basi cum angulis posterioribus rotundatum, antrorsum subangustatum, supra leviter convexum, utrinque crebrius sat profunde punctatum, medio longitudinaliter laevi, spatio laevi biserialim punctato, sulculis anticis intermediis sat fortibus, ad mediam frontem usque adscendentibus, lateribus obliquis ad oculorum marginem inferiorem fere omnino obsoletis. Thorax coleopteris paulo angustior, latitudine sesqui longior, basin versus subangustatus, lateribus pone medium subsinuatis, angulis anterioribus rotundatis, posterioribus obtusis, parum convexus, seriebus dorsalibus minus regularibus, 7 — 9-punctatis, serie laterali litua 8 — 9-punctata, punctis minus subtilibus. Scutellum obsolete punctatum. Elytra thoracis longitudine, dorso minus dense subseriatim profundius punctata, secundum marginem lateralem laevia, margine inflexo iterum punctato, rufo-brunnea, nitida. Abdomen lineare, parce subtiliterque punctulatum, segmentis singulis margine summo brunneo. Pedes nigri, tibiis rufo-piceis, tarsis dilutius rufis.

- \* 44. *X. humeralis*: *Nigro-aeneus, nitidus, elytris angulo humerali pallido, capite subquadrato, utrinque fortiter punctato, thorace seriebus dorsalibus irregularibus 6-punctatis.* — Long.  $3\frac{1}{2}$  lin.

Habitat in Americae meridionalis ins. St. Johannis, Dom. Moritz.

Statura omnino *X. punctulati*, at paulo maior, nigro-aeneus, pernitidus. Antennae capite sesqui longiores, crassiusculae, articulo tertio secundo paulo longiore, 4 — 10 sensim paululum crassioribus, 5 — 10 transversis, rufo-piceae, articulo ultimo apice ferrugineo. Palpi rufi. Caput thorace latius, eiusque prope longitudine, orbiculato-subquadratum, basi truncatum, angulis posterioribus rotundatis, lateribus subrectis, antrorsum haud angustatum, supra parum convexum, utrinque vage paraeque profunde punctatum, spatio medio longitudinali laevi, sulculis anticis intermediis sat profundis, subflexuosis, ad mediam frontem usque adscendentibus, parum conniventibus, lateribus obliquis ad oculi utriusque marginem inferiorem valde obsoletis. Thorax coleopteris parum angustior, latitudine tertia parte longior, basin versus leniter angustatus, lateribus utrinque subsinuatus, apice utrinque oblique truncatus, angulis anterioribus subrotundatis, leviter convexus, seriebus dorsalibus irregularibus, postice punctis nonnullis lateraliter adiectis auctis, cir-

citer sexpunctatis, serie laterali valde flexuosa circiter 9-punctata, punctis fortioribus. Scutellum laeve. Elytra thoracis longitudine, dorso crebrius profundiusque subseriatim punctata, secundum marginem lateralem laevia, margine laterali inflexo subtiliter punctato, angulo humerali luteo. Abdomen parce subtiliter punctatum, tenuiter longius griseo-pilosum, segmento quinto sextoque margine apicali summo testaceo. Pedes picei, coxis femoribusque nigro-aeneis, geniculis tarsisque rufo-piceis. Alae parum infuscae.

45. *X. rufescens*: *Piceus, nitidus, abdomine capiteque nigro-aeneis, hoc utrinque parce profundeque punctato, thorace seriebus dorsalibus 6—7-punctatis, elytris dorso vage punctatis.* — Long.  $2\frac{3}{4}$  lin.

Habitat in montibus prope Puerto Cabello Columbiae, Dom. Moritz.

Statura fere *X. glabri*, brunneo-piceus, nitidus, nitore aeneo perfusus, capite nigro-subaeneo, nitidissimo. Antennae capite sesqui longiores crassiusculae, articulo secundo tertioque subaequalibus, 4 — 10 transversis, sensim paulo crassioribus, rufo-piceae. Palpi dilute rufo-picei. Caput thorace parum latius et paulo brevius, basi cum angulis posterioribus rotundatum, lateribus subrectis, antrorsum haud angustatum, subdepressum, utrinque punctis parcis profundisque notatum, sulculis anticis intermediis sat profundis, leviter flexuosis, vix conniventibus, ad mediam frontem usque adscendentibus, lateralibus obliquis ad oculorum marginem inferiorem brevioribus, obsoletis. Thorax coleopteris paulo angustior, latitudine sesqui longior, basin versus leviter angustatus, lateribus pone medium vix sinuatis, angulis anterioribus obtusis, subrotundatis, leviter convexus, seriebus dorsalibus minus regularibus, sat fortiter 6- vel 7- punctatis, serie laterali litua 7- vel 8-punctata. Elytra thoracis longitudine, dorso et lateribus crebrius et profundius vage punctata. Abdomen obscure aeneum, nitidum, apice summo piceo-testaceo, tenuiter griseo-pilosum. Pedes dilute rufo-picei.

- \* 46. *X. punctulatus*: *Niger, nitidus, elytris nigro-virescentibus, capite dense punctato-ruguloso, fronte medio laeviore, thorace seriebus dorsalibus 6 — 11-punctatis.* — Long. 3 —  $3\frac{1}{2}$  lin.

Boisd. et Lacord. Faun. Ent. Paris. I. 415. 6. — Er. Col. March. I. 426. 6.

*Staph. punctulatus* Payk. Mon. Staph. 30. 22. Faun. Suec. III. 380. 17. — Fab. Ent. Syst. I. II. 528. 45. Syst. El. II. 600. 56. — Panz. Ent. Germ. 357. 35. — Gyll. Ins. Suec. II. 353. 68. — Sahlb. Ins. Fenn. I. 333. 67.

*Gyrophyn. punctulatus* Mannerh. Brachelytr. 33. 4. — Runde Brachelytr. Hal. 11. 4. — Nordm. Symb. 117. 7.

*Staph. elongatus* Fourc. Ent. Par. I. 171. 27. — Latr. Hist. nat. Crust. et Ins. IX. 330. 84. — Grav. Mon. 98. 98.

*Staph. elongatus* var. 5 Grav. Micr. 45. 66.

*Staph. fracticornis* Müll. Prodr. 99. 118. — Goetze Ent. Beitr. I. 728. 31.

Var. a: *Elytris antennis pedibusque rufo-piceis.*

*Staph. punctulatus* var. b Payk. Faun. Suec. III. 380. 71.

*Staph. elongatus* var. 1 — 4 Grav. Micr. 45. 66.

*Staph. elongati* var. intermed. Grav. Mon. 99.

*Staph. ochraceus* Gyll. Ins. Suec. II. 352. 67. — Sahlb. Ins. Fenn. I. 332. 64.

*Gyrophyn. ochraceus* Mannerh. Brachelytr. 33. 2. — Runde Brachelytr. Hal. 11. 2. — Nordm. Symb. 113. 2.



*Xantholin. ochraceus* Zetterst. Ins. Lappon. 66. 2.

*Var. b: Testaceo-piceus, elytris, antennis, pedibus anoque testaceis.*

Habitat in omni Europa sat frequens.

X. lineari aequalis, individuus licet maioribus paulo latioribus, niger, nitidus. Antennae capite paulo longiores, articulo secundo tertioque subaequalibus, quarto quintoque sequentibus paulo angustioribus, 6—10 transversis, aequalibus, rufo-piceae, articulo primo saepius nigricante. Caput vix thoracis latitudine eodemque paulo brevius, basi medio subsinuatum, utrinque oblique rotundatum, angulis posterioribus denticulo minutissimo notatis, lateribus subrectis, antrorsum leviter angustatum, minus nitidum, fortiter crebreque punctatum, punctis saepe confluentibus subrugulosum, fronte media laeviore, sulculis anticis intermediis sat fortibus, leviter conniventibus, lateralibus obliquis ad oculi utriusque marginem inferiorem valde obsoletis, parce pubescens, ore plerumque piceo. Thorax coleopteris perparum angustior, latitudine sesqui longior, basin versus vix angustatus, lateribus pone medium subsinuatis, angulis anterioribus rotundatis, leviter convexus, seriebus dorsalibus plerumque 8- vel 9-, laterali flexuosa 8—12-punctatis. Elytra thorace parum longiora, minus crebre, introrsum vage, extrorsum seriatim punctata, secundum marginem lateralem laevia, lateribus inflexis dense subtiliter punctatis, iam nigro-subaenea, iam fusco-testacea. Abdomen parce subtiliter obsoleteque punctulatum, densius griseo-pubescens, parce nigro-pilosellum. Pedes vel nigro- vel rufo-picei.

Variat statura maiore latioreque et minore angustioreque. Specimina maiora colore plerumque nigra, antennis pedibusque obscuris, elytris nigro-aeneis, thoracis in serie dorsali punctis saepius paucioribus, specimina minora statura simul angustiore, thoracis in serie dorsali punctis interdum ultra 8, antennis pedibusque dilute rufo-piceis, elytris rarius nigro-aeneis, plerumque testaceo-piceis vel piceo-testaceis, nonnunquam corpore piceo-testaceo, capite nigro-piceo, antennis, pedibus, elytris anoque dilutioribus differre videntur, at speciminibus numerosis sedule examinatis haec differentiae minus stabiles mihi repertae, ut speciebus distinguendis sufficere videantur.

\* 47. X. *Hesperius*: *Subdepressus, niger, nitidus, elytris nigro-subaeneis, margine apicali testaceo, capite parce subtiliter punctato, thorace seriebus dorsalibus 10—12-punctatis.* — Long.  $3\frac{1}{2}$  lin.

Habitat Ulyssipone in Lusitania, Dom. Com. Hoffmannsegg et Dom. Schüppel, in Andalusia, Dom. Waltl.

Statura X. punctulati et maximis eius speciminibus magnitudine aequalis, at paulo magis depressus, niger, subnitidus. Antennae capite paulo longiores, articulo secundo tertioque aequalibus, 4—10 sensim paulo crassioribus, transversis, rufae, articulo primo medio piceo. Palpi rufi. Mandibulae piceae. Caput basi thorace paululum latius, eoque vix brevius, basi medio subsinuatum, utrinque cum angulis posterioribus rotundatum, lateribus subrectis, antrorsum leviter angustatum, supra parum convexum, parce subtilius punctatum, plaga media longitudinali laevi, sulculis anticis intermediis profundioribus, modice conniventibus, lateralibus obliquis ad oculi utriusque marginem inferiorem sat distinctis, minus brevibus, interne puncto maiusculo terminatis. Thorax coleopteris paulo angustior, latitudine plus sesqui lon-

gior, basin versus leviter angustatus, lateribus angulisque anterioribus subrectis, parum convexus, seriebus dorsalibus 10- vel 12-, lateralibus flexuosis fere 12-punctatis, punctis omnibus subtilioribus. Elytra thorace paulo longiora, dorso interne vage, externe subseriatim punctata, secundum marginem lateralem laevia, lateribus inflexis crebre subtiliter punctatis, nigro-subaenea, nitida, margine apicali luteo-testaceo. Abdomen parce subtiliter obsoleteque punctulatum, tenuiter griseo-pubescens, segmentis singulis margine tenui apicali testaceo. Pedes picei, tibiis tarsisque rufis.

Obs. Capitis thoracisque punctura X. lineari affinis, a quo autem thoracis serie laterali flexuosa haud dispersa, elytris longioribus, apice testaceis, externe evidenter seriatim punctatis distinguitur.

\* 48. *X. obscurus*: *Nigro-piceus, nitidus, elytris pedibusque piceo-testaceis, capite utrinque crebre punctato, thorace seriebus dorsalibus 9—12-punctatis.* — Long. 3 lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Statura omnino X. punctulati et minimis eius individuis aequalis, niger, nitidus. Antennae capite sesqui longiores, articulo secundo tertioque aequalibus, 5 — 10 sensim paululum brevioribus at parum crassioribus, sensum paulo magis transversis, rufo-piceae, articulo primo nigricante. Palpi rufi. Caput oblongum, thoracis fere longitudine latitudineque, basi cum angulis posterioribus rotundatum, lateribus subrectis, antrorsum subangustatum, supra parum convexum, utrinque minus fortiter sat crebre punctatum, spatio medio longitudinali laevissimo, sulculis anticis intermediis sat profundis, leviter convergentibus, lateralibus obliquis ad oculi utriusque marginem inferiorem fere omnino obsoletis. Thorax coleopteris paulo angustior, latitudine sesqui longior, basin versus vix angustatus, lateribus subrectis, angulis anterioribus obtusiusculis, parum convexus, seriebus dorsalibus irregulariter 9 — 12-punctatis, serie laterali litua circiter 14-punctata, punctis omnibus minus subtilibus. Scutellum laeve. Elytra thorace perparum longiora, dorso sat crebre fortiterque subseriatim punctata, secundum marginem lateralem laevia, margine ipso punctorum serie sat regulari notato, lateribus inflexis crebre subtilius punctatis, piceo-testacea, nitida. Abdomen parcius subtiliter punctulatum, tenuiter cinereo-pubescens, ano segmentorumque singulorum margine summo obscure testaceis. Pedes rufo-testacei.

\* 49. *X. attenuatus*: *Niger, nitidus, pedibus testaceis, elytris piceis, apice pallidioribus, capite utrinque crebre punctato, thorace seriebus dorsalibus 7—9-punctatis.* — Long.  $2\frac{2}{3}$  lin.

Habitat in Americae insulis St. Thomae et Puerto-Rico, Dom. Moritz, St. Vincentii, Mus. Dom. Hope, in Brasilia, Dom. Melly.

Statura omnino Leptacini batychroi, et maioribus eius individuis magnitudine aequalis, piceus, pernitidus, capite nigro. Antennae capite paulo longiores, articulo secundo tertioque brevioribus, aequalibus, 4 — 10 brevibus, transversis, inter se aequalibus, totae rufo-piceae. Palpi rufo-testacei. Os piceum. Caput thoracis prope longitudine eoque parum latius, basi leviter rotundatum, angulis posterioribus denticulo prominulo minuto notatis, lateribus subrectis, antrorsum vix angustatum, supra leviter convexum, utrinque

grosse punctatum, spatio medio longitudinali laevisimo, sulcis anticis intermediis profundis, ad mediam frontem usque adscendentibus, leviter convergentibus, sulculis lateralibus obliquis ad oculi utriusque marginem inferiorem breviusculis obsoletisque. Thorax coleopteris parum angustior, latitudine tertia parte longior, lateribus pone medium vix sinuatis, angulis anterioribus obtusis, basin versus vix angustatus, parum convexus, seriebus dorsalibus et serie laterali litua 7—9-punctatis, punctis minus subtilibus. Scutellum laeve. Elytra thorace paulo longiora, dorso interius vage, exterius subseriatim punctata, secundum marginem lateralem laevia, margine ipso punctorum serie sat regulari notato, margine inflexo confertim subtiliter punctato, picea, apice pallida, pellucida. Abdomen parce subtiliter obsoleteque punctulatum, tenuiter griseo-pubescent, nigro-piceum, segmentis singulis margine piceo-testaceis. Pedes rufo-testacei.

Obs. Ad Leptacinum batyehroum habitu, colore et punctura prope accedit, at huius generis est.

4) Thorax punctorum seriebus dorsalibus rectis, integris, lateralibus confusis.

\* 50. *X. procerus*: *Niger, nitidus, elytris, pedibus, ano, ventre pectoreque rufis, capite ovato, utrinque subtiliter punctato, thorace seriebus dorsalibus 15-punctatis.* — Long. 6 lin.

Habitat in Germania boreali, Dom. Schmidt, in Hispania, Mus. Dr. Watl, in Sardinia, Mus. Reg. Taurin.

Forma X. linearis, at magnus, X. glabrato longior et gracilior. Antennae capite paulo longiores, crassiusculae, articulo tertio secundo sesqui longiore, 4—10 sensim paululum crassioribus, leviter transversis, fusco-piceae, articulo basali rufo-piceo, apicali ferrugineo. Caput thorace paulo latius eique longitudine subaequale, ovatum, antrorsum leviter angustatum, supra leviter convexus, utrinque minus dense subtiliter punctatum, medio longitudinaliter laeve, sulculis anticis intermediis brevibus subparallelis, lateralibus obliquis ad oculi utriusque marginem inferiorem subtilioribus, longioribus, interne puncto maiusculo terminatis, nitidum, nigrum, ore rufescente. Thorax latitudine duplo prope longior, basin versus subangustatus, lateribus pone medium subsinuatis, angulis anterioribus obtusis, subrotundatis, supra leviter convexus, niger, nitidus, seriebus dorsalibus irregularibus, subtiliter circiter 15-punctatis, lateribus vage subtiliterque multipunctatis, subtus rufus. Scutellum laeve piceum. Elytra thoracis longitudine, crebrius vage punctata, rufa. Abdomen subtilissime punctulatum, tenuiter griseo-pubescent, apicem versus subincrassatum, supra nigrum, segmentorum marginibus castaneis, ano late rufo, subtus rufum, segmentis anterioribus basi nigris. Pectus et pedes rufi.

\* 51. *X. tricolor*: *Rufo-testaceus, nitidus, capite, thoracis dorso postico abdomineque supra nigro-piceis, capite utrinque subtiliter punctato, thorace seriebus dorsalibus 12-punctatis.* — Long. 4 lin.

Zetterst. Faun. Lappon. I. 66. 4. Ins. Lappon. 66. 4. — Er. Col. March. I. 427. 7. Staph. tricolor Fab. Mant. Ins. I. 221. 30. — Payk. Mon. Staph. 33. 15. Faun. Suec. III. 378. 15. — Latr. Hist. nat. Crust. et Ins. IX. 330. 86. — Gyll. Ins. Suec. II. 355. 70. — Sahlb. Ins. Fenn. I. 334. 69.



- Paederus tricolor* Fab. Ent. Syst. I. II. 537. 7. Syst. El. II. 609. 7. — Panz. Ent. Germ. 363. 8.  
*Gyrophyn. tricolor* Mannerh. Brachelytr. 33. 7. — Runde Brachelytr. Hal. 11. 6. — Nordm. Symb. 118. 11.  
*Staph. elegans* Grav. Micr. 46. 68. Mon. 103. 104.  
*Staph. affinis* Marsh. Ent. Brit. 517. 57.

Habitat in Europa.

Statura fere *X. linearis* at paulo maior et minus gracilis. Antennae capite paulo longiores, articulo secundo tertioque subaequalibus, 4 — 10 sensim paulo crassioribus et magis transversis, totae rufae vel rufo-piceae. Palpi rufi. Caput magnum, thoracis longitudine eodemque paulo latius, oblongo-ovatum, antrorsum leviter angustatum, nigro-piceum, nitidum, supra leviter convexum, utrinque subtiliter sparsim punctatum, spatio medio longitudinali laevissimo, sulculis apicalibus intermediis sat profundis, parallelis, brevioribus, lateralibus obliquis ad oculi utriusque marginem inferiorem longioribus, at subtilibus, interne puncto maiusculo terminatis. Thorax latitudine media duplo prope longior, basin versus leviter angustatus, lateribus vix sinuatus, apice truncatus, angulis anterioribus subrotundatis, leviter convexus, rufo-testaceus, basin versus plerumque magis minusve piceus, nitidus, seriebus dorsalibus subtiliter 12-punctatis et ultra, lateribus subtiliter disperse punctatus. Scutellum piceum, laeve. Elytra thorace distincte breviora, minus subtiliter crebrius punctata, rufo-testacea, nitida. Abdomen subtiliter parcius punctatum, tenuiter griseo-pubescent, nitidum, supra nigrum, segmento quinto apice sextoque toto testaceis, subtus cum pectore pedibusque rufo-testaceum.

Variat thorace rufo-testaceo, unicolore.

✕ \* 52. *X. linearis*: Nigro-subaeneus nitidus, capite utrinque subtiliter punctato, thorace seriebus dorsalibus 12-punctatis. — Long.  $3\frac{1}{2}$  lin.

Er. Col. March. I. 428. 8.

*Staph. linearis* Oliv. Ent. III. 42. 19. 21. t. 4. f. 38. — Fab. Ent. Syst. Suppl. 180. 41 — 42. Syst. El. II. 599. 51. — Marsh. Ent. Brit. 516. 54.

*Staph. longiceps* Grav. Micr. 177. 36. — Latr. Hist. nat. Crust. et Ins. IX. 334. 94. — Gyll. Ins. Suec. II. 350. 66. — Sahlb. Ins. Fenn. I. 331. 63.

*Xantholin. longiceps* Zetterst. Faun. Lappon. I. 80 1. Ins. Lappon. 66. 1.

*Gyrophyn. longiceps* Mannerh. Brachelytr. 33. 1. — Runde Brachelytr. Hal. 11. 1. — Nordm. Symb. 114. 3.

*Staph. punctulatus* Schrank Faun. Boica I. 649. 889. — Grav. Micr. 177. 37. Mon. 98. 96.

Var. Corpore piceo-testaceo, elytris pedibusque dilutioribus.

*Staph. ochraceus* Grav. Micr. 43. 65. Mon. 97. 95. — Latr. Hist. nat. Crust. et Ins. IX. 329. 83.

*Xantholin. ochraceus* Boisd. et Lacord. Faun. Ent. Paris. I. 416. 8.

Habitat in Europa.

Elongatus, tenuis, niger, nitidus, capite, thorace elytrisque nigro-subaeneis, his saepe fusco-piceis; variat etiam saepe corpore toto piceo, elytris pedibusque testaceis, vel fusco-testaceo, elytris pedibusque dilutioribus. Antennae capite sesqui longiores, crassiusculae, articulo secundo tertioque subaequalibus, 4 — 10 sensim paulo crassioribus et magis transversis, rufo-

piceae, apice ferrugineo. Palpi rufi. Labrum rufum. Caput thoracis longitudine eodemque vix latius, oblongo-ovatum, antrorsum leviter angustatum, supra leviter convexum, utrinque subtiliter disperse punctulatum, spatio medio longitudinali laevi, sulculis anticis intermediis brevibus, sat profundis, parallelis, lateralibus obliquis ad oculi utriusque marginem inferiorem subtilioribus et paulo longioribus. Thorax coleopteris paulo angustior, latitudine antica plus sesqui longior, basin versus leniter angustatus, lateribus pone medium subsinuatis, apice truncatus, angulis anterioribus rotundatis, leviter convexus, seriebus dorsalibus subregularibus, 12-punctatis et ultra, lateribus disperse punctatis, punctis omnibus sat subtilibus. Scutellum laeve. Elytra thorace paululum breviora, subtiliter subseriatim punctata, secundum marginem lateralem laevia, tenuiter griseo-pubescentia. Abdomen parce punctulatum, tenuiter griseo-pubescentia. Pedes picei vel testacei.

### **Leptacinus Er.**

*Er. Col. March. I. p. 429.*

*Antennae geniculatae.*

*Labrum margine laterali membranaceo.*

*Palpi articulo ultimo subulato.*

*Ligula apice emarginata.*

Corpus valde elongatum, lineare, subdepressum, alatum. Caput oblongum, basi profunde constrictum, collo quasi brevi cylindrico thoraci affixum, exsertum, porrectum, oculis parvis, rotundatis, apicem versus positis, vix prominulis. Labrum angustum, transversum, medio profunde sinuatum, limbo laterali membranaceo, apice setosum. Mandibulae breviores, medio dentatae. Maxillae (t. IV. f. 22. b) mala interiore membranacea, interne barbata, exteriore cornea, apice membranacea barbataque. Palpi maxillares articulo primo parvo, sequentibus duobus subaequalibus, quarto minuto, subulato. Labium (t. IV. f. 22. a.) mento transversum, ligula brevi, rotundata, medio emarginata, membranea, paraglossis ligulam superantibus, membranaceis. Palpi labiales articulo secundo primo paululum longiore, ultimo minuto, subulato. Antennae basi inter se non minus quam oculis approximatae, breves, geniculatae, articulo primo leviter elongato, secundo tertioque aequalibus, obconicis, reliquis brevibus, pubescentibus, ultimo subovato. Thorax oblongus, basin versus subangustatus, basi rotundatus, apice truncatus, angulis anterioribus subrectis, subdeflexis. Scutellum triangulare. Elytra apice truncata, sutura imbricata. Abdomen lineare. Pedes breves, coxis intermediis vel modice distantibus, vel approximatis, tibiis spinulosis, anticis externe subincrassatis, extus spinuloso-muricatis, tarsis aut omnibus simplicibus aut anticis dilatatis.

#### **O b s e r v a t i o n e s.**

1. Habitu omnino generis praecedentis, et sectio secundo nonnisi palporum et ligulae structura ab eodem distincta. Notatu dignum videtur, pedum intermediarum insertionem et tarsorum anticorum structuram, in illo genere constantissimas, in hoc tam propinquo variare.

2. Sexus differentia latet.

3. Victus et mores cum genere praecedente conveniunt.

\* *Coxae intermediae approximatae. Tarsi antici simplices.*

\* 1. *L. brevicornis: Niger, nitidus, pedibus rufis, capite punctulato.*  
*Long. 3½ lin., lat. ½ lin.*

Habitat in Austria, Dr. Walzl.

Corpus valde elongatum, tenue, subcylindricum, nigrum, pernitidum. Antennae capite breviores, articulo secundo tertioque brevibus, aequalibus, 4—10 brevissimis, sensim paulo crassioribus, subcompressis, fortiter transversis, penultimis latitudine triplo brevioribus, ultimo praecedente duplo fere longiore, apice abrupte acuminato, pilosellae, rufo-piceae. Caput thorace paulo longius latiusque, basi truncatum, angulis posterioribus leviter rotundatis, lateribus rectis, antrorsum subangustatum, supra leviter convexum, densius subtiliter punctatum, spatio medio angusto laeviore, sulculis duobus anticis inter antennis brevibus obsoletioribus, spatio interiacente elevato, prominulo, canaliculato, sulculis obliquis ante oculos nullis, subtus parcius punctulatum, canaliculatum, utrinque secundum marginem lateralem sulcatum. Thorax coleopteris parum angustior, latitudine sesqui longior, basin versus subangustatus, lateribus medio subsinuatis, angulis anterioribus late rotundatis, leviter convexus, subtilissime parce punctulatus, dorso utrinque punctis nonnullis maioribus, lateribus sulculo leviter flexuoso tenui obsoleto imsculptus. Elytra thorace paulo longiora, subtilissime obsoleteque vage punctulata. Abdomen laevigatum, parce nigro-pilosellum. Pedes rufi, tarsi testaceis.

\* 2. *L. filarius: Niger, nitidus, elytris pedibusque piceis, capite laevi.*  
*— Long. 3 lin.*

Habitat in Columbia, Dom. Moritz.

Affinis praecedenti, at minus cylindricus, niger, nitidus. Antennae capitis longitudine, articulo secundo tertio paulo maiore, obconicis, quarto paulo brevior et crassior, subcylindrico, 5—10 fortiter transversis, subcompressis, ultimo praecedente sesqui longiore, ovato, dense pilosellae, rufo-piceae, ultimo apice ferrugineo. Palpi rufo-picei. Caput thoracis longitudine eodemque distincte latius, basi truncatum, angulis posterioribus leviter rotundatis, lateribus rectis, antrorsum subangustatum, latitudine parum longius, ideoque subquadratum, subdepressum, sulculis anticis inter antennis profundioribus, at brevissimis, distantibus, parallelis, spatio interiacente plano, sulculis lateralibus obliquis ante oculos sat distinctis, ad mediam frontem adscendentibus, in ipso oculi utriusque margine lineola brevi, supra oculum utrumque lineola longitudinali, et supra hanc lineolam puncto oblongo impressum, ceterum laevissimum. Thorax antice coleopteris vix angustior, latitudine plus sesqui longior, basin versus leniter angustatus, lateribus medio subsinuatis, utrinque oblique truncatus, angulis anterioribus obtusis, subdepressus, paulo ante medium dorsi punctis 4 transversim positus, et paulo infra haec exterius utrinque puncto singulo, summo apice utrinque punctis duobus impressis notatus. Elytra thoracis longitudine, laevia, testaceo-picea. Abdomen laevigatum, parce tenuiterque griseo-pilosellum. Pedes picei, tarsi testaceis.



**\*\* Coxae intermediae distantes. Tarsi antici simplices.**

\* 3. *L. parum punctatus*: Niger, nitidus, elytris piceo-testaceis, angulo apicali pallido, capite utrinque fortiter punctato, thorace seriebus dorsalibus 5 — 6-punctatis. — Long.  $2\frac{3}{4}$  lin.

*Staph. parumpunctatus* Gyll. Ins. Suec. IV. 481. 67 — 68. — Sahlb. Ins. Fenn. I. 333. 66.

*Xantholin. parumpunctatus* Boisd. et Lacord. Faun. Ent. Paris. I. 414. 5.

*Gyrohypn. parumpunctatus* Mannerh. Brachelytr. 33. 5. — Runde Brachelytr. Hal. 11. 5. — Nordm. Symb. 117. 9.

Habitat in Germania, Suecia, Dom. Schüppel.

Statura omnino sequentis, at paulo maior, niger, nitidus. Antennae capite parum longiores, articulo secundo tertio vix maiore, his obconicis, 4—10 cylindricis, transversis, inter se subaequalibus, ultimo apice abrupte acuminato, rufo-piceae. Palpi rufi. Caput thoracis longitudine eodemque basi parum latius, antrorsum angustatum, basi medio leviter emarginatum, angulis posterioribus rotundatis, parum convexum, utrinque punctis magnis oblongis minus crebre impressum, spatio lato medio longitudinali interstitisque laevissimis, sulcis anticis intermediis et lateralibus ante oculos longitudine aequalibus et aequaliter fortibus profundisque. Thorax coleopteris paulo angustior, latitudine longior, basin versus subangustatus, lateribus pone medium subsinuatis, apice utrinque oblique truncatus, angulis anterioribus obtusis, subrotundatis, parum convexus, seriebus dorsalibus fortiter 5 — 6-punctatis, serie laterali litua 5-punctata. Elytra thoracis longitudine, parce subtilius seriatim punctata, nigro-picea, apicem versus saepe sensim piceo-testacea, angulo apicis exteriore pallido. Abdomen parce subtiliter punctatum, tenuiter griseo-pubescens pilosellumque, subtus segmentis ultimis apice piceo-testaceis. Pedes testacei.

\* 4. *L. batychrus*: Niger, nitidus, elytris piceo-testaceis, angulo apicali pallido, capite utrinque crebre punctato, thorace seriebus dorsalibus 12 — 16-punctatis. — Long. 2 —  $2\frac{1}{2}$  lin.

*Er. Col. March. I. 429. 1.*

*Staph. batychrus* Gyll. Ins. Suec. IV. 480. 67 — 68. — Sahlb. Ins. Fenn. I. 332. 65.

*Gyrohypn. batychrus* Mannerh. Brachelytr. 33. 3. — Runde Brachelytr. Hal. 11. 3. — Nordm. Symb. 117. 8.

*Staph. linearis* Grav. Micr. 43. 64. Mon. 97. 94. — Latr. Hist. nat. Crust. et Ins. IX. 329. 82.

*Gyrohypn. linearis* Nordm. Symb. 120. 20.

*Xantholin. episcopalis* Boisd. et Lacord. Faun. Ent. Paris. I. 416. 7.

*Xantholin. minutus* Boisd. et Lacord. Faun. Ent. Paris. I. 417. 9.

Habitat in Europa et in Asia minore.

Habitu fere Xantholini punctulati, at minimis eius individuis duplo, triplo vel quadruplo minor, magnitudine varians, subdepressus, niger, nitidus. Antennae capite paulo longiores, articulo secundo tertioque subaequalibus, obconicis, 4—10 cylindricis, vix transversis, inter se aequalibus, ultimo subovato, apice leniter acuminato. Caput thoracis longitudine, eodemque basi paulo latius, basi leviter, angulis posterioribus fortius rotundatis, lateribus subrectis, antrorsum leniter angustatum, minus depressum, utrinque

crebrius fortiter punctatum, interstitiis spatioque medio longitudinali laevis-  
simis, sulcis anticis intermediis leviter convergentibus, profundis, lateralibus  
obliquis ante oculos sat distinctis, licet minus fortibus, utrisque ad mediam  
frontem prope adscendentibus. Thorax coleopteris parum angustior, latitu-  
dine plus sesqui longior, basin versus leniter angustatus, lateribus pone  
medium subsinuatis, angulis anterioribus obtusis, subrotundatis, parum con-  
vexus, seriebus dorsalibus sat regularibus, 12-punctatis et ultra, serie late-  
rali litua multipunctata. Scutellum laeve. Elytra thoracis longitudine, dorso  
minus crebre, introrsum vage, extrorsum seriatim, lateribus inflexis subtiliter  
subseriatim punctata, piceo-testacea, basi obscuriora, saepe nigricantia, apice  
sensim dilutiora, saepius lutescentia, angulo apicali exteriori saepius late  
pallido. Abdomen subtiliter parce punctatum, parce tenuiter griseo-pube-  
scens, segmentis duobus ultimis apice piceo-testaceis. Pedes piceo-testacei,  
tarsis dilutioribus.

- \* 5. *L. tenellus*: *Piceus, nitidus, elytris piceo-testaceis, capite oblongo, utrinque subtiliter punctato, thorace seriebus dorsalibus 10-punctatis.* — Long. 2 lin.

Habitat in Madagascar, Dom. Goudot.

Habitu omnino Xantholini linearis, at multo minor, subdepressus, te-  
staceo-piceus, nitidus, capite nigro. Antennae articulo secundo obconico,  
tertio hoc brevior, subcylindrico, 4 — 10 brevibus transversisque, sensim  
paulo crassioribus, ultimo maiusculo, ovato, totae dilute rufo-piceae. Palpi  
testacei. Caput thoracis fere longitudine eoque parum latius, basi trunca-  
tum, angulis posterioribus rotundatis, lateribus subrectis, antrorsum vix an-  
gustatum, subdepressum, utrinque parcius subtiliter punctatum, spatio medio  
longitudinali laevis-  
simo, sulculis duobus apicalibus intermediis sat profundis,  
modice convergentibus, sulculis ante oculos distinctis, licet brevibus subti-  
libusque. Thorax coleopteris angustior, latitudine duplo prope longior,  
basin versus leniter angustatus, lateribus pone medium subsinuatis, angulis  
anterioribus obtusis, parum convexus, seriebus dorsalibus serieque laterali  
litua subtiliter circiter 10-punctatis. Scutellum laeve. Elytra vix thoracis  
longitudine, parce subtiliterque seriatim punctata. Abdomen parce subtilis-  
sime obsoleteque punctulatum, tenuiter griseo-puberulum. Pedes testacei.

6. *L. debilis*: *Nigro-piceus, nitidus, antennis, elytris pedibusque fusco-testaceis, capite utrinque parce punctato, thorace seriebus dorsalibus 8-punctatis.* — Long. 1½ lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Statura et summa affinitas praecedentis. Antennae capite sesqui lon-  
giores, articulo tertio secundo duplo fere minore, apicem versus leviter in-  
crassatae, articulis 4 — 10 sensim paulo crassioribus et magis transversis,  
ultimo ovato, fusco-testaceae, articulo primo testaceo, ultimo ferrugineo.  
Os fusco-testaceum. Caput thorace paulo latius, latitudine parum longius,  
basi truncatum, angulis posterioribus rotundatis, lateribus rectis, antrorsum  
subangustatum, subdepressum, utrinque punctis parvis minus subtilibus im-  
pressum, spatio medio longitudinali lato laevi, sulcis anticis intermediis sat  
profundis, parallelis, ad mediam frontem fere adscendentibus, lateralibus obli-

quis ante oculos brevibus et valde obsoletis, nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine longior, basin versus subangustatus, lateribus pone medium subsinuatis, angulis anterioribus obtusis, subrotundatis, parum convexus, niger, nitidus, seriebus dorsalibus sat regularibus, 8-punctatis, serie laterali litua circiter 6-punctata. Scutellum laeve, piceum. Elytra thoracis longitudine, introrsum vage, extrorsum biseriatim punctata, piceo-testacea, nitida. Abdomen laeve, tenuiter griseo-pilosellum, piceum, segmentorum marginibus dilutioribus. Pectus piceum. Pedes testacei.

\* 7. *L. subtilis*: *Piceus, nitidus, antennis pedibusque testaceis, capite thoraceque lateribus elytrisque subtilissime punctulatis.* — Long. 2 lin.

Habitat in Columbia, Dom. Moritz.

Statura tenui praecedentis, piceus, nitidus, capite, thorace elytrisque fusco-subaeneis. Antennae capite sesqui longiores, apicem versus sensim fortius incrassatae, articulo secundo tertioque obconicis, hoc paulo brevior, 4 — 10 fortiter transversis, sensim crassioribus, arcte cohaerentibus, ultimo praecedente plus duplo longiore, apice obtuse acuminato, rufo-testaceae. Palpi testacei. Caput thorace parum latius, oblongum, basi cum angulis posterioribus rotundatum, lateribus subrectis, antrorsum vix angustatum, supra leviter convexum, utrinque parce subtilissime punctatum, sulculis anticis intermediis sat distantibus, parallelis, lateralibus obliquis ante oculos obsoletis, utrisque brevibus. Thorax coleopteris paulo angustior, latitudine plus duplo fere longior, basin versus leviter angustatus, lateribus anterieus leviter rotundatis, posterius subsinuatis, angulis anterioribus omnino nullis, transversim convexus, utrinque vage subtilissime punctatus. Elytra thoracis longitudine, obsolete subtilissime vage punctata, parce subtiliterque pubescentia. Abdomen parce subtilissime punctatum, segmentis singulis tenuiter testaceo-marginatis. Pedes testacei.

8. *L. clavicornis*: *Niger, nitidus, antennis, pedibus capiteque rufo-piceis, hoc thoraceque lateribus subtiliter punctatis, elytris vage subtiliter punctatis, margine apicali testaceo.* — Long. 2½ lin.

Habitat in valle Araguensi Columbiae, Dom Moritz.

Habitu praecedentium, at paulo maior. Antennae capite paulo longiores, articulo secundo tertioque obconicis, apicem versus fortiter incrassatae, articulis 4 — 10 sensim crassioribus et fortius transversis, ultimo conico, piceo-rufae. Palpi testacei. Caput thorace parum latius, oblongum, basi truncatum, angulis posterioribus late rotundatis, antrorsum subangustatum, parum convexus, utrinque crebrius subtiliter punctatum, sulcis anticis intermediis brevissimis, sulculis lateralibus ante oculos omnino obsoletis, piceo-rufum, nitidum. Thorax coleopteris paulo angustior, latitudine duplo fere longior, basin versus leviter angustatus, lateribus pone medium subsinuatus, angulis anterioribus subrectis, leviter convexus, nigro-piceus, nitidus, utrinque vage subtiliter punctatus, spatio angusto longitudinali medio laevi. Scutellum laeve, piceum. Elytra thoracis longitudine, vage subtilius punctata, nigra, nitida, margine apicali testaceo. Abdomen subtiliter punctulatum, nigrum, nitidulum, subtiliter tenuiterque pubescens, segmentis anterioribus margine apicali summo piceo-testaceo. Pedes testacei.



\*\*\* Coxae intermediae distantes. Tarsi antici dilatati.

- \* 9. *L. nothus*: *Subdepressus, niger, antennis pedibusque rufo-ferrugineis, capite confertissime punctato-subrugoso, thorace elytrisque dense punctatis, illo linea media longitudinali laevi.* — Long.  $2\frac{1}{2}$  — 3 lin.

Habitat in Croatia meridionali, Dom. Schüppel, in Mesopotamia, Dom. Helfer.

Corpus elongatum, lineare, subdepressum, nigrum, nitidum, totum breviter tenuiterque griseo-pubescens. Antennae capite sesqui longiores, articulo primo sat elongato, secundo tertioque modice elongatis, aequalibus, 4—10 transversis, ultimo penultimo sesqui longiore, cylindrico, apice rotundato, totae rufo-ferrugineae. Palpi testacei. Os piceum. Caput thorace vix latius, oblongum, basi utrinque oblique truncatum, angulis posterioribus rotundatis, lateribus subrectis, antrorsum vix angustatum, supra parum convexum, confertissime punctatum, longitudinaliter subrugulosum, opacum, linea tenui obsoletaque longitudinali sublaevi, sulculis anticis intermediis brevissimis lateralibusque obliquis ante oculos perbrevibus obsoletis. Thorax coleopteris parum angustior, latitudine duplo longior, basin versus subangustatus, apice cum angulis anterioribus fortiter rotundatus, lateribus medio sinuatus, subdepressus, crebre punctatus, linea media longitudinali laevi. Scutellum leviter impressum. Elytra vix thoracis longitudine, dense punctata. Abdomen densius subtiliter punctulatum, apice piceum. Pedes breviusculi, rufo-picei, tarsi fulvis, anticis dilatatis.

\*\*\*\* Coxae intermediae approximatae. Tarsi antici dilatati.

- \* 10. *L. tenuicornis*: *Piceo-testaceus, opacus, abdomine fusco, segmentorum marginibus testaceis, capite thoraceque obscure aeneis, utrinque subtiliter punctatis.* — Long. 4 lin.

*Gyrophyn. tenuicornis Nordm. Symb. 125. 27.*

Habitat in Promontorio bonae spei, Dom. Bergius.

Maximis Xantholini punctulati individuus paulo maior. Antennae capitis longitudine, apicem versus haud incrassatae, articulo primo sat elongato, tertio leviter elongato, secundo sesqui longiore, 4 — 10 sensim paululum brevioribus, ultimo ovato, apice subacuminato, rufae, apice summo ferrugineo. Palpi rufi. Mandibulae piceae. Caput thorace vix latius eodemque paulo brevius, basi et lateribus leviter, angulis posterioribus fortius rotundatum, antrorsum parum angustatum, supra leviter convexum, obscure aeneum, parum nitidum, fronte utrinque subtilius parce punctata, sulculis duobus anticis intermediis brevioribus, parallelis, sat distinctis, lateralibus obliquis ante oculos obsoletioribus, inter oculos punctis quatuor maioribus transversim positus minus profundis impressum. Thorax coleopteris paulo angustior, latitudine sesqui longior, basin versus subangustatus, lateribus rectis, apice truncatus, angulis anterioribus rectis, leviter transversim convexus, obscure aeneus, limbo tenuiore indeterminate fusco-piceo, parum nitidus, utrinque parcius disperse subtiliter punctatus, spatio medio longitudinali laevi. Scutellum fusco-testaceum, laeve. Elytra thorace paulo breviora, fusco-testacea, aeneo-sub-micantia, aequaliter crebrius punctata. Abdomen subtilissime punctulatum, nigro-piceum, opacum, segmentis margine fusco-testaceis. Pectus rufo-piceum. Pedes rufo-testacei.

## Subtribus secunda.

### Genuini.

*Stigmata prothoracica detecta.*

*Antennae basi distantes.*

Labrum transversum, bilobum, limbo exteriore coriaceo membranaceo.

Mandibulae magis minusve prominentes.

Maxillae malis coriaceis, exteriore basi cornea, hac apice, illa intus barbatis;

utraque breviuscula plerisque,  
exteriore elongata, Haematodi, Cordylaspidi, Scariphaeo, Palaestrino.

Palpi maxillares filiformes plerisque;

articulo ultimo truncato, subsecuriformi Scariphaeo, Palaestrino;

articulo ultimo subulato Heterothopi.

Ligula minuta, rotundata:

medio incisa vel biloba Haematodi, Cordylaspidi, Scariphaeo, Staphylino, Ocypodi;

integra, Belonucho, Philontho, Heterothopi, Acylophoro.

Paraglossae ligulam longe superantes.

Palpi labiales filiformes plerisque;

articulo tertio magis minusve securiformi Staphylini fam. XI, Ocypodis fam. II et III;

articulo tertio hemisphaerico Staphylini fam. XII;

articulo tertio subulato Heterothopi.

Antennae rectae, filiformes vel apicem versus leniter incrassatae, articulis tribus primis leviter elongatis, ultimo saepius apice truncato vel emarginato et inferne acuminato plerisque;

compressae, apicem versus attenuatae Cordylaspidi;

in clavam fusiformem coactae Haematodi;

serratae, articulo ultimo tricuspidato Staphylini fam. X;

fractae, articulo primo fortius elongato Acylophoro.

Caput basi constrictum, collo quasi crassiusculo thoraci insertum.

Thorax basi rotundatus, apice truncatus.

Elytra truncata, sutura integerrima.

Abdomen aut parallelum aut apicem versus leniter angustatum.

Pedes plerumque breviusculi, tibiis saepissime aut omnibus aut posterioribus spinulosis, tarsis aut omnibus simplicibus aut anticis minus magisve dilatatis, posterioribus articulis 4 primis longitudine decrescentibus, teretibus plerisque;

totis depressis Haematodi, Cordylaspidi, Scariphaeo.

**Haematodes Laporte.***Laporte Etud. Ent. p. 113.**Platycnemus Nordm. Symb. p. 135.**Antennae rectae, breves, compressae, clavatae.**Palpi articulo ultimo acuminato.**Maxillae mala exteriore elongata.**Ligula apice emarginata.**Pedes intermediï basi distantes.**Tarsi posteriores depressi.*

Corpus maioris magnitudinis, oblongum, convexum, alatum. Caput thorace vix angustius, suborbiculatum, supra convexum, basi modice contractum, thoraci immissum, oculis parvis, obliquis, haud prominulis. Labrum corneum, bilobum, margine membranaceo setoso. Mandibulae subfalcatae, validiores, medio dentatae. Maxillae (t. IV. f. 16. b.) mala interiore oblonga, coriacea, interne barbata, exteriore elongata, cornea, apice ciliato-barbata. Palpi maxillares breves, malam exteriorem haud excedentes, articulo primo minuto, secundo tertioque subaequalibus, obconicis, apice setosis, quarto parum longiore, subacuminato. Labium (t. IV. f. 16. a.) mento transverso, ligula sat prominula, rotundata, apice leviter sinuata, membranea, paraglossis divergentibus, ligulam superantibus, membraneis, introrsum dense ciliatis. Palpi labiales articulo primo minuto, secundo crassiusculo, obconico, tertio huic longitudine subaequali, leviter ovato. Antennae perbreves, articulo primo breviusculo, obconico, piloso, reliquis glabris, arcte coactis, compressis, clavam oblongam fusiformem formantibus, ultimo apice emarginato. Thorax coleopterorum latitudine, transversus, transversim convexus, basi fortiter rotundatus, lateribus deflexis, rectis, apice truncatus, utrinque subsinuatus, angulis posterioribus obtusis, anterioribus deflexis, acuriusculis. Scutellum sat magnum, rotundato-triangulare. Elytra apice extrorsum leviter rotundata, introrsum oblique truncata. Abdomen breviusculum, apice attenuatum. Pedes breves, crassi, coxis intermediis modice distantibus, tibiis femoribus recipiendis profunde canaliculata, tibiis compressis, externe spinulis numerosis muricatis, tarsis perbrevibus, depressis, articulo primo sequentibus vix longiore, anticis articulis 4 primis leviter dilatatis, subtus utrinque fasciculato-setosis.

**O b s e r v a t i o.**

Habitu quodammodo ad Staphylinum hirtum accedit, at thorax latior et convexior, ceterum antennarum structura et tibiis externe muricatis hoc genus sat insigne.

- \* 1. *H. bicolor*: *Convexus, niger, nigro-pilosus, capite, thorace, elytris anoque rufis.* — Long. 3 lin.

*Laporte Etud. Ent. I. p. 113. t. 3. f. 6.*

*Platycnemus lateritius Nordm. Symb. 135. 1. t. 2. f. 4.*

Habitat in Brasilia meridionali, Sellow.

Antennae capite breviores, piceae, nitidae, articulis compressis, angulis fusco-pubescentibus. Mandibulae nigrae. Caput thorace angustius, orbiculatum, convexum, rufum, nitidum, nigro-pilosum, punctatum, margine postico frontisque medio laevibus. Thorax rufus, nitidus, margine laterali nigro,



dorso laevisissimus, ad latera vage punctulatus et nigro-pilosus. Scutellum punctatissimum, nigrum. Elytra thorace vix longiora, leviter convexa, parcius punctata, rufa, nitida, rufo-pubescentia. Abdomen punctulatum, segmentis quatuor anterioribus nigris, nigro-pubescentibus, ultimis duobus rufis, rufo-pubescentibus. Pectus nigrum. Pedes breves, validi, nigri. Alae fuscae.

Mas abdominis segmento inferiore sexto apice leviter emarginato, quinto medio macula magna triangulari rufo-villosa instructo distinctus.

### **Cordylaspis Nordm.**

*Nordm. Symb. p. 17.*

*Smilax Laport. Etud. Ent. p. 116.*

*Antennae rectae, compressae, acuminatae.*

*Palpi articulo ultimo minuto, acuminato.*

*Maxillae mala exteriori elongata.*

*Ligula apice emarginata.*

*Pedes intermedii basi approximati.*

*Tarsi posteriores depressi.*

Corpus maioris magnitudinis, oblongum, dilatatum, utrinque attenuatum, alatum. Caput thorace minus et angustius, transversim suborbiculatum, basi profunde constrictum, collo quasi brevi angustoque thoraci inmissum, oculis haud prominulis. Labrum transversum, bilobum, corneum, membranaceo-limbatum. Mandibulae angustae, acutae, parum arcuatae, muticae. Maxillae (t. IV. f. 17. b.) mala interiore coriacea, intus barbata, exteriori elongata, cornea, apice coriacea barbataque. Palpi maxillares mala exteriori paulo longiores, articulo primo minuto, secundo tertioque aequalibus, leviter elongatis, quarto his duplo brevior, acuminato. Labium (t. IV. f. 17. a.) mento transverso, ligula brevi, biloba, membranea, paraglossis elongatis, linearibus, coriaceis, intus ciliatis, ligulam longe superantibus. Palpi labiales articulo primo brevi, obconico, secundo elongato, crassiusculo, cylindrico, intus spinuloso, tertio hoc duplo brevior tenuioreque. Antennae breves, compressae, apicem versus leviter attenuatae, utrinque pubescentes, articulo primo tertioque reliquis maioribus, ultimo apice emarginato. Thorax transversus, coleopterorum latitudine, antrorsum leviter angustatus, basi lateribusque leviter rotundatus, apice late emarginatus, angulis posterioribus obtusis, anterioribus acutiusculis, leviter transversim convexus. Scutellum triangulare. Elytra thoracis longitudine, apice introrsum oblique truncata, angulo exteriori extrorsum oblique truncato. Abdomen breviusculum, apicem versus sensim angustatum. Pedes breves, coxis intermediis contiguis, femoribus tibiisque compressis, illis tibiis recipientibus, his extus spinulosis, tarsis depressis, articulis arcte coactis, primo subelongato, anticis articulis 4 primis leviter dilatatis coniunctis ovatis, subtus utrinque squamoso-pubescentibus.

### **Observations.**

1. Hoc cum praecedente et sequente genere a genuino Staphylinio differt maxillarum mala exteriori elongata et tarsis posterioribus latiusculis, depressis, (conf. t. IV. f. 23. c), inter se autem corporis habitu, antennarum, palporum et tibiarum structura distinguuntur. *Cordylaspis* praeterea mandibulis muticis discedit.

2. Femora compressa, tibiae retractas partim occultantia, *Drepanum* (Carrorum genus) in animum revocant.

3. Nomen generis a Com. Laporte de Castelnau propositum praetermittendum mihi visum est, quo iam dudum pro notissimo plantarum generi usitati sunt Botanici.

4. Sexus differentiam haud cognovi, at feminas solas mihi observatas esse credo.

5. De victu nihil constat.

\* 1. *C. pilosus*: *Piceus*, capite thorace cyanescentibus nitidulis elytrisque subtiliter tuberculatis utrinque pilosis. — Long. 6 lin.

*Staph. pilosus* Fab. Mant. Ins. I. 220. 10. Ent. Syst. I. II. 523. 18. Syst. El. II. 594. 20. — Oliv. Ent. III. 42. 24. 30. t. 1. f. 3.

*Smilax Americanus* Laporte Etud. Ent. I. p. 117.

*Cordylaspis tuberculatus* Nordm. Symb. 17. 1. t. 1. f. 6.

Habitat in America meridionali: Columbia, Cayenna, Surinamo, Brasilia.

Corpus piceum, opacum, ano rufescente, capite thoraceque supra subcyaneis, nitidis. Antennae capite vix longiores, dilute piceae. Caput fortius parce punctatum, pilosum, fronte modice convexa. Thorax latitudine dimidio brevior, coleopterorum basi paulo latior, leviter transversim convexus, dorso obsolete canaliculatus, parce disperseque punctatus, glaber, versus latera densius subtilius punctatus, fusco-pubescent, margine laterali longius densiusque nigro-piloso. Scutellum subtiliter punctulatum. Elytra subtiliter confertim punctata, fusco-pubescentia, tuberculis minutis sparsis parce obsita, tuberculo quoque seta reclinata nigra instructo, lateribus nigro-pilosa. Abdomen depressum, confertim subtiliter punctulatum, fusco-pubescent, segmento quinto apice sextoque toto piceo-rufis. Pedes dilute rufo-piceo, tibiis tarsisque nigricantibus. Alae fusco-hyalinae.

Obs. Errore quodam vel potius calami lapsu Dom. Nordmann pro patria huius insecti Sumatram affert.

## Scariphaeus Er.

*Antennae rectae, moniliformes, subcompressae.*

*Palpi articulo ultimo truncato.*

*Maxillae mala exteriore elongata.*

*Ligula apice emarginata.*

*Pedes intermediarii basi distantes.*

*Tarsi posteriores depressi.*

Corpus maioris magnitudinis, oblongum, alatum. Caput thorace paulo angustius, transversim suborbiculatum, basi profunde constrictum, collo quasi angusto brevique thoraci immisum, oculis haud prominulis. Labrum breve, transversum, bilobum, corneum, membranaceo-limbatum. Mandibulae parum falcatae, extus canaliculatae, intus medio subdentatae. Maxillae (t. IV. f. 18. b.) mala interiore coriacea, intus barbata, exteriore elongata, cornea, apice coriacea barbataque. Palpi maxillares breves, mala exteriore paulo longiores, articulo primo parvo angustiore, reliquis tribus crassiusculis, subaequalibus, quarto oblique truncato. Labium (t. IV. f. 18. a.) mento transversum, ligula brevi, leviter emarginata, membranacea, paraglossis membranaceis, intus dense ciliatis, ligulam superantibus. Palpi labiales filiformes, articulis sensim paulo longioribus, ultimo apice summo truncato. Antennae breviusculae, articulo primo leviter elongato, obconico, secundo tertioque subaequalibus, obconi-

cis, reliquis transversis, fere perfoliatis, ultimo apice fortiter emarginato. Thorax transversus, coleopterorum latitudine, basi lateribusque rotundatus, apice truncatus, angulis posterioribus nullis, anterioribus subrectis, deflexis, modice convexus, lateribus basique explanato-marginatus. Scutellum triangulare. Elytra thoracis fere latitudine, truncata, angulo exteriori oblique truncato. Abdomen apicem versus leniter angustatum. Pedes breves, crassi, coxis intermediis approximatis, femoribus subtus pro tibiis recipiendis profunde canaliculatis, tibiis omnibus externe spinulosis, tarsis brevibus, depressis, articulis arcte coactis, primo subelongato, (tars. postic. t. IV. f. 18. c.) anticis articulis 4 primis leviter dilatatis, coniunctis ovatis, subtus dense spongioso-tomentosis.

#### Observationes.

1. Habitu a genuino Staphylino parum discedit, quibus antennarum quoque structura magis appropinquatur, licet tarsorum, maxillarumque structura arcte Cordylaspidi et Haematodi iungitur, ab hoc autem femoribus basi approximatis, ab illo mandibulis medio dentatis praeter antennarum structuram distinguitur.

2. Temere Dom. Runde hoc insectum Quedio dilatato sub genere Velleio associat, cum ne antennae quidem serratae sint, aliis omnibus characteribus praetermissis, et equidem hanc ob rem ex descriptione citata aegre illud cognovissem, nisi typus a celeberrimo Prof. Germar amice mihi communicatus sit.

3. Sexus differentia ut in genuinis Staphylinis se habet.

\* 1. *S. luridipennis*: *Niger, nitidus, capite fortius punctato thoraceque cyaneis, elytris anoque rufis.* — Long. 7 lin.

*Velleius luridipennis* Runde *Brachelytr.* Hal. p. 1. not.

Habitat in Brasilia, Dom. Virmond.

Staph. caesareo brevior, latitudine autem ei fere aequalis, niger, nitidus. Antennae capite vix longiores, articulis secundo tertioque aequalibus, obconicis, quarto quintoque modice transversis, 6 — 10 his paulo latioribus et brevioribus, fortiter transversis, fere perfoliatis, modice compressis, ultimo paulo angustiore, oblique emarginato, articulis 1 — 5 nigris, reliquis obscure ferrugineis. Caput supra leviter convexum, parce fortius punctatum, cyaneum, nitidum, parce tenuiter pubescens. Thorax latitudine paulo brevior, suborbiculatus, apice truncatus, modice convexus, laevissimus, glaber, cyaneus, subnitidus. Scutellum confertissime subtiliter punctatum, nigrum, tenuiter fulvo-pubescens. Elytra thorace vix longiora, confertim subtiliter punctata, testaceo-rufa, nitida, pube depressa fulva dense vestita. Abdomen coleopteris paulo angustior, nigrum, nitidum, segmentis duobus ultimis rufis, supra glabrum et utrinque parce disperse punctatum, subtus crebrius punctatum, puncto quoque pilo brevi appresso instructo, segmentis anterioribus 4 nigro-, ultimis duobus rufo-pubescentibus. Pedes nigri.

Mas abdominis segmento sexto apice triangulariter inciso distinguitur.

#### Palaestrinus Er.

*Antennae rectae.*

*Palpi articulo ultimo apice truncato.*

*Maxillae mala exteriori elongata.*

*Pedes intermedii basi distantes.*

*Tarsi posteriores teretes.*



*Corpus elongatum, alatum. Caput thorace paulo latius, oblongum, supra convexum, basi profunde constrictum, oculis parvis, haud prominulis. Labrum transversum, bilobum, corneum, margine pilosum, absque limbo coriaceo vel membranaceo. Mandibulae validae, medio dentibus duobus fortibus armatae. Maxillae mala exteriori elongata, cornea, apice coriacea barbataque. Palpi maxillares breviusculi, articulo primo minuto, secundo tertioque obconicis, hoc paulo crassiore, ultimo praecedente breviori, at vix crassiore, fere semigloboso, apice truncato. Palpi labiales articulo ultimo praecedente longiore, cylindrico, apice truncato. Antennae filiformes, articulo ultimo apice emarginato et inferne acuminato. Thorax oblongus, convexus. Scutellum triangulare. Elytra apice truncata, angulo apicis exteriori rotundato. Abdomen longiusculum, subcylindricum, lateribus tenuiter marginatum. Pedes tenuiores, tibiis posterioribus spinulosis, anticis incrassatis, externe laevibus, interne tomentosis, tarsis posterioribus modice elongatis, articulis 4 primis longitudine decrescentibus, tarsis anticis minutis, articulis 4 primis leviter dilatatis, subtus longius pubescentibus.*

#### O b s e r v a t i o n e s.

1. Maxillarum mala exteriori elongata cum praecedentibus tribus generibus convenit, discedit autem tarsorum structura. Pedum anticorum palporumque maxillarum structura insigne videtur hoc genus, exemplo singulo viso haud accuratius mihi examinatum

2. Mas abdominis segmento inferiore sexto apice emarginato cognoscendus. Femina latet.

1. *P. Sykesii*: *Niger, capite thoraceque obscure aeneis, aureo-tomentosis, elytris punctato-rugosis, rufo-aureo-pubescentibus, abdomine fasciis duabus aureo-tomentosis, pedibus rufo-piceis. — Long. 7 lin.*

Habitat in India orientali, Mus. Dom. Hope.

Antennae capite sesqui longiores, nigrae, articulo tertio modice elongato, secundo sesqui longiore, 4—10 subaequalibus, leviter obconicis, crassitie paulo longioribus, ultimo apice oblique emarginato et inferne acuminato. Mandibulae nigrae. Caput thorace paulo latius, subovatum, supra valde convexum, creberrime punctatum vel punctato-subrugosum, nigro-aeneum, nitidulum, tomento aureo obtectum. Thorax coleopteris plus dimidio angustior, latitudine paulo longior, basin versus leviter angustatus, valde convexus, subtiliter punctato-rugosus, linea tenui media longitudinali laevi, obscure aeneus, nitidus, dense aureo-tomentosus. Elytra thorace paulo longiora, punctato-rugosa, obscure violacea, nitidula, pube longiore depressa rubro-aurea densius vestita. Abdomen nigrum, supra basi fasciisque duabus in segmenti quarti quintique basi sitis dense aureo-tomentosis, subtus minus dense aureo-villosum. Pectus nigrum, aureo-pubescentius. Pedes piceo-castanei.

Species mihi invisae, forte huius generis.

2. *P. (?) aureus*: *Capite, thorace elytrisque villosa-aureis, abdomine nigro, fasciis cinereis.*

*Staph. aureus* Fab. Mant. Ins. I. 219. 1. Ent. Syst. I. II. 519. 1. Syst. El. II. 589. 1. — Oliv. Ent. III. 42. 7. 1. t. 1. f. 2.

Habitat in Siam.

Statura Staph. hirti. Antennae pilosae, nigrae, articulo ultimo ovato, testaceo. Caput magnum, villosum, ferrugineo-aureum. Clypeus porrectus, rotundatus, ciliatus, integer. Thorax et elytra pilis brevibus dense villosa, ferrugineo-aurea. Pectus villosa-aureum. Abdomen atrum, fasciis duabus latis villosa-cinereis, nigro-punctatis. Pedes anteriores rufi, femoribus supra nigris.

## Staphylinus L.

*Er. Col. March. I. p. 430.*

*Antennae rectae.*

*Palpi maxillares filiformes.*

*Ligula apice emarginata.*

*Pedes intermedii basi distantes.*

*Tarsi posteriores teretes.*

Corpus maioris magnitudinis, elongatum, alatum. Caput suborbiculatum, vel subquadratum, thoraci plerumque magnitudine subaequale, exsertum, collo quasi angusto thoraci immissum. Labrum corneum, transversum, bilobum, membranaceo-coriaceo-ve-limbatum. Mandibulae plerumque sat fortes. Maxillae mala interiore trapezoidea, introrsum barbata, exteriori basi obconica cornea, apice coriacea barbataque, palpis longe brevioribus. Palpi maxillares filiformes, articulo primo parvo, secundo subelongato, ultimo apice subacuminato. Labium (t. IV. f. 14.) mento transversum, ligula brevi, medio sinuata, membranacea, paraglossis coriaceis, interne ciliatis, ligulam longe superantibus. Palpi labiales articulo ultimo plerumque longioribus, aut subacuminato aut apice truncato. Antennae ante mandibularum basin insertae, capite thoraceque plerumque paulo brevioribus, articulo primo leviter elongato. Thorax basi rotundatus, apice truncatus, angulis anterioribus plerumque rectis, subdeflexis. Scutellum rotundato-triangulari. Elytra apice aut rotundata aut introrsum oblique truncata. Abdomen plerumque parallelum. Pedes coxis intermediis magis minusve distantibus, tarsis posterioribus filiformibus, articulo primo subelongato, anticis plerumque articulis 4 primis dilatatis, patellam ovalem formantibus, subtus spongioso-tomentosis.

### Observationes.

1. Distinguitur hoc genus, speciebus locuplex, formas varias continens, a praecedentibus praecipue tarsis posterioribus haud complanatis, maxillarum mala exteriori elongata, a proxime sequenti imprimis pedibus intermediis magis minusve distantibus. Coxae intermediae nonnunquam quidem valde approximatae, acetabula autem semper interstitio disiuncta.

2. Mas abdominis segmento inferiore sexto apice emarginato vel inciso, nonnunquam quoque capite grandiore distinctus.

3. Victus in cadaveribus, in fimo, in sylvis sub musco. In fimo potius insectorum fimetariorum rapina degere quam fimo ipso delectari videntur. Staph. hirtum in stercore Onthophagum nuchicornem devorantem ipse observavi.

Distinxi in hoc genere Familias duodecim, quorum plures iam ab auctoribus variis pro generibus peculiaribus descriptae sunt, quae vero recipere dubitavi, cum characteres mihi leviores et minus constantes videantur. Nihilominus has familias plerasque naturales existimo, et notas earum distinctivas accuratius eruere conatus sum.

## Fam. I.

*Emus* Leach. *Curtis Britt. Ent. XII. t. 534. Steph. Illustr. Britt. Ent. V. p. 203*

*Antennae apicem versus subincrassatae, articulis 6 — 10 transversis, articulo ultimo integro.*

*Palpi omnes filiformes, maxillares articulo quarto tertio brevior.*

*Pedes breves, tibiis omnibus spinulosis, tarsis anticis dilatatis.*

Corpus subтус longius pubescens, supra dense villosum. Antennae breviores, articulo primo modice elongato, secundo tertioque obconicis, quarto quintoque subglobosis, 6 — 10 paulo latioribus, transversis, inter se aequalibus, ultimo breviter ovato, apice subcompresso et utrinque impresso, articulis 5 primis leviter puberulis, reliquis pube subtilissima cinerea dense obductis. Caput sat magnum, leviter deflexum, oculis minoribus parum convexus. Labrum limbo latiore membranaceo instructum. Mandibulae validae, apice leviter falcatae, acutae, medio fortiter dentatae. Palpi maxillares articulo secundo tertio paulo longiore, quarto tertio paulo brevior, acuminato, labiales articulis duobus primis aequalibus, ultimo paulo longiore, apice subacuminato. Thorax basin versus angustatus. Elytra angulo apicali exteriori oblique truncato. Abdomen coleopterorum latitudine, parallelum. Pedes breviusculi, validi, intermediis basi satis distantes, tibiis omnibus dense pubescentibus, crebre spinulosis, spinis apicalibus mediocribus, tarsis posterioribus breviusculis latiusculisque, articulo primo leviter elongato, sequentibus tribus sensim paulo minoribus, apice fere bifidis, tarsis anticis articulis 4 primis dilatatis, patellam ovatam subtus tomentoso-spongiosam formantibus.

Mas capite saepius maiore, abdominis segmento inferiore sexto apice leviter emarginato et pedum structura distinctus.

Obs. Affinis Fam. sequenti, distincta autem praeter corporis vestimentum tarsis brevibus, et antennarum articulis 6 — 10 aequalibus.

X \* 1. *St. hirtus*: *Niger, hirsutus, capite, thorace abdominisque apice flavo-villosis, elytris fascia postica cinerea.* — *Long. 7 — 11 lin.*

*Linn. Faun. Suec. n. 839. Syst. Nat. I. II. 589. 2. — Fab. Syst. Ent. 264. 1. Spec. Ins. I. 334. 1. Mant. Ins. I. 219. 2. Ent. Syst. I. II. 519. 2. Syst. El. II. 589. 2. — Sulz. Hist. Ins. p. 69. t. 7. f. 16. — Müll. Prodr. 97. 1091. — Goeze Ent. Beitr. I. 417. 1. — Fourcr. Ent. Paris. I. 165. 7. — De Vill. Ent. I. 409. 1. — Payk. Mon. Staph. 1. 1. Faun. Suec. III. 368. 1. — Schrank Faun. Boica. I. 639. 860. — Harrer Besch. d. Ins. I. 249. 410. — Rossi Faun. Etr. Mant. 95. 214. Ed. Hellw. I. 441. 214. — Walken. Faun. Paris. I. 268. 1. — Oliv. Ent. III. 42. 7. 2. t. 1. f. 6. — Panz. Faun. Germ. 4. 19. Ent. Germ. 351. 1. — Mart. Engl. Ent. t. 42. f. 40. — Grav. Micr. 159. 1. Mon. 125. 148. — Latr. Hist. nat. Crust. et Ins. IX. 292. 1. t. 79. f. 4. Gen. Crust. et Ins. I. 286. 1. — Marsh. Ent. Britt. 496. 1. — Don. Britt. Ins. XVI. t. 552. — Gyll. Ins. Suec. II. 281. 2. — Nordm. Symb. 38. 18. — Er. Col. March. I. 431. 1.*

*Emus hirtus* Mannerh. *Brachelytr. 20. 1. — Curt. Britt. Ent. XII. t. 534. — Boisd. et Lacord. Faun. Ent. Paris. I. 360. 2. — Runde Brachelytr. Hal. 2. 1.*

*Staph. Bombylius* De Geer *Ins. IV. 20. 5.*

*Le Staph. Bourdon* Geoffr. *Hist. Ins. I. 363. 7.*

*Staph. quintus* Schaeff. *Icon. I. t. 36. f. 6. Abhandl. I. t. 2. f. 12.*

*Bombylius corpore longissimo, cornibus quasi Scarabaeus armatus* *Petiver Op. I. t. 138. f. 20.*

Habitat in Europa.



Niger, parum nitidus, supra subtiliter punctatissimus. Antennae capite vix longiores, minus crassae, articulo tertio secundo paulo longiore, 6—10 praecedentibus crassioribus, transversis, ultimo apice utrinque impresso. Caput thorace vix latius, supra modice convexum, pilis densis flavo-aureis villosum, subtus glabrum. Thorax coleopteris paulo angustior, latitudine paulo brevior, basin versus leviter angustatus, basi angulisque posterioribus rotundatis, lateribus subrectis, apice truncatus, angulis anterioribus acutiusculis, parum convexus, pube erecta densa flavo-aurea uti caput vestitus, limbo posteriore libero, atro-pubescente. Scutellum nigro-tomentosum. Elytra thorace paulo longiora, basi nigro-pubescentia, apice latius cinereo-tomentosa, maculis punctisque nigris interdum immixtis vix variegata. Abdomen supra segmentis tribus primis nigro-pubescentibus, reliquis tribus tomento spisso flavo-aureo obductis, subtus violaceum, nitidum, segmento quarto macula utrinque laterali, quinto fascia integra aureo-villosis. Pedes validiusculi, nigri, nigro-pubescentes.

Mas trochanteribus posticis unco apice truncato bicuspique armatis, tibiis posticis leviter sinuatis insignis.

### Fam. II.

*Creophilus Kirby Mannerh. Brachel. p. 20. — Steph. Illustr. Britt. Ent. V. p. 202.*

*Antennae subclavatae, articulis 7—10 transversis, sensim crassioribus.*

*Palpi omnes filiformes, maxillares articulo quarto tertio brevior.*

*Pedes tibiis posterioribus spinulosis, tarsis anticis dilatatis.*

Caput et thorax glabra, scutellum, elytra et abdomen pubescentia vel tomentosa. Antennae capite vix vel parum longiores, articulo primo longiusculo, secundo tertioque obconicis, quarto, quinto sextoque lenticularibus, saepius transversis, septimo, octavo, nono et decimo his distinctius crassioribus, transversis, saepius sensim paulo crassioribus, ultimo praecedente paulo angustiore, breviusculo, obtuse acuminato, apice subcompresso et plerumque impresso, articulis 1—6 breviter pilosellis, 7—11 pube brevissima grisea dense obductis. Caput porrectum, oculis minutis, oblongis, parum prominulis. Labrum breve, limbo angustiore coriaceo fortiter barbato cinctum. Mandibulae validae, leviter falcatae, apice acutae, medio fortiter dentatae. Palpi maxillares articulo secundo tertioque subaequalibus, quarto minore, acuminato, labiales articulo secundo primo dimidio brevior, tertio longior, primo aequali, apice subacuminato. Elytra angulo apicali exteriori rotundato. Abdomen coleopterorum latitudine, fere parallelum. Pedes breviusculi, minus validi. Tibiae anticae interne, posteriores undique spinulosae, spinis apicalibus validioribus. Tarsi antici articulis 4 primis sensim minus dilatatis, posteriores graciliores, articulo primo leviter elongato, sequentibus tribus leviter decrescentibus.

Mas capite subquadrato, vel thoracis latitudine, vel saepius thorace latiore, mandibulis longioribus validioribusque, thorace basin versus magis angustato, angulis anterioribus acutioribus, abdominis segmento inferiore sexto apice triangulariter inciso.

Femina capite subtriangulari, thorace angustiore.

\* 2. *St. maxillosus*: Niger, nitidus, thoracis angulis posterioribus rotundatis, elytris fascia cinerea, abdomine subtus segmentis 4 vel 5 cinereo-tomentosis, pectore nigro-pubescente. — Long. 6 — 10 lin.

Linn. Faun. Suec. n. 891. Syst. Nat. I. II. 683. 3. — Ström Act. Nidros. III. 403. 25. — Fub. Syst. Ent. 265. 3. Spec. Ins. I. 334. 3. Mant. Ins. I. 220. 5. Ent. Syst. I. II. 521. 9. Syst. El. II. 592. 11. — Müll. Faun. Frid. 23. 222. — Goeze Ent. Beitr. I. 715. 3. — De Vill. Ent. I. 410. 3. — Payk. Mon. Staph. 5. 2. Faun. Suec. III. 369. 2. — Rossi Faun. Etr. I. 246. 607. Ed. Hellw. I. 304. 607. — Schrank Faun. Boic. I. 639. 861. — Cederh. Faun. Ingr. 117. 357. — Walken. Faun. Paris. I. 269. 4. — Martyn Engl. Ent. t. 29. f. 4. 5. — Panz. Ent. Germ. 353. 8. Faun. Germ. 27. 2. — Oliv. Ent. III. 42. 9. 5. t. 1. f. 5. — Grav. Micr. 2. 1. Mon. 126. 150. — Latr. Hist. nat. Crust. et Ins. IX. 293. 2. — Donov. Britt. Ins. III. t. 96. f. 3. — Marsh. Ent. Britt. 498. 5. — Gyll. Ins. Suec. II. 279. 1. — Zetterst. Faun. Lappon. I. 69. 7. Ins. Lapp. 60. 8. — Sahlb. Ins. Fenn. I. 308. 1. — Er. Col. March. I. 432. 2.

*Emus maxillosus* Boisd. et Lacord. Faun. Ent. Paris. I. 358. 1.

*Creophilus maxillosus* Mannerh. Brachelytr. 20. 1. — Runde Brachelytr. Hal. 1. 1. — Nordm. Symb. 21. 1.

*Staph. nebulosus* Fourc. Ent. Paris. 165. 5.

*Staph. anonymus* Sulz. Kennz. d. Ins. 17. 22. t. 7. f. 49.

*Staph. balteatus* De Geer Ins. IV. 18. 4. t. 1. f. 7. 8. — Müll. Prod. 98. 1114. — Harrer Besch. d. Ins. I. 251. 414.

*Staph. fasciatus* Fuesly Verz. Schweiz. Ins. 21. 410. — Goeze Ent. Beitr. I. 724. 4.

*Le Staph. nebuleux* Geoffr. Hist. Ins. I. 362. 5.

*Staph. tertius* Schaeff. Icon. I. 20. 1.

Habitat in Europa.

Corpus totum nigrum, capite thoraceque nitidis. Antennae subclavatae, articulis 2 et 3 aequalibus, obconicis, 4—6 sensim paulo brevioribus, 7—10 sensim crassioribus, transversis, ultimo apice compresso et utrinque impresso. Caput maris thorace maius, feminae eodem minus, subtilissime punctulatum, fronte media laevi, basi utrinque crebre punctatum et nigro-pilosellum. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus et basi rotundatus, apice truncatus, angulis anterioribus acutiusculis, posterioribus rotundatis, leviter convexus, laevis, glaberrimus, limbo angusto punctulato, margine anteriore utrinque nigro-griseoque-puberulo. Scutellum dense punctatum, griseo-pubescentis, apice macula nigro-tomentosa. Elytra punctata, dorso serie longitudinali punctorum maiusculorum notata, nigro-pubescentia, fascia lata inaequali nigro-punctata cinereo-tomentosa. Abdomen supra pube nigra cinereaque variegatum, subtus segmentis quatuor primis tomento spisso cinereo obtectis, secundo quartoque puncto utrinque laterali nigro, quinto sextoque pube nigra interdum maculatim cinerascens vestita. Pectus nigro-pubescentis. Pedes nigri.

\* 3. *St. arcticus*: Niger, nitidus, thoracis angulis posterioribus subrotundatis, elytris fascia cinerea, abdomine subtus segmento secundo tertioque cinereo-tomentosis, pectore nigro-pubescente. — Long. 9 lin.

Habitat in Kamtschatka, Dom. Chamisso.

Niger, capite thoraceque nitidis. Antennae subclavatae, articulis 2 et 3 aequalibus, obconicis, 4—6 sensim paulo brevioribus, 7—10 sensim crassioribus, transversis, ultimo apice compresso, utrinque impresso. Caput parcius subtiliter punctatum, fronte media laeviore, pone oculos utrinque

punctis nonnullis maioribus impressum, basi utrinque cinereo-villosulum. Thorax coleopteris paulo angustior, latitudine anteriore dimidio brevior, basi et lateribus modice rotundatis, his pone medium subsinuatis, apice truncatus, angulis posterioribus obtusissimis, subrotundatis, leviter convexus, laevissimus, limbo angusto punctulato, basi utrinque crebre punctato et cinereo-villoso. Scutellum nigro-tomentosum. Elytra thorace sesqui longiora, dense subtiliter punctata, nigra, nigro-pubescentia, fascia lata cinereo-pubescente. Abdomen supra nigro-pubescentius, segmento secundo tertioque utrinque cinereo-tomentosis, sexto pilis cinereis pubi nigrae immixtis canescente, subtus nigro-pubescentius, segmento secundo tertioque cinereo-tomentosis. Pedes nigri.

Obs. Intermedius videtur inter *St. maxillosum* et *villosum*, ab hoc pectore nigro-pubescente, capite subtilius punctato, antennis sensim crassioribus, thoracis angulis posterioribus minus rotundatis, ab illo thorace apice utrinque cinereo-villosulo, angulis posterioribus minus rotundatis, abdominis segmentis ventralibus duobus tantum cinereo-tomentosis distinguendus. Specimen singulum femineum a def. Chamisso allatum.

\* 4. *St. villosus*: *Niger, nitidus, thorace antice utrinque cinereo-villoso, angulis posterioribus obtusis, elytris fascia cinerea, subtus pectore abdominisque segmentis 4 primis cinereo-tomentosis.* — Long. 9 lin.

Grav. Micr. 160. 2. Mon. 126. 149.

*Creophilus villosus* Nordm. Symb. 21. 2.

Habitat in America septentrionali, sat frequens, in Mexico, Dom. Deppe, in Cuba, Dom. Otto.

Statura et summa affinitas *C. maxilloso*, at vere distinctus. Antennae subclavatae, articulo secundo tertioque aequalibus, obconicis, 4 — 6 sensim paulo crassioribus, 7 — 10 abrupte crassioribus, fortiter transversis, ultimo apice emarginato. Caput maris thoracis latitudine vel eodem paulo latius, feminae angustius, crebrius subtiliusque punctatum, fronte medio paulo laeviore, basi utrinque cinereo-villosulum. Thorax coleopteris paulo angustior, basin versus angustatus, latitudine antica tertia parte brevior, basi et lateribus rotundatis, his pone medium subsinuatis, apice truncatus, angulis posterioribus obtusis, subrotundatis, anterioribus maris acutiusculis, feminae obtusiusculis, leviter convexus, laevissimus, limbo angusto punctulato, apice utrinque cinereo-villoso. Scutellum dense punctatum, nigro-pubescentius, apice macula nigro-tomentosa. Elytra thorace sesqui longiora, punctata, nigro-pubescentia, fascia inaequali cinereo-tomentosa. Abdomen segmentis tribus primis cinereo-tomentosis, nigro-variegatis, reliquis nigro-pubescentibus, pilis paucis cinereis immixtis vix variegata, subtus segmento secundo, tertio quartoque tomento spisso cinereo obductis, quarto apice, quinto sextoque nigro-pubescentibus, immaculatis. Pectus dense cinereo-pubescentius. Pedes nigri.

Obs. *Creophilus fasciatus* Laporte Etud. Ent. p. III, „Maxilloso similis, at paulo latior, capite magis transverso, thorace antice magis dilatato, elytrorum fascia griseo-tomentosa angusta, abdominis fascia ventrali angustiore, nonnisi segmenta duo occupante” long. 7 lin. ex America septentrionali forte ad huius speciei marem referendus.



- \* 5. *St. cinerarius*: *Niger, nitidus, elytris fascia cinerea, subtus pectore abdominisque segmentis 4 primis cinereo-tomentosis, his utrinque nigro-punctatis.* — Long. 9 lin.

Habitat in Aegypto, Dom. Prof. Ehrenberg.

C. maxilloso simillimus, at pectore cinereo-pubescente distinctus. Antennae subclavatae, articulo secundo tertioque obconicis aequalibus, 4 — 6 sensim paulo brevioribus, 7 — 10 abrupte paulo crassioribus, transversis, ultimo apice compresso et utrinque impresso. Caput glabrum, omnium subtilissime obsoleteque punctulatum, supra oculos punctis nonnullis maiusculis notatum, basi utrinque crebre punctatum et teguiter cinereo-pubescent. Thorax coleopteris paulo angustior, basin versus angustatus, latitudine antica tertia parte brevior, lateribus subrectis, pone medium subsinuatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus maris acutis, feminae subrectis, laevissimus, glaber, limbo tenui, apice utrinque latiore, punctulato et cinereo-pubescente. Scutellum dense punctatum, cinereo-pubescent, macula minore apicali nigro-tomentosa. Elytra thorace tertia parte longiora, subtiliter punctata, nigro-pubescentia, fascia inaequali cinereo-tomentosa, nigro-punctata. Abdomen supra pubescent, segmento primo ultimisque duobus nigris, cinereo-variegatis, secundo, tertio quartoque nigro-variegatis, subtus segmento primo apice sequentibusque tribus totis tomento spisso cinereo vestitis, his singulis puncto laterali nigro, ultimis duobus nigris, cinereo-variegatis. Pectus dense cinereo-pubescent. Pedes nigri.

Obs. *St. villosus* proximus, at capite minus evidenter punctato, thoracis angulis posterioribus magis rotundatis et abdomine supra magis variegato, subtus segmento quarto aequae ac praecedentia tomentoso distinctus.

- \* 6. *St. incanus*: *Niger, nitidus, thoracis angulis posterioribus obtusis, elytris fascia cinerea, abdomine subtus segmentis 4 primis aequaliter albido-pubescentibus.* — Long. 6 — 10 lin.

Klug Col. Madag. 49. 40.

Habitat in Madagascar, Dom. Goudot.

Praecedentibus valde affinis, niger, nitidus. Antennae subclavatae, articulis 2 et 3 aequalibus, obconicis, 4 — 7 sensim paululum brevioribus, 7 — 10 sensim paulo crassioribus, fortiter transversis, ultimo subgloboso, apice compresso. Caput maris thorace latius, subquadratum, feminae thorace angustius, omnium subtilissime punctulatum, glabrum. Thorax coleopteris paulo angustior, basin versus angustatus, latitudine antica tertia parte brevior, basi modice rotundatus, apice truncatus et utrinque subsinuatus, lateribus vix rotundatis, pone medium subsinuatis, angulis anterioribus acutiusculis, posterioribus obtusis, leviter convexus, laevis, intra angulos posteriores foveola minuta impressus, glaber, margine impunctato, anteriore etiam non pubescente. Scutellum dense punctatum, nigro-pubescent. Elytra punctata, tenuiter nigro-pubescentia, paululum infra medium fascia angusta undulata cinereo-pubescente. Abdomen supra nigro-pubescent, segmentis secundo, tertio quartoque cinereo-variegatis, subtus segmento primo apice cinereo-ciliato, secundo tertio quartoque tomento denso caesio-albido obductis, quinto sextoque parcius nigro-pubescentibus, immixtis pilis nonnullis albidis. Pectus parce cinereo-pubescent. Pedes nigri.

Obs. St. maxilloso proximus, a quo praecipue thorace paulo longiore, limbo impunctato, apice haud pubescente, basi fortius rotundato, angulis posterioribus autem non rotundatis, elytrorum fascia angusta, et abdominis tomento ventrali subvirescente discedit. A cel. Klug femina descripta, marem Dom. Goudot nuper attulit.

- X \* 7. *St. variegatus*: *Niger, nitidus, abdomine elytrisque pubescentibus, cinereo-variegatis, his sutura apiceque fulvo-hirtis.* — Long.  $6\frac{1}{2}$  — 10 lin  
*Creophilus variegatus* Mannerh. *Brachelytr.* 20. 2. — Nordm. *Symb.* 21. 3.

Habitat in Brasilia.

Statura omnino C. maxilloso, niger, nitidus. Antennae clavatae, articulo tertio secundo paulo longiore, 4 — 6 sensim paulo brevioribus, 7 — 10 sensim at fortiter crassioribus, ultimo apice subcompresso, nigrae, articulis intermediis (4 — 7) testaceis. Caput maris thorace maius, omnium subtilissime punctulatum, feminae eodem minus, crebrius subtiliter punctatum, glabrum, basi utrinque cinereo-pubescent. Thorax coleopteris angustior, latitudine antica in inare paulo, in femina vix brevior, basin versus in mare fortius, in femina parum angustatus, basi rotundatus, lateribus subrectis, prope basin sinuatis, angulis posterioribus prominulis licet obtusis, anterioribus acutiusculis, supra parum convexus, laevissimus, apice utrinque subtiliter punctatus, intra angulum posteriorem utrinque foveolatus, glaber, apice utrinque cinereo-pubescent, basi tenuiter fulvo-ciliatus. Scutellum cinereo-pubescent, macula magna atro-holosericea. Elytra thorace sesqui longiora, vage punctata, nigro-pubescentia, cinereo-variegata, sutura apiceque fulvo-hirtis, margine laterali testaceo. Abdomen nigro-pubescent, cinereo-variegatum, subtus dense cinereo-pubescent. Pectus et pedes parce cinereo-pubescentia.

Obs. Insectum in Brasilia non infrequens pro Anglico descripsit Dom. Stephens nomine *Creophili ciliaris*. (Illustr. V. p. 202).

- X \* 8. *St. erythrocephalus*: *Niger, thorace nitido, angulis posterioribus rotundatis, elytris nigro-cyaneis, capite rufo, macula frontali oreque nigris.* — Long. 8 — 9 lin.

*Fab. Syst. Ent.* 265. 6. *Spec. Ins.* I. 335. 6. *Mant. Ins.* I. 220. 9. *Ent. Syst.* I. II. 523. 17. *Syst. El.* I. 593. 19. — *Oliv. Ent.* III. 42. 12. 9. t. 2. f. 9. — *Goeze Ent. Beitr.* I. 724. 6. — *Latr. Hist. nat. Crust. et Ins.* IX. 293. 3. — *Boisduv. Faun. Ent. Ocean.* 55. 2.

*Creophilus erythrocephalus* Nordm. *Symb.* 23. 4.

Habitat in Nova Hollandia.

Statura omnino C. maxilloso et minoribus eius individuis magnitudine aequalis. Antennae leviter clavatae, articulo tertio secundo paulo longiore, 4 — 6 leviter transversis, subaequalibus, 7 — 10 abrupte paulo crassioribus, transversis, ultimo apice subcompresso, nigrae. Os nigrum. Caput maris thoracis magnitudine, feminae eodem paulo minus, laevissimum, glaberrimum, fulvo-rufum, macula parva nigra in media fronte. Thorax coleopteris paulo angustior, latitudine antica dimidio brevior, lateribus et basi aequaliter rotundatus, illis pone medium subsinuatis, angulis posterioribus rotundatis, anterioribus subrectis, leviter convexus, laevissimus glaberrimusque, apice utrinque parce punctulato tenuiterque nigro-pubescente, niger, nitidus. Scu-

tellum dense punctatum, nigro-pubescent, macula apicali nigro-holosericea. Elytra confertim subtilius punctata, nigro-cyanea, parum nitida, nigro-pubescentia, callo humerali oblongo nigro, nitido, laevissimo, glabro. Abdomen, pectus et pedes nigra, opaca, nigro-pubescentia. Alae fusco-violaceae.

\* 9. *St. lanio*: *Niger, thorace nitido, angulis posterioribus obtusis, capite rufo, macula frontali oreque nigris.* — Long. 8 — 10 lin.

*Staph. oculus var. Grav. Mon. p. 127.*

Habitat in Terra Van-Diemenii.

Statura et summa affinitas praecedentis, at thoracis angulis posterioribus haud rotundatis et elytris non cyaneis constanter diversus. Niger. Antennae leviter clavatae, articulo secundo tertioque aequalibus, 4 — 6 sensim paulo brevioribus, leviter transversis, 7 — 10 leviter abrupte crassioribus, transversis, ultimo apice compresso et utrinque subimpresso, nigrae. Os nigrum. Caput maris thorace paulo latius, feminae eodem minus, parce subtilissimeque punctulatum, glabrum, rufum, nitidum, macula plerumque leviter transversa nigra in media fronte notatum. Thorax coleopteris paulo angustior, basin versus leviter angustatus, latitudine antica dimidio brevior, basi et lateribus leviter rotundatus, his pone medium subsinuatis, apice truncatus, angulis anterioribus rectis, posterioribus obtusis, nullo modo rotundatis, leviter convexus, apice utrinque subtiliter punctulatus et nigro-pubescent, ceterum laevissimus glaberrimusque, nitidus. Scutellum dense punctatum, nigro-pubescent. Elytra thorace sesqui longiora, confertim subtiliter punctata, nigro-pubescentia, opaca, callo humerali oblongo laevigato, glabro, nitido. Abdomen, pectus et pedes opaca, nigro-pubescentia.

\* 10. *St. oculus*: *Niger, nitidus, capite pone oculos utrinque rufo.* — Long. 9 lin.

*Fab. Syst. Ins. 265. 4. Spec. Ins. I. 335. 4. Mant. Ins. I. 220. 6. Ent. Syst. I. II. 521. 10. Syst. El. II. 592. 12. — Goeze Ent. Beitr. I. 724. 5. — Oliv. Ent. III. 42. 11. 8. t. 2. f. 19. — Grav. Mon. 126. 151. — Boisd. Faun. Ent. Ocean. 54. 1.*

*Creophilus oculus Nordm. Symb. 23. 5.*

Habitat in Nova Seelandia, Hübner.

Statura omnino et summa affinitas praecedentium. Corpus nigrum. Caput maris thorace paulo maius, vertice utrinque parcius punctato, ceterum laevissimum, nitidum, supra lateribus pone oculos rufis, subtus rufum, ore nigro. Thorax coleopteris parum angustior, latitudine antica dimidio brevior, basin versus leviter angustatus, lateribus et basi rotundatis, illis pone medium subsinuatis, angulis posterioribus obtusis, anterioribus subrotundatis, modice convexus, laevissimus, nitidus. Scutellum subtilius punctulato-rugulosum. Elytra thorace sesqui longiora, subtiliter punctato-rugulosa, subtiliter pubescentia, subopaca, callo humerali laevigato, glabro, nitido. Abdomen, pectus et pedes dense nigro-pubescentia, opaca.

Obs. Praecedenti affinis, praeter capitis colorem autem thorace convexiore, angulis anterioribus summo apice rotundatis, et elytris rugulosis distinctus. Antennae in specimine singulo, quod exstat in Museo, (masculo,) mutilatae.



## Fam. III.

*Brachydirus Nordm. Symb. p. 31.*

*Antennae breves, tenues, subclavatae, articulo ultimo integro, acuminato. Palpi omnes filiformes, maxillares articulo ultimo praecedente paulo longiore. Pedes graciles, tibiis setulosus, tarsi anticis fortiter dilatatis.*

Corpus totum tenuiter pubescens. Antennae capite vix vel parum longiores, articulo primo modice, secundo tertioque leviter elongatis, quarto quintoque obconicis, sequentibus transversis, sensim crassioribus, ultimo apice acuminato. Caput fortiter depressum, suborbiculatum, oculis maiusculis, modice prominulis. Labrum breve, transversum, limbo latiore membraneo cinctum. Mandibulae breviores, apice falcatae, tenues, acutae, basi dilatatae, muticae. Maxillarum malae latiusculae, membranaceae. Palpi tenues, maxillares articulo tertio secundo brevior, quarto iterum longiore, secundo aequali, apice acuminato, labiales articulo secundo primo paululum longiore, interne parcius setuloso, tertio paulo longiore, apice subacuminato. Thorax transversim subquadratus. Elytra apice cum angulo exteriori rotundata. Abdomen apicem versus sensim attenuatum. Pedes graciliores, at minus elongati, tibiis setis longioribus instructis, spinis apicalibus longioribus, gracilibus, tarsi anticis fortiter dilatatis, articulis 4 primis patellum ovalem, tibia duplo vel triplo latiore, subtus dense tomentoso-spongiosam formantibus, posterioribus gracilibus, articulis 4 primis longitudine decrescentibus, primo leviter elongato.

Mas nonnisi abdominis segmento inferiore sexto apice inciso distinctus.

Obs. Capite depresso, antennis brevibus subclavatis et tarsis anticis fortiter dilatatis facile distinguitur haec familia. Antennae omnibus testaceae vel flavae. Omnes Americam meridiionalem habitant.

\* 11. *St. xanthocerus*: *Niger, nitidus, antennis, ore anoque testaceis.* — *Long. 4 lin.*

*Brachydirus xanthocerus Nordm. Symb. 131. 1. t. 2. f. 3.*

Habitat in Brasilia.

*Niger, nitidus.* Antennae articulo secundo tertioque longitudine subaequalibus, quarto hoc paulo brevior, elongato-obconicis, 5 — 10 subobconicis, sensim leviter crassioribus, decimo leviter transverso, ultimo ovato, apice acuminato, totae flavo-testaceae. Os flavo-testaceum, mandibulis obscurioribus. Caput depressum, thoracis latitudine, creberrime fortius punctatum, tenuiter griseo-pubescens. Thorax coleopteris angustior, latitudine haud brevior, basi angulisque posterioribus rotundatis, lateribus subsinuatis, antice leviter emarginatus, angulis anterioribus acutiusculis, supra parum convexus, fortiter vage punctatus, linea media longitudinali laevi, parce griseo-pilosus. Scutellum punctulatum, griseo-pubescens. Elytra crebre minus profunde punctata, basi apiceque griseo-, medio nigro-pubescentia. Abdomen apicem versus sensim attenuatum, parce subtiliterque punctulatum, nigro-, segmento secundo tertioque griseo-pubescens, segmento quinto apice sextoque toto rufo-testaceis, stylis analibus nigris. Pedes postici nigri, anteriores picei, femoribus anterioribus apice tibiisque anticis basi testaceis,

pedes intermedii maris intus densius pubescentes, femoribus basi dilatatis. Alae fuscae.

Variat pedibus anterioribus totis femoribusque posticis testaceis.

- \* 12. *St. velitaris*: *Niger, nitidus, cinereo-pubescent, capite, thorace, ano pedibusque anterioribus testaceis.* — Long. 4 lin.

Habitat in Brasilia, Dom. Virmond.

Statura omnino praecedentis. Antennae articulo tertio secundo paulo longiore, quarto tertio dimidio brevior, his elongato-obconicis, quinto quarto paulo brevior, obconico, 6 — 10 transversis, sensim paulo crassioribus, ultimo breviter ovato, apice acuminato, totae flavae. Caput thoracis latitudine, depressum, creberrime fortiterque punctatum, basi summa laeve, testaceum, minus nitidum, parce pilosum. Thorax coleopteris paulo angustior, latitudine paululum brevior, basi angulisque posterioribus rotundatis, lateribus ante medium rotundatis, pone medium subsinuatis, antice late leviterque emarginatus, angulis anterioribus acutiusculis, leviter convexus, fortiter profundeque vage punctatus, linea media longitudinali laevi, rufo-testaceus, nitidus, margine nigro-pilosus. Scutellum testaceo-piceum, punctulatum, griseo-pubescent. Elytra thorace tertia parte longiora, crebre punctata, nigra, nitida, pube longiore depressa grisea vestita. Abdomen parce punctulatum, pilis longis depressis griseis vestitum, segmento secundo tertioque griseo-pubescentibus, nigrum, segmento quinto apice sextoque toto testaceis, stylis analibus brunneis. Pectus nigrum, mesosterno testaceo. Pedes anteriores cum coxis toti flavo-testacei, postici nigri, geniculis tarsisque piceis. Alae fuscae.

- \* 13. *St. testaceus*: *Testaceus, abdomine nigro, ano testaceo.* — Long. 5 lin.

*Fab. Syst. El. II. 595. 27.*

Habitat in Columbia, Dom. Moritz.

Antennae capite fere breviores, articulo secundo tertioque modice elongatis, subaequalibus, quarto quintoque subobconicis, hoc parum brevior, 6 — 10 brevibus, sensim crassioribus, transversis, ultimo crassitie haud longiore, apice fortiter acuminato, testaceae, apice fuscrescentes. Caput thoracis latitudine, depressum, anterieus confertissime, posterius crebre fortiterque punctatum, testaceum, parce pilosum. Thorax coleopteris angustior, latitudine paulo brevior, basi angulisque posterioribus rotundatis, lateribus angulisque anterioribus subrectis, antice truncatus, leviter convexus, fortius vage punctatus, linea media longitudinali laevi, testaceus, nitidus, parce fusco-testaceoque pilosus. Scutellum testaceum, apice punctatum flavoque pubescens. Elytra thorace paulo longiora, crebrius punctata, testacea, pube longiore depressa testacea densius vestita. Abdomen parce subtiliterque punctulatum, parce subtiliterque nigro-pubescent, segmentis secundo, tertio quartoque apice cinereo-pubescentibus, nigrum, segmento quinto apice sextoque toto testaceis, stylis analibus nigris, basi testaceis. Pectus testaceum. Pedes toti testacei. Alae fuscae.

- \* 14. *St. facialis*: *Testaceus, nitidus, capite antice abdomineque nigris, ano testaceo.* — Long. 3½ lin.

Habitat in Cayenna, Dom. Buquet.

Antennae vix capitis longitudine, articulo secundo tertioque leviter elongatis, longitudine aequalibus, hoc autem tenuiore, quarto quintoque subobconicis, hoc paulo brevior, 6 — 10 sensim crassioribus, transversis, ultimo conico, acuminato, crassitie paulo longiore, testaceae. Caput thorace paulo latius, depressum, supra parte anteriore maiore confertissime punctata, opaca, nigra, posteriore minore, parce punctata, nitida, testacea, subtus totum testaceum. Thorax coleopteris angustior, latitudine paulo brevior, basi rotundatus, apice truncatus, lateribus leviter sinuatus, angulis posterioribus obtusis, anterioribus acutiusculis, subdepressus, fortiter profundeque vage punctatus, linea media longitudinali laevi, rufo-testaceus, nitidus, parce nigropilosus. Scutellum rufo-testaceum, apice subtiliter punctulatum. Elytra thorace tertia parte longiora, fortius punctata, testacea, testaceo-pubescentia, parce nigro-pilosa. Abdomen parce subtiliterque punctulatum, nigro-pubesceus, segmento secundo, tertio quartoque, pube grisea immixta, nigrum, segmento quinto supra margine summo apicali sextoque toto testaceis. Pedes testacei. Pedes toti testacei.

#### Fam. IV.

*Antennae tenues, filiformes, ab articulo sexto inde subserratae, articulo ultimo integro, acuminato.*

*Palpi omnes elongati, filiformes, apice subacuminatae, maxillares articulo ultimo praecedente paulo longiore.*

*Pedes graciles, tibiis omnibus setulosis, tarsis anticis simplicibus.*

Corpus parce pubescens. Antennae longiores, graciles, basi satis distantes, articulis 5 primis elongatis, teretibus, articulo primo reliquis parum crassiore, tertio secundo sesqui longiore, quarto secundo aequali vel parum longiore, quinto hoc paulo brevior, 6—10 apice inferius magis prominulis ideoque leviter serratis, ultimo oblongo, apice acuminato, totae tenuiter pilosellae. Caput porrectum, suborbiculatum, oculis modice prominulis. Labrum latum, at breve, limbo latiusculo membraneo cinctum. Mandibulae longiores, falcatae, tenues, acutae, medio unidentatae. Maxillae malis parvis. Labium ligula brevi, medio sinuata. Palpi tenues, modice elongati, maxillares articulo quarto tertio sesqui longiore, labiales articulo primo brevior, secundo tertioque aequalibus. Thorax coleopteris angustior. Elytra angulo apicali exteriori oblique truncato. Abdomen apicem versus sensim leniter angustatum. Pedes modice elongati, graciles, intermediis basi sat late distantes, tibiis omnibus setulosis, spinis apicalibus subtilioribus, tarsis posterioribus articulo primo leviter elongato, sequentibus tribus leviter decrecentibus, quinto iterum elongato, anticis in utroque sexu simplicibus vel maris articulis tribus primis singulis leviter dilatatis.

Mas abdominis segmento inferiore sexto apice levius triangulariter inciso distinguendus.

Obs. Habitu quibusdam Philonthis non dissimiles sunt haec species, inter se concinne congruentes, at illi magis quam huic generi alienae videntur, licet corporis forma, palpis tenuibus elongatis et tarsis anticis simplicibus Ph. rufipenni, Baltimorensi etc, evidenter appropinquantur. Characteribus secundariis omnino inconstantibus, habitualibus saepe ambiguis, tanto magis characteres primarii observandi sunt. Ligula mihi emarginata videtur, qua a Philonthis discedunt haec insecta. Constituunt igitur in hoc genere familiam peculiarem, tarsis anticis simplicibus



eximiam, antennis subserratis sequenti propinquam, habitu graciliore, thorace parcius fortiusque punctato, ano vel toto abdomine testaceo, elytris cyaneis, fasciis e pube cinerea ornatis insignem. Species omnes hucusque cognitae Americam meridionalem habitant.

- \* 15. *St. comptus*: *Niger, ano testaceo, elytris cyaneis, fasciis duabus cinereo-pubescentibus.* — Long. 5 lin.

Habitat in Brasilia, Dom. Beske.

*Niger, nitidus.* Antennae capite sesqui longiores, articulo ultimo oblongo-ovato, apice sensim acuminato, totae nigrae. Caput thorace multo latius, suborbiculatum, supra parum convexum, parce disperseque grosse punctatum, parce nigro-pilosum. Thorax coleopteris longe angustior, basin versus sensim leviterque angustatus, latitudine apicis paulo longior, lateribus anterieus rotundatus, posterius leviter sinuatus, basi cum angulis posterioribus rotundatus, apice truncatus, transversim convexus, parce disperseque fortiter punctatus, linea longitudinali media laevi, parce nigro-pilosus. Scutellum laeve, opacum, apice cinereo-pubescentibus. Elytra thorace paulo longiora, dense punctata, cyanea, nitida, fasciis duabus cinereo-pubescentibus, altera pone basin, altera interne leviter arcuata ante apicem. Abdomen parcius punctulatum, nigrum, nitidum, parce nigro-pubescentibus, segmentis duobus apicalibus testaceis, testaceo-pubescentibus. Pectus lateribus cinereo-pubescentibus. Pedes graciles, setulosi, nigri.

Mas abdominis segmento inferiore quinto linea media transversali breviter dense longeque testaceo-pilosa instructo, tarsis anticis simplicibus.

- \* 16. *St. ornatus*: *Niger, nitidus, ano rufo-testaceo, elytris cyaneis, fasciis duabus, abdomine fascia singula cinereo-tomentosis.* — Long.  $5\frac{1}{2}$  lin.

Habitat in Cayenna, Dom. Buquet.

Statura omnino praecedentis, niger, nitidus. Antennae tenues, capite duplo fere longiores, nigrae, articulis ultimis duobus ferrugineis, ultimo oblongo, apice obtusius acuminato. Caput thorace latius, orbiculatum, subdepressum, utrinque fortiter punctatum, spatio medio longitudinali laevi, inter oculos transversim irregulariter punctato. Thorax coleopteris dimidio angustior, basin versus subangustatus, lateribus pone medium subsinuatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, latitudine apicis haud longior, modice transversim convexus, utrinque fortius crebriusque punctatus, spatio medio longitudinali laevi, lateribus subtiliter cinereo-pubescentibus. Scutellum opacum, dense punctulatum, apice cinereo-tomentosum. Elytra confertim subtilius punctata, cyanea, nitida, fasciis duabus dense cinereo-tomentosis, altera basali, altera apicali, angulis posterioribus exterioribus nigro-pilosellis. Abdomen subtiliter punctulatum, supra parce nigro-pubescentibus, segmento secundo fascia apicali cinereo-tomentosa, subtus segmentis secundo tertioque cinereo-pubescentibus, segmentis duobus apicalibus totis rufo-testaceis. Pectus dense cinereo-pubescentibus. Pedes graciles, setulosi, nigri, coxis femoribusque subtus cinereo-pubescentibus.

17. *St. fetialis*: *Niger, nitidus, coxis femoribusque anterioribus anoque rufo-testaceis, elytris cyaneis, fasciis duabus cinereo-pubescentibus.* — *Long.*  $4\frac{1}{3}$  lin.

Habitat in Mexico, collect. Dom. Chevrolat.

Habitu omnino praecedentium, at paulo minor, niger, nitidus. Antennae capite thoraceque paulo breviores, nigrae, articulo ultimo ferrugineo, apice abrupte acuminato. Mandibulae et palpi elongati, picei. Caput thorace paulo latius, suborbiculatum, fortiter punctatum, spatio angusto medio laeviore, parce griseo-pubescentis et nigro-pilosellum. Thorax coleopteris dimidio angustior, latitudine apicis haud brevior, basin versus subangustatus, lateribus anterieus leviter rotundatus, posterius subsinuatus, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus obtusiusculis, transversim convexus, sat crebre punctatus, linea media longitudinali laevi, parcius cinereo-nigroque puberulus. Scutellum nigrum, crebre punctulatum, cinereo-pubescentis. Elytra thorace paulo longiora, densius minus fortiter punctata, cyanea, nitida, tenuiter nigro-pubescentia, fasciis duabus e pube cinerea, altera basali, altera ante apicem. Abdomen parce obsolete punctatum, parce nigro-pubescentis, nitidum, segmentis ultimis duobus testaceis. Pedes nigri, anteriores coxis, trochanteribus femoribusque testaceo-rufis.

Mas tarsi anticis simplicibus.

- \* 18. *St. scenicus*: *Niger, abdomine rufo-testaceo, elytris cyaneis, fasciis duabus cinereo-pubescentibus.* — *Long.*  $5\frac{1}{2}$  lin.

Habitat in Cayenna, Dom. Buquet.

Statura praecedentium. Antennae capite duplo fere longiores, nigrae, articulo apicali obscure ferrugineo, apice obtuse acuminato. Caput thorace paulo latius, suborbiculatum, subdepressum, nigrum, nitidum, utrinque parce profundius punctatum, fronte medio laevi, antice parcius grosse punctata. Thorax coleopteris dimidio fere angustior, basin versus leviter angustatus, lateribus anterieus leviter rotundatus, posterius subsinuatus, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrectis, latitudine apicis paulo brevior, modice transversim convexus, niger, nitidus, utrinque obsoletius parce subseriatim punctatus, ad marginem lateralem cinereo-pubescentis. Scutellum nigrum, nitidum, apice punctulatum denseque cinereo-pubescentis. Elytra crebrius punctata, lacte cyanea, subnitida, fasciis duabus cinereo-pubescentibus, altera basali, altera paulo infra medium. Abdomen rufo-testaceum, nitidum, subtilius punctulatum, parcius testaceo-pubescentis. Pectus nigrum, cinereo-pubescentis. Pedes nigri, graciles, setulosi.

- \* 19. *St. venustus*: *Niger, nitidus, abdomine pedibusque rufo-testaceis, elytris cyaneis, fasciis duabus cinereo-pubescentibus.* — *Long.* 4 lin.

Habitat in Cayenna, Dom. Buquet.

Antennae capite duplo longiores, testaceae, articulis 7 — 9 fuscis, ultimo apice obtuse acuminato. Palpi pallide flavo-testacei. Caput thorace vix latius, suborbiculatum, subdepressum, nigrum, nitidum, utrinque fortiter punctatum, fronte medio laevi, antice parce grosse punctata. Thorax coleopteris vix dimidio angustior, basin versus haud angustatus, lateribus an-

tice leviter rotundatus, postice subsinuatus, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus obtusiusculis, latitudine paululum brevior, modice transversim convexus, niger, nitidus, utrinque parce subseriatim punctatus. Scutellum nigrum, dense punctulatum, dense cinereo-pubescentibus. Elytra crebrius punctata, coerulea, fasciis duabus cinereo-pubescentibus, altera basali, altera paulo infra medium. Abdomen testaceo-rufum, summa basi nigrum, laeve, parce ferrugineo-pubescentibus. Pectus nigrum. Pedes graciles, setulosi, testacei, coxis omnibus nigris.

Mas abdominis segmento superiore sexto apice trisinuato, sinu intermedio maiore, medio denticulo instructo, septimo elongato, fisso, portione utraque apice truncata, intus spinula, lateribus processu deflexo, interne spinuloso terminato, segmento inferiore sexto longitudinaliter bi-impresso, apice utrinque sinuato, medio profunde inciso, tarsis anticis articulis tribus primis leviter dilatatis, singulis oblongis.

Femina abdomine integro, tarsis anticis simplicibus.

### Fam. V.

*Leistotrophus* Perty *Delect. An. art. p. 30.*

*Schizochilus* Gray *An. Kingd. XIV. p. 310.*

*Antennae tenues, articulis basalibus valde elongatis, apice leviter incrassatae, ab articulo sexto inde subserratae, articulo ultimo apice emarginato.*

*Palpi omnes filiformes, maxillares articulo ultimo praecedente longe breviores.*

*Pedes elongati, tibiis infra medium pubescentibus, tarsis anticis in utroque sexu leviter dilatatis.*

Corpus tomento brevi densius obductum. Caput magnum, nutans, fronte verticali, planiuscula, oculis mediocribus, convexus. Antennae ab oculis distantes, tenues, articulis 3 primis teretibus, primo sequentibus duobus coniunctis paulo longiore, secundo tertio dimidio prope brevior, quarto quintoque leviter elongato-obconicis, hoc paulo brevior, 6 — 10 sensim paulo brevioribus et crassioribus, subcompressis, interne subserratis, ultimo ovato, acuminato. Labrum magnum, latum, distinctius membraneo-limbatum. Mandibulae falcatae, tenues, acutae, medio dente bicuspi armatae. Palpi maxillares articulo tertio secundo minore, quarto minuto, tenui, acuminato, labiales articulo secundo maiore, interne pilis rigidis ciliato, tertio minuto, tenui, acuminato. Thorax parvus, coleopteris multo angustior. Abdomen coleopteris angustius, apicem versus sensim angustatum. Pectus tumens. Coxae intermediae interstitio lato distantes. Pedes elongati, graciliores, femoribus anticis basi leviter incrassatis, tibiis omnibus inermibus, a medio ad apicem usque densius pubescentibus vel breviter barbatis, spinis apicalibus subtilioribus, tarsis gracilibus, posterioribus articulis 4 primis decreascentibus, primo leviter elongato, antici articulis 4 primis in utroque sexu singulis leviter dilatatis, inter se aequalibus, triangularibus, subtus apice membraneo-lobatis.

Mas capite maiore, mandibulis valde elongatis, coxis intermediis dentatis, tibiis intermediis leviter flexuosis, abdominis segmento inferiore sexto apice leviter emarginato distinctus.

Obs. Insectum unicum continet haec familia, plurimis notis admodum sin-



gulare, ut genus proprium et eximium constituere habeatur. Accuratius inspectum et examinatum minus ab hoc genere discedere videtur, et praecipue summa affinitate cum Fam. proxime sequente coniungitur, neque magis ab ea distat, quam haec a ceteris huius generis familiis discedit. Corporis indumento etiam cum illa omnino convenit. Antennae tenuiores, apicem versus subserratae in familia etiam proxime praecedente observantur.

\* 20. *St. versicolor*: *Niger, tomento fusco nebulosus, elytris inaequalibus, extrorsum subcostatis, pedibus testaceo-annulatis.* — Long. 7 — 9 lin.

*Grav. Mon.* 119. 138. — *Encycl. Meth. X.* 477. 1. — *Nordm. Symb.* 27. 1.

*Leistotrophus Gravenhorsti* *Perty Delect. An. art. p.* 30. t. 7. f. 1. — *Laport. Etud. Ent. p.* 112.

*Schizochilus Brasiliensis* *Gray An. Kingd. XIV. p.* 304. t. 32. f. 3.

*Staph. infuscatus* *Nordm. Symb.* 29. 2.

Habitat in America meridionali: in Parà Brasiliae, in silvis humidis in fructibus putrescentibus legit Sieber, prope Rio Janeiro, Dom. v. Olfers, in Surinamo, Dom. Cordua, in Columbia, Dom. Moritz, in Mexico, Dom. Deppe.

Antennae capite sesqui fere longiores, articulis 5 primis tenuibus elongatis, sequentibus leviter incrassatis, introrsum subserratis, testaceae, ante apicem fusciscentes. Palpi nigro-picei. Corpus nigro-obscure. Caput subdepressum, nutans, maris permagnum, mandibulis longis, falcatis, supra parce punctulatum, subtiliter carinatum, tomentosum, subnebulosum, subtus atrum, glabrum, nitidum. Thorax parvus, capite multo angustior, latitudine antica brevior, basin versus valde angustatus, basi angulisque posterioribus aequaliter rotundatis, angulis anterioribus acutis, lateribus subcompressis, leviter convexus, olivaceo-fuscoque nebuloso-tomentosus. Scutellum magnum, nebuloso-tomentosum. Elytra ampla, nigro-subaenea, margine humerali testaceo, inaequalia, versus latera subcarinata, tomento olivaceo, nigro fuscoque variegata. Abdomen angustatum, supra tomento fusco, olivaceo nigroque variegato, apice tomento olivaceo-subaureo aequali vestitum, subtus nigrum, tomento argenteo micante versicolor. Pectus amplum, utrinque tuberosum. Pedes elongati, nigri, antici femoribus apice, tibiis tarsisque testaceis, posteriores femoribus annulo basali apiceque, tibiis basi tarsisque testaceis. Alae fulvo-hyalinae.

Obs. *Staphyl. infuscatus* Nordm. Mexicanus, specimen femineum, nonnisi statura minore et alis fuscis apice late dilutius flavescenti-hyalinis discrepat, aegre specie discernendus.

## Fam. VI.

*Antennae filiformes, articulo ultimo integro, acuminato.*

*Palpi omnes filiformes, maxillares articulo ultimo praecedente minore.*

*Pedes tibiis posterioribus tenuiter spinulosis, tarsis anticis dilatatis.*

Corpus tomento brevi dense obductum. Antennae mediocres, tenuiores, apicem versus haud incrassatae, articulo primo modice elongato, sequentibus duobus coniunctis vix longitudine aequali, secundo tertio brevioribus, sequentibus sensim paulo brevioribus, penultimis leviter transversis, ultimo apice oblique emarginato, abrupte acuminato. Caput leviter deflexum, thoracis latitudine vel paululum latius, subquadratum, fronte planiuscula, oculis me-

diocribus modice convexis. Labrum sat magnum, late membraneo-limbatum. Mandibulae longiores, falcatae, acutae, medio dente bicuspi armatae. Palpi maxillares articulo tertio secundo longitudine aequali, quarto hoc multo minore, tenuiore, acuminato. Palpi labiales articulis duobus primis subaequalibus, tertio secundo paulo tenuiore, at vix brevior, acuminato. Abdomen coleopteris parum vel vix angustius, apicem versus subangustatum. Pedes mediocres, graciliores, femoribus basi subincrassatis, tibiis posterioribus parcius tenuiterque spinulosis, intermediis leviter arcuatis, spinis apicalibus longioribus, tarsis posterioribus articulis 4 primis sensim decreascentibus, primo subelongato, anticis articulis 4 primis modice dilatatis, tibia paulo latioribus, singulis transversis, latitudine decreascentibus, subtus dense tomentosis.

Mas nonnisi abdominis segmento inferiore sexto apice emarginato distinctus.

Obs. Ingeniose hanc Fam. cum Emo coniunxit Com. Mannerheim, et iis magis quam Fam. VIII. propinqua videtur, licet pedibus gracilioribus, tibiis anticis haud spinulosis et antennarum structura satis distinguitur. Propius accedit ad proxime praecedentem Fam., St. murinus tamen minus quam St. nebulosus, et maxime St. cingulatus, antennarum articulis inferioribus magis elongatis et tibiis posterioribus minus spinulosis.

\* 21. *St. cingulatus*: Niger, tomento fusco-nebuloso vestitus, pectore anoque aurichalceo-tomentosis, palpis pedibusque testaceis, femoribus subtus nigris. — Long. 7–8 lin.

Grav. Micr. 166. 14. Mon. 121. 140. — Nordm. Symb. 31. 5.

*Emus speciosus* Mannerh. *Brachelytr.* 21. 3. — Nordm. Symb. 31. 4.

Habitat in America septentrionali.

Statura et summa affinitis St. nebulosi. Niger, capite, thorace elytrisque tomento depresso cinereo-fuscoque nebuloso obductis. Antennae capite plus sesqui longiores, articulo tertio secundo duplo prope longiore, sequentibus duobus leviter elongatis, teretibus, sexto subobconico, crassitie longiore, 7–10, obconicis, crassitie haud brevioribus, testaceae, extrorsum fuscescentes. Palpi testacei. Caput thorace vix latius, suborbiculatum. Thorax coleopteris angustior, latitudine antica fere longior, basin versus sensim angustatus, lateribus subsinuatus, basi angulisque posterioribus rotundatis, apice truncatus, angulis anterioribus antrorsum prominulis, acutis, parum convexus. Scutellum macula gemina nigro-tomentosa notatum. Elytra thorace parum longiora, margine laterali inflexo testaceo. Abdomen supra segmentis quatuor primis tomento fusco, nigro-nebuloso, quarto aurichalceo-versicolore, quinto sextoque tomento denso depresso aequali aurichalceo nitido vestitis, subtus segmento singulo basi aurichalceo-pubescente. Pectus dense aurichalceotomentosum, nitidum. Pedes testacei, femoribus subtus, coxis trochanteribusque nigris.

Obs. Errore quodam apud Nordmannum St. spinosus nominatus.

X \* 22. *St. nebulosus*: Niger, tomento fusco-nebuloso vestitus, palpis pedibusque testaceis, his basi nigris. — Long. 6–8 lin.

Fab. Ent. Syst. I. II. 520. 3. Syst. El. II. 590. 3. — Payk. Faun. Suec. III. 375. 11. — Panz. Ent. Germ. 352. 2. — Grav. Micr. 13. 14. Mon. 121. 139. — Latr. Hist. nat. Crust. et Ins. IX. 295. 5. — Gyll. Ins. Suec. II. 282. 3. — Zetterst. Ins. Lappon. 59. 1. — Nordm. Symb. 30. 3. — Er. Col. March. I. 432. 3.

- Emus nebulosus* Mannerh. *Brachelytr.* 20. 2. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 361. 3. — *Runde Brachelytr. Hal.* 2. 2.  
*Staph. tessellatus* Fourc. *Ent. Par.* I. 165. 6.  
*Staph. hybridus* Marsh. *Ent. Britt.* 500. 9.  
*Staph. murinus* Harrer *Beschr. d. Ins.* I. 252. 415. — *Panz. Faun. Germ.* 66. 16.  
*Staph. villosus* var. *maior* Degeer *Ins.* IV. 12. 1. t. 16. f. 1-6.  
*Le Staph. velouté* Geoffr. *Hist. Ins.* I. 362. 6.  
*Staph. secundus* Schaeff. *Icon.* I. t. 4. f. 11.

Habitat in Europa.

Antennae capite plus sesqui longiores, articulo tertio secundo sesqui longiore, sequentibus tribus crassitie longioribus, testaceae, extrorsum fuscae. Palpi testacei. Mandibulae rufo-piceae. Caput thorace paulo latius, suborbiculatum. Thorax coleopteris angustior, basin versus angustatus, latitudine antica haud brevior, lateribus subsinuatus, basi angulisque posterioribus rotundatis, angulis anterioribus acutiusculis, parum convexus. Elytra thorace parum longiora. Caput, thorax et elytra tomento denso depresso cinereo, fusco-subnebuloso obducta. Scutellum macula magna nigro-tomentosa tectum. Abdomen supra tomento fusco, fulvo-nitente, maculatim nigro-variegato, apicem versus rariore, vestitum, subtus nigrum, segmento singulo fascia basali argenteo-pubescente versicolore. Pectus fusco-cinereo-pubescentibus. Pedes coxis, trochanteribus et femoribus nigris, his apice, tibiis tarsisque testaceis.

\*23. *St. marginalis*: Nigro-subaeneus, tomento depresso fusco-nebuloso, vestitus, palpis nigris, pedibus testaceis, basi nigris. — Long. 6 lin.

*Emus marginalis* Gené *Act. Reg. Scienc. Acad. Taurin.* XXXIX. 171. 10. t. 1. f. 4.

Habitat in Sardinia, Prof. Gené, in Lusitania, Com. Hoffmannsegg.

Statura et summa affinitas omnino *St. murini*, at vere distinctus. Antennae articulo tertio secundo sesqui longiore, sequentibus crassitie haud longioribus, testaceae, apice fusciscentes. Palpi nigri. Caput, thorax et elytra viridi-aenea, vix nitida, confertissime punctata, tomento depresso fulvo, fusco cinereoque maculatim variegatis, elytris margine laterali inflexo summoque apicali testaceis. Caput thoracis latitudine, subdepressum. Thorax coleopteris paulo angustior, latitudine antica paulo brevior, basin versus angustatus, apice leviter emarginatus, lateribus subrectis, angulis posterioribus obtusissimis, fere rotundatis, anterioribus acutis, parum convexus. Scutellum macula magna nigro-tomentosa. Elytra thorace paululum longiora. Abdomen supra segmentis anterioribus fusco-tomentosis, nigro-maculosis, ultimis duobus, ut subtus omnibus, basi cinereo-pubescentibus. Pectus dense aureo-pubescentibus. Pedes nigri, femoribus apice, tibiis tarsisque testaceis.

Obs. Intermedius quasi inter *St. nebulosum* et *murinum*, huic statura et corporis forma, illi antennarum pedumque colore simillimus, ab utroque elytrorum margine laterali inflexo testaceo distinctus. Pectus confertius quam in *St. murino* punctatum.

\*24. *St. murinus*: Niger, tomento depresso fusco-nebuloso vestitus, palpis pedibusque nigris. — Long. 6 lin.

*Linn. Faun. Suec. n.* 840. *Syst. Nat.* I. II. 683. 2. — *Fab. Syst. Ent.* 265. 2. *Spec. Ins.* I. 334. 2. *Mant. Ins.* I. 219. 3. *Ent. Syst.* I. II. 520. 4. *Syst. El.* II. 590.



4. — Goeze *Ent. Beitr.* I. 714. 2. — De Vill. *Ent.* I. 409. 2. — Payk. *Mon. Staph.* 14. 8. *Faun. Suec.* III. 376. 12. — Panz. *Ent. Germ.* 352. 3. — Ceder. *Faun. Ingr.* 116. 354. — Walken. *Faun. Paris.* I. 269. 2. — Oliv. *Ent.* III. 42. 15. 14. t. 6. f. 51. — Marsh. *Ent. Britt.* 499. 8. — Grav. *Micr.* 14. 15. *Mon.* 122. 142. — Latr. *Hist. nat. Crust. et Ins.* IX. 294. 4. — Gyll. *Ins. Suec.* II. 283. 4. — Zetterst. *Faun. Lappon.* I. 67. 1. *Ins. Lappon.* 59. 2. — Sahlb. *Ins. Fenn.* I. 308. 2. — Nordm. *Symb.* 31. 10. — Er. *Col. March.* I. 433. 4.

*Emus murinus* Mannerh. *Brachelytr.* 21. 7. — Boisd. et Lacord. *Faun. Ent. Paris.* I. 363. 6. — Runde *Brachelytr.* Hal. 2. 4.

*Staph. villosus* var. *minor* Degeer *Ins.* IV. 12. 1.

*Staph. murinus* Scop. *Ent. Carn.* 100. 302. — Schrank *Enum. Ins. Aust.* 229. 433. Müll. *Faun. Frid.* 223. 21. *Prod.* 98. 1115. — Preysl. *Böhm. Ins.* I. 18. 12. — Rossi *Faun. Etr.* I. 247. 608. *Ed. Helwe.* I. 305. 608. (*forte*)

Habitat in Europa.

Niger, capite, thorace elytrisque tomento fusco, fulvo cinereoque maculatim variegato subnebulosis. Antennae capite sesqui longiores, articulo tertio secundo sesqui longiore, sequentibus duobus crassitie paulo longioribus, testaceae, apicem versus fusciscentes. Palpi nigri. Caput thoracis latitudine, transversum, subdepressum. Thorax coleopteris paulo angustior. basin versus sub-angustatus latitudine antica paulo brevior, apice vix emarginatus, lateribus rectis, angulis posterioribus obtusissimis, fere rotundatis, angulis anterioribus acutis, parum convexus. Scutellum macula gemina nigro-pubescente. Elytra thorace vix longiora. Abdomen supra segmentis anterioribus fusco-tomentosis, nigro-maculosis, ultimis duobus ut subtus omnibus basi fascia cinerea-pubescente. Pectus tenuiter cinereo-pubescent. Pedes nigri.

25. *St. leucomus*: *Nigro-aeneus, tomento brevi olivaceo-obtuctus, albo-subnebulosus, pectore elytrisque albido-tomentosis, his basi olivaceo-nebulosis.* — Long. 6 lin.

Habitat in Nepal, Mus. Dom. Hope.

Corpus nigro-aeneum, tomento denso brevi obtectum. Antennae capite plus sesqui longiores, articulis primis elongatis, tertio secundo sesqui longiore, 4—6 teretibus, modice elongatis, 7—10 obconicis, crassitie haud brevioribus, ultimo apice sat fortiter emarginato, nigrae, articulo primo rufopiceo. Palpi picei. Caput thorace paulo latius, subtiliter punctatissimum, olivaceo-subtiliter tomentosum, albo-subnebulosum. Thorax coleopteris duplo fere angustior, latitudine antica paulo longior, basin versus angustatus, lateribus subrectis, basi leviter rotundatus, apice truncatus, angulis omnibus subrotundatis, fortiter transversim convexus, subtiliter punctatissimus, dense olivaceo-pubescent, albo-subvariegatus, carinula media tenui laevi medio late interrupta. Scutellum olivaceo-pubescent, macula atro-tomentosa. Elytra thorace tertia parte longiora, albo-tomentosa, subnebulosa, circa scutellum late fusco-olivaceo-pubescentia. Abdomen coleopteris angustius, olivaceo-pubescent, supra tomento atro bifarium submaculato, segmento ultimo albo-villoso, subtus segmentis 4 primis sextoque albido-tomentosis, quinto fusco-pubescente. Pectus totum dense albido-tomentosum. Pedes modice elongati, nigri, albido-pubescentes, tibiis posterioribus parce spinulosis, infra medium densius pubescentibus, anticis apicem versus sensim subincrassatis, inermibus, castaneis.

Obs. A praecedentibus notis nonnullis discrepat, praecipue fronte thoraceque convexis, hoc prope subcylindrico, angulis anterioribus obtusis, tibiis anticis dilatatis, et tibiis posterioribus apice densius pubescentibus, antennarum articulis 6 primis elongatis, teretibus, nec non abdomine angustiore ambiguus quasi inter hanc et praecedentem familiam videtur, at tibiis posterioribus licet parce spinulosis, tarsorum anticorum et oris structura propius ad genuinas huius familiae species accedit.

Species mihi invisae, forte huius familiae.

26. *St. Sibiricus*: *Niger, capite thoraceque fulvo-, abdomine albedo-tomentosis, elytris fuscis, fascia obliqua albida.* — Long. 6 lin.

*Gehl. Ledebour. It. App. III. 71. 16.*

Habitat prope Barnaul in Sibiria.

Opacus, tomento brevi tectus, adspersus pilis longis nigris. Caput orbiculare, nigrum, subtilissime alutaceum. Antennae capite longiores, griseo-tomentosae. Thorax longitudine capiteque paulo angustior, antice truncatus, lateribus rectus, postice angustior et rotundatus, supra tenuissime alutaceus, antice convexus, ultra medium carinulatus. Scutellum nigro-tomentosum. Elytra thorace latiora, nec longiora, apice truncata, supra fusco-tomentosa, subtilissime alutacea, punctis sparsis maioribus impressis, fascia lata, a parte marginis posteriore ad apicem sutura descendente, albedo-tomentosa, hinc inde fusco-punctata. Pygidium tomento albedo, fulvo nigroque variegatum. Pectus subtus fulvo-, abdomen albedo-tomentosum. Pedes nigro-fulvoque tomentosissimi. (Gehl.)

Obs. Praecedenti forte affinis.

27. *St. inauratus*: *Pubescens, niger, angustior, capite, thorace elytrisque e fusco et aureo nebulosis, thorace postice haud carinato.*

*Emus inauratus Mannerh. Brachelytr. 21. 7.*

Habitat in Dauria.

St. murino angustior, magis linearis, thorace postice haud carinato et pube aurea micante in capite, thorace et elytris conspersus. (Mannerh.)

### Fam. VII.

*Antennae filiformes, articulo ultimo integro, apice acuminato.*

*Palpi omnes filiformes, maxillares articulo quarto tertio aequali.*

*Pedes breviusculi, tibiis posterioribus parce tenuiterque spinulosis, tarsis anticis fortius dilatatis.*

Corpus tenuiter subtiliterque pubescens, capite thoraceque subglabris. Antennae breviores, apicem versus haud incrassatae, articulo primo sequentibus ambobus paulo breviori, vix clavato, secundo tertioque teretibus, hoc longiore, sequentibus sensim paulo brevioribus, magis minusve obconicis, ultimo ovato, apice acuminato. Caput porrectum, plerumque subquadratum, depressum, fronte plana, oculis mediocribus, parum prominulis. Labrum minutum, transversum, minus late coriaceo-limbatum, limbo barbato. Mandibulae breviores, apice falcatae, acutae, medio dente valido munitae. Palpi maxillares articulo secundo paulo longiore, tertio quartoque aequalibus, hoc acuminato, labiales articulis tribus sensim longioribus, tertio cylindrico, apice subacuminato. Thorax coleopteris angustior. Elytra angulo apicali exteriore rotundato. Abdomen apicem versus parum angustatum. Pedes bre-

viusculi, validiusculi, femoribus anticis basin versus modice dilatatis, tibiis omnibus dense pubescentibus, posterioribus parce spinulosis, spinis apicalibus subtilioribus, tarsis posterioribus articulis tribus intermediis subaequalibus, primo leviter elongato, anticis in utroque sexu articulis quatuor primis fortiter dilatatis, patellam ovatam vel suborbiculatam, subtus dense spongioso-tomentosam formantibus.

Mas abdominis segmento inferiore sexto apice emarginato distinctus.

Obs. Habitus fere Philonthi, capite thoraceque prope glabris, reliquo corpore subtiliter pubescente, thoraceque medio nonnunquam biserialiter punctato, eidem similis, at characteris omnino huius generis, et capite depresso subquadrato ad Fam. praecedentem accedit. A Fam. sequente praecipue antennarum articulo, ultimo apice acuminato differt.

\* 28. *St. sapphirinus*: *Niger, nitidus, abdomine segmentis ultimis duobus rufis, elytris cyaneis, antennis testaceis.* — Long. 6 lin.

Habitat in Parà interiore, Sieber, in Columbia, Dom. Moritz.

*Niger, nitidus.* Antennae capite sesqui longiores, articulo tertio secundo duplo prope longiore, 4 — 10 sensim paulo brevioribus, penultimo etiam haud transverso, ultimo oblongo-ovato, apice valde acuminato, totae testaceae. Palpi testacei. Labrum piceum. Caput thoracis latitudine, subquadratum, depressum, subtilius crebriusque punctatum, linea media longitudinali laeviore, parce nigro-pubescentis. Thorax coleopteris paulo angustior, latitudine haud brevior, basi angulisque posterioribus rotundatus, lateribus subrectis, apice truncatus, angulis anterioribus rectis, leviter convexus, utrinque crebrius subtiliusque punctatus, linea media longitudinali laevi, parce nigro-pubescentis. Scutellum dense punctatum, nigro-pubescentis. Elytra thorace parum longiora, densius punctata, nigro-pubescentia, cyanea, nitida. Abdomen densius punctulatum, nigro-pubescentis, segmentis ultimis duobus totis testaceo-rufis, rufo-pubescentibus. Pedes nigri, dense nigro-pubescentes, tarsis anticis obscure ferrugineis. Alae fuscae.

Mas abdominis segmento inferiore quinto lineola brevi transversa dense longeque fulvo-pilosa notato, sexto apice profunde inciso.

\* 29. *St. calidus*: *Niger, nitidus, abdomine segmentis ultimis duobus rufis, elytris virescenti-coeruleis, labro antennisque testaceis.* — Long. 6 lin.

Habitat in Columbia, Dom. Moritz.

Statura praecedentis, at paulo gracilior, niger, nitidus. Antennae capite plus sesqui longiores, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus, penultimis vix transversis, ultimo ovato, apice acuminato, totae dilute testaceae. Maxillae rufae. Palpi testacei. Mandibulae piceae. Labrum testaceum. Caput thoracis prope longitudine, subquadratum, depressum, crebre fortius punctatum, fronte medio longitudinaliter laevi, antice autem sat crebre punctata, parce tenuiterque nigro-pilosellum. Thorax coleopteris dimidio prope angustior, latitudine haud brevior, lateribus rectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis posterioribus rectis, modice transversim convexus, utrinque minus crebre profundius punctatus, linea lata media longitudinali laevi, parce nigro-pilosellus (puncto quoque pilifero). Scutellum crebre punctatum, nigro-pu-



bescens. Elytra thorace paulo longiora, parcius punctata, tenuiter nigro-pubescentia, virescenti-coerulea, pernitida. Abdomen crebrius at parum profunde punctatum, parce nigro-pubescentis, segmentis ultimis duobus totis rufis, parce rufo-pubescentibus. Pedes nigri, tarsi anticis ferrugineis. Alae fuscae.

Obs. Differt a praecedente statura graciliore, capite thoraceque minoribus, fortius et minus dense punctatis, labro testaceo, elytris laete virescenti-coeruleis ut in *St. cyaneletrio*.

\* 30. *St. hilaris*: *Niger, nitidus, abdomine segmentis ultimis duobus rufis, elytris cyaneis, antennis testaceis, ante apicem fuscis*. — Long.  $5\frac{1}{2}$  lin.

Habitat in Cayenna, Dom. Buquet.

Statura et summa affinitas *St. sapphirini*, niger, nitidus. Antennae capite plus sesqui longiores, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus, penultimis etiam haud transversis, ultimo ovato, apice fortiter acuminato, rufo-testaceae, articulis 5 — 9 fuscis, ultimis duobus ferrugineis. Palpi rufo-testacei. Labrum rufum. Caput thoracis latitudine, transversim subquadratum, depressum, parcius subtiliusque punctatum, fronte media laevi. Thorax coleopteris paululum angustior, latitudine haud brevior, basi angulisque posterioribus rotundatis, lateribus angulisque anterioribus rectis, apice truncatus, parum convexus, utrinque parce subtiliterque punctatus, linea media longitudinali laevi, ut caput parce subtiliterque nigro-pilosus (puncto quoque pilifero). Scutellum punctulatum, parcius nigro-pubescentis. Elytra thorace tertia parte longiora, subtiliter densius punctata, longius nigro-pubescentia, cyanea, nitida. Abdomen crebre punctulatum, longius nigro-pubescentis, segmentis ultimis duobus totis rufis, rufo-pubescentibus. Pedes validiusculi, nigri, nigro-pubescentes, tarsi anticis concoloribus.

\* 31. *St. tepidus*: *Niger, nitidus, abdomine segmentis ultimis duobus testaceis, elytris cyaneis, antennis fuscis, basi testaceis*. — Long.  $5\frac{1}{2}$  lin.

Habitat in Cayenna, Dom. Buquet.

Statura et summa affinitas praecedentis. Niger, nitidus. Antennae capite vix sesqui longiores, articulo tertio secundo vix longiore, 4 — 10 sensim paulo brevioribus et crassioribus, 7 — 10 transversis, ultimo breviter ovato, apice acuminato, dilute fuscae, articulis 4 primis testaceis. Palpi et labrum testacea. Caput thoracis latitudine, transversim subquadratum, depressum, supra versus latera et inter oculos punctis parvis profunde impressis, parce pilosus. Thorax coleopteris dimidio prope angustior, latitudine paululum brevior, lateribus modice rotundatus, pone medium subsinuatus, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, leviter transversim convexus, sat profunde parce punctatus, punctis exterioribus sparsis, dorso in series duas longitudinales digestis, spatio intermedio latiore laevi, versus latera parce nigro-pilosus. Scutellum crebre punctatum, tenuiter pubescens. Elytra thorace paulo longiora, densius punctata, cyanea, nitida, longius nigro-pubescentia. Abdomen crebre punctatum, nigro-pubescentis, segmentis ultimis duobus totis laete testaceis, testaceo-pubescentibus. Pedes nigri, tarsi anticis concoloribus.

- \* 32. *St. iopterus*: *Niger, nitidus, ano flavo, elytris violaceis, antennis testaceis.* — Long. 6 lin.

Habitat in Brasilia.

Statura et summa similitudo *St. sapphirini*. *Niger, nitidus*. Antennae graciliores, articulo tertio secundo plus sesqui longiore, 4 — 6 sensim brevioribus, 7 — 10 subaequalibus, leviter obconicis, crassitie haud brevioribus, ultimo breviter ovato, apice fortiter acuminato, totae dilute testaceae. Labrum et palpi testacea. Caput thoracis latitudine, subquadratum, depressum, disperse punctatum, punctis in media fronte rarioribus, parce nigro-pilosellum. Thorax coleopteris paulo angustior, latitudine haud brevior, basi rotundatus, lateribus subrectis, apice truncatus, angulis posterioribus obtusis, fere rotundatis, anterioribus rectis, modice convexus, utrinque crebrius punctatus, nigro-puberulus, linea media longitudinali laevi. Scutellum dense punctatum, nigro-pubescent. Elytra thorace longiora, profundius punctata, violacea, nitida, longius nigro-pubescentia. Abdomen densius punctulatum, parce nigro-pubescent, segmenti penultimi dimidio apicali segmentoque ultimo toto cum stylis analibus testaceo-flavis et testaceo-pubescentibus. Pedes nigri, tarsi anticis ferrugineis. Alae fusco-hyalinae.

- \* 33. *St. cyanelytrius*: *Niger, nitidus, ano testaceo, elytris coeruleis, antennis basi testaceis.* — Long. 6½ lin.

*Perty Delect. an. art. p. 30. t. 7. f. 3.*

Habitat in Brasilia, Dom. Virmond.

*Niger, nitidus*. Antennae capite sesqui longiores, graciliores, articulo tertio secundo sesqui longiore, 4 — 10 subobconicis, sensim paulo brevioribus, penultimis crassitie haud brevioribus, ultimo subovato, apice subito acuminato, fusco-nigrae, articulis tribus primis testaceis. Palpi testacei. Caput thoracis latitudine, subquadratum, supra subdepressum, disperse punctatum, punctis in fronte rarioribus, media fronte laevi, parce nigro-pilosellum. Thorax coleopteris dimidio fere angustior, latitudine haud brevior, basin versus subangustatus, basi rotundatus, lateribus pone medium subsinuatus, apice truncatus, angulis posterioribus obtusis, anterioribus acutiusculis, leviter transversim convexus, utrinque disperse punctatus, linea media longitudinali laevi, parce nigro-pilosellus (puncto quoque pilifero). Scutellum crebre punctatum, tenuiter nigro-pubescent. Elytra thorace sesqui longiora, minus profunde punctata, nigro-pubescentia, virescenti-coerulea vel laete cyanea, nitida. Abdomen nitidum, metallico-versicolor, parce punctulatum parceque nigro-pubescent, ano usque ad segmenti quinti dimidium testaceo. Pedes nigri, nigro-pubescentes, tarsi anticis concoloribus. Alae fuscae.

- \* 34. *St. chrysopygus*: *Niger, nitidus, ano flavo, elytris coeruleis, antennis testaceis.* — Long. 6 lin.

*Nordm. Symb. 45. 26.*

Habitat in Brasilia, Dom. v. Olfers.

Statura et summa similitudo sequentis. *Niger, nitidus*. Antennae capite plus sesqui longiores, graciliores, articulo tertio secundo sesqui lon-

giore, 4—6 teretibus, crassitie longioribus, licet sensim brevioribus, 7—10 inter se subaequalibus, leviter obconicis, crassitie vix brevioribus, ultimo ovato, apice abrupte acuminato, totae rufo-testaceae. Palpi rufo-testacei. Labrum ferrugineum. Caput thoracis latitudine, subquadratum, depressum, media fronte laevi. Thorax coleopteris dimidio angustior, latitudine vix brevior, basi rotundatus, lateribus anterieus rotundatis, posterius sinuatis, apice truncatus, angulis posterioribus obtusis, anterioribus subrectis, leviter transversim convexus, utrinque disperse parce punctatus, linea media longitudinali laevi. Scutellum dense punctatum, nigro-pubescent. Elytra thorace sesqui prope longiora, densius minus profunde punctata, tenuiter nigro-pubescentia, coerulea, nitida. Abdomen parce punctatum, parce nigro-pubescent, nitidum, metallico-versicolor, segmenti penultimi dimidio apicali, ultimoque toto cum stylis analibus testaceo-flavis, testaceo-pubescentibus. Pedes nigri, nigro-pubescentes, tarsi anticis ferrugineis. Alae fuscae.

\* 35. *St. pyraeter*: *Niger, nitidus, capite thoraceque subviolaceis, elytris violaceis, antennarum articulis duobus primis anoque testaceis.* — Long. 5 lin.

*Staph. haemorrhoidalis Germ. Spec. Ins. 54. 56.*

Habitat in Brasilia, Dom. Prof. Germar.

Statura fere *St. sapphirini*, at paulo minor, niger, nitidus. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus et crassioribus, 7—10 transversis, ultimo breviter ovato, apice abrupte acuminato, pilosellae, nigrae, articulis duobus primis rufo-testaceis. Palpi rufo-testacei. Labrum nigro-piceum. Caput thoracis latitudine, subquadratum, depressum, supra utrinque parce, supra oculos crebre punctatum, medio late longitudinaliter laeve, nigro-violaceum, utrinque parce pilosellum, subtus nigrum. Thorax coleopteris paulo angustior, latitudine vix longior, lateribus rectis, basi cum angulis posterioribus rotundatus, apice truncatus, utrinque subsinuatus, angulis anterioribus acutiusculis, sat convexus, crebre fortiterque punctatus, linea media longitudinali laevi, nigro-violaceus. Scutellum crebre punctatum, nigrum, nigro-pubescent. Elytra thorace longiora, crebrius punctata, longius nigro-pubescentia, laete violacea, nitida. Abdomen crebre punctatum, parcius nigro-pubescent, segmentis ultimis duobus testaceis, testaceo-pubescentibus, penultimo basi nigro. Pedes nigri, tarsi anticis concoloribus. Alae fusco-hyalinae.

\* 36. *St. chrysurus*: *Niger, nitidus, abdominis segmentis ultimis duobus rufis, elytris cyaneis.* — Long. 3 lin.

*Philonithus chrysurus Nordm. Symb. 108. 98.*

Habitat in Parà interiore Brasiliae, Sieber.

Statura omnino praecedentis at quadruplo fere minor, niger, nitidus. Antennae capite paulo longiores, crassiusculae, articulo tertio secundo sesqui prope longiore, 4—10 sensim paulo crassioribus, 6—10 transversis, ultimo subovato, apice subito acuminato, totae nigrae. Caput thoracis latitudine, transversim subquadratum, subdepressum, punctis raris dispersis profunde impressum. Thorax coleopteris paulo angustior, latitudine haud brevior, basin versus subangustatus, lateribus ante medium leviter rotundatis,



pone medium leviter sinuatis, apice truncatus, basi cum angulis posterioribus rotundatus, apice utrinque oblique truncatus, angulis anterioribus acutiusculis, subdepressus, dorso seriebus duabus minus regularibus 9 — 10-punctatis, utrinque punctis parvis dispersis notatus. Scutellum punctulatum. Elytra thorace paulo longiora, parcius punctata, longius nigro-pubescentia. Abdomen fortius crebriusque punctulatum, nigro-pubescentis, segmentis ultimis duobus totis rufis, rufo-pubescentibus. Pedes breves, nigri, femoribus anticis compressis, tarsis piceis. Alae fusco-hyalinae.

\* 37. *St. faustus*: *Niger, nitidus, abdominis segmentis ultimis duobus flavo-testaceis, elytris cyaneis, antennarum basi, coxis femoribusque anticis testaceis.* — Long. 4 lin.

Habitat in Cayenna, Dom. Buquet.

Subdepressus, niger, nitidus. Antennae capite plus sesqui longiores, articulo secundo tertioque modice elongatis, hoc longiore, 4 — 10 sensim paulo brevioribus et crassioribus, 7 — 10 transversis, ultimo ovato, apice abrupte fortiter acuminato, fuscae, articulis duobus primis testaceis, tertio quartoque rufo-piceis. Palpi rufi. Labrum ferrugineum. Caput thorace paululum latius, transversim suborbiculatum, depressum, parce fortiter vage punctatum, fronte medio laeviore. Thorax coleopteris angustior, latitudine paulo brevior, basin versus subangustatus, lateribus antierius rotundatis, pone medium subsinuatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus obtusiusculis, subdepressus, dorso utrinque serie minus regulari 8—9-punctata, lateribus parce disperseque punctatus, punctis omnibus sat profundis, parce nigro-pilosellus. Scutellum dense punctatum, nigro-pubescentis. Elytra thorace longiora, levius punctata, nigro-pubescentia, cyaneæ, nitida. Abdomen parce punctatum parceque nigro-pubescentis, segmentis ultimis duobus totis testaceo-flavis, flavo-pubescentibus. Pedes nigri, coxis femoribusque anticis testaceis, tarsis anticis ferrugineis.

### Fam. VIII.

*Antennae filiformes, articulo ultimo apice emarginato.*

*Palpi omnes filiformes, maxillares articulo quarto tertio aequali.*

*Pedes breves, tibiis posterioribus spinulosis, tarsis anticis dilatatis.*

Corpus subtiliter pubescens tomentosumve, scutello plerumque dense tomentoso. Antennae saepius breviores, apicem versus haud incrassatae, articulo primo modice elongato, secundo tertioque obconicis, 6—10 saepius transversis, ultimo apice truncato vel emarginato et inferne acuminato, articulis 4 primis breviter pilosellis, sequentibus pube subtilissima brevissimaque densius vestitis. Caput porrectum, oculis minoribus obliquis, ovatis, minus prominulis, apici magis quam basi capitis approximatis. Labrum breve, limbo angustiore coriaceo, densius barbato cinctum. Mandibulae validae, at minus elongatae, falcatae, medio fortiter dentatae. Palpi breviusculi, maxillares articulo secundo paulo longiore, tertio quartoque longitudine subaequalibus, hoc apice acuminato, labiales articulis tribus sensim paulo longioribus, secundo interne setuloso, tertio apice subacuminato. Elytra angulo apicali exteriore rotundato. Abdomen iam parallelum iam apicem versus sensim magis minusve angustatum. Pedes breviusculi, tibiis posterioribus spinulosis, spinis

apicalibus fortioribus, tarsis posterioribus brevioribus, articulo primo leviter elongato, sequentibus tribus sensim paululum brevioribus, tarsis anticis articulis 4 primis dilatatis, patellam ovatam, subtus dense tomentoso-spongiosam formantibus.

Mas plerumque nonnisi abdominis segmento inferiore sexto apice emarginato distinctus.

Obs. Species plurimas continet haec Fam., quae genuina in hoc genere videtur. Abdomen saepius maculis tomentosis ornatum.

- \* 38. *St. Chrysis: Rufus, capite flavo, thorace postice elytrisque aureo-tomentosis, abdomine nigro-aeneo, segmentis singulis late rufo-marginatis. — Long. 10 lin.*

*Grav. Mon. 124. 147. — Nordm. Symb. 31. 7.*

*Staph. auricomus Brull. D'Orbigny It. Am. mer. Ins. t. 5. f. 6.*

Habitat in Bahía, Gomez.

Antennae capite sesqui longiores, apice subattenuatae, articulo tertio secundo paulo longiore, 4 — 7 cyathiformibus, sensim paulo brevioribus, 8 — 10 transversim subobconicis, ultimo transverso, apice emarginato, interne abrupte acuminato, piceae, apice et articulo primo obscure rufis. Palpi rufo-brunnei. Mandibulae nigro-piceae. Labrum rufo-brunneum. Caput thorace paulo angustius, orbiculatum, convexum, subtilissime punctulatum, dilute flavum, aeneo-micans, subtiliter denseque flavo-aureo-pubescent, macula rhombea inter antennis subimpressa obscure aenea opaca, infra antennis et subtus rufo-brunneum. Thorax basi coleopteris paulo angustior, latitudine parum brevior, antrorsum subangustatus, basi, angulis posterioribus lateribusque eodem arcu rotundatis, antrorsum subangustatus, apice truncatus, angulis anterioribus acutiusculis, basi et lateribus marginatus, dorso modice convexus, nigro-subaeneus, opacus, antice nigro-, postice fulvo-aureo- dense tomentosus. Scutellum atro-tomentosum. Elytra thorace haud longiora, confertim punctato-subrugulosa, nigro-brunnea, pube depressa rufo-aurea dense vestita. Abdomen apicem versus sensim angustatum, parce punctulatum, tenuiter nigro-pilosum, nitidum, nigro-subaeneum, segmentis singulis late sinuatim rufo-marginatis, sexto toto rufo. Pectus rufo-brunneum. Pedes rufi.

Mas abdominis segmento inferiore quinto arcu longe ciliato impresso insignis.

- \* 39. *St. regalis: Rufo-testaceus, capite, thorace elytrisque aureo-tomentosis, thorace, elytrorum fascia abbreviata, abdominis maculis fasciisque violaceis. — Long. 7 lin.*

Habitat in Brasilia, Dom. Prof. Germar.

Statura omnino praecedentis, at dimidio minor. Antennae capite paulo longiores, articulo tertio secundo paulo longiore, quarto quintoque obconicis, sequentibus leviter transversis, ultimo apice truncato, inferne abrupte acuminato, totae nigrae. Palpi rufi. Mandibulae rufae, apice nigrae. Caput thoracis latitudine, supra minus convexum, crebre subtiliter punctatum, totum rufo-testaceum, supra pube depressa aurea nitida densiore, et pilis subtilibus longioribus nigris parcis vestitum. Thorax coleopteris dimidio

angustior, basin versus subangustatus, latitudine antice haud longior, basi et lateribus leviter rotundatus, apice truncatus, angulis posterioribus subrotundatis, anterioribus rectis, modice convexus, confertissime punctatus, linea tenui elevata laevi, purpureo-violaceus, nitidulus, pube depressa aurea nitida dense obductus. Scutellum nigrum, nigro-tomentosum. Elytra thorace longiora, confertissime subtilius punctata, rufo-testacea, pube longiore depressa aurea nitida dense vestita, fascia media, ad suturam abbreviata violacea, nigro-pubescente. Abdomen apicem versus sensim angustatum, parce subtiliter punctatum, tenuiter aureo-pubescentius, segmentis 4 primis rufo-testaceis, singulo supra et infra maculis ternis nigris, transversim positis, intermedia maiore transversa, lateralibus marginalibus minutis rotundatis, segmento quinto nigro-subviolaceo, summa basi rufo-testaceo, summo margine apicali flavo. Pectus rufo-testaceum, dense aureo-tomentosum. Pedes rufo-testacei.

Obs. Sequentibus duobus coloribus assimilis, thorace autem aureo-tomentoso primo intuitu distinctus.

\* 40. *St. biplagiatus*: Rufo-testaceus, thorace cupreo, nitido, elytris aureo-pubescentibus, macula media transversa cyanea, abdomine nigro, segmentis singulis rufo-marginatis. — Long. 10 lin.

Perty Delect. An. art. p. 31. t. 7. f. 4.

Habitat Cassapavae in Brasilia, Sellow.

Antennae capite paulo longiores, crassiusculae, articulo primo elongato, tertio secundo sesqui longiore, 5 — 10 fortius transversis, interne subserratis, ultimo transverso, apice leviter emarginato et inferne acuminato, nigrae, articulo primo piceo. Palpi rufo-picei. Mandibulae rufae, apice nigrae. Caput thorace latius, transversim suborbiculatum, supra leviter convexus, dilutius testaceum, metallico-subnitidum, punctatum, tenuiter fulvo-pubescentius. Thorax coleopteris angustior, latitudine paulo brevior, basi fortiter, lateribus vix rotundatus, apice truncatus, angulis anterioribus rectis, posterioribus rotundatis, lateribus et basi marginatus, dorso modice convexus, crebre punctatus, linea media longitudinali laevi, cupreus, nitidus, violaceo-micans, parce tenuiter nigro-puberulus. Scutellum nigrum, atro-tomentosum. Elytra thorace paulo longiora, subtiliter punctulata, longius fulvo-pubescentia, testacea, macula magna media transversa seu fascia ad suturam abbreviata nigro-subviolacea. Abdomen apicem versus subangustatum, nigro-subcyaneum, nitidum, parcius punctatum, parce pubescens, segmentis singulis late sinuatum rufo-marginatis, ultimo toto rufo. Pectus rufum, aureo-tomentosum, antice nigricans. Pedes rufo-testacei. Alae fuscae.

Obs. Antennae ab oculis remotae, vix magis inter se quam ab his distantes.

\* 41. *St. cyanicollis*: Rufo-testaceus, fronte puncto violaceo, thorace cyaneo, nitido, elytris aureo-pubescentibus, macula transversa cyanea, abdomine nigro, segmentis singulis rufo-marginatis. — Long. 7 lin.

Laporte Etud. Ent. I. 115. 5.

*Staph. interruptus* Brull. D'Orbigny It. Am. mer. Ins. t. 5. f. 7.

Habitat in Brasilia, Sellow.

Praecedente dimidio minor. Antennae capite paulo longiores, articulo



tertio secundo paulo brevior, 5—10 transversis, ultimo transversò, apice leviter emarginato et inferne acuminato, fuscae, articulis primis rufo-piceis. Oris partes rufo-testaceae, mandibulae apice nigrae. Caput thorace vix latius, supra minus convexum, crebre fortius punctatum, testaceum, nitidum, fronte paulo ante oculos macula minuta media violacea notata, tenuiter aureo-pubescent, parce nigro-pilosum. Thorax coleopteris parum angustior, basin versus haud angustatus, latitudine paulo longior, basi rotundatus, lateribus subrectis, apice truncatus, angulis anterioribus deflexis rectis, posterioribus obtusis, subrotundatis, modice convexus, crebre profundius punctatus, linea media longitudinali laevi, cyaneus, nitidus, violaceo-micans, parce tenuiterque nigro-pubescent, lateribus longius nigro-pilosellus. Scutellum nigrum, nigro-tomentosum. Elytra thoracis latitudine, confertim subtilius licet sat profunde punctata, testacea, dense aureo-pubescentia, parce nigro-pilosa, macula transversa, vel fere obliqua, biloba, seu fascia interne abbreviata nigro-violacea. Abdomen apicem versus subangustatum, crebrius punctatum, parcius fulvo-pubescent et nigro-pilosellum, nigrum, nitidum, segmentis 4 anterioribus margine apicali lato sinuatoque, quinto margine tenui apicali, sexto toto rufis. Pectus rufo-testaceum, tenuiter fulvo-pubescent. Pedes rufo-testacei.

Mas abdominis segmento inferiore quinto arcu longius densiusque viloso notato insignis.

Obs. Praecedenti similis, antennarum autem ab oculis minus distantium structura, thorace longiore, fronte minus convexa, macula violacea notata, etc. sat distinctus.

\* 42. *St. chrysocephalus*: *Niger, tomento fusco-nebuloso vestitus, ventre argenteo-villoso, capite femorumque annulo testaceis.* — Long. 5—7 lin.

*Fourcr. Ent. Paris. I. 165. 8. — Latr. Hist. nat. Crust. et Ins. IX. 296. 7. — Grav. Mon. 124. 146. — Nordm. Symb. 31. 12.*

*Emus chrysocephalus* *Mannerh. Brachelytr. 21. 4. — Boisd. et Lacord. Faun. Ent. Paris. I. 362. 5.*

*Staph. pubescens* *Rossi Faun. Etr. Mant. 97. 217. Ed. Hellw. I. 442. 217. — Walken. Faun. Paris. I. 269. 3. — Oliv. Ent. III. 42. 16. 15. t. 2. f. 15.*

*Le Staph. à tête jaune Geoffr. Hist. Ins. I. 363. 8.*

Habitat in Gallia, Italia, Dalmatia.

Statura et summa affinitas speciei sequentis, plerumque tamen paulo maior. Antennae capite paulo longiores, articulo tertio secundo paulo longiore, 6—10 sensim paulo brevioribus et fortius transversis, ultimo apice oblique emarginato et inferne acute acuminato, testaceae, articulis ultimis 5 fuscis. Palpi testacei. Mandibulae nigrae, basi testaceae. Caput maris thorace latius, feminae eiusdem latitudinis, suborbiculatum, testaceum, pubes densa flavescenti-grisea, albido-versicolore vestitum, punctis duobus minutis inter oculos nigris. Thorax coleopteris dimidio fere angustior, latitudine antica parum longior, lateribus ante medium subangulato-rotundatis, pone medium sinuatis; basi leviter rotundatus, apice truncatus, angulis posterioribus obtusissimis, anterioribus subrectis, parum convexus, niger, ferrugineo-subvariegatus, tomento subfasciculato fusco nebulosus. Scutellum maculis duabus oblongis nigro-tomentosis. Elytra thorace paulo longiora, nigra, tomento fusco reticulatim nebulosa, margine humerali testaceo. Ab-

domen nigrum, supra segmentis singulis apice medio ferrugineis, tomento dense fusco-variegato, subtus tomento spisso cinereo-albido vestitum. Pectus nigrum, cinereo-pubescent. Pedes nigri, annulo testaceo ante femur apicem ornati.

Mas capite latiore et tibiis intermediis leviter curvatis distinctus.

Obs. Sequenti similis et affinis, capite autem testaceo insignis. Europae meridionali propius.

- ✕ \* 43. *St. pubescens*: Niger, tomento denso fusco-nebuloso vestitus, ventre argenteo-villoso, capite flavescenti-villoso, femoribus annulo testaceo. — Long. 5 — 7 lin.

De Geer Ins. IV. 17. 2. Fab. Ent. Syst. I. II. 520. 5. Syst. El. II. 590. 6. — Payk. Mon. Staph. 16. 9. Faun. Suec. III. 376. 13. — Panz. Ent. Germ. 352. 4. — Cederh. Faun. Ingr. 117. 355. — Grav. Micr. 12. 13. Mon. 123. 144. — Latr. Hist. nat. Crust. et Ins. IX. 295. 6. — Marsh. Ent. Britt. 500. 10. — Gyll. Ins. Suec. II. 284. 5. — Zetterst. Faun. Lappon. 1. 67. 2. Ins. Lappon. 59. 3. — Sahlb. Ins. Fenn. I. 309. 3. — Nordm. Symb. 31. 9. — Er. Col. March. I. 434. 5.

*Emus pubescens* Mannerh. Brachelytr. 21. 5. — Boisd. et Lacord. Faun. Ent. Paris. I. 361. 4. — Runde Brachelytr. Hal. 2. 3.

Habitat in Europa.

Antennae capite paulo longiores, articulo tertio secundo paulo longiore, 6 — 10 transversis, subaequalibus, ultimo transverso, apice emarginato et inferne acuminato, nigrae, articulis basalibus testaceis, supra nigris. Palpi nigri. Mandibulae nigrae, basi testaceae. Labrum testaceum. Corpus nigrum. Caput maris thorace latius, feminae eiusdem latitudine, orbiculatum, nigrum, villis densis flavo-griseis obductum, macula minuta in media fronte punctisque nonnullis sparsis laevissimis. Thorax coleopteris angustior, latitudine antica vix longior, lateribus ante medium leviter rotundatis, pone medium subsinuatis, basi rotundatus, apice truncatus, angulis posterioribus obtusis, subrotundatis, anterioribus subrectis, supra tomento subfasciculato fusco-nebulosus, margine basali summo ferrugineo. Scutellum maculis duabus oblongis nigro-tomentosis. Elytra thorace paulo longiora, simili modo uti thorax fusco-tomentosa, margine humerali testaceo. Abdomen supra totum tomento fusco-variegato obductum, subtus cum pectore tomento spisso cinereo-albo, argenteo-micante vestitum. Pedes nigri, albido-pubescentes, femoribus ante apicem annulo testaceo.

Mas capite latiore et tibiis intermediis curvatis distinctus.

Obs. Corporis tomento similis omnino *St. nebuloso* et *murino*, licet antennarum, palporum, labri et tarsorum structura satis distinctus, neque iis affinis.

- \* 44. *St. chloropterus*: Capite thoraceque cupreis, elytris viridibus, abdomine ferrugineo, basi obscuro, pedibus testaceis, femoribus fusco-annulatis. — Long.  $4\frac{1}{2}$  lin.

Panz. Faun. Germ. 36. 20. — Grav. Micr. 166. 15. Mon. 121. 141. — Fab. Syst. El. II. 590. 5. — Latr. Hist. nat. Crust. et Ins. IX. 306. 22. — Nordm. Symb. 33. 14.

*Emus chloropterus* Boisd. et Lacord. Faun. Ent. Paris. I. 364. 7.

Habitat in Austria.

Antennae capite duplo longiores, articulo tertio secundo paulo lon-

giore, quarto quintoque oblongis, cylindricis, 6 — 10 subobconicis, sensim paululum brevioribus, licet omnibus crassitie longioribus, ultimo praecedente paulo brevior, apice truncato et inferne acuminato, testaceae. Os testaceum. Caput subquadratum, thoracis latitudine, subdepressum, confertim subtilius punctatum, cupreo-aeneum, nitidulum, parcius griseo-pubescent. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus parallelis, subrectis, basi fortius rotundatus, apice truncatus, angulis posterioribus obtusissimis, subrotundatis, anterioribus subrectis, perparum convexus, confertim punctatus, basi linea longitudinali subelevata laevi abbreviata, cupreo-aeneus, nitidulus, griseo-pubescent. Scutellum nigro-tomentosum. Elytra thorace vix longiora, obsolete punctata, viridia, opaca, margine laterali apicalique tenui testaceo, pube grisea nebulosa vestita. Abdomen ferrugineum, opacum, griseo-pubescent, segmentis anterioribus basi nigricantibus, maculis binis nigro-holosericeis. Pectus nigrum, griseo-pubescent. Pedes testacei, coxis nigris, femoribus omnibus facie antica macula media fusca notatis.

- \* 45. *St. ochropygus*: Niger, capite, thorace elytrisque coeruleis, his fascia undulata cinereo-pubescente, scutello atro-tomentoso, abdomine apice testaceo, supra fusco-tessellato. — Long. 7 lin.

Nordm. Symb. 46. 28.

*Emus nebulosus* Percheron Genera Ins. III. Nr. 1. (Coleopt.) tab. 5.

Habitat in Parà interiore Brasiliae, Sieber; in Cayenna, Dom. Buquet.

Statura omnino *St. chalocephali*. Antennae capite vix longiores, articulo secundo tertioque obconicis subaequalibus, sequentibus transversis, 6 — 10 brevioribus, inter se aequalibus, ultimo leviter transverso, apice truncato et inferne acute acuminato, nigro-piceae, apice summo ferrugineo. Palpi picei. Caput thorace paulo angustius, subtriangulare, subdepressum, supra sat crebre punctatum, fusco-pubescent, coeruleum, parum nitidum, subtus nigrum, nitidum. Thorax basi coleopteris latitudine aequalis, latitudine haud brevior, antrorsum leviter angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, utrinque subsinuatus, angulis anterioribus acutiusculis, parum convexus, dense punctatus, postice lineola media longitudinali laevi, fusco-pubescent, coeruleus, parum nitidus. Scutellum dense atro-tomentosum. Elytra thorace vix longiora, subtiliter alutacea, parcius punctulata, cyanea, opaca, fascia media valde undulata dense cinereo-pubescente, basi etiam undulatum cinereo-pubescente, interstitiis subtilissime nigro-tomentosis. Abdomen nigrum, segmentis ultimis duobus testaceis, reliquis supra tomento brevissimo fusco variegatis, subtessellatis, subtus parce pubescentibus. Pectus nigrum, lateribus subtiliter cinereo-pubescent. Pedes nigri vel picei.

Mas femoribus anterioribus basi leviter sensim dilatatis, tibiis intermediis subarcuatis.

Variat femoribus omnibus rufo-piceis.

- \* 46. *St. antiquus*: Niger, capite thorace elytrisque obscure aeneis, pube fusco-grisea subvariegatis, scutello nigro-pubescente, abdomine supra fusco-subtessellato, ano rufo-testaceo. — Long. 5 — 7 lin.

Nordm. Symb. 35. 16.

Habitat in Columbia, Brasilia.



Habitu omnino *St. chalcocephali*, plerumque autem paulo minor. Antennae capite paulo longiores, articulo secundo tertio paulo brevior, 6 — 10 transversim subobconicis, inter se aequalibus, ultimo praecedente paulo longiore, apice emarginato et inferne acuminato, nigrae, articulo primo subtus rufo-piceo. Palpi picei. Caput thorace minus et angustius, subtriangulare, subdepressum, supra dense fortius punctatum, carinula minuta frontali laevi, obscure aeneum, subopacum, densius fusco-pubescent. Thorax coleopteris paulo angustior, latitudine vix longior, antrosum subangustus, lateribus perparum rotundatis, pone medium nullo modo sinuatis, basi rotundatus, apice truncatus, angulis anterioribus rectis, posterioribus obtusis, subdepressus, confertissime fortius punctatus, carinula postica polita, obscure aeneus, opacus, densius fusco-pubescent, lateribus parce longius pilosus. Scutellum atro-tomentosum. Elytra thorace vix longiora, subtiliter alutacea, obscure aenea, opaca, pube densa subtili fusca cinerea-que nebula. Abdomen nigrum, segmento quinto apice sextoque toto rufo-testaceis, supra tomento tenui fusco-griseo nigroque variegatum. Pectus nigrum. Pedes nigri, femoribus anticis supra linea longitudinali lutea. Alae fuscae.

Antennae feminae capite vix longiores.

- \* 47. *St. latro*: *Niger, capite thoraceque punctatissimis, opacis, nigro-pubescentibus, scutello atro-tomentoso, elytris obscure aeneis, abdomine supra bifariam nigro-maculato, ano rufo-testaceo.* — Long.  $7\frac{1}{2}$  lin.

Habitat in Brasilia, Dom. Prof. Germar.

Affinis praecedenti. Antennae capite sesqui longiores, articulo tertio secundo sesqui longiore, 4 — 10 subaequalibus, leviter obconicis, crassitie paululum longioribus, ultimo apice leviter emarginato et inferne acuminato, totae nigrae. Palpi picei. Caput thoracis latitudine, subquadratum, basi emarginatum, angulis posterioribus retrorsum leviter prominentibus, supra depressiusculum, confertissime punctatum, pube breviuscula erecta nigra parcius vestitum, nigrum, opacum, infra sublaeve glabrumque, nitidum. Thorax coleopteris vix angustior, latitudine haud brevior, lateribus rectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, supra parum convexus, confertissime punctatus, pube brevi erecta nigra vestitus, niger, opacus. Scutellum nigrum, atro-tomentosum. Elytra thorace paulo longiora, confertissime subtiliter punctata, aenea, subopaca, e pube brevi densiore nigra griseaque obsolete variegata. Abdomen apicem versus sensim angustatum, parcius obsoletius punctatum, nigrum, supra opacum, tenuiter nigro-pubescent, segmentis 5 primis singulo maculis duabus triangularibus approximatis atro-tomentosis, tomento subtiliore fulvo-grisea cinctis ornatis, apice aureo-pubescente, margine apicali pallido, infra nitidum, metallico-versicolor, tenuiter nigro-pubescent, segmento sexto toto rufo-testaceo, subtiliter aureo-pubescente. Pectus nigrum, subnitidum, subtiliter nigro-pubescent. Pedes nigri.

- \* 48. *St. exulans*: *Niger, capite, thorace elytrisque obscure aeneis, scutello atro-tomentoso, elytris abdominisque dorso griseo-subtessellatis.* — Long. 5 lin.

Habitat in America septentrionali, Dom. Prof. Germar.

Statura omnino *St. antiqui* et minoribus eius individuis aequalis, niger, parum nitidus. Antennae capite paulo longiores, articulo tertio secundo sesqui longiore, 6—10 transversis, at sensim paulo brevioribus, ultimo apice emarginato et inferne acuminato, piceae, articulis tribus primis nigris. Caput thorace minus et angustius, subtriangulare, subdepressum, supra crebre fortius punctatum, linea media longitudinali antice abbreviata laevi, griseo-pubescentis, nigro-aeneum, nitidum, ore nigro, subtus parce punctatum, nigrum, nitidum, subtiliter nigro-pubescentis. Thorax basi coleopteris parum angustior, latitudine haud brevior, apicem versus leviter angustatus, lateribus leviter, basi cum angulis posterioribus fortiter rotundatus, apice truncatus, angulis anterioribus subrectis, parum convexus, creberrime fortius punctatus, linea media longitudinali medio subinterrupta laevi, griseo-puberulus, nigro-aeneus nitidus. Scutellum atro-tomentosum. Elytra thorace paulo longiora, subtilissime rugoso-punctulata, nigro-aenea, opaca, e pube subtili depressa obsolete griseo-nigroque tessellata. Abdomen e pube subtili depressa supra nigro-griseoque variegatum, subtus segmentis singulis basi cinereo-pubescentibus. Pedes nigri, geniculis tarsisque piceis.

\* 49. *St. scriptus*: *Niger, dense fulvo-pubescentis, antennis pedibusque testaceis, elytris ferrugineis abdomineque tomento fusco-variegatis.* — Long. 7 lin.

*Nordm. Symb.* 32. 13.

Habitat ad Promont. Bonae Spei, Dom. Prof. Lichtenstein.

Antennae capite paulo longiores, articulo secundo tertioque obconicis, subaequalibus, quarto quintoque subglobosis, 6 — 10 transversis, aequalibus, ultimo iterum subgloboso, apice profunde transversim impresso. Palpi et labrum testacea. Mandibulae nigrae. Caput thoracis prope latitudine, antrosum parum angustatum, supra subdepressum, subtiliter punctulatum, dense aureo-pubescentis, nigro-subaeneum, margine antico orbitalique ferrugineo, subtus nigrum, nitidum, glabrum. Thorax coleopteris paulo angustior, latitudine paululum longior, lateribus parallelis subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrectis, leviter convexus, confertissime subtiliter punctatus, linea media longitudinali elevata laevissima medio interrupta, pube densa aurea obtectus, nigro-subaeneus, margine omni tenuiter ferrugineo. Scutellum nigro-tomentosum. Elytra thorace vix longiora, ferruginea, subtiliter punctata, fulvo-pubescentia, maculis liturisque fuscis variegata. Abdomen supra nigro-subaeneum, segmentis singulis margine apicali ferrugineis, obsolete punctulatum, punctis maioribus parvis sparsum, fulvo-pubescentis, pube fusca maculosum, subtus nigrum, segmentis singulis basi dense argenteo-pubescentibus. Pectus nigrum, fulvo-pubescentis. Pedes rufo-testacei, coxis maculaque magna oblonga femorum basali nigris.

\* 60. *St. maculosus*: *Nigro-aeneus, capite thoraceque ferrugineo-variegatis, elytris ferrugineis, fulvescenti-pubescentis, elytris abdomineque fusco-maculosis.* — Long. 8 — 10 lin.

*Grav. Micr.* 165. 13. *Mon.* 123. 145. — *Nordm. Symb.* 31. 6.

*Staph. viduatus* *Fab. Syst. El. II.* 591. 7.

*Staph. erythropennis* *Mannerh. Brachelytr.* 22. 6. — *Nordm. Symb.* 62. 47.

Habitat in America septentrionali.

Antennae capite paulo longiores, articulo tertio secundo paulo longiore, quarto subgloboso, 6—10 transversis, aequalibus, ultimo apice profundius emarginato, inferne acuminato, testaceae, articulo tertio fusco, vel fuscae basi testaceae. Caput subquadratum, thoracis latitudine, supra subdepressum, confertissime punctatum, fulvescenti-pubescentia, aeneum, parum nitidum, magis minusve ferrugineo-variegatum, interdum ferrugineo-bivittatum. Thorax coleopteris parum angustior, latitudine antica vix brevior, basin versus subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus acutis, leviter convexus, confertim punctatus, linea media longitudinali, antrorsum mox obsoleta, elevata laevi, fulvescenti-pubescentia, obscure aeneus, vix nitidus, basi apiceque ferrugineo-variegatus, interdum ferrugineo-bivittatus. Scutellum aeneum, maculis duabus longitudinalibus nigro-holosericeis. Elytra thorace vix longiora, subtiliter punctata, fulvescenti-pubescentia, fusco-maculosa. Abdomen subtiliter punctulatum, supra nigro-subaeneum, segmentis singulis apice obscure ferrugineis, pube subtiliori densa fusco-nebulosa vestitum, biseriatim nigro-maculatum, subtus nigrum, flavescenti-pubescentia, segmentis singulis basi densius flavescenti-griseo-pubescentibus. Pectus nigrum, flavescenti-pubescentia. Pedes rufo-testacei, femoribus supra coxisque anterioribus nigris.

\* 51. *St. fuscomaculatus*: *Niger, capite thoraceque aeneo-ferrugineoque variegatis, elytris ferrugineis, pedibus testaceis, femoribus nigro-maculatis.* — Long.  $6\frac{2}{3}$  lin.

*Emus fuscomaculatus* Laporte *Etud. Ent.* I. p. 113.

Habitat in Mexico, Dom. C. Ehrenberg.

Praecedenti affinis, at multo minor et minus robustus. Antennae capite paulo longiores, articulo tertio secundo sesqui longiore, sequentibus sensim paulo brevioribus, penultimis tribus vel quatuor transversis, ultimo subovato, apice emarginato et inferne acuminato, nigrae, articulo primo testaceo, apicali ferrugineo. Palpi picei. Mandibulae nigrae. Caput thoracis latitudine, antrorsum modice angustatum, supra parum convexus, confertissime punctatum, fulvescenti-pubescentia, ferrugineum vel testaceum, fronte magis minusve aeneo-variegata, lateribus nigro-aeneis, subtus nigrum, nitidum, glabrum. Thorax elytris parum angustior, latitudine haud brevior, lateribus parallelis, rectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, supra parum convexus, confertissime punctatus, linea longitudinali media laevi medio obsoleta, tenuius fulvescenti-pubescentia, vel aeneus, vittis duabus irregularibus ferrugineis, vel testaceis, vittis tribus longitudinalibus inaequalibus nigro-aeneis, lateralibus brevioribus, arcuatis. Scutellum nigro-subaeneum, nigro-tomentosum, basi utrinque cinereo-pubescentia. Elytra thorace paulo longiora, subtiliter punctata, ferruginea vel testacea, fulvescenti-pubescentia, fusco-substrigosa. Abdomen nigrum, griseo-pubescentia, subtus segmentis singulis macula laterali basali testacea, segmentis retractis occulta. Pectus nigrum, flavescenti-pubescentia. Pedes testacei, coxis nigris, femoribus anterioribus disco anteriore et posteriore nigro, posticis basi lineola dorsali nigra. Alae fusco-hyalinae.



- \* 52. *St. fossor*: *Niger, capite thoraceque brunneo-rufis, tibiis, tarsis elytrisque rufo-testaceis, his apice dilutioribus, flavo-pubescentibus, scutello atro-tomentoso, abdomine variegato.* — — Long. 8 lin.

*Scop. Ann. hist. nat. V. 109. 105. — Goeze Ent. Beitr. I. 725. 17. — Fab. Ent. Syst. I. II. 523. 16. Syst. El. II. 593. 18. — Panz. Ent. Germ. 353. 12. Faun. Germ. 27. 6. — Grav. Micr. 10. 9. Mon. 117. 136. — Latr. Hist. nat. Crust. et Ins. IX. 299. 12. — Nordm. Symb. 60. 43.*

*Emus fossor* Boisd. et Lacord. *Faun. Ent. Paris. I. 367. 11.*

*Staph. erythropterus* var. 2 *Scop. Ent. Carn. 101. 306.*

*Staph. fodiens* Grav. *Mon. 116. 134. — Nordm. Symb. 62. 45.*

Habitat in Europa temperata.

Antennae capitis thoracisque prope longitudine, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus, penultimo vix transverso, ultimo apice inferne acuminato, nigrae, apicem versus picescentes, articulo ultimo ferrugineo. Palpi picei, apice rufi. Caput thoracis latitudine, suborbiculatum, supra parum convexum, confertissime subtiliter punctatum, brunneo-rufum, opacum, brevissime fulvo-pubescent, subtile nigrum, nitidum. Thorax coleopteris angustior, latitudine antica paulo longior, basin versus subangustatus, lateribus ante medium leviter rotundatis, pone medium subsinuatis, basi leviter, angulis posterioribus fortius rotundatis, apice truncatus, angulis anterioribus subrectis, deflexis, parum convexus, subtiliter obsolete carinatus, confertim subtiliter punctatus, opacus, brunneo-rufus, pube brevissima fulvo-sericans. Scutellum atro-tomentosum. Elytra thorace vix longiora, confertissime subtiliter punctata, opaca, dilutius brunneo-rufa, fulvo-pubescentia, apice oblique dilutiora et densius flavo-aureo-pubescentia. Abdomen nigrum, supra subtilissime brevissimeque nigro-tomentosum, segmentis singulis basi macula minuta triangulari aureo-tomentosa terminata, subius segmentis 2 — 4 fascia basali cinereo-pubescente, nitida. Pectus nigrum. Pedes nigri, tibiis tarsisque rufo-testaceis.

Obs. *Staph. fodiens* Grav. e Carolina a *St. fossore* Europaeo nullo modo distinctus videtur. Gravenhorst ipse dubius hanc speciem separavit, neque ulla ex eius descriptione differentia patet. Constat varias huius familiae species Europae et Americae septentrionali communes esse. Fortasse etiam specimen collectionis Boiscianae, a Gravenhorstio descriptum, patria falsa notatum fuit.

- \* 53. *St. erythropterus*: *Niger, opacus, antennarum basi apiceque, elytris pedibusque rufis, scutello maculisque posterioribus abdominis orichalceo-tomentosis.* — Long. 6 — 7 lin.

*Linn. Faun. Suec. n. 842. Syst. Nat. I. II. 683. 4. — Ström. Act. Nidros. III. 404. 26. t. 6. f. 8. — De Geer. Ins. IV. 21. 6. — Müll. Faun. Frid. 23. 223. Prodr. 97. 1092. — Goeze Ent. Beitr. I. 716. 4. — De Vill. Ent. I. 411. 4. — Oliv. Ent. III. 42. 12. 10. t. 2. f. 14. — Cederh. Faun. Ingr. 117. 358. — Er. Col. March. I. 434. 6.*

*Staph. erythropterus* var. *Fab. Ent. Syst. I. II. 522. 14.*

*Staph. castanopterus* Grav. *Micr. 10. 10. Mon. 116. 132. — Latr. Hist. nat. Crust. et Ins. IX. 298. 10. — Gyll. Ins. Suec. II. 295. 14. — Zetterst. Faun. Lappon. I. 68. 4. Ins. Lappon. 60. 5. — Sahlb. Ins. Fenn. I. 312. 12. — Mannerh. Brachelytr. 22. 3. — Nordm. Symb. 55. 36. — Runde Brachelytr. Hal. 2. 2.*

Habitat in Europa.

*Niger, opacus.* Antennae capite thoraceque paululum breviores, articulo secundo tertio paulo brevior, 4—10 sensim paulo brevioribus, 4—7 subcylindricis, 8—10 obconicis, hoc vix transverso, ultimo apice inferne acuminato, fuscae, basi apiceque rufae. Palpi rufi. Caput thorace latius, orbiculatum, supra subdepressum, subtiliter confertim punctatum, fronte posterius subtilissime carinata, anterieus inter oculos pedunculoque orichalceo-pubescentibus. Thorax coleopteris angustior, latitudine paulo longior, basin versus subangustatus, lateribus modice subsinuatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrectis, leviter deflexis, leviter transversim convexus, confertim subtiliter punctatus, uti caput brevissime subtiliter nigro-pubescentis. Scutellum orichalceo-tomentosum. Elytra thoracis longitudine, subtiliter punctata, rufa, rufo-pubescentia. Abdomen confertim subtilissime punctatum, subtiliter nigro-pubescentis, supra segmento primo, quarto, quinto sextoque utrinque macula, subtus segmentis ultimis tribus fascia basali interrupta aurichalceo-tomentosa ornatis. Pectus parapleuris orichalceo-tomentosis. Pedes rufi, coxis nigris.

Obs. Hunc verum *Staph. erythropterum* Linnei esse e descriptione sua satis patet.

X \* 54. *St. caesareus: Niger, opacus, elytris pedibusque rufis, antennis testaceis, apice fuscis, scutello atro-tomentoso, thoracis margine posteriore maculisque abdominis orichalceo-tomentosis.* — Long. 8—9 lin.

*Cederh. Faun. Ingr. 335. 1055. t. 3. f. c. — Er. Col. March. I. 435. 7.*

*Staph. erythropterum* Fab. *Syst. Ent.* 265. 5. *Spec. Ins.* I. 335. 5. *Mant. Ins.* I. 220. 8. *Ent. Syst.* I. II. 522. 14. *Syst. El.* II. 593. 16. — *Schrank Enum. Ins. Aust.* 230. 435. *Faun. Boica.* I. 640. 862. — *Harrer Besch. d. Ins.* I. 150. 114. — *Foucr. Ent. Par.* I. 166. 9. — *Walken. Faun. Paris.* I. 270. 7. — *Donov. Britt. Ins.* IX. t. 308. — *Rossi Faun. Etrusc.* I. 247. 609. *Ed. Hellw.* I. 306. 609. — *Payk. Mon. Staph.* 7. 3. *Faun. Suec.* III. 369. 3. — *Panz. Faun. Germ.* 27. 4. *Ent. Germ.* 353. 10. t. 12. f. 10. — *Marsh. Ent. Britt.* 499. 6. — *Grav. Micr.* 10. 11. *Mon.* 116. 133. — *Latr. Hist. nat. Crust. et Ins.* IX. 297. 9. *Gen. Crust. et Ins.* I. 297. 2. — *Gyll. Ins. Suec.* II. 293. 13. — *Zetterst. Faun. Lappon.* I. 68. 3. *Ins. Lappon.* 59. 4. — *Sahlb. Ins. Fenn.* I. 311. 11. — *Mannerh. Brachelytr.* 22. 2. — *Nordm. Symb.* 55. 37. — *Runde Brachelytr. Hal.* 2. 1.

*Emus erythropterum* Boisd. et Lacord. *Faun. Ent. Paris.* I. 365. 8.

*Staph. erythropterum* var. 1. *Scop. Ent. Carn.* 101. 306.

*Le Staph. à étuis couleur de rouille Geoffr. Hist. Ins.* I. 364. 9.

*Staph. primus* Schaeff. *Icon.* I. t. 2. f. 2. *Elem. Ent.* t. 117.

Habitat in Europa.

*Niger, opacus.* Antennae capite thoraceque breviores, articulo tertio leviter elongato, secundo paulo longiore, sequentibus subobconicis, sensim paulo brevioribus, penultimis duobus vel tribus transversis, ultimo apice leviter emarginato et inferne acuminato, testaceae, apice fuscae. Palpi testacei. Caput thoracis latitudine, suborbiculatum, supra subdepressum, subtilius confertissime punctatum, breviter subtiliterque nigro-pubescentis, fronte posterius subtilissime carinata, antice, pone oculos pedunculoque orichalceo-pubescentis. Thorax coleopteris paulo angustior, latitudine perparum longior, lateribus vix rotundatis, pone medium subsinuatis, basi cum angulis posterioribus rotundatus, apice utrinque subsinuatus, angulis anterioribus subrotundatis, leviter transversim convexus, subtiliter confertissime puncta-

tus, longitudinaliter subtilissime carinatus, brevissime subtiliter nigro-pubescentis, margine posteriore orichalceo-tomentoso. Scutellum atro-tomentosum, margine laterali basi orichalceo-pubescente. Elytra thorace paulo longiora, confertissime subtiliter punctata, rufa, rufo-pubescentia. Abdomen subtilissime confertissimeque punctatum, subtiliter nigro-pubescentis, supra segmento primo margine apicali, sequentibus quatuor macula laterali obliqua, subtus omnibus, primo excepto, macula utrinque transversali basali orichalceo-tomentosa. Pectus apice orichalceo-pubescentis, lateribus utrinque maculis duabus orichalceo-tomentosis. Pedes rufi, coxis nigris.

Variat antennis totis testaceis.

Obs. Staph. aeneocephalus Panz. Faun. Germ. 27. 5, elytris fuscis, haud differe videtur.

\*55. *St. vulpinus*: Niger, supra ferrugineo-pubescentis, capite supra obscure rufo, antennarum basi, pedibus, elytris anoque rufis, scutello atro-tomentoso, abdomine flavo-tomentoso-submaculato. — Long.  $8\frac{1}{2}$  lin.

Nordm. Symb. 53. 34.

Staph. immaculatus Mannerh. Brachelytr. 22. 7. — Nordm. Symb. 62. 48. (forte)

Habitat Philadelphiae, Kinn.

Statura omnino St. maculosi et parum minor. Antennae capite paulo longiores, articulo tertio secundo paulo brevior, sequentibus sensim paulo brevioribus, penultimo autem non transverso, ultimo apice truncato et inferne acute acuminato, fuscae, basi rufae. Palpi rufi. Mandibulae nigrae. Caput thoracis latitudine, oblongum, antrorsum subangustatum, supra perparum convexum, fortius confertim punctatum, supra os impressum, obscure rufum, macula minuta mediae frontis nigricante, ferrugineo-pubescentis, subtus nigrum, nitidum, glabrum. Thorax coleopteris parum angustior, latitudine paulo brevior, lateribus subrectis, parallelis, basi cum angulis posterioribus fortiter rotundatus, apice truncatus, angulis anterioribus rectis, leviter transversim convexus, confertim punctatus, subtiliter carinatus, niger, opacus, dense ferrugineo-pubescentis. Scutellum nigro-tomentosum. Elytra thorace parum longiora, subtiliter dense punctata, rufa, ferrugineo-pubescentia. Abdomen nigrum, supra subtilissime nigro-tomentosum, segmentis 4 primis macula utrinque obliqua, minus distincta, pubescente orichalcea, margine laterali apicalique ferrugineo-pubescentibus, subtus flavo-pubescentibus, apice ferrugineo-marginatis, segmento sexto toto rufo. Pectus orichalceo-pubescentis. Pedes rufi, coxis nigris, anticis apice rufescentibus.

Obs. Descriptio Mannerheimiana supra citata nimis manca: „Linearis, rufo-piceus, opacus, abdomine obscuriore, subtiliter punctulatus, antennis, capite, pedibus, elytris anoque rufo-cinnabarinis, pectore segmentorumque abdominalium marginibus flavo-aureis. — Statura St. lutarii, sed maior, magis linearis et aliter coloratus.”

\*56. *St. castaneus*: Niger, supra ferrugineo-pubescentis, capite obscure rufo, antennarum basi, elytris, pedibus anoque rufis, scutello atro-tomentoso. — Long. 8 lin.

Nordm. Symb. 54. 35.

Habitat in Mexico, Dom. Deppe.



Statura et summa affinitas praecedentis, niger, opacus. Antennae capite parum longiores, articulo tertio secundo sesqui prope longiore, quarto subgloboso, 5 — 10 sensim paulo brevioribus, 7 — 10 transversis, ultimo apice truncato et inferne acuminato, nigrae, basi rufae. Palpi rufi. Caput thoracis prope latitudine, antrorsum subangustatum, supra subdepressum, subtiliter confertissime punctatum, obscure rufum, ferrugineo-pubescent, sub-  
tus nigrum, nitidum. Thorax coleopteris paululum angustior, latitudine parum brevior, lateribus subrectis, basi cum angulis posterioribus fortiter rotundatus, apice truncatus, angulis anterioribus rectis, transversim convexus, confertissime subtiliter punctatus, postice carinula subtili polita, ferrugineo-pubescent. Scutellum atro-tomentosum. Elytra thorace vix longiora, confertissime subtiliter punctata, rufa, rufo-pubescentia. Abdomen supra nigrum, subtiliter fusco-pubescent, e pube nigra bifariam maculatum, margine laterali elevato segmentique quinti margine apicali obscure rufo, sub-  
tus parcius pubescens, segmentis singulis rufo-marginatis, segmento sexto toto obscure rufo. Pectus fulvo-pubescent. Pedes rufi, coxis nigris, anticis rufescentibus.

\* 57. *St. phoenicurus*: *Niger, nigro-pubescent, scutello atro-tomentoso, elytris anoque rufis, rufo-pubescentibus.* — Long. 7 lin.

Nordm. Symb. 57. 40.

Habitat in Mexico, Dom. Deppe.

Statura omnino *St. stercorarii* et maioribus eius individuis aequalis. Niger, opacus. Antennae capite sesqui longiores, articulo secundo tertio dimidio prope brevior, 4—10 subobconicis, sensim paulo brevioribus, pen-  
ultimis vix transversis, ultimo apice truncato et inferne acuminato, nigrae, articulo ultimo obscure ferrugineo. Palpi picei. Caput thoracis latitudine, antrorsum subangustatum, supra perparum convexus, confertim minus subtiliter punctatum, carinula longitudinali tenuissima antice abbreviata, nigro-pubescent. Thorax coleopterorum prope latitudine, latitudine vix brevior, lateribus parum, basi cum angulis posterioribus fortiter rotundatis, apice truncatus, angulis anterioribus rectis, leviter transversim convexus, minus subtiliter confertissime punctatus, subtiliter carinatus, nigro-pubescent. Scutellum atro-tomentosum. Elytra thorace vix longiora, confertissime subtiliter punctata, rufa, rufo-pubescentia. Abdomen subtiliter punctulatum, nigro-pubescent, supra e pube densiore bifariam nigro-maculatum, segmento sexto toto rufo. Pedes nigri.

X

\* 58. *St. stercorarius*: *Niger, capite thoraceque nigro-pubescentibus, antennarum basi elytris pedibusque rufis, scutello atro-tomentoso, abdomine cinereo-sericeo-maculato.* — Long. 6 — 8 lin.

Oliv. Ent. III. 42. 18. t. 3. f. 23. — Latr. Hist. nat. Crust. et Ins. IX. 299. 11. — Marsh. Ent. Britt. 499. 7. — Grav. Micr. 11. 12. Mon. 115. 130. — Gyll. Ins. Suec. II. 296. 15. — Sahlb. Ins. Fenn. I. 312. 13. — Mannrh. Brachelytr. 22. 4. — Nordm. Symb. 55. 38. — Runde Brachelytr. Hal. 2. 3. — Er. Col. March. I. 436. 8.

*Emus stercorarius* Boisd. et Lacord. Faun. Ent. Paris. I. 366. 10.

Habitat in Europa.

*St. erythroptero* plerumque paulo minor, praecipue angustior et magis

parallelus, niger, opacus. Antennae capite paulo longiores, articulo tertio secundo longiore, 6 — 10 transversis, subaequalibus, ultimo apice truncato et inferne acuminato, rufae, extrorsum piceae. Palpi rufi. Caput thoracis latitudine, antrorsum leviter angustatum, supra subdepressum, sat crebre punctatum, dense nigro-pubescent. Thorax coleopterorum fere latitudine, latitudine paulo brevior, lateribus subrectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, supra parum convexus, confertim punctatus, carinula postica polita iam infra medium abbreviata, dense nigro-pubescent. Scutellum atro-tomentosum. Elytra thoracis longitudine, subtiliter punctulata, rufa, rufo-pubescentia. Abdomen supra subtilissime brevissimeque nigro-tomentosum, segmentis quatuor primis macula obliqua laterali, ultimis duobus fascia lata basali cinereo-sericeo-tomentosis, subtus segmentis omnibus fascia basali cinereo-sericeo-pubescente. Pedes breves, rufi, coxis nigris, anticis subtus rufis.

- \* 59. *St. lutarius*: Niger, capite thoraceque supra aeneis, fulvo-pubescentibus, antennis, elytris, tibiis tarsisque rufis, scutello atro-tomentoso, abdomine cinereo-sericeo-maculato. — Long. 8 lin.

Grav. Mon. 115. 131. — Mannerh. Brachelytr. 23. 8. — Nordm. Symb. 58. 41.

*Staph. flavopunctatus* Latr. Hist. nat. Crust. et Ins. IX. 297. 8.

Habitat in Austria.

Niger, capite thoraceque supra aeneis. Antennae capite vix longiores, articulo tertio secundo aequali, sequentibus transversis, sensim magis obconicis, ultimo transverso, apice emarginato et inferne acuminato, rufo-testaceae, apice fuscescentes. Palpi testacei. Caput thorace paulo minus et angustius, subtriangulare, supra subdepressum, dense punctatum, fulvo-pubescent, subtus nigrum, parce nigro-pubescent. Thorax coleopterorum prope latitudine, latitudine paulo brevior, antrorsum subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, supra parum convexus, dense punctatus, fulvo-pubescent. Scutellum macula magna rotundata apicali atro-holosericæ. Elytra thorace paulo longiora, subtiliter crebre punctata, rufa, fulvo-pubescentia. Abdomen breviter subtiliterque nigro-tomentosum, supra segmentis tribus primis basi maculis tribus, quarto quintoque fascia, subtus segmentis omnibus, ultimo excepto, fascia basali cinereo-tomentosis, argenteo-nitidulis. Pectus nigrum, cinereo-pubescent. Pedes nigri, femoribus apice, tibiis tarsisque omnibus rufis.

Variat femoribus magis minusve nigris.

- \* 60. *St. chaltocephalus*: Niger, capite thoraceque supra aeneis, fulvescenti-pubescentibus, elytris, tibiis tarsisque rufis, scutello atro-tomentoso, abdomine cinereo-sericeo-maculato. — Long. 7 lin.

Fab. Syst. El. II. 593. 17. — Er. Col. March. I. 436. 9.

*Staph. aeneocephalus* Fab. Ent. Syst. I. II. 522. 17. — Panz. Ent. Germ. 353. 11.

*Staph. ochropterus* Germ. Spec. Ins. 34. 57. Faun. Ins. Europ. VIII. 14. — Nordm. Symb. 59. 42.

*Emus Carinthiacus* Boisd. et Lacord. Faun. Ent. Paris. I. 365. 9.

Habitat in Germania, Borussia, Gallia.

Niger, capite thoraceque supra aeneis. Antennae capite sesqui longiores, articulo tertio secundo sesqui prope longiore, sequentibus sensim paulo brevioribus, 7—10 transversis, ultimo apice truncato et inferne acuminato, nigrae, apice interdum picescentes. Palpi rufi. Caput thorace minus et angustius, subtriangulare, supra subdepressum, confertim sat fortiter punctatum, fulvo-pubescent. Thorax basi coleopterorum latitudine, latitudine vix brevior, antrorsum subangustatus, lateribus perparum, basi et angulis posterioribus fortiter rotundatis, apice truncatus, angulis posterioribus subrectis, perparum convexus, confertim sat fortiter punctatus, basi carinula abbreviata polita, fulvo-pubescent. Scutellum atro-tomentosum. Elytra thorace vix longiora, subtiliter punctulata, rufa, fulvo-pubescentia. Abdomen supra subtilissime brevissimeque nigro-tomentosum, segmentis anterioribus quatuor basi maculis tribus, ultimis duobus fascia cinereo-sericeo-tomentosis, illis inter maculas brunneo-tomentosis, subtus segmento singulo fascia basali cinereo-sericeo-tomentosa. Pedes nigri, tibiis tarsisque rufis.

Obs. Staph. latebricola specim. Boruss. Grav. Mon. p. 115. ad hanc speciem pertinet.

\* 61. *St. latebricola*: Niger, capite thoraceque supra aeneis, fulvescenti-pubescentibus, elytris pedibusque rufis, scutello atro-tomentoso, abdomine cinereo-sericeo-maculato. — Long.  $5\frac{1}{2}$  lin.

Grav. Mon. 113. 129. — Nordm. Symb. 56. 39. — Er. Col. March. I. 437. 10.

Habitat in Germania, Brunsvigae in acervis formicarum rufarum lectus a Dahlho.

Niger, capite thoraceque supra aeneis. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 6—10 transversim subobconicis, aequalibus, ultimo crassitie haud brevior, apice truncato et inferne acuminato, piceae, basi nigrae. Palpi piceae. Caput thorace vix angustius, subquadratum, fulvescenti-pubescent. Thorax coleopterorum prope latitudine, latitudine vix brevior, lateribus subrectis, antrorsum subangustatus, apice truncatus, angulis anterioribus rectis, basi cum angulis posterioribus rotundatus, leviter convexus, dense profunde punctatus, carinula postica laevi, fulvescenti-pubescent. Scutellum atro-tomentosum. Elytra thorace vix longiora, subtiliter punctulata, rufa, fulvo-pubescentia. Abdomen supra subtilissime brevissimeque nigro-tomentosum, segmentis quatuor primis basi maculis tribus, ultimis duobus, ut subtus omnibus fascia cinereo-sericeo-tomentosis. Pedes rufi, coxis trochanteribusque nigris, femoribus anticis basi piceis.

Obs. Praecedenti valde affinis, differt autem statura minore et angustiore, thorace minus lato, capite minus triangulari, minus confertim punctato, femoribus posterioribus totis rufis.

\* 62. *St. fulvipes*: Niger, opacus, capite, thorace elytrisque obscure cyaneis, abdomine segmentis ultimis duobus basi cinereo-tomentosis, antennis basi apiceque pedibusque rufis. — Long. 6—7 lin.

Scop. Ent. Carn. 99. 301. — Goeze Ent. Beitr. I. 725. 12. — Er. Col. March. I. 438. 11.

Staph. bicinctus Rossi Faun. Etr. I. Mant. 96. 216. Ed. Helw. 442. 216. — Block Ins. Vall. Plauens. 115. 3. f. 3.



*Staph. erythropus* Payk. Faun. Suec. III. 372. 7. — Gyll. Ins. Succ. II. 290. 11. — Sahlb. Ins. Fenn. I. 310. 8. — Germ. Faun. Ins. Europ. VI. 14. — Mannerh. Brachelytr. 24. 22. — Nordm. Symb. 73. 71. — Runde Brachelytr. Hal. 4. 15.  
*Staph. tricolor* Grav. Micr. 7. 6. Mon. 110, 113. — Latr. Hist. nat. Crust. et Ins. IX. 305. 20.

*Staph. azureus* Gebl. Ledebour. It. Append. III. 70. 4.

Habitat in Europa et Sibiria, ubique rarus.

Statura fere *St. chalconecephali*, at paulo angustior. Antennae capite thoraceque parum breviores, articulo tertio secundo parum longiore, sequentibus sensim paulo brevioribus, penultimis etiam crassitie haud brevioribus, ultimo apice subemarginato et inferne acuminato, nigrae, articulis duobus basalibus duobusque apicalibus rufis. Palpi rufi. Caput thorace minus, subtriangulare, supra subdepressum, crebre sat profunde punctatum, nigro-subcyaneum vel subvirescens, parum nitidum, nigro-pubescent. Thorax coleopterorum latitudine, latitudine paululum brevior, lateribus vix rotundatus, antrorsum subangustatus, apice truncatus, angulis anterioribus subrectis, basi cum angulis posterioribus rotundatus, parum convexus, creberrime punctatus, nigro-cyaneus, brevis nigro-pubescent. Scutellum nigrum, nigro-tomentosum. Elytra thorace haud longiora, punctato-subrugulosa, nigro-pubescentia, cyanea vel virescentia, nigro-pubescentia. Abdomen nigrum, nigro-pubescent, supra segmentis ultimis duobus fascia e tomento subtili albedo-cinereo, argenteo-micante. Pectus nigrum. Pedes rufi, coxis nigris.

Obs. Abdomine contracto fasciae haud conspicuae.

\* 63. *St. Buquetii*: Niger, capite, thorace elytrisque nigro-cyaneis, opacis, subtiliter atro-tomentosis, alis fuscis. — Long. 10 — 12 lin.

Laporte Etud. Ent. I. 114. 2.

*Staph. atrox* Nordm. Symb. 38. 19.

Habitat in Brasilia.

Magnus, robustus, niger, subtus subnitidus, supra opacus, capite, thorace elytrisque obscure cyaneis, tomento subtili brevissimo nigro-sericante vestitus. Antennae capite paulo longiores, tenuiores, subcompressae, articulo tertio secundo sesqui longiore, 4—10 sensim paulo brevioribus, 8—10 leviter transversis, ultimo apice profunde emarginato et inferne acuminato, nigrae, articulis 4 primis nigro-pilosellis, reliquis dense subtilissime cinereo-pubescentibus. Caput thoracis latitudine, antrorsum subangustatum, supra subdepressum, subtiliter confertissime punctatum. Thorax coleopteris paulo angustior, latitudine paulo brevior, lateribus rectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, parum convexus, subtiliter confertissime punctatus, postice carinula tenuissima parum distincta. Scutellum spissius atro-tomentosum. Elytra thorace longiora, omnium confertissime subtilissime alutacea. Abdomen supra densius nigro-pubescent, tomento atro bifariam maculatum. Pedes nigri. Alae fuscae, violaceo-nitidae.

Obs. Prof. Nordmann errore quodam Mexico huius insecti patriam affert.

- \* 64. *St. ferox*: *Niger, capite, thorace elytrisque cyaneis, subtiliter nigro-tomentosis, alis fulvis. — Long. 10 lin.*

*Nordm. Symb. 39. 20.*

Habitat Jalappae in Mexico, Dom. Deppe.

Statura et summa similitudo praecedentis. *Niger*, subtus subnitidus, supra capite, thorace elytrisque saturate cyaneis, opacis, subtiliter tenuiterque nigro-holosericeis. Antennae capite paulo longiores, articulo tertio secundo sesqui longiore, 4—10 sensim brevioribus, 8—10 fortiter transversis, ultimo transverso, apice profunde emarginato et inferne acuminato, nigrae. Caput thoracis fere latitudine, subquadratum, supra subdepressum, confertissime subtiliter punctatum. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus antice rectis, postice cum angulis posterioribus et basi rotundatis, apice truncatus, utrinque subsinuatus, angulis anterioribus acutiusculis, leviter transversim convexus, subtiliter confertissime punctatus, postice carinula tenui laevi. Scutellum atro-holosericeum. Elytra thorace paululum longiora, confertissime subtilissime punctato-rugulosa. Abdomen supra densius nigro-pubescent, tomento atro bifariam maculato. Pedes nigri. Alae fulvo-luteae, nitidae.

- \* 65. *St. fulvomaculatus*: *Niger. capite, thorace elytrisque cyaneis, opacis, subtiliter atro-tomentosis, elytris macula magna laterali aurantiaca, alis fulvis. — Long. 12 — 14 lin.*

*Nordm. Symb. 39. 21.*

Habitat Jalappae in Mexico, Dom. Deppe.

Statura omnino et similitudo *St. Buquetii*, niger, capite, thorace elytrisque supra saturate cyaneis, parum nitidis, supra subtiliter nigro-tomentosus. Antennae capite haud longiores, subcompressae, articulo tertio secundo paulo longiore, sequentibus omnibus transversis, sensim brevioribus, ultimo apice emarginato et inferne acuminato, nigrae. Palpi nigro-picei. Mandibulae nigrae. Caput thoracis latitudine, transversim subquadratum, antrorsum subangustatum, supra parum convexus, confertim subtilissime punctatum. Thorax coleopteris paulo angustior, latitudine haud brevior, basin versus subangustatus, lateribus pone medium subsinuatis, basi fortiter rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, modice transversim convexus, confertissime subtiliter punctatus. Scutellum nigrum, nigro-tomentosum. Elytra thoracis longitudine, confertissime subtiliter punctato-subrugulosa, macula magna laterali ovata aurantiaca notata. Abdomen apicem versus sensim angustatum, creberrime subtiliter punctulatum, tenuiter nigro-pubescent, dorso e tomento densiore bifariam nigro-maculato. Pedes nigri. Alae fulvae, fusco-marginatae.

Mas abdominis segmento inferiore quinto scopo medio longitudinali atro-piloso instructo.

- \* 66. *St. nobilis*: *Niger, capite thoraceque cyaneis, nitidis, elytris fulvis, margine laterali nigro. — Long. 9 lin.*

*Nordm. Symb. 41. 22.*

*Staph. chrysopterus* Brullé *D'Orbigny It. Am. mer. Ins. t. V. f. 8. (forte)*

Habitat Cassapavae in Brasilia, Sellow.

Corpus nigrum, subnitidum. Antennae capitis longitudine, articulo tertio secundo longiore, 4—10 fortiter transversis, 5—10 aequalibus, ultimo apice profunde emarginato et inferne acuminato, nigrae, articulis 4 primis nigro-pilosellis, reliquis tomento subtilissimo fusco-cinereo dense obductis. Caput thorace paulo angustius, apicem versus sensim angustatum, subtriangulare, cyaneum, nitidum, supra perparum convexum, densius subtiliusque punctatum, tenuiter nigro-pubescentibus pilosusque, subtus nigrum. Thorax coleopterorum latitudine vel basi eadem paululum superans, latitudine baseos brevior, antrorsum angustatus, lateribus parum, basi fortiter rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrectis, posterioribus obtusis, subrotundatis, supra leviter transversim convexus, cyaneus, nitidus, subtiliter punctatus, tenuiter nigro-pubescentibus pilosusque. Scutellum nigrum, dense nigro-pubescentibus. Elytra thorace paulo longiora, fulvo-aurea, pube depressa fulva nitida dense vestita, margine basali aeneo, margine laterali inflexo nigro, lateribus longe nigro-pilosella, apice nigro-ciliata. Abdomen punctulatum, nigro-pubescentibus, subtus metallico-versicolor. Pedes validiusculi, nigri, nigro-pubescentes. Alae fuscae, nitidae.

- \* 67. *St. erythrocnemus*: *Niger, opacus, capite, thorace elytrisque nigro-subaeneis, fulvo-pubescentibus, his margine laterali fusco-testaceo, scutello atro-tomentoso, tibiis tarsisque rufis.* — Long. 9 lin.

Nordm. Symb. 42. 24.

Habitat ad Promont. Bonae Spei, Krebs.

Statura et prope magnitudo *St. maculosi*, niger, opacus. Antennae capite vix longiores, articulo tertio secundo paulo longiore, 4—10 sensim brevioribus, 6—10 transversis, ultimo apice profunde emarginato et inferne acuminato, nigrae. Palpi picei. Caput basi prope thoracis latitudine, antrorsum angustatum, supra parum convexum, confertissime punctatum. Thorax coleopterorum prope latitudine, latitudine vix brevior, lateribus subrectis, parallelis, basi cum angulis posterioribus fortiter rotundatus, apice truncatus, angulis anterioribus rectis, supra parum transversim convexus, confertissime punctatus, linea media longitudinali laevi. Elytra thorace vix longiora, subtiliter punctata. Caput, thorax et elytra supra pube crassiuscula densa fulvo-aurea vestita, nigro-subaenea, elytra margine laterali fusco-testaceo. Scutellum atro-tomentosum. Abdomen supra subtiliter nigro-tomentosum, e tomento atro bifariam maculatum, subtus segmentis singulis fascia basali cinereo-pubescente micante. Pedes nigri, tibiis tarsisque testaceo-rufis. Alae fulvo-hyalinae.

- \* 68. *St. dimidiatus*: *Niger, opacus, capite thoraceque supra aenei., dense ferrugineo-pubescentibus, scutello atro-tomentoso, antennis, tibiis tarsisque rufis.* — Long. 9 lin.

Laporte Etud. Ent. 115. 4.

*Staph. velutinus* Nordm. Symb. 44. 25.

Habitat ad Promont. Bonae Spei, Krebs.

Statura omnino praecedentis, at minus robustus, ater, opacus. Antennae capite parum longiores, articulo tertio secundo paulo longiore, 4—10 sensim brevioribus, 6—10 transversis, ultimo apice emarginato et inferne



acuminato, totae testaceo-rufae. Palpi rufi. Caput basi thoracis latitudine, apicem versus leviter angustatum, supra leviter convexum, confertissime punctatum, aeneo-cupreum, dense ferrugineo-pubescentia, subtus nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus subrectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, supra leviter transversim convexus, confertissime punctatus, linea media longitudinali laevi, aeneo-cupreus, dense ferrugineo-pubescentia, lateribus nigro-pilosellus. Scutellum atro-tomentosum. Elytra thorace paulo longiora, confertissime subtilissimeque punctatopunctulosa, nigro-pubescentia. Abdomen dense nigro-pubescentia, dorso e tomento atro bifariam maculatum. Pedes nigri, tibiis tarsisque testaceo-rufis.

- \* 69. *St. Dauricus: Niger, capite thoraceque subaeneis, fulvo-pubescentibus, scutello atro-tomentoso, elytris, tibiis tarsisque testaceis.* — Long. 8 — 9 lin.

*Mannerh. Brachelytr.* 22. 5. — *Nordm. Symb.* 62. 46.

Habitat in Sibiria, sub cortice arborum corruptarum et sub fimo sicco, Pallas. Dedit Dom. Schüppel.

Statura fere *St. maculosi* et minoribus eius individuis aequalis, at paulo brevior, niger, parum nitidus, capite thoraceque supra aeneis. Antennae capite vix longiores, articulo tertio secundo haud longiore, sequentibus sensim paulo brevioribus, 6 — 10 transversis, ultimo his angustiore, apice emarginato et inferne acuminato, nigrae. Caput maris thoracis latitudine, feminae eodem angustius et minus, antrorsum angustatum, subtriangulare, supra subdepressum, confertissime punctatum, nigro-subaeneum, pube brevi fulva dense vestitum, subtus nigrum, nitidum. Thorax coleopteris parum angustior, latitudine vix brevior, basin versus subangustatus, lateribus perparum, angulis posterioribus et basi fortiter rotundatus, apice truncatus, angulis anterioribus rectis, supra leviter transversim convexus, confertissime subtilius punctatus, posteriorius carinula abbreviata polita notatus, nigro-subaeneus, minus nitidus, pube brevi fulva pubescens. Scutellum atro-tomentosum. Elytra thorace paulo longiora, confertim subtiliter punctata, fulvo-pubescentia, rufa, obsolete fusco-maculosa. Abdomen nigro-pubescentia, segmentis duobus primis fulvo-ciliatis. Pectus flavescenti-pubescentia. Pedes nigri, tibiis tarsisque rufis.

- \* 70. *St. fossator: Niger, opacus, nigro-pubescentia, elytris macula apicis aureo-tomentosa.* — Long. 7 lin.

*Grav. Micr.* 164. 12. *Mon.* 117. 135. — *Nordm. Symb.* 52. 33.

Habitat in America septentrionali, Knoch.

Statura et magnitudo omnino *St. chalcoccephali*, niger, opacus, pube brevi depressa nigra vestitus. Antennae articulo tertio secundo paulo longiore, 6 — 10 transversis, aequalibus, ultimo apice emarginato et inferne acuminato, nigrae. Palpi picei. Caput thorace paulo angustius, subtriangulare, supra subdepressum, fortius confertim punctatum. Thorax basi coleopteris parum angustior, latitudine vix brevior, antrorsum subangustatus, lateribus perparum, basi cum angulis posterioribus fortius rotundatus, apice truncatus,

angulis anterioribus rectis, subdepressus, confertim fortius punctatus, postice carinula brevi polita. Scutellum atro-tomentosum. Elytra thorace vix longiora, parcius subtiliter punctata, opaca, macula magna apicali, suturam haud attingente, aureo-tomentosa. Abdomen supra parce subtiliterque punctulatum, subtus parce pubeseens. Pedes nigri.

Obs. Elytra in nostro specimine obscure fusca, in specime altero melius conservato autem, a Dom. Hope benevole communicato, corpori concolora.

\* 71. *St. aurifluus*: *Niger, capite thoraceque supra cyaneis, elytris brunneis, apice aureo-tomentosis. — Long. 7 lin.*

Habitat in Brasilia, Dom. Prof. Germar.

Statura omnino praecedentis. Antennae capite sesqui longiores, tenuiores, articulo secundo tertioque modice elongatis, hoc longiore, sequentibus sensim paulo brevioribus, penultimis tribus transversis, ultimo apice truncato et inferne acuminato, nigrae. Caput thoracis latitudine, antrorsum vix angustatum, angulis posterioribus tumidulis, supra subdepressum, crebre punctatum, cyaneum, subnitidum, subtiliter tenuiterque nigro-pubescentibus, ore nigro, subtus nigrum, nitidum. Thorax coleopteris angustior, latitudine hand brevior, lateribus subrectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus obtusiusculis, supra parum convexus, dense minus subtiliter punctatus, posteriorius linea tenui abbreviata subelevata laevi, cyaneus, subnitidus, densius nigro-pubescentibus, lateribus parce nigro-pilosellus. Scutellum atro-tomentosum. Elytra thorace paulo longiora, obsoletius punctata, brunnea, humeris et lateribus inflexis dilutioribus, dorso nigricantia, subtiliter nigro-pubescentia, apice aureo-tomentosa. Abdomen coleopteris angustius, apicem versus sensim attenuatum, obsolete punctulatum, nigrum, nitidum, supra parce nigro-pubescentibus, e tomento subtilissimo atro bifariam maculato, subtus metallico-versicolor, segmentis singulis basi subtiliter tenuiterque griseo-pubescentibus. Pectus nigrum, tenuiter subtiliterque nigro-pubescentibus. Pedes tenuiores, nigri.

\* 72. *St. gentilis*: *Niger, capite thoraceque violaceis, scutello atro-tomentoso, elytris aureo-tomentosis, basi maculisque nonnullis disci nigricantibus. — Long. 6 — 7 lin.*

Habitat in Brasilia, Dom. Virmond.

Praecedenti affinis. Antennae capite paulo longiores, articulo tertio secundo sesqui longiore, sequentibus sensim brevioribus, 6—10 transversis, ultimo apice emarginato, nigrae. Palpi picei. Caput basi thoracis latitudine, antrorsum leviter angustatum, supra perparum convexus, crebre sat fortiter punctatum, parcius nigro-pubescentibus, violaceum, nitidulum, ore nigro, subtus nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine brevior, lateribus rectis, parallelis, basi cum angulis posterioribus fortiter rotundatus, apice truncatus, angulis anterioribus rectis, supra parum convexus, crebre punctatus, nigro-pubescentibus, violaceus, nitidulus. Scutellum atro-tomentosum. Elytra thorace longiora, subtiliter confertissime punctatopunctulosa, brunnea, dense fulvo-aureo-tomentosa, nitida, tenuiter nigro-pubescentibus, basi maculaque marginali punctoque dorsali medio nigris, dense nigro-pubescentibus. Abdomen nigrum, nigro-pubescentibus pilosellumque,

supra e tomento brevissimo obsolete bifariam atro-maculato, subtus nitidulum. Pectus nigrum, pube brevi griseo-sericans. Pedes nigri.

\* 73. *St. caliginosus*: *Niger, opacus, nigro-tomentosus, ano sanguineo, antennarum articulo primo subtus testaceo.* — Long.  $5\frac{1}{2}$  lin.

Habitat in Mexico.

Affinis *St. tomentosus*, at paulo minor, totus niger, opacus, tomento brevi nigro obductus. Antennae capite parum longiores, articulo secundo tertioque aequalibus, 4 — 10 subobconicis, inter se aequalibus, longitudine vix crassioribus, ultimo apice emarginato et inferne acuminato, nigrae, articulo primo subtus testaceo. Palpi rufo-picei. Caput thorace paulo angustius, subtriangulare, supra leviter convexum, confertissime punctatum, fronte subtilissime carinata, subtus laeve, glabrum, nitidum. Thorax coleopteris vix angustior, latitudine haud brevior, antrorsum vix angustatus, lateribus rectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, leviter convexus, confertissime subtiliter punctatus, subtilissime carinatus. Scutellum dense aterrimo-tomentosum. Elytra thoracis longitudine, confertissime subtiliter ruguloso-punctata. Abdomen supra omnium subtilissime rugulosum, e tomento aterrimo bifariam maculato, subtus confertim subtiliter punctatum, supra segmento penultimo dimidio ultimoque toto, subtus segmentis duobus ultimis obscure sanguineis. Pedes nigri, tarsis concoloribus. Alae fulvo-hyalinae.

\* 74. *St. sepulchralis*: *Niger, opacus, nigro-pubescent, antennis concoloribus, tarsis medio piceis.* — Long.  $5\frac{1}{2}$  lin.

Habitat in Mexico, Dom. Prof. Germar.

Statura omnino praecedentis, niger, opacus, dense subtilius nigro-pubescent. Antennae capite paulo longiores, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus, penultimis tribus leviter transversis, ultimo iterum paulo longiore, apice truncato et inferne acuminato. Palpi rufo-picei. Caput thorace paulo angustius, antrorsum angustatum, subtriangulare, supra subdepressum, confertissime fortius punctatum, pube brevi nigra densius vestitum, parce pilosellum, subtus parce subtiliter punctatum, glabrum, nitidum. Thorax coleopteris vix angustior, latitudine haud longior, antrorsum nullo modo angustatus, lateribus subrectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, parum convexus, confertim minus subtiliter punctatus, posteriorius carinula subtili valde abbreviata laevi nitida instructus, pube brevi depressa nigra densius obductus. Scutellum atro-tomentosum. Elytra thorace prope breviora, densius subtiliter punctata, dense nigro-pubescentia. Abdomen supra parce obsoleteque punctulatum, pube brevi subtilique nigra dense aequaliter obductum, subtus nitidum, confertim punctatum, parce tenuiterque pubescens. Pectus densius nigro-pubescent, opacum. Pedes nigri, tarsis medio rufescenti-piceis.

Obs. *St. tomentosus* simillimus et proxime affinis, statura autem paulo minore, corpore minus tomentoso quam pubescente, abdomine maculis atro-tomentosis destituto, capite thoraceque fortius punctatis, antennarum articulis secundo et tertio minus elongatis, tarsis medio rufescentibus distinguendus.



- \* 75. *St. fuscicornis*: *Niger, opacus, nigro-tomentosus, antennis, palpis anoque brunneis.* — *Long. 6 lin.*

*Germ. Ins. Spec. 33. 35.*

*Staph. lugubris Nordm. Symb. 49. 31.*

Habitat in Brasilia meridionali, Sellow.

Statura praecedentis, plerumque autem paulo maior, niger, opacus, pube depressa subtiliore nigro-fusca densius obductus. Antennae capite paulo longiores, articulo tertio secundo sesqui prope longiore, sequentibus sensim paulo brevioribus, penultimis vix transversis, ultimo praecedente paulo longiore, apice leviter emarginato et inferne acuminato, totae ferrugineae. Palpi castanei. Caput thoracis prope latitudine, antrorsum posterius leviter, antice subito fortiter angustatum, supra subdepressum, dense punctatum, subnitidum, subtus parce punctulatum, glabrum, nitidum. Thorax coleopterorum latitudine, latitudine vix brevior, antrorsum vix angustatus, lateribus rectis, basi rotundatus, angulis posterioribus obtusis, apice truncatus, angulis anterioribus rectis, leviter convexus, confertissime subtilius punctatus, carinula tenui postica laevi. Elytra thoracis longitudine, confertissime subtiliter punctato-rugulosa. Abdomen supra parcius obsoletiusque punctulatum, aequaliter nigro-pubescent, subtus nitidulum, parcius punctatum parceque pubescens, segmento quinto margine summo apicali sextoque toto brunneis. Pedes nigri, tibiis tarsisque interdum brunneis, tibiis anticis fulvo-pubescentibus. Alae fusco-hyalinae.

Obs. Antennis totis ferrugineis ab affinis distinctus.

- \* 76. *St. tomentosus*: *Niger, opacus, nigro-tomentosus, tarsis concoloribus, antennis apice ferrugineis.* — *Long. 6 lin.*

*Grav. Micr. 161. 4. Mon. 111. 118.* — *Nordm. Symb. 50. 32.*

Habitat in America septentrionali, Knoch.

Statura omnino *St. stercorarii*, totus niger, opacus, tomento brevi fusco-nigro vestitus. Antennae capite sesqui longiores, articulo secundo tertioque leviter elongatis, hoc paulo longiore, sequentibus sensim paulo brevioribus, penultimis tribus leviter transversis, ultimo iterum paulo longiore, apice subemarginato et inferne acuminato, nigrae, articulis penultimis subtus ultimoque toto obscure ferrugineis. Palpi nigri, articulo ultimo rufo-piceo. Caput thorace paulo angustius, antrorsum angustatum, subtriangulare, supra subdepressum, confertissime minus subtiliter punctatum. Thorax basi coleopterorum prope latitudine, latitudine haud brevior, antrorsum subangustatus, lateribus subrectis, pone medium subsinuatis, basi et angulis posterioribus rotundatis, apice truncatus, angulis anterioribus rectis, supra modice convexus, confertissime subtiliter punctatus. Elytra thoracis longitudine, subtiliter confertissime subruguloso-punctata. Abdomen supra confertissime subtilissimeque punctatum, parce punctulatum, segmentis anterioribus basi punctis duobus atro-tomentosis notatis, subtus parcius punctulatum, parce pubescens, subnitidum. Pedes nigri, tarsis concoloribus.

- \* 77. *St. afer*: *Niger, opacus, nigro-tomentosus, palpis tarsisque piceis.* — *Long. 6½ lin.*

Habitat ad Promont. Bonae Spei, Dom. Westermaun.

Statura omnino *St. chalconcephali*, niger, opacus, dense nigro-tomentosus. Antennae capite sesqui longiores, articulo tertio secundo sesqui prope longiore, sequentibus sensim paulo brevioribus, penultimis tribus leviter transversis, ultimo apice truncato et inferne acuminato, totae nigrae. Palpi rufo-picei. Caput thorace angustius et minus, subtriangulare, supra depressiusculum, confertissime minus subtiliter punctatum. Thorax basi coleopterorum latitudine, latitudine paulo brevior, antrorsum leviter angustatus, lateribus rectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, parum convexus, confertissime punctatus. Elytra thorace paululum longiora, subtiliter confertissime punctata. Abdomen confertissime punctatum, supra aequaliter tomentosum, subtus parum nitidum, parcius pubescens. Pedes nigri, tarsi picei.

Obs. Differt a praecedente antennis nigris et ano concolore, a *St. tomentosus* abdomine supra aequaliter tomentoso, immaculato, a *St. sepulchrali* statura latiore, thorace brevior, antrorsum angustato.

X

\*78. *St. cinnaopterus*: Niger, capite thoraceque aeneis, elytris, tibiis, ano abdominisque segmentorum marginibus rufis, scutello atro-tomentoso. — Long. 6 lin.

Grav. Micr. 164. 11. Mon. 113. 127. — Mannerh. Brachelytr. 23. 9. — Nordm. Symb. 59. 40.

Var. *Capite thoraceque badio-rufis.*

*Staph. zonatus* Grav. Micr. 162. 7. Mon. 113. 128. — Nordm. Symb. 61. 44.

*Staph. badius* Mannerh. Brachelytr. 23. 10. — Nordm. Symb. 59. 39.

Habitat in America septentrionali.

Antennae capite sesqui fere longiores, articulo secundo tertioque leviter elongatis, hoc paulo longiore, sequentibus sensim paulo brevioribus, penultimis vix transversis, ultimo apice truncato et inferne acuminato. Palpi rufi. Caput thoracis latitudine, latitudine paulo longius, antrorsum vix angustatum, subdepressum, supra fortius confertim punctatum, tenuiter ferrugineo-pubescens, vel obscure aeneum, subnitidum, vel obscure rufum, subtus nigrum, nitidum, parce subtiliterque punctatum. Thorax coleopteris paulo angustior, latitudine vix longior, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, leviter transversim convexus, fortius confertim punctatus, linea media longitudinali laevi, ferrugineo-pubescens, vel obscure aeneus, basi magis minusve rufescens, vel totus obscure rufus. Scutellum nigrum, atro-tomentosum. Elytra thorace vix longiora, subtilius confertim punctato-subrugulosa, rufo-castanea, ferrugineo-pubescens. Abdomen parce punctulatum, longius pubescens, nigrum, opacum, segmentis singulis rufo-marginatis, sexto toto rufo. Pectus nigrum. Pedes coxis nigris, femoribus picei, tibiis tarsisque rufo-testaceis.

\*79. *St. hottentottus*: Niger, capite thoraceque aeneis, subnitidis, elytris brunneis abdomineque pubescenti-subvariegatis, scutello atro-tomentoso. — Long. 5½ lin.

Nordm. Symb. 37. 17.

Habitat ad Promont. Bonae Spei.

Statura omnino *St. cinnamopteri*. Antennae capite sesqui longiores, articulo secundo tertioque leviter elongatis, hoc paulo longiore, sequentibus sensim paulo brevioribus, penultimis vix transversis, ultimo subovato, apice emarginato et inferne acuminato, obscure piceae, basi nigrae. Palpi picei. Caput thoracis fere latitudine, subquadratum, apice subito angustatum, supra subdepressum, aeneum, nitidulum, crebre punctatum, linea media longitudinali laevi, densius griseo-pubescentibus, ore nigro, subtus nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine vix brevior, basin versus subangustatus, lateribus perparum, basi et angulis posterioribus fortiter rotundatis, apice truncatus, angulis anterioribus rectis, supra leviter convexus, sat crebre punctatus, linea media longitudinali subelevata postice distincta, antrorsum obsoletiore, griseo-pubescentibus, aeneus, nitidulus. Scutellum nigrum, macula magna nigro-velutina. Elytra thoracis latitudine, subtiliter punctulata, rufo-brunnea, pube subtili cinerea maculatis densiore vestita. Abdomen subtiliter punctulatum, segmentis singulis supra punctis quatuor maioribus transversim positae impressis, nigrum, nitidulum, pubescens, supra anterieus fusco-subnebulosum, segmentis 5 primis in medio baseos macula minuta triangulari flavescenti-pubescente notatis. Pectus nigrum. Pedes nigri, tarsi, interdum etiam tibiis, piceis.

\* 80. *St. cupricollis*: *Niger, griseo-pubescentibus, capite thoraceque aeneis, subnitidis, elytris fusco-subaeneis, opacis, densius fusco-griseo-pubescentibus, scutello fusco-tomentoso. — Long. 6 lin.*

*Nordm. Symb. 34. 15.*

Habitat in Brasilia, in prov. Minarum, Dom. Beske.

Statura omnino *St. lutarii*, at paulo minor. Antennae capite parum longiores, articulo secundo tertioque aequalibus, obconicis, quarto subgloboso, 5 — 10 transversis, sensim paulo brevioribus, ultimo subovato, apice emarginato et inferne acuminato, obscure piceae, basi nigrae. Palpi picei. Caput thorace paulo angustius, subtriangulare, subdepressum, supra aeneum, nitidulum, sat crebre punctatum, griseo-pubescentibus, ore nigro, subtus nigrum, nitidum. Thorax coleopteris vix angustior, latitudine paulo brevior, lateribus subrectis, parallelis, basi et angulis posterioribus rotundatis, apice truncatus, utrinque subsinuatus, angulis anterioribus acutiusculis, leviter transversim convexus, aequaliter sat crebre punctatus, postice lineola longitudinali media subelevata laevissima, griseo-pubescentibus, aeneus, nitidus. Scutellum nigrum, fusco-tomentosum. Elytra thoracis fere latitudine, subtiliter confertissime punctato-alutacea, sparsim subtilius punctulata, fusco-subaenea, opaca, densius fusco-griseo-pubescentibus. Abdomen subtiliter punctulatum, nigrum, supra densius fusco-pubescentibus, segmentis quatuor anterioribus macula minuta triangulari in medio baseos flavescenti-pubescente, subtus nitidulum, metallico-versicolor, parcius pubescens, segmentis singulis basi tenuiter griseo-pubescentibus. Pectus nigrum, cinereo-pubescentibus. Pedes nigri, tarsi piceis.

81. *St. praelongus*: *Niger, capite, thorace elytrisque obscure viridi-subaeneis, illis glabris, crebre punctatis, linea media laevi, his abdomineque fusco-pubescentibus, scutello atro-tomentoso. — Long. 7 lin.*

*Mannerh. Brachelytr. 24. 21. — Nordm. Symb. 73. 69.*



Habitat in America septentrionali, collect. Dom. Chevrolat.

Statura omnino praecedentis. Antennae capite vix longiores, articulo secundo tertioque leviter elongatis, hoc longiore, sequentibus sensim paulo brevioribus, quarto oblongo, penultimis vix transversis, ultimo subovato, apice inferne acuminato, nigrae. Palpi nigro-picei. Caput thorace angustius, subtriangulare, dense sat fortiter punctatum, linea media longitudinali laevi, supra obscure viridi-aeneum, nitidulum, tenuiter nigro-pilosellum, ore nigro, subtus nigrum, nitidum, glabrum. Thorax coleopteris angustior, latitudine paulo longior, lateribus subrectis, parallelis, basi rotundatus, angulis posterioribus obtusis, apice truncatus, angulis anterioribus rectis, modice transversim convexus, supra dense sat fortiter punctatus, linea media longitudinali sat lata subelevata laevi, obscure viridi-aeneus, nitidus. Scutellum macula rotundata atro-tomentosa notatum. Elytra thoracis latitudine, obscure aenea, opaca, subtiliter confertissime punctato-rugulosa, tenuiter nigro-pubescentia. Abdomen crassiusculum, supra nigrum, subnitidum, subtiliter punctatum, tenuiter fusco-griseo-pubescent, supra bifariam e pube densiore atro-punctatum, subtus rufo-piceum, metallico versicolor, tenuiter pubescens. Pectus nigrum, griseo-pubescent. Pedes nigri.

82. *St. cyanomelas*: *Niger, nitidus, capite, thorace elytrisque supra cyaneis, nitidis, tenuiter nigro-pubescent, scutello densius atro-tomentoso.* — Long, 7 lin.

Habitat in Mexico, collect. Dom. Chevrolat.

Statura omnino *St. lutarii*, niger, nitidus, capite, thorace elytrisque supra cyaneis, nitidis, abdomine metallico-versicolore, totus tenuiter subtiliterque nigro-pubescent, Antennae capite paulo longiores, articulo tertio modice elongato, secundo sesqui longiore, sequentibus sensim paulo longioribus, quarto oblongo, penultimis 3 — 4 leviter transversis, ultimo apice leviter emarginato et inferne obtuse acuminato, nigrae. Caput thorace paulo angustius, antrorsum subangustatum, sat dense fortius punctatum, ore nigro, subtus nigrum, nitidum, glabrum. Thorax coleopteris parum angustior, latitudine vix brevior, antrorsum vix angustatus, lateribus rectis, basi cum angulis posterioribus fortiter rotundatus, apice truncatus, angulis anterioribus rectis, supra parum convexus, sat dense aequaliter punctatus. Scutellum pube densa depressa nigra obtectum. Elytra thoracis latitudine, subtiliter punctulata, interstitiis subtilissime alutaceis. Abdomen parcius subtiliter punctatum. Pedes nigri.

- \* 83. *St. violaceus*: *Niger, subnitidus, capite thoraceque subcyaneis, elytris violaceis, abdomine supra anterieus griseo-pubescente.* — Long, 5 — 6 lin.

*Grav. Micr.* 162. 6. *Mon.* 110, 115. — *Nordm. Symb.* 48. 30.

Habitat in America septentrionali, Knoch.

Antennae capite sesqui longiores, graciliores, articulo secundo tertioque leviter elongatis, hoc longiore, 4 — 10 subobconicis, sensim brevioribus, omnibus crassitie haud brevioribus, ultimo apice leviter emarginato et inferne acuminato, nigrae. Palpi picei. Corpus nigrum, capite thoraceque subcyaneis, nitidulum. Caput thoracis latitudine, subquadratum, tenuiter sub-

tiliterque nigro-pubescent, subdepressum, supra dense fortius punctatum, linea longitudinali tenuissima laevi, antice oblitterata, ore nigro, subtus parce punctatum, nigrum, nitidum, glabrum. Thorax coleopteris paulo angustior, latitudine paulo brevior, lateribus levissime, basi cum angulis posterioribus fortiter rotundatus, apice truncatus, angulis anterioribus rectis, subprominulis, supra modice transversim convexus, dense fortius punctatus, linea tenui longitudinali laevi, subtiliter tenuiterque nigro-pubescent, lateribus longius nigro-pilosellus. Scutellum nigrum, opacum, confertissime punctatum, apice atro-tomentosum. Elytra thorace paulo longiora, subtilissime punctato-rugulosa, violacea, opaca, densius subtiliter nigro-pubescentia, immixta pube rara brevi grisea. Abdomen parcius punctulatum, nigro-pubescent, segmentis tribus primis supra pube densiore grisea micante vestitis, reliquis parcius nigro-pubescentibus. Pedes graciliores, nigri.

- \* 84. *St. chrysaster*: *Niger, capite, thorace elytrisque supra coeruleis, nitidulis, abdomine apice testaceo, segmentis ultimis duobus supra aureo-tomentosis. — Long. 5 lin.*

Habitat in Brasilia, Dom. Prof. Germar.

Statura omnino *St. latebricolae*, niger, nitidus, capite, thorace elytrisque supra coeruleis aut virescenti-coeruleis, nitidulis. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 6 — 10 transversis, subaequalibus, ultimo ovato, apice truncato et inferne acuminato, nigrae. Caput thorace latitudine, subquadratum, supra subdepressum, sat profunde minus confertim punctatum, parce subtiliterque nigro-pubescent, ore nigro, subtus nigrum, nitidum, laeve glabrumque. Thorax coleopteris paululum angustior, latitudine vix brevior, lateribus parallelis, rectis, basi rotundatus, apice truncatus, angulis anterioribus rectis, posterioribus obtusis, subrotundatis, parum convexus, sat fortiter dense punctatus, carinula postica laevi, parce subtiliterque nigro-pubescent. Scutellum nigrum, confertissime punctulatum, nigro-pubescent. Elytra thorace paulo longiora, subtilissime alutacea, subtiliter punctulata, tenuiter subtiliterque nigro-pubescentia. Abdomen nigrum, nitidum, parce pubescens, segmento quinto margine apicali, sexto toto testaceis, his duobus supra totis dense aureo-tomentosis. Pedes nigri.

Species mihi invisae, forte huius familiae.

85. *St. femoratus*: *Nigro-subaeneus, pube obscura nebulosus, ano femoribusque anterioribus testaceo-ochraceis, his macula nigra notatis. — Long. 6 — 6½ lin.*

*Fab. Syst. El. II. 594. 24. — Grav. Mon. 122. 143. — Nordm. Symb. 31. 11.*

Habitat in Carolina.

Corpus subaeneum ut *St. murino*. Antennae capite paulo longiores, fuscae, articulis duobus primis rufis. Palpi articulis nigris, apice rufis. Caput thorace minus, subtriquetrum, linea media longitudinali laevi obsoleta. Thorax capitis colore atque punctura, linea media laevi longitudinali, angulis anticis obtusioribus quam in *St. murino*, paulo acutioribus quam in *St. pubescente*. Coleoptera subgriseo-nebulosa, subquadrata. Abdomen pilosellum, nigrum, dorso subrufo-varium, ventre subsericans, sub-aeneo-ni-

tens, segmento ultimo ochraceo. Pectus subsericans. Pedes nigricantes, femoribus ochraceis, lateribus, imprimis femorum posticorum nigris. (sec. Grav.)

Obs. Simillimus videtur *St. antiquo*, at praecipue antennis basi rufis pedumque colore distinctus.

86. *St. chrysocomus*: *Niger, opacus, pubescens, subtiliter punctulatus, thorace violaceo-micante, antennis, capite, elytris pedibusque fulvo-aureis.* — Long. 6 — 7 lin.

*Mannerh. Brachelytr. 21. 1. — Nordm. Symb. 41. 23.*

*Staph. chrysopterus Gebl. Nouv. Mem. d. l. Soc. Imp. des Nat. de Mosc. II. 41. 1.*

Habitat in Dauria (Mannerh.), prope Nertschinsk Sibiriae occidentalis (Gehl.).

*St. xanthoptero* vix angustior, sed multo brevior. Caput thorace multo latius, subquadrangulare, creberrime punctulatum, fulvum, pilis aureo-micantibus dense adpersum. Antennae thorace breviores, fulvae. Thorax antice parum lator, latitudine longior, antice truncatus, lateribus fere rectis, niger, subtiliter punctulatus, violaceo-micans, parce pubescens. Scutellum rotundatum, holosericeo-atrum. Elytra thorace latiora, subquadrata, angulo exteriori rotundata, subtilissime punctulata, fulva, dense aureo-sericea. Abdomen nigrum, tenue pubescens, segmentorum ventralium marginibus aureo-micantibus. Pedes fulvi, aureo-sericei. (Mannerh.)

Obs. *Staph. chrysopterus* Gebler thorace nigro-aeneo, linea longitudinali in medio baseos, abdominis segmento secundo basi aureo-pubescente, coxis nigris, a specie *Mannerheimiana* haud diversus videtur.

87. *St. Herculeanus*: *Cyaneus, tomentosus, subtilis subvirescens, thorace angulis anticis fortiter productis.* — Long. 15 lin.

*Laporte Etud. Ent. I. 114. 2. t. 3. f. 1.*

Habitat in Brasilia.

Cyaneo-subviolaceus, tomentosus. Antennae nigrae. Thorax basi rotundatus, lateribus subrectis, angulis anterioribus productis, rotundatis, margine antico medio rotundato, utrumque profunde sinuato, supra sulco longitudinali medio posteriore distincte impressus. Elytra subtilissime granulata. Corpus subtus et pedes subvirescentia. (Lap.)

Obs. Forte prope *St. Buquetii* inserendus. Speciebus Fam. X. corpore tomentoso alienus videtur.

88. *St. lugubris*: *Thorace toto punctato, ater, atro-pubescent.* — Long. 15 lin.

*Encycl. Meth. X. 477. 2.*

Habitat in Cayenna.

Maximus, totus niger, opacus, pube brevi concolore obductus, infra subnitidus. Caput angulis posterioribus prominulis. (Enc.)

89. *St. haemorrhoidalis*: *Nigro-violaceus, abdomine apice rufo.*

*Oliv. Ent. III. 42. 11. 7. t. 2. f. 20.*

Habitat in Cayenna.



Magnitudo Staph. erythrocephali. Antennae nigrae, capite paulo longiores. Caput, thorax et elytra nigro-violacea, nitida, nigro-pubescentia. Abdomen nigrum, apice rufum. Pedes nigri. Alae nigricantes. (Ol.)

90. *St. Gorgi*: Nigro-subaeneus, pubescens, scutello atro-tomentoso, elytris viridi-pubescentibus, macula gemina basali punctisque duobus marginalibus atro-tomentosis. — Long. 11 lin.

Laporte Etud. Ent. I. 113. 1. t. 3. f. 2.

Habitat in India orientali. (Decan.)

Nigro-subaeneus, confertissime punctatus, pubescens. Thorax brevis, linea media longitudinali subelevata. Scutellum nigrum, tomentosum. Elytra laete viridia, pube concolore vestita, macula interrupta baseos punctisque duobus marginis exterioris atro-tomentosis. Alae violaceae. Corpus subtus et pedes nigra, tarsis rufescentibus. (Lap.)

### Fam. IX.

*Antennae filiformes, articulo ultimo apice emarginato.*

*Palpi omnes filiformes, maxillares articulo quarto tertio longiore.*

*Pedes graciliores, tibiis posterioribus spinulosis, tarsis anticis dilatatis.*

Corpus parcius pubescens, capite thoraceque parce pilosellis. Antennae sat elongatae, apicem versus haud incrassatae, articulo primo sequentibus ambobus multo brevioribus, crassiusculo, his elongato-obconicis, 4 — 10 leviter obconicis, crassitie longioribus, ultimo oblongo-ovato, apice oblique emarginato et inferne acuminato, tenuiter pilosellae. Caput rotundatum, supra leviter convexum, oculis mediocribus, leviter convexis. Labrum breve, limbo membranaceo minus dilatato. Mandibulae apice falcatae, acutae, basin versus sensim latiores, muticae. Palpi modice elongati, maxillares articulo secundo elongato, tertio hoc duplo prope brevioribus, quarto iterum elongato, secundo aequali, apice subacuminato, labiales articulo primo brevioribus, secundo tertioque aequalibus, illo interne parce setuloso, hoc apice iam subacuminato (*St. luridipes*, anceps, terminalis,) iam truncato (*St. segmentarius*, fasciatus). Thorax basin versus subangustatus. Elytra apice oblique truncata. Abdomen apicem versus leviter angustatum. Pedes graciles, modice elongati, tibiis posterioribus parce spinulosis, spinis apicalibus mediocribus, tarsis posterioribus articulo primo modice elongato, sequentibus tribus subaequalibus, anticis articulis 4 primis maris fortiter, feminae levius dilatatis, subtus dense longius tomentosus.

\* 91. *St. luridipes*: Niger, capite, thorace elytrisque aeneis, subopacis, pedibus fusco-luteis. — Long. 4 lin.

Habitat in Columbia, Dom. Moritz.

Antennae capitis thoracisque fere longitudine, articulo tertio secundo sesqui longiore, 4 — 10 subobconicis, sensim paulo brevioribus, penultimis etiam crassitie haud brevioribus, ultimo subovato, apice oblique truncato et inferne acuminato, nigrae. Palpi picci, apice rufescentes. Corpus nigrum, nitidum. Caput thoracis fere latitudine, orbiculatum, supra convexiusculum, profunde punctatum, fronte medio late laevi, antice inter antennas late le-

viterque impressa, obscure aeneum, parum nitidum, parce nigro-pilosellum, infra nigrum, nitidum, laeve. Thorax coleopteris dimidio angustior, latitudine paululum longior, basin versus modice angustatus, lateribus ante medium leviter rotundatis, dein sinuatis, basi rotundatus, apice truncatus, angulis posterioribus obtusis, subrotundatis, anterioribus acutiusculis, parum convexus, obscure aeneus, opacus, utrinque subtilius punctatus, puncto quoque pilum nigrum erectum emittente, linea media longitudinali laevi, nitidiore. Scutellum obscure aeneum, dense punctatum, flavescenti-pubescentia. Elytra thorace vix longiora, dense punctata, obscure aenea, subnitida, basi apiceque flavescenti-pubescentia. Abdomen nigrum, nitidum, concolor, punctulatum, parcius nigro-pubescentia. Pedes obscure lutei, coxis fuscis. Alae infuscae.

\* 92. *St. anceps*: *Niger, capite, thorace elytrisque aeneis, opacis, abdominis segmento penultimo rufo, pedibus luteis.* — Long. 4 lin.

Habitat in Columbia, Dom. Moritz.

Statura et summa affinitas praecedentis. Antennae capite thoraceque paulo longiores, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus, penultimis etiam crassitie longioribus, 6 — 10 leviter obconicis, ultimo oblongo-ovato, apice truncato et inferne acuminato, nigrae, articulo ultimo ferrugineo, primo subtus rufo-piceo. Palpi picei, articulo ultimo testaceo. Mandibulae rufo-picei. Corpus nigrum, nitidum. Caput thoracis latitudine, orbiculatum, supra convexusculum, fortiter punctatum, fronte medio laevi, antice inter antennas leviter impressa, obscure aeneum, subnitidum, parce nigro-pilosellum, infra nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine vix longior, basin versus subangustatus, lateribus anterieus subrotundatis, posterius leviter sinuatis, basi rotundatus, apice truncatus, angulis posterioribus obtusis, subrotundatis, anterioribus acutiusculis, supra parum convexus, obscure aeneus, opacus, utrinque minus fortiter punctatus, puncto quoque pilum subtilem nigrum emittente, linea media longitudinali laevi. Scutellum subtiliter punctulatum, dense flavescenti-pubescentia. Elytra thorace paulo longiora, dense punctata, obscure aenea, subnitida, fusco-, basi apiceque flavescenti-pubescentia. Abdomen punctulatum, parce longius nigro-pubescentia, nigrum, nitidum, segmento quinto rufo-testaceo, basi nigro, subtus etiam segmentis 4 primis margine apicali testaceis. Pedes lutei, coxis fuscis.

\* 93. *St. terminalis*: *Niger, capite thoraceque nigro-subaeneis, opacis, ano rufo-testaceo, pedibus piceis.* — Long. 4 lin.

Habitat in Brasilia, Dom. Virmond.

Statura et summa affinitas praecedentium. Corpus nigrum. Antennae capite thoraceque paulo longiores, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo brevioribus, penultimis etiam crassitie paulo longioribus, ultimo ovato, apice oblique truncato et inferne acuminato, totae nigrae. Palpi, mandibulae et labrum picea. Caput thoracis latitudine orbiculatum, supra parum convexum, nigro-subaeneum, nitidum, fortiter punctatum, fronte medio laevi, antice late leviter impressa, parce nigro-pilosellum, basi utrinque griseo-pubescentia, subtus nigrum, nitidum. Thorax

coleopteris dimidio angustior, latitudine paulo longior, basin versus subangustatus, lateribus anterieus leviter rotundatis, dein subsinuatis, basi rotundatus, apice truncatus, angulis posterioribus obtusis, anterioribus acutiusculis, parum convexus, nigro-subaeneus, opacus, utrinque minus fortiter punctatus, puncto quoque pilo erecto nigro instructo, linea media longitudinali laevi, subnitida. Scutellum dense punctulatum, flavescenti-pubescentia. Elytra thorace parum longiora, dense punctata, parum nitida, dense nigro-, basi anguste flavescenti-pubescentia. Abdomen segmentis singulis basi confertim subtiliter punctatis, nigro-pubescentia, segmenti quinti parte dimidia apicali segmentoque sexto toto rufo-testaceis, subtus segmentis reliquis margine apicali rufo-piceo. Pedes testaceo-picei.

\* 94. *St. fasciatus*: Capite, thorace scutelloque nigro-aeneis, parum nitidis, elytris pectoreque dilute rufis, abdomine rufo-nigroque fasciato, pedibus testaceis. — Long. 4 lin.

*Philonthus fasciatus* Nordm. Symb. 112. 102.

Habitat in Mexico, Dom. Deppe.

Praecedenti proximus. Antennae capite thoraceque longiores, articulo tertio secundo sesqui longiore, 4—10 sensim paulo brevioribus, penultimis etiam crassitie longioribus, 6—10 leviter obconicis, ultimo subovato, apice oblique emarginato et inferne acuminato, apice subferrugineae, articulo primo subtus rufo-piceo. Palpi piceae, apice testaceae. Mandibulae rufo-piceae. Caput thorace paululum latius, orbiculatum, supra leviter convexum, obscure aeneum, nitidulum, fortiter punctatum, puncto quoque pilum longum depressum griseum emittente, fronte medio late laevi, antice late levissime impressa, parce nigro-pilosellum, subtus nigrum, nitidum. Thorax coleopteris dimidio prope angustior, latitudine vix longior, basin versus subangustatus, lateribus anterieus leviter rotundatis, pone medium subsinuatis, basi rotundatus, apice truncatus, angulis posterioribus obtusis, subrotundatis, anterioribus obtusiusculis, subdepressus, obscure aeneus, subopacus, punctatus, puncto quoque pilo erecto nigro instructo, linea media longitudinali laevi, subnitida. Scutellum nigro-aeneum, dense punctulatum, flavo-pubescentia. Elytra thorace paulo longiora, dense punctata, dilute rufa, flavo-pubescentia. Abdomen parcius punctulatum, pubescens, supra segmentis duobus, subtus tribus primis rufis, reliquis nigris, margine apicali rufo, ultimo nigro, basi rufo. Pectus rufum. Pedes testacei, coxis rufis.

\* 95. *St. segmentarius*: Capite thoraceque aeneis, opacis, scutello, elytris pectoreque rufis, abdomine rufo-nigroque fasciato, pedibus testaceis. — Long. 4 lin.

Habitat in Columbia, Dom. Moritz.

Statura et summa affinitas praecedentium. Antennae capitis thoracisque longitudine, articulo tertio secundo sesqui longiore, sequentibus subobconicis, sensim paulo brevioribus, penultimis etiam crassitie longioribus, ultimo ovato, apice leviter emarginato et inferne acuminato, nigro-piceae. Palpi picei, apice rufi. Mandibulae rufae. Caput thoracis latitudine, orbiculatum, supra convexusculum, fortiter punctatum, fronte medio laevi, antice inter antennas late leviter impressa, nigro-aeneum, sub-opacum, parce nigro-



pilosellum, subtus nigrum, nitidum. Thorax coleopteris dimidio fere angustior, latitudine paululum longior, basin versus angustatus, lateribus anterioribus leviter rotundatis, posterius subsinuatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrectis, parum convexus, nigro-aeneus, opacus, utrinque minus fortiter punctatus, puncto quoque pilo nigro instructo, linea media longitudinali laevi, subnitida. Scutellum rufum, subtiliter punctulatum, flavo-pubescentia. Elytra thorace parum longiora, densius punctata, dilute rufa, opaca, fulvo-pubescentia. Abdomen subtiliter punctulatum, segmento primo rufo, sequentibus tribus rufis, macula transversa basali, sensim maiore, nigra, quinto nigro, apice rufo, sexto toto nigro. Pectus rufum, mesosterno piceo. Pedes cum coxis dilute testacei. Alae infuscae.

### Fam. X.

*Triacrus Nordm. Symb. p. 19.*

*Antennae ab articulo quinto inde subserratae, articulo ultimo tricuspi. Palpi omnes filiformes, maxillares articulis ultimis duobus subaequalibus. Pedes tibiis posterioribus spinulosi, tarsis anticis dilatatis.*

Corpus tenuiter, caput et thorax supra tenuissime subtilissimeque pubescentia. Antennae breviores, basi satis distantes, articulo primo modice elongato, secundo obconico, tertio hoc sesqui longiore, elongato-obconico, quarto cyathiformi, apice longitudine prope crassiore, sequentibus crassioribus, brevioribus, perfoliatis, deorsum fortius prominentibus ideoque subserratis, ultimo tripartito, portione superiore maiore porrecta, inferioribus minoribus, inter se aequalibus, leviter deflexis, omnibus acuminatis. Caput porrectum, sat magnum, orbiculatum, supra parum convexum, oculis minoribus, vix prominulis. Labrum minutum, transversum, membranaceo-limbatum. Mandibulae validae, apice incurvae, acutae, medio bidentatae. Palpi minus elongati, maxillares articulo secundo tertioque aequalibus, quarto his longitudine aequali, at paulo tenuiore, apice subacuminato, labiales articulis sensim paulo longioribus. Thorax coleopteris angustior. Elytra angulo apicali exteriori oblique truncato. Abdomen apicem versus sensim attenuatum. Pedes breviores, validi, tibiis posterioribus subtiliter spinulosi, spinis apicalibus validioribus, tarsis posterioribus articulis 4 primis sensim paulo brevioribus, primo leviter elongato, anticis articulis 4 primis dilatatis, patellam ovalem, tibia parum latiore, subtus tomentoso-spongiosam formantibus.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso.

\*96. *St. superbus*: *Niger, capite thoraceque cyaneis, elytris violaceis, ano rufo, antennis subserratis. — Long. 13 lin.*

*Triacrus dilatus Nordm. Symb. 19. 1. t. 1. f. 7.*

Habitat in Brasilia, Dom. v. Langsdorf.

Magnus. Corpus nigrum, nitidulum. Antennae capite sesqui longiores, nigrae, articulis tribus primis obconicis, quarto subgloboso, 4 — 10 transversis, perfoliatis, introrsum magis quam extrorsum dilatatis ideoque subserratis, ultimo excavatione profunda tripartito, lobis duabus basalibus

minoribus, aequalibus, deorsum spectantibus. Mandibulae nigrae. Caput thorace paulo latius, suborbiculatum, supra subdepressum, cyaneum, subnitidum, vage punctulatum, parce subtilissime pubescens, subtus nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine brevior, lateribus medio leviter rotundatis, posterius subsinuatis, basi modice emarginatus, angulis posterioribus obtusiusculis, apice truncatus, utrinque subsinuatus, angulis anterioribus acutiusculis, supra modice transversim convexus, cyaneus, subnitidus, subtiliter vage punctulatus, parce subtilissime nigro-pubescens. Scutellum nigrum, punctatissimum, subtiliter nigro-pubescens. Elytra thorace longiora, densius subtiliter punctato-subrugulosa, violacea, subtiliter nigro-pubescentia. Abdomen punctulatum, tenuiter nigro-pubescens, segmento ultimo cum stylis analibus rufo, rufo-pubescente. Pedes nigri, nigro-pubescentes.

Variat capite thorace maiore, mandibulis capite longitudine aequalibus, et capite thorace aequali, mandibulis capite brevioribus.

Obs. Prof. Nordmann hanc speciem in collectione Langsdorffiana nomine Staph. dilati Leach signatam fuisse tradit: revera autem loco illo a Latreillio huic insecto eximio tantummodo nota fuit affixa, qua Staph. dilatatus F. ad animum revocetur, quocum antennis serratis comparetur, et cui Dom. Leach genus Velleius condidisset. Nomen igitur illud, tali errore huic speciei impertitum, delendum putavi.

### Fam. XI.

*Trigonophorus Nordm. Symb. p. 8. t. 1. f. I.*

*Antennae apicem versus subincrassatae, articulo ultimo apice emarginato.*

*Palpi maxillares filiformes, articulo quarto tertio subaequali, apice acuminato, labiales articulo tertio subsecuriformi.*

*Pedes validi, tibiis posterioribus spinulosis, tarsis anticis dilatatis.*

Corpus subtiliter pubescens, capite thoraceque glaberrimis. Antennae breviores, articulo primo parum elongato, secundo tertioque subobconicis, penultimis 5 transversis, ultimo apice emarginato et inferne acuminato, articulis 4 primis breviter pilosellis, reliquis pube brevissima subtilissimaque cinerea dense obductis. Caput porrectum, sat magnum, thoracis prope latitudine, oculis parvis, oblongis, vix prominulis. Labrum mediocre, transversum, angustius coriaceo-limbatum, limbo barbato. Mandibulae breviores, validae, apice falcatae, acutae, medio fortiter dentatae. Palpi maxillares articulo secundo paulo longiore, quarto tertio fere aequali, apice acuminato, labiales articulis tribus sensim longioribus, ultimo apicem versus magis minusve incrassato, apice truncato, leviter securiformi. Thorax coleopteris paulo angustior. Elytra angulo apicali exteriore rotundato. Pedes breviusculi, validi, tibiis intermediis leviter curvatis, posterioribus spinulosis, spinis apicalibus validioribus, tarsis posterioribus articulo primo modice elongato, sequentibus tribus sensim paulo minoribus, anticis articulis quatuor primis dilatatis, patellam ovalem, subtus dense tomentoso-spongiosam formantibus.

Mas abdominis segmento inferiore sexto apice leviter inciso distinctus.

Obs. Habitus specierum partim Fam. VIII. partim Fam. II., a quibus palporum labialium articulo ultimo apice truncato, subsecuriformi distinguuntur. Caput et thorax laevissima, reliquum corpus parce et subtilissime pubescens.

- \* 97. *St. columbinus*: *Cyaneus, nitidus, antennis pedibusque nigris, thorace laevissimo, margine laterali punctatissimo.* — Long. 13 lin.

Habitat in Brasilia, Dom. Virmond.

Corpus magnum, cyaneum, nitidum. Antennae capite vix longiores, articulo tertio secundo paulo longiore, quarto quintoque subglobosis, 6—10 transversis, sensim paulo brevioribus, ultimo penultimo plus sesqui longiore, apice truncato et inferne acuminato, nigrae, basi cyanescentes. Palpi labiales articulo ultimo cylindrico, apice truncato. Oris partes nigrae. Caput thoracis latitudine, ovatum, supra leviter convexum, oculis parvis, ante medium sitis, laevissimum, lateribus prope basin crebre punctatum. Thorax coleopteris paulo angustior, latitudine fere brevior, lateribus leviter rotundatus, posterius subsinuatus, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus antrorsum prominentibus, acutis, modice convexus, laevissimus, limbo, basi apiceque medio interrupto, punctatissimo. Scutellum dense subtiliter punctatum, nigro-pubescent. Elytra thorace longiora, violacea, subopaca, confertissime subtiliter subruguloso-punctata, pube brevi nigra vestita. Abdomen confertissime subtiliter punctatum, nigro-pubescent. Pedes nigri. Alae fusco-violaceae.

- \* 98. *St. myrtillinus*: *Cyaneus, nitidus, antennis pedibusque nigris, capite thoraceque parce subtiliterque punctatis.* — Long. 10 lin.

*Trigonophorus myrtillinus* Nordm. Symb. 8. 1. t. 1. f. 1.

Habitat in Brasilia, Dom. v. Olfers.

Habitu omnino praecedentis, at duplo minor, cyaneus, nitidus. Antennae capite vix longiores, articulo tertio secundo longiore, quarto subgloboso, 5—10 transversis, sensim paulo brevioribus, ultimo subovato, apice emarginato et inferne acuminato, nigrae. Mandibulae et palpi nigri, palpi labiales articulo tertio leviter securiformi. Caput thoracis latitudine, subovatum, supra parum convexum, parce subtiliterque punctatum. Thorax coleopteris angustior, latitudine vix longior, basin versus subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus antrorsum productis, acutis, modice transversim convexus, undique disperse subtiliterque parce punctatus. Scutellum subtiliter punctatum, nigro-pubescent. Elytra thorace paululum longiora, subtiliter punctata subtiliterque nigro-pubescentia. Abdomen subtilius punctatum, parce subtiliterque nigro-pubescent. Pedes nigro-subcyanei, nigro-pubescentes.

- \* 99. *St. badiipennis*: *Laete viridis, nitidus, pedibus cyaneis, elytris luteis.* — Long. 8 lin.

*Creophilus badiipennis* Nordm. Symb. 25. 6.

Habitat in Brasilia, Dom. v. Olfers.

Statura fere *St. maxilloso*. Antennae capite haud longiores, articulo secundo tertio paululum maiore, his obconicis, quarto quintoque subglobosis, 6—10 paulo latioribus et sensim paulo brevioribus, transversis, ultimo subovato, apice inferne leviter emarginato, nigrae. Mandibulae nigrae. Palpi et labrum rufo-picea, palpi labiales articulo tertio leviter securiformi. Cor-



pus lucide viride, nitidum, abdomine apice sub-coerulescente. Caput thorace paulo angustius, subovatum, supra parum convexum, punctis paucis subtilibus adpersum, nitidissimum, subtus nigrum. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus rectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, utrinque subsinuatus, angulis anterioribus acutiusculis, leviter transversim convexus, nitidissimus, apicem versus punctis raris subtilibus impressus. Scutellum punctatum, viridi-cyaneum, tenuiter luteo-pubescentia. Elytra thorace parum longiora, subtiliter punctata, lutea, tenuiter luteo-pubescentia. Abdomen crebrius subtiliter punctulatum, subtiliter nigro-pubescentia. Pectus cyaneo-nitidum. Pedes cyanei, nitidi, tibiis nigris, tarsis piceis.

\* 100. *St. Chloris*: *Laete viridi-metallicus, nitidus, elytris cyaneis, pedibus nigris.* — Long. 6 lin.

*Creophilus Chloris* Nordm. Symb. 26. 8.

Habitat in Brasilia, Dom. v. Olfers.

Habitu omnino *St. maxilloso*, laete lucideque metallice viridis, nitidus. Antennae capitis longitudine, articulo secundo tertioque obconicis, aequalibus, quarto subgloboso, quinto leviter transverso, 6 — 10 fortius transversis, sensim paulo brevioribus et latioribus, ultimo ovato, apice leviter acuminato, nigrae. Os nigrum. Palpi labiales articulo ultimo cylindrico, apice truncato. Caput thorace paulo angustius, ovatum, leviter convexus, supra oculos utrinque punctis parvis impressum. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus subrectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, utrinque subsinuatus, angulis anterioribus rectis, modice transversim convexus, laevissimus. Scutellum cyaneum, subtiliter punctulatum, subtiliter pubescens. Elytra thorace paulo longiora, laete cyanea, subnitida, parce minus subtiliter punctata, subtiliter nigro-pubescentia. Abdomen punctulatum, tenuiter nigro-pubescentia. Pedes nigri.

\* 101. *St. Erithacus*: *Viridi-coerulescens, nitidus, elytris cyaneis, ano rufo, pedibus nigris.* — Long. 6 lin.

*Creophilus Erithacus* Nordm. Symb. 25. 7.

Habitat in Brasilia, Dom. v. Olfers.

Habitu omnino praecedentium. Laete viridi-coerulescens, nitidus. Antennae capite vix longiores, nigrae, articulo secundo tertioque aequalibus, quarto transversim obconico, quinto transverso, 6 — 10 abrupte paulo crassioribus, transversis, inter se aequalibus, ultimo ovato, apice compresso, acuminato. Palpi rufo-picei, labiales articulo ultimo cylindrico, apice truncato. Caput thorace paulo angustius, apicem versus subangustatum, pone oculos utrinque punctis paucis sparsis impressum. Thorax coleopteris parum angustior, latitudine vix brevior, lateribus rectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, utrinque subsinuatus, angulis anterioribus acutiusculis, leviter transversim convexus, laevissimus, glaberrimus. Scutellum cyaneum, subtiliter punctulatum, subtiliter tenuiterque nigro-pubescentia. Elytra thorace paulo longiora, parce subtiliterque punctata, cyanea, opaca, parce subtiliterque nigro-pubescentia. Abdomen pun-

ctulatum, parce subtiliterque nigro-pubescent, segmento ultimo rufo. Pedes nigri, femoribus subviolaceis.

Obs. Prope ad Fam. II. accedit habitu, palporum maxillarum articulo ultimo brevior, acuminato, labialium articulo tertio apice haud latiore, discedit autem antennarum structura, capite maris neque thorace latiore neque subquadrato, et elytris abdomineque parce et vix perspicue pubescentibus.

\* 102. *St. purpurascens*: *Niger, capite thoraceque purpureo-cupreis, nitidis, elytris cyaneis, opacis.* — Long.  $7\frac{1}{2}$  lin.

Nordm. Symb. 47. 21.

Habitat in Brasilia, Dom. v. Langsdorf.

Corpus elongatum, nigrum, subnitidum. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus, penultimis transversis, ultimo subgloboso, apice inferius acuminato, nigrae. Palpi picei, labiales articulo tertio crassiusculo, cylindrico, apice truncato. Mandibulae nigrae. Caput et thorax supra purpureo-cuprea, splendida. Caput thoracis latitudine, subquadratum, supra depressum, versus margines punctis parvis impressum, subtus nigrum, nitidum. Thorax coleopteris angustior, basin versus subangustatus, latitudine longior, lateribus pone medium subsinuatis, basi rotundatus, angulis posterioribus obtusis, apice truncatus, angulis anterioribus subrectis, leviter transversim convexus, intra angulos anteriores punctis nonnullis, dorso ante medium punctis quatuor transversim positos impressus. Scutellum nigrum, punctis paucis impressum. Elytra thorace paululum longiora, subtilissime rugulosa, subtiliter punctata, opaca, saturate cyanea, parce tenuiterque nigro-pubescentia. Abdomen parce punctulatum, parce nigro-pubescent. Pedes minus validi, nigri. Alae fulvo-hyalinae.

## Fam. XII.

*Tympanophorus* Nordm. Symb. p 9. t. 1. f. 2.

*Antennae apicem versus sensim subincrassatae, articulo ultimo apice emarginato.*

*Palpi maxillares articulo ultimo penultimo brevior, acuminato, labiales articulo ultimo hemisphaerico, apice truncato.*

*Pedes breviusculi, tibiis omnibus dense spinulosis, tarsis anticis simplicibus.*

Corpus punctatum, tenuiter pubescens. Antennae capite plus sesqui longiores, articulo primo leviter elongato, secundo tertioque aequalibus, obconicis, sequentibus subcylindricis, sensim paulo brevioribus et crassioribus, ultimo transverso, apice truncato et inferne acuminato, articulis 4 primis tenuiter pilosellis, reliquis pube brevissima grisea dense obductis. Caput porrectum, orbiculatum, thorace minus. Labrum breve, transversum, membranaceo-limbatum. Mandibulae breviores et debiliores, medio dentatae. Palpi maxillares articulo secundo tertioque subaequalibus, quarto multo minore, tenuiore, fortiter acuminato, fere subulato, labiales articulis duobus primis minutis, tertio valde incrassato, hemisphaerico, apice truncato, tympaniformi. Thorax coleopteris angustior, angulis anterioribus deflexis. Elytra angulo apicali exteriori oblique truncato. Abdomen crassiusculum. Pedes breviusculi, tibiis omnibus densius breviter spinulosis, spinis apicalibus va-

Idioribus, tarsis posterioribus articulo primo modice elongato, sequentibus tribus sensim paulo brevioribus, ultimo minus elongato, praecedentibus ambobus haud longiore, anticis feminae simplicibus.

Mas latet.

\*103. *St. canaliculatus*: *Niger, opacus, capite, thorace elytrisque dense punctatis.* — *Long. 6 lin.*

*Tympanophorus canaliculatus Nordm. Symb. 10. 1.*

Habitat in Brasilia, Dom. v. Olfers.

Niger, parum nitidus, capite nigro-subaeneo. Antennae capite plus sesqui longiores, piceae, articulis 6 — 10 transversis. Caput thorace angustius et multo minus, suborbiculatum, subdepressum, fortius crebre punctatum, griseo-subpubescens. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus et basi leviter rotundatus, apice truncatus, angulis posterioribus rotundatis, anterioribus obtusiusculis, parum convexus, sat crebre fortius punctatus, linea longitudinali media laevi, postice canaliculata. Scutellum subtiliter punctulatum, griseo-pubescens. Elytra thorace paulo longiora, dense punctata, cinereo-pubescentia. Abdomen crassiusculum, nitidum, subtiliter punctulatum, griseo-pubescens, subtus segmentis singulis apice tenuiter piceo-marginatis. Pedes nigri. Alae fuscae, costa nigra.

### **Ocypus Kirby.**

*Er. Col. March. I. p. 439.*

*Antennae rectae.*

*Palpi maxillares filiformes.*

*Ligula apice emarginata.*

*Pedes intermedii basi approximati.*

*Tarsi posteriores teretes.*

Corpus maioris magnitudinis, elongatum, lineare, aliis alatum, aliis apterum. Caput sat magnum, thoracis circiter magnitudine, suborbiculatum vel subquadratum, porrectum, exsertum, collo quasi brevi thoraci immissum, oculis parvis, haud prominulis. Labrum transversum, bilobum, corneum, lateribus latius, antice angustius membranaceo-limbatum. Mandibulae plerumque sat validae, aliis medio dentatae, aliis graciliores, falcatae, edentatae, basi lacinia elongata, lineari, inferne ciliata instructae. Maxilla malis brevibus, coriaceis, interiore interne, exteriori apice barbata, hac basi cornea. Palpi maxillares articulo primo minuto, secundo tertioque subaequalibus, quarto praecedenti aut aequali aut eodem parum breviori, apice aut subacuminato, aut plerumque truncatulo. Labium mento transversum, ligula membranacea, biloba, paraglossis membranaceis, interne ciliatis, ligulam paulo superantibus. Palpi labiales articulis duobus primis brevioribus, tertio aut cylindrico, apice truncato, aut leviter fortiusve securiformi. Antennae filiformes, articulo primo subelongato, articulis a quarto vel a quinto inde magnitudine decrescentibus, ideoque apicem versus subangustatae. Thorax coleopteris parum angustior, basi rotundatus, apice truncatus, angulis posterioribus obtusis, magis minusve rotundatis, anterioribus plerumque rectis, subdeflexis. Scutellum triangulare. Elytra apice extrorsum leviter rotun-



data, introrsum oblique truncata. Abdomen lineare. Pedes breviusculi, coxis intermediis contiguis, tibiis posterioribus spinulosis, tarsis posterioribus filiformibus, articulo primo leviter elongato, anticis articulis 4 primis dilatatis, subtus dense spongioso-tomentosis.

#### O b s e r v a t i o n e s.

1. Genus praecedenti admodum affine, praeter notas plures leviores praecipue ab illo distinctum coxis intermediis contiguis, acetabulis earum medio confluentibus.

2. Mas abdominis segmento inferiore sexto apice emarginato distinguendus.

3. Victus in silvis et arvis sub musco et sub lapidibus. Species omnes Europaeae, plures Alpêtres, singula simul in America septentrionali obvia.

4. Larva O. olentis observata est a Dom. Heer. (Obs. ent.) et a Dom. Blanchard (Guer. Mag. Zool.). Corpus eius elongatum, apicem versus sensim attenuatum. Caput sat magnum, rotundato-subquadratum. Antennae capite breviores, 4 articulae, articulo primo minuto, secundo tertio paulo longiore, apice interne processu aucto, quarto minuto, subulato. Clypeus apice denticulatus. Mandibulae sat magnae, falcatae. Maxillae cylindricae, processu minuto, acuminato, palpo 4-articulato, articulo primo brevissimo, crasso, secundo tertioque aequalibus, teretibus, quarto minuto, conico. Labium obconicum, apice medio acuminatum, palpis bi-articulatis, articulo primo terete, longiore, secundo conico. Prothorax capite paulo angustior, quadrangulus, latitudine dimidio brevior, lateribus leviter rotundatus. Meso- et metathorax transversus, prothoraci paulo latiores. Pedes breviusculi, tibiis brevioribus, spinulosis, tarsis unguiculo simplice. Abdomen segmentis sensim angustioribus, dorso areis binis corneis. Anus tubulosus, descendens. Caudae anales setaceae, 3-articulae. — Victitat in silvis, in terra, in cubiculis mandibularum ope effossis, mandibulas exserens, Cicindelorum larvarum more insecta, quae ei appropinquantur, arripens, lacerans, devorans, noctu interdum circumvagans. (Conf. Heer l. c. p. 18.) Nympha thoracis margine antico pilis longis 12 coronato.

Secundum antennarum, mandibularum palporumque structuram sectiones quatuor in hoc genere constituendae sunt, quae quodammodo familias naturales referunt, quarum autem notae pro characteribus genericis rite valere non videntur. Prima earum palpis labialibus filiformibus, ultima mandibulis muticis, secunda antennarum articulo ultimo integro, acuminato, tertia palporum maxillarium articulo ultimo in maribus subsecuriformi insignis.

#### Fam. I.

Goërius Leach Steph. *Illust. Britt. Ent. V.* p. 208.

*Trichoderma* Steph. *ibid.* p. 435.

*Antennae articulo ultimo apice emarginato.*

*Palpi labiales articulo ultimo cylindrico.*

*Mandibulae medio dentatae.*

Corpus elongatum, saepius apterum. Caput sat magnum, suborbiculatum, basi profunde constrictum, oculis parvis, haud prominulis. Antennae basi minus late distantes, apicem versus subattenuatae, articulo ultimo apice oblique truncato. Mandibulae breviores, validae, falcatae, acutae, medio dentatae. Palpi maxillares articulo quarto paulo minore, subacuminato, labiales articulis duobus primis brevioribus, ultimo cylindrico. Elytra saepius thorace breviora. Pedes tibiis intermediis densius, posticis parcius spinulosis, tarsis anticis fortius dilatatis.

Obs. Genus *Trichoderma* ex O. picipenni a Dom. Steph. conditum. Characterem elytra tessellata praebent.

- \* 1. *P. olens*: *Alatus, niger, opacus, supra subtilissime atro-tomentosus, antennis apice ferrugineis, elytris thoracis longitudine.* — Long. 11 — 13 lin.

*Er. Col. March. I.* 439. 1.

*Staph. olens* Müll. *Faun. Fridr.* 23. 228. *Prodr.* 97. 1090. — *Fab. Mant. Ins. I.* 219. 4. *Ent. Syst. I. II.* 520. 6. *Syst. El. II.* 591. 8. — *Goeze Ent. Beitr. I.* 726. 19. — *De Vill. Ent. I.* 424. 66. — *Payk. Mon. Staph.* 9. 4. *Faun. Suec. III.* 371. 5. — *Oliv. Ent. III.* 42. 9. 4. t. 1. f. 1. — *Cederh. Faun. Ingr.* 117. 358. — *Walken. Faun. Paris. I.* 270. 6. — *Panz. Ent. Germ.* 352. 5. *Faun. Germ.* 27. 1. — *Grav. Micr. A. 2. Mon.* 111. 117. — *Latr. Hist. nat. Crust. et Ins. IX.* 302. 16. — *Marsh. Ent. Britt.* 497. 2. — *Shaw General Zool. VI.* 108. t. 39. — *Gyll. Ins. Suec. II.* 285. 6. — *Mannerh. Brachelytr.* 23. 15. — *Nordm. Symb.* 63. 50. — *Runde Brachelytr. Hal.* 3. 8.

*Emus olens* *Boisd. et Lacord. Faun. Ent. Paris. I.* 367. 12.

*Staph. unicolor* *Herbst Archiv.* 149. 4.

*Staph. maxillosus* *Schrank Enum. Ins. Austr.* 230. 434. — *Fourcr. Ent. Par. I.* 164. 1.

*Staph. maior* *Degeer. Ins. IV.* 17. 3. — *De Vill. Ent. I.* 420. 38.

*Le grand Staphyl. noir lisse* *Geoffr. Hist. Ins. I.* 360. 1. t. 7. f. 1.

Habitat in Europa omni temperata meridionalique et in Sibiria, in Mauritania, Dom. Wagner, in Teneriffa, Dom. Richter.

Magnus, totus niger, opacus, tomento brevissimo nigro supra vestitus. Antennae capite sesqui longiores, articulo tertio secundo vix longiore, 4 — 10 sensim paulo breviores, prioribus subcylindricis, sequentibus sensim paulo magis obconicis, ultimo apice oblique truncato et inferne acute acuminato, nigrae, apice ferrugineae. Palpi articulo ultimo piceo. Caput thoracis latitudine vel eodem paulo latius, orbiculatum, supra confertissime subtiliter punctatum, opacum, subtus parce punctatum, nitidum. Thorax coleopterorum prope latitudine, latitudine paulo brevior, lateribus leviter, basi fortius rotundatus, apice truncatus, angulis anterioribus subrectis, deflexis, posterioribus obtusis, leviter transversim convexus, confertissime subtiliter punctatus. Elytra thoracis longitudine vel eodem paulo longiora scutellumque confertissime subtiliter punctata. Abdomen supra subtilissime punctulatum, subtus parcius subtiliter punctatum, tenuiter pubescens. Alae flavescenti-hyalinae.

- \* 2. *O. cyaneus*: *Alatus, niger, subopacus, capite, thorace elytrisque supra obscure coeruleis, elytris thoracis longitudine.* — Long. 7—10 lin.

*Er. Col. March. I.* 440. 2.

*Staph. cyaneus* *Payk. Mon. Staph.* 13. 7. *Faun. Suec. III.* 370. 4. — *Fab. Ent. Syst. I. II.* 521. 11. *Syst. El. II.* 592. 13. — *Oliv. Ent. III.* 42. 14. 13. t. 1. f. 4. — *Walken. Faun. Paris. I.* 270. 5. — *Rossi Faun. Etr. Mant.* 96. 215. *Ed. Helw. I.* 442. 215. — *Panz. Faun. Germ.* 27. 3. — *Grav. Micr.* 6. 5. *Mon.* 110. 114. — *Latr. Hist. nat. Crust. et Ins. IX.* 304. 19. — *Marsh. Ent. Britt.* 501. 11. — *Gyll. Ins. Suec. II.* 286. 7. — *Sahlb. Ins. Fenn. I.* 309. 4. — *Kirby et Spence Introd. I. t. 1. f. 2.* — *Mannerh. Brachelytr.* 24. 16. — *Nordm. Symb.* 69. 60. — *Runde Brachelytr. Hal.* 3. 9.

*Emus cyaneus* *Boisd. et Lacord. Faun. Ent. Paris. I.* 368. 13.

*Staph. azurescens* *Mannerh. Brachelytr.* 23. 15. — *Nordm. Symb.* 69. 59.

*Staph. ophthalmicus* *Scop. Ent. Carn.* 99. 300. — *Goeze Ent. Beitr. I.* 725. 11.

*Staph. coerulescens* *Fourcr. Ent. Par. I.* 164. 2. — *De Vill. Ent. Par. I.* 421. 39.

*Staph. atro-coerulescens* Goetze Ent. Beitr. I. 729. 38.

*Le Staph. bleu* Geoffr. Ins. I. 361. 2.

Habitat in Europa.

O. simili plerumque paulo maior, niger, parum nitidus, capite, thorace elytrisque supra nigro-coeruleis, undique pube brevi subtilique nigro-cyaneo sericans. Antennae capite duplo prope longiores, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus, penultimis subobconicis, ultimo apice oblique truncato et inferne acuminato, nigrae, apice summo obscure ferrugineo. Palpi apice picei. Caput thoracis latitudine vel eodem paulo latius, orbiculato-subquadratum, creberrime subtilius punctatum, carinula frontali subtilissima, parum distincta, saepe omnino obsoleta, subtus parce punctatum, nigrum, nitidum. Thorax coleopteris paululum angustior, latitudine haud brevior, basi rotundatus, lateribus leviter rotundatis vel subrectis, apice truncatus, angulis posterioribus obtusis, subrotundatis, anterioribus deflexis, iam rectis vel subrectis, iam obtusiusculis, leviter transversim convexus, confertissime subtilius punctatus, carinula postica tenuissima, parum distincta, saepe omnino obsoleta. Scutellum nigro-tomentosum. Elytra thoracis longitudine, subtilius confertissime punctata. Abdomen subtiliter punctato-subrugulosum. Alae flavescenti-hyalinae.

Obs. 1. Variat magnitudine, exemplis maximis praecedenti duplo minoribus, minimis O. simili aequalibus; colore interdum subvirescente; capitis thoracisque punctura, iam fortiore, iam subtiliore et magis conferta, carinula media longitudinali magis minusve obsoleta vel omnino nulla; thoracis lateribus iam leviter rotundatis, iam fere rectis, angulis anterioribus iam obtusiusculis iam rectis; quae autem differentiae nullo modo stabiles videntur, et species ab auctoribus et mercatoribus nonnullis propositae (St. cyanescens Ziegl. subcyaneus et violaceus Dahl, azurescens Dej) rite distingui nequeunt. St. azurescens secundum Com. Mannerheim carinula frontis thoracisque omnino deficiente a genuino O. cyaneo differt, quae autem in indigenis saepius quam in speciminibus ex Europa meridionali obsolescit.

2. Pinguedine corruptus totus niger evadit. Huc referendus videtur *Staph. mordax* Block Ins. Vall. Plauens. 115. 2. f. 2. — *Staph. rugifrons* Runde Brachelytr. Hal. 4. 14. certe huc pertinet, quod ex specimine ab auctore descripto ipso a Dom. Zschorn mihi benevole communicato compertum habeo. (Conf. Col. March. p. 440.)

X \* 3. *O. Italicus: Apteris, niger, subtiliter ferrugineo-pubescent capite thoraceque confertissime subtiliter punctatis, hoc basi rotundato, elytris thorace brevioribus, brunneis. — Long. 9—11 lin.*

Habitat in montibus Appeninis, Dom. Schüppel.

Statura et summa affinitis O. macrocephali, niger, supra fere opacus, subtus subnitidus. Antennae capite paulo longiores, tenuiores, articulo tertio secundo sesqui longiore, sequentibus sensim brevioribus, ultimo apice oblique emarginato et inferne acuminato, nigro-piceae, apice ferrugineae. Palpi picei, apice rufescentes. Caput magnum, plerumque thorace latius, suborbiculatum, supra subdepressum, omnium confertissime subtiliter punctatum, linea tenuissima longitudinali laevi, anterie obsoleta, dense subtiliter ferrugineo-pubescent, subtus parce punctatus, glaber, nitidus. Thorax coleopterorum latitudine, latitudine vix longior, basin versus subangustatus, lateribus ante medium subangulatis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrectis, posterioribus obtusis, parum convexus,



confertissime subtiliter punctatus, linea tenuissima media longitudinali, anteri-  
us obsoleta, laevi, dense subtiliter ferrugineo-pubescent. Scutellum con-  
fertim subtiliter punctatum, subtiliter nigro-pubescent. Elytra thorace dimi-  
dio breviora, depressa, confertissime subtiliter punctata, rufo-brunnea, dense  
ferruginea pubescentia. Abdomen densius subtilissime punctatum, opacum,  
subtiliter ferrugineo-pubescent, subtus subnitidum, tenuiter nigro pubescent.  
Pedes nigri.

Obs. Simillimam *O. macrocephalo* hanc speciem, distinctam autem capitis tho-  
racisque punctura confertissima, hoc basin versus subangustato, basi fortius rotundato,  
Dom. Schüppel nomine Staph. Italici Gené signatam a Dom Frivaldsky accepit.

- \* 4. *O. macrocephalus*: *Apterus, niger, ferrugineo-pubescent, capite tho-  
raceque densius subtiliter punctatis, hoc basi subtruncato, elytris thorace  
brevioribus, rufo-brunneis.* — Long. 10—11 lin.

*Staph. macrocephalus* Grav. Micr. 160. 3. Mon. 111. 116. — Latr. Hist. nat.  
Crust. et Ins. IX. 305. 21. — Nordm. Synb. 64. 52.

Habitat in Hercyniae Austriaeque montibus, in Carpathibus et in Vol-  
hynia.

Statura fere *O. similis*, at multo maior, niger, capite thoraceque sub-  
aeneis, subnitidis, undique pube brevi fusco-ferruginea vestitus. Antennae  
capite plus sesqui longiores, articulo tertio secundo sesqui prope longiore,  
4—10 sensim paulo brevioribus, subcylindricis, penultimis leviter obconicis,  
ultimo apice oblique truncato et inferne acuminato, nigrae, apice rufo-testa-  
ceae. Palpi picei, apice rufescentes. Caput magnum, thorace latius, sub-  
orbiculatum, supra subdepressum, densius subtiliter punctatum, subtus laeve,  
glabrum, nitidum. Thorax coleopterorum latitudine, subquadratus, latitudine  
vix longior, basi subtruncatus, lateribus parallelis, ante medium subangulatis,  
apice truncatus, angulis posterioribus obtusis, subrotundatis, anterioribus le-  
viter deflexis, obtusiusculis, leviter convexus, densius subtiliter punctatus,  
basin versus obsolete canaliculatus. Scutellum piceum. Elytra thorace di-  
midio breviora, depressa, crebrius subtiliter obsoleteque punctulata, rufo-  
brunnea, opaca. Abdomen subtiliter punctulatum, piceum, segmentis singulis  
margine postico rufo-piceo. Pedes picei.

- \* 5. *O. megacephalus*: *Apterus, nigro-aeneus, capite thoraceque dense  
fortius punctatis, hoc basi fortiter rotundato, elytris thorace longe bre-  
vioribus, rufo-testaceis.*

*Staph. megacephalus* Nordm. Synb. 63. 51.

Habitat in Carinthiae montibus, Dahl.

Statura omnino praecedentis, at minor. Antennae capite sesqui lon-  
giores, articulo tertio secundo sesqui longiore, sequentibus sensim paulo  
brevioribus, subcylindricis, penultimis leviter transversis, ultimo apice obli-  
que truncato et inferne acuminato, nigrae, apice testaceae. Palpi rufo-picei.  
Caput thorace latius, subquadrato-orbiculatum, basi emarginatum, supra pa-  
rum convexum, obscure aeneum, nitidulum, dense fortius punctatum, linea  
media longitudinali subelevata laevi, subtiliter griseo-pubescent, lateribus  
parce pilosellum, subtus parce punctatum parceque pubescent, nigrum, niti-  
dum. Thorax coleopterorum latitudine, latitudine paulo longior, basin versus

subangustatus, basi cum angulis posterioribus fortiter rotundatus, lateribus subrectis, apice utrinque oblique truncatus, angulis anterioribus subrectis, leviter convexus, creberrime minus subtiliter punctatus, linea angusta media longitudinali subelevata laevi, obscure aeneus, nitidulus, tenuiter griseo-pubescent, lateribus parce pilosellus. Scutellum nigrum, nigro-pubescent. Elytra thorace dimidio breviora, depressa, subtiliter punctata, lateritia, testaceo-pubescentia. Abdomen piceo-nigrum, subtilissime punctulatum, supra tomento brevissimo fusco immixtis pilis fulvis vestitum, subtus tenuiter griseo-pubescent, segmentis singulis margine apicali dilutiore. Pedes nigro-picei, tarsi rufescentibus.

\* 6. *P. alpestris*: *Apterus, niger, opacus, elytris thorace dimidio brevioribus*. — Long. 7—9 lin.

Habitat in alpibus Carinthiacis, Dahl, Tyrolensibus, Stentz.

O similem colore et magnitudine aequans, ceterum autem abunde distinctus, O. macrocephalo proximus. Totus niger, opacus, subtiliter nigro-pubescent. Antennae capite sesqui longiores, articulo tertio secundo sesqui longiore, sequentibus sensim minoribus, praecipue brevioribus, cylindricis, penultimis leviter obconicis, ultimo apice oblique emarginato et inferne acuminato, apice testaceae. Palpi rufo-picei. Caput magnum, thorace multo latius, orbiculatum, supra perparum convexus, dense subtiliter punctatum, linea longitudinali tenuissima laevi antius obsoleta, subtiliter nigro-fuscescent subtus laevis, glaber, nitidus. Thorax coleopterorum latitudine, latitudine vix longior, basin versus subangustatus, lateribus subrectis, basi leviter rotundatus, apice truncatus, angulis posterioribus obtusis, subrotundatis, anterioribus rectis, leviter transversim convexus, subtiliter dense punctatus, basi obsolete canaliculatus, subtiliter nigro-fuscescent pubescent. Scutellum nigro-tomentosum. Elytra thorace dimidio breviora, depressa, confertissime punctata. Abdomen subtiliter punctulatum, segmentis singulis margine apicali piceo. Pedes apice picei.

Obs. 1. Ab O simili capite magno et elytris thorace multo brevioribus facile distinguendus, habitu omnino O. macrocephali eique nimis forte affinis, at statura minore et thorace basin versus subangustato, basi distincte rotundato, angulis anterioribus rectis constanter differre videtur. Elytra quoque plerumque concolora, rarius cum pedibus picea.

2. Specimina Tyrolensia Carinthiacis magnitudine paulo inferiora, capite minus lato, magis subquadrato quam suborbiculato, ceterum autem omnino congruentia, et cum non nisi gradu, non autem proportionem differant, haud rite distinguenda. Occurrunt etiam inter Carinthiacae specimina debiliora, his simillima, et inter Tyrolensia exempla validiora, ab illis nullo modo differentia.

3. Varietatem multo minorem, O. morioni aequalem, capite thorace haud latiore, subquadrato, nomine Staph. tenuipennis Dahl ex Austria missam dedit Dom. Schuppel, praeter staturam debilem nullo modo a O. alpestri discedentem, ab O. simili elytris brevioribus et punctura minus conferta distinctam.

\* 7. *O. similis*: *Apterus, niger, opacus, elytris thorace paulo brevioribus*. — Long. 8—9 lin.

Er. Col. March. I. 440. 3.

Staph. similis Fab. Ent. Syst. I. II. 521. 7. Syst. El. II. 591. 9. — Panz. Ent. Germ. 352. 6. — Grav. Micr. 5. 3. Mon. 111. 119. — Latr. Hist. nat. Crust. et

*Ins. IX. 302. 17. — Gyll. Ins. Suec. II. 287. 8. — Mannerh. Brachelytr. 24. 17.*

*— Nordm. Symb. 68. 56. — Runde Brachelytr. Hal. 3. 10.*

*Emus similis* Boisd. et Lacord. Faun. Ent. Paris. I. 368. 14.

*Staph. lugens* Nordm. Symb. 67. 54.

*Staph. nitens* Schrank Enum. Ins. Austr. 231. 436. Faun. Boica I. 640. 863.

*Staph. niger* Degeer Ins. IV. 22. 7.

*Staph. globulifer* Fourc. Ent. Par. I. 164. 3.

Habitat in Europa et in Sibiria.

Elongatus, niger, opacus, subtiliter breviterque nigro-pubescent. Antennae capite sesqui longiores, articulo tertio secundo parum longiore, sequentibus sensim minoribus, prioribus cylindricis, penultimis leviter obconicis, ultimo apice truncato et inferne acuminato, nigrae, apice ferrugineae. Caput thoracis latitudine, subquadratum, angulis omnibus rotundatis, supra subdepressum, confertissime subtiliter punctatum, linea tenui longitudinali laevi, anterieus obsoleta, supra parcius punctatum, nitidum, glabrum. Thorax coleopterorum latitudine, latitudine paulo brevior, lateribus rectis, basi rotundatus, apice utrinque oblique truncatus, angulis anterioribus subrectis, posterioribus obtusissimis, subrotundatis, leviter transversim convexus, confertissime subtiliter punctatus, linea tenui longitudinali laevi subelevata. Elytra thorace paulo breviora, depressa scutellumque confertissime subtiliter punctata. Abdomen subtilissime punctulatum. Pedes nigri.

Obs. 1. Specimina cum hac specie omnino convenientia nom. St. subterranei, brachypteri et hortensis a Dahlio ex Austria missa dedit Dom. Schüppel.

2. *Staph. lugens* Nordm. nullo modo ab O. simili distinctus.

\* 8. *O. masculus*: *Alatus, niger, opacus, palpis antennisque rufo-piceis, his basi nigris, elytris thoracis longitudine. — Long. 7 lin.*

*Staph. masculus* Nordm. Symb. 67. 55.

Habitat in Lusitania, Com. Hoffmannsegg.

Primo intuitu O. simili affinis videtur, at minor, et satis distinctus, niger, opacus, supra tomento tenui fusco-nigro vestitus. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, sequentibus sensim paulo minoribus, subobconicis, ultimo subovato, apice oblique truncato et inferne acuminato, basi nigrae, apice rufo-piceae. Palpi rufo-picei. Caput thoracis latitudine, subquadratum, supra subdepressum, subtiliter confertim punctatum. Thorax coleopterorum prope latitudine, latitudine vix longior, lateribus subrectis, basi fortiter cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, subdepressus, subtiliter confertissime punctatus. Scutellum atro-holosericum. Elytra thoracis prope latitudine, subtiliter punctato-subrugulosa. Abdomen parcius subtiliter punctulatum. Pedes nigri, tibiis piceis, tarsis rufescentibus.

\* 9. *O. picipes*: *Alatus, niger, opacus, antennis, palpis pedibusque piceis, thorace elytris subcrassiore, his thorace subbrevioribus. — Long. 7 lin.*

*Staph. picipes* Nordm. Symb. 68. 58.

Habitat in Austria et in Istria, Dahl, in ins. Jonicis, Parreyss.

Praecedenti affinis, niger, opacus, tomento subtili nigro-fusco obductus.



Antennae capite vix sesqui longiores, articulo tertio secundo sesqui longiore, sequentibus sensim paulo brevioribus, cylindricis, penultimis leviter obconicis, ultimo subovato, apice emarginato et inferne acuminato, rufo-piceae, apice ferrugineae. Palpi rufi. Caput thoracis prope latitudine, subquadratum, angulis omnibus fortiter rotundatis, supra parum convexum, confertim subtiliter punctatum. Thorax coleopteris prope latior, latitudine haud brevior, lateribus subrectis, parallelis, basi fortiter rotundatus, apice truncatus, angulis anterioribus subrectis, deflexis, posterioribus obtusissimis, leviter transversim convexus, confertissime subtiliter punctatus, postice linea longitudinali elevata, subtilissima, vix observanda. Scutellum nigro-pubescent. Elytra thoracis prope longitudine, confertissime subtiliter ruguloso-punctata. Abdomen apicem versus sensim attenuatum, subtilissime punctulatum, supra puncto fulvo-pubescente in medio baseos singuli segmenti. Pedes coxis posterioribus nigris, anticis femoribusque omnibus rufo-piceis, tibiis tarsisque rufis.

Obs. 1. Thorace crassiusculo et abdomine apicem versus sensim attenuato habitu quodammodo a praecedentibus discedit.

2. A Dahlio nom. Staph. picipedis et fuscipedis missus.

\* 10. *O. brunnipes*: *Apterus, niger, opacus, capite thoraceque nitidis, antennis basi apiceque pedibusque rufis, elytris thorace dimidio brevioribus.* — Long. 6 lin.

*Er. Col. March. I. 411. 4.*

*Staph. brunnipes* Fab. *Spec. Ins. I. 336. 8. Mant. Ins. I. 220. 12. Ent. Syst. I. II. 524. 23. Syst. El. II. 595. 26.* — *De Vill. Ent. I. 419. 27.* — *Oliv. Ent. III. 42. 15. 11. t. 1. f. 7.* — *Latr. Hist. nat. Crust. et Ins. IX. 326. 75.* — *Marsh. Ent. Britt. 502. 13.* — *Grav. Mon. 112. 124.* — *Gyll. Ins. Suec. II. 289. 10.* — *Sahlb. Ins. Fenn. I. 310. 7.* — *Mannerh. Brachelytr. 24. 23.* — *Nordm. Symb. 73. 72.* — *Runde Brachelytr. Hal. 4. 16.*

*Emus brunnipes* Boisd. et Lacord. *Faun. Ent. Paris. I. 371. 20.*

Habitat in Germania, Gallia, Anglia, Suecia.

Statura omnino *O. fuscata*, *niger*, *opacus*, subtiliter breviterque nigro-pubescent, capite thoraceque nitidis. Antennae capite duplo longiores, articulo tertio secundo paulo longiore, 4 — 8 subaequalibus, cylindricis, sequentibus sensim paulo minoribus, penultimis duobus leviter obconicis, ultimo apice truncato et inferne acuminato, basi rufae, medio nigrae, apice testaceae. Palpi rufi. Mandibulae rufo-piceae. Caput thoracis latitudine, suborbiculatum, supra leviter convexum, posterius creberrime, anterius parce punctatum, linea media longitudinali laevi, tenuiter subtiliterque nigro-pubescent, subtus parce punctatum, glabrum. Thorax coleopterorum latitudine, latitudine haud brevior, antrosum subangustatus, lateribus rectis, basi fortius rotundatus, apice truncatus, angulis anterioribus subrectis, posterioribus obtusis, subrotundatis, modice convexus, sat crebre minus subtiliter punctatus, linea tenui media longitudinali laevi, parce subtiliterque pubescent. Elytra thorace dimidio breviora scutellumque subtiliter confertissime punctata, subtiliter nigro-pubescentia, opaca. Abdomen dense subtiliter punctulatum. Pedes rufi, coxis piceis. Alae brevissimae.

- \* 11. *O. alpicola*: *Apterus, niger, opacus, capite thoraceque nitidis, sub-aeneis, antennis basi pedibusque rufis, illis apice flavis, elytris thorace paulo brevioribus.* — Long.  $6\frac{1}{2}$  lin.

Habitat in Carinthiae Carniolaeque alpibus.

*O. brunnipedi* proxime affinis, at paulo maior, niger, opacus, capite thoraceque nigro-subaeneis, nitidis. Antennae capite duplo longiores, articulo tertio secundo paulo longiore, 4 — 8 subaequalibus, cylindricis, sequentibus sensim paulo minoribus, penultimis duobus leviter obconicis, ultimo apice oblique truncato et inferne acuminato, nigrae, basi rufae, apice flavae. Palpi rufi. Mandibulae piceae. Caput thoracis latitudine, suborbiculatum, supra parum convexum, minus subtiliter sparsim punctatum, anteriùs sublaeve, interstitiis subtiliter punctatis, subtiliter tenuiterque nigro-pubescent, subtus parce punctatum, tenuiter pubescens. Thorax coleopteris prope latior, latitudine vix longior, lateribus subrectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus obtusiusculis, modice convexus, inaequaliter sat crebre subtilius punctatus, linea media longitudinali laevi obsoletiore, tenuiter pubescens. Elytra thorace parum breviora, depressa scutellumque subtiliter confertissime punctata, subtiliter nigro-pubescentia, opaca. Abdomen subtilissime minus confertim punctatum, supra opacum, subtus subnitidum. Pedes rufi, coxis rufo-piceis. Alae brevissimae.

Obs. A praecedente differt antennis apice dilutioribus, capite subtiliter inter-punctato, thorace minus crebre punctato, antrorsum nullo modo angustato, longius pubescente.

- \* 12. *O. fuscatus*: *Alatus, niger, opacus, capite thoraceque nigro-aeneis, nitidis, elytris thoracis longitudine.* — Long. 6 lin.

Er. Col. March I. 441. 5.

*Staph. fuscatus* Grav. Micr. 164. 10. Mon. 112. 123.

*Staph. crassicollis* Grav. Micr. 7. 7. — Latr. Hist. nat. Crust. et Ins. IX. 302. 15.

*Staph. subpunctatus* Gyll. Ins. Suec. IV. 474. 9. — Sahlb. Ins. Fenn. I. 310. 6.

— Mannerh. Brachelytr. 24. 19. — Nordm. Symb. 72. 67. — Runde Brachelytr. Hal. 3. 12.

*Staph. morio* Gyll. Ins. Suec. II. 288. 9. — Zetterst. Faun. Lappon. I. 69. 6. Ins. Lappon. 60. 7.

*Staph. punctulatus* Marsh. Ent. Britt. 501. 7.

*Staph. politus* Degeer Ins. IV. 22. 7.

Habitat in Europa.

Niger, opacus, subtiliter nigro-pubescent, capite thoraceque glabris, supra nigro-aeneis, pernitidis. Antennae capite sesqui longiores, articulo tertio secundo sesqui longiore, sequentibus sensim brevioribus, primis eorum subglobosis, penultimis transversim obconicis, ultimo apice truncato et inferne acuminato, nigrae, apice summo obscure ferrugineae. Palpi rufi. Caput thorace paulo angustius, suborbiculatum, supra parum convexum, undique creberrime, disco parce subtilius punctatum. Thorax coleopterorum latitudine, latitudine haud brevior, antrorsum subangustatus, lateribus rectis, basi rotundatus, apice truncatus, angulis anterioribus obtusiusculis, posterioribus obtusis, subrotundatis, transversim convexus, parcius subtiliusque punctatus. Scutellum creberrime punctulatum, densius pubescens. Elytra tho-

racis longitudine, omnium confertissime subtiliter punctata, dense subtiliterque nigro-pubescentia, opaca. Abdomen parcius subtiliter punctatum, subtilissime nigro-tomentosum. Pedes nigri, tarsi piceis. Alae fusco-hyalinae.

X

\* 13. *O. picipennis*: *Alatus, niger, capite thoraceque aeneis, nitidis, densius subtiliter punctatis, tenuiter pubescentibus, elytris tarsisque piceis, abdomine cinereo-lineato.* — Long. 7 — 8 lin.

*Er. Col. March. I. 442. 6.*

*Staph. picipennis* *Fab. Ent. Syst. I. II. 521. 8. Syst. El. II. 591. 10.* — *Panz. Ent. Germ. 353. 7.* — *Runde Brachelytr. Hal. 2. 4.*

*Staph. aeneocephalus* *Payk. Mon. Staph. 11. 6. Faun. Suec. III. 374. 10.* — *Grav. Micr. 8. 8. Mon. 113. 126.* — *Gyll. Ins. Suec. II. 291. 12.* — *Zetterst. Faun. Lappon. I. 68. 5. Ins. Lappon. 60. 6.* — *Sahlb. Ins. Fenn. 310. 9.* — *Mannerh. Brachelytr. 23. 11.* — *Nordm. Symb. 70. 61.*

*Emus aeneocephalus* *Boisd. et Lacord. Faun. Ent. Paris. I. 373. 22.*

*Staph. penetrans* *Müll. Prodr. 97. 1102.* — *Goeze Ent. Beitr. I. 727. 26.* — *Har-rer Besch. d. Ins. I. 251. 412.*

*Staph. quartus* *Schaeff. Icon. I. t. 35. f. 9.*

*Var. Elytris nigro-piceis.*

*Staph. tristis* *Fab. Ent. Syst. I. II. 524. 21. Syst. El. II. 594. 23.*

*Staph. sericeus* *Marsh. Ent. Britt. 508. 29.*

*Staph. leucophthalmus* *Marsh. Ent. Britt. 511. 39.*

*Staph. chalconcephalus* *Mannerh. Brachelytr. 23. 12.*

*Emus chalconcephalus* *Boisd. et Lacord. Faun. Ent. Paris. I. 374. 23.*

Habitat in Europa.

Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, sequentibus sensim brevioribus, prioribus subcylindricis, reliquis sub-obconicis, ultimo apice oblique truncato et inferne acuminato, rufo-piceae, basi nigricantes. Palpi rufo-picei. Caput thoracis fere latitudine, orbiculato-subovatum, supra parum convexum, dense subtiliterque punctatum, subtiliter griseo-pubescent, aeneum, nitidum, subtus nigrum, nitidum, parce nigro-pubescent. Thorax coleopterorum latitudine, latitudine vix longior, antrosum haud angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, transversim convexus, dense subtiliterque punctatus, linea media longitudinali tenui obsolete carinata, subtiliter griseo-pubescent, aeneus, nitidus. Scutellum nigrum, macula rotundata nigro-tomentosa. Elytra thoracis longitudine, subtiliter omnium confertissime punctata, densius plagiatis subtiliter cinereo-pubescentia, plerumque piceo-castanea, rarius fusco-nigricantia. Abdomen subtiliter punctulatum, punctis raris maioribus sparsis impressum, nigrum, opacum, supra pube densiore grisea 5-lineatum, linea intermedia distinctiore, angusta. Pectus nigrum. Pedes nigri, tibiae interdum piceis, tarsi rufo-piceis.

\* 14. *O. cupreus*: *Alatus, niger, capite thoraceque aeneis, nitidulis, confertissime subtiliter punctatis, hoc carinato, elytris fuscis, tarsis piceis, abdomine cinereo-lineato.* — Long. 5 — 7 lin.

*Er. Col. March. I. 443. 7.*



- Staph. cupreus* Rossi Faun. Etr. I. 248. 612. t. 7. f. 13. Ed. Hellr. I. 307. 612. t. 7. f. 13. — Fourc. Ent. Paris. I. 173. 33.
- Staph. aeneocephalus* Degeer Ins. IV. 22. 8. — De Vill. Ent. I. 423. 58; — Latr. Hist. nat. Crust. et Ins. IX. 300. 13.
- Staph. aeneicollis* Gyll. Ins. Suec. IV. 475. 12—13. — Sahlb. Ins. Fenn. I. 311. 10. — Mannerh. Brachelytr. 23. 13. — Nordm. Symb. 70. 63. — Runde Brachelytr. Hal. 3. 5.
- Emus aeneicollis* Boisd. et Lacord. Faun. Ent. Paris. I. 24. 74.
- Staph. sericeicollis* Menetr. Cat. 143. 570. — Nordm. Symb. 70. 64.
- Staph. strigatus* Nordm. Symb. 70. 65.
- Staph. puberulus* Runde Brachelytr. Hal. 3. 7.

Habitat in Europa.

Præcedenti similis, at vere distinctus, apud nos eodem dimidio vel duplo minor, specimina Italica nonnunquam illum aequantur. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus, prioribus subcylindricis, reliquis leviter obconicis, ultimo apice truncato et inferne acuminato, rufae, medio, interdum etiam basi nigropiceae. Palpi rufi. Caput thorace angustius, suborbiculatum, supra parum convexum, confertim subtilius punctatum, posterius linea tenui longitudinali subelevata, densius griseo-pubescent, aeneum, nitidulum, subtus nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine paulo longior, apicem versus sensim subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus obtusiusculis, transversim convexus, confertim subtilius punctatus, linea media longitudinali laevi elevata, densius griseo-pubescent, aeneus, nitidulus. Scutellum nigrum, griseo-pubescent, macula magna nigro-tomentosa. Elytra thoracis longitudine, omnium confertissime subtiliter punctata, aequaliter cinereo-pubescentia, fusco-picea. Abdomen dense subtiliter punctulatum, punctis maioribus raris sparsis impressum, opacum, supra pube densiore grisea 5-lineatum, lineis lateribus obsoletioribus, intermedia latiore distinctioreque. Pectus nigrum. Pedes nigri, tibiis tarsisque piceis.

- \* 15. *O. fulvipennis*: *Alatus, niger, capite thoraceque nigro-aeneis, nitidis, antennis, elytris pedibusque rufis, abdomine griseo-sublineato.* — Long. 5 — 6 lin.

- Staph. chaltocephalus* Nordm. Symb. 70. 62. — Runde Brachelytr. Hal. 3. 6.
- Staph. picipennis* Nordm. Symb. 71. 66.
- Emus picipennis* Boisd. et Lacord. Faun. Ent. Paris. I. 374. 25.

Habitat in Austria.

Statura omnino præcedentis. Antennae capite plus sesqui longiores, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus, subobconicis, ultimo apice truncato et inferne acuminato, rufo-testaceae, medio interdum piceae. Palpi rufi. Caput thorace paulo angustius, suborbiculatum, supra parum convexum, subtiliter densius punctatum, subtiliter griseo-pubescent, nigro-aeneum, nitidum, subtus laeve, glabrum, nigrum, nitidum. Thorax coleopterorum prope latitudine, latitudine paululum longior, antrorsum vix angustatus, lateribus levissime, basi fortiter rotundatus, apice truncatus, angulis anterioribus rectis, posterioribus obtusissimis, modice transversim convexus, dense subtilius punctatus, linea media longitu-

dinali tenui haud elevata, subtiliter tenuiterque griseo-pubescentia, obscure aeneus, nitidus. Scutellum griseo-pubescentia, apice nigro-tomentosum. Elytra thoracis longitudine, confertissime subtiliter punctata, rufo-testacea, aequaliter fulvo-pubescentia. Abdomen subtiliter punctulatum, punctis raris sparsis maioribus impressum, nigrum, opacum, supra griseo-pubescentia, sublineatum. Pectus nigrum. Pedes rufi, coxis nigris, anticis piceis.

Obs. Lapsu forte typographico a Nordm. l. c. apud Staph. picipennem Australasia pro huius insecti patria indicata est.

## Fam. II.

*Physetops Mannerh. Brachelytr. p. 32. — Nordm. Symb. p. 14. t. 1. f. 4.*

*Antennae articulo ultimo integro.*

*Palpi labiales articulo ultimo securiformi.*

*Mandibulae medio dentatae.*

Corpus elongatum, subcylindricum, alatum. Caput thorace paulo latius, oblongum, leviter convexum, basi minus profunde constrictum, oculis parvis leviter prominulis. Antennae basi distantes, articulo ultimo oblongo, acuminato. Labrum perbreve. Mandibulae prominentes, validae, leviter falcatae, acutae, medio fortiter dentatae. Palpi maxillares articulo quarto praecedente paulo minore, subacuminato, labiales articulis duobus primis breviusculis, ultimo leviter securiformi. Thorax oblongus, basin versus angustatus. Pedes tibiis intermediis crebre, posterioribus creberrime spinulosis, tarsis anticis leviter dilatatis.

Obs. Statura cylindrica praecipue a congeneribus discedit, vix autem rite genere distinguendus, labro paulo brevior, mandibulis prominentibus, antennarum articulo ultimo integro, acuminato et tarsis anticis leviter dilatatis diversus, palporum structura cum Fam. sequente conveniens.

\* 16. *O. Tataricus: Alatus, subcylindricus, niger, nitidus, capite thoraceque crebre punctatis, elytris thorace longioribus. — Long. 10 lin.*

*Staph. Tataricus Pall. It. App. n. 74. — Goeze Ent. Beitr. I. 723. 3. — Sturm Catal. 61. IX. t. 1. f. 9. — Guér. Iconogr. Regn. An. Ins. t. 9. f. 5.*

*Physetops Tartaricus Mannerh. Brachelytr. 33. 1. — Nordm. Symb. 14. 1.*

Habitat in desertis Russiae meridionalis.

Elongatus, subcylindricus, niger, nitidus. Antennae capite parum longiores, articulo tertio secundo sesqui longiore, quarto quinto paululum brevior, 5—10 sensim paulo brevioribus, penultimis duobus leviter transversis, ultimo praecedente duplo prope longiore, acuminato, nigrae, apice obscure ferrugineae. Caput oblongo-subquadratum, licet angulis rotundatis, crassum, fronte modice convexa, crebre punctata, lateribus subrugosa. Thorax capitis fere longitudine, latitudine antica paulo longior, basin versus angustatus, coleopteris angustior, lateribus pone medium sinuatis, basi rotundatus, apice truncatus, angulis omnibus obtusis rotundatisque, anterioribus fortiter deflexis, subcylindricus, supra subtilius crebriusque punctatus, lateribus punctato-rugulosus. Scutellum creberrime punctatum. Elytra thorace longiora, subtilissime parce griseo-pubescentia, subtilius minus crebre punctata, interstitiis subtiliter rugulosis. Abdomen subtiliter griseo-pubescentia,

apice nigro-pilosellum, subtiliter punctulatum punctisque maioribus sparsis impressum. Pedes nigri, tibiis tarsisque piceis.

Obs. „In deserto Tatarico circa lacum salsum Inderiensem primum observatus, at frequentissimus in montoso tractu orae meridionalis Chersonesi tauricae, ubi circa vias, praesertim fimo allectus recenti, perfrequens cursitat, sero praesertim autumnu. Cornicula odorifera ex caudae apice exserit alba, papillaria, fragrantissima, odore modo corniculorum larvae Machaonis, modo rosaceo, Cerambycis fere moschati.” Pall. mspt.

### Fam. III.

*Tasgius Steph. Illustr. Britt. Ent. V. p. 213. — Curtis Britt. Ent. X. t. 438.*

*Antennae articulo ultimo apice truncato.*

*Palpi labiales articulo tertio securiformi.*

*Mandibulae medio dentatae.*

Corpus elongatum, alatum. Caput basi profunde constrictum, collo quasi angusto thoraci insertum, suborbiculatum et subdepressum. Antennae apicem versus subattenuatae, articulo ultimo apice oblique truncato et inferne acuminato. Labrum latiusculum. Mandibulae longiores, validiores, falcatae, acutae, medio fortiter dentatae. Palpi breviores, maxillares articulo ultimo maris tertio fere aequali, apice truncato, subsecuriformi, feminae tertio minore, subacuminato, labiales in utroque sexu articulis duobus primis breviusculis, tertio leviter securiformi. Thorax basin versus subangustatus. Pedes tibiis intermediis crebre, posticis parce spinulosis, tarsis anticis fortiter dilatatis.

\* 17. *O. pedator*: *Alatus, niger, capite thoraceque nitidis, antennis pedibusque rufis, elytris thorace longioribus.* — Long. 9—10 lin.

*Staph. pedator Grav. Micr. 163. 9 Mon. 113. 125. — Nordm. Symb. 65. 53.*

*Astrapaenus rufipes Latr. Gen. Crust. et Ins. I. 285. 2.*

*Emus rufipes Boisd. et Lacord. Faun. Ent. Paris. I. 371. 19.*

*Tasgius rufipes Curt.-Britt. Ent. X. t. 438.*

Habitat in Gallia, Italia, Dalmatia.

Habitu omnino *O. atri*, et duplo maior at minus depressus, *O. cyaneo* aequalis, at angustior, niger, subnitidus, capite thoraceque nitidioribus. Antennae capitis thoracisque longitudine, articulo tertio secundo paulo longiore, sequentibus sensim minoribus, cylindricis, ultimo oblongo-ovato, apice truncato et inferne acuminato, obscure rufae, medio obscuriores. Palpi rufi. Mandibulae piceae. Caput thorace fere angustius, orbiculatum, supra parum convexum, minus subtiliter creberrime punctatum, parce pilosum. Thorax coleopterorum prope latitudine, latitudine paulo longior, basin versus subangustatus, lateribus rectis, basi rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, posterioribus obtusissimis, transversim convexus, crebre minus subtiliter punctatus, linea tenui longitudinali laevi, subelevata, pube subtili rara depressa vix conspicua vestitus, lateribus parce pilosus. Scutellum confertissime punctulatum, tenuiter pubescens. Elytra thorace paulo longiora, confertissime subtilius punctata, tenuiter nigro-pubescentia, cyanescenti-nigra, minus nitida. Abdomen subtiliter confertissime punctatum,



sparsim fortius punctatum, densius nigro-pubescens, parcius nigro-pilosellum. Pectus piceum. Pedes rufi, coxis omnibus concoloribus.

Obs. Palpi labiales articulo ultimo in utroque sexu subsecuriformi, maxillares articulo ultimo maris apice truncato, subsecuriformi, feminae subacuminato.

\* 18. *O. ater*: *Alatus, niger, nitidus, antennis apice ferrugineis, capite thoraceque parcius punctatis, elytris thorace sublongioribus.* — Long. 8 lin.

*Er. Col. March. I. 444. 8.*

*Staph. ater* Grav. *Micr. 161. 5. Mon. 112. 120.*

*Staph. fuscatus* Gyll. *Ins Suec. IV. 474. 9–10.* — Nordm. *Symb. 69. 58.*

*Staph. obscurus* Marsh. *Ent. Brit. 514. 50.*

*Emus nigripes* Boisd. et Lacord. *Faun. Ent. Paris. I. 370. 18.*

*Tasgius confinis* Curt. *Britt. Ent. X. fol. 438. n. 2.*

*Staph. Uralensis* Mannerh. *Brachelytr. 24. 20.* — Nordm. *Symb. 72. 68. (forte)*

Habitat in Europa infrequens; in montibus Uralensibus, Dom. Prof. Eversmann; in America septentrionali, Knoch.

Elongatus, subdepressus, niger, capite thoraceque nitidis. Antennae capite plus sesqui longiores, articulo tertio secundo sesqui longiore, sequentibus sensim brevioribus minoribusque, subcylindricis, penultimis leviter obconicis, ultimo apice oblique truncato et inferne acuminato, nigrae, apice obscure ferrugineae. Palpi picei. Caput thoracis fere latitudine, suborbiculatum, supra parum convexum, parcius fortiusque punctatum, linea media longitudinali laevi, parce subtiliterque nigro-pubescens. Thorax coleopteris paulo angustior, latitudine paululum longior, basin versus subangustatus, lateribus rectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, parum convexus, parcius punctatus, linea media longitudinali posterius subelevata laevi. Elytra thorace paulo longiora, depressa scutellumque confertim subtiliter punctata, minus nitida, densius subtiliter nigro-pubescens. Abdomen confertim punctulatum, minus nitidum. Pedes nigri, tarsis piceis.

Obs. 1. Palpi labiales sat fortiter securiformes in utroque sexu, maxillares maris subsecuriformes, feminae cylindricae, apice truncatae.

2. Synonymiam huius speciei confusam extricavi: specimina Europaea ab *O. atro* Americano, cuius exemplum a Gravenhorstio descriptum in Museo regio conservatur, nullo modo distincta, quare non potui quin *Staph. nigripedem* Dej. (*Staph. fuscatum* Gyll.) huic subiungerem. *Staph. Uralensem* Mannerh. non sine dubio allegavi, qui „convexus” et „thorace postice foveolis duabus longitudinalibus parum profundis punctatis impresso” describitur. Foveolas hasce autem thoracis linea longitudinali posterius elevata exortas esse suspicor: thorax etiam hoc loco crebrius punctatus est, et apud *O. morionem* quoque hoc modo nonnunquam thorax postice bi-impressus videtur.

#### Fam. IV.

*Ocypus* Steph. *Illustr. Britt. Ent. V. p. 211.*

*Anodus* Nordm. *Symb. p. 11. t. 1. f. 1.*

*Antennae articulo ultimo apice truncato.*

*Palpi labiales articulo tertio securiformi.*

*Mandibulae elongatae, falcatae, edentatae.*

Corpus subdepressum, omnibus alatum. Caput depressiusculum, thorace

saepe latius, oculis parvis, haud prominulis. Antennae basi magis distantes, graciles, apicem versus leviter attenuatae, articulis singulis crassitie tenuioribus, pilosellae, ab articulo quarto inde pube brevissima cinerea dense obducta. Labrum perbreve. Mandibulae elongatae, leviter falcatae, tenues, apice acuminatae, simplicissimae. Palpi breves, maxillares articulo quarto tertio subaequali, apice acuminato, labiales articulis duobus primis breviusculis. Thorax basin versus subangustatus. Abdomen parallelum, depressum. Pedes tibiis intermediis crebrius, posticis parcius spinulosi, tarsis anticis in utroque sexu fortiter dilatatis.

- \* 19. *O. morio*: *Alatus, niger, opacus, antennis apice ferrugineis, thorace angustiore, subtiliter carinato, elytris thoracis longitudine.* — Long. 5—7 lin.

*Er. Col. March* I. 444. 9

*Staph. morio* Grav. *Micr.* 6. 4. *Mon.* 112. 121. — *Latr. Hist. nat. Crust. et Ins.* IX. 303. 18. — *Gyll. Ins. Suec.* IV. 373. 8-9. — *Sahlb. Ins. Fenn.* I. 309. 5. — *Mannerh. Brachelytr.* 24. 18. — *Runde Brachelytr. Hal.* 3. 11.

*Emus morio* Boisd. et Lacord. *Faun. Ent. Paris.* I. 369. 15.

*Anodus morio* Nordm. *Symb.* 13. 3.

*Staph. edentulus* Block *Ins. Vall. Plauens.* 115. 1. f. 1.

*Staph. similis* Payk. *Mon. Staph.* 10. 5. *Faun. Suec.* III. 371. 6.

*Emus angustatus* Boisd. et Lacord. *Faun. Ent. Paris.* I. 369. 16.

*Emus fuscatus* Boisd. et Lacord. *Faun. Ent. Paris.* I. 372. 21.

*Staph. obscurus* Runde *Brachelytr. Hal.* 4. 13.

Habitat in Europa.

Elongatus, subdepressus, niger, opacus, capite thoraceque subnitidis, subtiliter nigro pubescens. Antennae capite duplo fere longiores, articulo tertio secundo sesqui longiore, sequentibus sensim paulo brevioribus, penultimis leviter transversis, ultimo praecedente paulo longiore, apice truncato et inferne acuminato, nigrae, apice ferrugineae. Palpi nigri, labiales articulo ultimo fortius securiformi. Mandibulae piceae. Caput thorace plerumque paulo latius, orbiculatum, supra subdepressum, minus subtiliter crebre punctatum, linea media longitudinali obsoleta, tenuiter subtiliterque pubescens. Thorax coleopteris angustior, latitudine vix longior, basin versus leviter angustatus, lateribus rectis, basi leviter rotundatus, apice truncatus, angulis anterioribus subrectis, posterioribus obtusis, subdepressus, confertim minus subtiliter punctatus, linea media longitudinali subelevata magis minusve distincta, tenuiter pubescens. Elytra thoracis longitudine vel eodem parum longiora, scutellumque confertissime subtiliter punctata, densius pubescentia. Abdomen dense subtiliter punctatum, densius pubescens. Pedes nigri, tarsis plerumque piceis, interdum picei, tarsis dilutioribus.

Obs. Variat statura maiore minoreve, capite thorace magis minusve latiore; feminae saepius capite thoracem latitudine haud superante instructae sunt.

- \* 20. *O. cerdo*: *Alatus, niger, opacus, confertissime punctatus, antennarum articulis tribus primis ultimoque, palpis tarsisque rufis, elytris thoracis longitudine.* — Long. 7—8 lin.

Habitat in Italia: in montibus Appeninis, Dom. Schüppel, in Sardinia, Dom. Gené.

Statura et summa affinitas *O. compressi*. Niger, vix nitidus, pube brevi subtili nigra dense vestitus. Antennae capite duplo fere longiores, articulo tertio secundo sesqui prope longiore, 4—7 subaequalibus, cylindricis, 8—10 sensim paulo minoribus, leviter obconicis, ultimo paulo crassiore, oblique truncato et apice inferne acuminato, nigrae, articulis tribus primis rufis, ultimo ferrugineo. Palpi rufi, labiales articulo tertio fortiter securiformi. Mandibulae nigro-piceae. Labrum nigrum. Caput subquadratum, thoracis prope latitudine, supra subdepressum, confertissime minus subtiliter punctatum. Thorax coleopteris vix angustior, latitudine sublongior, basin versus subangustatus, lateribus rectis, basi modice rotundatus, apice truncatus, angulis anterioribus subrectis, posterioribus obtusis, supra parum convexus, confertissime subtilius punctatus, pone medium linea longitudinali laevi subtilissima, vix observanda. Elytra thoracis longitudine et scutellum confertissime subtilius punctata. Abdomen densius subtiliter punctulatum. Pedes nigri, tibiis piceis, tarsis rufis.

Obs. Ab *O. morione* capite thoraceque confertissime punctatis, antennarum basi rufa, ab *O. compresso* capite thoraceque minus subtiliter punctatis, pedibus nigris, antennarum articulo ultimo solo testaceo facile distinguendus.

✕

\* 21. *O. compressus*: *Alatus, niger, opacus, pedibus, palpis antennisque rufis, his medio fuscis, elytris thoracis longitudine.* — Long. 7 lin.

*Staph. compressus* Marsh. Ent. Brit. 503. 15.

*Emus rufipalpis* Boisd. et Lacord. Faun. Ent. Paris. I. 370. 17.

Habitat in Anglia, Hollandia, Gallia, Hungaria.

*O. morioni* affinis et maioribus eius speciminibus aequalis, niger, opacus, subtiliter nigro-pubescent. Antennae capite plus sesqui longiores, articulo tertio secundo plus sesqui longiore, sequentibus sensim paulo minoribus, penultimis leviter transversis, ultimo apice oblique truncato et inferne acuminato, rufae, medio fuscae. Palpi rufi, labiales articulo ultimo fortiter securiformi. Mandibulae piceae. Labrum rufo-piceum. Caput thoracis latitudine, suborbiculatum, basi truncatum, supra subdepressum, confertissime subtiliter punctatum, subtilissime nigro-pubescent, subtus laeve, nitidum. Thorax coleopteris paululum angustior, latitudine vix longior, basin versus subangustatus, lateribus rectis, basi rotundatus, apice truncatus, angulis anterioribus rectis, posterioribus obtusis, parum convexus, confertissime subtiliter punctatus, dense subtiliter nigro-pubescent, opacus. Elytra thoracis longitudine scutellumque subtiliter confertissimeque subruguloso-punctata, dense subtiliter nigro-pubescentia. Abdomen subtiliter punctulatum, subtiliter nigro-tomentosum. Pedes laete rufi, coxis nigris, anticis intus rufescentibus.

\* 22. *O. falcifer*: *Alatus, niger, capite, thorace elytrisque supra coerules, antennis basi apiceque pedibusque rufis.* — Long.  $4\frac{1}{2}$ —7 lin.

*Anodus falcifer* Nordm. Symb. 12. 2.

*Anodus messor* Nordm. Symb. 11. 1.

Habitat in Italia, Dahl.

Statura omnino *O. morionis*, at magnitudine magis quam ille instabilis. Antennae capite plus sesqui longiores, articulo tertio secundo sesqui longiore, sequentibus sensim paulo brevioribus, penultimis leviter obconicis, ultimo ovato, apice inferne acuminato, fuscae, basi apiceque rufae. Palpi



rufi, labiales articulo ultimo leviter securiformi. Mandibulae nigrae. Caput thorace plerumque latius, subquadrato-orbiculatum, supra subdepressum, parcius punctatum, linea media longitudinali laevi, parce subtilissime nigro-pubescent, coeruleum, nitidum, subtus nigrum. Thorax coleopteris angustior, basin versus leviter angustatus, latitudine apicis parum longior, lateribus rectis, basi leviter rotundatus, apice utrinque oblique truncatus, angulis anterioribus rectis, posterioribus obtusis, subrotundatis, supra parum convexus, crebre punctatus, linea media longitudinali laevi subcarinata, subtiliter tenuiterque nigro-pubescent, coeruleus, nitidus. Scutellum confertim punctulatum, nigrum, subtiliter pubescens. Elytra thorace paulo longiora, depressa, confertim subtiliter punctata, densius subtiliter nigro-pubescentia, coerulea, minus nitida. Abdomen dense subtilius punctatum, densius subtiliter nigro-pubescent, nigrum, subnitidum. Pectus nigrum. Pedes laete rufi, coxis posterioribus nigro-piceis.

Obs. I. Individua maiora capite latiore et magis subquadrato, minora capite minore et magis orbiculato, minima capite thorace haud latiore instructa sunt.

2. Anodus falcifer et messor Nordm. nonnisi sexu differunt: ille mas hic femina.

3. Staph. morsitans Rossi Faun. Etr. I. 249. 614. Ed. Hellw. I. 308. 614, huic proximus videtur, at palpi nigri et caput, thorax et elytra nigro-aenea descripta: an autem revera distinctus?

### **Belonuchus Nordm.**

*Nordm. Symb. p. 129. t. 2. f. 2.*

*Antennae rectae.*

*Palpi filiformes.*

*Ligula rotundata, integra.*

*Femora antica et postica subtus biseriatim spinulosa.*

Corpus subdepressum, elongatum, saepe lineare, alatum. Caput exsertum, porrectum, basi profunde constrictum, oculis vix prominulis. Labrum transversum, medio incisum, limbo membranaceo, setoso. Mandibulae prominulae, tenues, decussatae, falcatae, apice valde acutae, muticae. Maxillae malis coriaceis, interiore intus ciliata, exteriori minuta, apice barbata. Palpi maxillares modice elongati, filiformes, articulo primo minuto, secundo ceteris paulo magis incrassato, quarto saepius tertio paulo longiore, apice leviter acuminato. Labium mento transversum, ligula minuta, rotundata, membranacea, paraglossis ligulam paulo superantibus, membranaceis. Palpi labiales filiformes, articulis omnibus subelongatis. Antennae filiformes, articulo primo modice elongato, ultimo apice oblique emarginato et inferne acuminato. Thorax coleopteris angustior, basin versus saepius angustatus, basi rotundatus, apice truncatus. Scutellum triangulare. Elytra truncata. Abdomen lineare. Pedes coxis intermediis distantibus, femoribus omnibus subtus leviter canaliculatis, anticis saepius leviter dilatatis posticisque subtus spinulis minutis biseriatis munitis, intermediis muticis, tibiis spinulosis, tarsis setulosis, posterioribus articulo primo leviter elongato, anticis in utroque sexu simplicibus.

### **Observationes.**

1. Genus *Philontho* proximum, non sine dubio mihi receptum et ab illo distinctum. Oris structura omnino eadem, et femora antica et postica subtus biseriatim spinulosa solam fere notam distinctivam praebent. Congruunt praeterea species in hoc genere enumeratae statura corporis magis minusve depressa, capite maiusculo, por-

recto, fronte antice sulco versus capitis medium evanescente impressa, mandibulis porrectis, muticis, apice falcatis, acutis, tibiis omnibus fortius spinulosis et tarsis anticis in utroque sexu simplicibus. Species quaedam thorace magis elongato (e. g. *B. haemorrhoidalis*) habitu *Xantholinum* simulant.

2. Corpus ut in *Philontho punctulatum*, pubescens, saepius parce pilosellum, capite thoraceque laevissimis, illo punctis 4 transversis inter oculos et pluribus supra oculos, hoc punctis praeter marginalia nonnullis dorsalibus et lateralibus impresso. Puncta dorsalia rarius singula, saepius utrinque 5 in series dorsales digesta sunt. Series dorsales infra et ultra hunc punctorum numerum nondum in hoc genere observatae.

3. Mas abdominis segmento inferiore sexto apice emarginato vel inciso distinctus, saepius etiam capite et mandibulis insignis.

4. Species omnes hucusque cognitae Americam meridiionalem habitant.

### 1) Thorax punctorum seriebus dorsalibus nullis.

\* 1. *B. haemorrhoidalis*: *Elongatus, depressus, niger, nitidulus, abdomine apice rufo*. — *Long. 4 — 5 lin.*

*Nordm. Symb. 130. 1.*

*Staph. haemorrhoidalis* *Fab. Syst. El. II. 596. 33.*

Habitat in America meridionali, in Brasilia, Dom. Beske, Sieber, in Columbia, Dom. Moritz.

Statura elongata fere *Xantholini*, depressus, niger, nitidus. Antennae capite plus sesqui longiores, apicem versus vix incrassatae, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo crassioribus, penultimis leviter transversis, nigrae. Palpi picei. Mandibulae modice elongati. Caput thorace latius, depressum, subquadratum, angulis posterioribus subdentatis, utrinque supra oculos punctis paucis profundis, inter oculos anterieus utrinque punctis duobus transversim positis impressum, fronte antice leviter canaliculata, canalicula ante apicem dilatata. Thorax coleopteris angustior, latitudine antica paulo longior, basin versus angustatus, lateribus pone medium sinuatis, angulis posterioribus rotundatis, apice utrinque oblique truncatus, angulis anterioribus subrectis, depressus, praeter puncta marginalia antice utrinque puncto unico dorsali, alteroque laterali impressus. Scutellum punctulatum. Elytra thorace paulo longiora, subtilius punctata, parce griseo-pubescens, pilis raris longioribus erectis, e punctis impressis maioribus nascentibus. Abdomen subtilius punctatum, nigro-pubescens, segmento quinto apice sextoque toto rufis. Pedes breviusculi, nigri. Alae fuscae.

2. *B. dichrous*: *Elongatus, depressus, nitidus, niger, thorace, pectore, elytris, ano pedibusque rufo-testaceis*. — *Long. 5 lin.*

Habitat in monte Orizaba in Mexico, collect. Dom. Chevrolat.

Statura et summa affinitas praecedentis. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, sequentibus duobus subcylindricis, crassitie paulo longioribus, 6 — 10 leviter obconicis, subaequalibus, nigrae, articulo ultimo ferrugineo. Palpi rufo-picei. Mandibulae modice elongatae, nigrae. Caput thorace latius, depressum, subquadratum, angulis posterioribus subdentatis, posterius utrinque punctis paucis profundis et inter oculos binis transversim positis impressum, fronte antice fortius ca-

naliculata, nigrum, nitidum. Thorax coleopteris angustior, latitudine antica vix longior, basin versus angustatus, lateribus pone medium subsinuatis, angulis posterioribus rotundatis, apice utrinque oblique truncatus, angulis anterioribus rectis, depressus, punctis nonnullis marginalibus ternisque utrinque discoidalibus, binis dorsalibus altero medio, altero ante apicem, tertioque lateri impressus, testaceo-rufus, nitidus. Scutellum subtiliter confertissime punctatum, rufum. Elytra dense subtiliter punctata punctisque paucis maioribus impressa, rufo-testacea, nitida, fulvo-pubescentia. Abdomen crebre subtiliter punctulatum, parcius nigro-pubescent, segmenti quinti margine apicali sextoque toto rufo-testaceis. Pectus totum rufo-testaceum. Pedes cum coxis rufo-testacei.

Obs. Differt a praecedente praeter colorem elytris subtilius, abdomine crebrius punctatis.

## 2) Thorax seriebus dorsalibus 5-punctatis.

\* 3. *B. Satyrus: Niger, nitidus, elytris rufis, ano late rufo-testaceo.* — Long. 4 — 5 lin.

Habitat in Columbia, Dom. Moritz.

Statura omnino Philonthi aenei, niger, nitidus, capite thoraceque supra sericeo-versicoloribus, elytris lateritiis, abdominis segmento quinto rufo-testaceo, basi nigro, sexto testaceo. Antennae capite sesqui longiores, apicem versus leviter incrassatae, articulo tertio elongato, secundo sesqui longiore, 4 — 10 sensim paulo brevioribus et crassioribus, nigrae, articulo secundo basi rufo. Palpi picei. Mandibulae longae, falcatae, decussatae, acutae, piceae. Caput maris thorace latius, feminae eius latitudine, suborbiculatum, utrinque supra oculos punctis pluribus, inter oculos utrinque duobus transversim positis fortibus profundisque, antice sulco lato longitudinali in frontis medio evanescente impressum, in mare intra antennarum insertionem utrinque cornicula prominulo munitum. Thorax coleopteris angustior, latitudine haud longior, basin versus subangustatus, lateribus pone medium subsinuatis, angulis posterioribus cum basi rotundatis, apice truncatus, angulis anterioribus subrectis, deflexis, parum convexus, seriebus dorsalibus punctis 5, antico et postico remotioribus, compositis, lateribus punctis praeter marginalia utrinque 6 impressus, punctis omnibus profundioribus. Scutellum subtiliter punctulatum, fulvo-pubescent. Elytra thorace parum longiora, subtiliter punctata, pube depressa fulva et pilis raris erectis fuscis vestita. Abdomen parce subtilissime punctulatum, tenuiter nigro-pubescent. Pedes nigri, tarsi picei. Alae fusco-hyalinae.

\* 4. *B. xanthopterus: Niger, nitidus, antennarum basi, pedibus, elytris pectorisque lateribus rufis, ano testaceo.* — Long. 5 lin.

*Philonthus xanthopterus* Nordm. *Symb.* 99. 83.

Habitat in Parà Brasiliae interiore, Sieber.

Statura omnino praecedentis. Antennae capite sesqui longiores, apicem versus leviter incrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, leviter obconicis, penultimis 4 leviter transversis, nigrae, articulis tribus primis testaceo-rufis, ul-



timo ferrugineo. Palpi testaceo-rufi. Mandibulae longiusculae, acutae, piceae. Labrum rufum. Caput nigrum, nitidum, thorace latius, transversim subquadratum, pone oculos utrinque punctis nonnullis vagis, inter oculos utrinque duobus transversim positis profundis, apice sulco lato, ante medium obsolescente, impressum, intra antennarum insertionem utrinque subdentatum. Thorax coleopteris angustior, latitudine paulo brevior, basin versus vix angustatus, lateribus pone medium subsinuatis, basi cum angulis posterioribus rotundatus, apice utrinque oblique truncatus, angulis anterioribus subrectis, leviter convexus, seriebus dorsalibus punctis 5, antico et postico paulo remotioribus, compositis, lateribus punctis praeter marginalia utrinque 6 impressus, punctis omnibus profundioribus, niger, nitidus. Scutellum piceum, creberrime punctulatum, fulvo-pubescentia. Elytra thorace paulo longiora, crebre subtilius punctata, testaceo-rufa, fulvo-pubescentia. Abdomen punctulatum, longius nigro-pubescentia, nigrum, nitidum, segmento quinto apice, sextoque toto testaceis. Pectus rufum, medio piceum. Pedes rufi, coxis anterioribus nigris.

\* 5. *B. mordens*: *Niger, nitidus, ano late testaceo, thorace lateribus subrotundato, elytris subtiliter punctatis.* — Long. 5 lin.

Habitat in Brasilia, Dom. Prof. Germar.

Statura omnino praecedentium. Corpus nigrum, nitidulum, capite thoraceque nitidis. Antennae capite sesqui longiores, articulo tertio secundo sesqui longiore, 4—10 subobconicis, sensim paulo brevioribus, fere transversis, ultimo oblongo, apice emarginato et inferne acuminato, fuscae, basi nigro-piceae, articulo ultimo ferrugineo. Palpi picei. Mandibulae longiusculae, acutae, piceae. Labrum rufo-piceum. Caput thorace paulo latius, subquadratum, depressiusculum, supra oculos utrinque punctis parvis, inter oculos utrinque duobus oblique positis profunde impressum, fronte antice late canaliculata. Thorax coleopteris angustior, latitudine paulo brevior, basin versus vix angustatus, lateribus leviter rotundatis, pone medium vix sinuatis, basi cum angulis posterioribus rotundatus, apice utrinque oblique truncatus, angulis anterioribus obtusiusculis, leviter convexus, seriebus dorsalibus punctis 5, postico paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 6 impressus. Scutellum crebre punctatum. Elytra thorace longiora, confertim subtilius punctata, pube depressa grisea pilisque raris longioribus erectis, e punctis maioribus nascentibus, vestita. Abdomen parce punctulatum, nigro-pilosellum, segmentis ultimis duobus totis testaceis. Pedes nigro-picei, tarsis piceis.

\* 6. *B. latro*: *Niger, nitidus, ano late rufo-testaceo, thorace lateribus recto, elytris fortius punctatis.* — Long. 4 lin.

Habitat — — E collectione Herbstiana.

Statura et summa affinitas praecedentis, at paulo minor, niger, nitidus, abdominis segmentis duobus apicalibus totis rufo-testaceis. Antennae capite plus sesqui longiores, articulo tertio secundo paulo longiore, 4—7 sensim paulo brevioribus et crassioribus, 7—10 aequalibus, 5—10 transversis, ultimo ovato, apice subemarginato et inferne acuminato, fuscae, basi piceae, apice obscure ferrugineae. Palpi picei. Mandibulae longiusculae,

acutae, piceae. Labrum rufo-piceum. Caput thoracis latitudine, subquadratum, depressiusculum, utrinque supra oculos punctis pluribus, inter oculos utrinque duobus transversim positis sat magnis profundisque impressum, antice canaliculatum, sulco latiusculo, ultra frontis medium adscendente. Thorax coleopteris angustior, latitudine haud brevior, lateribus rectis, basi rotundatus, apice truncatus, angulis anterioribus subrectis, posterioribus obtusis, supra leviter convexus, seriebus dorsalibus punctis 5, apicali paulo minore et remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus fortibus et sat profundis. Scutellum crebre punctatum. Elytra thorace paulo longiora, minus dense profundeque punctata, pube depressa nigra pilisque raris erectis vestita. Abdomen punctulatum, nigro-, segmentis ultimis duobus fulvo-pilosellum. Pedes picei. Alae fusco-hyalinae.

Obs. A praecedente differt statura paulo minore, capite utrinque fortius punctato, thorace longiore, lateribus recto, elytris brevioribus, minus subtiliter punctatis.

7. *B. avidus*: *Niger, nitidus, ano rufo-testaceo, antennis articulo ultimo ferrugineo, thorace lateribus recto, elytris fortius punctatis.* — Long.  $3\frac{1}{2}$  lin.

Habitat Novae Valentiae in Columbia, Dom. Moritz.

Praecedenti proximus, at dimidio minor, niger, nitidus, ano testaceo. Antennae capite thoraceque paulo breviores, apicem versus parum incrassatae, articulo secundo tertioque aequalibus, 4 — 10 crassiusculis, subcylindricis, crassitie vix longioribus, inter se aequalibus, ultimo ovato, apice truncato et inferne acuminato, nigrae, articulo ultimo obscure ferrugineo. Palpi elongati, nigro-picei. Caput thoracis latitudine, subquadratum, inter oculos punctis quatuor transversim positis, supra oculos utrinque punctis nonnullis dispersis impressum, fronte antice profunde sulcata. Thorax coleopteris dimidio angustior, latitudine paululum longior, basin versus subangustus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrectis, leviter convexus, seriebus dorsalibus punctis 5, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 6 impressus, punctis omnibus sat profundis. Scutellum crebre punctulatum. Elytra thorace paulo longiora, densius sat profunde punctata, longius griseo-pubescentia. Abdomen minus confertim punctatum, longius nigro-pubescent, segmento penultimo margine lato apicali ultimoque toto laete rufo-testaceis. Pedes nigri, tarsi apice piceis.

8. *B. agilis*: *Depressus, niger, nitidus, ano late rufo, thorace basin versus angustato, elytris subtilius punctatis.* — Long. 5 lin.

Habitat in Cuba, collect. Dom. Chevrolat.

Statura depressa et summa similitudo *B. haemorrhoidalis*, niger, nitidus, parce nigro-pilosellus, capite thoraceque sericeo-micantibus, abdomine segmentis ultimis duobus totis rufis. Antennae capite sesqui longiores, articulo secundo tertioque elongatis, aequalibus, quarto subgloboso, 5 — 10 transversim subobconicis, aequalibus, ultimo subovato, apice emarginato et inferne acuminato, totae nigrae, articulo ultimo concolore. Palpi picei. Caput thorace paulo latius, depressum, quadratum, punctis 4 transversim positis

inter oculos et nonnullis in fronte superiore sparsis, sat profundis, impressum, fronte antice subtiliter canaliculata et inter antennis obsolete impressa. Thorax coleopteris dimidio angustior, latitudine paulo longior, basin versus angustatus, lateribus anterie leviter rotundatus, basi subtruncatus, apice utrinque oblique truncatus, angulis anterioribus obtusiusculis, posterioribus obtusis, depressus, seriebus dorsalibus punctis 5, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum crebre punctatum. Elytra thorace paulo longiora, minus dense subtilius punctata, subtiliter nigro-pubescentia. Abdomen sublaeve, parce obsoleteque punctatum. Pedes toti nigri.

- \* 9. *B. gagates*: *Subdepressus, niger, nitidus, antennis palpis pedibusque concoloribus, capite suborbiculato.* — Long. 3 — 4 lin.

Habitat in Puerto Rico, Dom. Moritz, in Cuba, Mus. Dom. Chevrolat.

Statura omnino *B. formosi*, totus aterrimus, pernitidus. Antennae capite thoraceque paulo breviores, apicem versus leviter incrassatae, articulo tertio secundo sesqui prope longiore, 4 — 10 crassiusculis, sensim brevioribus et crassioribus, penultimis transversis, ultimo subovato, apice oblique emarginato et inferne acuminato, totae nigrae. Mandibulae palpique elongati nigri. Caput thorace paulo latius, suborbiculatum, supra oculos utrinque punctis nonnullis, inter oculos 4 transversim positis sat profundis impressum, fronte antice sulcata. Thorax coleopteris angustior, latitudine paulo longior, basin versus subangustatus, lateribus anterie subrotundatis, pone medium subsinuatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus obtuse rotundatis, parum convexus, seriebus dorsalibus punctis 5, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum crebre punctatum. Elytra thorace paulo longiora, subtilius minus confertim punctata, tenuiter nigro-pubescentia. Abdomen parce punctatum parceque nigro-pubescentia et pilosellum. Pedes nigri. Alae fusco-hyalinae.

- \* 10. *B. minax*: *Subdepressus, niger, nitidus, antennarum basi apiceque pedibusque testaceis, capite suborbiculato.* — Long. 3 — 4 lin.

Habitat in Americae insulis St. Thomae et St. Johannis, Dom. Moritz.

Elongatus, subdepressus, niger, nitidus, capite thoraceque nitidissimis. Antennae capite thoraceque breviores, articulo secundo tertioque aequalibus, articulis 4 — 10 sensim paulo brevioribus at vix crassioribus, penultimis crassitie paulo brevioribus, ultimo ovato, apice oblique truncato et inferne acuminato, nigrae, articulis basalibus tribus rufo-testaceis, apicalibus duobus ferrugineis. Palpi rufo-testacei. Mandibulae elongatae, falcatae, acutae, rufo-piceae. Caput thoracis latitudine, vel maris saepe thorace latius, suborbiculatum, supra oculos utrinque punctis sat profundis pluribus vagis, inter oculos 4 transversim positis impressum, fronte antice longitudinaliter sulcata. Thorax coleopteris angustior, latitudine haud longior, basin versus subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrectis, parum convexus, seriebus dorsalibus punctis 5 compositis, lateribus punctis praeter marginalia utrinque 6 impressus, punctis omnibus profundioribus. Scutellum crebre



punctatum. Elytra thorace paulo longiora, subtilius minus confertim punctata, tenuiter nigro-pubescentia. Abdomen nigro-piceum, segmentis omnibus apice piceo-marginatis, subtiliter punctulatum, parcius nigro-pubescentia. Pedes rufo-testacei. Alae fusciscenti-hyalinae.

\* 11. *B. planiceps*: *Elongatus, depressus, niger, antennarum basi pedibusque testaceis, capite subquadrato.* — Long.  $2\frac{2}{3}$  lin.

Habitat in Columbia, Dom. Moritz.

Praecedenti affinis, at minor et magis depressus, niger, capite thoraceque nitidis. Antennae capite sesqui longiores, crassiusculae, articulo secundo tertioque aequalibus, 4—10 crassiusculis, 5—10 crassitie aequalibus, sensim paulo brevioribus et magis transversis, ultimo ovato, apice subemarginato et inferne acuminato, nigrae, apice concolores, articulis tribus primis rufo-testaceis. Palpi rufo-testacei. Mandibulae arcuatae, acutae, rufo-piceae. Caput thorace latius, subquadratum, depressum, utrinque inter oculos punctis duobus oblique positis, supra oculos punctis pluribus grossiusculis, minus profundis, vage impressum, fronte apice longitudinaliter sulcata. Thorax coleopteris dimidio fere angustior, latitudine paulo longior, basin versus leviter angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice utrinque oblique truncatus, angulis anterioribus subrectis, depressus, seriebus dorsalibus punctis 5, apicali vix remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis minus subtilibus. Elytra thorace paulo longiora et scutellum dense subtilius punctata, tenuiter griseo-pubescentia. Abdomen parcius subtiliter punctulatum, parcius griseo-pubescentia, segmento quarto quintoque pube rigidiora depressa obsitis, subtus segmentorum marginibus piceis. Pedes rufo-testacei, coxis anterioribus tibiisque omnibus piceis. Alae hyalinae.

12. *B. mundus*: *Elongatus, subdepressus, niger, elytris rufo-testaceis, pectore rufo, mandibulis palpis pedibusque rufo-piceis.* — Long. 3 lin.

Habitat in Americae meridionalis ins. St. Vincentii, Mus. Dom. Hope.

Statura omnino praecedentium. Antennae capite sesqui longiores, apicem versus modice incrassatae, articulo tertio secundo sesqui longiore, 4—10 sensim crassioribus at vix brevioribus, ultimo subovato, apice oblique truncato et inferne acuminato, nigro-piceae, articulo ultimo obscure ferrugineo. Mandibulae palpique elongati, rufo-picei. Caput thorace latius, subquadratum, punctis 4 transversim positis inter oculos et utrinque supra oculos pluribus vagis sat fortibus impressum, fronte anterieus fortius sulcata, nigrum, pernitidum. Thorax coleopteris dimidio angustior, latitudine paulo longior, basin versus subangustatus, lateribus subrectis, basi cum angulis posterioribus leviter rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, parum convexus, seriebus dorsalibus punctis 5 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 6 impressus, punctis omnibus sat fortibus profundisque, niger, pernitidus. Scutellum punctatissimum, nigrum. Elytra thorace vix longiora, depressa, minus dense subtilius punctata, subtiliter testaceo-pubescentia, rufo-testacea, nitida. Abdomen parcius punctatum, tenuiter griseo-pubescentia, nigrum, nitidum. Pectus rufum. Pedes dilute picei, coxis concoloribus.

\* 13. *B. formosus*: *Rufus, capite, thorace, abdominis apice antennisque nigris, his basi apiceque pedibusque testaceis. — Long. 3 lin.*

*Staph. formosus* Grav. Mon. 72. 61.

*Staph. rufipennis* Fab. Syst. El. II. 597. 39.

Habitat in Columbia, Dom. Moritz, Cayennae, Dom. Banon, in Carolina sec. Gravenhorst.

Antennae capite duplo fere longiores, articulo secundo tertioque aequalibus, 4—10 sensim paulo brevioribus, 5—10 crassitie aequalibus, penultimo leviter transverso, ultimo subovato, apice oblique truncato et inferne acuminato, nigrae, articulo primo laete, secundo tertioque obscure testaceis, ultimo ferrugineo. Palpi testacei. Mandibulae longiusculae, falcatae, acutae, rufo-piceae. Caput thoracis latitudine, vel eodem paulo latius, rotundato-subquadratum, punctis 4 transversim positis inter oculos et pluribus vagis utrinque supra oculos, omnibus sat fortibus, impressum, fronte antice late sulcata, nigrum, nitidum. Thorax coleopteris angustior, latitudine parum longior, basin versus vix angustatus, lateribus rectis, basi cum angulis posterioribus fortius rotundatus, apice truncatus, angulis anterioribus subrectis, leviter convexus, seriebus dorsalibus punctis 5 compositis, lateribus punctis praeter marginalia utrinque 6 impressus, punctis omnibus sat fortibus profundisque, niger, nitidus. Scutellum creberrime punctatum, piceum. Elytris thoracis longitudine, subtilius punctata, rufa, nitida, pube depressa flavescente pilisque raris erectis fuscis vestita. Abdomen parcius punctulatum, longius tenuiter griseo-pubescent, rufum, nitidum, segmentis ultimis duobus nigris, nigro-pubescentibus. Pectus rufum. Pedes rufo-testacei. Alae fuscae.

Obs. In describendo hoc insecto Fabricius abdominis segmenta 4 anteriora rufa neglexit, quare nonnisi specimine, quod auctor ipse descripsit, inspecto et examinato, hoc synonymum cognovi. Gravenhorstii descriptio exacte convenit, nisi quod antennarum basin testaceam haud memorat, nihilominus eius citatum pro certo habeo, et nomen eius praetulit, quod ille prius quam Fabricius alii speciei (Philonth. rufipenni) tribuerit. In Catalogo Dejeaniano recentiore haec species nomine Cafii apicalis signata est.

### **Philonthus Leach.**

*Curt. Britt. Ent. XIII. t. 610. — Nordm. Symb. p. 73. — Er. Col. March. 1. p. 445. Philonthus Steph. Illustr. V. p. 226. — Cafius ibid. p. 245. — Curt. Britt. Ent. VII. p. 322. — Mannerh. Brachelytr. p. 31. — Bisnius Steph. l. c. p. 247. — Gabrilius Steph. ibid. p. 249. — Remus Holme Transact. Ent. Soc. Lond. II. I. p. 64.*

*Antennae rectae.*

*Palpi filiformes.*

*Ligula rotundata, integra.*

*Pedes femoribus omnibus muticis.*

Corpus plerisque mediae magnitudinis, elongatum, lineare, plerisque alatum. Caput aliis suborbiculatum, aliis ovatum, exsertum, collo quasi crassiusculo thoraci immissum. Labrum transversum, medio incisum, margine membranaceo setosoque. Mandibulae falcatae, medio leviter dentatae. Maxillae (t. IV. f. 12. b.) malis parvulis, coriaceis, interiore intus, exteriori

apice barbata. Palpi maxillares modice elongati, articulo primo minuto, secundo reliquis paulo magis elongato, ultimo apice magis minusve acuminato. Labium (t. IV. 12. a.) mento brevi, transverso, ligula semiorbiculari, rotundata, membranacea, paraglossis coriaceis, interne ciliatis, ligulam superantibus. Palpi labiales filiformes, articulis tribus sensim paulo longioribus. Antennae filiformes, articulo primo leviter elongato, ultimo apice truncatulo vel magis minusve emarginato et inferne acuminato. Thorax coleopteris plerumque paulo angustior, basi rotundatus, apice truncatus, angulis posterioribus obtusis vel rotundatis, anticis deflexis, magis minusve rotundatis. Scutellum triangulare. Elytra truncata. Abdomen lineare. Pedes breviusculi, coxis intermediis aliis approximatis, aliis magis minusve distantibus, tibiis aut omnibus aut posterioribus spinulosis, tarsis setulosis, posterioribus articulo primo ultimoque leviter elongatis, anticis aut in utroque sexu aut in masculis solis dilatatis, aut in utroque sexu simplicibus.

#### Observationes.

1. Corpus punctulatum, pubescens, capite thoraceque politis, glaberrimis, plerumque punctis parvis notatis, illo punctis 2 vel plerumque 4 inter oculos et pluribus supra utrumque oculum, hoc seriebus dorsalibus e punctis plerumque 3, 4, 5, 6 aut pluribus et nonnullis in utroque latere inter seriem dorsalem et marginem, plerisque in anteriore parte sitis, notato. Nonnullis (divis. 8.) caput et thorax crebrius punctati et puncto quoque pilifero subtiliter pubescentia, spatio medio longitudinali laevissimo. A Staphylino genuino et ab Ocypode distinguitur hoc genus imprimis ligula integra, reliqui enim characteres non satis stabiles.

2. Characteribus mancis plura ab auctoribus condita sunt genera, quae huic subiunxi:

**Cafius**, species quasdam marium littoribus proprias continens, corpore elongato, depressiusculo, ab auctoribus recentioribus plerisque receptum et antennarum articulo ultimo ovato, apice haud emarginato a reliquis distinctum. In Ph. xantholomate, huius generis typo, antennarum articulus terminalis revera vix distincte apice emarginatus, attamen non omnino ovatus sed apice abrupte subacuminatus; in Ph. fucicola apice truncatulus et inferne subacuminatus, ut in multis aliis; in Ph. fonticola, qui Ph. xantholomatis habitum insigniter prae se fert et corporis structura et pubescentia cum eo congruit, apice profundius emarginatus. — Ph. xantholomate pro Cafii typo prosito, huic adiungendi essent Ph. variegatus; fucicola, cicatricosus, australis et fonticola, auctores autem parum congruunt in Cafii generis limitibus determinandis. Com. Mannerheim. Ph. nanum et aterrimum, Com. Dejean in Catalogo collectionis suae recentiore species plerasque, quibus genus Belonuchus Nordm. componitur et nonnullos alios Philonthos (Ph. cribricollem) cum Ph. xantholomate coniungunt. Dom. Lacordaire denique, potius characterem quam typum propositum respiciens, tantum aberravit, ut Othium fulvipennem pro mero Cafio cognosceret (V. s. p. 295.).

**Bisnius**, pro cuius typo Ph. cephalotes proponitur, et cui Ph. sordidus, fimetarius, fuscus, placidus et blandus summa affinitate adiunguntur, a Dom. Stephens palporum articulo ultimo acuminato a Philontho genuino distinguitur. Potius praeter corporis staturam elongatam species laudatae tarsis anticis in utroque sexu simplicibus, quamvis hoc characterem Dom. Stephens cum Philontho Bisnium componit, inter se conveniunt, differentia autem in palporum acumine adeo exigua et nulla, ut qui eam observare possit nesciam.

**Gabrius** contra tarsis anticis simplicibus a Philontho distinguitur ab auctore modo laudato: licet species sub hoc genere enumeratae pleraeque (Ph. nanus, aterrimus, quibus forte Ph. nitidulus et vernalis adnumerandi sunt) hac nota inter se conveniunt, tamen nullo modo ab aliis permixtis huius generis speciebus eadem nota distinguuntur. Ph. cinerascens etiam Gabrio adscriptus, tarsis anticis in utroque sexu dilatatis gaudet.

**Remus**, genus a Dom. Holme propositum thoracis crebre punctati, quo cum Lathrobiis conveniret, et palporum filiformium causa, quibus ab iis distin-



guatur. Palpi omnino huius generis, sculptura e characteribus genericis reicienda Species ab auctore descripta Ph. cinerascenti simillima et summa affinitate cum eodem iuncta.

**Philonthus** denique ipse a Dom. Stephens antennarum apice abrupte acuminato, tarsis anticis mediocriter dilatatis, articulo ultimo eorum elongato, oculis mediocribus, palporum articulo ultimo truncato distinguitur.

3. Philonthi ab auctoribus plerisque Staphylinis genuinis adscripti sunt. Com. Dejean hos sub Emi nomine cum Ocypode coniunxit, illis nomen Linneanum retinens. Prof Nordmann Philonthi genus recepit, Quedios vero nonnumquam distinxit, nonnullos autem Philonthos minores corpore elongato, quippe quos Com. Mannerheim Cafio substituit, Gyrohypno (Xantholino) adiecit.

4. Species nonnullae in huius generis calce descriptae, (Ph. proluxus, procerulus, elongatulus, sobrinus, humilis, cognatus, ludicrus et palmula), palporum maxillarum articulo ultimo fortius acuminato, acumine elongato, acutissimo, a reliquis magis recedere videntur, ceterum capite thoraceque crebre punctatis, hoc seriebus dorsalibus nullis et tarsis anticis leviter dilatatis ad Ph. cinerascensem propius accedunt, nec genus peculiare iis condendum putavi, quod character eam laudato, solo, quo a Philonthis distinguuntur, species enumeratae nonnullae aliquantulum inter se discedunt.

5. Mas abdominis segmento inferiore sexto apice emarginato vel plerumque inciso, saepius quoque tarsis anticis fortius dilatatis distinguitur. Nonnullae etiam species capite in masculis validiore insignes, mares earum autem capitis forma quodammodo variant secundum individui staturam, maioribus caput maius, latius, et antrosum magis dilatatum, minoribus caput minus et magis orbiculare vel ovatum esse solet, ita ut minima specimina eius proportionem a femineis vix distinguantur. Caveas igitur, ne speciei eiusdem individua ob capitis disparem figuram pro speciebus distinctis habeas.

6. Victus in cadaveribus, in stercore, sub foliis deciduis, musco et lapidibus. Per omnem terrarum orbem occurrit hoc genus.

7. Philonthi aenei larvam observavit Dom. Bouché et hoc modo descripsit: „Linearis, apicem versus angustata, depressa, grisea, parce ferrugineo-setulosa. Caput lenticulare, piceum, subtilissime rugulosum, nitidum, glabrum. Antennae subuliformes, 5-articulatae, articulo primo et secundo brevibus, cylindricis, tertio secundo triplo longiore, obconico, tenui, quarto eadem figura, at dimidio brevior, quinto minuto, conico, quarto interne ad apicem processu parvo elliptico aucto. Clypeus brevis, rotundatus, apice 6-denticulatus. Labrum breve, membranaceum, inflexum. Mandibulae longae, tenues, falcatae, muticae, parce setosae. Maxillae oblongo-lineares, processu brevi elliptico, palpo lato, 5-articulato, articulo basali brevi. Labium parvum, rotundatum, palpis subulatis, 2-articulatis. Prothorax quadrangulus, latitudine paulo longior, lateribus leviter marginatus, capite paulo angustior. Meso- et metathorax transversus, prothorace latiora, quadrangula. Pedes longi, tenues, femoribus basi seta longa, tenui, subtus spinulis ferrugineis biseriatis instructis, tibiis brevibus, spinulosis, spinulis ferrugineis, tarsis minutis, ferrugineis, unguiculis simplicibus et spinis binis terminatis. Abdomen segmentis aequalibus, supra et infra maculis binis quadratis obscuris, segmento anali angustato, caudis duabus elongatis subulatis, triarticulatis, ferrugineo-setulosis, segmento anali sextuplo longioribus. Anus tubulosus, verticaliter descendens, parce rufo-setosus, segmento anali quadruplo longiore. Stigmata testacea, elevata. Long 10 lin. — Vicinitas autumnus et hyeme in fimo et sub vegetabilibus putridis. Dipteriorum larvis aliisque insectis strenue insidiatur. Carne etiam crudo pascitur. (Bouché Hist. nat. Ins. I. 179. I. t. 7. f. 29 — 41.)

8. Frusta conatus genus hoc speciebus copiosum, corporis habitu et pluribus characteribus quodammodo instabile, in sectiones familiares naturales distribuere, divisiones a Gyllenhalio primo propositas secundum thoracis puncturam recipere non dubitavi, quamquam non solum species maxime affines divelluntur, sed punctorum etiam in thoracis seriebus dorsalibus numerus in quibusdam individuis aut minutus aut auctus observatur. Cum autem talia individua rarius occurrant, ut punctorum illorum numerum omnino inconstantem et variabilem censeamus, non

est, quod divisiones ex iis derivatas relinquam: speciebus facilius eruendis sine dubio prosunt.

1) Thorax punctorum seriebus dorsalibus nullis.

\* 1. *Ph. splendens*: *Niger, nitidus, elytris virescenti-aeneis, thorace lateribus subsinuato, disco laevissimo.* — Long.  $4\frac{1}{2}$  — 6 lin.

*Nordm. Symb.* 73. 1. — *Er. Col. March.* I. 447. 3.

*Staph. splendens* *Fab. Ent. Syst.* I. II 523. 19. *Syst. El.* II. 594. 21. — *Payk. Faun. Suec.* III. 373. 9. — *Panz. Ent. Germ.* 353. 13. — *Cederh. Faun. Ingr.* 118. 359. — *Grav. Micr.* 16. 16. *Mon.* 87. 82. — *Latr. Hist. nat. Crust. et Ins.* IX. 306. 23. — *Gyll. Ins. Suec.* II. 297. 16. — *Sahlb. Ins. Fenn.* I. 313. 14. — *Zetterst. Faun. Lappon.* I. 69. 8. *Ins. Lappon.* 60. 9. — *Mannerh. Brachelytr.* 25. 24. — *Boisd. et Lacord. Faun. Ent. Paris* I. 387. 3. — *Runde Brachelytr. Hal.* 5. 18.

*Staph. aeneus* *Müll. Prodr.* 98. 1110. — *Harrer Besch. d. Ins* I. 251. 413.

*Staph. sextus* *Schöff. Icon.* I. t. 30. f. 12.

*Staph niger* *Müll. Faun. Frid.* 23. 229. *Prodr.* 98. 1109. — *Goeze Ent. Beitr.* I. 726. 20. (*forte*)

Habitat in Europa.

*Ph. aeneo* affinis, plerumque autem maior, niger, nitidus, elytris aeneis. Antennae capite sesqui longiores, articulo tertio modice elongato, secundo paulo longiore, sequentibus sensim paululum brevioribus, intermediis crassioribus, ultimo ovato, apice oblique emarginato, totae nigrae. Palpi nigri, apice picei. Caput feminae thoraci latitudine aequale, maribus maioribus eodem latius, supra parum convexum, pone oculos utrinque parce fortiterque punctatum, inter oculos punctis 4 transversim positis foveolaque minuta obsoletaque media notatum. Thorax coleopteris paulo angustior, latitudine fere brevior, basin versus subangustatus, basi rotundatus, lateribus anterioribus rotundatis, pone medium subsinuatis, apice truncatus, angulis anterioribus subrectis, posterioribus obtusis, modice convexus, disco impunctatus, lateribus utrinque punctis duobus tribusve impressus. Scutellum nigrum, confertim punctatum. Elytra thorace longiora, densius punctata, nigro-pubescentia. Abdomen parcius punctulatum, parcius nigro pubescens, ano longius piloso. Pedes nigri, tarsis apice piceis, tibiis omnibus densius spinulosis, coxis intermediis distantibus.

Tarsi antici maris vix dilatati, feminae simplices.

Obs. Habitus omnino *Ph. aenei*, plerumque autem maior et thoracis disco impunctato distinctus.

\* 2. *Ph. intermedius*: *Niger, nitidus, capitè thoraci latitudine aequali, thoraceque lateribus subrotundato viridi-aeneis, nitidissimis, elytris aeneis, parcius profundiusque punctatis.* — Long. 4 — 5 lin.

*Er. Col. March.* I. 447. 2.

*Staph. intermedius* *Boisd. et Lacord. Faun. Ent. Paris.* I. 338. 4.

*Philonth. laminatus* *Nordm. Symb.* 74. 3.

Habitat in Germania, Gallia, Italia, Graecia, Mesopotamia.

*Intermedius* quasi inter *Ph. laminatum* et *splendentem*, niger, nitidus, antennis pedibusque concoloribus, capite thoraceque viridi-aeneis, nitidissimis, elytris aeneis, abdomine dorso subaeneo-nitidulo. Antennae capite duplo fere longiores, articulo tertio modice elongato, secundo paulo lon-

giore, sequentibus sensim paulo brevioribus, penultimis leviter transversis, ultimo apice profunde emarginato. Palpi apice picei. Caput thoracis latitudine, vel maribus maioribus eodem latius, omnium subtilissime sparsim punctulatum, supra oculos utrinque inaequaliter punctatum, inter oculos punctis 4 transversim positis maiusculis impressum. Thorax coleopteris paulo angustior, latitudine vix brevior, antrorsum haud angustatus, basi fortius, lateribus leviter rotundatis, his ante basin subsinuatis, apice truncatus, angulis posterioribus obtusis, anterioribus deflexis, acutiusculis, parum convexus, ad latera utrinque punctis tribus impressus. Scutellum aeneum, dense punctatum. Elytra thorace paulo longiora, parcius punctata. Abdomen parce punctulatum, longius nigro-pubescent. Pedes tibiis omnibus densius spinulosis, tarsis anticis in utroque sexu modice, licet in mare fortius dilatatis; coxis intermediis leviter distantibus.

Abdominis segmentum inferius quartum in mare apice prolongatum, quinti maximam partem obtegens, in femina simplex.

Obs. Ph. laminato affinis, distinctus autem capite maiore, thorace apicem versus haud angustato, elytris fortius et minus confertim punctatis et colore magis aeneo.

\* 3. *Ph. laminatus*: Niger, nitidus, capite thorace angustiore, thoraceque lateribus rotundato viridi-aeneis, nitidissimis, elytris viridibus, confertim subtiliter punctatis. — Long.  $4\frac{1}{2}$  lin.

*Er. Col. March. I.* 446. 1.

*Staph. laminatus* Creutz. *Ent. Vers.* 128. 9. t. 3. f. 31. — *Panz. Faun. Germ.* 67. 20. — *Grav. Micr.* 16. 17. *Mon.* 86. 81. — *Latr. Hist. nat. Crust. et Ins.* IX. 307. 24. — *Gyll. Ins. Suec. II.* 298. 17. — *Sahlb. Ins. Fenn. I.* 313. 15. — *Zetterst. Faun. Lappon. I.* 70. 9. *Ins. Lappon.* 60. 10. — *Mannerh. Brachelytr.* 25. 25. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 389. 5. — *Runde Brachelytr. Hal.* 5. 19.

*Philonth. viridanus* Nordm. *Symb.* 74. 3.

*Staph. aeneus* Marsh. *Ent. Britt.* 511. 37.

Habitat in Europa.

Niger, nitidus, antennis palpis pedibusque concoloribus, capite thoraceque viridi-aeneis, nitidissimis, elytris viridibus, minus nitidis. Antennae capite plus duplo longiores, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus, penultimis autem haud transversis, ultimo ovato, apice emarginato. Caput in utroque sexu thorace angustius, suborbiculatum, omnium subtilissime punctulatum, utrinque supra oculos fortiter punctatum, inter oculos punctis 4 transversim positis impressum. Thorax coleopteris parum angustior, latitudine paululum brevior, basi lateribusque aequaliter rotundatus, antrorsum leviter angustatus, apice subemarginatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, supra leviter convexus, latera versus utrinque punctis tribus impressus. Scutellum dense punctatum. Elytra thorace paulo longiora, confertim subtiliter punctata, nigro-pubescentia. Abdomen subtiliter minus confertim punctulatum, parce nigro-pubescent. Pedes tibiis omnibus dense spinulosis, tarsis anticis in utroque sexu leviter dilatatis, coxis intermediis leviter distantibus.

Abdominis segmentum inferius quartum maris apice prolongatum, quintum fere omnino obtegens, feminae simplex.



Variat elytris cyanescentibus.

Obs. *Staph. splendens* var. *Gray. Mon. p. 87.* ad hanc varietatem pertinet

- \* 4. *Ph. cribratus*: *Niger, nitidus, pedibus piceis, elytris crebre profundeque punctatis, thorace laevissimo, capite orbiculato.* — *Long. 4—5 lin.*

Habitat in Istria, Gallia meridionali, Sardinia, Lusitania.

Statura omnino *Ph. atrati*, at paulo maior, niger, subnitidus, capite thoraceque nitidissimis. Antennae capite cum thorace paulo breviores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus, penultimis haud transversis, ultimo apice oblique truncato et inferne acuminato, nigropiceae, apice ferrugineae. Palpi et mandibulae picei. Caput thoracis latitudine, orbiculatum, lateribus supra utrumque oculum parce punctatum, inter oculos punctis 4 transversim positis impressus. Thorax coleopteris angustior, latitudine paulo longior, neque basin neque apicem versus angustatus, lateribus anterieus leviter rotundatis, posteribus subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus subrotundatis, posterioribus obtusis, convexus, intra angulum anteriorem punctis nonnullis impressus. Scutellum crebre punctatum. Elytra thorace paulo longiora, sat crebre profundeque punctata, parce pubescentia. Abdomen minus dense subtilius punctatum, parce pubescens. Pedes fusco-picei, tibiis posterioribus parcius spinulosis, tarsis anticis in utroque sexu fortiter dilatatis, coxis intermediis approximatis.

## 2) Thorax dorso utrinque 1-punctatus.

- \* 5. *Ph. ruficollis*: *Niger, nitidus, thorace anoque rufis.* — *Long. 4 lin.*

*Staph. ruficollis Gray. Mon. 71. 58.*

Habitat in Terra Van-Diemenii, Dom. Schayer.

Niger, nitidus. Antennae capite plus sesqui longiores, articulo tertio secundo vix longiore, 4—10 subcylindricis, sensim paulo brevioribus et crassioribus, penultimis crassitie haud brevioribus, ultimo ovato, apice abrupte acuminato, nigro-piceae, articulis ultimis duobus ferrugineis. Palpi nigri, apice summo testaceo. Labrum transversum, integrum. Caput thorace paulo angustius, suborbiculatum, basi haud constrictum, supra oculos punctis utrinque tribus, inter oculos puncto utrinque singulo in ipso oculi margine interiore impressum, ceterum laevissimum. Thorax basi coleopterorum latitudine, antrorsum modice angustatus, latitudine brevior, lateribus leviter, basi cum angulis posterioribus fortius rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, transversim convexus, punctis nonnullis marginalibus et in disco ante medium punctorum pari singulo notatus, ceterum laevissimus, saturate rufus seu sanguineus, nitidissimus. Scutellum obscure sanguineum, crebrius punctatum, nigro-pubescent. Elytra thorace paulo longiora, minus confertim fortius punctata, parcius nigro-pubescentia, nigerrima, nitida. Abdomen minus subtiliter crebrius punctatum, nigro-pubescent, segmento quinto margine apicali sextoque toto rufis. Pedes nigri, coxis anterioribus cum trochanteribus sanguineis, coxis intermediis approximatis, tibiis omnibus parce spinulosis, tarsis piceis, anticis in utroque sexu dilatatis.

Obs. Habitus omnino *Quedii*, at stigmata antica haud oblecta.

- \* 6. *Ph. hybridus*: *Subdepressus, niger, capite subquadrato thoraceque nitidissimis, antennarum basi elytrorumque margine apicali obscure testaceis, pedibus piceis.* — Long.  $2\frac{1}{2}$  lin.

Habitat in Terra Van-Diemenii, Dom. Schayer.

Corpus depressiusculum, nigrum, subnitidum, capite thoraceque nitidissimis. Antennae capite sesqui prope longiores, articulo tertio secundo paulo tenuiore et vix longiore, sequentibus sensim paulo brevioribus, vix autem crassioribus, penultimis leviter transversis, ultimo oblongo-ovato, apice abrupte acuminato, nigrae, articulis duobus primis testaceis. Palpi picei, maxillares articulo quarto tenuiore, acuminato. Labrum breve, medio leviter sinuatum. Caput sat magnum, thoracis latitudine, subquadratum, basi distincte constrictum, subdepressum, supra oculos punctis utrinque tribus maioribus, inter oculos puncto utrinque singulo in ipso oculi margine interiore impressum, ceterum laevissimum, infra parce punctatum. Thorax coleopteris paulo angustior, basin versus angustatus, latitudine antica paulo brevior, lateribus modice, basi cum angulis posterioribus fortius rotundatus, apice truncatus, angulis anterioribus deflexis, acutiusculis, leviter transversim convexus, praeter puncta marginalia punctorum pari discoidali singulo ante medium impressus. Scutellum crebre punctatum, cinereo-pubescent. Elytra thorace sesqui longiora, depressa, crebrius subtiliter punctata, cinereo-pubescentia, margine apicali indeterminate testaceo. Abdomen apicem versus sensim angustatum, crebre minus subtiliter punctatum, cinereo-pubescent, lateribus parce nigro-pilosellum. Pedes breviusculi, picei, apice dilutiores, coxis intermediis approximatis, tibiis posterioribus parcius spinulosis.

Mas tarsis anticis modice dilatatis, intermediis articulo primo incrassato. Femina latet.

### 3) Thorax seriebus dorsalibus 3-punctatis.

- \* 7. *Ph. cinctulus*: *Niger, nitidus, antennis pedibusque piceis, elytris aeneis, capite orbiculato.* — Long. 5 lin.

*Staph. cinctulus* Grav. *Micr.* 167. 16. *Mon.* 89. 84.

Habitat in Bengala, Dom. Daldorf.

Statura et magnitudo *Ph. intermedi*, niger, nitidus, capite thoraceque nitidissimis, elytris aeneis. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus, vix autem crassioribus, ultimo ovato, apice abrupte acuminato, ferrugineae, articulis tribus primis piceo-rufis. Caput thorace vix angustius, suborbiculatum, utrinque supra oculos inaequaliter punctatum, inter oculos punctis utrinque duobus transversim positis impressum. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus parallelis, anterieus leviter rotundatis, pone medium subsinuatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, rotundatis, transversim convexus, seriebus dorsalibus punctis ternis, duobus prope apicem, tertio longe pone medium compositis, praeterea lateribus punctis pluribus (5—7) impressus. Scutellum nigrum, crebre punctulatum, griseo-pubescent. Elytra thorace paulo longiora, parcius et minus subtiliter punctata, parce griseo-pubescentia. Abdomen metallico-versicolor, subtiliter punctulatum, griseo-pu-

berulum. Pedes rufo-picei, tibiis omnibus crebre spinulosis, coxis intermediis leviter distantibus.

Mas latet. Femina tarsis anticis leviter dilatatis.

Obs. Abdominis cingula albida, quae a Gravenhorstio laudantur, membranis intersegmentariis, quae in hoc specimine extensae erant, exorta sunt.

\* 8. *Ph. laevicollis*: Niger, nitidus, capite, thorace elytrisque nigro-aeneis, abdominis segmentorum marginibus pedibusque piceis, elytris parce obsoleteque punctatis. — Long.  $3\frac{1}{2}$  — 4 lin.

*Staph. laevicollis* Boisd. et Lacord. Faun. Ent. Paris. I. 397. 21.

Habitat in Carinthia, Dom. Schüppel.

Statura omnino *Ph. atrati*, niger, capite thoraceque supra nigro-aeneis, pernitidis, elytris aeneis, nitidulis. Antennae capitis thoracisque prope longitudine, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paululum brevioribus et crassioribus, penultimis crassitie haud brevioribus, ultimo subovato, apice oblique emarginato, nigrae. Palpi picei. Caput thorace paulo angustius, subovatum, parce omnium subtilissime punctulatum, supra oculos utrinque punctis nonnullis distantibus, inter oculos punctis utrinque duobus transversim positis impressum. Thorax coleopteris paulo angustior, latitudine vix longior, lateribus leviter rotundatus, apicem versus paululum magis quam basin versus angustatus, basi rotundatus, apice leviter late emarginatus, angulis anterioribus subrotundatis, posterioribus obtusis, leviter convexus, seriebus dorsalibus tripunctatis, puncto postremo in disci medio, antico ad marginem anteriorem sito, praeterea ad latera praeter marginalia puncto singulo notatus, punctis omnibus subtilibus. Scutellum punctulatum, nigrum, parce griseo-pubescens. Elytra thorace paulo longiora, confertissime subtiliter alutacea, parce vageque obsolete punctata, lateribus et margine apicali cinereo-pubescentibus. Abdomen crebre minus subtiliter punctatum, pube depressa fusco-grisea densius vestitum, segmentis singulis margine apicali piceis. Pedes picei, tibiis omnibus spinulosis, coxis intermediis modice distantibus.

Tarsi antici maris modice, feminae leviter dilatati.

4) Thorax seriebus dorsalibus 4-punctatis.

X \* 9. *Ph. cyanipennis*: Niger, nitidus, elytris cyaneis, thorace lateribus rotundato, capite suborbiculato. — Long. 5 —  $5\frac{1}{2}$  lin.

*Staph. cyanipennis* Fab. Ent. Syst. I. II. 525. 29. Syst. El. II. 597. 37. — Walcken. Faun. Paris. I. 271. 12. — Grav. Micr. 168. 17. Mon. 86. 78. — Boisd. et Lacord. Faun. Ent. Paris. I. 387. 2. — Ründe Brachelytr. Hal. 7. 35.

*Staph. amoenus* Oliv. Ent. III. 42. 26. 32. t. 4 f. 36. — Latr. Hist. nat. Crust. et Ins. IX. 308. 27.

*Staph. coeruleipennis* Mannerh. Brachelytr. 27. 43.

*Philonth. coeruleipennis* Nordm. Symb. 80. 25.

Habitat in Gallia, in Austria, in Chersoneso Taurica et in Pennsylvania.

Statura omnino *Ph. splendidis*, niger, nitidus. Antennae capite cum thorace paulo breviores, articulo tertio elongato, secundo sesqui longiore, quarto duplo brevior, terete, 5 — 10 subobconicis, sensim paulo breviori-



bus at parum crassioribus, ultimo ovato, apice oblique emarginato et inferne acuminato, nigrae, articulo apicali subferrugineo. Mandibulae et palpi elongati, picei. Caput maris thoracis fere latitudine, transversim quadrato-orbiculatum, feminae thorace angustius, ovali-suborbiculatum, fronte depressa, utrinque supra oculos parce vageque punctatum, inter oculos utrinque subimpressum et punctis duobus transversim positis notatum. Thorax coleopteris paulo angustior, latitudine brevior, basi, angulis posterioribus et lateribus rotundatis, his deflexis, antrorsum subangustatus, apice truncatus, angulis anterioribus obtusis, modice transversim convexus, antice subcompressus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, utrinque praeter margines punctis circiter 5 impressus, omnibus subtilioribus. Scutellum nigrum, crebre punctulatum, tenuiter pubescens. Elytra thorace tertia parte longiora, laete cyanea vel virescenti-cyanea, nitidula, crebre subtiliter punctulata, nigro-pubescencia. Abdomen apicem versus sensim attenuatum, parce subtiliter punctulatum, parce nigro-pubescens, dorso coerulescenti-nitidulum. Pedes graciliores, nigri, coxis intermediis modice distantibus, tibiis omnibus spinulosis, tarsis piceis, anticis in utroque sexu simplicibus.

Obs. Specimina Americana cum Europaeis exacte congruunt, nisi quod, observante D. Prof. Gravenhorst, puncta inter oculos intermedia in illis distincta, in his obsoleta esse solent. Occurrunt inter Europaea individua, quibus hoc punctum in altero latere distinctius, in altero obsoletum est.

\* 10. *Ph. furvus*: *Niger, capite thoraceque nitidissimis, elytris margine inflexo luteo, thorace lateribus subsinuato, capite orbiculato.* — Long. 5 lin.

*Nordm. Symb.* 88. 50.

Habitat in Mexico, Dom. Deppe.

Statura et summa affinitas *Ph. aenei*, niger, nitidulus, capite thoraceque nitidissimis, elytrorum margine laterali inflexo solo luteo. Antennae capituli thoracisque fere longitudine, articulo tertio secundo sesqui longiore, sequentibus sensim paulo brevioribus, subobconicis, penultimis crassite haud brevioribus, ultimo ovato, apice oblique emarginato et inferne acuminato. Caput maiusculum, thorace vix angustius, orbiculatum, utrinque supra oculos punctis pluribus fortioribus, inter oculos utrinque punctis duobus oblique positis. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus leviter rotundatis, postice subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus subrotundatis, posterioribus obtusis, supra leviter convexus, seriebus dorsalibus punctis 4, antico paulo remotiore, submarginali, compositis, lateribus punctis utrinque praeter marginalia 5 impressus, punctis omnibus modice profundis. Scutellum confertissime punctatum, tenuiter nigro-pubescens. Elytra densius subtiliter punctata, nigro-pubescencia. Abdomen parcius punctulatum, parce nigro-pubescens. Pedes nigri, coxis intermediis modice distantibus, tibiis omnibus spinulosis.

Tarsi antici maris leviter dilatati, feminae simplices.

\* 11. *Ph. nigerrimus*: *Niger, capite thoraceque nitidissimis, elytris creberrime subtiliter punctatis, thorace lateribus subsinuato, capite orbiculato.* — Long. 5 lin.

Habitat in Mexico, Dom. C. Ehrenberg.

Statura omnino Ph. aenei, totus nigerrimus, capite thoraceque nitidissimis. Antennae capite cum thorace breviores, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus, prioribus eorum subcylindricis, penultimis subobconicis, his crassitie haud brevioribus, ultimo subovato, apice subemarginato et inferne acuminato, totae nigrae. Caput thorace vix angustius, orbiculatum, utrinque supra oculos punctis nonnullis et inter oculos punctis 4 transversim positissimis impressum. Thorax coleopteris parum angustior, latitudine vix brevior, antrorsum haud angustatus, lateribus rotundatis, postice subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, supra leviter transversim convexus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus utrinque punctis circiter 4 praeter marginalia impressus, omnibus subtilioribus. Elytra thorace paulo longiora et scutellum parum nitida, subtiliter confertissime punctata, nigro-pubescentia. Abdomen nitidulum, densius subtiliter punctulatum, nigro-pubescentia. Pedes concolores, coxis intermediis parum distantibus, tibiis omnibus spinulosis.

Tarsi anticis maris modice dilatati. Femina latet.

\* 12. *Ph. testaceipennis*: Nigerrimus, nitidus, elytris rufo-testaceis, thorace lateribus subsinuato, abdomine supra parce punctato. — Long. 5 lin.

Habitat in Mexico.

Habitu omnino Ph. nitidi, nigerrimus, nitidus, elytris solis rufo-testaceis. Antennae capite sesqui longiores, medio subincrassatae, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo brevioribus, penultimis haud transversis, ultimo oblongo-ovato, apice emarginato. Mandibulae parum prominentes. Caput thoracis prope latitudine, suborbiculatum, inter oculos punctis 4 transversim positissimis, supra oculos utrinque punctis nonnullis sparsis impressum. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus leviter rotundatis, postice subsinuatis, basi rotundatus, apice leviter emarginatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, supra leviter convexus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus punctis utrinque 3 praeter marginalia impressus, omnibus minus subtilibus. Scutellum crebre subtiliter punctatum, nigro-pubescentia. Elytra thorace paulo longiora, subtilius minus dense punctata, rufo-testacea, nitida, tenuiter fulvo-pubescentia. Abdomen parce leviter punctatum, parce nigro-pubescentia, apice densius longiusque nigro-pilosum. Pedes nigri, coxis intermediis modice distantibus, tibiis omnibus spinulosis.

Tarsi antici feminae simplices. Mas latet.

\* 13. *Ph. nitidus*: Niger, nitidus, elytris rufis, thorace lateribus subsinuato, abdomine supra crebre punctato. — Long. 5 — 6 lin.

Nordm. Symb. 90. 52. — Er. Col. March. I. 448. 4.

Staph. nitidus Fab. Mant. Ins. I. 220. 13. Ent. Syst. I. II. 524. 24. Syst. El. II. 596. 30. — Payk. Mon. Staph. 40. 30. Faun. Suec. III. 390. 30. — Panz. Ent. Germ. 354. 16. — Cederh. Faun. Ingr. 118. 361. — Schrank Faun. Boica I. 640. 864. — Gyll. Ins. Suec. II. 315. 31. — Zetterst. Faun. Lappon. I. 74. 17. Ins.

*Lappon.* 63. 23. — *Sahlb. Ins. Fenn.* I. 320. 29. — *Germ. Faun. Ins. Europ.* XIII. 12. — *Mannerh. Brachelytr.* 27. 42.

*Staph. coenosus* Grav. *Mon.* 51. 15. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 386. 1.

Habitat in Europa.

Statura omnino *Ph. aenei* et maximis eius individuis magnitudine aequalis, niger, nitidus, elytris testaceo-rufis. Antennae capite cum thorace breviores, articulo tertio elongato, secundo paulo longiore, 4—10 sensim paulo brevioribus, subobconicis, penultimis leviter transversis, ultimo apice emarginato et inferne acuminato. Caput orbiculatum, feminae thoracis latitudine, maris eodem paulo latius, supra oculos utrinque dense punctatum, fronte utrinque punctis paucis fortioribus et inter oculos punctis 4 transversim positis impressa. Thorax coleopteris parum angustior, latitudine paulo brevior, lateribus antice rotundatis, postice subsinuatis, basi modice rotundatus, apice truncatus, angulis anterioribus deflexis, obtusiusculis, posterioribus obtusis, transversim convexus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus utrinque punctis circiter 9 praeter marginalia notatus, omnibus fortioribus. Scutellum disco confertim punctatum. Elytra thorace paulo longiora, confertim subtiliterque punctata, rufescenti-pubescentia. Abdomen densius punctatum, fusco-pubescent. Pedes nigri, coxis intermediis modice distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

Variat thoracis seriebus dorsalibus vel utrisque 5-punctatis, vel altera 4-, altera 5-punctata.

\* 14. *Ph. asphaltinus*: *Niger, nitidus, tibiis tarsisque rufo-piceis, elytris subtiliter punctatis, thorace lateribus subsinuato, capite magno subquadrato.* — *Long.* 5 lin.

Habitat in Germania meridionali, Dom. Godet.

Statura *Ph. aenei*, at paulo gracilior. Corpus totum cum elytris nigerrimum, nitidum, capite thoraceque nitidissimis. Antennae capite duplo longiores, articulo tertio secundo sesqui longiore, 4—10 subobconicis, sensim paulo brevioribus at vix crassioribus, ultimo apice oblique leviter emarginato et inferne fortiter acuminato, nigro-piceae, articulo ultimo ferrugineo. Mandibulae et palpi picei. Caput magnum, thoracem latitudine superans, subquadratum, utrinque supra oculos vage subtiliterque punctatum, fronte inter oculos superius punctis 6, arcuatim dispositis, anterie punctis 4 transversim positis, omnibus subtilioribus, impressa. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus leviter subrotundatis, postice subsinuatis, basi leviter rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, leviter transversim convexus, seriebus dorsalibus punctis 4 distantibus compositis, lateribus punctis circiter 7 praeter marginalia impressus, omnibus sat subtilibus. Scutellum subtiliter punctatum. Elytra thorace multo longiora, minus dense subtiliter punctata, subtiliter parceque nigro-pubescentia. Abdomen parce subtiliter punctatum, tenuiter nigro-pubescent. Pedes graciles, nigri, coxis intermediis perparum distantibus, tibiis omnibus parce subtiliterque spinulosis tarsisque piceo-rufis.

Tarsi antiqui maris simplices. Femina latet.



- \* 15. *Ph. carbonarius*: *Niger, nitidus, elytris nigro-aeneis, capite subquadrato, thorace lateribus subsinuato, utrinque impresso, abdomine parcius punctato, parce nigro-pubescente.* — Long. 4 — 5½ lin.

*Er. Col. March. I. 448. 5.*

*Staph. nigrinus Runde Brachelytr. Hal. 7. 33.*

*Var. Pedibus piceis, abdominis segmentis ventralibus rufo-marginatis.*

*Staph. carbonarius Gyll. Ins. Suec. II. 319. 35. — Mannerh. Brachelytr. 28. 15.*

Habitat in Europä.

Statura et summa similitudo *Ph. aenei*. *Niger, nitidus*. Antennae capite thoraceque breviores, articulis tribus primis modice elongatis, tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, subobconicis, penultimis leviter transversis, ultimo apice emarginato et inferne acuminato, nigrae, articulo apicali ferrugineo, interdum totae ferrugineae. Palpi nigri. Caput maris thoracis latitudine vel eodem paulo maius, subquadratum, feminae thorace angustius, orbiculato-subquadratum, supra oculos utrinque vage fortiter punctatum, fronte anterieus inter oculos utrinque punctis duobus transversim positis impressa. Thorax coleopteris paulo angustior, latitudine vix brevior, antrorsum subangustatus, lateribus antrorsum leviter rotundato-ampliatus, pone medium subsinuatus, basi rotundatus, apice truncatus, angulis omnibus rotundatis, leviter convexus, versus latera utrinque oblique leviter impressus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis omnibus minus profundis. Scutellum dense punctatum, nigrum. Elytra thorace paulo longiora, crebre punctata, nigro-aenea, nigro-pubescentia. Abdomen dorso parcius punctulatum, parce nigro-pubescentia, subaeneo-, interdum etiam metallico-versicolor. Pedes graciliores, coxis intermediis modice distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

Abdominis segmentum inferius quintum in mare apice leviter emarginatum.

Obs. 1. Cum *Ph. aeneo* plerumque confusus, distinctus autem statura saepius paulo minore, elytris obscurioribus abdomineque parcius distinctiusque punctulato nigro-pubescentibus, tarsisque anticis in utroque sexu simplicibus. Caput maris speciminibus maioribus maius, minoribus minus, ut in *Ph. aeneo*.

2. Nomen a Gyllenhalio usitatum retinui, quamvis *Staph. carbonarius* Grav. ad aliam speciem, *Ph. varium*, pertinet. Specimen a Gyllenhalio ipso e Suecia sibi missum dedit Dom. Schüppel.

- \* 16. *Ph. aeneus*: *Niger, capite thoraceque nigro-aeneis, elytris viridi-aeneis, capite subquadrato, thorace lateribus subsinuato, utrinque leviter impresso, abdomine subtilius punctulato, cinereo-pubescente.* — Long. 4 — 6 lin.

*Nordm. Symb. 81. 26. — Er. Col. March. I. 449. 6.*

*Staph. aeneus Rossi Faun. Etrusc. I. 249. 613. Ed. Hellw. I. 307. 613. — Grav. Micr. 17. 18. Mon. 86. 80. — Latr. Hist. nat. Crust. et Ins. IX. 307. 25. — Gyll. Ins. Suec. II. 314. 30. — Sahlb. Ins. Fenn. I. 319. 28. — Zetterst. Faun. Lappon. I. 73. 16. Ins. Lappon. 62. 22. — Mannerh. Brachelytr. 27. 48. — Botsd. et Lacord. Faun. Ent. Paris. I. 389. 6. — Runde Brachelytr. Hal. 6. 27.*

*Staph. laticeps Zetterst. Faun. Lappon. I. 73. 15. Ins. Lappon. 62. 21.*

*Staph. metallicus* Boisd. et Lacord. Faun. Ent. Paris. I. 390. 7.

*Staph. similis* Marsh. Ent. Britt. 497. 3.

Var. *Pedibus rufo-piceis*.—

*Staph. atratus* Boisd. et Lacord. Faun. Ent. Paris. I. 392. 11.

Habitat in Europa Asiaque confini et in America septentrionali frequens.

Niger, capite thoraceque nigro-aeneis, nitidis, elytris viridi-aeneis. Antennae capite thoraceque paulo longiores, articulis tribus primis modice elongatis, tertio secundo paulo longiore, 4—10 subobconicis, sensim paulo brevioribus et crassioribus, penultimis transversis, ultimo subovato, apice emarginato et inferne acuminato, nigrae, articulo apicali ferrugineo, interdum fusco. Caput maribus maioribus magnum, thorace latius, transversim subquadratum, minoribus minus, interdum vix thoracis latitudine, pone oculos minus angulatum, feminae thorace paulo angustius, orbiculato-subquadratum, supra oculos vage fortiter punctatum, inter oculos punctis 4 transversim positis impressum. Thorax coleopteris paulo angustior, latitudine paulo brevior, antrorsum subangustatus, lateribus ante medium leviter rotundato-ampliat, pone medium subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus subrotundatis, posterioribus obtusis, parum convexus, versus latera utrinque oblique leviter impressus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 2, rarius 3 vel 4 impressus, punctis omnibus modice profundis. Scutellum nigrum, dense subtiliterque punctatum, griseo-pubescens. Elytra thorace paulo longiora, dense punctata, griseo-pubescentia. Abdomen densius subtiliter punctatum, griseo-pubescens. Pedes nigri, tibiis omnibus spinulosis tarsisque saepe piceis, coxis intermediis modice distantibus.

Abdominis segmentum inferius quintum maris apice leviter emarginatum. Tarsi antici maris fortiter dilatati, feminae simplices.

Obs. 1. *Staph. aeneus* auctorum plerorumque ad specimina maiora mascula capite grandi spectat, *Staph. laticeps* Zett. ad eadem, *Staph. aeneus* Zetterst. contra et *Staph. metallicus* Boisd. et Lacord. ad individua minora mascula capite thorace haud latiore pertinent. Pro *Staph. atrato* auctores modo citati huius speciei specimen femineum, pedibus rufo-piceis, descripsisse videntur.

2. Specimen a Gravenhorstio Mon. p. 52. pro *Staph. coenosi* varietate? descriptum, e collectione citata conservatum, huius speciei individuum est immaturum, elytris pedibusque fuscis.

\* 17. *Ph. scutatus*: Niger, capite orbiculato, thorace elytrisque obscure aeneo-virescentibus, nitidis, pedibus piceis, elytris dense subtiliter punctatis. — Long. 6 lin.

Habitat in Germania meridionali, Hungaria, Sardinia, Persia.

Statura omnino *Ph. atrati*, at paulo maior, niger, capite, thoraceque nigro-virescentibus, nitidissimis, elytris concoloribus, nitidis. Antennae capite thoraceque parum breviores, graciliores, articulo tertio secundo perparum longiore, sequentibus sensim paulo brevioribus, subcylindricis, reliquis subobconicis, penultimis crassitie haud brevioribus, ultimo oblongo-ovato, apice inferne abrupte acuminato, nigrae. Palpi picei. Caput orbiculatum, thorace paulo angustius, leviter convexum, parce subtilissime punctatum, supra oculos utrinque punctis nonnullis maioribus, inter oculos punctis 4 transversim positis impressum. Thorax coleopteris paulo angustior, latitudine paulo

brevior, antrorsum subangustatus, basi et lateribus leviter rotundatis, his postice subsinuatis, apice truncatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, supra modice convexus, omnium subtilissime parcius punctulatus, seriebus dorsalibus punctis 4, antico et postico paulo remotioribus, compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis impressis subtilioribus. Scutellum nigrum, creberrime punctulatum, nigro-pubescentia. Elytra thorace tertia parte longiora, dense subtiliter punctata, longius griseo-pubescentia. Abdomen punctulatum, longius griseo-pubescentia. Pedes picei, coxis intermediis approximatis, tibiis omnibus spinulosis, tarsis anticis in utroque sexu leviter dilatatis.

Abdominis segmentum inferius quartum in mare apice prolongatum, lamina eius segmentum quintum obtegente.

Obs. 1. Abdominis maris structura omnino cum Ph. laminato et intermedio congruens, at thorax seriebus dorsalibus 4-punctatis instructus.

2. Specimen in Persia septentrionali prope Lankoran ad littora maris Caspici occidentalia lectum Dom. Ménétries pro Staph. rotundicollis suo communicavit, at descriptio eius (Catal. rais. 145. 581.) nimis manca, et nomine eodem a Com. Dejean in Catalogo collectionis suae apud Ph. carbonarium citato, sub nomine illo plures commutatas fuisse species suspicandum esse videtur.

\* 18. *Ph. atratus*: *Niger, nitidus, elytris aeneo-viridibus, parcius punctatis, capite orbiculato, thorace paulo minore.* — Long. 3 — 4 lin.

*Nordm. Symb.* 82. 30. — *Er. Col. March.* I. 450. 7.

*Staph. atratus* Grav. *Micr.* 21. 23. *Mon.* 84. 74. — *Latr. Hist. nat. Crust. et Ins.* IX. 310. 29. — *Gyll. Ins. Suec.* II. 319. 34. — *Sahlb. Ins. Fenn.* I. 321. 32. — *Zetterst. Ins. Lappon.* 83. 25. — *Mannerh. Brachelytr.* 28. 29.

*Staph. metallicus* Grav. *Micr.* 168. 78. — *Latr. Hist. nat. Crust. et Ins.* IX. 323. 64.

*Var. Elytris viridi-coeruleis.*

*Staph. coerulescens* Boisd. et Lacord. *Faun. Ent. Paris.* I. 393. 13.

Habitat in Europa.

Ph. polito paulo minor et angustior, niger, capite thoraceque nitidissimis, interdum subaeneis, elytris metallico-viridibus, antennis, palpis, pedibusque nigris. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo brevioribus at vix crassioribus, penultimis crassitie haud brevioribus, ultimo ovato, apice truncato et inferne acuminato. Caput in utroque sexu thorace minus et angustius, orbiculatum, supra leviter convexus, pone oculos utrinque parce fortiter punctatum. Thorax basi fere coleopteris latitudine aequalis, antrorsum subangustatus, latitudine paulo brevior, basi modice, lateribus leviter rotundatis, postice subsinuatis, apice truncatus, angulis anterioribus rotundatis, posterioribus obtusis, leviter convexus, seriebus dorsalibus punctis aequaliter distantibus, licet antico paulo magis extrorsum sito, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus fortioribus. Scutellum nigrum, confertissime punctatum, parce griseo-pubescentia. Elytra thoracis longitudine, minus subtiliter parcius punctata, parce griseo-pubescentia. Abdomen supra parcius punctulatum, parce griseo-pubescentia. Pedes coxis intermediis approximatis, tibiis omnibus spinulosis, tarsis anticis in utroque sexu, licet in mare fortius dilatatis.

Variat elytris magis minusve coerulescentibus.



Obs. 1. Ab affinibus praecipue elytrorum punctura minus conferta, fortiore distinctus.

2. Staph. metallicus Grav., cuius specimen singulum typicum benevole mihi communicavit celeberrimus auctor, nonnisi thoracis serie altera 5-punctata a Ph. atrato genuino differt.

\* 19. *Ph. atramentarius: Nigerrimus, nitidus, femoribus piceis, capite ovato, thorace oblongo, antrorsum subangustato, abdomine confertim subtiliter punctato. — Long. 4 lin.*

Habitat in Columbia, Dom. Moritz.

Ph. polito gracilior, nigerrimus, nitidus, capite thoraceque pernitidis. Antennae capite thoraceque prope longiores, minus graciles, articulo tertio secundo sesqui prope longiore, 4 — 10 subcylindricis, sensim paulo brevioribus, at non crassioribus, omnibus crassitie multo longioribus, ultimo oblongo-ovato, apice emarginato et inferne acuminato, nigrae. Palpi picei. Caput thorace paulo angustius, subovatum, leviter convexum, utrinque supra oculos punctis parvis, inter oculos utrinque punctis duobus transversim positis impressum. Thorax coleopteris duplo prope angustior, latitudine sesqui longior, apicem versus subangustatus, lateribus rectis, ad apicem subrotundatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, leviter convexus, seriebus dorsalibus punctis 4 remotis et aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 4 impressus. Scutellum subtiliter punctatum, tenuiter nigro-pubescent. Elytra thorace paulo longiora, crebrius subtiliter punctata, densius subtiliter nigro-pubescentia. Abdomen supra dense subtiliter punctatum, densius subtiliter nigro-pubescent, subtus lucidum, parcius subtiliter punctatum, parce nigro-pubescent. Pedes graciliores, coxis intermediis approximatis, coxis anterioribus testaceo-piceis, femoribus piceis, tibiis omnibus spinulosis. Alae fusco-hyalinae.

Tarsi antiqui maris leviter dilatati. Femina latet.

\* 20. *Ph. coelites: Niger, nitidus, capite ovato thoraceque violaceis, elytris parcius punctatis, cyaneis, abdomine parce punctato. — Long. 4 lin.*

Habitat in Brasilia, Dom. Prof. Germar.

Statura omnino Ph. flagrantis, Ph. decoro minor et gracilior, niger, nitidus, capite thoraceque supra violaceis, pernitidis, elytris laete nitideque cyaneis. Antennae capite thoraceque vix breviores, articulo tertio secundo paululum longiore, 4 — 10 sensim paulo brevioribus et crassioribus, penultimo crassitie haud longiore, ultimo ovato, apice emarginato et inferne acuminato, nigrae. Palpi nigri. Caput thorace angustius, oblongo-ovatum, utrinque supra oculos punctis nonnullis, inter oculos puncto singulo ad utriusque oculi marginem anteriorem notatum, apice sulco abbreviato impressum. Thorax coleopteris duplo prope angustior, latitudine sesqui longior, antrorsum vix angustatus, lateribus rectis, antrorsum subrotundatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, leviter convexus, seriebus dorsalibus punctis 4 sat remotis, aequaliter distantibus, compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis omnibus subtilioribus. Scutellum nigrum, nitidum, parce punctatum. Elytra thorace paulo longiora, parcius subtilius-

que punctata. Abdomen apicem versus sensim angustatum, parce subtilius punctatum, parce nigro-pubescenti, lateribus et apice nigro-pilosellum. Pedes graciles, nigri, coxis intermediis modice distantibus, tibiis omnibus spinulosis, tarsis piceis.

Tarsi antici feminae simplices. Mas latet.

\*21. *Ph. flagrans*: Niger, nitidus, capite thoraceque nigro-violaceis, elytris viridi-cupreis, splendidis. — Long.  $4\frac{1}{2}$  lin.

Habitat in Mexico; in Columbia etiam a Dom. Moritz lectus.

Statura omnino et summa affinitas praecedentis, nigerrimus, nitidus, capite thoraceque supra nigro-violaceis, nitidis, elytris viridi-aeneis, dorso cupreis, splendidis. Antennae capite thoraceque parum breviores, apicem versus subincrassatae, articulo secundo tertio vix brevior, 4 — 10 subcylindricis, sensim paulo brevioribus et crassioribus, penultimis crassitie haud longioribus, ultimo ovato, apice leviter emarginato et subacuminato, nigrae. Palpi nigri. Caput thoracis apice vix angustius, subovatum, utrinque supra oculos punctis paucis, inter oculos utrinque puncto singulo notatum, fronte antius utrinque secundum oculi marginem lateralem leviter longitudinaliter impressa, apice medio profunde sulcata. Thorax coleopteris multo angustior, antrosum subangustatum, latitudine paulo longior, lateribus subrectis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, posterioribus obtusis, parum convexis, seriebus dorsalibus punctis 4 distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum nigrum, subtiliter parce punctulatum, tenuiter pubescens. Elytra thorace vix longiora, minus confertim subtiliter punctata, tenuiter cinereo-pubescentia. Abdomen apicem versus sensim attenuatum, punctis raris obsoletis impressum, latera versus parce pubescens. Pedes elongati, graciles, coxis intermediis leviter distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

\*22. *Ph. ambiguus*: Niger, capite, thorace elytrisque obscure virescentibus, abdomine versicolore, elytris creberrime punctatis, capite minore, suborbiculato. — Long. 4 lin.

Habitat in Columbia, Dom. Moritz.

Statura fere *Ph. politus*. Antennae capite cum thorace paulo breviores, crassiusculae, articulo tertio secundo sesqui prope longiore, 4 — 10 sensim paulo brevioribus et crassioribus, prioribus subcylindricis, penultimis subobconicis, leviter transversis, ultimo ovato, apice oblique emarginato et inferne acuminato, nigrae. Palpi picei. Caput thorace minus et angustius, suborbiculatum, supra leviter convexum, utrinque pone oculos profunde punctatum, antice inter oculos utrinque punctis duobus transversim positae fortibus notatum, nigro-virescenti, nitidum. Thorax coleopteris paulo angustior, latitudine haud brevior, antrosum haud angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus fortiter rotundatis, leviter convexus, nigro-virescenti, politus, seriebus dorsalibus punctis 4, antico paulo remotiore, compositis, lateribus punctis praeter marginalia 4 impressus, punctis tribus posterioribus serierum dorsalium sat fortibus, reliquis modice profundis. Scutellum nigrum, crebre

punctatum, nigro-pubescent. Elytra thorace paulo longiora, nigro-subaenea, subnitida, dense fortius punctata, pube crassiore depressa nigra vestita. Abdomen temere punctatum, pube depressa rara rigida nigra obsitum, nigrum, nitore viridi-aureo splendide versicolor. Pedes nigri, coxis intermediis leviter distantibus, tibiis omnibus spinulosis, tarsis picescentibus, anticis in utroque sexu simplicibus.

23. *Ph. fibularius*: *Niger, capite subovato, thorace elytrisque viridi-aeneis, his creberrime punctatis, abdomine versicolore, tibiis piceo-testaceis.* — Long.  $3\frac{3}{4}$  lin.

Habitat in Mexico, coll. Dom. Chevrolat.

Habitu omnino *Ph. politus*, at paulo minor, praecedente gracilior, niger, nitidus, capite thoraceque nigro-aeneis, pernitidis, elytris virescenti aeneis, nitidulis. Antennae capite thoraceque paululum breviores, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, penultimo crassitie haud longiore, ultimo apice oblique truncato et inferne acuminato, nigrae. Palpi picei. Caput thorace paulo angustius, suborbiculatum, pone oculos utrinque crebre fortius punctatum, inter oculos punctis 4 transversim positus impressum. Thorax coleopteris angustior, latitudine vix longior, lateribus rectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, convexus, seriebus dorsalibus punctis 4, anterioribus tribus magis distantibus, compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis omnibus subtilioribus. Scutellum obscure viridi-aeneum, confertissime punctatum, densius griseo-pubescent. Elytra thorace paulo longiora, minus fortiter dense punctata, densius griseo-pubescentia. Abdomen apicem versus sensim attenuatum, minus dense profunde punctatum, pube depressa longiore et fortiore nigra vestitum, splendidius metallico-versicolor. Pedes graciliores, picei, coxis intermediis modice distantibus, tibiis omnibus spinulosis, testaceis.

Tarsi antici maris leviter dilatati. — Femina latet.

Obs. In specimine singulo descripto in serie thoracis dorsali altera basin versus punctum quintum exstat, a quarto remotum et subtilius, forte fortuitum.

- X \* 24. *Ph. decorus*: *Niger, capite ovato thoraceque nigro-virescentibus, elytris subtiliter punctatis, nigro-aeneis, opacis, abdomine segmentis ventralibus ferrugineo-marginatis.* — Long. 5 lin.

Nordm. Symb. 82. 31. — Er. Col. March. I. 451. 8.

*Staph. decorus* Grav. Micr. 19. 20. Mon. 85. 77. — Gyll. Ins. Suec. II. 316. 32. — Sahlb. Ins. Fenn. I. 320. 30. — Mannerh. Brachelytr. 27. 44. — Boisd. et Lacord, Faun. Ent. Paris. I. 391. 9. — Runde Brachelytr. Hal. 7. 28.

Habitat in Europa.

*Ph. politus* habitu affinis, at paulo maior. Antennae elongatae, graciles, articulo tertio secundo vix sesqui longiore, 4 — 10 sensim paulo brevioribus, licet omnibus crassitie longioribus, ultimo ovato, apice truncato et inferne abrupte acuminato, nigrae. Palpi rufo-picei, basi nigricantes. Corpus nigrum. Caput et thorax nigro-virescentia, subnitida. Caput thorace angustius et minus, subovatum, supra leviter convexum, basi utrinque crebre



punctatum, utrinque supra oculos punctis paucis maioribus, antice inter oculos punctis 4 transversim positis impressum. Thorax coleopteris angustior, latitudine paulo longior, antrorsum subangustatus, basi modice, lateribus leviter rotundatis, his antè basin subsinuatis, apice truncatus, angulis anterioribus deflexis, rotundatis, posterioribus obtusis, leviter convexus, seriebus dorsalibus punctis 4, antico remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilibus. Scutellum nigrum, subtiliter punctulatum, fusco-pubescent. Elytra thorace parum longiora, obscure aenea, opaca, dense subtiliterque punctata, densius subtiliterque fusco-pubescentia. Abdomen supra densius subtiliter punctulatum, tenuiter fusco-pubescent, subtus segmentis singulis apice tenuiter ferrugineo-marginatis. Pedes nigri, coxis intermediis approximatis, tibiis omnibus spinulosus tarsisque rufo-piceis, tarsis anticis in utroque sexu simplicibus.

Varietatem colore capitis thoracisque cupreo, splendido, elytrorum cyanescenti-iridi, opaco insignem Dom. Prof. Gené in Sardinia legit.

\* 25. *Ph. lucens*: Niger, capite ovato, thorace angustiore thoraceque nitidis, elytris confertim subtilius punctatis, opacis. — Long.  $4\frac{1}{2}$  lin.

Nordm. Symb. 81. 28. — Er. Col. March. I. 452. 9.

*Staph. lucens* Mannerh. Brachelytr. 28. 48.

Habitat in Germania.

*Ph. polito* magnitudine aequalis, paulo tamen latior, niger, parum nitidus, capite thoraceque politis, interdum virescenti-subaeneis. Antennae capite thoraceque paulo breviores, articulo tertio secundo vix longiore, 4—10 sensim paulo brevioribus et crassioribus, prioribus subcylindricis, penultimis transversim subobconicis, ultimo ovato, apice inferius abrupte acuminato, nigrae, concolores, Palpi nigri. Caput thorace angustius et multo minus, supra oculos punctis nonnullis subtilioribus, inter oculos duobus transversim positis utrinque impressum. Thorax latitudine paulo brevior, basi coleopteris latitudine aequalis, antrorsum leviter angustatus, basi et lateribus rotundatus, apice truncatus, angulis anterioribus rotundatis, posterioribus obtusis, subrotundatis, modice convexus, seriebus dorsalibus punctis 4 aequaliter distantibus, antico paulo magis extrorsum sito, compositis, lateribus punctis praeter marginalia utrinque 2 versus medium et 5 intra angulum anteriorem impressus, punctis omnibus subtilioribus. Elytra thorace paulo longiora et scutellum confertissime subtiliter punctata, perparum nitida, subtiliter tenuiterque nigro-pubescentia. Abdomen subtilius minus dense punctatum, parce breviterque nigro-pubescent, metallico-versicolor. Pedes nigro-picei, coxis intermediis leviter distantibus, tibiis omnibus spinulosus, tarsis anticis in utroque sexu simplicibus.

Obs. 1. A *Ph. polito* statura paulo brevior, colore obscuriore et antennarum articulo primo concolore facile distinguendus.

2. Descriptio Mannerheimiana nimis manca, nonnisi ex auctoritate Prof. Nordmanni citata, qui collectionem celeberrimi huius auctoris lustravit.

\* 26. *Ph. politus*: Niger, supra subvirescent, capite ovali, thorace angustiore thoraceque nitidis, antennarum articulo primo subtus testaceo. — Long. 5 lin.

Nordm. Symb. 81. 27. — Er. Col. March. I. 452. 10.

*Staph. politus* Fab. Syst. Ent. 266. 7. Spec. Ins. I. 335. 7. Mant. Ins. I. 220. 11. Ent. Syst. I. II. 524. 20. Syst. El. II. 594. 22. — Panz. Ent. Germ. 354. 14. — Oliv. Ent. III. 42. 25. 31. t. 2. f. 10. — Grav. Micr. 17. 19. Mon. 85. 76. — Latr. Hist. nat. Crust. et Ins. IX. 308. 26. Gen. Crust. et Ins. I. 287. 3. — Gyll. Ins. Suec. II. 317. 33. — Zetterst. Faun. Lappon. I. 74. 18. Ins. Lappon. 63. 24. — Sahlb. Ins. Fenn. I. 320. 31. — Mannerh. Brachelytr. 28. 46. — Boisd. et Lacord. Faun. Ent. Paris. I. 392. 10. — Runde Brachelytr. Hal. 7. 29.

Habitat in Europa.

Niger, capite thoraceque supra obscure viridi-subaeneis, politis, elytris obscure viridi-aeneis, subnitidis, palpis pedibusque nigris. Antennae capite thoracisque longitudine, graciles, articulo tertio secundo paululum longiore, 4—10 sensim paulo brevioribus at vix crassioribus, omnibus crassitie longioribus, ultimo ovato, apice oblique emarginato, nigrae, articulo primo testaceo, supra nigro. Caput thorace minus et angustius, subovatum, supra oculos utrinque parce punctatum, inter oculos punctis utrinque duobus transversim positis notatum. Thorax latitudine haud brevior, basi coleopterorum fere latitudine, antrorsum modice angustatus, basi fortius, lateribus leviter rotundatus, apice subemarginatus, angulis anterioribus leviter deflexis, subrotundatis, posterioribus obtusis, modice convexus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum nigrum, confertim punctulatum, subtiliter fusco-pubescent. Elytra thorace vix longiora, dense subtiliterque punctata, subtiliter fusco-pubescentia. Abdomen subnitidum, subtilius punctatum, tenuiter nigro-pubescent. Pedes graciliores, coxis intermediis approximatis, tibiis omnibus parcius spinulosis.

Tarsi antici maris modice dilatati, feminae simplices.

Obs. 1. *Staph. politus* Lin., secundum collectionem Linneanam ad *Ph. aeneum* pertinet, at descriptio Linneana hanc speciem non magis quam multas alias significat, immo plures ab eo confusas esse patet. Paykul itidem plures species confusas pro *St. polito* descripsit. Loci citati igitur sequentes a *Ph. polito* nostro excludendi:

*Staph. politus* Linn. Faun. Suec. n. 843. Syst. Nat. I. II. 683. 5. — Ström. Act. Nidros. III. n. 27. — Müll. Faun. Frid. 23. 224. Prodr. 98. 1108. — Goeze Ent. Beitr. I. 717. 5. — De Vill. Ent. I. 412. 5. — Scop. Ent. Carn. 100. 304. — Schrank Enum. Ins. Austr. 231. 437. Faun. Boica. I. 641. 865. — Walk. Faun. Paris. I. 271. 11. — Cederh. Faun. Ingr. 118. 360. — Payk. Mon. Staph. 42. 31. Faun. Suec. III. 391. 31. — Marsh. Ent. Brit. 504. 18.

*Staph. aeneus* Degeer Ins. IV. 23. 9. — Fourcr. Ent. Par. I. 168. 17.

*Le Staph. à etuis bronzés* Geoffr. Hist. Ins. I. 367. 17.

Citatum Fabricianum, specimine in auctoris collectione ipso examinato, indubium.

2. *Staph. fuscipennem* Mannerh. Brachelytr. 28. 47. nonnisi elytris fusco-castaneis et pedibus piceis a *Ph. polito* diversum, revera distinctum esse, mihi non satis persuasum est, cum varietates aliarum specierum (*Ph. varii*, albidipedis et ebenini) analogae nobis sint.

\* 27. *Ph. marginatus*: Niger, nitidus, thoracis lateribus pedibusque flavis, capite ovato. — Long. 4 lin.

Curt. Britt. Ent. XIII. t. 610. — Nordm. Symb. 84. 38. — Er. Col. March. I. 453. 11.

*Staph. marginatus* Fab. Syst. Ent. 266. 8. Spec. Ins. I. 336. 9. Mant. Ins. I. 221. 15. Ent. Syst. I. II. 526. 30. Syst. El. II. 597. 38. — Ström. Act. Nidros. IV. n. 13. — Müll. Faun. Fridr. 23. 230. Prodr. 97. 1100. — Goeze Ent. Beitr. I. 726. 21. — Payk. Mon. Staph. 44. 32. Faun. Suec. III. 392. 31. — Oliv. Ent. III. 42. 26. 33. t. 3. f. 29. — Marsh. Ent. Brit. 512. 40. — Grav. Micr. 24. 32.



*Mon.* 71. 56. — *Latr. Hist. nat. Crust. et Ins.* IX. 313. 37. — *Gyll. Ins. Suec.* II. 322. 38. — *Zetterst. Faun. Lappon.* I. 75. 22. *Ins. Lappon.* 64. 31. — *Sahlb. Ins. Fenn.* I. 322. 36. — *Mannerh. Brachelytr.* 28. 54. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 397. 22. — *Runde Brachelytr. Hal.* 7. 34.

Habitat in Europa.

St. polito paulo minor et gracilior. Antennae capite thoraceque parum breviores, graciliores, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus at haud crassioribus, omnibus crassitie longioribus, ultimo apice emarginato, nigrae, articulo primo flavo, supra nigro. Palpi utrique nigri, articulis duobus primis flavis. Caput ovatum, thorace angustius et minus, supra oculos utrinque parce punctatum, inter oculos utrinque punctis duobus oblique positis impressum, nigrum, nitidum. Thorax latitudine haud brevior, basi coleopteris paulo angustior, antrorsum leviter angustatus, basi modice, lateribus leviter rotundatus, apice truncatus, angulis anterioribus rotundatis, posterioribus obtusis, leviter convexus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis omnibus subtilioribus, niger, nitidus, margine laterali late flavo. Scutellum nigrum, confertim punctulatum, griseo-pubescent. Elytra thorace paulo longiora, densius subtiliter punctata, griseo-pubescentia, virescenti-nigra, subnitida. Abdomen parce subtiliterque punctulatum, nigrum, supra metallico-versicolor, subtus segmentis singulis apice ferrugineo-marginatis. Pectus nigrum. Pedes flavi, coxis anterioribus concoloribus, intermediis approximatis, tibiis anticis breviter, posterioribus longius spinulosus, tarsis nigricantibus.

Tarsi antici maris fortius, feminae leviter dilatati.

\* 28. *Ph. umbratilis*: Niger, nitidus, capite orbiculato, elytris nigro-aeneis, subtiliter punctatis, pedibus obscure testaceis. — Long. 3 lin.

*Er. Col. March.* I. 454. 12.

*Staph. umbratilis* Grav. *Micr.* 170. 21. *Mon.* 69. 46.

*Staph. subfuscus* Gyll. *Ins. Suec.* II. 326. 42. — *Sahlb. Ins. Fenn.* I. 323. 39. — *Mannerh. Brachelytr.* 29. 57. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 395. 18. — *Runde Brachelytr. Hal.* 8. 36.

*Philonth. subfuscus* Nordm. *Symb.* 84. 41.

*Philonth. umbratilis* Nordm. *Symb.* 84. 47.

Habitat in Europa et in America septentrional.

Habitu fere *Ph. atrati*, *Ph. vario* aequalis, niger, nitidus, elytris nigro-subaeneis. Antennae capite thoraceque vix longiores, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paululum crassioribus at haud brevioribus, crassitie longioribus, ultimo oblongo, inferne abrupte acuminato, piceo-nigrae. Palpi picei. Caput maris thoracis latitudine, feminae eodem paulo angustius, orbiculatum, supra oculos utrinque parce fortiter punctatum, inter oculos punctis utrinque duobus impressum. Thorax coleopteris paulo angustior, latitudine vix brevior, antrorsum haud angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, parum convexus, seriebus dorsalibus punctis 4 profundioribus, anterioribus duobus paulo magis approximatis, compositis, lateribus punctis praeter marginalia utrinque 4 impressus, sericeo-versicolor. Scutellum confertissime subtiliter punctulatum, griseo-pubescent.



Elytra thorace vix longiora, dense subtiliter punctata, minus subtiliter griseo-pubescentia. Abdomen supra subtilius punctatum, densius griseo-pubescentis, subtus segmentis ultimis apice ferrugineo-marginatis. Pedes fusco-testacei, coxis intermediis leviter distantibus, tibiis omnibus spinulosis.

Tarsi antici maris fortiter dilatati, feminae simplices.

Obs. Specimen Americanum a Gravenhorstio e collectione Knochiana descriptum, nunc in Museo regio asservatum, ab Europaeis non differt nisi antennis basipiceis, palpis pedibusque dilutioribus, elytris margine summo suturali et apicali abdominisque segmentis omnibus margine apicali piceo, specie nullo modo ab his distinguendum.

\* 29. *Ph. piceatus*: Niger, nitidus, elytris nigro-subaeneis, pedibus fusco-testaceis, abdomine infra segmentis rufo-marginatis, capite orbiculato. — Long.  $3\frac{1}{2}$  lin.

Nordm. Symb. 85. 45.

Habitat in Mexico, Dom. Deppe.

Affinis praecedenti, at duplo maior, niger, nitidus, capite thoraceque politis, elytris nigro-subaeneis. Antennae nigrae. Palpi fusco testacei. Caput thorace vix angustius, maiusculum, orbiculatum, utrinque supra oculos punctis paucis, inter oculos punctis utrinque duobus transversim positiss notatum. Thorax coleopteris latitudine fere aequalis, longitudine nonnihil lator, antrorsum subangustatus, lateribus subrectis, basi rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, transversim convexus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis omnibus fortioribus. Scutellum nigrum, dense punctulatum, griseo-pubescentis. Elytra thorace paulo longiora, crebre punctata, densius longiusque emereo-pubescentia. Abdomen laevigatum, vel punctis rarissimis, praecipue segmentorum margines versus, impressum, nigrum, pernitidum, metallico-versicolor, segmentis ventralibus rufo-marginatis. Pedes fusco-testacei, coxis intermediis approximatis, tibiis anticis breviter, posterioribus longius spinulosis.

Tarsi antici maris fortiter dilatati; femina latet.

Obs. 1. Differt a praecedente statura duplo maiore, elytris densius punctatis, abdomine prope laevigato.

2. Antennae in nostro specimine unico mutilatae: articuli 5 primi nigri, tertius secundo paulo longior.

\* 30. *Ph. cautus*: Niger, nitidus, antennarum basi, elytrorum margine summo, ano pedibusque testaceis, capite orbiculato. — Long. 3 lin.

Habitat in Carolina meridionali, Dom. Zimmermann, a Dom. Prof. Germar benevole communicatus.

Statura Ph. umbratilis, habitu Ph. discoideo propior, niger, capite thoraceque pernitidis. Antennae capite paulo longiores, apicem versus haud crassiores, articulo tertio secundo paulo longiore, sequentibus cylindricis, 4—10 sensim paulo brevioribus, penultimis transversis, ultimo apice leviter emarginato et inferne subacuminato, fuscae, articulo primo testaceo, secundo piceo-testaceo. Palpi picei. Mandibulae rufo-piceae. Caput sat magnum, thoracis latitudine, suborbiculatum, perparum convexus, supra oculos utrin-

que punctis paucis, inter oculos punctis 4 transversim positis, intermediis magis distantibus, omnibus subtilibus, impressum. Thorax basi coleopteris vix angustior, latitudine baseos vix longior, antrorsum modice angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, rotundatis, satis convexus, seriebus dorsalibus punctis 4 sat distantibus, apicali subtili et a reliquis magis remoto, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum crebrius subtiliusque punctulatum, griseo-pubescentia. Elytra crebrius et minus subtiliter punctata, densius griseo-pubescentia, margine laterali inflexo summoque apicali determinato testaceis. Abdomen parcius subtiliusque punctulatum, longius griseo-pubescentia, segmento ultimo toto testaceo, ventralibus margine apicali late testaceis. Pedes breviusculi, testacei, coxis concoloribus, intermediis approximatis, tibiis omnibus spinulosis.

Tarsi antici maris fortius dilatati, femina latet.

\* 31. *Ph. varius*: *Niger, nitidus, elytris viridi-aeneis, capite thorace angustiore, subovato.* — Long. 3 —  $3\frac{1}{2}$  lin.

Nordm. Symb. 84. 37. — Er. Col. March. F. 455. 14.

*Staph. varius* Gyll. Ins. Suec. II. 321. 37. — Zetterst. Faun. Lappon. I. 74. 19.

Ins. Lappon. 63. 28. — Sahlb. Ins. Fenn. I. 321. 35. — Mannerh. Brachelytr. 28. 53. — Boisd. et Lacord. Faun. Ent. Paris I. 395. 17. — Runde Brachel. Hal. 7. 32.

*Staph. varians* Grav. Micr. 20. 21. — Latr. Hist. nat. Crust. et Ins. IX. 309. 28.

*Var. a, Pedibus piceo-testaceis.*

*Staph. carbonarius* Grav. Micr. 23. 31. Mon. 67. 42. — Latr. Hist. nat. Crust. et Ins. IX. 313. 36.

*Var. b, Elytris apice rutilis.*

*Var. c, Elytris pedibusque rufis.*

Habitat in Europa.

Habitu fere *Ph. politus*, at multo minor, niger, nitidus, capite thoraceque politis, elytris viridi-aeneis, nitidis. Antennae capitis thoracisque longitudine, articulo tertio secundo paululum longiore, sequentibus subcylindricis, dein sensim subobconicis, omnibus crassitie longioribus, ultimo oblongo-ovato, apice truncato et inferne acuminato, nigrae. Palpi nigri. Caput thorace multo minus et angustius, subovatum, supra oculos utrinque punctis parvis, inter oculos utrinque punctis duobus impressum. Thorax basi fere coleopterorum latitudine, latitudine haud brevior, antrorsum leviter angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice subemarginatus, angulis anterioribus deflexis, rotundatis, transversim convexus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis subtilioribus. Scutellum nigrum, confertim punctulatum. Elytra thoracis longitudine, minus dense subtilius punctata, griseo-pubescentia. Abdomen subtiliter parceque punctatum, griseo-pubescentia. Pedes plerumque nigri, interdum fusci vel fusco-testacei, coxis intermediis leviter distantibus, tibiis omnibus spinulosis.

Tarsi antici maris fortiter, feminae leviter dilatati.

Variat elytris minus magisque rufescentibus, vel nonnunquam totis rufis.

Obs. 1. *Staph. lucidus* Grav. Micr. 21. 24, a Gravenhorstio Mon. 82. *Staph. finetario subiunctus*, at vix rite: speciminum enim duorum a cel. auctore descriptorum alterum fuit *Ph. varius*, alterum *Ph. quisquiliarius*.

2. *Staph. nitens* Grav. Micr. 26. 36. Mon. 78. 71. (Latr. Hist. nat. Crust. et Ins. IX. 315. 41.) forte ad hanc speciem, forte etiam ad *Ph. albigredem* pertinet, mihi omnino dubius.

\* 32. *Ph. gilvipes*: *Niger, nitidus, elytris viridi-aeneis, antennis pedibusque testaceis, capite minuto, oblongo-ovato.* — Long. 3 lin.

*Er. Col. March. I.* 456. 15.

Habitat in Germania, Dom. Schüppel.

*Ph. vario* statura affinis, niger, capite thoraceque nitidissimis, elytris virescenti-aeneis, subnitidis. Antennae capite thoraceque parum breviores, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo brevioribus et crassioribus, penultimis crassitie vix brevioribus, ultimo oblongo-ovato, apice truncato et inferne acuminato, fusco-testaceae, articulis duobus primis testaceis. Palpi testacei. Caput parvulum, thorace multo angustius, oblongo-ovatum, supra oculos utrinque punctis 4 vel 5, inter oculos utrinque punctis 2 transversim positis impressum. Thorax basi coleopteris latitudine aequalis, longitudine paululum lator, antrorsum angustatus, lateribus leviter, basi cum angulis posterioribus fortius rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, convexus, seriebus dorsalibus punctis 4 remotis, aequaliter distantibus, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis subtilibus, angulis anterioribus summo margine rufis. Scutellum nigrum, crebre punctulatum. Elytra thorace paulo longiora, subtiliter dense punctata, densius griseo-pubescentia. Abdomen crebrius punctulatum, griseo-pubescentis, segmentis ventralibus rufo-marginatis. Pedes testacei, coxis anticis nigricantibus, intermediiis approximatis, tibiis omnibus spinulosus.

Tarsi antici maris modice dilatati, feminae simplices.

Obs. A *Ph. vario* antennis pedibusque testaceis, elytris magis virescentibus abdomineque densius subtilisque punctatis, capite angustiore, a *Ph. albipede* statura duplo maiore et elytris viridibus distinctus.

\* 33. *Ph. bimaculatus*: *Niger, nitidus, capite subovato, elytris aeneis, macula apicali rufa.* — Long. 3 lin.

*Nordm. Symb.* 92. 59. — *Er. Col. March. I.* 454. 13.

*Staph. bimaculatus* Grav. Micr. 38. 55. Mon. 63. 32. — Latr. Hist. nat. Crust. et Ins. IX. 321. 60. — Boisd. et Lacord. Faun. Ent. Paris. I. 400. 26. — Runde Brachelytr. Hal. 9. 44.

*Staph. nitidicollis* Boisd. et Lacord. Faun. Ent. Paris. I. 408. 28.

*Staph. dubius* Grav. Mon. 74. 63. (forte)

Habitat in Borussia, Germania, Gallia, Italia.

Statura et magnitudo ferè *Ph. varii*, niger, nitidus, capite thoraceque politis, elytris viridi-aeneis, nitidis, macula apicali lateritia minore maioreve, nonnunquam per totum elytri discum effusa. Antennae capite thorace paulo breviores, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo brevioribus, penultimis crassitie haud brevioribus, ultimo ovato, apice inferne abrupte acuminato, nigrae. Palpi nigri. Caput thorace



minus et angustius, subovatum, supra oculos utrinque parce fortius punctatum, inter oculos punctis 4 transversim positis impressum. Thorax coleopteris vix angustior, latitudine fere brevior, antrorsum subangustatus, basi modice, lateribus leviter rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusissimis, modice convexus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis fortioribus. Scutellum nigrum, confertim punctulatum. Elytra thoracis longitudine, crebrius fortiusque punctata, griseo-pubescentia. Abdomen parcius subtiliter punctatum, griseo-pubescentia. Pedes nigri, tibiis tarsisque saepe rufo-piceis, coxis intermediis approximatis, tibiis anticis breviter, posterioribus longius spinulosis.

Tarsi antici maris fortiter, feminae leviter dilatati.

Variat thoracis seriebus dorsalibus 5-punctatis, minus vero regularibus.

Obs. Staph. nitidicollis Boisd. et Lacord, nonnisi macula per maximam elytrorum partem effusa a genuino differt. Staph. dubius Grav. ad eadem individua pertinere videtur.

\* 34. *Ph. albipes*: *Niger, nitidus, abdominis segmentorum ventralium marginibus pedibusque testaceis, elytris nigro-piceis, crebre punctatis, capite ovato.* — Long.  $2\frac{1}{2}$  lin.

*Er. Col. March. I.* 457. 16.

*Staph. albipes* Grav. *Micr.* 28. 40. *Mon.* 76. 68. — *Latr. Hist. nat. Crust. et Ins. IX.* 316. 45. — *Gyll. Ins. Suec. II.* 327. 43. — *Zetterst. Faun. Lappon. I.* 75. 23. *Ins. Lapp.* 64. 33. — *Sahlb. Ins. Fenn. I.* 323. 14. — *Mannerh. Brachelytr.* 29. 58. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 405. 37.

Habitat in Germania, Suecia, Gallia.

Habitu Ph. varii, at plus duplo minor, niger, capite thoraceque nitidis. Antennae capite thoraceque breviores, articulo tertio secundo vix longiore, 4 — 10 subobconicis, sensim paululum brevioribus et crassioribus, penultimis crassitie haud brevioribus, ultimo oblongo-ovato, apice emarginato et inferne subacuminato, nigrae, articulis duobus primis piceis. Palpi nigri. Caput thorace angustius minusque, ovatum, supra oculos utrinque punctis nonnullis, inter oculos utrinque punctis duobus transversim positis notatum. Thorax basi coleopterorum latitudine, latitudinis fere longitudine, antrorsum perparum angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, modice convexus, seriebus dorsalibus punctis 4 sat distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus, sericeo-versicolor. Scutellum nigrum, confertim punctulatum. Elytra nigro-picea, nitidula, subtilius dense punctata. Abdomen parcius subtiliter punctatum, cum elytris pube densiore, longiore et minus subtili, depressa, grisea vestitum, segmentis inferioribus apice ferrugineo-marginatis. Pedes testacei, coxis intermediis leviter distantibus, tibiis omnibus spinulosis.

Tarsi antici maris leviter dilatati, feminae simplices.

Variat I, Antennis basi piceo-testaceis. Specimina Suecica dedit Dom. Schüppel.

2, Antennis basi pedibusque flavo-testaceis, elytris obscure rufis. Specimen singulum Dom. Schüppel Berolini legit.

3, Thoracis serie dorsali altera 4-, altera 5-punctata.

- \* 35. *Ph. lepidus*: *Niger, nitidus, antennarum basi, elytris pedibusque rufis, capite ovato.* — Long. 3 lin.

*Er. Col. March. I.* 457. 17.

*Staph. lepidus* Grav. *Micr.* 31. 45. *Mon.* 74. 64. — *Latr. Hist. nat. Crust. et Ins. IX.* 318. 50. — *Gyll. Ins. Suec. II.* 328. 44. — *Sahlb. Ins. Fenn. I.* 323. 41. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 401. 29.

*Staph. flavopterus* Grav. *Mon.* 74. 65.

*Philonth. flavopterus* Nordm. *Symb.* 89. 51.

*Staph. fulvipes* Runde *Brachelytr. Hal.* 10. 52.

Habitat in Germania, Suecia, Gallia.

Apterus, niger, capite thoraceque pernitidis. Antennae capitis thoracis-que prope longitudine, articulo tertio secundo paulo longiore, 4—10 subobconicis, sensim paululum brevioribus at non crassioribus, omnibus crassitie longioribus, ultimo oblongo, apice leviter emarginato et inferne acuminato, nigrae vel nigro-piceae, articulo primo, interdum etiam secundo, rufo-testaceo. Palpi nigro-picei. Caput thorace minus angustiusque, ovatum, supra oculos utrinque punctis nonnullis, inter oculos utrinque punctis duobus transversim positissimum impressum. Thorax coleopteris latitudine aequalis, latitudine vix brevior, antrorsum subangustatus, lateribus parum, basi cum angulis posterioribus fortius rotundatus, apice truncatus, angulis posterioribus deflexis, subrotundatis, convexus, seriebus dorsalibus punctis 4 remotis compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis omnibus subtilioribus. Scutellum nigrum, confertim subtiliter punctatum. Elytra thorace fere breviora, confertim subtiliter punctata, lateritia, griseo-pubescentia. Abdomen subtiliter punctatum, densius subtiliter griseo-pubescentes, parce nigro-pilosum, segmentis inferioribus apice tenuiter ferrugineo-marginatis. Pedes cum coxis anterioribus rufo-testaceis, coxis intermediis approximatis, tibiis anticis breviter, posterioribus longius spinulosis.

Tarsi antiqui maris fortius dilatati, feminae simplices.

- \* 36. *Ph. nitidulus*: *Niger, nitidus, pedibus testaceis, elytris brevibus, profunde punctatis, piceis, capite ovato.* — Long. 2 lin.

*Nordm. Symb.* 85. 43. — *Er. Col. March. I.* 458. 18.

*Staph. nitidulus* Grav. *Micr.* 27. 37. *Mon.* 69. 51. — *Latr. Hist. nat. Crust. et Ins. IX.* 315. 42. — *Gyll. Ins. Suec. II.* 330. 46. — *Sahlb. Ins. Fenn. I.* 323. 42. — *Mannerh. Brachelytr.* 29. 60. — *Runde Brachelytr. Hal.* 8. 37.

*Staph. denigrator* Grav. *Mon.* 92. 88.

*Philonth. denigrator* Nordm. *Symb.* 85. 44.

Habitat in Germania, Suecia.

Apterus, niger, capite thoraceque pernitidis. Antennae capite thorace-que breviores, crassiusculae, articulo tertio secundo haud longiore, 5—10 subaequalibus, crassitie paulo brevioribus, ultimo ovato, apice truncato et abrupte subacuminato, dilutius saturatiusve rufo-piceae. Palpi picei, articulo ultimo apice fortius acuminato. Caput thorace angustius, ovatum, supra oculos utrinque punctis paucis, inter oculos punctis 4 transversim positissimum aequaliter distantibus, impressum. Thorax coleopteris latitudine aequalis, latitudine prope longior, antrorsum leviter angustatus, lateribus levissime, basi cum angulis posterioribus fortius rotundatus, apice truncatus, angulis anterioribus deflexis, rotundatis, fortius transversim convexus, seriebus dor-

salibus punctis 4 sat distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilibus. Scutellum obsolete punctulatum. Elytra thorace breviora, minus dense fortiter punctata, parce griseo-pubescentia, piceo-subaenea, nitida. Abdomen parcius subtiliusque punctulatum, parce griseo-pubescentia, segmento ultimo subtus apice rufo-piceo. Pedes cum coxis anterioribus testacei, coxis intermediis tenui spatio distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

- \* 37. *Ph. hepaticus*: *Niger, thoracé abdominisque segmentorum marginibus fusco-piceis, elytris fuscis, apice pallidis, antennarum basi pedibusque flavo-testaceis, capite ovato. — Long. vix 2 lin.*

Habitat in America septentrionali, e coll. Knochiana; a Dom. Zimmermann quoque missus.

Statura fere *Ph. albipedis*, at paulo minor. Antennae capite cum thorace paulo breviores, articulo tertio secundo paulo longiore, sequentibus subobconicis, sensim paululum brevioribus, non autem crassioribus, penultimis crassitie haud brevioribus, ultimo oblongo-ovato, apice truncato et inferne acuminato, fuscae, articulis 2 primis testaceis. Palpi picei. Caput thorace angustius, subovatum, supra oculos utrinque punctis nonnullis, inter oculos punctis 4 transversim positis, intermediis remotioribus, impressum, nigro-piceum, nitidum. Thorax basi coleopteris paulo angustior, latitudine haud brevior, antrorsum leviter angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, transversim convexus, seriebus dorsalibus punctis 4 aequaliter distantibus et sat remotis compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis minus subtilibus, rufo-piceis, nitidus. Elytra thorace paulo longiora et scutellum fusca vel fusco-testacea, illa dorso infusca, margine apicali pallido, subtilius crebre punctata, griseo-pubescentia. Abdomen subtiliter punctulatum, griseo-pubescentia, nigro-fuscum, segmentorum marginibus rufo-piceis vel fusco-testaceis, infra apice testaceum. Pedes flavo-testacei, coxis intermediis leviter distantibus, tibiis anticis subtiliter, posterioribus fortius spinulosis.

Tarsi antici maris fortiter, feminae leviter dilatati.

- \* 38. *Ph. vilis*: *Nigro-piceus, nitidus, antennarum basi, pedibus abdominisque segmentorum ventralium marginibus testaceis, thorace piceo, capite subovato. — Long. 2 lin.*

Habitat in Americae meridionalis ins. St. Thomae, Dom. Moritz.

Praecedenti affinis, coloribus *Ph. ventrali* similis, at multo minor, praesertim angustior, statura *Ph. aterrimi*, nigro-piceus, nitidus, capite thoraceque nitidissimis. Antennae capite thoraceque breviores, crassiusculae, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus, non autem crassioribus, penultimis crassitie vix brevioribus, ultimo subovato, apice truncato et inferne abrupte acuminato, nigrae, articulis duobus basalibus testaceis. Os cum palpis obscure testaceum. Caput thorace paulo angustius, breviter subovatum, utrinque supra oculos punctis paucis subtilioribus, inter oculos punctis 4 transversim positis, intermediis remotioribus, impres-



sum. Thorax coleopteris paulo angustior, latitudine haud longior, antrorsum leviter angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrotundatis, leviter convexus, seriebus dorsalibus punctis 4 aequaliter distantibus et sat remotis compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus minus fortibus, piceus. Scutellum nigrum, crebre punctatum, griseo-pubescentia. Elytra thorace paulo longiora, nigro-picea, nitida, parcius minus profunde punctata, longius griseo-pubescentia. Abdomen parce subtilissime punctulatum, griseo-pubescentia, segmentorum ventralium marginibus rufo-testaceis. Pedes cum coxis anterioribus flavo-testacei, coxis intermediis approximatis, tibiis omnibus parcius spinulosis, tarsis anticis in utroque sexu fortiter dilatatis.

\* 39. *Ph. orphanus*: *Niger, nitidus, antennarum basi abdominisque segmentorum ventralium marginibus testaceis, pedibus flavis, thorace oblongo, capite subovato.* — Long. 2 lin.

Habitat ad Caracas Columbiae, Dom. Moritz.

Statura omnino *Ph. aterrimi* et maximis eius individuis parum maior, niger, nitidus. Antennae capite thoraceque breviores, articulo tertio secundo parum longiore, 4 — 10 sensim brevioribus at vix crassioribus, penultimis subtransversis, ultimo oblongo-ovato, apice oblique truncato et inferne subacuminato, nigrae, articulis duobus primis piceo-testaceis. Mandibulae rufo-piceae. Palpi nigro-picei. Caput thoracis antico latitudine prope aequale, subovatum, modice convexum, supra oculos utrinque punctis paucis, inter oculos punctis 4 transversim positiss, intermediis paulo remotioribus, impressum. Thorax latitudine paulo longior, coleopteris dimidio prope angustior, apicem versus subangustatus, lateribus subrectis, basi rotundatus, apice truncatus, angulis omnibus fortius rotundatis, leviter convexus, seriebus dorsalibus punctis 4 sat remotis et aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis omnibus subtilioribus. Scutellum subtilissime punctulatum, tenuiter pubescens. Elytra thorace paulo longiora, minus confertim subtilius punctata, tenuiter griseo-pubescentia. Abdomen parcius subtiliter punctatum, tenuiter griseo-pubescentia, subius segmentis singulis margine postico rufo-testaceis. Pedes flavo-testacei, coxis anterioribus concoloribus, intermediis tenui spatio distantibus, tibiis anticis subtilissime, posterioribus distincte spinulosis.

Tarsi antici maris fortius, feminae leviter dilatati.

Variat minus maturus nigro-piceus, elytris fusco-testaceis.

✕ 40. *Ph. xantholoma*: *Niger, elytris opacis, margine inflexo luteo, abdomine cinereo-lineato, pedibus fusco-testaceis, thorace basin versus angustato.* — Long. 3½ lin.

*Staph. xantholoma* Grav. Mon. 41. 3. — Gyll. Ins. Suec. II. 323. 39. — Germ. Faun. Ins. Europ. XIII. 13. — Zetterst. Ins. Lappon. 64. 32.

*Cafius xantholoma* Curtis Britt. Ent. VII. fol. 322. — Mannerh. Brachelytr. 31. 1. *Gyrohypn. xantholoma* Nordm. Symb. 126. 28.

Habitat in Germaniae septentrionalis, Sueciae, Angliae, Lusitaniae Italiaeque littoribus.

Statura fere *Ph. cephalotis*. Antennae capite cum thorace breviores, articulo tertio secundo parum longiore, 4 — 10 sensim paulo brevioribus, crassitie autem aequalibus, ultimo subovato, apice abrupte acuminato, nigropiceae. Palpi picei, apice rufescentes. Mandibulae rufo-piceae. Caput mainseculum, thorace plerumpue latius, subquadratum, utrinque supra oculos punctis magnis at parum profundis parvis impressum, fronte inter oculos utrinque punctis duobus transversim positis notata, medio leviter foveolata, nigrum, nitidum. Thorax coleopteris duplo fere angustior, latitudine parum longior, basin versus subangustatus, lateribus leviter rotundatis, pone medium subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, posterioribus obtusis, parum convexus, punctis serierum dorsalium posterioribus tribus grossis, antico remotiore, minuto, lateribus intra angulum anteriorem punctis nonnullis impressus, nigro-piceus, nitidus. Elytra thorace sesqui fere longiora et scutellum subtilissime confertissimeque punctata, nigra, opaca, densius griseo-pubescentia, illa margine laterali inflexo luteo. Abdomen subtilissime confertissimeque punctatum, opacum, nigrum, cinereo-pubescentia, dorso lineis duabus macularibus e pube cinerea ornatum. Pectus nigrum, opacum. Pedes fusco-testacei, coxis intermediis tenui spatio distantibus, anticis femoribusque obscurioribus, piceis, tibiis omnibus spinulosis, tarsis anticis in utroque sexu leviter dilatatis.

Abdominis segmentum inferius quintum in mare medio longitudinaliter impressum et leviter emarginatum.

\* 41. *Ph. variegatus*: *Niger, thoracis limbo sanguineo, elytris opacis, margine inflexo luteo, abdomine ferrugineo, supra nigro-cinereoque variegato, pedibus testaceis, thorace basin versus angustato.* — Long.  $3\frac{1}{2}$  lin.

Habitat in Sicilia, Dom. Grohmann.

Statura et summa affinitas praecedentis. Antennae capite thoraceque breviores, articulo secundo tertioque subaequalibus, 4 — 10 subaequalibus, crassitie vix longioribus, ultimo ovato, apice abrupte acuminato, nigrae. Palpi picei, apice dilute rufescentes. Caput thoracis latitudine, subquadratum, supra oculos utrinque obsolete punctatum, inter oculos punctis utrinque duobus transversim positis notatum et medio longitudinaliter obsolete impressum, nigrum, nitidum. Thorax latitudine paulo longior, basin versus leviter angustatus, basi coleopteris duplo prope angustior, lateribus ante medium leviter rotundatis, pone medium leviter sinuatis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, posterioribus obtusis, leviter convexus, serierum dorsalium punctis tribus posterioribus maioribus, postico et antico subtiliore remotioribus, lateribus intra angulum anteriorem punctis paucis impressus, rufo-sanguineus, disco lato et macula minuta in utroque angulo anteriore summo nigris, nitidus. Scutellum confertissime subtiliter punctatum, griseo-pubescentia, rufum, opacum. Elytra thorace tertia parte longiora, confertissime subtiliter punctata, dense subtiliterque cinereo-pubescentia, nigra, opaca, margine laterali inflexo summoque apicali luteis. Abdomen confertim subtiliter punctatum, subtus aequaliter subtiliter pubescens, supra maculis biseriatis e pube densiore cinerea et triseriatis nigris alternantibus variegatum seu tessellatum. Pectus nigrum, densius cinereo-pubescentia. Pedes testacei, coxis intermediis tenui spatio distantibus, femoribus posterioribus fuscis, tibiis omnibus spinulosis.

- \* 42. *Ph. fucicola*: *Brunneus, pectore capiteque nigricantibus, pedibus ferrugineis, abdomine biserialim subfoveolato, thorace basin versus angustato.* — Long. 5 lin.

*Nordm. Symb.* 84. 42.

*Cafius fucicola* *Curtis Britt. Ent. VII.* 322.

Habitat in Anglia, Dom. Leach.

Statura omnino *Ph. xantholomatis*, at maior et alius. Antennae capite cum thorace breviores, articulo tertio secundo paululum brevior, 4 — 10 sensim paululum brevioribus, penultimis longitudine haud crassioribus, ultimo ovato, apice subtruncato et inferne subacuminato, nigricantes, basi brunneae, articulo apicali ferrugineo. Caput oblongo-subquadratum, thorace paulo latius, basi lateribus grosse punctatum, fronte inter oculos medio leviter impressa, supra nigro-piceum, subnitidum, ore brunneo, infra ferrugineum. Thorax coleopteris multo angustior, latitudine paululum longior, basin versus angustatus, lateribus anterieus rotundatis, pone medium leviter sinuatis, basi parum, angulis posterioribus fortius rotundatis, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, leviter convexus, serierum dorsalium punctis tribus posterioribus grossis, quarto antico remoto, subtiliore, lateribus secundum anguli anterioris marginem punctis 4 maioribus impressus, omnium subtilissime alutaceus, supra brunneus, nitidus, infra ferrugineus. Elytra thorace paulo longiora et scutellum creberrime subtilius punctata, tenuiter pubescentia, fusca, opaca. Abdomen crebre punctulatum, segmentis singulis dorso obsolete bi-impressis, supra brunneum, subnitidum, infra ferrugineum. Pectus nigricans. Pedes rufo-ferruginei, coxis intermediis leviter distantibus, tibiis omnibus spinulosis, tarsis anticis (feminae) leviter dilatatis.

Variat interdum thoracis serie dorsali altera 5-punctata.

- \* 43. *Ph. cicatrisosus*: *Elongatus, piceus, antennis, pedibus abdomineque ferrugineis, thorace basin versus angustato, capite magno, quadrato.* — Long. 5 lin.

Habitat in Sicilia.

Statura elongata et depressa omnino praecedentis. Antennae capite thoraceque breviores, articulo tertio secundo sesqui prope longiore, 4 — 10 sensim brevioribus, primis eorum cylindricis, reliquis subobconicis, penultimo crassitie vix brevior, ultimo ovato, apice truncatulo et inferne subacuminato, ferrugineae, basi piceae. Palpi ferruginei. Caput thorace latius, magnum, depressum, oblongo-subquadratum, piceum, nitidum, margine laterali et verticali punctato, fronte utrinque longitudinaliter punctis grossis inaequalibus parum profundis et obsolescentibus variolosa, medio canalicula brevissima obsolete impressa. Thorax coleopteris angustior, latitudine apicis parum longior, basin versus fortius angustatus, lateribus ante medium leviter rotundatis, pone medium subsinuatis, basi parum, angulis posterioribus fortius rotundatis, apice truncatus, angulis anterioribus rotundatis, depressus, serierum dorsalium punctis posterioribus approximatis, grossis at parum profundis et obsoletioribus, irregularibus, uno alterove saepe extincto, antico remoto, pariter grossiusculo, lateribus utrinque ad angulum anteriorem et pone medium ad marginem lateralem punctis nonnullis irregularibus, parum profundis notatus, piceus, nitidus. Scutellum crebrius punctatum, sub-



tiliter pubescens. Elytra thorace paulo longiora, picea aut ferruginea, confertim fortius punctata, subtiliter pubescentia. Pectus fusco-piceum. Abdomen ferrugineum, subtiliter punctulatum, subtiliter griseo-pubescens, dorso obsolete variegatum. Pedes graciliores, ferruginei, coxis femoribusque piceis, coxis intermediis approximatis, tibiis omnibus spinulosis, tarsis anticis in utroque sexu modice dilatatis.

- \* 44. *Ph. albilabris*: *Niger, nitidus, pedibus flavis, antennis apice ferrugineis, thorace oblongo, capite magno, orbiculato.* — Long. 3 lin.

*Nordm. Symb.* 86. 46.

Habitat in Japonia, Dom. v. Langsdorf.

Statura fere *Ph. fimetarii*, niger, nitidus, capite thoraceque nitidissimis, elytris vix aenescentibus. Antennae capite cum thorace parum breviores, sat elongatae, articulo tertio secundo plus sesqui longiore, 4—10 elongato-obconicis, sensim paululum crassioribus, at vix brevioribus, ultimo oblongo, apice oblique truncato et inferne acuminato, nigrae, basi piceae, apice ferrugineae. Mandibulae longae et palpi picei. Caput thorace fere latius, orbiculatum, leviter convexum, utrinque supra oculos punctis nonnullis, inter oculos punctis utrinque duobus transversim positis, omnibus sat fortibus profundisque impressum. Thorax coleopteris angustior, latitudine paulo longior, lateribus rectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, modice convexus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis omnibus profundioribus. Scutellum confertim punctatum, griseo-pubescens. Elytra thorace tertia parte longiora, minus confertim subtilius punctata, densius griseo-pubescentia. Pedes pallide flavi, coxis omnibus nigris, intermediis leviter distantibus, tibiis anticis breviter, posterioribus longius spinulosis.

Tarsi antici maris modice dilatati. Femina latet.

- X \* 45. *Ph. cephalotes*: *Elongatus, niger, elytris aeneis, fortius punctatis, pedibus fusco-piceis, capite latiore, suborbiculato.* — Long. 3 — 4 lin.

*Nordm. Symb.* 82. 35. — *Er. Col. March.* I. 439. 19.

*Staph. cephalotes* Grav. *Micr.* 22. 27. *Mon.* 71. 55. — *Latr. Hist. nat. Crust. et Ins.* IX. 311. 32. — *Gyll. Ins. Suec.* II. 320. 36. — *Sahlb. Ins. Fenn.* I. 321. 34. — *Zetterst. Ins. Lappon.* 63. 27. — *Mannerh. Brachelytr.* 28. 52. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 90. 8.

Habitat in Europa; in Syria et Aegypto etiam a Prof. Ehrenberg lectus.

Elongatus, linearis, niger, capite thoraceque pernitidis, elytris aeneis, nitidulis. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, 4—10 subobconicis, sensim paulo brevioribus, penultimo crassitie fere breviori, ultimo ovato, apice subemarginato et inferne acuminato, nigro-piceae. Palpi picei. Caput feminae minoribusque individuis masculis thoracis fere latitudine, maribus maioribus thorace magis minusve latius, suborbiculatum, supra oculos utrinque punctis paucis, inter oculos punctis quatuor transversim positis notatum. Thorax coleopteris paulo angustior, latitudine vix longior, basin versus subangustatus, lateribus anterioribus leviter rotundatis, posterius subsinuatis, basi rotundatus, apice trun-

catus, angulis anterioribus leviter deflexis, rotundatis, posterioribus obtusis, basin versus subangustatus, parum convexus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus modice profundis. Scutellum confertim subtiliter punctulatum, griseo-pubescentia. Elytra thorace paulo longiora, densius profundiusque punctata, griseo-pubescentia. Abdomen subtiliter punctatum, densius longiusque griseo-pubescentia, segmentis ventralibus ferrugineo-marginatis. Pedes picei, coxis intermediis approximatis, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

Variat thoracis seriebus dorsalibus 5-punctatis.

Obs. Caput apud masculos secundum corporis magnitudinem magnitudine variat, ut in *Ph. aeneo*, *carbonario*, etc., ita ut quanto validius sit specimen, tanto grossiore capite, quo debilius, eo minore capite instructum sit; masculi igitur minores, qui frequentius occurrunt, capitis forma et proportionem parum a femina discedunt.

\* 46. *Ph. fimetarius*: *Elongatus, niger, nitidus, elytris viridi-aeneis, subtilius punctatis, pedibus flavis, capite suborbiculato*. — Long. 3 lin.

*Er. Col. March. I. 460. 21.*

*Staph. fimetarius* Grav. *Micr. 175. 32. Mon. 80. 72.* — *Latr. Hist. nat. Crust. et Ins. IX. 325. 72.* — *Gyll. Ins. Suec. II. 324. 40.* — *Zetterst. Faun. Lappon. I. 75. 21. Ins. Lappon. 63. 30.* — *Sahlb. Ins. Fenn. I. 322. 37.* — *Mannerh. Brachelytr. 29. 55.* — *Boisd. et Lacord. Faun. Ent. Paris. I. 406. 38.*

*Staph. rigidicornis* Grav. *Micr. 23. 29. Mon. 69. 49.* — *Latr. Hist. nat. Crust. et Ins. IX. 312. 34.*

*Philonth. rigidicornis* Nordm. *Symb. 82. 33.*

Habitat in Europa.

Habitu praecedentis, niger, capite thoraceque pernitidis, elytris lucide viridi-aeneis, nitidis. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, 4—10 sensim perparum brevioribus et crassioribus, crassiusculis, crassitie haud brevioribus, ultimo ovato, apice oblique emarginato et inferne acuminato, nigrae. Palpi picei. Caput in utroque sexu thoracis latitudine, suborbiculatum, supra oculos utrinque punctis paucis fortibus, inter oculos utrinque punctis duobus transversim positis et medio impressione lata at parum profunda notatum. Thorax coleopteris angustior, latitudine paulo longior, lateribus parallelis, subrectis, vel pone medium levissime sinuatis, basi rotundatus, angulis posterioribus obtusis, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, parum convexus, seriebus dorsalibus punctis 4, posterioribus tribus remotioribus, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus sat magnis profundisque. Scutellum punctulatum, griseo-pubescentia. Elytra thorace tertia parte longiora, densius subtiliusque punctata, longius griseo-pubescentia, segmentis singulis supra infraque ferrugineo-marginatis. Pedes laete flavi, coxis anticis picescentibus, intermediis tenui spatio distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

\* 47. *Ph. sordidus*: *Elongatus, niger, nitidus, elytris aeneis, parcius profunde punctatis, pedibus piceis, capite suborbiculato*. — Long. 3 lin.

*Nordm. Symb. 84. 40.* — *Er. Col. March. I. 459. 20.*

*Staph. sordidus* Grav. *Micr. 176. 33.* — *Gyll. Ins. Suec. II. 325. 41.* — *Zetterst.*

*Faun. Lappon. I. 75. 20. Ins. Lappon. 63. 21. — Sahlb. Ins. Fenn. I. 322. 38. — Mannerh. Brachelytr. 29. 55. — Boisd. et Lacord. Faun. Ent. Paris. I. 396. 19. Philonth. pachycephalus Nordm. Symb. 82. 35.*

Habitat in Europa.

Ph. cephaloti affinis, at paulo minor, niger, capite thoraceque pernitidis, elytris aeneis, nitidis. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus et crassioribus, subobconicis, penultimis crassitie vix brevioribus, ultimo ovato, apice oblique truncato et inferne acuminato, nigrae. Caput thoracis fere latitudine, suborbiculatum, leviter convexum, supra oculos utrinque punctis paucis sparsis, inter oculos punctis 4 transversim positis impressum. Thorax coleopteris paulo angustior, basin versus subangustatus, latitudine vix longior, lateribus subrectis vel pone medium levissime sinuatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, leviter rotundatis, parum convexus, seriebus dorsalibus punctis 4 sat remotis, aequaliter distantibus, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus profundioribus. Scutellum confertum punctulatum, tenuiter griseo-pubescent. Elytra thorace parum longiora, parcius foriter profundeque punctata, parce griseo-pubescentia. Abdomen subtiliter punctatum, griseo-pubescent, segmentis ventralibus ferrugineo-marginatis. Pedes plerumque piceo-testacei, interdum picei vel nigro-picei, coxis intermediis leviter distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

Obs. Caput in hac specie plerumque thoracis latitudine, in utroque sexu aequale; invidua mascula solito paulo maiora et capite paulo grossiore instructa a Prof. Nordmann nomine Ph. pachycephali pro peculiari specie descripta sunt. Differunt a Ph. cephalote elytrorum punctura rariore profundioreque et capite magis orbiculato.

\* 48. *Ph. fuscus: Elongatus, niger, nitidus, thorace, elytris pedibusque rufis, abdominis segmentis rufo-marginatis. — Long. 3 lin.*

*Nordm. Symb. 90. 54. — Er. Col. March. I. 461. 22.*

*Staph. fuscus Grav. Micr. 29. 41. Mon. 59. 26. — Latr. Hist. nat. Crust. et Ins. IX. 317. 46. — Gyll. Ins. Suec. II. 329. 45. — Mannerh. Brachelytr. 29. 59.*

*Staph. subuliformis Grav. Micr. 29. 42. Mon. 61. 30. — Latr. Hist. nat. Crust. et Ins. IX. 317. 47.*

*Staph. terminatus Grav. Micr. 30. 43. — Latr. Hist. nat. Crust. et Ins. IX. 317. 48.*

*Staph. fragilis Grav. Micr. 30. 44. — Latr. Hist. nat. Crust. et Ins. IX. 317. 49.*

Habitat in Germania, Suecia.

Statura praecedentium. Antennae tenues, capite parum longiores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus et crassioribus, 5—10 transversim obconicis, ultimo subovato, apice oblique emarginato et inferne acuminato, rufae, medio magis minusve piceae, apice ferrugineae. Labrum et palpi rufa, mandibulae rufo-piceae. Caput feminae minoribusque individuis masculis thoracis vix latitudine, maribus maioribus thorace paulo latius, orbiculatum, leviter convexum, supra oculos utrinque punctis paucis, inter oculos punctis 4 transversim positis impressum, nigrum, nitidum. Thorax coleopteris paululum angustior, latitudine vix longior, lateribus subrectis, parallelis, basi rotundatus, angulis posterioribus obtusis, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, parum convexus, seriebus dorsalibus punctis 4, posterioribus tribus remotio-



ribus, compositis, lateribus punctis praeter marginalia utrinque circiter 5 impressus, punctis omnibus subtilioribus, rufus, nitidus, dorso antice plerumque nigricans. Scutellum nigricans, dense subtiliter punctulatum, griseo-pubescentis. Elytra thorace parum longiora, dense subtiliter punctata, subtiliter griseo-pubescentia, rufo-picea. Abdomen subtiliter punctulatum, subtiliter griseo-pubescentis, nigrum, segmentis singulis rufo-marginatis. Pectus nigrum. Pedes cum coxis anterioribus rufi, coxis intermediis leviter distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

Variat thorace toto rufo.

Obs. Caput magnitudine variat ut in *Ph. cephalote*, at in hac specie caput feminae thorace angustius, masculorum validiorum thorace parum latius et magis orbiculatum.

\* 49. *Ph. placidus*: *Elongatus, niger, nitidus, pedibus elytrisque rufis, his circa scutellum nigricantibus.* — Long. 3 lin.

*Var. Elytris rufis, plaga communi lata, apice abbreviata, nigra.*

Habitat in Chersoneso Taurica, Dom. Schüppel et Parreyss, in Hungaria, Dom. Frivaldsky.

Statura omnino sequentis, niger, nitidus, capite thoraceque nitidissimis, elytris rufis. Antennae capite cum thorace breviores, articulo tertio secundo paulo longiore, 4 — 10 paulo brevioribus at haud crassioribus, penultimis haud transversis, ultimo ovato, apice oblique emarginato et inferne acuminato, ferrugineae, basi nigro-fuscae. Palpi dilute rufi. Caput orbiculatum, maiusculum, plerumque thoracis latitudine vel masculis maioribus paulo latius, supra oculos punctis parvis dispersis, inter oculos punctis 4 transversim positis subtilioribus impressum. Thorax coleopteris parum angustior, latitudine haud brevior, lateribus parallelis, subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, leviter convexus, seriebus dorsalibus punctis 4 aequaliter distantibus compositis, lateribus punctis praeter marginalia saepius 5 impressus, punctis omnibus sat subtilibus. Scutellum nigrum, crebre punctatum, tenuiter griseo-pubescentis. Elytra thorace parum longiora, crebrius subtiliusque punctata, parce tenuiterque griseo-pubescentia, rufa, nitida, circa scutellum tenui spatio nigricantia. Abdomen parce subtiliterque punctatum, griseo-pubescentis, segmentis ventralibus obscure rufo-marginatis. Pedes rufi, coxis omnibus nigris, intermediis leviter distantibus, femoribus posterioribus saepe nigricantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

Obs. Varietatis supra memoratae specimen singulum masculum, genuinis paulo maior, capite validiore, thorace paulo longiore, ceterum nullo modo distinctum, a Dom. Dr. Frivaldsky e Hungaria communicatum est.

\* 50. *Ph. blandus*: *Nitidus, rufus, capite, pectore, elytris abdominisque apice nigris, capite orbiculato.* — Long. 3 lin.

*Staph. blandus Grav. Mon. 72. 60.*

Habitat in America septentrionali, Knoch.

Statura omnium praecedentium. Antennae capite thoraceque multo breviores, articulo tertio secundo paulo longiore, sequentibus sensim paulo

crassioribus at vix brevioribus, penultimis subtransversis, ultimo subovato, apice oblique emarginato et inferne acuminato, nigrae, basi obscure testaceae, apice pallidae. Palpi obscure testacei. Caput maiusculum, thoracis latitudine, orbiculatum, supra oculos punctis parvis dispersis, inter oculos 4 transversim positis subtilioribus impressum, nigrum, nitidum. Thorax coleopteris angustior, latitudine paulo longior, lateribus parallelis, subrectis, vel ante basin levissime sinuatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, modice convexus, punctis serierum et nonnullis utrinque inter series et marginem lateralem sitis subtilioribus, testaceo-rufus, nitidus. Scutellum punctatum, nigrum. Elytra thorace parum longiora, nigra, nitida, parcius punctata, tenuiter griseo-pubescentia. Abdomen parce subtiliterque punctulatum, nitidulum, rufum, segmentis ultimis tribus nigricantibus. Pectus nigrum. Pedes cum coxis rufo-testacei, coxis intermediis tenui spatio distantibus, tibiis omnibus spinulosis, tarsis anticis maris simplicibus.

Obs. In specimine unico, quod in Museo regio exstat, thorax serie punctorum dorsali altera 4-, altera 5-punctata, punctis lateralibus in altera latere 6, in altero 8 impressus est; cum praecedentibus summa affinitate iuncta haec species videtur.

### Species mihi invisae.

51. *Ph. suturalis*: Niger, nitidus, antennis et pedibus fusco-piceis, elytris rufis, crebre punctatis, sutura nigra. — Long.  $4\frac{1}{3}$  lin.

Nordm. Symb. 91. 57.

Habitat in Sibiria et Russia orientali.

Ph. polito paulo brevior, capite thoraceque maioribus, antennis multo brevioribus. Antennae capite paulo longiores, extrorsum nonnihil crassiores, fusco-pubescentes. Palpi mandibulaeque nigro-picei. Caput magnum, thorace tamen vix latius, subquadrangulum, angulis rotundatis, nigrum, nitidissimum, laeve, inter oculos puncta 4, et in angulis verticis puncta plura pilifera, maiuscula, profunde impressa. Thorax magnus, antice truncatus, angulis compressis, lateribus versus basin subobliquis et paululum sinuatis, postice obtuse rotundatus, supra sat convexus, tenuissime marginatus, niger, nitidissimus, in angulis anticis puncta quatuor maiora, quorum tria oblique versus latus, quartum ad marginem anticum impressum, series dorsales profunde punctatae. Scutellum nigrum, subnitidum, creberrime et subtilissime punctatum. Elytra thorace latiora et longiora, supra depressa, rufo-testacea, creberrime et subtiliter punctulata, flavescenti-pubescentia, margine suturali laevi, nitido, nigra. Alae dilute fuscescentes. Abdomen elytris fere duplo longius, nigrum, nitidum, crebre et rude punctulatum, pubescens et nigro-pilosum, segmento ultimo in mare sat profunde emarginato. Pedes toti nigro-picei, tarsis anticis in mare paulo dilatatis. (Nordm.)

Obs. An a Ph. bimaculato satis distinctus?

52. *Ph. microcephalus*: Nitidus, niger, pedibus rufis, capite suborbiculato, thorace seriebus dorsalibus 4-punctis. — Long. 3 lin.

*Staph. microcephalus* Grav. Mon. 70. 52.

Habitat Lutetiae.

Caput latitudine paulo longius, thorace minus, nigrum, antennis capituli longitudine et palpis fuscis. Thorax coleopteris paulo minor, 2, 3, 4 punctus. Coleoptera quadrata, fusciscenti-nigricantia, subsericantia. Abdomen nigrum, griseo-sericans, segmentis marginibus ventralibus summis rufis. Pedes rufi, capitibus femorum obscuris. (Grav.)

5) Thorax seriebus dorsalibus 5-punctatis.

\* 53. *Ph. flavipennis*: Niger, nitidus, elytris lucide testaceis, summa basi violaceis. — Long. 5 lin.

Habitat Bogotae in Columbia interiore, a Dom. Buquet cum Staph. flavipennis Dej. nomine communicatus.

Nigerrimus, nitidus, capite thoraceque nitidissimis, elytris solis testaceis, metallico-nitentibus, basi angusta violacea. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, 6 — 10 transversis, ultimo apice emarginato et inferne acuminato, nigrae. Palpi nigri. Mandibulae longiusculae, falcatae, acutae. Caput maris thorace paulo latius, feminae eiusdem latitudine, subquadratum, utrinque supra oculos punctis parvis, inter oculos posterius punctis utrinque tribus oblique positis, anterieus duobus transversim positis, sat profundis, notatum, fronte antice subimpressa. Thorax coleopteris multo angustior, latitudine vix brevior, neque basin neque apicem versus angustatus, lateribus leviter rotundatis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, posterioribus obtusis, supra parum convexus, seriebus dorsalibus 5-punctatis, lateribus punctis praeter marginalia utrinque 5 vel 6 impressus, punctis omnibus minus profundis. Scutellum crebre punctatum, nigrum, subtiliter griseo-pubescent. Elytra thorace plus sesqui longiora, subtiliter parceque punctata, flavescenti-pubescentia. Abdomen supra segmentis singulis lateribus subtilius punctulatis, tenuiter nigro-pubescentibus, medio laevigatis, infra punctulatum. Pedes graciles, coxis intermedii distantibus, tibiis omnibus subtilius spinulosis, tarsis anticis in utroque sexu simplicibus.

\* 54. *Ph. candens*: Niger, nitidus, elytris crebrius punctatis, cupreis, fulgidis. — Long. 4 — 5 lin.

Habitat Bogotae in Columbia interiore, nom. Staph. candentis Dej. a Dom. Buquet et Dejean communicatus.

Nigerrimus, nitidus, capite thoraceque nitidissimis, elytris fulgidis, cupreis, lateribus sensim virescenti-cupreis. Antennae capitis thoracisque fere longitudine, articulo tertio secundo paulo longiore, 4 — 10 elongato-subobconicis, sensim paulo brevioribus, ultimo ovato, apice oblique emarginato et inferne acuminato. Mandibulae longiusculae, falcatae, acutae. Caput thorace latius, maris interdum multo latius, ovatum, fronte supra oculos punctis paucis et inter oculos punctis 4, omnibus transversim positis, impressum, inter antennis in mare foveolatum. Thorax coleopteris duplo fere angustior, latitudine vix longior, basin versus subangustatus, lateribus anterieus leviter rotundatis, posterius subsinuatis, basi leviter rotundatis, apice truncatus, angulis anterioribus deflexis, rotundatis, posterioribus obtusis, leviter convexus, seriebus dorsalibus punctis 5, apicali remotiore, compositis,



lateribus punctis utrinque 5 vel 6 impressus, punctis omnibus subtilioribus. Scutellum subtilius crebre punctatum. Elytra thorace paulo longiora, crebrius punctata, pube parca brevi depressa grisea vix conspicua vestita. Abdomen apicem versus sensim angustatum, subtiliter parce punctulatum, parce nigro pubescens. Pedes elongati, graciles, coxis intermediis distantibus, tibiis omnibus subtiliter spinulosis, tarsis anticis in utroque sexu simplicibus.

\* 55. *Ph. gratiosus*: *Niger, nitidus, elytris parce punctatis, cupreo-purpureis, fulgidis.* — *Long.*  $3\frac{1}{2}$  lin.

Habitat Bogotae in Columbia interiore, Dom. Com. Dejean.

Praecedenti affinis, at duplo minor, nigerrimus, nitidus, capite thoraceque nitidissimis, elytris splendide purpureo-cupreis. Antennae capite thoraceque paulo breviores, articulo tertio secundo sesqui prope longiore, quarto subovato, 5 — 10 subobconicis, crassiusculis, crassitie haud brevioribus, subaequalibus, ultimo ovato, apice abrupte subacuminato. Caput thoracis latitudine, ovatum, posterius utrinque parce fortiter punctatum, inter oculos punctis 4 transversim positis notatum, antice inter antennis sulco longitudinali abbreviato impressum. Thorax coleopteris duplo angustior, latitudine paulo longior, lateribus pone medium subsinuatis, basi leviter rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, posterioribus obtusis, convexus, seriebus dorsalibus punctis 5, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 vel 6 impressus, punctis subtilioribus. Scutellum crebre punctulatum, tenuiter pubescens. Elytra thorace sesqui longiora, parce punctata, tenuiter pilosella. Abdomen apicem versus sensim angustatum, punctis raris subtilibus impressum, parce nigro-pilosum. Pedes elongati, graciles, coxis intermediis distantibus, tibiis omnibus subtilius spinulosis, tarsis anticis in utroque sexu simplicibus.

X \* 56. *Ph. ebeninus*: *Niger, nitidus, elytris viridi-aeneis, abdomine parcius punctato, capite suborbiculato.* — *Long.*  $2\frac{1}{2}$  — 4 lin.

*Philonth. varians* Er. Col. March. I. 461. 23.

*Var. maior*:

*Staph. ebeninus* Grav. Micr. 170. 22. Mon. 67. 21. — Latr. Hist. nat. Crust. et Ins. IX. 323. 66. — Gyll. Ins. Suec. II. 326. 52. — Sahlb. Ins. Fenn. I. 325. 48. — Mannerh. Brachelytr. 30. 66. — Boisd. et Lacord. Faun. Ent. Paris. I. 394. 15.

*Philonth. ebeninus* Nordm. Symb. 93. 63.

*Staph. brevicornis* Grav. Micr. 22. 28. Mon. 69. 50. — Latr. Hist. nat. Crust. et Ins. IX. 312. 33.

*Philonth. brevicornis* Nordm. Symb. 94. 69.

*Var. minor*:

*Staph. varians* Grav. Mon. 82. 73. — Gyll. Ins. Suec. II. 342. 58. — Zetterst. Faun. Lappon. I. 77. 27. Ins. Lappon. 65. 37. — Sahlb. Ins. Fenn. I. 329. 55. — Mannerh. Brachelytr. 30. 73. — Boisd. et Lacord. Faun. Ent. Paris. I. 394. 16. — Runde Brachelytr. Hal. 9. 47.

*Philonth. varians* Nordm. Symb. 93. 65.

*Staph. concinnus* Grav. Micr. 21. 25.

*Var. Pedibus fusco-testaceis:*

*Staph. ochropus* Grav. *Mic.* 39. 57. *Mon.* 67. 39. — *Latr. Hist. nat. Crust. et Ins.* IX. 322. 62. — *Gyll. Ins. Suec.* II. 336. 51. — *Mannerh. Brachelytr.* 29. 65. — *Runde Brachelytr. Hal.* 8. 40.

*Philonth. ochropus* Nordm. *Symb.* 93. 62.

Habitat in Europa, Sibiria, Asia minore, Syria, Aegypto, frequens.

Variat magnitudine, individuis maioribus *Ph. atrato*, minoribus *Ph. vario* aequalibus, niger, capite thoraceque pernitidis, elytris viridi-aeneis, nitidis. Antennae capitis thoracisque longitudine, articulo tertio secundo paulo longiore, 4—10 subobconicis, sensim perparum brevioribus crassioribusque, penultimo haud transverso, ultimo subovato, apice emarginato et inferne subacuminato, nigrae. Caput suborbiculatum, individuis maioribus thoracis latitudine, minoribus eodem minus et angustius, supra oculos utrinque punctis parvis vagisque, inter oculos punctis utrinque duobus transversim positis impressum. Thorax prope coleopterorum latitudine, latitudine vix brevior, antrorsum subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, modice convexus, seriebus dorsalibus punctis 5, apicali remotiore compositis, lateribus utrinque punctis praeter marginalia 4 impressus, punctis omnibus modice profundis. Scutellum dense subtiliter punctulatum, subtiliter pubescens. Elytra thoracis longitudine, subtilius densiusque punctata, tenuiter griseo-pubescentia. Abdomen subtilius et minus dense punctulatum, parce pubescens. Pedes nigri, saepe picei vel interdum subtestacei, coxis intermediis interstitio angusto distinctis, tibiis omnibus spinulosis. Alae infuscaetae.

Tarsi antici maris fortiter, feminae leviter dilatati.

Variat 1, Pedibus fusco-testaceis.

2, Elytris rufis.

3, Thoracis seriebus dorsalibus altera 5-, altera 4-punctata.

Obs. 1. Individuis minoribus, ubique copiose obviis sexus habitu simillimi, maioribus, minus frequentibus, mares capite maiore, thoracis latitudine, instructi sunt, simili modo ut in *Ph. aeneo*, *carbonario*, *cephalote*, etc.; aliae differentiae speciminibus plurimis examinatis comparatisque autem desiderantur, quamobrem plures ab auctoribus distinctas species huic subiunxi. *Staph. ebeninus* et *brevicornis* Gr. ad individua maxima spectant, *Staph. ochropus* ad varietatem pedibus fusco-testaceis pertinet. Magis discedere videtur var. sub n. 2 indicata, cuius exemplum singulum, minimis huius speciei individuus magnitudine aequale, Dom. Schüppel Berolini legit, quod autem praeter elytrorum colorem nullo modo ab eadem differt, et cum varietates analogas iam apud *Ph. varium* et *albipedem* observaverim, eo magis hanc pro specie peculiari habere dubito.

2. *Staph. variantem* auctorum veterum ab hoc satis distinctum esse patet, quare hoc nomen illi restitui.

\* 57. *Ph. corvinus*: Niger, nitidus, elytris concoloribus, abdomine supra parcius punctato, capite orbiculato. — Long. 3½ lin.

*Er. Col. March.* I. 462. 24.

Habitat in Germania.

Statura omnino praecedentis, maioribusque eius, non autem maximis individuus magnitudine aequalis, his vero paulo gracilior, nigerrimus, nitidus, elytris concoloribus, capite thoraceque pernitidis. Antennae capitis thoracisque longitudine, articulo tertio secundo paulo longiore, 4—10 oblongo-subobconicis, sensim paulo brevioribus at vix crassioribus, omnibus crassitie

longioribus, ultimo oblongo ovato, apice subemarginato et inferne acuminato. Caput thoracis latitudine, suborbiculatum, modice convexum, supra oculos utrinque fortiter parce vageque punctatum, fronte inter oculos utrinque punctis maioribus, superius tribus oblique positis, anterieus duobus transversim positis impressa. Thorax coleopteris latitudine prope aequalis, latitudine vix brevior, antrorsum subangustatus, lateribus subrectis, basi rotundatus, angulis posterioribus obtusis, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, modice convexus, seriebus dorsalibus punctis 5, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 4 vel 5 impressus, punctis omnibus modice profundis. Scutellum subtiliter punctulatum, tenuiter pubescens. Elytra thoracis longitudine, subtiliter minus dense punctata, tenuiter nigro-pubescentia. Abdomen parcius subtiliusque punctatum, parcius nigro-pubescentis. Pedes nigri, coxis intermediis interstitio angusto distantibus, tibiis omnibus spinulosis, tarsis, nonnunquam etiam tibiis piceis.

Tarsi antici maris fortiter, feminae leviter dilatati.

Obs. A praecedente constanter elytrorum colore nigerrimo, a sequente abdominis punctura distinctus.

\*58. *Ph. fumigatus*: *Niger, nitidus, elytris nigro-subaeneis, pedibus fuscis, tibiis tarsisque ferrugineis, abdomen supra crebrius subtiliter punctulato, capite orbiculato.* — Long.  $3\frac{1}{2}$  lin.

*Er. Col. March. I.* 463. 25.

*Staph. atrati* var. *intermed. Grav. Mon. p.* 85.

Habitat in Germania, Gallia.

Statura omnino praecedentium, minoribus *Ph. ebenini* individuis paulo maior, niger, capite thoraceque pernitidis, elytris nigro-aeneis, nitidulis. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, sequentibus subobconicis, sensim paululum brevioribus et crassioribus, penultimis crassitie haud brevioribus, ultimo apice truncato et inferne subacuminato, nigrae. Palpi picei. Caput maiusculum, thorace tamen paulo angustius, suborbiculatum, supra oculos utrinque punctis parvis vagisque, inter oculos punctis utrinque duobus transversim positis impressum. Thorax coleopteris haud angustior, latitudine haud brevior, antrorsum subangustatus, lateribus subrectis, basi rotundatus, angulis posterioribus obtusis, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, leviter convexus, seriebus dorsalibus punctis 5, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia subtiliora utrinque 5 impressus, punctis omnibus minus profundis. Scutellum dense punctulatum, tenuiter griseo-pubescentis. Elytra thoracis longitudine, densius fortiusque punctata, tenuiter griseo-pubescentia. Abdomen apicem versus sensim angustatum, praecipue supra confertim subititerque punctulatum, pube subtili densiore depressa vestitum, versicolor, subtus segmentis singulis tenuiter ferrugineo-marginatis. Pedes nigro-picei, coxis intermediis leviter distantibus, tibiis anticis muticis, posterioribus spinulosis tarsisque ferrugineis, tarsis anticis in utroque sexu, licet in mare fortius, dilatatis.

Obs. Differt ab affinibus praecipue thorace latiore, abdomine confertim subtiliter punctulato, tibiis anticis haud spinulosis.



- \* 59. *Ph. umbrinus*: *Nigro-piceus, nitidus, pedibus fusco-testaceis, abdomine supra parce punctato, segmentis ventralibus rufo-marginatis.* — Long. 3 lin.

*Mas* Capite orbiculato.

*Staph. umbrinus* Grav. Micr. 169. 22. Mon. 69. 47.

*Fem.* Capite subovato.

*Staph. moestus* Grav. Mon. 47. 9.

*Philonth. moestus* Nordm. Symb. 97. 73.

Habitat in America septentrionali.

Statura omnino *Ph. ebenini* et magnis eius individuis magnitudine aequalis. Antennae capite thoraceque multo breviores, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo brevioribus et crassioribus, penultimis leviter transversis, ultimo ovato, apice emarginato et inferne acuminato, fusco-piceae, articulo ultimo subferrugineo. Mandibulae et palpi picei. Caput maris thoracis apice vix angustius, orbiculatum, feminae thorace multo angustius, ovatum, utrinque supra oculos parce punctatum, fronte inter oculos punctis maioribus utrinque posterioribus tribus oblique, anterioribus duobus transversim positis impressa, nigrum, nitidissimum. Thorax basi coleopteris haud angustior, longitudine paulo latior, antrorsum angustatus, antice subcompressus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, rotundatis, modice convexus, seriebus dorsalibus punctis 5, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus profundioribus, nigro-piceus vel piceus, nitidissimus. Elytra thoracis longitudine et scutellum nigro-picea, vel picea, nitida, crebre punctata, griseo-pubescentia. Abdomen parce punctulatum, griseo pubescens, nigrum, nitidulum, segmentorum ventralium marginibus rufo-piceis. Pedes cum coxis anterioribus obscure testacei, coxis intermediis leviter distantibus, tibiis omnibus spinulosis.

Tarsi antici maris fortiter, feminae leviter dilatati.

Variat thoracis serie dorsali altera 4-, altera 5-punctata.

- \* 60. *Ph. figulus*: *Niger, nitidus, abdomine supra parcius punctato, versicolore, segmentis ventralibus anoque rufis, femoribus testaceis, capite subovato.* — Long. 4 lin.

Habitat in Brasilia, Dom. Westermann.

Statura praecedentium, niger, nitidus, capite thoraceque pernitidis. Antennae capitis thoracisque fere longitudine, crassiusculae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus, obconicis, ultimo apice truncato et inferne acuminato, nigrae, apice obscure ferrugineae. Palpi picei. Caput thorace paulo angustius, breviter subovatum, supra leviter convexum, utrinque supra oculos punctis paucis, inter oculos binis, oculo approximatis, oblique positis, impressum. Thorax basi coleopteris paulo angustior, latitudine baseos paulo longior, antrorsum modice angustatus, lateribus rectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrectis, transversim convexus, seriebus dorsalibus punctis 5 aequaliter distantibus compositis, lateribus punctis

praeter marginalia utrinque 3, longitudinaliter dispositis impressus, punctis omnibus fortioribus, piliferis. Scutellum confertim subtilius punctulatum, densius nigro-pubescentis. Elytra thorace paulo longiora, densius subtiliusque punctata, minus subtiliter nigro-pubescentia. Abdomen supra parcius, infra crebrius minus subtiliter punctulatum, parcius minus subtiliter nigro-pubescentis, supra laete metallico-versicolor, segmento quinto apice sextoque toto rufo-ferrugineis, praeterea segmentis ventralibus quatuor primis determinate rufo-marginatis. Pedes coxis anterioribus piceo-testaceis, femoribus omnibus testaceis, anticis dilutioribus, tibiis tarsisque omnibus nigris, tibiis omnibus spinulosis.

Tarsi antici maris fortius dilatati; femina latet.

\* 61. *Ph. suspectus*: *Niger, nitidus, elytris subvirescentibus, pedibus anterioribus obscure testaceis, capite suborbiculato*. — Long.  $3\frac{2}{3}$  lin.

Habitat prope Caracas Columbiae, Dom. Moritz.

Statura omnino *Ph. ebenini* et maximis eius individuis magnitudine aequalis, niger, nitidus. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 4 — 10 crassiusculis, crassitie subaequalibus, sensim paulo brevioribus, ultimo apice emarginato et inferne subacuminato, nigro-piceae. Palpi picei. Caput thoracis prope latitudine, suborbiculatum, leviter convexum, fronte supra punctis parvis, vagis, inter oculos punctis quatuor transversim positis, intermediis maioribus, dilatatis, multo magis distantibus, notata, antice inter antennas longitudinaliter leviter impressa. Thorax coleopteris paululum angustior, latitudine sublongior, lateribus parallelis, subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, modice convexus, seriebus dorsalibus punctis 5, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia subtiliora utrinque 5, duobus eorum margini approximatis, impressus, punctis sat profundis. Scutellum punctulatum, tenuiter pubescens. Elytra thoracis longitudine, minus crebre punctata, tenuiter pubescentia, nigro-subvirescentia, nitida. Abdomen parcius punctatum, tenuiter nigro-pubescentis, subtus segmentis singulis margine posteriore rufo-piceo. Pedes anteriores obscure testacei, postici nigri, coxis intermediis modice distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

\* 62. *Ph. corruscus*: *Niger, nitidus, elytris rufis, capite suborbiculato*. — Long.  $3\frac{1}{2}$  lin.

*Nordm. Symb.* 98. 81. — *Er. Col. March.* I. 463. 26.

*Staph. corruscus Grav. Micr.* 33. 47. *Mon.* 51. 14. — *Latr. Hist. nat. Crust. et Ins.* IX. 319. 52.

*Staph. nitidus Marsh. Ent. Britt.* 511. 38.

*Staph. planus Boisd. et Lacord. Faun. Ent. Paris.* I. 401. 27.

Habitat in Europa temperata.

Statura omnino *Ph. ebenini* et mediis eius individuis magnitudine aequalis, niger, nitidus, capite thoraceque pernitidis, elytris saturate lateritiis. Antennae capitis thoracisque longitudine, articulo secundo tertioque aequalibus, 4—10 subobconicis, sensim paululum brevioribus crassioribusque, penultimis crassitie haud brevioribus, ultimo apice leviter emarginato et inferne

subacuminato, nigrae. Palpi nigri. Caput thorace minus et angustius, orbiculatum, supra oculos utrinque punctis paucis, inter oculos utrinque duobus, transversim positis, fortibus impressum. Thorax basi coleopterorum fere latitudine, latitudine vix brevior, antrorsum subangustatus, lateribus subrectis, basi rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, leviter convexus, seriebus dorsalibus punctis 5, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia subtiliora utrinque 5 impressus, punctis modice profundis. Scutellum confertim punctulatum, longius griseo-pubescent. Elytra thoracis longitudine, minus subtiliter et minus dense punctata, tenuiter griseo-pubescentia. Abdomen parcius subtiliusque punctulatum, parce tenuiterque griseo-pubescent. Pedes nigri, coxis intermediis interstitio angusto distantibus, tibiis omnibus spinulosis, piceis, tarsis rufescentibus.

Tarsi antici maris modice, feminae vix dilatati.

\* 63. *Ph. Caucasicus*: *Niger, nitidus, elytris rufis, basi inaequaliter nigris, antennis pedibusque fusco-piceis, capite suborbiculato.* — Long. 3 lin.

*Nordm. Symb.* 99. 22.

Habitat in Asia occidentali.

Statura omnino *Ph. ebenini* et minoribus eius individuis aequalis, niger, nitidus, capite thoraceque pernitidis, elytris rufis, dorso macula magna basali ad humerum dilatata et cum sutura descendente nigra. Antennae capite thoraceque multo breviores, articulo secundo tertioque aequalibus, 4—10 crassiusculis, sensim paulo brevioribus, penultimis leviter transversis, ultimo ovato, apice emarginato et inferne acuminato, nigro-piceae, articulis tribus primis nigris. Palpi nigro-picei. Caput thorace paulo angustius, suborbiculatum, leviter convexus, utrinque punctis supra oculos parvis dispersis, inter oculos binis transversim positis impressum. Thorax coleopteris parum angustior, latitudine baseos vix longior, apicem versus vix angustatus, lateribus leviter rotundatis, ante basin subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, posterioribus obtusis, modice convexus, seriebus dorsalibus punctis 5, apicali remotiore, compositis, lateribus punctis praeter marginalia subtiliora utrinque 4 impressus, punctis minus subtilibus. Scutellum subtilius punctulatum, griseo-pubescent. Elytra thorace paulo longiora, minus confertim et minus subtiliter punctata, griseo-pubescentia. Abdomen crebrius subtiliter punctulatum, densius longiusque griseo-pubescent. Pedes picei, coxis intermediis modice distantibus, tibiis omnibus spinulosis.

Tarsi antici feminae leviter dilatati. Mas latet.

Obs. Similis *Ph. bimaculato*, at antennis brevioribus, thorace antrorsum vix angustato, elytris basi fusco-nigris neque aeneis, marginè laterali inflexo rufo distinguendus.

64. *Ph. turbatus*: *Niger, nitidus, elytris, palpis, tibiis tarsisque rufis, capite suborbiculato.* — Long. 3 lin.

Habitat in Sardinia, Mus. Reg. Taurin.

Statura omnino *Ph. ebenini*, niger, nitidus, capite thoraceque pernitidis.



Antennae breviusculae, capite sesqui longiores, crassiusculae, fuscae, articulis tribus primis nigris. Palpi rufi. Mandibulae rufo-piceae. Caput thoracis prope latitudine, suborbiculatum, supra oculos utrinque punctis pluribus sat fortibus notatum. Thorax basi coleopterorum prope latitudine, antrorsum subangustatus, lateribus leviter rotundatis, latitudine brevior, modice convexus, punctis ordinariis sat fortibus. Scutellum subtiliter punctulatum, parcius griseo-pubescent. Elytra thorace paulo longiora, minus confertim minusque subtiliter punctata, lateritia, nitida, fulvo-pubescentia. Abdomen crebre minus subtiliter punctatum, nigro-pubescent, segmentorum marginibus concoloribus, sexto subtus apice piceo. Pedes nigro-picei, tibiis tarsisque rufis, tarsis anticis feminae simplicibus.

Obs. Affinis Ph. corrusco, at minor, praecipue brevior et distinctus antennis brevioribus, palpis, tibiis tarsisque rufis. Specimen singulum ad examinandum communicavit Dom. Prof. Gené.

\*65. *Ph. sanguinolentus*: Niger, nitidus, elytris sutura maculaque humerali sanguineis, coxis anticis testaceis, capite suborbiculato. — Long. 4 lin.

Nordm. Symb. 97. 76. — Er. Col. March. I. 464. 27.

*Staph. sanguinolentus* Grav. Micr. 36. 53. Mon. 64. 34. — Latr. Hist. nat. Crust. et Ins. IX. 320. 58. — Gyll. Ins. Suec. II. 338. 54. — Sahlb. Ins. Fenn. I. 326. 50. — Mannerh. Brachelytr. 30. 68. — Boisd. et Lacord. Faun. Ent. Paris. I. 399. 24. — Runde Brachelytr. Hal. 8. 42.

Var. *Elytrorum maculis confluentibus*.

*Staph. contaminatus* Grav. Micr. 174. 31. — Latr. Hist. nat. Crust. et Ins. IX. 325. 71. — Runde Brachelytr. Hal. 9. 45.

Habitat in Europa.

Statura omnino et magnitudine Ph. ebenini individuís maioribus aequalis, niger, capite thoraceque pernitidis, elytris nigris, sutura maculaque humerali sanguineis, plerumque discretis, saepe dilatatis et confluentibus, interdum strictioribus et macula humerali obsoleta. Antennae capite thoraceque breviores, articulo tertio secundo paulo longiore, sequentibus oblongo-subobconicis, inter se subaequalibus, ultimo apice truncato et inferne subacuminato, nigrae. Palpi nigri. Caput maiusculum, thorace paululum angustius, orbiculatum, supra oculos utrinque punctis parvis, inter oculos punctis 4 transversim positis notatum. Thorax prope coleopterorum latitudine, latitudine haud brevior, lateribus parallelis, subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, modice convexus, seriebus dorsalibus punctis 5, apicali remotiore, compositis, lateribus punctis praeter marginalia subtiliora utrinque 5 impressus, punctis profundioribus. Scutellum confertim punctulatum, griseo-pubescent. Elytra thorace vix longiora, confertim subtilius punctata, tenuiter griseo-pubescentia. Abdomen confertim subtilissime punctulatum, puber griseo-pubescentia. Abdomen confertim subtilissime punctulatum, puber subtili densiore subtoementosa fusco-grisea vestitum. Pedes nigri, tibiis tarsisque obscure rufo-piceis, coxis anticis rufo-testaceis, coxis intermediis modice distantibus, tibiis anticis parce subtiliterque, posterioribus longius fortiusque spinulosis, tarsis anticis in utroque sexu, licet in mare fortius, dilatatis.

66. *Ph. stragulatus*: *Niger, nitidus, elytris rufo-testaceis, plaga lata communi longitudinali nigra, coxis anticis piceo-testaceis, capite subovato. — Long. 3 lin.*

Habitat Massiliae, collect. Dom. Aubé.

Habitu omnino *Ph. fumarii*, at paulo minor, niger, nitidus, capite thoraceque pernitidis. Antennae capitis thoracisque longitudine, nigrae. Palpi nigri. Caput thorace paulo angustius, subovatum, utrinque supra oculos parcius punctatum. Thorax basi coleopteris parum angustior, antrorsum leviter angustatus, latitudine baseos vix brevior, transversim convexus, punctis ordinariis minus profundis. Scutellum subtiliter punctulatum. Elytra thorace vix longiora, confertim subtiliter punctata, tenuiter griseo-pubescentia, rufo-testacea, nitida, plaga longitudinali communi sat lata, apicem versus sensim leniter angustata, nigra. Abdomen dense subtiliter punctatum, fusco-pubescent. Pedes picei, antici coxis piceo-testaceis, femoribus testaceo-piceis.

Tarsi antici maris leviter dilatati; femina latet.

- \* 67. *Ph. bipustulatus*: *Niger, nitidus, elytris macula postica sanguinea, coxis anticis nigris, immaculatis, capite ovato. — Long. 3½ lin.*

*Nordm. Symb.* 98. 78. — *Er. Col. March.* I. 464. 28.

*Staph. bipustulatus* *Panz. Faun. Germ.* 27. 10. — *Latr. Hist. nat. Crust. et Ins.* IX. 321. 59. t. 79. f. 5. — *Grav. Mon.* 63. 33. — *Gyll. Ins. Suec.* II. 339. 55. — *Sahlb. Ins. Fenn.* I. 327. 52. — *Mannerh. Brachelytr.* 30. 70. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 399. 25. — *Runde Brachelytr. Hal.* 8. 43.

*Staph. bipustulatus* *var. 3 Grav. Micr.* 37. 54.

Habitat in Europa, in Asia minore, Syria.

Statura omnino *Ph. ebenini* et minoribus eius individuis paulo maior, niger, capite thoraceque pernitidis. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, 4 — 10 obconicis, sensim paulo brevioribus, at vix crassioribus, omnibus crassitie sublongioribus, ultimo ovato, apice emarginato et inferne acuminato, piceae. Caput thorace angustius minusque, ovatum, supra oculos utrinque punctis parvis, inter oculos utrinque punctis duobus transversim positis impressum. Thorax basi prope coleopteris angustior, latitudine paululum longior, antrorsum vix angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrotundatis, modice convexus, seriebus dorsalibus punctis 5, apicali remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis minus subtilibus. Scutellum confertim punctulatum, nigro-pubescent. Elytra thorace parum longiora, confertim subtilius punctata, nigro-pubescentia, macula rufa, infra medium sita, notata. Abdomen subtilius parciusque punctatum, nigro-pubescent. Pedes nigri, coxis anticis plerumque immaculatis, rarius interne testaceis, intermedii interstitio angusto distantibus, tibiis omnibus spinulosis. Alae albo-hyalinae.

Tarsi antici maris fortiter, feminae leviter dilatati.

Variat elytris immaculatis.

Obs. 1. *Ph.* varianti affinis, at capite minus ovato, thorace antrorsum minus an-

gustato, elytrorum abdominisque punctura paulo fortiore, elytrorum macula ante apicem neque in medio sita et coxis anticis concoloribus distinguendus.

2. *Staph. bipustulatus* L., „magnitudine minoris pediculi et antennis clavatis” potius ad *Aleocharam* quandam pertinet. (Conf. quae supra apud Al. nitidam p. 168. monui.) *Staph. bipustulatus* Fab. Syst. 266. 11. Spec. Ins. I. 336. 12. Mant. Ins. I. 221. 18, nonnisi diagnosi et citato Linneano descriptus. *Staph. bipustulatus* Fab. Ent. Syst. I. II. 526. 34, Syst. El. II. 598. 44, „elytrorum puncto postico ferrugineo” distinctius ad *Aleocharam* quandam, *A. bipustulatum*, tristem nitidamve spectare, huic autem generi alienus videtur. In collectione Fabriciana de hac specie nihil mihi memoratum.

- \* 68. *Ph. aeruginosus*: *Niger, nitidus, coxis anticis intus testaceis, alis fuscis, abdomine fortius punctulato, segmentis ventralibus rufo-marginatis, capite ovato.* — Long. 4 lin.

*Nordm. Symb.* 95. 71.

Habitat in America meridionali: in Brasilia, Dom. Sellow, in Columbia, Dom. Moritz.

Statura et summa affinitas *Ph. variantis*, at maximis eius individuis fere maior et robustior, totus nigerrimus, capite thoraceque nitidissimis, elytris et abdomine nitidulis, coxarum anticarum parte interiore, interdum etiam femoribus anterioribus testaceis. Antennae capitis thoracisque longitudine, graciles, articulo tertio secundo paulo longiore, sequentibus elongato-subobconicis, subaequalibus, ultimo oblongo, apice oblique emarginato et inferne acuminato, fusco-nigrae. Palpi nigro-picei. Caput thorace minus et angustius, ovatum, utrinque supra oculos punctis paucis, inter oculos binis oblique positis impressum. Thorax basi coleopteris angustior, latitudine baseos haud longior, antrorsum leviter angustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, modice convexus, seriebus dorsalibus punctis 5, postremo paululum remotiore, compositis, lateribus punctis praeter marginalia subtilia utrinque 5 impressis, punctis modice profundis. Elytra thorace paulo longiora et scutellum crebre subtiliter punctata, densius griseo-pubescentia. Abdomen iridescens, parcius et fortius punctulatum, pilis appressis nigris parcius vestitum, segmentis ventralibus rufo-marginatis. Pedes graciliores, coxis intermediis approximatis, tibiis omnibus spinulosis. Alae fuliginosae.

Tarsi antici maris modice, feminae leviter dilatati.

- \* 69. *Ph. feralis*: *Nigerrimus, nitidus, alis fuscis, abdomine subtilissime punctulato, capite parvulo, ovato.* — Long.  $3\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Statura et summa affinitas praecedentis, at dimidio minor. Corpus totum nigerrimum, capite thoraceque nitidissimis, elytris et abdomine nitidulis. Antennae capite thoraceque paulo longiores, articulo tertio secundo paulo longiore, 4—10 sensim brevioribus at haud crassioribus, omnibus crassitie longioribus, subobconicis, ultimo oblongo-ovato, apice truncato et inferne acuminato. Caput thorace multo minus et angustius, ovatum, convexus, utrinque supra oculos punctis paucis dispersis et inter oculos binis oblique positis notatum. Thorax latitudine baseos tertia parte longior, basi coleopteris dimidio angustior, apicem versus leviter angustatus, lateribus subrectis,



basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, transversim convexus, seriebus dorsalibus punctis 5 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis omnibus subtilioribus. Scutellum subtiliter punctulatum, tenuiter nigro-pubescent. Elytra thoracis longitudine, crebrius subtiliusque punctulata, densius nigro-pubescentia. Abdomen concolor, subtilissime parce punctulatum, pube parca longiore depressa nigra tenuiter vestitum, segmentorum ventralium marginibus summis obscure rufescentibus. Pedes nigro-picei, coxis anticis intus piceis, intermediis leviter distantibus, tibiis omnibus spinulosis. Alae fuscae, basi dimidiatim albo-hyalinae.

Tarsi antici maris fortius dilatati, feminae simplices. Segmentum abdominis inferius sextum feminae apice emarginatum.

Obs. Proximus praecedenti, distinctus autem statura minore, thorace longiore, abdominis punctura subtilissima.

\*70. *Ph. varians*: *Niger, nitidus, elytris macula media sanguinea, coxis anticis testaceis, capite ovato.* — Long. 3 —  $3\frac{1}{2}$  lin.

*Staph. varians* Payk. Mon. 45. 33. Faun. Suec. III. 393. 33.

*Staph. opacus* Grav. Mon. 64. 35. — Gyll. Ins. Suec. II. 340. 56. — Zetterst. Faun. Lappon. I. 77. 25. Ins. Lappon. 64. 35. — Sahlb. Ins. Fenn. I. 328. 53. — Mannerh. Brachelytr. 30. 71. — Boisd. et Lacord. Faun. Ent. Paris. I. 404. 34.

*Philonth. opacus* Nordm. Symb. 98. 79. — Er. Col. March. I. 465. 29.

*Staph. bipustulatus* var. 1, 2, 4 — 7 Grav. Micr. 37. 54.

*Staph. bimaculatus* Marsh. Ent. Britt. 525. 78.

Var. a, *Elytris immaculatis.* — Long. 3 —  $3\frac{1}{2}$  lin.

*Staph. varians* Fab. Ent. Syst. I. II. 524. 22. Syst. El. II. 594. 25. — Oliv. Ent. III. 42. 27. 35. t. 3. f. 46. — Panz. Ent. Germ. 354. 15.

*Staph. opacus* Grav. Micr. 26. 35. — Latr. Hist. nat. Crust. et Ins. IX. 315. 40.

*Staph. aterrimus* Marsh. Ent. Britt. 513. 44.

Var. b, *Elytris immaculatis, antennis articulo primo subtus flavo.* — Long.  $3\frac{1}{2}$  lin.

*Philonth. scybalarius* Nordm. Symb. 94. 70.

*Philonth. fuscicornis* Nordm. Symb. 96. 72.

Var. c, *Niger, immaculatus.* — Long. 2 —  $2\frac{1}{2}$  lin.

*Staph. simplex* Marsh. Ent. Britt. 505. 20. (forte)

Var. d, *Niger, elytris fuscis, apice ferrugineis, pedibus fuscis.* — Long. 2 —  $2\frac{1}{2}$  lin.

*Staph. agilis* Grav. Mon. 77. 70. — Gyll. Ins. Suec. II. 341. 57. — Sahlb. Ins. Fenn. I. 328. 54. — Zetterst. Faun. Lappon. I. 77. 26. Ins. Lappon. 64. 36. — Mannerh. Brachelytr. 30. 72.

*Philonth. agilis* Nordm. Symb. 98. 80.

*Staph. parvicornis* Grav. Micr. 23. 30, Mon. 69. 48. — Latr. Hist. nat. Crust. et Ins. IX. 313. 35.

*Staph. discoideus* Boisd. et Lacord. Faun. Ent. Paris. I. 403. 33. (forte)

Habitat in Europa; ad Promont. Bonae Spei, Dom. Bergius, in Americae mer. ins. St. Johannis, Dom. Moritz, St. Vincentii, Mus. Dom. Hope, Cuba, collect. Dom. Chevrolat.

Habitu et statura omnino *Ph. varii*, *niger*, capite thoraceque pernitidis. Antennae capite thoraceque paulo breviores, articulo secundo tertioque aequalibus, 4—10 subobconicis, sensim paululum brevioribus at non crassioribus, omnibus crassitie longioribus, ultimo oblongo, apice emarginato et inferne acuminato, nigrae, articulo primo plerumque toto nigro, interdum (var. b.) subtus testaceo. Caput parvum, thorace multo minus angustiusque, ovatum, utrinque supra oculos punctis nonnullis, inter oculos punctis 4 transversim positis, intermediis magis distantibus, impressum. Thorax basi coleopterorum prope latitudine, latitudine vix brevior, antrorsum fortius angustatus, lateribus subrectis, basi cum angulis posterioribus fortius rotundatus, apice truncatus, angulis anterioribus deflexis, rotundatis, leviter convexus, seriebus dorsalibus punctis 5, anteriore et posteriore paulo magis remotis, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum confertim punctulatum, nigro-pubescent. Elytra thorace parum longiora, densius subtiliter punctata, nigro-pubescentia, minus nitida, macula vel litura media obscure rufa, saepe obsoleta. Abdomen subtiliter punctatum, nigro-pubescent, subtus segmentis singulis interdum rufo-piceo-marginatis. Pedes nigri, coxis anticis raro totis nigris, plerumque vel interne vel totis testaceis, femoribus anticis etiam interdum basi testaceis, coxis intermediis approximatis, tibiis omnibus spinulosis. Alae albo-hyalinae.

Tarsi antici maris fortiter, feminae leviter dilatati.

Obs. In Museo Hafniensi (coll. Sehestedt), a Fabricio citato, nom. *St. variantis* plures huius generis species (ebeninum, sanguinolentum, varium, ventralem etc. confusas vidi. Patet autem *Staph. variantem* Paykuli, Fabricii ipsius et Olivieri, cum e reliqua descriptione, tum praecipue ex alis albis memoratis, ad hanc speciem pertinere, nomen igitur restitui, huic speciei magnopere variabili egregie adaptatum. Varietates indicatae, primo intuitu saepius nimium differentes, accuratius examinatae et plurimis specimenibus comparatis, certo limite nullo distinctae videntur. *Phil. fuscicornis* Nordm. Capensis ab indigenis non magis discedit, quam specimina plura e variis Americae meridionalis insulis allata.

\* 71. *Ph. flavolimbatu*s: *Niger*, *nitidus*, *elytrorum limbo laterali et apicali, antennarum basi pedibusque flavis, capite ovato*. — Long. 2 lin.

Habitat in Columbia, Dom. Moritz.

Habitu omnino *Ph. variantis*, at minimis eius individuis minor, *niger*, *nitidus*. Antennae capite thoraceque multo breviores, articulo secundo tertioque aequalibus, sequentibus sensim paulo brevioribus at vix crassioribus, omnibus crassitie longioribus, ultimo ovato, apice subemarginato et inferne subacuminato, fuscae, articulis duobus primis testaceis. Caput thorace angustius, utrinque supra oculos punctis paucis, inter oculos punctis 4 transversim positis, intermediis remotioribus, impressum. Palpi nigro-picei. Thorax latitudine haud longior, basi coleopteris paulo angustior, antrorsum angustatus, lateribus subrectis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, posterioribus obtusis, transversim convexus, seriebus dorsalibus punctis 5, antico et postico paulo remotioribus, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum obsolete punctulatum. Elytra thoracis longitudine, sat crebre punctata, longius griseo-pubescentia, nigro-subvirescentia, margine laterali inflexo limboque apicali, interne dilatato, sutu-

ram autem haud attingente, flavis. Abdomen crebre subtiliter punctatum, longius griseo-pubescens, subtus segmentis singulis margine postico testaceo. Pedes testacei, coxis anterioribus concoloribus, intermediis approximatis, tibiis omnibus spinulosis.

Tarsi antici maris fortius, feminae leviter dilatati. Abdominis segmentum inferius sextum feminae apice emarginatum.

\* 72. *Ph. infirmus*: *Niger, nitidus, antennis, elytris pedibusque fuscescenti-testaceis, capite minuto, ovato.* — Long.  $2\frac{1}{2}$  lin.

Habitat in Arabia felice; Dom. Prof. Ehrenberg.

Statura omnino variantis et minimis eius individuis aequalis, nigro-piceus, capite thoraceque nitidissimis, antennis, elytris et pedibus cum coxis anterioribus dilute fuscescenti-rufo-testaceis. Antennae capite cum thorace multo breviores, articulo secundo tertioque aequalibus, 4 — 10 crassiusculis, subobconicis, sensim paulo brevioribus, penultimis crassitie prope brevioribus, ultimo ovato, apice leviter emarginato et inferne acuminato. Mandibulae et palpi picei. Caput thorace multo angustius, ovatum, utrinque supra oculos punctis paucis, inter oculos punctis 4 transversim positis, intermediis remotioribus, notatum. Thorax basi coleopteris paulo angustior, latitudine parum brevior, antrorsum angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, convexus, seriebus dorsalibus punctis 5, antico posticoque paululum remotioribus, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus minus subtilibus. Scutellum nigricans, punctulatum, griseo-pubescens. Elytra thorace paulo longiora, crebre subtilius punctata, densius griseo-pubescentia. Abdomen apicem versus attenuatum, crebre subtiliter punctulatum, pube brevi depressa fuscescenti-grisea dense vestitum, segmentorum marginibus supra anguste, subtus late rufo-piceis. Pedes coxis intermediis leviter distantibus, tibiis omnibus spinulosis.

Tarsi antici feminae simplices. Mas latet.

\* 73. *Ph. debilis*: *Niger, nitidus, elytris pedibusque fuscis, capite suborbiculato.* — Long. 2 —  $2\frac{1}{2}$  lin.

*Er. Col. March. I.* 467. 31.

*Staph. debilis* Grav. *Micr.* 35. 52. — *Latr. Hist. nat. Crust. et Ins. IX.* 329. 57.

*Staph. lucidus* Grav. *Micr.* 21. 24. — *Latr. Hist. nat. Crust. et Ins. IX.* 311. 30.

*Staph. fimetarii* var. Grav. *Mon. p.* 82.

*Staph. agilis* Boisd. et Lacord. *Faun. Ent. Paris. I.* 407. 42.

Habitat in Germania, Gallia; in Aegypto, Dom. Prof. Ehrenberg, in Carolina, Dom. Zimmermann.

*Ph. variantis* var. d similis, at alius, habitu potius *Ph. ebenini*, cui magnitudine semper inferior est, niger, capite thoraceque pernitidis. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, 4 — 10 crassiusculis, penultimis crassitie paulo brevioribus, ultimo ovato, apice truncato et inferne acuminato, nigro-fuscae, basi saepius piceae. Mandibulae rufo-piceae. Palpi picei. Caput thorace paulo minus et angustius, suborbiculatum, supra oculos utrinque punctis paucis, inter oculos punctis



4 transversim positis, intermediis late distantibus, impressum. Thorax coleopteris paulo angustior, latitudine haud brevior, antrorsum subangustatus, lateribus subrectis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, posterioribus obtusis, leviter convexus, seriebus dorsalibus punctis 5, aequaliter distantibus, compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis omnibus subtilioribus. Scutellum subtiliter punctatum. Elytra thorace paulo longiora, densius subtiliusque punctata, fusca, minus subtiliter griseo-pubescentia. Abdomen subtiliter punctulatum, pube longiore depressa grisea densius vestitum, subtus segmentorum marginibus rufo-piceis. Pedes dilute fusci, coxis intermediis leviter distantibus, tibiis anticis breviter, posterioribus fortius spinulosis. Alae albo-hyalinae.

Tarsi antici maris fortius, feminae modice dilatati.

Variat thoracis serierum dorsalium altera 4-punctata.

Obs. A Ph. variantis var. d praecipue distinguitur capite minus ovato, abdomine confertius subtiliter punctato, densius griseo-pubescente.

\* 74. *Ph. ventralis*: Nigro-piceus, nitidus, antennarum basi, pedibus abdominisque segmentorum ventralium marginibus testaceis, capite orbiculato. — Long.  $2\frac{1}{2}$  lin.

Nordm. Symb. 93. 60. — Er. Col. March. I. 468. 32.

*Staph. ventralis* Grav. Micr. 174. 29. Mon. 60. 28. — Gyll. Ins. Suec. II. 334. 49. — Sahlb. Ins. Fenn. I. 325. 45. — Mannerh. Brachelytr. 29. 63. — Boisd. et Lacord. Faun. Ent. Paris. I. 437. 41.

*Staph. immundus* Grav. Mon. 66. 36.

*Staph. anthrax* Grav. Micr. 176. 34. Mon. 67. 38.

*Philonth. anthrax* Nordm. Symb. 97. 74.

Habitat in Europa; in America septentrionali, Knoch., in Amer. mer. ins. St. Vincentii, Mus. Dom. Hope.

Statura omnino Ph. debilis, nigro-piceus, thorace plerumque dilutiore, fusco-piceo, elytris nigricantibus, capite thoraceque pernitidis. Antennae capite thoraceque breviores, articulo tertio secundo paulo longiore, 4 — 10 crassiusculis, sensim paululum brevioribus, non autem crassioribus, penultimis crassitie paulo brevioribus, ultimo ovato, apice leviter emarginato et inferne acuminato, fusco-nigrae, articulo primo testaceo. Palpi fusco-testacei. Caput thorace parum minus et angustius, orbiculatum, supra oculos utrinque punctis parvis, inter oculos punctis 4 transversim positis, intermediis remotioribus, impressum. Thorax coleopteris paulo angustior, latitudine vix longior, lateribus parallelis, subrectis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, posterioribus obtusis, leviter convexus, seriebus dorsalibus punctis 5, antico et postico remotioribus, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus sat magnis profundisque. Scutellum subtiliter punctulatum. Elytra thorace vix longiora, minus dense minusque subtiliter punctata, griseo-pubescentia. Abdomen parce subtiliterque punctulatum, longius griseo-pubescentes, subtus segmentorum marginibus late rufo-testaceis. Pedes rufo-testacei, coxis anterioribus concoloribus, intermediis leviter distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus. Alae albo-hyalinae.

- \* 75. *Ph. discoideus*: Niger, nitidus, antennis, pedibus elytrorumque limbo testaceis, capite orbiculato. — Long. 2 — 2½ lin.

Nordm. Symb. 97. 75. — Er. Col. March. I. 469. 33.

*Staph. discoideus* Grav. Micr. 38. 56. Mon. 58. 24. — Latr. Hist. nat. Crust. et Ins. IX. 322. 61. — Gyll. Ins. Suec. II. 331. 47. — Sahlb. Ins. Fenn. I. 324. 43. — Mannerh. Brachelytr. 29. 61. — Germ. Faun. Ins. Europ. XIII. 4. — Runde Brachelytr. Hal. 8. 38.

*Staph. conformis* Boisd. et Lacord. Faun. Ent. Paris. I. 398. 23.

*Staph. testaceus* Grav. Mon. 60. 27. (forte)

Habitat in Europa; in Cuba, collect. Dom. Chevrolat.

Statura fere praecedentis, niger, capite thoraceque pernitidis. Antennae capite thoraceque multo breviores, tenuiores, articulo secundo tertioque aequalibus, 4 — 10 sensim paululum brevioribus, non autem crassioribus, penultimis crassitie haud brevioribus, ultimo ovato, apice leviter emarginato et inferne subacuminato, totae rufo-testaceae. Os fusco-testaceum. Caput thoracis fere latitudine et magnitudine, suborbiculatum, vel in maribus interdum subquadratum, supra oculos utrinque punctis parvis profundis, inter oculos punctis 4 transversim positis, intermediis magis distantibus, impressum. Thorax coleopterorum fere latitudine, latitudine vix brevior, lateribus parallelis, subrectis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, leviter convexus, seriebus dorsalibus punctis 5, antico posticoque remotioribus, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis ordinariis maioribus profundioribusque. Scutellum crebre punctulatum, griseo-pubescent. Elytra thoracis longitudine, densius fortiusque punctata, minus subtiliter griseo-pubescentia, nigro-fusca, marginibus, imprimis margine laterali et suturali testaceis. Abdomen subtiliter parcius punctulatum, densius griseo-pubescent, segmentorum marginibus supra piceis, subtus testaceis. Pedes fusco-testacei, coxis intermediis leviter distantibus, tibiis omnibus spinulosis.

Tarsi antici maris fortiter, feminae leviter dilatati.

Variat thoracis serie dorsali altera 4-punctata.

- \* 76. *Ph. palliatus*: Niger, nitidus, antennarum basi, elytris pedibusque rufo-testaceis, coxis nigris, capite orbiculato. — Long. 2 lin.

*Staph. palliatus* Grav. Mon. 53. 16.

Habitat in America septentrionali.

Niger, capite thoraceque nitidissimis, elytris testaceo-rufis. Antennae articulis duobus basalibus testaceis, reliquis fusciscentibus, tertio secundo paulo longiore. Mandibulae et palpi rufo-picei. Caput thorace parum angustius, orbiculatum, utrinque supra oculos parce punctatum, inter oculos punctis 4 transversim positis, intermediis remotioribus. Thorax coleopteris vix angustior, latitudine paulo longior, antrorsum haud angustatus, lateribus leviter, basi fortius rotundatus, apice truncatus, angulis anterioribus deflexis posterioribusque subrotundatis, convexus, seriebus dorsalibus punctis 5 sat distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus profundioribus. Scutellum nigrum, parce punctulatum, tenuiter griseo-pubescent. Elytra thoracis longitudine, parcius punctata, parce griseo-pubescentia. Abdomen crassiusculum, parcius punctatum,

griseo-puberulum, segmentis ventralibus piceo-marginatis. Pedes testacei, coxis omnibus nigris.

Tarsi antici feminae modice dilatati. Mas latet

\*77. *Ph. rubidus*: *Niger, antennarum basi, pedibus elytrisque rufis, his basi summa nigricantibus, capite suborbiculato.* — Long. 3 lin.

*Staph. dimidiatus* Boisd. et Lacord. Faun. Ent. Paris. I. 402. 31.

*Philonth. dimidiatus* Er. Col. March. I. 467. 30.

Habitat in Germania, Gallia.

Statura omnino *Ph. ebenini*, at minimis eius individuis vix magnitudine aequalis, niger, capite thoraceque pernitidis. Antennae capitis thoracisque prope longitudine, articulo tertio secundo paulo longiore, 4—10 oblongo-subobconicis, sensim paulo brevioribus, non autem crassioribus, omnibus crassitie longioribus, ultimo oblongo, apice obsolete emarginato et subacuminato, fuscae, articulo primo toto, secundo basi rufis. Palpi rufi. Caput thorace paulo minus et angustius, suborbiculatum, utrinque supra oculos punctis pluribus, antice inter oculos punctis binis oblique positis impressum. Thorax coleopteris paulo angustior, latitudine vix longior, antrorsum vix angustatus, lateribus et basi leviter rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, parum convexus, seriebus dorsalibus punctis 5, primo paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 vel 6 impressus, punctis omnibus minus subtilibus. Scutellum nigrum, dense subtiliter punctulatum, griseo-pubescentia. Elytra thorace paulo longiora, densius subtiliter punctata, griseo-pubescentia, laete rufa, basi summa nigricante. Abdomen densius subtiliter punctatum, pube brevi grisea densius vestitum. Pedes rufo-testacei, coxis concoloribus, intermediis approximatis, tibiis omnibus spinulosis, fuscescentibus.

Tarsi antici maris fortiter dilatati, feminae simplices.

Obs. Ob speciem huius generis a Sahlbergio prius descriptam nomen mutavi.

X \*78. *Ph. quisquiliarius*: *Elongatus, niger, nitidus, elytris virescentibus, antennarum basi pedibusque flavis, capite suborbiculato.* — Long.  $3\frac{1}{2}$  lin.

*Nordm. Symb.* 93. 61. — *Er. Col. March.* I. 469. 31.

*Staph. quisquiliarius* Gyll. Ins. Suec. II. 335. 50. — *Sahlb. Ins. Fenn.* I. 325.

46. — *Mannerh. Brachelytr.* 29. 64. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 406.

39. — *Runde Brachelytr. Hal.* 8. 39.

Habitat in Europa; in Syria et Aegypto etiam a Prof. Ehrenberg lectus.

*Ph. ebenino* angustior et minimis eius individuis aequalis, elongatus, niger, capite thoraceque pernitidis. Antennae capitis thoracisque longitudine, articulo tertio secundo paulo longiore, 4—10 subobconicis, sensim paulo brevioribus et crassioribus, omnibus crassitie longioribus, ultimo ovato, apice emarginato et inferne subacuminato, nigrae, articulo primo testaceo, interdum dorso nigricante. Palpi piceo-testacei, apice nigricantes. Caput thorace paulo minus angustiusque, suborbiculatum, supra oculos utrinque parce profundeque punctatum, inter oculos punctis 4 transversim positis, intermediis fortius distantibus, notatum. Thorax coleopteris paulo angustior,



latitudine haud brevior, antrorsum subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, parum convexus, seriebus dorsalibus punctis 5, primo paululum remotiore, compositis lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus profundioribus. Scutellum dense punctulatum, griseo-pubescentia. Elytra thorace paulo longiora, confertim subtilius punctata, longius griseo-pubescentia, nigro-virescentia, nitidula. Abdomen densius subtiliter punctulatum, tenuiter griseo-pubescentia. Pedes flavi, coxis anticis nigro-piceis, femoribus posticis fusciscentibus, coxis intermediis approximatis, tibiis omnibus spinulosis. Alae albo-hyalinae.

Tarsi antici maris fortiter dilatati, feminae simplices.

- \* 79. *Ph. alcyoneus*: *Elongatus, niger, nitidus, capite thoraceque nigro-subaeneis, capite suborbiculato.* — Long. 3 lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura elongata omnino *Ph. quisquiliarii*, niger, nitidus, capite thoraceque supra nigro-subaeneis, pernitidis. Antennae capitis thoracisque prope longitudine, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo brevioribus et crassioribus, penultimis crassitie haud brevioribus, ultimo ovato, apice leviter emarginato et inferne subacuminato, nigrae. Palpi nigri. Caput feminae thoracis latitudine, suborbiculatum, maris thorace paulo latius, orbiculato-subquadratum, supra oculos utrinque parce fortiter profundeque punctatum, inter oculos punctis utrinque duobus oblique positis notatum. Thorax coleopteris paulo angustior, latitudine paulo longior, basin versus subangustatus, lateribus anterioribus subrotundatis, basi rotundatis, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, posterioribus obtusis, leviter convexus, seriebus dorsalibus punctis 5, aequaliter distantibus, compositis, lateribus punctis utrinque 5 impressus, punctis omnibus sat magnis profundisque. Scutellum punctulatum, longius albedo-pubescentia. Elytra thorace vix longiora, minus dense punctata, viridi-coerulea, nitida, tenuiter albedo pubescentia. Abdomen parcius punctatum, pube pallide grisea certo situ incanum. Pedes nigri, coxis intermediis approximatis, tibiis omnibus spinulosis, tarsis anticis in utroque sexu leviter dilatatis. Alae hyalinae.

- \* 80. *Ph. rufimanus*: *Elongatus, niger, nitidus, elytris cyaneis, coxis femoribusque anterioribus rufo-testaceis, capite suborbiculato.* — Long. 3 lin.

Habitat in Carinthia, Dom. Schmidt.

Statura elongata *Ph. quisquiliarii*, niger, nitidus, capite thoraceque pernitidis. Antennae capitis thoracisque prope longitudine, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo brevioribus et crassioribus, omnibus crassitie longioribus, ultimo ovato, apice emarginato et inferne subacuminato, totae nigrae. Palpi nigri. Caput thoracis latitudine, suborbiculatum, leviter convexum, posterius punctis parvis, inter oculos utrinque punctis duobus, oblique positis, fortioribus impressum. Thorax coleopteris dimidio angustior, latitudine paulo longior, lateribus parallelis, subrectis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, leviter convexus, seriebus dorsalibus

punctis 5, apicali paululum remotiore, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus fortioribus. Scutellum nigrum, punctulatum, tenuiter albido-pubescent. Elytra thorace paulo longiora, minus dense subtilius punctata, laete virescenti-cyanea, tenuiter albido-pubescentia. Abdomen parcius subtiliter punctatum, cinereo-pubescent. Pedes nigri, coxis femoribusque anterioribus laete rufo-testaceis, coxis intermediis approximatis, tibiis omnibus spinulosis.

Tarsi antici maris fortiter, feminae leviter dilatati.

\* 81. *Ph. laetus*: *Niger, nitidus, thorace abdominisque basi rufis, antennis apice pedibusque albidis, femoribus tibiisque apice nigris.* — Long.  $3\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz, in Brasilia, Dom. Prof. Germar.

Statura gracilis fere et colores Paederi. Antennae capite thoraceque parum breviores, crassiusculae, articulo tertio secundo sesqui prope longiore, 4 — 10 cylindricis, sensim paulo brevioribus et crassioribus, ultimo subovato, apice minus abrupte acuminato, nigrae, basi piceae, apice late albido-luteae. Palpi nigro-picei. Mandibulae tenues, falcatae, rufo-piceae. Caput thorace latius, orbiculatum, leviter convexum, nigrum, nitidum, supra oculos utrinque punctis parvis, inter oculos punctis 4 transversim positis, intermediis late distantibus, impressum, fronte apice sulcata. Thorax coleopteris duplo angustior, latitudine plus sesqui longior, basin versus subangustatus, lateribus ante medium leviter rotundatis, pone medium subsinuatis, basi rotundatus, apice utrinque oblique truncatus, angulis anterioribus rotundatis, posterioribus obtusis, convexus, seriebus dorsalibus punctis 5, apicali paululum remotiore minoreque, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, omnibus fortibus profundisque, laete rufus, nitidus. Scutellum dense punctulatum, nigrum. Elytra thorace vix longiora, fortiter punctata, tenuiter nigro-pubescentia, nigra, nitida. Abdomen segmentis singulis basi fortius punctulatis, apice laevibus, nitidis, primis tribus obscure rufis, reliquis nigris, nigro-pilosellum. Pectus nigrum. Pedes elongati, graciles, coxis anterioribus testaceis, intermediis distantibus, femoribus albidis, apice nigris, tibiis anticis muticis, nigris, basi piceis, posterioribus spinulosis, albidis, apice nigris, tarsis testaceis, anticis in utroque sexu simplicibus.

82. *Ph. angularius*: *Rufus, capite thoraceque nigro-aeneis, nitidissimis, abdominis postico nigro, coleopterorum angulis posterioribus femoribusque piceis.* — Long. 3 lin.

Habitat in Mexico, collect. Dom. Chevrolat.

Statura fere praecedentis. Antennae capitis thoracisque prope longitudine, articulo tertio secundo paulo longiore, 4 — 10 subcylindricis, crassiusculis, penultimo crassitie haud longiore, ultimo apice oblique truncato, nigrae, apice ferrugineae. Palpi elongati, picei. Mandibulae elongatae, grales, rufo-piceae. Caput thorace latius, suborbiculatum, posterius utrinque parce punctatum, inter oculos punctis 4 transversim positis, intermediis magis distantibus impressum, fronte anterieus sulcata, nigro-aeneum, nitidissimum. Thorax coleopteris dimidio angustior, latitudine paulo longior, basin versus leviter angustatus, lateribus posterius subsinuatis, basi rotundatus,

apice truncatus, angulis anterioribus minus deflexis, posterioribus subrotundatis, modice convexus, seriebus dorsalibus punctis 5 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque circiter 7 impressus, punctis omnibus sat fortibus profundisque, nigro-aeneus, nitidissimus. Scutellum nigro-aeneum, punctatissimum. Elytra thorace paulo longiora, minus dense sat fortiter punctata, tenuiter griseo-pubescentia, rufa, nitida, angulo posteriore exteriore nigricante. Abdomen parce punctatum, longius pubescens, nitidum, segmentis 3 anterioribus rufis, posterioribus nigris, penultimo margine postico late testaceo. Pectus rufum. Pedes coxis anterioribus nigris, posticis rufis, femoribus piceis, tibiis albidis, apice tarsisque rufo-testaceis, coxis intermediis distantibus, tibiis anticis subtiliter, posterioribus longius parce spinulosis.

Tarsi antici feminae simplices. Mas latet.

Obs. Habitus fere Belonuchi formosi, et femora antica subtus subtilissime spinulosi, at femora postica omnino mutica.

\* 83. *Ph. vernalis*: *Elongatus, niger, nitidus, antennarum basi pedibusque testaceis, capite ovato.* — Long.  $2\frac{2}{3}$  lin.

Nordm. Symb. 93. 59. — Er. Col. March. I. 470. 35.

*Staph. vernalis* Grav. Mon. 75. 67. — Gyll. Ins. Suec. II. 332. 48. — Sahlb. Ins. Fenn. I. 324. 44. — Mannerh. Brachelytr. 29. 62. — Boisd. et Lacord. Faun. Ent. Paris. I. 404. 36.

Habitat in Europa, sub lapidibus et foliis deciduis.

Corpus elongatum, nigrum, nitidum, capite thoraceque pernitidis. Antennae capite thoraceque breviores, articulo tertio secundo paulo longiore, sequentibus subobconicis, sensim paulo brevioribus crassioribusque, penultimis crassitie vix brevioribus, ultimo apice subemarginato et inferne subacuminato, nigrae, articulo primo testaceo. Oris partes testaceae. Caput thorace minus et angustius, oblongo-ovatum, supra oculos utrinque punctis parvis, inter oculos utrinque punctis binis approximatis oblique positis impressum. Thorax coleopterorum fere latitudine, latitudine paulo longior, antrorsum subangustatus, lateribus parum, basi modice rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, modice convexus, seriebus dorsalibus punctis 5 sat distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum punctulatum, griseo-pubescentia. Elytra thorace paulo breviora, dense fortius punctata, minus subtiliter griseo-pubescentia, nigro-picea. Abdomen subtiliter punctulatum, griseo-pubescentia. Pedes testacei, coxis anterioribus concoloribus, intermediis approximatis, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

Obs. Virus olet.

\* 84. *Ph. splendidulus*: *Elongatus, piceus, nitidus, antennis, ore pedibusque testaceis, thorace oblongo, capite ovato.* — Long. 2 —  $2\frac{1}{2}$  lin.

Er. Col. March. I. 470. 36.

*Staph. splendidulus* Grav. Micr. 41. 60. Mon. 93. 89. — Latr. Hist. nat. Crust. et Ins. IX. 327. 78. — Gyll. Ins. Suec. II. 348. 64. — Sahlb. Ins. Fenn. I. 330. 61. *Cafius splendidulus* Mannerh. Brachelytr. 32. 3.



*Gyrophyn. splendidulus* Nordm. *Symb.* 126. 30.

*Staph. nanus* Grav. *Mon.* 96. 93. — *Gyll. Ins. Suec. II.* 347. 63. — *Sahlb. Ins. Fenn. I.* 330. 60. — *Zetterst. Faun. Lappon. I.* 79. 30. *Ins. Lappon.* 65. 40.

*Cafius nanus* Mannerh. *Brachelytr.* 32. 2. — *Runde Brachelytr. Hal.* 10. 1.

*Gyrophyn. nanus* Nordm. *Symb.* 126. 29.

*Staph. parvulus* Grav. *Micr.* 42. 63. *Mon.* 95. 91. — *Latr. Hist. nat. Crust. et Ins. IX.* 328. 81.

Habitat in Europae truncis putridis.

Parvus, elongatus, linearis, iam nigro-, iam testaceo-piceus, nitidus. Antennae capite sesqui longiores, crassiusculae, articulo tertio secundo vix longiore, 4—10 sensim paulo brevioribus at vix crassioribus, 6—10 transversis, ultimo subovato, apice subacummato et prope acumen leviter impresso, totae testaceae. Palpi testacei, articulo ultimo fortiter acuminato. Caput thoracis vix longitudine, ovatum, supra oculos utrinque parce punctatum, inter oculos punctis utrinque duobus transversim positis notatum, fronte inter antennis obsolete longitudinaliter impressa, ore testaceo. Thorax coleopteris paulo angustior, latitudine sesqui prope longior, basin versus subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, leviter convexus, seriebus dorsalibus punctis 5 sat distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilibus. Scutellum subtiliter punctulatum. Elytra thoracis longitudine, minus dense fortius punctata, tenuiter griseo-pubescentia, corpore dilutiora. Abdomen subtiliter punctulatum, tenuiter pubescens, segmentis margine dilutioribus. Pedes testacei, coxis anterioribus concoloribus, intermediis approximatis, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

Obs. Xantholinum simulat, corpore etiam spirae instar involuto quiescens. *Staph. splendidulus*, *nanus* et *parvulus* Grav., speciminibus ab auctore descriptis ipsis inter se comparatis, specie distingui nequeunt, neque *St. splendidum* et *nanum* Gyll., exemplis ab auctore ipso missis et a Dom. Schüppel mihi traditis, examinatis, nisi colore immaturo saturove differunt. — A Ph. aterrimo facilius distinguitur corpore magis elongato linearique, thoracis seriebus dorsalibus 5-punctatis, antennis totis testaceis etc.

### Species mihi invisae.

85. *Ph. immundus*: *Niger, nitidus, antennarum basi, pedibus segmentorumque ventralium marginibus rufo-testaceis, capite subovato.*

*Nordm. Symb.* 93. 64.

*Staph. immundus* Gyll. *Ins. Suec. II.* 337. 53. — *Sahlb. Ins. Fenn. I.* 326. 49.

Habitat in Suecia.

Statura et fere magnitudo Ph. ebenini (maioris), Ph. ventrali colore similis, sed plus duplo maior, caput magis ovatum, etc. Caput subovatum, vel latitudine paulo longius, thorace vix angustius, nigrum, nitidissimum, laeve, punctis paucis in angulis verticis et inter oculos, ut in prioribus; os et palpi picei. Antennae longitudine thoracis, filiformes, nigro-fuscae, articulo primo toto ferrugineo. Thorax antice truncatus, angulis deflexis, compressis, lateribus et postice rotundatus, supra convexus, niger, nitidissimus, laevis, punctis eodem ordine ac numero ut in prioribus. Elytra thorace paulo latiora, sed vix longiora, apice truncata, supra subdepressa, nigra, vix vel parum

virescentia, crebre rugoso-punctata, pubescentia, margine laterali piceo. Alae amplae, albo-hyalinae. Abdomen pone elytra longitudine reliqui corporis, nigrum, violaceo-micans, punctulatum, pubescens. Corpus subtus nigro-piceum, punctulatum, segmentorum ventralium marginibus anoque late rufo-ferrugineis; anus maris late emarginatus. Pedes elongati, validi, rufo-ferruginei, tarsi antici dilatati. Masculos tantum vidi. (Gyll.)

Obs. Affinis videtur *Ph. umbrino*. A *Com. Dejean Ph. fumario* subiungitur, huic autem thoracis series dorsales 6-punctatae et alae fortiter infuscae.

86. *Ph. dimidiatus*: *Niger, nitidus, elytris postice rufo-testaceis, pedibus totis nigro-piceis, capite thorace paulo angustiore, orbiculato, antennis extrorsum crassioribus, thorace brevioribus.*

*Nordm. Symb.* 97. 77.

*Staph. dimidiatus* *Sahlb. Ins. Fenn. I.* 326. 51. — *Mannerh. Brachelytr.* 30. 69.

Habitat in Fennia meridionali.

Magnitudo et statura *Ph. sanguinolenti*, colore coxarum et antennis brevioribus ab illo imprimis distinctus. Caput latitudine fere thoracis, orbiculatum, nigrum, nitidum, laeve, punctis in angulo verticis et quatuor inter oculos ut in praecedentibus impressis. Antennae thorace breviores, extrorsum paulo crassiores, articulis tribus baseos nigris, nitidis, longioribus, ceteris fusco-pubescentibus, brevibus, longitudine latioribus, ultimo longiore. Thorax antice truncatus, lateribus vix dilatatus, postice rotundatus, supra convexus, niger, nitidus, inter angulum anticum puncta duo, dein iterum duo, et in seriebus dorsalibus 5 sat profunde impressa, aequali spatio a se invicem remota. Scutellum triangulare, acuminatum, depressum, nigrum, crebre punctatum. Elytra thorace nonnihil latiora, vix illo longiora, apice oblique truncata, supra depressa, sat crebre et profunde punctata, tenue pubescentia, nitida, nigra, macula magna postica, dimidium elytri occupante, suturam non attingente, sed iuxta marginem exteriorem parum adscendente, rufo-testacea. Abdomen pone elytra longitudine fere reliqui corporis, nigrum, subopacum, pubescens, subtiliter punctulatum. Corpus subtus nigrum, subopacum, subtiliter punctulatum. Pedes elongati, nigro-picei, tarsi paulo dilutioribus. (*Sahlb.*)

87. *Ph. celer*: *Niger, nitidus, antennis fuscis, pedibus rufo-testaceis, segmentorum marginibus ventre rufis, thorace seriebus dorsalibus 5-punctis, capite subovato. — Long. 3 lin.*

*Nordm. Symb.* 94. 68.

*Staph. celer* *Grav. Mon.* 66. 37.

Habitat Lutetiae.

Caput subovatum, thorace paulo angustius, nigrum, palpis antennisque fuscis, his capite duplo longioribus. Thorax coleopteris paululum minor, nigricans, 5, 3, 1 punctus, punctis distinctis. Coleoptera subdepilia, piceo-nigra. Pedes tarsi tenuibus. (*Grav.*)

Obs. *Ph. ventrali* affinis videtur, species mihi perquam dubia.

88. *Ph. irregularis*: Niger, nitidus, elytris fusco virescentibus, pedibus fusco-piceis, capite thorace maiore, subquadrato, thoracis seriebus dorsalibus irregularibus.

Nordm. Symb. 93. 66.

*Staph. irregularis* Mannerh. *Brachelytr.* 30. 74.

Habitat Petropoli.

Ph. ebenino colore similis, capite autem maiore, subquadrato et elytris minus crebre punctatis. In thorace etiam prope seriem dorsalem extus punctum maiusculum utrinque impressum, unde series minus regulares. (Mannerh.)

Obs. An a Ph. ebenino revera distinctus, an forte varietas fortuita?

6) Thorax seriebus dorsalibus 6-punctatis.

- \* 89. *Ph. nigrita*: Nigerrimus, thorace griseo-subsericante, elytris subtiliter confertissime punctatis, capite ovato. — Long. 3 lin.

Nordm. Symb. 101. 86. — Er. Col. March. I. 471. 37.

*Staph. nigrita* Grav. Mon. 68. 44.

Habitat in Germania, Suecia.

Habitu omnino Ph. variantis et maximis eius individuis aequalis, nigerimus, capite thoraceque pernitidis, hoc certo situ griseo-subsericante. Antennae capite thoraceque longiores, graciliores, articulo tertio secundo paulo longiore, sequentibus omnibus elongato-obconicis, 4 — 10 sensim paulo brevioribus, ultimo praecedente paulo longiore, apice truncato et abrupte subacuminato, nigrae, articulo ultimo brunneo. Palpi nigri. Caput thorace multo minus et angustius, ovatum, supra oculos utrinque parce punctatum, inter oculos utrinque punctis duobus approximatis oblique positis impressum. Thorax basi coleopterorum prope latitudine, latitudine baseos vix longior, antrorsum leviter angustatus, lateribus vix, basi leviter rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, modice convexus, seriebus dorsalibus punctis 6, apicali paulo remotiore compositis, lateribus punctis utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum subtiliter punctulatum, subtiliter pubescens. Elytra thorace paulo longiora, confertim subtiliter punctata, tenuiter subtiliterque nigro-pubescentia. Abdomen subtiliter minus confertim punctatum, tenuiter nigro-pubescentia. Pedes nigri, coxis intermediis approximatis, tibiis omnibus spinulosis, geniculis piceis, tarsis rufo-piceis.

Tarsi antici maris modice, feminae leviter dilatati.

Variat thoracis seriebus dorsalibus 4-punctatis.

Obs. E Suecia nonnunquam pro Ph. virgine missus.

- \* 90. *Ph. thoracicus*: Niger, thorace rufo, antennarum basi, pedibus elytrisque testaceis, capite ovali. — Long.  $3\frac{1}{2}$  lin.

*Staph. thoracicus* Grav. Micr. 170. 23. Mon. 71. 57.

Habitat in America septentrionali.

Statura et magnitudo omnino Ph. nigritae. Antennae capitis thoracisque longitudine, articulo tertio secundo paulo longiore, sequentibus subob-



conicis, sensim crassioribus, 4 — 10 sensim paulo brevioribus, penultimis crassitie vix longioribus, ultimo apice truncato et inferne subacuminato, fuscae, articulis 3 primis ultimoque testaceis. Palpi testacei. Mandibulae rufo-piceae. Caput thorace multo angustius, oblongo-ovatum, utrinque supra oculos punctis paucis, inter oculos punctis 4 transversim positis impressum, nigrum, nitidum. Thorax basi coleopterorum prope latitudine, latitudine baseos vix brevior, antrorsum angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, modice convexus, seriebus dorsalibus punctis 6, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilibus, rufus, nitidus. Elytra thorace vix longiora et scutellum crebre subtiliter punctata, fuscescenti-testacea, griseo-pubescentia. Abdomen subtilius punctulatum, griseo-pubescentis, nigrum, nitidulum, segmentorum marginibus rufo-piceis. Pectus nigrum. Pedes cum coxis anterioribus flavo-testacei, coxis intermediis approximatis, tibiis omnibus spinulosus.

Tarsi antici feminae simplices. Mas latet.

X

\* 91. *Ph. fumarius*: *Niger, nitidus, pedibus fulvis, elytris virescentibus, dense subtiliter punctatis, capite ovato.* — Long.  $3\frac{1}{2}$  lin.

*Nordm. Symb.* 100. 84. — *Er. Col. March.* I. 472. 38.

*Staph. fumarius* Grav. *Mon.* 67. 43.

Habitat in Germania.

Habitu *Ph. variantis*, at duplo fere maior, niger, capite thoraceque nitidissimis, hoc certo situ griseo-subsericante. Antennae capite thoraceque longiores, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo brevioribus crassioribusque, omnibus crassitie longioribus, ultimo oblongo, apice subtruncato et inferne subacuminato, nigrae, articulis 3 primis obscure piceis, ultimo brunneo. Palpi piceo-testacei. Caput thorace minus et angustius, ovatum, supra oculos utrinque punctis paucis, inter oculos punctis 4 transversim positis, intermediis magis distantibus, impressum. Thorax basi coleopterorum prope latitudine, latitudine baseos vix brevior, antrorsum leviter angustatus, lateribus parum, basi modice rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, modice convexus, seriebus dorsalibus punctis 6, plerumque aequaliter distantibus, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum nigrum, subtiliter punctulatum, tenuiter pubescens. Elytra thorace paulo longiora, confertim subtiliter punctata, griseo-pubescentia, nigro-virescentia, nitidula. Abdomen densius subtiliter punctulatum, densius griseo-pubescentis, subtus segmentorum marginibus obscure ferrugineis. Pedes fulvi, coxis omnibus nigris, intermediis approximatis, tibiis omnibus spinulosus. Alae fuscae.

Tarsi antici maris modice, feminae levissime dilatati.

92. *Ph. lomatus*: *Niger, nitidus, antennarum articulo primo, palpis, pedibusque testaceis, elytris confertim subtiliter punctatis, viridi-aeneis, margine laterali tenuique apicali obscure testaceis, capite ovato.* — Long.  $3\frac{1}{2}$  lin.

Habitat in America septentrionali, Mus. Dom. Chevrolat.

Statura omnino praecedentium, niger, nitidus. Antennae capite thorace paulo longiores, graciles, articulo tertio secundo paulo longiore, 4 — 10 elongato-subobconicis, sensim paulo brevioribus at vix crassioribus, ultimo oblongo, apice oblique truncato et inferne acuminato, nigrae, articulo primo testaceo, ultimo brunneo. Caput thorace multo angustius, ovatum, utrinque supra oculos punctis paucis, inter oculos punctis utrinque duobus transversim positis impressum. Thorax basi coleopterorum prope latitudine, latitudine baseos vix longior, antrorsum angustatus, lateribus leviter, basi fortius rotundatus, apice truncatus vel subemarginatus, angulis anterioribus leviter deflexis, subrotundatis, posterioribus obtusis, modice convexus, seriebus dorsalibus punctis 6, apicali remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilibus. Scutellum subtiliter punctulatum, griseo-pubescentia. Elytra thorace vix longiora, subtiliter confertim punctata, densius griseo-pubescentia, obscure viridi-aenea, nitidula, margine laterali tenuique apicali saturate testaceo. Abdomen subtiliter punctulatum, longius griseo-pubescentia, segmentorum marginibus rufo-piceis. Pedes testacei, coxis omnibus piceis, intermediis approximatis, tibiis omnibus parce spinulosis.

Tarsi antici maris fortiter dilatati, femina latet.

\* 93. *Ph. virgo*: Niger, nitidus, pedum geniculis tarsisque piceis, elytris subtiliter punctulatis, capite ovato. — Long. 2 $\frac{3}{4}$  lin.

Nordm. Symb. 101. 85.

*Staph. virgo* Grav. Micr. 169. 19. Mon. 69. 45. — Latr. Hist. nat. Crust. et Ins. IX. 323. 65. — Gyll. Ins. Suec. II. 315. 61. — Sahlb. Ins. Fenn. I. 330. 58. — Zetterst. Faun. Lappon. I. 78. 28. Ins. Lappon. 65. 38. — Mannerh. Brachelytr. 31. 77. — Boisd. et Lacord. Faun. Ent. Paris. I. 404. 35.

Habitat in Lusitania, Com. Hoffmannsegg, in Sardinia, Dom. Prof. Gène, in Suecia, Dom. Prof. Zetterstedt.

Statura *Ph. micantis*, at plerumque paulo maior, *Ph. nigrita* multo minor, niger, capite thoraceque nitidis, haud sericeo-micantibus. Antennae capitis thoracisque prope longitudine, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim leviter crassioribus, non autem brevioribus, omnibus crassitie longioribus, ultimo oblongo, apice inferne abrupte subacuminato, nigrae. Palpi nigri. Caput thorace angustius, breviter subovatum, utrinque supra oculos punctis pluribus, inter oculos punctis utrinque duobus transversim positis impressum. Thorax coleopteris dimidio angustior, latitudine vix longior, antrorsum perparum angustatus, lateribus parum, basi modice rotundatus, apice truncatus, angulis anterioribus leviter deflexis, posterioribus subrotundatis, modice transversim convexus, seriebus dorsalibus punctis 6, apicali plerumque paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis minus subtilibus. Elytra thorace paulo longiora et scutellum subtiliter punctulata, minus subtiliter griseo-pubescentia, subnitida. Abdomen densius subtiliter punctulatum, longius densiusque griseo-pubescentia, lateribus nigro-pilosellum, nitidulum, segmento ventrali ultimo apice subtestaceo. Pedes nigro-picei, coxis intermediis approximatis, tibiis omnibus spinulosis, geniculis tarsisque dilutioribus. Alae hyalinae.

Tarsi antici maris fortiter, feminae leviter dilatati.

- \* 94. *Ph. turbidus*: *Niger, nitidus, antennarum basi apiceque, palpis pedibusque fulvis, elytris dense subtiliter punctatis, capite oblongo-ovato.* — *Long. 4 lin.*

Habitat in Madagascar, Dom. Goudot.

Statura omnino *Ph. fumarii*, at paulo maior, niger, nitidus, capite thoraceque pernitidis, hoc certo situ griseo-subsericante. Antennae capitis thoracisque longitudine, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus et crassioribus, primis eorum cylindricis, sequentibus subobconicis, penultimis crassitie paulo longioribus, ultimo oblongo-ovato, apice subemarginato et inferne acuminato, fuscae, articulis tribus primis ultimoque saturate testaceis. Palpi testacei. Caput thorace angustius, oblongo-ovatum, leviter convexum, supra oculos utrinque punctis pluribus dispersis, inter oculos utrinque duobus transversim positis impressum. Thorax coleopteris angustior, latitudine baseos tertia fere parte longior, antrorsum leviter angustatus, lateribus levissime, basi modice rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, modice convexus, seriebus dorsalibus punctis 6 compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum crebre punctatum, tenuiter nigro-pubescent. Elytra thorace tertia parte longiora, dense subtiliter punctata, subtiliter nigro-pubescentia. Abdomen crebrius subtiliter punctatum, tenuiter nigro-pubescent, nitidulum, metallico-subversicolor. Pedes fulvi, coxis anterioribus concoloribus, intermediis approximatis, tibiis omnibus spinulosis.

Maris tarsi antici fortiter dilatati, abdominis segmentum inferius quintum apice emarginatum. Femina latet.

Variat immaturus piceo-testaceus, capite elytrisque nigricantibus.

- \* 95. *Ph. micans*: *Niger, opacus, capite thoraceque nitidis, sericantibus, pedibus testaceis, capite ovato.* — *Long. 2½ lin.*

*Nordm. Symb. 102. 88. — Er. Col. March. I. 473. 40.*

*Staph. micans Grav. Micr. 25. 34. Mon. 76. 69. — Latr. Hist. nat. Crust. et Ins. IX. 314. 39. — Gyll. Ins. Suec. II. 344. 60. — Sahlb. Ins. Fenn. I. 329. 57. — Zetterst. Faun. Lappon. I. 78. 29. Ins. Lappon. 65. 39. — Mannerk. Brachelytr. 31. 76. — Boisd. et Lacord. Faun. Ent. Paris. I. 406. 40.*

*Staph. pullus Runde Brachelytr. Hal. 9. 48.*

*Staph. luridus Runde Brachelytr. Hal. 9. 50.*

*Staph. varians var. γ Payk. Mon. Staph. p. 46. Faun. Suec. III. p. 394.*

Habitat in Europa.

Habitu *Ph. variantis* et minoribus eius individuis aequalis, niger, opacus, capite thoraceque solis nitidis, hoc distinctius sericeo-micante. Antennae capitis thoracisque longitudine, articulo secundo tertio paulo longiore, 4—10 elongato-obconicis, sensim paulo crassioribus at vix brevioribus, ultimo oblongo, apice emarginato et inferne subacuminato, nigrae. Palpi picei. Mandibulae piceae. Caput thorace minus et angustius, subovatum, leviter convexum, supra oculos utrinque punctis pluribus, inter oculos utrinque duobus, transversim positis impressum. Thorax basi coleopteris paulo angustior, latitudine baseos vix longior, antrorsum subangustatus, lateribus levissime, basi modice rotundatus, apice truncatus, angulis anterioribus leviter deflexis



posterioribusque rotundatis, leviter convexus, seriebus dorsalibus punctis 6, apicali saepius paulo remotiore, compositis, lateribus praeter marginalia punctis utrinque 4 impressus, punctis omnibus subtilioribus. Elytra thorace longiora et scutellum subtilissime confertissimeque punctata, pube subtili densa cinerascens. Abdomen confertim subtilissime punctatum, subtiliter dense cinereo-pubescent, subtus segmentorum marginibus late rufo-ferrugineis. Pedes testacei, coxis nigris, intermediis approximatis, tibiis omnibus spinulosis. Alae hyalinae.

Tarsi antici maris fortiter, feminae vix dilatati.

Obs. Staph. pullus et luridus Runde, mihi ad examinandum a Dom. Zschorn benevole communicati, certe ad hanc speciem pertinent, Staph. velox Runde l. c. p. 9. n. 49. quoque eidem subiungendus videtur; Staph. pusillus eiusdem etiam auctoris l. c. p. 10. f. 54. partim ad Phil. micantem partim ad Phil. debilem revocandus est.

\* 96. *Ph. fulvipes*: Niger, antennarum basi, elytris pedibusque laete testaceo-rufis, capite ovato. — Long.  $2\frac{1}{2}$  lin.

Nordm. Symb. 102. 89. — Er. Col. March. I. 473. 41.

*Staph. fulvipes* Fab. Ent. Syst. I. II. 526. 31. Syst. El. II. 597. 40. — Panz. Ent. Germ. 355. 21. Faun. Germ. 29. 9. (male) — Grav. Micr. 24. 33. Mon. 75. 66. — Latr. Hist. nat. Crust. et Ins. IX. 314. 38. — Gyll. Ins. Suec. II. 343. 59. — Sahlb. Ins. Fenn. I. 329. 56. — Mannerh. Brachelytr. 30. 75. — Boisd. et Lacord. Faun. Ent. Paris. I. 402. 30.

*Staph. mendax* Runde Brachelytr. Hal. 10. 51.

*Staph. flavopterus* var. *A* Fourcr. Ent. Par. I. 166. 10. (forte)

*Le Staph. noir à étuis fauves et corselet strié* Geoffr. Hist. Ins. I. p. 365.

Habitat in Europa frequens; in America septentrionali, Mus. Dom. Chevrolat.

Statura *Ph. micantis*, niger, capite thoraceque nitidis, haud sericantibus. Antennae capitis thoracisque longitudine, articulo tertio secundo paululum longiore, 4 — 10 subcylindricis, sensim paulo brevioribus et crassioribus, penultimis crassitie parum longioribus, ultimo oblongo, apice truncato et inferne acuminato, nigrae, articulis tribus primis testaceis. Palpi testacei. Caput thorace minus et angustius, ovatum, supra oculos utrinque punctis parvis, inter oculos utrinque duobus transversim positissimis impressum. Thorax basi coleopteris paulo angustior, latitudine baseos parum longior, antrosum subangustatus, lateribus levissime, basi fortius rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, leviter convexus, seriebus dorsalibus punctis 6, primo parum remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Scutellum confertim punctulatum, nigrum, griseo-pubescent. Elytra thorace paululum longiora, subtiliter confertim punctata, laete testaceo-rufa, subtiliter testaceo-pubescentia. Abdomen confertim subtiliter punctatum, tenuiter griseo-pubescent. Pedes testacei, coxis anterioribus concoloribus, intermediis approximatis, tibiis omnibus subtili spinulosis.

Tarsi antici maris fortius, feminae vix dilatati.

Variat antennarum articulis 3 primis piceis.

- \*97. *Ph. brunneus*: *Piceus, nitidus, antennarum basi, pedibus abdominisque segmentorum marginibus testaceis, thorace rufescenti-piceo.* — Long.  $2\frac{1}{2}$  lin.

*Staph. brunneus* Grav. *Micr.* 172. 26. *Mon.* 57. 22.

*Staph. sericans* Grav. *Micr.* 171. 25. *Mon.* 59. 25.

Habitat in America septentrionali.

Statura omnino *Ph. debilis*. Antennae capite thoraceque paulo breviores, articulo tertio secundo paululum longiore, 4 — 10 sensim paulo brevioribus at vix crassioribus, penultimis crassitie subbrevioribus, ultimo apice truncato et inferne subacuminato, fuscae, basi testaceae, apice ferrugineae. Os obscure testaceum. Caput maris thoracis fere latitudine, orbiculatum, feminae thorace angustius, ovatum, utrinque supra oculos punctis paucis, inter oculos punctis 4, intermediis magis distantibus, notatum, nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine baseos vix longior, antrorsum subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, modice transversim convexus, seriebus dorsalibus punctis 6, apicali paulo remotiore compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus minus subtilibus, plerumque rufo-, rarius nigro-piceus, nitidus. Elytra thorace vix longiora et scutellum crebre subtilius punctata, minus subtiliter griseo-pubescentia, fusco-picea. Abdomen subtiliter punctulatum, densius griseo-pubescentia, piceum, segmentis omnibus rufo-marginalis. Pectus piceum. Pedes cum coxis anterioribus testacei, coxis intermediis approximatis, tibiis omnibus spinulosis.

Tarsi antici maris fortiter, feminae leviter dilatati.

Obs. *Staph. brunneus* et *sericans* Grav. nullo puncto differunt. Thoracem sericeo-micantem, quem Grav. apud *St. sericantem* suum laudat, equidem in eodem specimine observare nequeo. Conf. Grav. *Mon.* 59. 25.

- \*98. *Ph. fimbriolatus*: *Piceus, nitidus, antennarum basi, pedibus elytrorumque margine tenui apicali testaceis, sutura abdominisque segmentorum marginibus ferrugineis, capite orbiculato.* — Long.  $2\frac{1}{4}$  lin.

Habitat in Madagascar, Dom. Goudot.

Praecedenti affinis, at minor, *Ph. aterrimo* parum minor, licet minus gracilis. Antennae capite thoraceque breviores, articulo secundo tertioque aequalibus, 4 — 10 sensim paulo brevioribus, at vix crassioribus, primis eorum subcylindricis, sequentibus subobconicis, penultimis crassitie vix brevioribus, ultimo subovato, apice subemarginato et inferne subacuminato, fuscae, articulis 3 primis testaceis, ultimo ferrugineo. Caput thoracis latitudine, suborbiculatum, utrinque supra oculos punctis pluribus sparsis, inter oculos utrinque duobus transversim positis notatum, fronte anterieus medio subimpressa, nigro-piceum, pernitidum, ore rufo-piceo, palpis testaceis. Thorax coleopteris dimidio prope angustior, latitudine tertia parte longior, lateribus parallelis, rectis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, leviter convexus, seriebus dorsalibus punctis 6, antico remotiore et subtiliore, reliquis sat fortibus, compositis, lateribus punctis praeter marginalia utrinque 5 subtilioribus impressus, piceus, pernitidus. Scutellum nigro-piceum, parce subtiliterque punctatum,

parce pubescens. Elytra thoracis longitudine, minus confertim profundius punctata, parce griseo-pubescentia, fusco-picea, nitida, humeris suturaeque rufescentibus, margine angusto apicali testaseo. Abdomen parce subtiliter punctulatum, parcus longiusque griseo-pubescentis, minus nitidum, nigrum, segmentis omnibus apice ferrugineo-marginatis. Pedes testacei, tibiis omnibus spinulosis.

Tarsi antici maris simplices, femina latet.

\* 99. *Ph. cribricollis*: Nitidus, niger, antennarum basi apiceque oreque testaceis, thorace, elytrorum abdominisque basi rufis, pedibus albidis. — Long.  $3\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz, in Mexico, coll. Dom. Chevrolat.

Statura et affinitas *Ph. laeti*. Antennae capite thoraceque parum breviores, validiores, articulo tertio secundo paulo longiore, 4 — 10 subcylindricis, sensim paulo brevioribus et crassioribus, penultimis crassitie perparum longioribus, ultimo apice inferne abrupte acuminato, brunneae, articulis tribus basalibus duobusque apicalibus testaceis. Mandibulae falcatae, acutae, cum toto ore rufo-testaceae. Caput thorace latius, orbiculatum, nigro-piceum, nitidissimum, utrinque parce grosseque punctatum, inter oculos punctis utrinque duobus transversim positis impressum, fronte antice longitudinaliter fortiter sulcata. Thorax coleopteris duplo angustior, latitudine sesqui longior, lateribus ante medium leviter rotundatis, pone medium subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus fortiter rotundatis, posterioribus obtusis, convexus, seriebus dorsalibus punctis 6, apicali parum remotiore, compositis, lateribus punctis praeter marginalia utrinque 6 impressus, punctis omnibus grossis et profundis, rufus, nitidissimus, metallico-lucidus. Scutellum crebre punctatum, rufum. Elytra thoracis longitudine, confertim fortiter punctata, tenuiter griseo-pubescentia, nigra, basi rufa. Abdomen nitidum, segmentis singulis basi subtilius punctulatis, tribus primis rufis, parce pubescentibus, reliquis nigris, densius nigro-pubescentibus, quinto margine apicali testaceo. Pectus nigrum. Pedes elongati, graciles, albidus, femoribus tibiisque apice fusciscentibus, coxis anterioribus testaceis, posticis rufis, coxis intermediis modice distantibus, tibiis omnibus subtiliter spinulosis, tarsis anticis in utroque sexu simplicibus.

Obs. Nomine *Cafii cribricollis* hanc speciem sibi impositam esse monuit D. Com. Dejean.

100. *Ph. angusticollis*: Niger, nitidus, antennis apice ferrugineis, thorace abdominisque segmentis 3 primis rufis, pedibus albidis, geniculis fuscis. — Long.  $3\frac{1}{2}$  lin.

Habitat in Mexico, collect. Dom. Chevrolat.

Statura et summa affinitas praecedentis. Antennae capite thoraceque parum breviores, validiusculae, articulo tertio secundo paulo longiore, 4 — 10 subcylindricis, sensim paulo brevioribus, at parum crassioribus, penultimo crassitie parum longiore, ultimo apice inferne abrupte acuminato, nigrae, articulo secundo basi testaceo, ultimis duobus ferrugineis. Mandibulae et palpi piceo-testacei. Caput thorace latius, orbiculatum, posterius punctis parcis profundis, inter oculos utrinque duobus transversim positis impressum, fronte an-



tice canaliculata, nigro-subaeneum, nitidissimum. Thorax coleopteris duplo prope angustior, latitudine longior, basin versus subangustatus, lateribus anterieus leviter rotundatis, posterius subsinuatis, basi leviter rotundatus, angulis posterioribus obtusis, apice cum angulis anterioribus rotundatus, modice convexus, seriebus dorsalibus punctis 6, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 6 vel 7 impressus, punctis omnibus sat fortibus profundisque, rufus, nitidus. Scutellum confertim punctatum, obscure rufum. Elytra thorace paulo longiora, minus crebre sat fortiter punctata, tenuiter nigro-pubescentia, nigra, nitida, summa basi obscure rufa. Abdomen segmentis anterioribus tribus sublaevibus, laete rufis, nitidis, posterioribus parcius punctulatis, nigro-pubescentibus, nigris, penultimo apice testaceo. Pectus nigrum. Pedes elongati, graciles, albidii, femoribus apice fuscis, coxis intermediis sat distantibus, tibiis omnibus subtiliter spinulosis, tarsis anticis in utroque sexu simplicibus.

\* 101. *Ph. tenuis*: Nitidus, niger, thorace, pedibus elytrisque rufis, his basi nigris, capite ovato. — Long.  $2\frac{1}{2}$  lin.

Nordm. Symb. 103. 91. — Er. Col. March. I. 474. 42.

*Staph. tenuis* Fab. Ent. Syst. I. II 528. 43. Syst. El. II. 599. 53. — Panz. Ent. Germ. 357. 33. — Grav. Micr. 39. 58. Mon. 72. 59. — Latr. Hist. nat. Crust. et Ins. IX. 322. 63. — Boisd. et Lacord. Faun. Ent. Paris. I. 403. 32. — Runde Brachelytr. Hal. 10. 53.

*Paederus dimidiatus* Panz. Faun. Germ. 27. 24. Ent. Germ. 362. 3.

Habitat in Germania.

Antennae capite thoraceque paulo breviores, articulo secundo tertioque subaequalibus, sequentibus sensim brevioribus et crassioribus, penultimis longitudine crassioribus, ultimo subovato, apice emarginato et inferne acuminato, nigrae, articulis duobus primis obscure rufis. Caput thorace paulo minus et angustius, ovatum, supra oculos utrinque punctis subtilibus pluribus, inter oculos utrinque duobus transversim positis impressum, nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine longior, lateribus parallelis, subrectis, basi cum angulis posterioribus rotundatus, basi truncatus, angulis anterioribus subrotundatis, convexus, seriebus dorsalibus punctis 6, apicali paululum remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilibus, rufus, nitidus. Scutellum confertim punctulatum, nigrum. Elytra thoracis longitudine, subdepressa, densius subtilissime punctata, subtiliter griseo-pubescentia, dimidiatim basi nigra, apice rufa. Abdomen dense subtiliter punctatum, nigrum, subnitidum, subtiliter densius griseo-pubescentis, dorso pilis dispersis erectis nigris instructum. Pectus nigrum. Pedes graciles rufi, coxis anterioribus concoloribus, intermediis approximatis, tibiis omnibus parce subtiliterque spinulosis.

Tarsi antici maris fortiter, feminae levissime dilatati.

\* 102. *Ph. pullus*: Niger, nitidus, elytris nigro-subvirescentibus, thorace oblongo, convexo, capite subovato. — Long.  $2\frac{1}{2}$  lin.

Nordm. Symb. 104. 92. — Er. Col. March. I. 475. 43.

Habitat in Germania, Gallia.

Statura omnino praecedentis, niger, nitidus, capite thoraceque pernitidis. Antennae vix capitis thoracisque longitudine, articulo secundo tertioque subaequalibus, 4—10 subcylindricis, sensim paulo brevioribus crassioribusque, penultimis crassitie paulo brevioribus, ultimo apice subemarginato et inferne acuminato, nigrae. Palpi nigri. Caput thorace paulo minus et angustius, ovatum, pone oculos lateribus utrinque confertim subtiliter punctatum, supra utrinque supra oculos punctis paucis, inter oculos utrinque duobus approximatis transversim positissubtilioribus impressum. Thorax coleopteris dimidio prope angustior, latitudine paulo longior, lateribus parallelis, subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, convexus, seriebus dorsalibus punctis 6, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilibus. Scutellum confertissime subtiliter punctatum, subtiliter griseo-pubescentia. Elytra thoracis longitudine, subtilius minus dense punctata, nigro-subvirescentia, subtus griseo-pubescentia. Abdomen subtiliter punctatum, pube grisea micante densius vestitum, dorso pilis erectis nigris sparsum. Pedes graciles, nigri aut nigro-picei, coxis intermediis approximatis, tibiis omnibus parce spinulosis.

Tarsi antiqui maris fortiter, feminae leviter dilatati.

- \* 103. *Ph. lustralis*: Nigro-subaeneus, nitidus, elytris fusco-aeneis, pedibus flavis, thorace oblongo, apice attenuato, capite subovato. — Long.  $2\frac{1}{2}$  lin.

Habitat in ins. Puerto Rico, Dom. Moritz.

Statura fere praecedentis, nigro-subaeneus, capite thoraceque nitidissimis, elytris fuscis, nitore aeneo perfusus, seu aeneis, basi fusciscentibus. Antennae capite thoraceque parum breviores, articulo tertio secundo longiore, 4—10 subcylindricis, sensim paulo brevioribus et crassioribus, penultimis crassitie vix longioribus, ultimo apice oblique truncato et inferne acuminato, nigrae, articulo primo piceo. Mandibulae et palpi piceo-testacei. Caput thoracis latitudine, subobovatum, supra oculos utrinque sparsim punctatum, anterieus inter oculos punctis 4 subtilibus, intermedium fortius distantibus, impressum. Thorax coleopteris duplo fere angustior, latitudine sesqui longior, lateribus rectis, parallelis, basi cum angulis posterioribus rotundatus, angulis anterioribus deflexis, fortiter rotundatis, convexus, seriebus dorsalibus punctis 6, apicali remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus minus subtilibus. Scutellum subtiliter punctulatum. Elytra thoracis longitudine, dense profundius punctata, minus subtiliter griseo-pubescentia. Abdomen crebrius subtiliusque punctatum, parcius griseo-pubescentia. Pedes flavo-testacei, coxis anterioribus concoloribus, intermediis approximatis, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

Abdominis segmentum inferius sextum maris apice profunde bisinuatum, feminae integrum.

- \* 104. *Ph. trepidus*: Elongatus, niger, nitidus, elytris obscure virescentibus, antennarum basi pedibusque testaceis, thorace oblongo, capite ovato. — Long. 3 lin.

Habitat in Americae ins. St. Johannis et Puerto Rico, Dom. Moritz.

Statura et affinitas *Ph. quisquiliarii*. Niger, capite thoraceque nitidissimis, elytris obscure viridi-aeneis, subnitidis. Antennae capitis thoracisque longitudine, articulo tertio secundo longiore, 4—10 subobconicis, sensim paulo brevioribus et crassioribus, penultimis crassitie haud brevioribus, ultimo oblongo, apice oblique subemarginato et inferne acuminato, nigrae, articulo primo testaceo, interdum medio piceo, ultimo plerumque ferrugineo. Palpi piceo-testacei. Mandibulae piceae. Caput thorace parum angustius, ovatum, utrinque supra oculos, lateribus verticeque crebrius punctatum, inter oculos punctis utrinque duobus transversim positis impressum. Thorax coleopteris dimidio angustior, latitudine plus sesqui longior, lateribus parallelis, subrectis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, leviter convexus, seriebus dorsalibus punctis 6, apicali remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 vel 6 impressus, punctis omnibus subtilioribus. Scutellum nigrum, subtiliter punctulatum, subtiliter nigro-pubescent. Elytra thorace paulo longiora, subtiliter dense punctata, densius subtiliter nigro-pubescentia. Abdomen subtiliter punctulatum, subtiliter fusco-pubescent. Pedes testacei, coxis anterioribus, femoribus posticis, interdum etiam tibiis piceis, coxis intermediis approximatis, tibiis omnibus spinulosis. Alae fuscae.

Tarsi antici maris modice, feminae leviter dilatati.

\* 105. *Ph. alumnus*: Niger, nitidus, antennis basi testaceis, pedibus fusco-testaceis, elytris confertim subtiliter punctatis, opacis, capite subovato. — Long.  $2\frac{1}{2}$  lin.

Habitat in ins. Puerto Rico, Dom. Moritz.

*Ph. micanti* affinis, at paulo gracilior, niger, nitidus, capite thoraceque haud sericeo-micantibus. Antennae capitis thoracisque longitudine, articulo tertio secundo paulo longiore, 4—10 subobconicis, sensim parum brevioribus et crassioribus, omnibus crassitie longioribus, ultimo oblongo, apice oblique truncato et inferne acuminato, nigrae, articulis duobus basalibus testaceis, supra infuscatis. Palpi piceo-testacei. Caput thorace paulo angustius, breviter subovatum, utrinque supra oculos parcius punctatum, inter oculos punctis utrinque duobus transversim positis impressum. Thorax coleopteris dimidio angustior, latitudine paulo longior, lateribus parallelis, subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, modice convexus, seriebus dorsalibus punctis 6 compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus subtilioribus. Elytra thorace paulo longiora et scutellum subtiliter dense punctata, densius subtiliter griseo-pubescentia, subopaca. Abdomen subtiliter crebrius punctulatum, nitidulum, densius griseo-pubescent, segmentis ventralibus piceo-marginatis. Pedes fusco-testacei, geniculis tarsisque dilutioribus, coxis omnibus nigris, coxis intermediis approximatis, tibiis omnibus spinulosis.

Tarsi antici maris fortiter, feminae modice dilatati.

Obs. *Staph. Havaniensis* Laporte, Etud. Ent. 116.7, e Cuba, forte ad hanc, forte etiam ad praecedentem speciem pertinet, descriptione manca ambigua.



- \* 106. *Ph. luxurians*: *Elongatus*, *niger*, *nitidus*, *capite thoraceque sub-aeneis*, *elytris cyaneis*, *antennis articulo apicali luteo*, *capite suborbiculato*. — *Long.* 3 lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura omnino *Ph. ebenini* et minoribus eius individuis aequalis, niger, nitidus, capite thoraceque supra nigro-aeneis, politis. Antennae capitis thoracisque prope longitudine, articulo tertio secundo paulo longiore, 4—10 subcylindricis, sensim brevioribus et paulo crassioribus, penultimis crassitie brevioribus, ultimo ovato, apice leviter emarginato et inferne acuminato, nigrae, articulo ultimo laete luteo. Palpi nigri. Caput thorace paulo angustius, suborbiculatum, supra oculos utrinque punctis pluribus dispersis, inter oculos utrinque duobus transversim positus impressum. Thorax latitudine paulo longior, coleopteris dimidio angustior, antrorsum subangustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus deflexis posterioribusque subrotundatis, modice convexus, seriebus dorsalibus punctis 5 aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus minus subtilibus. Scutellum punctulatum, nigrum, nigro-pubescent. Elytra thoracis longitudine, minus confertim punctata, cyanea, nitida, parce subtiliterque nigro-pubescentia. Abdomen nitidum, supra crebrius, subtus parce punctatum, parce nigro-pubescent. Pedes nigri, coxis intermediis approximatis, tibiis omnibus spinulosis, tarsis piceis.

Tarsi antici feminae leviter dilatati. Mas latet.

- \* 107. *Ph. exiguus*: *Niger*, *nitidus*, *pedibus testaceis*, *elytris viridi-aeneis*, *parcius punctatis*, *capite ovato*. — *Long.* 2 lin.

*Nordm. Symb.* 105. 93. — *Er. Col. March.* I. 475. 44.

Habitat in Germania.

Habitu fere *Ph. micantis*, at dimidio minor, niger, capite thoraceque pernitidis, hoc nullo modo sericeo-micante. Antennae capite thoraceque breviores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus et crassioribus, penultimis crassitie parum brevioribus, ultimo ovato, apice leviter emarginato et inferne acuminato, nigrae, articulis tribus primis fusco-piceis. Palpi testacei. Mandibulae rufo-piceae. Caput thorace paulo minus et angustius, oblongo-ovatum, supra oculos utrinque punctis nonnullis, inter oculos utrinque duobus approximatis oblique positus impressum. Thorax coleopteris angustior, latitudine paulo longior, antrorsum subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus modice deflexis, subrotundatis, modice convexus, seriebus dorsalibus punctis 6 compositis, lateribus punctis praeter marginalia utrinque 6, interioribus eorum 4 in seriem alteram longitudinalem dispositis, impressus, punctis omnibus subtilioribus. Scutellum subtiliter punctulatum. Elytra thoracis longitudine, minus confertim subtilius punctata, minus subtiliter griseo-pubescentia, viridi-aenea, nitidula. Abdomen subtiliter punctulatum, griseo-pubescent, subtus apice testaceum. Pedes testacei, coxis anticis piceis, intermediis leviter distantibus, tibiis omnibus spinulosis.

Tarsi antici feminae simplices. Mas latet.

\* 108. *Ph. astutus*: *Elongatus, niger, nitidus, antennis basi piceis, ore pedibusque testaceis, thorace oblongo, capite ovato.* — Long. 3 lin.

Habitat in Austria, Dom. Ullrich.

*Ph. aterrimo* affinis, at plus duplo maior, et magis elongatus, niger, nitidus. Antennae capite thoraceque parum breviores, articulo tertio secundo sesqui prope longiore, 4 — 10 subobconicis, sensim paulo brevioribus et crassioribus, penultimis crassitie parum longioribus, ultimo ovato, apice subemarginato et inferne acuminato, fuscae, basi piceae. Os rufo-testaceum, palpis testaceis, maxillaribus articulo ultimo fortiter acuminato. Caput thoracis fere latitudine, subovatum, pone oculos utrinque punctis pluribus, inter oculos utrinque duobus approximatis oblique positis impressum, fronte medio obsolete canaliculata. Thorax coleopteris dimidio angustior, latitudine sesqui longior, basin versus subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, leviter convexus, seriebus dorsalibus punctis 6 compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus minus subtilibus. Scutellum crebre punctulatum. Elytra thorace paulo longiora, crebre punctata, minus subtiliter griseo-pubescentia, nigro-picea. Abdomen subtiliter punctulatum, longius griseo-pubesceus, segmentorum marginibus piceis, apice summo testaceum. Pedes cum coxis anterioribus rufo-testacei, femoribus posticis infuscatis, coxis intermediis approximatis, tibiis omnibus spinulosis. Alae hyalinae.

Tarsi antiqui feminae simplices. Mas latet.

Obs. Differt a *Ph. aterrimo* statura maiore et magis elongata, thoracis angulis anterioribus magis rotundatis, elytris longioribus et crebrius minusque fortiter punctatis.

X \* 109. *Ph. aterrimus*: *Elongatus, niger, antennis basi piceis, pedibus testaceis, capite ovato.* — Long. 2 lin.

*Er. Col. March. I.* 476. 45.

*Staph. aterrimus* Grav. *Micr.* 41. 62. — *Latr. Hist. nat. Crust. et Ins. IX.* 328. 80. — *Gyll. Ins. Suec. II.* 349. 65. — *Sahlb. Ins. Fenn. I.* 331. 62. — *Zetterst. Faun. Lappon. I.* 79. 31. *Ins. Lappon.* 65. 41. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 408. 43.

*Cafius aterrimus* Mannerh. *Brachelytr.* 32. 5. — *Runde Brachelytr. Hal.* 11. 2.

*Gyrophypnus aterrimus* Nordm. *Symb.* 126. 31.

*Staph. nigritulus* Grav. *Micr.* 41. 61. *Mon.* 94. 90. — *Latr. Hist. nat. Crust. et Ins. IX.* 328. 79.

*Cafius nigritulus* Mannerh. *Brachelytr.* 32. 6.

*Gyrophypnus nigritulus* Nordm. *Symb.* 126. 32.

*Cafius pumilus* Mannerh. *Brachelytr.* 32. 4.

*Philonth. trossulus* Nordm. *Symb.* 102. 90.

Habitat in Europa frequens; ex America septentrionali etiam a Dom. Zimmermann missus.

Minutus, niger, nitidus. Antennae capite thoraceque paulo breviores, articulo secundo tertioque subaequalibus, 4 — 10 sensim paulo brevioribus et crassioribus, penultimis crassitie haud longioribus, ultimo ovato, apice truncato et inferne subacuminato, nigrae, basi piceae. Palpi piceo-testacei. Caput maris thoracis latitudine vel interdum eodem latius, marium debilio-

runi et feminae thorace paulo angustius, depressiusculum, utrinque supra oculos punctis pluribus sparsis, inter oculos utrinque duobus oblique positis impressum, fronte maris medio obsolete canaliculata. Thorax coleopteris dimidio angustior, latitudine sesqui longior, basin versus subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, leviter convexus, seriebus dorsalibus punctis 6, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus minus subtilibus. Scutellum punctulatum. Elytra thorace paulo longiora, minus dense sat fortiter punctata, tenuiter minus subtiliter griseo-pubescentia. Abdomen subtilius punctulatum, dense griseo-pubescentia, subtus segmentis singulis piceo-marginatis. Pedes testacei, coxis intermediis approximatis, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

Variat thoracis serie dorsali altera 5-punctata.

Obs. 1. Magnitudine et capitis forma instabilis: caput maribus maioribus maius, thorace paulo latius, antrorsum subdilatatum, minoribus thoracis latitudine, oblongo-subquadratum, minimis thorace paulo angustius, oblongo-ovatum ut in femina.

2. *Ph. trossulus* Nord. a femina non distinctus, *Caf. pumilus* Manerh. ad individua feminea minus matura spectare videtur.

### Species mihi invisae.

110. *Ph. puella*: Niger, subnitidus, ore tarsisque fusco-ferrugineis, elytris nigris, subopacis, subtiliter punctulatis, thoracis lateribus multipunctatis, capite subovato.

Nordm. *Symb.* 101. 87. — *Er. Col. March.* 1. 472. 39.

Habitat Berolini.

*Ph. virgini* simillimus et valde affinis. Caput, palpi, oculi, antennae et elytra ut in *Ph. virgine* omnino. Thorax nonnihil brevior et latior, supra paulo minus convexus, angulis anticis minus quam in illo deflexis, sericeo-micans, lateribus obliquis; postice obtuse rotundatus, intra angulum anticum et versus medium laterum puncta 12 — 14 sine ordine sparsa et series dorsales punctis 6 — 7 impressis, omnibus sat conspicuis et profundis. Pedes nigro-picei, tarsis dilutioribus, ceterum ut in *Ph. virgine*. Species haec nisi quod ad puncturam thoracis attinet vix a *Ph. virgine* differt. — Individuum unicum, femina, tempore vernali, anno 1830, in quisquiliis lectum. (Nordm.)

111. *Ph. (?) stenoides*: Nitidus, oculis magnis prosilientibus. — Long. fere 6 lin.

Habitat in Java.

Caput forma tota capitis *Stenorum*, orbiculatum, et oculis magnis globosis fere transversum, thorace latius, punctum, viridi-subcupreo-aeneum, subtus nigrum, clypeo, labro et primo antennarum articulo (reliqui articuli desunt) testaceis, mandibulis falcatis, longis, acutis, nigris, maxillis nigris, apice rufo-testaceis, palpis longis, gracilibus, filiformibus, articulis duobus ultimis aequali longitudine et crassitie, haud clavatis, articulis omnibus nigris, summo apice rufo-testaceo. Thorax convexus, quadratus, basi rotun-



datum et paulo angustior quam apice, coleopteris angustior et brevior, nitidus, viridi-aeneus, in angulis apicis multipunctus, seriebus dorsalibus 6-punctis, et praeterea punctis quibusdam vagis maioribus impressus. Scutellum pallide viridi-sericans. Coleoptera quadrata, elytris planiusculis, punctis, viridicupreo-aenescentibus, macula magna disci pallida viridi-sericante. Abdomen nigrum, pectore sub-flavo-sericante, segmentorum marginibus testaceis. Pedes graciliores quam reliquis Staphylinis, capitibus femorum nigris, femoribus anterioribus testaceis, margine superiore nigro, posticis testaceis, geniculis nigris, tibiis anterioribus testaceis, margine superiore nigro, posticis testaceis, apice nigro, tarsis obscuris, anticis fossoriis. (Grav.)

Obs. An huius generis?

7) Thorax seriebus dorsalibus multipunctatis.

- \* 112. *Ph. helvolus*: Testaceus, nitidus, thorace utrinque parce punctato, seriebus dorsalibus multipunctatis. — Long 6 — 6½ lin.

Habitat in Madagascar, Dom. Goudot.

Corpus totum testaceum, nitidum, capite thoraceque rufo testaceis, abdominis leviter versicoloris segmentis basi fusco-testaceis. Antennae capite sesqui longiores, articulo tertio secundo sesqui longiore, 4 et 5 subcylindricis, longitudine decrescentibus, sexto praecedenti aequali, subobconico, 7 — 10 obconicis, crassitie brevioribus, ultimo subovato, apice compresso, testaceae, ante apicem fuscescentes. Caput maris magnum, thorace latius, transversim subquadratum, antrorsum dilatatum, antice inter oculos transversim impressum, crebrius punctatum, mandibulis valde elongatis, falcatis, acutis, piceis, — feminae thoracis latitudine, subquadratum, utrinque vage punctatum, supra antennarum insertionem utrinque oblique transversim impressum, impressione punctata, mandibulis mediocribus, piceis. Thorax coleopteris angustior, latitudine vix brevior, lateribus leviter, basi cum angulis posterioribus fortius rotundatus, apice truncatus, angulis anterioribus subrectis, modice convexus, seriebus dorsalibus irregulariter 11—12-punctatis, lateribus punctis praeter marginalia fere totidem, in lineam flexuosam dispositis, impressus. Scutellum crebre punctatum. Elytra thorace sesqui longiora, minus fortiter crebrius punctata, fulvo-pubescentia. Abdomen parcius subtiliter punctulatum, parce pubescens. Pedes dilutiores, coxis intermediis sat distantibus, tibiis omnibus subtiliter spinulosis.

Tarsi antici maris dilatati, feminae simplices, femora intermedia maris basi cum trochanteribus dilatati, trochanteres postici apice dentati, femora postica pone medium dente acuto fortiter hamato armata.

113. *Ph. mirabilis*: Rufus, nitidus, capite pedibusque nigris, antennis abdomineque apice albidis, ante apicem nigris, elytris splendidis, viridibus, fascia media cuprea. — Long. 5 lin.

Habitat Bogatae in Columbia interiore, Staph. mirabilis Dej. nomine a Dom. Buquet. communicatus.

Species pulcherrima, statura Ph. Baltimorensi affinis. Antennae capite sesqui longiores, articulis duobus primis testaceis, apice piceis, sequentibus tribus nigris, reliquis sex lutescenti-albidis. Palpi testacei. Caput thoracis

latitudine, suborbiculatum, utrinque parcius inaequaliter punctatum, fronte medio longitudinaliter late laevi, inter oculos punctis nonnullis vagis notata, nigrum, pernitidum. Thorax coleopteris duplo angustior, latitudine paulo longior, basin versus subangustatus, lateribus subrectis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, posterioribus obtusis, modice convexus, seriebus dorsalibus irregularibus, 10 — 11-punctatis, lateribus punctis praeter marginalia utrinque fere 16, partim subseriatis, impressus, punctis omnibus subtilioribus, rufus, nitidus. Scutellum rufum, parce subtiliter punctulatum. Elytra ampla, thorace sesqui longiora, parcius subtiliter punctata, parce nigro-griseoque-pubescentia, laete splendideque viridia, fascia media lata cupreo-aurea fulgente. Abdomen coleopteris angustius, apicem versus angustatum, infra parcius punctulatum, supra segmentis tribus primis laevissimis, quarto et quinto basi dense punctulatis, rufum, segmento quinto basi nigro, apice dimidiato-, sexto toto albido-flavo. Pectus rufum. Pedes nigri, coxis anticis testaceis, posterioribus rufis, intermediis modice distantibus, tibiis anticis inermibus, posterioribus spinulosis. Alae fusco-hyalinae.

Tarsi antici maris modice dilatati; femina latet.

\* 114. *Ph. bicolor*: Rufus, capite thoraceque cyaneis, abdomine cingulo nigro ante apicem. — Long. 6 lin.

*Staph. bicolor* Laporte Etud. Ent. 115. 6.

Habitat in Cayenna, Dom. Banon.

Statura omnino sequentis. Antennae breviusculae, capite paulo longiores, apicem versus sensim leviter incrassatae, articulo tertio secundo longiore, quarto subgloboso, 5 — 10 transversis, sensim crassioribus, ultimo breviter subacuminato, apice abrupte acuminato, articulis 4 primis rufis, reliquis nigris. Palpi rufi. Mandibulae piceae. Labrum rufum. Caput thorace paulo angustius, suborbiculatum, supra depressiusculum, punctis raris dispersis impressum, laete cyaneum, parum nitidum, subtus nigrum, nitidum. Thorax coleopteris dimidio angustior, latitudine parum longior, lateribus anterieus leviter rotundatis, pone medium subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, posterioribus subrotundatis, parum convexus, seriebus dorsalibus circiter 9-punctatis, lateribus punctis praeter marginalia utrinque fere 15 dispersis notatus, punctis omnibus fortioribus, laete cyaneus, subnitidus. Scutellum parcius punctatum, nigrum. Elytra thoracis latitudine, profundius dense punctata, rufo-testacea, tenuiter fulvo-pubescentia. Abdomen parcius punctatum, tenuiter fulvo-pubescent, rufum, segmento quarto apice, quinto basi nigris, hoc apice sextoque toto testaceis. Pectus rufum, antice nigricans. Pedes rufi, coxis anterioribus obscurioribus, intermediis sat distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu leviter dilatatis.

\* 115. *Ph. analis*: Niger, nitidus, capite, thorace elytrisque cyaneis, abdomine segmentis ultimis duobus rufis. — Long. 7 lin.

Habitat in Cayenna, Dom. Buquet; in Columbia, Dom. Moritz.

Niger, nitidus, capite, thorace elytrisque supra virescenti-coeruleis, ano rufo. Antennae capite sesqui fere longiores, apicem versus sensim leviter

incrassati, articulo tertio paulo longiore, quarto subgloboso, 6 — 10 transversis, sensim brevioribus at vix crassioribus, ultimo brevior, apice abrupte acuminato, nigrae. Palpi rufo-picei. Mandibulae breves, piceae. Caput thorace paulo angustius, suborbiculatum, depressiusculum, fronte posterius laevi, antierius punctis raris impressa, pone oculos utrinque crebrius punctatum. Thorax coleopteris dimidio angustior, latitudine paulo longior, basin versus subangustatus, lateribus antierius leviter rotundatis, pone medium subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, posterioribus subrotundatis, transversim convexus, seriebus dorsalibus minus regularibus, circiter 9-punctatis, lateribus punctis praeter marginalia utrinque fere 18 irregulariter dispersis impressus, punctis omnibus subtilioribus. Scutellum nigrum, crebre punctatum, parce pubescens. Elytra thoracis longitudine, crebre profundius punctata, tenuiter subtiliterque nigro-pubescens. Abdomen parce subtiliterque punctulatum, parce nigro-pubescens, segmentis ultimis duobus rufis, rufo-pubescentibus. Pedes breviusculi, nigri, coxis intermediis distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu leviter dilatatis. Alae fuscae.

Obs. Antennis brevioribus, apicem versus crassioribus et abdomine latiore cum praecedente Staphylini familiae secundae habitu non dissimilis, at ligula integra et huius generis mihi videtur.

\* 116. *Ph. xanthopygus*: *Niger, nitidus, abdomine segmentis duobus ultimis rufo-testaceis, capite subquadrato.* — Long. 6 lin.

*Staph. xanthopygus Nordm. Symb. 45. 27.*

Habitat in Brasilia, Dom. Virmond, in Columbia, Dom. Moritz, in Mexico, Dom. Deppe.

Subdepressus, niger, capite thoraceque nitidis, elytris abdomineque subnitidis, abdominis segmentis ultimis duobus solis testaceo-rufis. Antennae capite cum thorace breviores, crassiusculae, apicem versus haud incrassatae, articulo tertio secundo sesqui longiore, 4 et 5 subglobosis, 6 — 10 transversis, ultimo breviter subovato, apice abrupte acuminato, nigrae. Palpi breviores, nigri. Caput orbiculato subquadratum, thoracis latitudine, lateribus et vertice crebrius punctatum, fronte punctis parvis sparsis notata, medio longitudinaliter laevissima, antice inter antennis linea arcuata punctis notata impressa. Thorax coleopteris parum angustior, latitudine haud brevior, basin versus modice angustatus, lateribus praecipue ante medium leviter rotundatus, pone medium subsinuatus, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrectis, leviter convexus, seriebus dorsalibus minus irregularibus, punctis fere 10 compositis, lateribus praeter margines utrinque punctis fere 16, antierius crebrioribus, impressus, punctis omnibus subtilioribus, angulis anterioribus limbo crebre subtiliterque punctato. Scutellum creberrime punctulatum, nigro-pubescens. Elytra thorace paulo longiora, dense subtiliter punctata, densius subtiliter nigro-pubescens. Abdomen subtiliter punctulatum, parcius nigro-, apice in segmentis rufis testaceo-pubescens. Pedes breves, validiusculi, coxis intermediis leviter distantibus, tibiis omnibus spinulosis, tarsis breviusculis, anticis in utroque sexu fortius dilatatis. Alae fuscae.



- \* 117. *Ph. opulentus*: Nigro-subaeneus, capite thoraceque purpureo-violaceis, nitidis, elytris testaceis, circa scutellum nigricantibus, capite subquadrato. — Long. 7 lin.

Habitat in Brasilia, Dom. Prof. Germar.

Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus at vix crassioribus, 7—10 crassitie paulo brevioribus, ultimo breviter subovato, apice emarginato et inferne subacuminato, nigrae. Palpi nigri. Caput thoracis latitudine, subquadratum, supra parum convexum, utrinque, praecipue supra oculos, crebrius profundiusque punctatum, fronte medio longitudinaliter laevi, antice punctis paucis notata, nitidum, violaceum vel purpureum, subtus cum ore nigrum. Thorax coleopteris paulo angustior, latitudine haud brevior, basin versus subangustatus, basi et lateribus anterioribus modice rotundatis, his pone medium leviter sinuatis, apice truncatus, angulis anterioribus leviter deflexis, subrectis, posterioribus obtusis, modice transversim convexus, seriebus dorsalibus minus regularibus fere 10-punctatis, lateribus praeter margines punctis utrinque fere 24, plerisque subseriatis, impressus, punctis omnibus subtilioribus, violaceus vel purpureus, nitidus. Scutellum parce subtiliter punctulatum, nigro-subaeneum, opacum, tenuiter pubescens. Elytra thorace paulo longiora, parcius subtiliter punctata, parcius longius testaceo-pubescentia, testacea, parum nitida, spatio angustiore semicirculari basali communi circa scutellum nigro-subaeneo. Abdomen parce subtiliter punctatum et parce nigro-pubescentis, apice summo fulvo-pilosum, nigro-subaeneum, subnitidum, segmento sexto limbo apicali testaceo. Pectus nigrum. Pedes nigri, coxis intermedjis approximatis, tibiis omnibus spinulosis.

Tarsi antici feminae fortius dilatati. Mas latet.

Variat abdominis segmento sexto apice concolore.

- \* 118. *Ph. herilis*: Nigro-subaeneus, capite thoraceque cyaneis, elytris fusco-virescentibus, abdominis segmentis duobus ultimis rufo-testaceis, thorace utrinque parce punctato, capite subquadrato. — Long. 5½ lin.

Habitat in Brasilia, Dom. Virmond.

Statura omnino praecedentis, at duplo minor. Antennae capite thoraceque breviores, articulo tertio secundo paulo longiore, quarto quintoque subcylindricis, crassitie paulo longioribus, 6—10 paulo crassioribus, transversis, ultimo subovato, apice emarginato et inferne acuminato, piceae. Palpi picei. Caput thoracis fere latitudine, subquadratum, supra depressiusculum, utrinque supra oculos et in margine verticali crebrius punctatum, fronte punctis nonnullis dispersis versus latera et apicem notata, violaceo-cyaneum, nitidum, subtus nigrum. Thorax coleopteris paulo angustior, latitudine apicis vix longior, basin versus leviter angustatus, lateribus ante medium rotundatis, posterius subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rectis, posterioribus subrotundatis, transversim convexus, violaceus, nitidus, seriebus dorsalibus sat regularibus, 7—8-punctatis, lateribus praeter margines utrinque punctis fere 15 irregulariter seriatis impressus, punctis omnibus subtilioribus, violaceo-cyaneus, nitidus. Elytra thorace paulo longiora et scutellum fusco-virescentia, subopaca, sericeo-micantia, parce subtiliter punctata, tenuiter fusco-pubescentia. Abdomen parce pun-

ctatum, nigro-aeneum, nitidulum, fusco-pubescent, segmento quinto basi excepta sextoque toto rufo-testaceis, testaceo-pubescentibus. Pectus nigrum. Pedes fusco-picei, coxis intermediis approximatis, tibiis omnibus spinulosis, tarsis anticis in utroque sexu modice dilatatis.

\* 119. *Ph. punctifer*: *Nigerrimus, nitidus, thorace utrinque crebrius punctato, capite orbiculato*. — Long. 4 lin.

Nordm. Symb. 107. 96.

Habitat in Mexico, Dom. Deppe.

*Ph. puncto* affinis, at paulo maior, praecipue latior, totus nigerrimus, capite thoraceque nitidis. Antennae capite thoraceque paulo breviores, apicem versus haud incrassatae, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus, penultimis crassitie haud longioribus, ultimo ovato, apice truncato et inferne acuminato. Caput thorace paulo angustius, orbiculatum, supra lateribus crebrius subtiliusque punctatum, fronte media late laevi, inter oculos transversim punctata. Thorax basi coleopteris parum angustior, latitudine paulo brevior, antrorsum subangustatus, lateribus leviter, basi fortius rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, transversim convexus, seriebus dorsalibus fere 16-punctatis, lateribus praeter margines punctis utrinque numerosis (ultra 40) dispersis impressus, punctis omnibus subtilioribus. Elytra thorace paulo longiora, subopaca, creberrime subtiliter punctata, subtiliter nigro-pubescentia. Abdomen nitidulum, subtiliter crebrius punctatum, nigro-pubescent. Pedes coxis intermediis approximatis, tibiis omnibus spinulosis, tarsis anticis dilatatis.

\* 120. *Ph. punctus*: *Niger, nitidus, thorace lateribus crebre punctato, capite orbiculato, crebre punctato, medio laevi*. — Long. 4 lin.

Er. Col. March. I. 477. 47.

*Staph. punctus* Grav. Micr. 20. 22. Mon. 85. 75. — Boisd. et Lacord. Faun. Ent. Paris. I. 393. 14. — Runde Brachelytr. Hal. 10. 55.

*Staph. punctatus* Latr. Hist. nat. Crust. et Ins. IX. 309. 29.

*Staph. multipunctatus* Mannerh. Brachelytr. 31. 79.

*Philonth. multipunctatus* Nordm. Symb. 106. 95.

*Staph. politus* Panz. Faun. Germ. 27. 7.

Habitat in Germania; in Sardinia etiam a Dom. Prof. Gené lectus.

Statura omnino *Ph. atrati*, niger, subnitidus, capite thoraceque pernitidis. Antennae capite thoraceque parum breviores, articulo tertio secundo paulo longiore, 4—10 subobconicis, sensim paulo brevioribus et crassioribus, prioribus elongatis, penultimis crassitie vix longioribus, ultimo subovato, apice emarginato et inferne acuminato, nigrae. Caput thorace parum angustius, orbiculatum, supra fortiter crebre punctatum, fronte medio laevi. Thorax coleopteris paulo angustior, latitudine haud brevior, antrorsum subangustatus, lateribus subrectis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, posterioribusque subrotundatis, leviter convexus, seriebus dorsalibus circiter 12-punctatis, lateribus punctis praeter marginalia utrinque numerosis (fere 30), aequaliter dispersis, minus subtilibus impressus. Scutellum confertim punctatum, nigro-pubescent. Elytra thorace paulo longiora, densius subtiliusque punctata, vix virescentia, tenuiter nigro-pubescentia.

Abdomen parcius punctulatum, tenuiter nigro-pubescens. Pedes nigri, coxis intermediis approximatis, tibiis omnibus spinulosis. Alae fuscae.

Tarsi antici maris fortiter, feminae leviter dilatati.

\* 121. *Ph. parumpunctatus*: Niger, nitidus, elytris nigro-aeneis, thorace lateribus parce punctato, capite orbiculato. — Long. 4 lin.

*Staph. punctus* Gyll. Ins. Suec. II. 346. 62. — *Sahlb. Ins. Fenn. I. 330. 59.* — *Mannerh. Brachelytr. 31. 78.*

*Philonth. punctus* Nordm. Symb. 106. 94.

Habitat in Suecia, sec. Gyll., in montibus Uralensibus, Dom. Prof. Eversmann.

Statura omnino *Ph. variantis* et maximis eius individuis aequalis, niger, nitidus, capite thoraceque pernitidis, elytris obscure aeneis, nitidulis. Antennae capite sesqui longiores, articulo tertio secundo vix longiore, 4—10 subobconicis, sensim paulo brevioribus et haud crassioribus, penultimo crassitie vix brevior, ultimo subovato, apice leviter emarginato et inferne acuminato, piceae. Palpi picei. Caput thoracis latitudine, orbiculatum, leviter convexum, supra oculos utrinque punctis pluribus, inter oculos punctis 4 transversim positis, inter se aequaliter distantibus, profundioribus, impressum. Thorax coleopteris paulo angustior, latitudine haud longior, antrorsum subangustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque rotundatis, modice convexus, seriebus dorsalibus punctis circiter 8 compositis, lateribus praeter margines utrinque punctis parvis (vix ultra 12) dispersis impressus, punctis omnibus minus subtilibus. Scutellum subtiliter punctulatum, parce nigro-pubescens. Elytra thorace paulo longiora, crebre profundius punctata, parcius longiusque griseo-pubescens. Abdomen parce subtiliter punctulatum, tenuiter griseo-pubescens. Pedes breviusculi, picei, coxis intermediis approximatis, tibiis omnibus spinulosis. Alae fuscae.

Tarsi antici maris modice dilatati, feminae simplices.

Obs. A praecedente satis distinctus statura paulo minore, capitis thoracisque punctura, elytris aeneis.

\* 122. *Ph. orphanus*: Nigro-piceus, nitidus, pedibus testaceis, capite subquadrato, thorace seriebus dorsalibus 7-, lateribus crebrius subtiliter punctatis. — Long. 2½ lin.

Habitat in Madagascar, Dom. Goudot.

Habitu fere *Ph. debilis*, nigro-piceus, nitidus. Antennae tenuiores, articulo tertio secundo paulo longiore, sequentibus teretibus, vix sensim brevioribus, fuscae, articulo primo rufo-piceo (articuli ultimi tres desunt). Palpi rufo-picei. Mandibulae piceae. Caput maiusculum, thorace paulo latius, subquadratum, parum convexum, utrinque parce subtiliter punctatum, spatio medio longitudinali lato laevi. Thorax coleopteris paulo angustior, latitudine paulo longior, basin versus subangustatus, lateribus ante medium leviter rotundatus, pone medium subsinuatus, basi rotundatus, apice truncatus, angulis anterioribus deflexis, rotundatis, posterioribus obtusis, subcylindricus, seriebus dorsalibus minus regularibus, fere 7-punctatis, lateribus punctis praeter marginalia pluribus (fere 16) utrinque impressus, punctis omni-



bus subtilibus. Scutellum punctis parvis subtilibus notatum. Elytra thorace sesqui fere longiora, parcius subtiliusque punctata, parce pubescentia. Abdomen crebre subtilius punctulatum, longius et minus subtiliter nigro-pubesceus, subtus segmentis singulis apice piceo-marginatis. Pedes rufo-testacei, coxis piceis, intermediis approximatis, tibiis omnibus subtiliter spinulosis.

Tarsi antici feminae simplices. Mas latet.

- \* 123. *Ph. dimidiatipennis*: *Elongatus, niger, antennarum articulis 3 primis, pedibus elytrisque rufis, his basi nigris, thorace seriebus dorsalibus 7 — 8-punctatis, lateribus parumpunctato.* — Long.  $2\frac{1}{2}$  lin.

Habitat in Sicilia, collect. Dom. Aubé, in Mesopotamia, Dr. Helfer.

Elongatus, linearis, subdepressus, niger, nitidus. Antennae capitis thoracisque prope longitudine, articulo secundo tertioque subaequalibus, 4—10 sensim brevioribus et crassioribus, penultimis transversis, ultimo subovato, apice leviter emarginato et inferne subacuminato, nigrae, articulis 3 primis rufo-testaceis. Palpi rufo-testacei. Mandibulae rufo-piceae. Caput thoracis latitudine, subovatum, utrinque crebrius punctatum, spatio medio longitudinali laevi. Thorax coleopteris angustior, latitudine paulo longior, lateribus parallelis, rectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, parum convexus, seriebus dorsalibus 7 — 8-punctatis, lateribus praeter margines punctis utrinque 4 vel 5 impressus. Scutellum punctulatum, tenuiter pubescens. Elytra thorace paulo longiora, crebrius sat distincte punctata, tenuiter griseo-pubescentia, testaceo-rufa, nitida, parte tertia basali nigra. Abdomen crebre subtiliter punctatum, minus nitidum, densius griseo-pubesceus, pube certo situ sericante, subtus segmentis singulis apice piceo-marginatis. Pedes rufo-testacei, coxis omnibus concoloribus, posticis antice nigricantibus, intermediis approximatis, tibiis omnibus parce spinulosis.

Tarsi antici feminae leviter dilatari. Mas latet.

Obs. A *Ph.* prolixo statura duplo maiore, minus depressa, capitis spatio medio longitudinali laevi latiore, thorace aliter punctato, abdomine subtiliter punctato, coxis anticis rufo-testaceis, etc. valde distinctus.

124. *Ph. labilis*: *Niger, elytris pectoreque rufis, opacis, antennarum basi, palpis pedibusque flavis, thorace seriebus dorsalibus multipunctatis, capite orbiculato.* — Long.  $2\frac{3}{4}$  lin.

Habitat in Guyana Venezuelensi, Dom. Moritz.

*Ph. fulvipedi* similis, at paulo gracilior, niger, capite thoraceque nitidis. Antennae capitis thoracisque longitudine, articulo tertio secundo paulo longiore, 4 — 10 oblongo-obconicis, sensim paulo brevioribus et crassioribus, ultimo oblongo, apice oblique truncato et inferne acuminato, fuscae, articulo primo toto, secundo tertioque basi testaceis. Palpi flavo-testacei. Caput thoracis prope latitudine, suborbiculatum, utrinque supra oculos parce punctatum. Thorax latitudine paululum longior, coleopteris dimidio angustior, antrorsum subangustatus, lateribus levissime, basi fortius rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rotundatis, posterioribus obtusis, leviter convexus, seriebus dorsalibus 8 — 9-punctatis, lateribus praeter marginem punctis utrinque 5 impressus, punctis omnibus subtilibus. Scu-

tellum obsolete punctatum, opacum, tenuiter griseo-pubescent. Elytra thorace paulo longiora, dense subtilissime punctata, laete rufa, opaca, dense subtiliter griseo-pubescentia. Abdomen dense subtiliter punctatum, densius griseo-pubescent, opacum, nigrum, subtus apice testaceum. Pectus obscure rufum. Pedes flavo-testacei, coxis anterioribus concoloribus; intermediis approximatis, tibiis omnibus spinulosis.

Tarsi antici feminae leviter dilatati. Mas latet.

- \* 125. *Ph. pacificus*: *Elongatus, niger, abdomine subtus, elytrorum margine inflexo et summo apicali pedibusque ferrugineis, capite oblongo-subquadrato, thorace seriebus dorsalibus 11-punctatis, lateribus parce punctatis.* — Long. 5 lin.

Habitat in Terra Van-Diemenii, Dom. Schayer.

Statura elongata subdepressa omnino *Ph. fucicolae*. Antennae capite sesqui longiores, apicem versus haud incrassatae, articulo tertio secundo tenuiore et sesqui longiore, 4—10 sensim brevioribus, penultimo crassitie vix longiore, ultimo ovato, apice truncatulo et inferne acuminato, fuscae, articulis tribus primis nigris. Palpi picei, articulo ultimo rufo. Mandibulae piceae. Caput thorace paulo latius, oblongo-subquadratum, parum convexum, utrinque crebrius vage fortiter profundeque punctatum, spatio medio longitudinali inaequali laevi, nigrum, nitidum. Thorax coleopteris plus dimidio angustior, latitudine sesqui longior, basin versus leviter angustatus, basi et lateribus ante medium leviter rotundatis, his pone medium subsinuatis, apice truncatus, angulis anterioribus rotundatis, posterioribus obtusis, leviter convexus, seriebus dorsalibus sat regularibus, lineae leviter impressae impositis, circiter 11-punctatis, lateribus punctis praeter marginalia utrinque fere 10, partim subseriatis, impressus, niger, nitidus, margine summo apicali et laterali inflexo sanguineo. Scutellum crebre punctatum, nigrum, opacum, subtiliter nigro-pubescent. Elytra thorace sesqui longiora, confertissime subtilius punctata, subrugulosa, nigra, opaca, limbo laterali inflexo et margine summo apicali ferrugineis, subtiliter nigro-pubescentia, apice cinereo-ciliata. Abdomen crebrius subtiliter punctatum, subnitidum, supra nigrum, segmentis 4 primis macula obsoleta, quinto apice, sexto toto subferrugineis, subtus totum ferrugineum, tenuiter subtiliterque supra nigro-, subtus ferrugineo-pubescent. Pectus nigrum. Pedes ferruginei, coxis intermediis leviter distantibus, tibiis omnibus subtiliter spinulosis.

Tarsi antici feminae leviter dilatati. Mas latet.

- \* 126. *Ph. fonticola*: *Ferrugineus, cinereo-sericeus, opacus, capite subquadrato thoraceque basin versus angustatis nitidis, supra nigropiceis, nitidis, elytris nigris, margine laterali, macula basali apicalibusque pallidis.* — Long. 3 lin.

Habitat in montibus Sinai, in regione montis S. Catharinae media ad fontes lectus a Dom. Prof. Ehrenberg.

Affinis *Ph. xantholomati*, statura autem brevior, crassiuscula. Antennae capite parum longiores, apice haud crassiores, articulo tertio secundo tenuiore at vix longiore, sequentibus omnibus transversis, sensim paulo brevioribus, ultimo ovato, apice profundius emarginato et inferne acuminato,



totae rufo-ferrugineae. Palpi rufi. Caput thorace fere latius, subquadratum, parum convexum, dense fortiter punctatum, spatio medio longitudinali, basin versus angustato, laevissimo, polito, nigro-piceum, nitidum. Thorax coleopteris paulo angustior, latitudine parum longior, basin versus angustatus, lateribus ante medium cum angulis anterioribus fortius, basi cum angulis posterioribus leviter rotundatis, leviter convexus, seriebus dorsalibus multipunctatis, duplicatis, interstitio subelevato, lateribus utrinque confertim punctatus, punctis omnibus piliferis, nigro-piceus, nitidus. Scutellum bruneum, creberrime punctulatum, griseo-pubescentia. Elytra thorace sesqui longiora, confertissime subtiliter punctata, subtiliter cinereo-pubescentia, opaca, brunnea, margine laterali inflexo, puncto ad basin scutelli, macula in angulo apicis exteriori, punctoque in angulo apicis interiore pallide flavis, puncto ad apicem scutelli ferrugineo. Abdomen confertissime subtiliter punctatum, pube densiore brevi tenui cinereo-sericea vestitum, supra e pube densiore bifariam submaculatum, supra obscure, subtus dilute ferrugineum. Pectus ferrugineum. Pedes ferruginei, femoribus extus nigricantibus, coxis intermediis leviter distantibus, tibiis omnibus spinulosis.

Tarsi antici feminae fortius dilatati. Mas latet.

- \* 127. *Ph. bistriatus*: *Elongatus, nigro-piceus, opacus, abdomine subtus pedibusque ferrugineis, elytrorum margine inflexo ochraceo, thorace lateribus crebre punctato, dorso lineis duabus multipunctatis impresso.*  
— Long.  $3\frac{1}{2}$  lin.

Habitat in Americae septentrionalis Insula Longa (Long-Island), Dom. Zimmermann.

Habitu prope *Ph. xantholomatis*, at angustior et paulo minor magisque depressus. Antennae capite plus sesqui longiores, apicem versus haud incrassatae, articulo tertio secundo subaequali, 4—10 sensim paulo brevioribus, ultimo subovato, apice truncatulo et inferne subacuminato, piceae, articulo primo nigricante. Mandibulae et palpi rufo-picei. Caput thorace vix latius, oblongo-subquadratum, antrorsum subangustatum, basi truncatum, angulis posterioribus leviter rotundatis, supra subdepressum, crebrius fortiusque profunde punctatum, plaga media longitudinali antrorsum dilatata laevi, nigrum, opacum, subtilissime nigro-pubescentia, parce fusco-pilosum. Thorax coleopteris dimidio prope angustior, latitudine tertia parte longior, basin versus leviter angustatus, basi rotundatus, lateribus rectis, pone medium subsinuatus, apice utrinque oblique truncatus, angulis anterioribus subrectis, posterioribus obtusis, subrotundatis, subdepressus, dorso striis duabus longitudinalibus parallelis impressis, apice evanescentibus, series dorsales e punctis numerosis continentibus, spatio interiecto laevissimo, nitidulo, lateribus sat crebre subaequaliter minus fortiter punctatis, opacis, tenuiter subtiliterque griseo-pubescentibus. Scutellum dense subtiliter punctatum. Elytra thorace sesqui longiora, dense subtiliter punctata, pube brevi depressa grisea sericante densius vestita, brunneo-picea, margine laterali inflexo ochraceo. Abdomen confertim subtiliter punctatum, pube depressa grisea sericante densius vestitum, supra nigro-piceum, segmento sexto et subtus totum rufo-piceum vel ferrugineum. Pedes cum coxis anterioribus ferruginei, tibiis omnibus spinulosis.

Tarsi antici feminae leviter dilatati. Mas latet.



128. *Ph. bilineatus*: *Elongatus, niger, antennarum basi, pedibus, abdominisque segmentorum ventralium marginibus piceis, elytris dense subtiliter punctatis, thorace lateribus crebre punctato, dorso lineis duabus longitudinalibus impresso. — Long. 3 lin.*

Habitat in Americae meridionalis ins. St. Johannis, Dom. Moritz.

Statura omnino et summa affinitas praecedentis. *Elongatus*, subdepressus, niger, capite thoraceque nitidis. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus at non crassioribus, crassitie paulo longioribus, ultimo ovato, apice truncato et inferne acuminato, nigrae, articulo primo piceo. Palpi picei. Caput thorace prope latius, subovatum, leviter convexum, fronte utrinque crebrius profunde punctata, spatio medio longitudinali angustiore et inaequali laevi. Thorax coleopteris plus dimidio angustior, basin versus leviter angustatus, latitudine apicis sesqui longior, lateribus anterieus leviter rotundatis, pone medium subsinuatis, basi rotundatus, apice utrinque oblique truncatus, angulis anterioribus rotundatis, posterioribus obtusis, leviter convexus, utrinque crebre punctatus, spatio medio longitudinali angustiore laevi, utrinque linea longitudinali impressa, punctorum fere 16 impressorum seriem continente, terminato. Scutellum crebre subtiliter punctulatum, pubescens. Elytra thorace tertia parte longiora, dense subtiliter punctata, densius subtiliter griseo-pubescentia. Abdomen crebrius subtiliter punctatum, densius griseo-pubescentis, subtus segmentis singulis apice rufo-piceis. Pedes picei, coxis intermediis approximatis, tibiis omnibus parcius spinulosus.

Tarsi antici feminae leviter dilatati. Mas latet.

Obs. Praecedenti proximus, at capite minus crebre punctato, thoracis angulis anterioribus rotundatis, elytrorum margine inflexo concolore, abdomine supra nitido etc. distinctus.

- 8) Thorax crebre punctatus, linea media longitudinali laevi.

- \* 129. *Ph. Baltimorensis*: *Rufo-castaneus, capite thoraceque nigris, nitidis, hoc lateribus subtiliter vage punctato. — Long. 5½ lin.*

*Nordm. Symb. 109. 99.*

*Staph. Baltimorensis Grav. Micr. 163. 8. Mon. 112. 122.*

Habitat in America septentrionali.

Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus et crassioribus, 5—10 transversis, ultimo apice truncato et inferne subacuminato, nigro-picei, articulis ultimis 4 vel 5 rufo-ferrugineis. Palpi rufo-picei. Caput maris thorace latius, feminae thoracis latitudine, orbiculatum, parce subtilius punctatum, spatio medio longitudinali laevi, nigrum, nitidum. Thorax coleopteris dimidio angustior, latitudine paulo longior, basin versus subangustatus, lateribus anterieus leviter rotundatis, posterius subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, obtusiusculis, posterioribus subrotundatis, convexus, utrinque parcius aequaliter subtiliterque punctatus, niger, nitidus. Scutellum crebre subtiliter punctatum, nigricans. Elytra thoracis latitudine, castaneo-rufa, nitidula, subtilius punctata, tenuiter griseo-pubescentia. Abdomen apicem versus sensim attenuatum, crebrius punctulatum, tenuiter griseo-pubescentis, rufo-castaneum, segmento sexto utrinque nigro-limbato.

Pectus nigro-piceum. Pedes rufo-, rarius nigro-picei, coxis intermediis sat distantibus, tibiis intermediis crebrius, anticis posticisque parce spinulosis.

Tarsi antici maris leviter dilatati, feminae simplices. Mas praeterea capite maiore, antrorsum dilatato, et mandibulis elongatis distinctus.

\* 130. *Ph. haematurus*: *Niger, nitidus, abdomine pone medium obscure sanguineo, thorace utrinque crebrius punctato.* — Long. 5 lin.

Habitat in America septentrionali.

Affinis praecedenti, at corpus nigrum, nitidum. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 4 — 10 leviter obconicis, subaequalibus, crassitie paulo longioribus, ultimo subovato, apice oblique submarginato et inferne acuminato, basi nigrae, dein fuscae, articulo apicali ferrugineo. Palpi picei. Caput thoracis latitudine, suborbiculatum, crebrius fortiusque punctatum, fronte spatio medio longitudinali laevi. Thorax coleopteris dimidio angustior, latitudine paulo longior, basin versus subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, sat convexus, utrinque crebrius punctatus, spatio medio longitudinali angusto laevi. Scutellum creberrime subtiliter punctatum. Elytra thorace vix longiora, subtilius punctata, nigro-pubescentia. Abdomen apicem versus attenuatum, crebrius fortiusque punctulatum, nigro-pubescentia, abdominis segmento quarto apice obscure sanguineo, quanto toto sanguineo, ultimo sanguineo, apice nigro. Pedes nigri, coxis intermediis sat distantibus, tibiis anticis posticisque parce, intermediis crebrius spinulosis.

Tarsi antici maris leviter dilatati, feminae simplices.

Obs. In specimine nostro masculo caput maris perparum maius quam feminae, et mandibulae parum magis elongatae.

\* 131. *Ph. rufipennis*: *Niger, nitidus, elytris, tibiis tarsisque rufis, antennis apice ferrugineis, thorace utrinque crebrius punctato.* — Long. 5 lin.

*Nordm. Symb.* 110. 100. — *Er. Col. March.* I. 477. 46.

*Staph. rufipennis* Grav. *Micr.* 40. 59. *Mon.* 104. 105. — *Latr. Hist. nat. Crust. et Ins.* IX. 327. 77.

Habitat in Germania, rarus; in Gallia a Dom. Aubé, in Sardinia a Dom. Prof. Gené lectus.

Statura *Ph. Baltimorensis*. Niger, nitidus, capite thoraceque pernitidis, elytris rufis. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo brevioribus et crassioribus, penultimis crassitie prope brevioribus, ultimo ovato, apice oblique submarginato et inferne subacuminato, nigro-piceae, articulo ultimo ferrugineo. Palpi rufi. Mandibulae rufo-piceae. Caput maris magnum, thorace paulo latius, orbiculato-subquadratum, feminae thoracis latitudine, suborbiculatum, fronte parcius fortiusque punctata, spatio medio longitudinali laevi. Thorax coleopteris dimidio angustior, latitudine paulo longior, basin versus subangustatus, lateribus subrectis, basi leviter rotundatus, apice truncatus, angulis anterioribus leviter, posterioribus fortius rotundatis, sat convexus, crebrius subtiliusque aequaliter punctatus, spatio medio longitudinali angustiore laevissimo. Scutellum dense subtiliter punctulatum, tenuiter pubescens. Elytra

thorace paulo longiora, parcius punctata, tenniter griseo-pubescentia. Abdomen apicem versus sensim attenuatum, parcius punctulatum, parce nigro-pubescent, lucidum. Pedes picei, tibiis tarsisque dilutioribus, coxis intermediiis sat distantibus, tibiis anticis posticisque parce, intermediiis crebrius spinulosis.

Tarsi antici maris leviter dilatati, feminae simplices. Mas praeterea capite maiore, antice angulato et mandibulis modice elongatis distinctus.

\* 132. *Ph. egregius*: *Rufus, capite nigro-aeneo, elytris viridibus, abdomine apice albido-flavo, pedibus antennisque nigris, his apice lutescentibus.* — Long. 5 lin.

Habitat Bogotae in Columbia interiore, a Dom. Buquet nomine Staph. ruficollis Dej. communicatus.

Statura fere Ph. decori. Antennae longiores, capite thoraceque haud breviores, apicem versus haud incrassatae, articulo tertio secundo paulo longiore, 4—10 cylindricis, sensim paulo brevioribus, omnibus crassitie longioribus, ultimo oblongo-ovato, apice inferne acuminato, nigrae, apice griseo-lutescentes. Palpi nigri, apice summo testaceo. Caput thoracis fere latitudine, orbiculatum, utrinque vage crebrius profundius punctatum, fronte medio longitudinaliter laevi, antierius punctis parvis dispersis notata, nigro-aeneum, nitidulum, collo rufo. Thorax coleopteris dimidio angustior, latitudine paulo longior, basin versus leviter angustatus, lateribus antierius leviter et basi rotundatus, apice truncatus, angulis antierioribus leviter deflexis, subrectis, posterioribus obtusis, leviter convexus, crebre subtilius punctatus, spatio medio longitudinali angustiore laevissimo, rufus, nitidus. Scutellum dense subtiliter punctatum, tenuiter pubescens, nigrum. Elytra thorace sesqui longiora, cyanescenti-viridia, nitida, densius subtiliter punctata, tenuiter nigro-pubescentia. Abdomen apicem versus subangustatum, subtiliter punctulatum, tenuiter nigro-pubescent, rufum, segmento quinto nigro, apice cum sexto flavo. Pedes cum coxis antierioribus nigri, coxis intermediiis approximat, tibiis omnibus parcius spinulosis.

Tarsi antici feminae simplices. Mas latet.

\* 133. *Ph. fervidus*: *Niger, nitidus, antennarum basi, capite, thorace abdominisque apice rufis, elytris viridibus.* — Long. 5 lin.

Habitat in Columbia, Dom. Moritz.

Statura omnino praecedentis, at antennae breviores et crassiores. Antennae capite sesqui longiores, crassiusculae, articulo tertio secundo paulo longiore, 4—10 sensim brevioribus et crassioribus, 6—10 transversis, ultimo apice inferne acuminato, nigrae, articulis tribus primis rufis. Caput thorace paulo angustius, suborbiculatum, supra parum convexus, fronte medio late laevi, lateribus et antice inter oculos fortiter irregulariter punctata, totum rufum, nitidum. Thorax coleopteris angustior, basin versus subangustatus, latitudine vix longior, lateribus ante medium leviter rotundatis, pone medium subsinuatis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis antierioribus leviter deflexis, subrectis, transversim conve-



xus, utrinque fortius crebriusque aequaliter punctatus, spatio medio longitudinali angustiore laevisimo, rufus, nitidus. Scutellum nigrum, crebre punctatum, parce nigro-pubescent. Elytra thorace paulo longiora, cyane-scenti-viridia, nitida, fortiter punctata, nigro-pubescentia. Abdomen nigrum, iridescent, segmento quinto apice rufescente, sexto rufo, nigro-pubescent, parce punctulatum, supra segmentis anterioribus tribus laevibus, utrinque lineola tenui obliqua impressis. Pectus nigrum. Pedes cum coxis nigri, coxis intermediis leviter distantibus, tibiis omnibus spinulosis, tarsis anticis in utroque sexu simplicibus.

\* 134. *Ph. viduus*: *Niger, subnitidus, elytris cyaneis, abdominis segmento ultimo subtus flavo.* — Long. 6 lin.

Habitat in Columbia, Dom. Moritz.

Corpus nigrum, subnitidum. Antennae capite thoraceque paulo breviores, apicem versus haud incrassatae, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus, subobconicis, omnibus crassitie longioribus, ultimo ovato, apice oblique truncato et inferne acuminato, totae nigrae. Palpi nigri. Mandibulae subelongatae, arcuatae, tenues, acutae. Caput thorace vix latius, suborbiculatum, utrinque crebre fortius punctatum, spatio medio longitudinali angustiore laevi, inter antennarum basin utrinque obsolete longitudinaliter impressum. Thorax coleopteris duplo fere angustior, latitudine sesqui longior, basin versus subangustatus, lateribus ante medium perparum rotundatis, pone medium leviter sinuatis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis, rectis, posterioribus subrotundatis, transversim convexus, fortius crebre punctatus, spatio medio longitudinali angustiore laevi. Scutellum dense punctulatum, nigrum, nigro-pubescent. Elytra thorace paulo longiora, subtiliter punctata, cyanea, subnitida, nigro-pubescentia. Abdomen parcius punctulatum, supra subaeneum, subtus segmento anali flavo. Pedes nigri, coxis intermediis leviter distantibus, tarsis omnibus spinulosis. Alae fuscae.

Tarsi antichi maris leviter dilatati, feminae simplices.

\* 135. *Ph. binotatus*: *Niger, nitidus, elytris angulo apicis exteriori late rufo, capite orbiculato.* — Long. 3½ lin.

*Staph. binotatus* Grav. Mon. 73. 62.

Habitat in Borussia, Kugelann.

Statura omnino *Ph. puncti* eoque vix minor, niger, nitidus, elytrorum angulo apicali exteriori solo late a suturae apice inde ultra marginis lateralis medium usque rufo. Antennae capitis thoracisque longitudine, articulo tertio secundo longiore, 4 — 10 elongato-obconicis, sensim paulo brevioribus et crassioribus, ultimo oblongo, apice subtruncato et inferne acuminato, nigrae. Palpi nigri. Caput thorace parum angustius, orbiculatum, crebrius punctatum, punctis in fronte rarioribus, sparsis. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus parallelis, subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrotundatis, modice convexus, crebre profundius punctatus, spatio medio longitudinali angusto laevisimo. Elytra thorace paulo longiora et scutellum crebrius subtiliusque punctata, griseo-pubescentia. Abdomen

crebrius punctulatum, griseo-pubescent. Pedes concolores, coxis intermediis approximatis, tibiis omnibus spinulosis.

Tarsi antici maris fortiter dilatati, femina latet.

\* 136. *Ph. ephippium*: *Niger, nitidus, pedibus elytrisque rufis, his macula communi scutellari nigra, capite orbiculato.* — Long.  $3\frac{1}{2}$  lin.

*Nordm. Symb.* 110. 101.

Habitat in Grusia, Dom. Marschall v. Bieberstein.

Affinis praecedenti, niger, capite thoraceque nitidis, elytris laete rufis, macula magna communi subquadrata circa scutellum nigra, pedibus dilute rufis. Antennae capitis thoracisque fere longitudine, articulo tertio secundo sesqui prope longiore, 4—10 oblongo-obconicis, sensim paulo brevioribus at vix crassioribus, ultimo oblongo, apice truncatulo et inferne acuminato, fusco-nigrae, basi piceae. Palpi dilute picei. Caput thorace angustius, orbiculatum, supra leviter convexum, crebrius punctatum, fronte medio longitudinaliter laevi, nonnisi inter oculos parce punctata. Thorax basi coleopteris angustior, latitudine paulo longior, antrorsum leviter angustatus, lateribus subrectis, basi rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, modice transversim convexus, supra crebre fortius punctatus, punctis ut in capite omnibus piliferis, spatio medio longitudinali angusto laevissimo. Scutellum densius punctulatum, nigrum, griseo-pubescent. Elytra subtilius crebriusque punctata, thorace tertia parte longiora, griseo-pubescentia. Abdomen parcius subtiliter punctulatum, griseo-pubescent. Pedes dilute rufi, coxis omnibus nigris, intermediis approximatis, tibiis omnibus spinulosis.

Tarsi anticis maris fortius, feminae leviter dilatati.

137. *Ph. leucotus*: *Niger, capite thoraceque nigro-subaeneis, nitidulis, pedibus elytrisque rubro-testaceis, his fascia subfusca, abdominis segmentis inferioribus 3 primis rufis, antennis articulis ultimis 5 albidis.* — Long. 5 lin.

Habitat in Nepal, Mus. Dom. Hope.

Antennae capitis thoracisque fere longitudine, articulo tertio secundo sesqui longiore, 4—10 sensim paulo brevioribus at vix crassioribus, prioribus elongatis, teretibus, penultimis oblongo-subobconicis, ultimo oblongo-ovato, apice inferne acuminato, nigrae, articulis 7—11 albis. Mandibulae et palpi modice elongati, rufo-picei. Caput thorace paulo angustius, orbiculatum, parum convexum, creberrime punctatum, fronte medio laevi, nigro-subaeneum, nitidulum. Thorax coleopteris parum angustior, latitudine vix brevior, antrorsum subangustatus, lateribus subrectis, basi rotundatus, apice subemarginatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, leviter convexus, confertissime punctatus, linea media longitudinali angusta subelevata laevi, nigro-subaeneus, nitidulus, tenuiter subtiliterque griseo-pubescent. Scutellum punctatissimum, tenuiter pubescens, nigrum. Elytra thoracis longitudine, confertissime punctata, fulvo-pubescentia, rubro-testacea, pone medium fascia arcuata obsoleta fusca. Abdomen subtilius dense punctatum, tenuiter griseo-pubescent, supra nigrum, subnitidum, segmento quinto margine apicali tenui albo, subtus nigrum, nitidum,

segmentis tribus primis rufis. Pectus nigrum, subnitidum. Pedes longiores, rubro-testacei, coxis omnibus nigris, coxis intermediis approximatis, tibiis anticis muticis, posterioribus parcius spinulosis. Alae fusco-hyalinae.

Tarsi antici maris leviter dilatati. Femina latet.

- \* 138. *Ph. elatus*: *Niger, nitidus, elytris viridi-metallicis, ano rufo, thorace elongato, valde convexo, confertim subtiliter punctato.* — Long.  $5\frac{1}{2}$  lin.

Habitat in Madagascar, Dom. Goudot.

Habitus omnino sequentis, at duplo fere maior. Caput deest. Thorax coleopteris paulo angustior, basin versus sensim angustatus, latitudine longior, basi cum angulis posterioribus leviter rotundatus, lateribus anterieus leviter rotundatis, posterius subsinuatis, apice truncatus, angulis anterioribus rotundatis, elevatus, valde convexus, crebre subtiliter punctatus, tenuiter subtilius nigro-pubescent, linea angusta media longitudinali laevi, niger, pernitidus. Scutellum subtiliter dense punctatum, nigrum, densius griseo-pubescent. Elytra thorace vix longiora, minus confertim fortius punctata, stria suturali impressa, pube parca longiore nigra vestita, splendide viridimetallica. Abdomen crebre fortius punctatum, pube parca longiore fortiore depressa nigra vestitum, nigrum, supra cupreo-versicolor, segmento sexto toto, quinto apice rufis. Pectus nigrum. Pedes nigri, griseo-pubescentes, tarsis piceis, tibiis posterioribus spinulosis.

Tarsi antici feminae modice dilatati. Mas latet.

- \* 139. *Ph. compressicollis*: *Elongatus, niger, nitidus, elytris chalybeis, thorace elongato, convexo, utrinque subtiliter punctato, capite orbiculato.* — Long.  $4\frac{1}{2}$  lin.

*Staph. compressicollis* Klug Col. Madag. 50. 41. t. 1. f. 12.

Habitat in Madagascar, Dom. Goudot.

Statura elongata et facies fere Paederi, niger, nitidus, elytris chalybeis, nitidis. Antennae capite thoraceque paulo breviores, apicem versus haud crassiores, articulo tertio secundo sesqui-prope longiore, 4—10 sensim paulo brevioribus et paulo magis obconicis, omnibus crassitie multo longioribus, ultimo oblongo-ovato, apice extrorsum acuminato, rufo-ferrugineae, articulo basali piceo. Palpi rufi. Caput thorace paulo latius, orbiculatum, supra parum convexum, subtiliter punctulatum, punctis in media fronte rerioribus. Thorax coleopteris duplo angustior, latitudine duplo longior, basin versus subangustatus, lateribus antice cum angulis anterioribus leviter rotundatis, pone medium subsinuatis, basi cum angulis posterioribus leviter rotundatus, fortius convexus, subtiliter crebrius punctatus, spatio medio longitudinali angusto laevi. Scutellum nigrum, creberrime subtiliter punctulatum. Elytra thoracis longitudine, convexiuscula, subtilius minus dense punctulata, tenuiter cinereo-pubescentia. Abdomen parce cinereo-pubescent, supra segmentis singulis basi subtiliter punctatis, apice sublaevibus, subtus segmentis basi anterioribus fortiter, posterioribus sensim subtilius punctatis. Pedes nigri, tarsis piceis, coxis intermediis approximatis, tibiis omnibus parce spinulosis.

Tarsi antici maris fortius dilatati. Femina latet.



- \* 140. *Ph. cinerascens*: *Niger, antennarum basi pedibusque testaceis, capite thoraceque utrinque crebre punctatis, elytris abdomineque subtilissime confertissimeque punctatis, subtiliter cinereo-pubescentibus.* — Long. 2 lin.

*Nordm. Symb.* 108. 97. — *Er. Col. March. I.* 478. 43.

*Staph. cinerascens* *Grav. Micr.* 49. 74. *Mon.* 117. 137. — *Latr. Hist. nat. Crust. et Ins.* IX. 334. 94. — *Gyll. Ins. Suec. II.* 361. 75. — *Sahlb. Ins. Fenn. I.* 336. 74. — *Mannerh. Brachelytr.* 31. 80.

Habitat in Europa.

Elongatus, linearis, subdepressus, niger, capite thoraceque nitidulis, elytris abdomineque opacis. Antennae capite paulo longiores, articulis duobus primis crassiusculis, tertio secundo longitudine aequali, 4—10 sensim paulo brevioribus et crassioribus, subobconicis, penultimo crassitie haud longiore, ultimo oblongo, apice oblique truncato et inferne acuminato, fuscae, articulo primo apiceque plerumque testaceis. Palpi maxillares articulo tertio brevi, quarto duplo longiore, fortiter acuminato. Os testaceum. Caput thoracis latitudine, oblongo-subquadratum, supra leviter convexum, confertissime subtiliter punctatum, linea angustiore media longitudinali laevi, basi apiceque evanescente. Thorax coleopteris angustior, latitudine tertia parte longior, lateribus parallelis, ante medium cum angulis anterioribus rotundatis, apice truncatus, basi rotundatis, angulis posterioribus obtusis, leviter convexus, crebrius punctatus, spatio medio angusto per totam longitudinem laevi. Elytra thorace tertia parte longiora, scutellum et abdomen confertissime subtiliter punctata, pube brevissima subtili cinerea dense vestita, abdominis segmentis omnibus apice ferrugineo-marginatis. Pedes obscure testacei, coxis omnibus nigris, intermediis approximatis, tibiis omnibus parce spinulosis, tarsis anticis in utroque sexu leviter dilatatis.

- \* 141. *Ph. sericeus*: *Elongatus, niger, opacus, subtiliter cinereo-pubescent, antennis pedibusque piceis, capite thoraceque utrinque crebre, elytris abdomineque subtilissime creberrimeque punctatis.* — Long. 2½ lin.

*Remus sericeus* *Holme Transact. Ent. Soc. Lond. II.* p. 58.

Habitat Lutetiae, Dom. Aubé, in Anglia, Dom. Rudd.

Elongatus, linearis, subdepressus, corpore toto cinerascanti-nigro, opaco. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus et crassioribus, subobconicis, penultimo crassitie vix longiore, ultimo oblongo, apice truncato et inferne acuminato, ferrugineae, articulis tribus primis nigro-piceis. Palpi maxillares articulo quarto tertio sesqui longiore, modice acuminato. Os piceum. Caput thoracis latitudine, oblongo-subquadratum, subdepressum, utrinque subtilius licet profunde sat crebre punctatum, linea media longitudinali angusta integra laevi, antierius tenuiter obsoleteque canaliculata, subtiliter tenuiterque cinereo-pubescent. Thorax coleopteris plus dimidio angustior, latitudine tertia parte longior, lateribus parallelis, rectis, basi rotundatus, apice truncatus, angulis anterioribus fortiter rotundatis, posterioribus obtusis, subdepressus, dense subtiliter punctatus, linea media longitudinali laevi, postice subelevata, subtiliter cinereo-pubescent. Elytra thorace paulo longiora, scutellumque confertim subtilissime punctata, pube subtili densa cinerea ob-

ducta. Abdomen confertim subtilissime punctatum, dense subtiliter cinereo-pubescentibus. Pedes picei, geniculis tarsisque piceo-testaceis, coxis intermediis approximatis, tibiis omnibus parcius spinulosis, tarsis anticis in utroque sexu leviter dilatatis.

Obs. Affinis praecedenti, statura autem magis elongata, capite thoraceque opacis, pubescentibus, hoc dense punctato etc. valde distinctus. Palporum maxillarium articulus tertius minus brevis, quartus minus elongatus et minus acutus.

- 142. *Ph. pruinus*: *Elongatus, niger, antennis rufo-testaceis, basi piceis, pedibus piceo-testaceis, capite thoraceque utrinque creberrime punctatis, elytris abdomineque subtiliter confertissimeque punctatis, subtiliter dense cinereo-pubescentibus.* — Long.  $2\frac{1}{2}$  lin.**

Habitat in Sardinia, Mus. Reg. Taurin., a Dom. Prof. Gené ad examinandum communicatus.

Statura omnino *Ph. cinerei*, capite autem maiore discedens, niger, vix nitidus, corpore pube subtili depressa cinerea subsericante dense vestito, capite thoraceque parce cinereo-pubescentibus. Antennae capite thoraceque paulo breviores, rufo-testaceae, articulis tribus primis nigro-piceis. Caput thorace paulo latius, subquadratum, depressum, dense profundius punctatum, linea media longitudinali laevi, medio subtiliter canaliculata, antice inter antennas, leviter excavatum, ore rufo, mandibulis validioribus prominentibus. Thorax latitudine paulo longior, basin versus subangustatus, basi coleopteris vix dimidio angustior, minus subtiliter confertim punctatus, linea media leviter elevata laevi. Scutellum confertissime subtiliter punctatum. Elytra thorace vix longiora, depressa, confertissime subtiliter punctata. Abdomen confertim subtiliter punctulatum. Pedes piceo-testacei.

Tarsi antici maris leviter dilatati. Femina latet.

Obs. Affinis *Ph. cinerascens*, at paulo maior, et capite thoraceque creberrime punctatis distinctus, *Ph. sericeo* propior, at capite thoraceque parce cinereo-pubescentibus elytrisque brevioribus etc. discedens.

- \* 143. *Ph. prolixus*: *Elongatus, niger, antennarum basi pedibusque testaceis, capite thoraceque utrinque crebre punctatis, elytris distincte punctatis, rufo-testaceis, basi nigricantibus.* — Long. 2 lin.**

Habitat in Austria, Dom. Ullrich.

*Elongatus*, linearis, subdepressus, niger, nitidus. Antennae capite plus sesqui longiores, articulo secundo tertioque subaequalibus, 4 — 10 sensim paululum crassioribus at vix brevioribus, penultimis crassitie subbrevioribus, ultimo ovato, apice truncato et inferne subacuminato, fuscae, articulis duobus basalibus rufo-testaceis, tertio rufo-piceo. Palpi maxillares articulo quarto tertio haud longiore, conico, acuminato, acuto. Os rufo-testaceum. Caput thorace fere latius, oblongum, subdepressum, crebre et sat profunde punctatum, relicto spatio medio longitudinali angusto, apice dilatato, laevi. Thorax coleopteris plus dimidio angustior, longitudine plus sesqui longior, basin versus subangustatus, lateribus subrectis, basi leviter rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, parum convexus, utrinque crebre aequaliter profundius punctatus, spatio medio longitudinali angusto laevissimo. Scutellum subtiliter punctulatum, nigrum. Elytra thorace sesqui longiora, distincte at minus dense punctata,

tenuiter minus subtiliter griseo-pubescentia, saturate rufo-testacea, basi usque ad tertiam elytri partem et lateribus nigricantia. Abdomen parcius subtiliter punctulatum, segmentis anterioribus basi transversim depressis et fortius punctatis, densius griseo-pubescentibus. Pedes rufo-testacei, coxis anterioribus picescentibus, intermediis approximatis, tibiis omnibus parce spinulosi, tarsis anticis in utroque sexu leviter dilatatis.

Obs. Affinis Ph. procerulo, differt autem statura magis elongata, capite oblongo, elytris ultra medium rufis.

\* 144. *Ph. procerulus*: *Elongatus, niger, antennarum basi pedibusque fusco-testaceis, thorace utrinque capiteque punctatis, elytris nigro-piceis, apice subrufescentibus.* — Long.  $1\frac{1}{2}$  — 2 lin.

*Er. Col. March. I.* 479. 49.

*Staph. procerulus* Grav. Mon. 95. 92. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 409. 45.

*Gyrophyn. procerulus* Mannerh. *Brachelytr.* 34. 16. — *Nordm. Symb.* 120. 22.

*Staph. planatus* Grav. Mon. 108. 112. (*forte*)

Habitat in Germania.

Statura fere Ph. cinerascens, at paulo angustior, niger, capite thoraceque nitidis. Antennae capitis thoracisque fere longitudine, validiusculae, articulo secundo tertioque subaequalibus, 4 — 10 sensim paulo crassioribus at vix brevioribus, penultimis crassitie paulo brevioribus, ultimo ovato, apice inferno acuminato, nigricantes, articulis tribus primis fusco-testaceis. Palpi maxillares articulo quarto tertio duplo longiore, conico, fortiter acuminato, acutissimo. Os testaceum. Caput thoracis latitudine, oblongo-subquadratum, parum convexum, fortiter crebrius punctatum, spatio medio medio longitudinali angustiore laevi, antice leviter impresso. Thorax coleopteris dimidio angustior, latitudine sesqui longior, basin versus subangustatus, lateribus rectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrectis, parum convexus, fortiter crebrius punctatus, spatio medio longitudinali angusto laevissimo, basi subelevato. Elytra thorace tertia parte longiora, subtiliter licet distincte densius punctata, minus subtiliter griseo-pubescentia, nigro-picea, margine apicali piceo-testaceo. Abdomen crebrius subtiliter punctulatum, densius griseo-pubescentibus, segmentis singulis apice piceo-marginatis. Pedes obscure testacei, coxis anterioribus fuscis, intermediis approximatis, tibiis omnibus parcius spinulosi, tarsis anticis in utroque sexu leviter dilatatis. Alae albo-hyalinae.

Obs. *Staph. planatum* Grav. non reperi in collectione Hellwig-Hoffmannseggiana, unde secundum descriptionem citatam non sine ullo dubio huc eum retuli.

\* 145. *Ph. elongatulus*: *Elongatus, niger, antennis pedibusque testaceis, capite thoraceque utrinque vage punctatis, elytris piceo-testaceis, distincte punctatis.* — Long.  $1\frac{2}{3}$  lin.

*Er. Col. March. I.* 479. 50.

Habitat in Germania.

Praecedenti affinis, niger, capite thoraceque nitidis. Antennae capitis thoracisque prope longitudine, articulo tertio secundo tenuiore et vix lon-



gitudine ei aequali, tertio quartoque subcylindricis, crassitie paulo longioribus, 5—10 crassioribus, transversim subobconicis, ultimo ovato, apice truncatulo et inferne obtuse acuminato, laete testaceae. Caput thorace paulo latius, depressiusculum, subquadratum, lateribus fortiter minus dense punctatum, spatio medio longitudinali late laevi, antice subimpresso. Thorax coleopteris dimidio angustior, latitudine tertia parte longior, basin versus subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus fortiter rotundatis, parum convexus, fortiter minus confertim subseriatim punctatus, spatio medio longitudinali angustiore laevissimo. Scutellum parce subtilissime punctulatum. Elytra thorace paulo longiora, minus confertim subtiliter licet distincte punctata, minus subtiliter griseo-pubescentia, fusco-picea, apice sensim dilutiora, testacea. Abdomen crebrius subtiliter punctulatum, densius griseo-pubescentis, segmentis singulis apice supra tenuiter, infra laete ferrugineo-marginatis. Pedes laete testacei, coxis anterioribus concoloribus, intermediis approximatis, tibiis omnibus subtiliter spinulosis. Alae albo-hyalinae.

Tarsi antici maris fortius, feminae leviter dilatatae.

- \* 146. *Ph. sobrinus*: *Elongatus, niger, antennarum basi apiceque pedibusque testaceis, capite thoraceque utrinque multi-punctatis, elytris distincte punctatis, margine apicali testaceo. — Long. 2 lin.*

Habitat in Pennsylvania, Dom. Zimmermann.

*Elongatus*, minus, depressus, niger, nitidus. Antennae capitis thoracisque fere longitudine, articulo tertio secundo paulo longiore et tenuiore, 4—10 sensim crassioribus at vix brevioribus, penultimo crassitie haud longiore, ultimo oblongo, apice subtruncato et inferne acuminato, fuscae, apice ferrugineae, articulis basalibus tribus flavo-testaceis. Palpi maxillares articulo quarto tertioque sesqui longiore, conico, apice fortiter acuminato acutoque. Os testaceum. Caput thoracis latitudine suborbiculatum, leviter convexum, parce punctatum, spatio medio longitudinali minus lato laevi, antice inter antennis foveola impressa. Thorax coleopteris dimidio angustior, latitudine tertia parte longior, basin versus subangustatus, lateribus ante medium leviter rotundatus, pone medium subsinuatus, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrotundatis, leviter convexus, utrinque punctatus, punctis minus crebris, hinc inde subseriatis, spatio medio longitudinali angusto laevissimo. Scutellum parce subtiliterque punctulatum, nigrum. Elytra thorace paulo longiora, distinctè parcius punctata, minus subtiliter griseo-pubescentia, nigro-picea, margine apicali indeterminate flavo-testaceo. Abdomen griseo-pubescentis, segmentis singulis basi crebrius, apice parcius obsoletiusque punctulatis, margine apicali piceis. Pedes cum coxis anterioribus testaceis, coxis intermediis approximatis, tibiis omnibus parce subtiliterque spinulosis.

Tarsi antici maris fortius, feminae leviter dilatati.

- \* 147. *Ph. humilis*: *Elongatus, niger, antennarum basi pedibusque testaceis, capite thoraceque utrinque crebre punctatis, elytris distincte punctatis, margine apicali testaceo. — Long. 1½ lin.*

Habitat in ins. Puerto Rico, Dom. Moritz.

Statura omnino praecedentis, at dimidio minor, niger, nitidus. Antennae capite thoraceque paulo breviores, articulo secundo tertioque subaequalibus, 4 — 10 sensim paulo crassioribus, at vix brevioribus, penultimis crassitie paulo brevioribus, ultimo oblongo, apice truncatulo et inferne subacuminato, nigro-fuscae, articulis tribus basalibus testaceis. Palpi maxillares articulo quarto tertio sesqui longiore, conico, acuminato, acuto. Os testaceum. Caput thoracis latitudine, orbiculatum, crebre profunde punctatum, spatio medio longitudinali angusto laevi, antice foveola minuta obsoleta notatum. Thorax coleopteris dimidio angustior, latitudine sesqui prope longior, basin versus subangustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice utrinque oblique truncatus, angulis anterioribus obtusiusculis, parum convexus, utrinque sat crebre profunde aequaliter punctatus, spatio medio longitudinali angusto laevisimo. Scutellum parce subtiliterque punctulatum, nigrum. Elytra thorace paulo longiora, parcius subtiliterque at distincte punctata, tenuiter griseo-pubescentia, nigra, margine apicali determinate testaceo. Abdomen densius subtiliter punctulatum, densius griseo-pubescentia, segmentorum marginibus rufo-piceis. Pedes cum coxis anterioribus testacei, coxis intermediis approximatis, tibiis omnibus parce spinulosis, tarsis anticis in utroque sexu fortius dilatatis. Alae albo-hyalinae.

\* 148. *Ph. agnatus*: *Elongatus, subdepressus, nig. r., antennarum basi pedibusque testaceis, capite thoraceque utrinque crebrius punctatis, punctis thoracis subseriatis, elytris distincte punctatis, margine apicali testaceo. — Long. 2 lin.*

Habitat in valle Araguensi Columbiae, Dom. Moritz.

*Ph. humili* paulo maior, niger, nitidus. Antennae capitis thoracisque prope longitudine, crassiusculae, articulo tertio secundo paulo longiore et tenuiore, 4 — 10 sensim crassioribus at vix brevioribus, penultimis crassitie paulo brevioribus, ultimo ovato, apice subemarginato et inferne acuminato, nigro-fuscae, basi testaceae. Palpi maxillares articulo ultimo tertio duplo longiore, conico, apice fortiter acuminato, acutissimo. Os testaceum. Caput thoracis latitudine, subovatum, crebre punctatum, spatio medio longitudinali angustiore laevi, antice obsolete impresso. Thorax coleopteris dimidio prope angustior, latitudine tertia parte longior, basin versus subangustatus, lateribus ante medium leviter rotundatis, pone medium subsinuatis, basi rotundatus, apice utrinque oblique truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, leviter convexus, crebrius subseriatim punctatus, spatio medio longitudinali angusto laevi. Scutellum parce subtiliterque punctulatum. Elytra thorace paulo longiora, minus dense subtilius punctata, tenuiter griseo-pubescentia, nigra, margine apicali determinate testaceo. Abdomen parce subtilissime punctatum, longius griseo-pubescentia, segmentorum marginibus rufo-piceis. Pedes flavi, coxis anterioribus concoloribus, intermediis approximatis, tibiis omnibus parce subtiliterque spinulosis.

Tarsi antiqui maris modice dilatati. Femina latet.

Obs. *Ph. humili* proximus, differt autem statura maiore, thorace lateribus minus crebre et subseriatim punctato, abdomine vix perspicue punctato.



149. *Ph. ludicrus*: *Elongatus, niger, nitidus, pedibus antennisque testaceis, his medio fuscis, capite thoraceque utrinque crebre punctatis, hoc rufo, disco fusco, elytris distincte punctatis, margine apicali testaceo.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Guyana Venezuelensi, Dom. Moritz.

Statura et summa affinitas praecedentium. Antennae capite thoraceque breviores, crassiusculae, articulo tertio secundo paulo brevior et tenuiore, 4 — 10 sensim crassioribus at paulo brevioribus, subobconicis, penultimis crassitie brevioribus, ultimo ovato, apice emarginato et inferne acuminato, nigrae, articulis tribus primis flavo-testaceis. Palpi maxillares articulo quarto tertio duplo longiore, conico, apice fortiter acuminato, acutissimo. Os testaceum. Caput thoracis latitudine, leviter convexum, subquadratum, crebre profundeque punctatum, spatio medio longitudinali angustiore laevissimo, nigrum, nitidum. Thorax latitudine sesqui prope longior, coleopteris dimidio angustior, basin versus subangustatus, lateribus subrectis, basi cum angulis posterioribus leviter rotundatus, apice utrinque oblique truncatus, angulis anterioribus subrotundatis, sat crebre profundeque punctatus, spatio medio longitudinali angusto laevissimo, nitidus, piceus, limbo omni indeterminate rufo. Scutellum parce subtiliter punctulatum, nigro-piceum, nitidum. Elytra thorace tertia parte longiora, minus dense subtilius punctata, tenuiter griseo-pubescentia, nigra, nitida, margine apicali determinate testaceo. Abdomen subtiliter punctulatum, griseo-pubescent, nitidulum, nigrum, segmentis singulis piceo-marginatis, subtus cum pectore rufo-piceum. Pedes flavi, coxis anterioribus concoloribus, intermediis leviter distantibus, tibiis omnibus parce subtiliterque spinulosis, tarsis anticis in utroque sexu modice dilatatis. Alae albo-hyalinae.

Variat antennis testaceis, articulis 3 — 6 fusciscentibus.

Obs. *Ph. humili* affinis, at magis elongatus et coxae intermediae distantes.

- \*150. *Ph. palmula*: *Elongatus, niger, antennis pedibusque testaceis, elytris abdomineque rufis, hoc ante apicem nigro, capite thoraceque utrinque crebre punctatis.* — Long.  $2\frac{1}{2}$  lin.

*Staph. palmula* Grav. Micr. 49. 73. Mon. 107. 110.

*Staph. cinctus* Latr. Hist. nat. Crust. et Ins. IX. 333. 92.

Habitat in Germania, Knoch.

Elongatus, subdepressus. Antennae capite sesqui longiores, articulo tertio secundo brevior et tenuiore, 4 — 10 sensim crassioribus, vix autem brevioribus, 6 — 10 sensim magis transversis, ultimo apice truncatulo et inferne acuminato, totae testaceae. Palpi maxillares articulo quarto tertio sesqui longiore, conico, fortiter acuminato et acuto. Os testaceum. Caput thorace paululum latius, subquadratum, depressiusculum, profunde minus crebre punctatum, spatio medio longitudinali laevi, antice dilatato et leviter impresso, parce griseo-pubescent, nigrum, nitidum. Thorax coleopteris dimidio angustior, latitudine paulo longior, lateribus parallelis, rectis, basi rotundatus, apice utrinque oblique truncatus, angulis anterioribus subrotundatis, posterioribus obtusis, parum convexus, utrinque fortius crebriusque vage punctatus, linea media lata longitudinali laevi, nigro-piceus, nitidus. Scutellum parce subtilissime punctatum, rufo-piceum. Elytra thorace paulo



longiora, crebrius subtiliusque punctata, tenuiter pubescentia, rufa, margine apicali testaceo. Abdomen basi coleopterorum latitudine, apicem versus leniter sensim incrassatum, tenuiter griseo-pubescent, supra laeve, subtus parce punctatum, rufum, nitidum, segmentis penultimis duobus nigris, apice ultimoque toto ferrugineis. Pectus nigro-piceum. Pedes testacei, coxis intermediis modice distantibus, tibiis spinulosus.

Tarsi antici maris leviter dilatati. Femina latet.

Obs. Corporis antrorsum angustati habitu inter affines insignis.

### **Heterothops Kirby.**

*Steph. Illustr. V. p. 256. — Er. Col. March. I. p. 480*

*Trichopygus Nordm. Symb. p. 137. t. 2. f. 5.*

*Antennae rectae.*

*Ligula rotundata, integra.*

*Palpi articulo ultimo subulato.*

Corpus minutum, oblongum, utrinque attenuatum, fere fusiforme, alatum. Caput thorace angustius minusque, ovatum, basi leviter constrictum, deflexum, oculis mediocribus, parum prominulis. Labrum transversum, apice setosum, medio incisum. Mandibulae breves, medio obtuse dentatae. Maxillae (t. IV. f. 13. b.) malis membranaceis, interiore intus, exteriore apice barbatis. Palpi maxillares modice elongati, articulo primo minuto, secundo elongato, tertio hoc longitudine aequali, crassiusculo, subovato, quarto minuto, subulato. Labium (t. IV. f. 13. a.) mento transversum, ligula membranacea, rotundata, paraglossis membranaceis, intus ciliatis, divergentibus, ligulam superantibus. Palpi labiales articulo primo minore, secundo sesqui longiore, crasso, cylindrico, tertio parvo, subulato. Antennae rectae, filiformes, apicem versus haud incrassatae, articulo ultimo apice truncatulo et inferne subacuminato. Thorax transversim convexus, basi fere coleopterorum latitudine, antrorsum leviter angustatus, basi fortiter rotundatus, apice truncatus. Scutellum triangulare. Elytra truncata. Abdomen apicem versus sensim attenuatum, vivis sat elongatum, stylis analibus longioribus exsertis. Pedes breviusculi, coxis intermediis contiguus, tibiis anticis muticis, posterioribus spinulosus, tarsis anticis articulis quatuor primis dilatatis, subtus spongioso-tomentosis, posterioribus setulosis, articulo primo ultimoque leviter elongatis.

### **Observationes.**

1. Corpus punctulatum, pubescens, capite thoraceque laevissimis, glabris, illo puncto singulo ad utriusque oculi marginem interiorem, nonnullis supra oculos et duobus verticalibus ante stricturam transversim positis, hoc punctis praeter pauca marginalia binis dorsalibus, paulo ante medium, et binis ad apicem in omnibus notatis. Habitu proxime ad Quedios minutos accedunt, a quibus autem stigmate prothoracico detecto et palporum structura differunt.

2. Mas abdominis segmento inferiore sexto apice leviter emarginato, tarsisque anticis fortius dilatatis ab altero sexu distinguitur.

3. Victus in quisquiliis, sub musco et sub foliis deciduis.

\* 1. *H. praevious*: Niger, pedibus rufo-piceis, capite breviter ovato, elytris thorace paulo longioribus, fusco-piceis, apice rufescentibus. — Long. 2 lin.

Er. Col. March. I. 480. 1.

*Trichopygus dissimilis* Nordm. Symb. 137. 2.

*Emus subuliformis* Boisd. et Lacord. Faun. Ent. Paris. I. 385. 42.

Habitat in Germania.

Niger, capite thoraceque politis. Antennae capite sesqui longiores, articulo tertio secundo paulo minore, 4—10 subobconicis, sensim paulo brevioribus, penultimis crassitie haud longioribus, ultimo oblongo-ovato, apice emarginato et inferne acuminato, fusco-nigrae, basi obscure rufo-piceae. Palpi picei. Caput thorace angustius, breviter ovatum, punctis verticalibus duobus, utrinque supra oculos tribus, longitudinaliter dispositis, profundioribus, inter oculos utrinque puncto singulo ad ipsius oculi marginem interiori-rem impressum. Thorax basi coleopterorum prope latitudine, latitudine baseos fere brevior, antrorsum angustatus, apice dimidio quam basi angustior, lateribus rectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, convexus, dorso ante medium punctorum pari et in ipso margine anteriore punctis duobus subtilioribus et paulo magis distantibus notatus, nigro-piceus, nitidus. Scutellum subtiliter punctulatum. Elytra thorace paulo longiora, densius punctulata, longius pubescentia, fusco-picea, apice rufescentia. Abdomen punctulatum, pube depressa fusco-grisea sericans, nigro-piceum, segmentorum marginibus apiceque rufo-ferrugineis. Pedes cum coxis rufo-picei.

Obs. Differt a *H. dissimili* statura paulo maiore, capite minus angusto, elytris longioribus, antennis brevioribus et crassioribus.

\* 2. *H. binotatus*: Niger, nitidus, antennarum basi pedibusque testaceis, capite suborbiculato, elytris thorace paulo longioribus, nigro-piceis, margine summo apicali subtestaceo. — Long.  $2\frac{1}{3}$  lin.

Habitat in Anglia, Dom. Rudd.

Praecedentis maximis individuis aequalis, niger, capite thoraceque politis. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, 4—10 subcylindricis, sensim paulo brevioribus, penultimis etiam crassitie longioribus, ultimo oblongo, apice leviter emarginato et inferne subacuminato, piceae, articulis tribus primis testaceis. Palpi nigro-picei. Caput thorace paulo angustius, suborbiculatum vel breviter ovatum, depressiusculum, punctis utrinque tribus longitudinaliter dispositis supra oculos, singulo inter utrumque oculum, fortioribus, duobus transversim positis verticalibus subtilioribus impressum. Thorax basi coleopterorum prope latitudine, latitudine baseos paulo brevior, antrorsum modice angustatus, apice vix tertia parte quam basi angustior, lateribus leviter, basi fortius rotundatus, apice truncatus, angulis anterioribus leviter deflexis posterioribusque subrotundatis, leviter convexus, dorso ante medium et apice punctis binis notatus, apicalibus subtilioribus et magis distantibus, nigro-piceus, nitidus. Scutellum crebre subtiliter punctulatum, tenuiter nigro-pubescentia. Elytra thorace paulo longiora, densius punctulata, nigro-pubescentia, nigro-picea, margine summo apicali piceo-testaceo. Scutellum dense subtiliter punctulatum, densius griseo-pubescentia, segmenti

sexti margine apicali testaceo, segmentorum ventralium reliquorum ferrugineo. Pedes testacei, coxis piceis.

Obs. Differt a praecedente thorace minus convexo, antrorsum minus angustato, antennis basi testaceis. Teste Dom. Rudd sub nomine eodem Steph. Illustr. V. p. 257 descriptus.

- \* 3. *H. dissimilis*: Niger, antennarum basi pedibusque testaceis, elytris thoracis longitudine, piceis, apice rufescentibus, capite oblongo-ovato. — Long.  $1\frac{3}{4}$  lin.

Er. Col. March. I. 480. 2.

*Tachyporus dissimilis* Grav. Micr. 125. 1. — Latr. Hist. nat. Crust. et Ins. X. 5. 1.

*Tachinus dissimilis* Grav. Mon. 23. 1.

*Staph. subuliformis* Gyll. Ins. Suec. II. 312. 29. — Sahlb. Ins. Fenn. I. 319. 27.

— Mannerh. Brachelytr. 27. 40.

Habitat in Germania, Suecia.

Niger, capite thoraceque politis. Antennae capite cum thorace paululum breviores, articulo tertio secundo paulo minore, 4 — 10 subcylindricis, sensim vix brevioribus, omnibus crassitie longioribus, ultimo oblongo, apice truncatulo et inferne subacuminato, fusco-nigrae, articulis duobus primis testaceis. Palpi testacei. Caput thorace duplo angustius, elongatum, punctis verticalibus duobus, supra oculos utrinque tribus longitudinaliter dispositis, et ad oculi utriusque marginem interiori puncto singulo subtilioribus impressum. Thorax coleopterorum fere latitudine, latitudine baseos haud brevior, antrorsum angustatus, antice tertia parte quam basi angustior, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, convexus, dorso ante medium et apice bipunctatus, punctis apicalibus haud subtilioribus at magis distantibus, nigro-piceus, nitidus. Scutellum punctulatum. Elytra thorace haud longiora, densius punctulata, longius pubescentia. Abdomen sat crebre punctulatum, apice segmentorumque singulorum margine apicali piceis infra rufis. Pedes testacei, coxis anticis concoloribus.

Variat aetate minus maturus thorace elytrisque rufo-piceis.

- \* 4. *H. 4-punctulus*: Niger, antennis concoloribus, pedibus piceo-testaceis, elytris thoracis longitudine, capite ovato. — Long.  $1\frac{3}{4}$  lin.

Er. Col. March. I. 481. 3.

*Tachinus 4-punctulus* Grav. Mon. 24. 2.

*Staph. subuliformis* Zetterst. Faun. Lappon. I. 72. 14. Ins. Lapp. 62. 20.

*Trichopygus punilio* Nordm. Symb. 138. 4.

Habitat in Germania.

Statura et summa affinitas praecedentis, niger, nitidulus, capite thoraceque politis. Antennae capitis thoracisque prope longitudine, articulo tertio secundoque subaequalibus, 4 — 10 subcylindricis, sensim paulo brevioribus, penultimis crassitie vix longioribus, ultimo oblongo, apice truncatulo et inferne subacuminato, piceo-nigrae, basi concolores. Palpi nigro-picei. Caput thorace duplo prope angustius, ovatum, supra oculos utrinque punctis tribus longitudinaliter dispositis, singulo ad oculi utriusque marginem interiori et duobus transversim positus verticalibus impressum. Thorax



basi coleopteris paulo angustior, latitudine baseos paulo longior, antrorsum leniter angustatus, apice vix quarta parte quam basi angustior, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, modice transversim convexus, dorso ante medium et apice bipunctatus, punctis apicalibus magis distantibus. Scutellum subtiliter punctulatum. Elytra thorace vix longiora, subtiliter parcius punctulata, parcius pubescentia, concolora. Abdomen subtiliter punctulatum, parcius pubescens, subtus apice piceum. Pedes piceo-testacei, coxis anticis concoloribus.

Obs. Differt a praecedente antennis elytrisque concoloribus, capite minus oblongo, elytris abdomineque subtilius punctulatis. Specimen in collectione Hoffmanns-eggiana pro typico asservatum (a Dahllo Brunsvigae lectum), ad speciem praecedentem pertinet: cui autem descriptio Gravenhorstiana contraria est. Quae igitur denominatio errore quodam exorta videtur, quod in speciebus adeo difficulter distinguendis facile accidere potuit, eamque retinere eo magis dubitavi, cum cel. Gravenhorst collectionem laudatam non citaverit.

- \* 5. *H. exilis*: Niger, ore, antennis pedibusque flavis, elytris thorace longioribus, capite breviter ovato. — Long.  $1\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Habitu omnino praecedentium, at duplo minor, niger, capite thoraceque politis. Antennae capitis thoracisque prope longitudine, articulo secundo crassiusculo, tertio duplo prope longiore, 4—10 subobconicis, sensim paulo brevioribus, omnibus crassitie longioribus, ultimo paulo crassiore, ovato apice subtruncato et inferne subacuminato, luteae, basi flavae. Os flavum. Caput breviter ovatum, prope oculi utriusque marginem superiorem et in margine oculi utriusque interiore, puncto singulo, vertice punctis 4 transversim positis subtilioribus impressum. Oculi sat magni, modice prominuli. Thorax basi coleopterorum latitudine, latitudine baseos vix brevior, antrorsum angustatus, apice tertia parte quam basi angustior, lateribus subrectis, basi rotundatus, apice truncatus, angulis anterioribus deflexis posterioribusque subrotundatis, modice convexus, dorso ante medium et apice bipunctatus, punctis apicalibus magis distantibus. Scutellum punctulatum. Elytra thorace perparum longiora, densius subtiliter punctulata, parce pubescentia, margine summo apicali rufescente. Abdomen densius punctulatum, apice marginibusque tenuibus segmentorum singulorum rufescentibus. Pedes flavi, coxis anterioribus concoloribus.

#### Species mihi invisae.

6. *H. nitens*: Convexior, niger, nitidissimus, antennis basi, segmentorum ventralium marginibus, uno pedilusque rufo-piceis, elytris ultra medium ab apice rufo-testaceis, nitidis. — Long.  $1\frac{1}{2}$  lin.

*Trichopygus nitens* Nordm. Symb. 138. 3.

Habitat in Chersoneso Taurica.

Praecedentibus similis et affinis, sed multo minor. Antennae fusco-piceae, articulis basalibus dilutioribus. Elytra convexa, laete rufo-testacea, versus scutellum paulo infuscata, nitidissima, subtilissime punctulata, fulvescenti-pubescentia. Pedes picei-ferruginei. (Nordm.)

**Acylophorus Nordm.**

Nordm. Symb. p. 127. t. 1. f. 19.

*Antennae fractae, articulo primo elongato.**Ligula rotundata, integra.*

Corpus minoris magnitudinis, fere fusciforme, alatum. Caput thorace multo minus, oblongo-ovatum, basi vix constrictum, inflexum, oculis haud prominulis. Labrum transversum, apice ciliatum, medio tenuiter incisum. Mandibulae parvae, acutae, basi fortiter bidentatae. Maxillae (t. IV. f. 9. b.) malis brevibus, coriaceis, apice barbatis. Palpi maxillares mediocres, articulo primo minuto, secundo elongato-obconico, tertio obconico. Labium (t. IV. f. 9. a.) mento transversum, corneo, ligula brevi, rotundata, coriacea, paraglossis minutis, ligulam haud superantibus. Palpi labiales articulo primo minuto, secundo obconico tertioque ovato vel subgloboso crassiusculis. Antennae tenuiusculae, fractae, articulo primo elongato, capite paulo brevior, apice subincrassatae, articulo ultimo globoso. Thorax basi coleopterorum latitudine, antrorsum angustatus, convexus, lateribus deflexis basique rotundatus, apice truncatus. Scutellum triangulare, sat magnuin. Elytra thoracis longitudine, truncata. Abdomen apicem versus sensim attenuatum, stylis analibus mediocribus, hirsutis. Pedes breviusculi, coxis intermediis contiguis, tibiis anticis muticis, posterioribus spinulosi, tarsis anticis breviusculis, crassiusculis, cylindricis, articulis parum distinctis, quatuor primis brevissimis, quinto his coniunctis fere longiore, unguiculis validis, tarsis posterioribus tenuiusculis, articulo primo elongato, quinto unguiculis minutis, debilibus.

**Observationes.**

1. Corpus punctulatum, pubescens, capite thoraceque laevissimis, glaberrimis, nonnisi punctis paucis notatis. Habitus huius generis, capite inflexo, antennis fractis et tarsorum anticarum structura in hac subtribui valde distincti, propius ad Tachyporinos accedit, neque etiam Gymnusa dissimilis videtur, corporis penitior structura autem et antennarum insertione huic tribui et subtribui adscribendum.

2. Palporum maxillarum structura species 5 mihi examinatae non omnino inter se congruunt.

3. Sexus differentia latet.

4. Victus speciei indigenae in silvis sub musco.

\* 1. *A. glabricollis*: Niger, nitidus, tarsis fulvis, elytris fortiter punctatis. — Long. 3 lin.

*Er. Col. March. I. 482. 1.*

*Staph. glabricollis* Boisd. et Lacord. Faun. Ent. Paris. I. 396. 20.

*Staph. rufilabris* Zetterst. Ins. Lappon. 65. 42.

*Acylophorus Ahrensii* Nordm. Symb. 127. 1.

Habitat in Europa.

Niger, nitidus, capite thoraceque politis. Antennae capitis thoracisque longitudine, articulo secundo tertio maiore, 4 — 10 sensim brevioribus et crassioribus, subobconicis, penultimis crassitie vix longioribus, ultimo subgloboso, apice compresso, obtuso, nigrae, articulo primo summa basi flavescens. Palpi nigro-picei, maxillares articulo ultimo ovato. Caput oblongum, lateribus supra oculos subtilissime punctulatum et subtiliter pubescens.

supra utrinque inter oculos puncto singulo fortiore, verticem versus punctis binis, oblique positis subtilioribus impressum. Thorax basi coleopterorum prope latitudine, latitudine baseos haud brevior, antrorsum angustatus, lateribus et basi modice rotundatus, apice truncatus, angulis anterioribus deflexis, posterioribus obtusis, modice convexus, dorso medio punctorum pari, et lateribus ante medium utrinque puncto singulo notatus. Scutellum parce punctatum parceque pubescens. Elytra thoracis longitudine, minus dense fortius punctata, pube parca depressa minus subtili nigra vestita. Abdomen parcius punctulatum, minus subtiliter longius pubescens, supra margine apicali segmenti quinti testaceo, subtus segmentorum 5 anteriorum piceo. Pedes nigri, tarsi fulvis. Alae fusco-hyalinae.

\* 2. *A. longicornis*: *Niger, nitidus, ano pedibusque rufis, elytris fortius punctatis*. — Long.  $3\frac{2}{3}$  lin.

Habitat in Cayenna, Dom. Buquet.

Praecedente sesqui fere maior, niger, nitidus. Antennae elongatae, capite thoraceque prope longiores, graciliores, articulo secundo tertioque subaequalibus, quarto sequente paulo minore, 5 — 8 sensim paulo brevioribus, omnibus teretibus, sequentibus duobus sensim brevioribus et crassioribus, subobconicis, ultimo subgloboso, praecedenti arctius applicato, nigrae, articulo primo basi rufo-piceo. Palpi rufi, maxillares articulo quarto cylindrico, apice acuminato. Caput ovatum, inter oculos puncto singulo utrinque ovatum. Thorax basi coleopterorum latitudine, latitudine baseos parum brevior, antrorsum modice angustatus, lateribus, basi et angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, modice convexus, punctorum pari altero in dorso paulo ante medium, altero ante apicem, et lateribus prope marginem ante medium puncto minuto singulo utrinque notatus. Scutellum punctatum, fusco-pubescens. Elytra thoracis longitudine, minus confertim fortius punctata, densius longiusque fusco-pubescentia. Abdomen minus confertim fortius punctulatum, fortiter nigro-pubescens, subtus coloribus lactis metallicis resplendens, segmento quinto sextoque apice testaceis. Pedes rufi, coxis nigris, anticiis interne rufescentibus.

\* 3. *A. femoralis*: *Niger, nitidus, pedibus piceis, femoribus tarsisque rufis, elytris parce fortius punctatis*. — Long.  $2\frac{3}{4}$  lin.

Nordm. *Symb.* 128. 2.

Habitat in Brasilia prope Rio Janeiro, Dom. v. Olfers.

*A. glabricolli* aequalis, niger, nitidus. Antennae capite thoraceque multo breviores, articulo secundo tertio duplo maiore, 3 — 10 sensim paulo brevioribus, 5 — 10 sensim paulo crassioribus, penultimis duobus transversis, ultimo subgloboso, piceae, articulo primo basi rufo. Mandibulae rufae. Palpi rufi, maxillares articulo ultimo incrassato, ovato, apice acuminato. Caput oblongo-ovatum, puncto singulo utrinque inter oculos impressum. Thorax basi coleopterorum latitudine, latitudine baseos paulo brevior, antrorsum leviter angustatus, basi, lateribus et angulis posterioribus rotundatus, apice leviter emarginatus, angulis anterioribus leviter deflexis, subrotundatis, modice convexus, dorso paulo ante medium et paulo ante apicem punctis binis, et lateribus prope marginem lateralem antierius puncto utrinque singulo



impressus. Scutellum laeve. Elytra thorace paulo longiora, minus confertim fortius punctata, pilo brevi crasso rigido declinato e quovis puncto enascente hispida. Abdomen parce subtilius punctatum, parcius nigro-pubesceus, segmentis inferioribus primis apice piceo-marginatis. Pedes femoribus tarsisque rufis, tibiis piceis.

- \* 4. *A. picipes*: *Niger, nitidus, pedibus piceis, elytris subtilius punctatis, griseo-pubescentibus.* — Long. 3 lin.

Habitat in Columbia, Dom. Moritz.

Statura omnino praecedentium, niger, subnitidus, capite thoraceque nitidissimis. Antennae capite thoraceque paulo breviores, apice leviter incrassatae, articulo secundo tertio paulo longiore at parum crassiore, quarto sequente brevior et tenuiore, 5 — 10 subobconicis, sensim brevioribus, 5 — 7 leniter, 8 — 10 fortius sensim crassioribus, penultimis duobus crassitie brevioribus, ultimo subgloboso, nigrae, articulo primo rufo, apice piceo. Palpi rufi, maxillares articulo ultimo conico, acuminato. Caput ovatum, lateribus supra oculos confertissime subtiliter punctulatum, dense griseo-pubesceus, supra inter oculos puncto singulo, verticem versus binis oblique positus utrinque impressum. Thorax basi coleopterorum latitudine, latitudine baseos paulo brevior, antrorsum angustatus, lateribus, basi et angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, modice convexus, laevis, dorso ante medium altero et ante apicem altero punctorum pari et lateribus prope marginem lateralem longe ante medium puncto utrinque singulo impressus. Scutellum punctulatum, griseo-pubesceus. Elytra thoracis longitudine, crebrius minus fortiter punctata, longius densiusque griseo-pubescentia. Abdomen punctulatum, nigro-griseoque pubesceus, metallico-versicolor, segmento penultimo margine apicali rufo-piceo. Pedes picei, anteriores rufescentes.

5. *A. pronus*: *Niger, nitidus, pedibus rufis, femoribus posterioribus piceis, elytris parcius subtiliusque punctatis, nigro-pubescentibus.* — Long. 2½ lin.

Habitat in America septentrionali, Mus. Dom. Chevrolat.

A. glabricolli dimidio minor, niger, nitidulus, capite thoraceque nitidissimis. Antennae articulo secundo tertio sesqui longiore, quarto quinto paulo minore, hoc cum sequentibus duobus terete, sensim paulo brevioribus, (articuli reliqui desunt), piceae, articulis singulis basi rufis. Palpi rufo-picei, maxillares articulo tertio ovato, apice acuminato. Caput oblongum, lateribus supra oculos subtilissime confertissimeque punctatum, inter oculos puncto singulo, verticem versus punctis binis oblique positus subtilioribus impressum. Thorax basi coleopterorum prope latitudine, latitudine baseos perparum brevior, antrorsum angustatus, lateribus basi et angulis posterioribus rotundatus, apice truncatus, apice leviter emarginatus, angulis anterioribus deflexis subrotundatis, convexus, anterieus subcompressus, punctorum pari, altero in medio dorsi, altero paulo ante apicem punctoque singulo utrinque prope marginem lateralem inter medium et apicem impressus. Scutellum sublaeve, sive parce subtiliter obsoleteque punctulatum. Elytra thorace vix longiora, parcius subtiliusque punctata, longius nigro-pubescentia. Abdomen parce subtilius punctulatum, longius nigro-pubesceus. Pedes rufi, coxis omnibus femoribusque posticis nigro-piceis.

Subtribus *tertia*.**O x y p o r i n i.***Stigmatica prothoracica oblecta.**Antennae basi distantes.*

Labrum transversum, bilobum, lobis rotundatis, lateribus membranaceum plerisque;  
 integrum aut subintegrum Qued. molochino, frontali, fuliginoso;  
 totum corneum, medio incisum Euryporo;  
 totum corneum, bilobum, lobis divergentibus Oxyporo.

Mandibulae haud prominentes, basi mediove dentatae plerisque;  
 muticae Astrapaeo;  
 fortiter prominentes, porrectae, decussatae, muticae Oxyporo.

Maxillae mala exteriore apice, interiore interne barbatis;  
 utraque coriacea, plerisque;  
 exteriore coriacea, interiore membranacea Oxyporo.

Palpi maxillares filiformes plerisque;  
 articulo quarto securiformi Astrapaeo.

Mentum corneum, transversum, apice truncatum, interstitio inter mentum et ligulam membranaceo, simplice plerisque;

brevissimum, apice late sinuatum, interstitio inter mentum et ligulam coriaceo, apice acute bidentato Oxyporo.

Ligula brevis vel brevissima;  
 integra Quedio, Europoro;  
 leviter incisa Astrapaeo;  
 elongata, profunde biloba, lobis acuminatis Oxyporo.

Paraglossae liberae, ligulam longe superantes plerisque;  
 nullae Oxyporo.

Palpi labiales filiformes Quedio;  
 articulo tertio securiformi Astrapaeo, Europoro;  
 articulo tertio lunato Oxyporo.

Antennae rectae, filiformes vel apicem versus leviter incrassatae, articulis tribus primis reliquis magis elongatis, glabris, ultimo apice saepius abrupte acuminato aut oblique truncato plerisque;

apicem versus dilatatae, compressae, articulis penultimis pluribus fortiter transversis Oxyporo.

Caput basi parum constrictum, collo quasi crasso thoraci insertum.

Thorax basi rotundatus, apice truncatus, margine laterali distincto Quedio, Astrapaeo, Euryporo;

inflexo, subtilissimo, indistincto Oxyporo.

Elytra truncata, sutura integerrima.

Abdomen plerumque apicem versus sensim attenuatum.

Pedes breviusculi, coxis intermediis  
 approximatis Quedio, Astrapaeo, Europoro;  
 late distantibus Oxyporo;

tibiis spinulosis Quedio, Astrapaeo, Euryporo;  
 muticis Oxyporo;  
 tarsis posterioribus articulis 4 primis decrescentibus, anticis  
 dilatatis Quedio, Astrapaeo;  
 simplicibus Euryporo, Oxyporo.

### Quedius Leach.

*Er. Col. March, I. p. 483.*

*Quedius* Steph. *Illustr. V. p. 215. Curt. Britt. Ent. XIV. t. 638. — Microsaurus*  
*Dej. Cat. p. 69. — Steph. Illustr. V. p. 435. — Velleius Mannerh. Brachelytr.*  
*p. 19. — Steph. Illustr. V. p. 201. — Boisd. et Lacord. Faun. Ent. Paris. I. p.*  
*356. — Raphirus Steph. Illustr. V. p. 241.*

*Palpi utrique filiformes.*

*Ligula rotundata, integra, paraglossis brevior.*

*Pedes intermedii approximati.*

Corpus plerumque mediocre, oblongum vel elongatum, alatum. Caput thorace minus, suborbiculatum vel subovatum, basi leviter constrictum, thoraci immissum, oculis minoribus maioribusve, minus magisve prominulis. Labrum transversum, bilobum vel medio incisum, nonnullis (e. g. *Q. molochino*) integrum, limbo laterali membranaceo. Mandibulae minores, haud prominentes, acutiusculae, intus bidenticulatae. Maxillae malis minutis, coriaceis, interiore intus, exteriore extus barbatis. Palpi maxillares parum elongati, filiformes, articulo primo minuto, secundo tertio paulo longiore, quarto tertio vel aequali vel plerumque brevior, vel interdum paulo longiore. Labium mento transversum, corneo, ligula perbrevis, rotundata, integra, membranea, paraglossis minutis, ligulam tamen superantibus, divergentibus, coriaceis, intus ciliatis. Antennae ante mandibularum basin insertae, filiformes, articulo ultimo subacuminato. Thorax coleopteris plerumque latitudine aequalis, suborbiculatus, apice truncatus. Scutellum triangulare. Elytra truncata. Abdomen plerumque apicem versus sensim angustatum. Pedes breviusculi, coxis intermediis contiguis, tibiis posterioribus spinulosis, tarsis anticis articulis quatuor primis dilatatis, patellam ovalem formantibus, subtus spongioso-tomentosis, posterioribus setosis, articulo primo leviter elongato.

### Observationes.

1. Proximum hoc genus, praecipue oris, antennarum pedumque structura, nec non corporis punctura et pubescentia videtur Philontho, ut praeter stigmata prothoracica obiecta vix ulla exstet nota distinctiva. Habitus capite basi minus constricto, thorace lateribus magis rotundato, abdomine saepius apicem versus sensim attenuato a Philontho aliquatenus distinctus et ad Tachyporinorum formam quodammodo vergit.

2. Genera plura ab Anglis auctoribus condita sunt, quae nonnisi levioribus et non satis constantibus notis temere distincta Quedio subiungo:

**Velleius**, a plerisque recentioribus auctoribus pro genere receptus, speciem singulam, statura et antennis serratis insignem continens. Antennarum figura sola in hac tribu generibus condendis non sufficit.

**Microsaurus**. Toto huic generi, Velleio excepto, Com. Dejean hoc nomen imposuit, quo Dom. Stephens lateralem a reliquis Quediis distinxit, quamvis nullo characterē ab iis scedere videtur.

**Raphirus** oculis magnis a Quedio genuino distinguitur, qua nota nonnullae quidem species satis insignes esse videntur, quam autem non satis constantem



reperio, ut sectionis etiam notam praebeat. Sectionis secundae species plurimae huc pertinent, e. g. *Q. attenuatus*, *boops*.

3. Mas abdominis segmento inferiore sexto apice emarginato a femina distinctus. Femina in nonnullis tarsis anticis minus dilatatis vel simplicibus a mare differt. Caput in utroque sexu plerumque aequale, in paucis speciebus (e. g. *Q. fulgido*) maribus maioribus maius, sicut in nonnullis *Philonthis*.

4. Victus in fimo, in quisquillis, sub musco, foliis deciduis, lapidibus.

5. *Q. fulgidi* larva a Dom. Bouché hoc modo descripta: „Linearis, posteriorius attenuata, supra leviter convexa, pallide flava, parce setosa. Caput oblongo-quadrangulare, angulis rotundatis, dilute piceum, nitidum, laeve, parce pilosum. Antennae conicae, 6-articulatae, articulo primo secundoque crassis, brevibus, cylindricis, penultimo intus apice processu parvo elliptico aucto. Mandibulae elongatae, tenues, falcatae. Clypeus apice sexdentatus. Labrum breve, inflexum. Maxillae subcylindricae, apice intus processu parvo elliptico, extus palpo subulato, 5-articulato. Labium breve, subcordatum, medio acumine parvo subulato instructum, palpis minutis, subulatis, 2-articulatis. Thorax parce pilosus. Prothorax latitudine longior, testaceus, nitidus, laevis. Meso- et metathorax latitudine breviores. Pedes dilute picei, ferrugineo-spinulosi. Abdomen sordide albo-flavescens, multi-setosum, setis brevibus, clavatis, clava muricata. Stigmata minuta, flava. Segmentum anale angustatum, apice caudis duabus 3-articulatis, subulatis, segmento duplo longioribus, articulo primo reliquis ambobus longitudine aequali. Anus tubulosus, perpendicularis, pedis officio inserviens, caudis longior, parce setosus. Long. 3—4 lin. Victitat per hiemem in vegetabilibus putrescentibus. Insidiatur imprimis Dipterorum larvis. Carne etiam crudo vescitur.” (Hist. nat. Ins. I. 180. 2. t. VIII. f. 1—8.)

*Q. fuliginosi* (tristis) larva a Dom. Waterhouse observata et sequenti modo descripta: „Caput et prothorax coriacea, saturate nigro-picea. Meso- et metathorax subcoriacei, flavescenti-albi. Abdomen molle, supra pallide fuscum. Long.  $6\frac{1}{2}$ —7 lin. Caput subquadratum, depressum. Antennae 4-articulatae, articulo basali brevi, secundo tertioque magis elongatis, ultimo gracili, acuto. Mandibulae longae, acutae, simplices. Labrum transversum, tuberculis numerosis instructum. Labium minutum. Palpi labiales 3-articulati. Maxillae elongatae, graciles, lobo singulo interne instructae. Palpi maxillares 4-articulati, articulo basali brevi, secundo tertioque mediocribus, subaequalibus, terminali acuto. Prothorax quadratus, capite angustior. Meso- et metathorax transversus, prothoraci latitudine subaequales. Abdomen segmentis omnibus longitudine subaequalibus, excepto ultimo, elongato, caudis duabus instructo. — Nympha artubus contiguis, abdomine recto, capite prothoraceque antorsum inflexis, illo truncum tangente. — Larva iisdem locis atque imago obvia. Versus Martii finem in nymphae staturum transit. Imago excluditur ultra medium Maii.” (Transact. Ent. Soc. Lond. I p. 32. t. III. f. 2.)

Praeter numerum articulorum antennarum at palporum, forte ab altero auctore haud satis accurate observato haec larvae satis convenire videntur.

\* Thorax processu pone coxas anticas corneo.

I. Thorax seriebus dorsalibus 3-punctatis.

\* 1. *Q. dilatatus*: Niger, parum nitidus, thoracis margine dilatato, antennis serratis. — Long. 8—10 lin.

*Er. Col. March. I.* 484. 1.

*Staph. dilatatus* Fab. Mant. Ins. I. 220. 7. Ent. Syst. I. II. 522. 12. Syst. El. II. 592. 14. — Payk. Faun. Suec. III. 389. 29. — Panz. Ent. Germ. 353. 9. — Latr. Hist. nat. Crust. et Ins. IX. 326. 74. — Grav. Mon. 89. 85. — Gyll. Ins. Suec. II. 299. 18. — Germ. Faun. Ins. Europ. VI. 13.

*Velleius dilatatus* Mannerh. Brachelytr. 19. 1. — Steph. Illustr. Britt. Ent. V. t. 26. f. 6. — Boisd. et Lucord. Faun. Ent. Paris. I. 357. — Nordm. Symb. 16. 1. — Rurde Brachelytr. Hal. I. 1.

*Staph. serraticornis* Schrank Faun. Boic. I. 641. 866.

*Staph. concolor* Marsh. Ent. Britt. 498. 4.

Habitat in Europa.

Magnus in hoc genere, niger, parum nitidus, thorace abdomineque se-

riceo-micantibus, elytris opacis. Antennae capite paulo longiores, medio crassiores, articulo tertio secundo plus duplo longiore, 4 — 10 introrsum productis, serratis, ultimo oblongo-ovato, integro, rufo-piceae, basi nigrae, apice ferrugineae. Caput thorace multo angustius, sat magnum, subovatum, supra oculos et in fronte ad oculi marginem interiorem puncto singulo utrinque impressum. Thorax latitudine tertia parte brevior, coleopteris paulo latior, lateribus et basi aequaliter rotundatus, apice truncatus, utrinque subsinuatus, medio convexus, margine laterali et basali late explanato, seriebus dorsalibus punctis tribus, apicali paulo remotiore, compositis, lateribus praeter plura marginalia singulo puncto inter medium et marginem utrinque impressus, punctis omnibus subtilibus. Scutellum punctulatum. Elytra thorace paulo longiora, subtiliter confertim punctata, breviter nigro-pubescentia. Abdomen apicem versus sensim angustatum, parcius punctulatum nigroque pubescens. Pedes concolores.

Tarsi antici maris fortiter, feminae modice dilatati.

Obs. In Vespae Crabronis nido aliquando lectus a Gyllenhal.

\* 2. *Q. lateralis*: Niger, nitidus, antennarum basi, palpis tarsisque rufo-piceis, elytris confertim punctatis, margine laterali testaceo. — Long. 5 lin.

Curt. Britt. Ent. XIV. t. 638. — Er. Col. March. I. 485. 2.

Staph. *lateralis* Grav. Micr. 35. 50. Mon. 42. 4. — Latr. Hist. nat. Crust. et Ins. IX. 319. 55. — Gyll. Ins. Suec. IV. 477. 20 — 21.

Philonth. *lateralis* Nordm. Symb. 76. 7.

Emus *lateralis* Boisd. et Lacord. Faun. Ent. Paris. I. 375. 26.

Habitat in Europa.

Statura omnino *Q. fulgidi*, at maior et praesertim latior, niger, capite, thorace scutelloque pernitidis. Antennae capite paulo longiores, tenuiores, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus, subcylindricis, ultimo oblongo, apice sensim acuminato, piceae, articulo basali rufo. Palpi rufo-picei. Caput thorace angustius, suborbiculatum, utrinque supra oculos punctis duobus, in fronte utrinque ad oculi marginem puncto singulo impressum. Oculi mediocres, parum prominuli. Thorax coleopteris paulo latior, latitudine vix brevior, lateribus et basi aequaliter rotundatis, supra leviter convexus, apice truncatus et utrinque subsinuatus, margine laterali pone medium explanato, seriebus dorsalibus punctis tribus aequaliter distantibus compositis, lateribus utrinque punctis praeter marginalia duobus impressus, punctis minus fortibus. Scutellum laevissimum. Elytra thorace parum longiora, dense subtiliter punctata, pube tenui nigra oblecta, subnitida, margine laterali inflexo testaceo. Abdomen breviusculum, apicem versus sensim angustatum, confertim subtilius punctulatum, tenuiter nigro-pubescent, cupreo-versicolor. Pedes picei, tarsi anticis in utroque sexu fortiter dilatatis.

\* 3. *Q. fulgidus*: Niger, nitidus, antennis pedibusque piceis, thorace lateribus subdilato. — Long. 3 — 5 lin.

Er. Col. March. I. 486. 3.

Staph. *variabilis* Gyll. Ins. Suec. II. 303. 21. — Zetterst. Faun. Lappon. I. 70.

10. Ins. Lappon. 61. 11. — Sahlb. Ins. Fenn. I. 314. 18. — Mannerh. Brachel. 25. 29.

Philonth. *variabilis* Nordm. Symb. 78. 15.

Staph. *nitidus* Grav. Micr. 31. 46. Mon. 42. 5. — Latr. Hist. nat. Crust. et Ins.

IX. 318. 51.

*Emus nitidus* Boisd. et Lacord. Faun. Ent. Paris. I. 377. 29.

*Elytris rufis.*

*Staph. fulgidus* Fab. Mant. Ins. I. 220. 14. Ent. Syst. I. II. 525. 25. Syst. El. II. 596. 32. — Panz. Ent. Germ. 354. 17.

*Staph. fuscipennis* Block Ins. Vall. Planens. 116. 4. f. 4.

*Staph. rufitarsis* Marsh. Ent. Brit. 512. 42.

*Staph. distinctus* Runde Brachelytr. Hal. 6. 22.

*Staph. iracundus* Say Transact. Am. Philosoph. Soc. Philadelph. Nov. Ser. IV. 449. 3.

*Emus floralis* Boisd. et Lacord. Faun. Ent. Paris. I. 380. 33.

*Philonth. assimilis* Nordm. Symb. 78. 76.

*Elytris concoloribus.*

*Staph. mesomelinus* Marsh. Ent. Brit. 510. 32.

*Staph. Grönländicus* Zetterst. Ins. Lappon. 61. 13.

*Emus occultus* Boisd. et Lacord. Faun. Ent. Paris. I. 379. 31.

Habitat in Europa et in America septentrionali.

Niger, nitidus, capite, thorace scutelloque pernitidis, elytris aut nigris, interdum nigro-piceis, sutura rufescente, aut rufis. Antennae capite sesqui longiores, articulo tertio secundo duplo prope longiore, 4 et 5 subcylindricis, 6—10 subobconicis, crassitie vix brevioribus, ultimo ovato, subcompresso, apice inferius abrupte acuminato, piceae, articulis basalibus nigricantibus. Palpi nigro-picei. Caput thorace angustius, subovale, in mare interdum capitis latitudine, suborbiculatum, omnium subtilissime punctulatum, supra utrinque supra oculos punctis tribus, in fronte utrinque ad oculi marginem interiorem puncto singulo impressum, lateribus supra oculos subtilius punctulatum. Oculi mediocres, parum prominuli. Thorax elytris latitudine aequalis, latitudine paulo brevior, basi lateribusque aequaliter rotundatis, antrosum leviter angustatus, leviter convexus, margine laterali subexplanato, seriebus dorsalibus punctis tribus approximatis compositis, lateribus punctis praeter marginalia utrinque pluribus impressis, praeterea omnium subtilissime vage obsoleteque punctulatus. Scutellum laevissimum. Elytra thorace paulo longiora, subtilius minus confertim punctata, tenuiter griseo-pubescentia. Abdomen apice angustatum, crebre subtilius punctulatum, pube subtili depressa grisea rariore vestitum, segmentis singulis piceo-marginatis. Pedes nigro-picei vel fusco-picei, tarsis rufo-piceis, anticis in utroque sexu fortius dilatatis. Alae fusco-hyalinae.

Variat 1, Niger, elytris rufis.

2, Niger, elytris concoloribus, pedibus saepius piceis.

3, Niger, pedibus piceis, abdomine rufo-brunneo, basi nigricante.

4, Dilute fusco-piceus, vel piceo-testaceus, capite elytris obscurioribus: immaturus.

5, Piceus, capite pectoreque nigricantibus, pedibus abdominisque segmentorum marginibus rufo-piceis, elytris rufis, ano testaceo.

Obs. 1. Coloribus instabilis, antennis basi nigricantibus et thoracis margine laterali subexplanato ab affinis facile distinguendus. Mas variat capite maiore minoreve, eodem modo ac Phil. aeneus alique.

2. Nomen Fabricianum, quo duo huius insecti exempla in auctoris collectione signata vidi, restitui. Typus Fabricianus elytris rufis a nonnullis auctori-



bus pro specie peculiari habetur, at vix rite, nullae enim praeter elytrorum colorem, in pluribus huius familiae speciebus eodem modo variantem, differentiae obviae sunt. Ex America septentrionali nobis quidem nonnisi individua elytris rufis allatae, at non est, quod illis in terris nonnisi eiusmodi specimina occurrere afferam. Staph. Grönländicus Westerm. e Grönländia, ab il. Prof. Zetterstedt inter Insecta Lapponica l. c. descriptus, a Dom. Westermann amice communicatus, nullo modo ab huius speciei individuis elytris concoloribus differt.

\* 4. *Q. cruentus*: *Niger, nitidus, antennarum basi, palpis, pedibus ano-*  
*que rufis, elytris parcius punctatis, rufis nigrisve.* — Long. 3–4 lin.

*Staph. cruentus* Oliv. Ent. III. 24. 27. 34. t. 5. f. 49.

*Staph. fulgidus* Marsh. Ent. Brit. 503. 16.

Habitat in Germania, Gallia, Italia, Volhynia.

Statura fere praecedentis, niger, nitidus, capite, thorace scutelloque pernitidis. Antennae capite sesqui longiores, articulo tertio secundo sesqui longiore, 4 et 5 cylindricis, crassitie haud brevioribus, 6 — 10 paulo crassioribus, transversis, subaequalibus, ultimo ovato, apice abrupte acuminato, nigro-piceae, articulis tribus primis testaceo-rufis. Palpi rufo-testacei. Caput thorace angustius, vel maris saepius eiusdem latitudine, orbiculatum, depressiusculum, supra vertice utrinque punctis duobus, supra oculos et inter oculos utrinque puncto singulo impressum, lateribus basi subtiliter punctulatis. Thorax coleopteris vix latior, latitudine summa paulo brevior, basi et lateribus cum angulis posterioribus aequaliter rotundatus, apicem versus leviter angustatus, apice truncatus, angulis anterioribus deflexis, rectis, convexus, seriebus dorsalibus punctis tribus, apicali paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 4 impressus, punctis minus fortibus. Scutellum laevissimum. Elytra thorace tertia parte longiora, minus fortiter parcius punctata, parcius griseo-pubescentia. Abdomen parallelum, parce subtiliter punctatum, parce longius griseo-pubescentia, segmentis 4 primis margine apicali, quinto apice, sexto toto rufis. Pedes rufo-testacei, coxis omnibus nigro-piceis, tarsis anticis in utroque sexu fortius dilatatis.

Variat 1, Elytris rufo-testaceis.

2, Elytris nigro-piceis, sutura picescente.

3, Elytris nigerrimis, abdominis segmentis tribus primis concoloribus, ultimis margine apicali testaceo, coxis nigris.

Obs. 1. Distinctus ab affinibus abdomine parallelo, antennis brevioribus, articulis earum 6 — 10 transversis, a praecedente praeter thoracis structuram antennarum basi pedibusque rufo-testaceis, a sequente coxis etiam anticis nigro-piceis nigrisve.

2. Var. 3 colore corporis nigerrimo et abdominis segmentorum ultimorum solo margine tenui apicali testaceo magis diversa, aliae autem differentiae desiderantur, quare pro specie peculiari rite haberi nequit. Specimina cum genuinis in Sardinia lecta communicavit Dom. Prof. Gené.

\* 5. *Q. xanthopus*: *Nigro-piceus, antennis, pedibus abdominisque seg-*  
*mentorum marginibus testaceis, elytris fortius punctatis.* — Long. 3½ lin.

*Er. Col. March. I.* 487. 4.

*Staph. variabilis* var. c Gyll. Ins. Suec. II. 303. — *Sahlb. Ins. Fenn I.* 314.

Habitat in Germaniae silvis frondosis.

*Statura omnino Q. fulgidi, at paulo minor, niger vel nigro-piceus, nitidus, capite, thorace scutelloque pernitidis. Antennae capite plus sesqui longiores, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo crassioribus at vix brevioribus, subobconicis, penultimis crassitie vix longioribus, ultimo ovato, apice abrupte acuminato, testaceae, apice interdum fusciscentes. Os rufo-piceum, palpis testaceis. Caput in utroque sexu thorace angustius, subovale, laevissimum, supra utrinque supra oculos punctis tribus, in fronte utrinque ad oculi marginem interiorem puncto singulo impressum. Oculi mediocres, parum prominuli. Thorax coleopterorum latitudine, latitudine summa vix brevior, basi et lateribus rotundatus, angulis posterioribus obtusis, subrotundatis, apice truncatus et utrinque subsinuatus, angulis anterioribus deflexis, acutiusculis, antrorsum subangustatus, leviter convexus, seriebus dorsalibus punctis tribus aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque duobus impressus, punctis minus fortibus. Scutellum laevissimum. Elytra thorace tertia parte longiora, crebrius fortiusque punctata, pube tenui haud depressa parcius vestita, nigro-picea, nitida, sutura saepe rufescente. Abdomen apice angustatum, parce subtiliter punctulatum, longius parciusque griseo-pubescent, nigro-pilosellum, segmentorum marginibus rufo-testaceis. Pedes cum coxis testacei, tarsis anticis in utroque sexu fortius dilatatis.*

Obs. Propinquus *Q. fulgido*, at antennis pedibusque testaceis, elytris longioribus, fortius punctatis, abdomine parcius punctato discedens, neque unquam cum eo obvenit; propior praecedenti, at antennarum structura, elytris crebrius fortiusque punctatis, abdomine apicem versus evidenter angustato et coxis omnibus testaceis diversus, proximus sequenti, at colore, elytris crebrius punctatis et antennarum articulo secundo tertio subaequali distinctus.

\* 6. *Q. scitus: Rufo-testaceus, nitidus, capite nigro, thoracis disco, pectore elytrorumque angulis posticis segmentisque singulis abdominis basi piceis, elytris parcius punctatis. — Long. 3 lin.*

*Er. Col. March. I. 487. 5.*

*Staph. scitus Grav. Mon. 50. 13. — Gyll. Ins. Suec. II. 306. 22. — Sahlb. Ins. Fenn. I. 316. 19. — Mannerh. Brachelytr. 26. 30. — Runde Brachelytr. Hal. 6. 23. Philonthus scitus Nordm. Symb. 79. 18.*

*Var. a: Elytris rufo-testaceis, immaculatis.*

*Staph. analis Fab. Mant. Ins. I. 221. 19. Ent. Syst. I. II. 526. 35. Syst. El. II. 598. 45. — Panz. Ent. Germ. 355. 25.*

*Var. b: Rufo-testaceus, capite solo piceo.*

*Staph. atricillus Grav. Mon. 55. 19.*

*Staph. pygmaeus Grav. Mon. 57. 23.*

Habitat in Europa, sub cortice Pinus.

*Q. fulgido minor et angustior. Antennae capite plus sesqui longiores, articulo secundo tertio vix brevior, 4 — 10 sensim paulo brevioribus et crassioribus, 6 — 10 subobconicis, penultimis crassitie vix brevioribus, ultimo ovato, apice abrupte acuminato, totae testaceae. Caput in utroque sexu thorace angustius, subovatum, supra vertice utrinque punctis duobus subtilioribus, supra oculos in fronte ad oculi marginem puncto utrinque singulo notatum, nigrum vel nigro-piceum, nitidum, ore infra antennis cum palpis testaceo, mandibulis piceis. Oculi mediocres, parum prominuli. Thorax*

coleopteris saepius paulo latior, latitudine summa paulo brevior, basi fortius, lateribus modice rotundatus, antrorsum subangustatus, apice truncatus et utrinque subsinuatus, angulis anterioribus acutiusculis, deflexis, posterioribus subrotundatis, leviter convexus, seriebus dorsalibus punctis 3, apicali remotiore, compositis, lateribus punctis praeter marginalia utrinque duobus impressus, punctis subtilioribus, nitidus, rufo-piceus, disco nigricante, nonnunquam etiam fero omnino niger, interdum totus rufo-testaceus. Scutellum laevissimum, picum. Elytra thorace tertia parte longiora, parcius minus subtiliter punctata, subtiliter parcius pubescentia, rufo-testacea, nitida, angulo apicis exteriorae saepe picescente. Abdomen elongatum, coleopteris angustius, apicem versus sensim attenuatum, parcius punctulatum, longius parciusque griseo-pubescent, undique pilis raris nigris longis erectis obsitum, rufo-testaceum, segmentis singulis plerumque basi fusciscentibus. Pectus plerumque piceum, rarius testaceum. Pedes cum coxis testacei, tarsi anticis in utroque sexu fortius dilatatis.

Obs. 1. Statura corporis minore, posterius attenuata, colore, licet quodammodo variabili, omnino dilutior facile a praecedentibus distinguitur.

2. Staph. analis Fab. secundum specimen in auctoris museo ipsum ad hanc speciem allatus: nomen autem recipere dubitavi, cum ab insecto (Boletobio anali) a Paykulo sub eodem nomine accuratius descripto et a Fabricio citato, omnino distinctum sit et distinctum esse e descriptione Fabriciana non satis pateat.

3. Staph. pygmaeum Grav. ab auctore benevole examinandum mihi communicatum, praeter magnitudinem duplo minorem vix distinctum esse me persuasum putavi.

\* 7. *Q. laevigatus*: Niger, nitidus, antennarum basi, segmentorum abdominis marginibus, pedibus elytrisque rufis, his glabris, sublaevibus. — Long. 4 lin.

Er. Col. March. I. 488. 6.

*Staph. laevigatus* Gyll. Ins. Suec. II. 306. 23. — Zetterst. Faun. Lappon. I. 71. 11. Ins. Lappon. 61. 14. — Sahlb. Ins. Fenn. I. 316. 20. — Mannerh. Brachelytr. 26. 31.

*Emus laevigatus* Boisd. et Lacord. Faun. Ent. Paris. I. 380. 32.

*Philonth. laevigatus* Nordm. Symb. 79. 19.

*Staph. glaber* Müll. Prodr. 98. 1106. (forte)

*Staph. flavopterus* Fourc. Ent. Par. I. 166. 10. (forte)

*Le Staph. noir à étuis fauves et lisses* Geoffr. Hist. Ins. I. 364. 10. (forte)

Habitat in Suecia, Germania, sub cortice Abietis, in Lapponia sylvatica (sec. Zetterst.) frequentissimus.

Statua *Q. fulgidi* individuis minoribus aequalis, at magis depressus, nigro-piceus, nitidulus. Antennae capite paulo longiores, articulo tertio secundo sesqui prope longiore, 4 — 7 sensim crassioribus, 6 — 10 transversis, 7 — 10 aequalibus, ultimo ovato, apice obtuse acuminato, nigrae, articulis tribus primis rufis. Labrum et palpi rufi. Caput thorace vix angustius, suborbiculatum, depressiusculum, utrinque vertice puncto singulo, supra oculos punctis duobus, fronte utrinque ad oculi marginem anteriorem puncto singulo impressum. Thorax coleopteris hand angustior, latitudine paulo brevior, basi et lateribus modice rotundatus, antrorsum non angustatus, apice truncatus, angulis anterioribus deflexis, subrectis, posterioribus subrotundatis, parum convexus, seriebus dorsalibus punctis tribus, apicali



remotiore, reliquis duobus approximatis, compositis, lateribus praeter marginem punctis nullis, ad marginem paulo ante medium puncto singulo maiusculo impressus. Scutellum laevissimum. Elytra thorace tertia parte longiora, rufa, glabra, subnitida, biseriatim obsolete punctata, serie altera suturali, altera dorsali. Abdomen parallelum, subtiliter parcius punctulatum, tenuiter pubescens, segmentis singulis rufo-marginatis. Pedes breves, cum coxis rufi.

Tarsi antici in mare paulo magis quam in femina dilatati.

Obs. Q. cruento magnitudine, corpore depressiusculo, antennarum structura et abdomine parallelo similis, elytris autem laevigatis et coxis rufis facile distinguendus.

\* 8. *Q. impressus*: Niger, nitidus, elytris biseriatim fortius punctatis, margine omni testaceo, fronte inter oculos bipunctata. — Long.  $3\frac{1}{2}$  lin.

Er. Col. March. I. 489. 7.

*Staph. impressus* Panz. Faun. Germ. 36. 21. — Grav. Micr. 35. 51. Mon. 39. 1. — Latr. Hist. nat. Crust. et Ins. IX. 320. 56. — Gyll. Ins. Suec. II. 307. 24. — Sahlb. Ins. Fenn. I. 317. 21. — Mannerh. Brachelytr. 26. 32.

*Emus impressus* Boisd. et Lacord. Faun. Ent. Paris. I. 381. 35.

*Philonth. impressus* Nordm. Symb. 79. 20.

*Staph. cinctus* Payk. Mon. Car. App. 137. 34–35. Faun. Suec. III. 395. 36.

*Staph. marginellus* Marsh. Ent. Brit. 505. 21.

*Staph. nitidus* var. 6 Grav. Micr. 32. 46.

Habitat in Europa.

Niger, nitidus, capite, thorace scutelloque pernigridis. Antennae capite sesqui longiores, minus graciles, articulo tertio secundo sesqui longiore, 4 — 10 cylindricis, sensim brevioribus at vix crassioribus, penultimis crassitie vix brevioribus, ultimo ovato, apice inferne abrupte acuminato, nigrae, apicem versus fuscесcentes. Palpi picei. Caput thorace paulo angustius, subovale, puncto singulo maiusculo supra oculos, alioque in oculi margine interno, duobus subtilioribus in vertice utrinque impressum. Oculi mediocres, parum prominuli. Thorax coleopteris vix angustior, latitudine paulo brevior, basi fortius, lateribus leviter rotundatus, antrorsum subangustatus, apice truncatus et utrinque subsinuatus, angulis anterioribus acutiusculis, posterioribus rotundatis, leviter convexus, seriebus dorsalibus punctis tribus, antico paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque fere 5 impressus, punctis minus subtilibus. Scutellum laevissimum. Elytra thorace paulo longiora, subtilissime punctulata, punctorum maiorum seriebus tribus, prima prope suturam, reliquis duabus in medio dorso, glabra, lateribus et margine apicali parcius pubescentibus, nigro-picea, nitida, limbo suturali, apicali lateraliue testaceo. Abdomen parce subtilius punctulatum, parcius longiusque griseo-pubescens, cupreo-versicolor, nitidulum, segmentis singulis nonnunquam tenuiter testaceo-marginatis. Pedes picei, geniculis tarsisque lutescentibus, tarsis anticis in utroque sexu fortius dilatatis.

Obs. *Staphylinus flavescens* Fab. Syst. Ent. 266. 9. Spec. Ins. I. 336. 10. Mant. Ins. I. 221. 16. Ent. Syst. I. II. 526. 32. Syst. El. II. 597. 41. — Müll. Prod. 98. 1113. — Goeze Ent. Beitr. I. 719. 13. — Panz. Ent. Germ. 355. 23. — Cederh. Faun. Ingr. 118. 362

forte ad hocce insectum revocandus; a Gravenhorstio quidem ad Philonthum di-

scoideum relatus est, magnitudo autem, quam in Syst. Entomologiae Philontho marginali aequalem affert Fabricius, magis huic speciei convenire videtur.

\* 9. *Q. pediculus*: *Niger, nitidus, elytris biserialim punctatis, piceo-marginatis, fronte inter oculos 4-punctata. — Long. 3 lin.*

*Philonth. pediculus Nordm. Symb. 79. 24.*

Habitat in Unalaschka, Def. Chamisso.

Praecedenti affinis, niger, nitidus. Antennae capite duplo prope longiores, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, 4 — 6 cylindricis, 7 — 10 subobconicis, penultimis crassitie haud brevioribus, ultimo ovato, apice acuminato, nigrae, articulo secundo summa basi testaceo. Caput thorace dimidio angustius, orbiculatum, parum convexum, supra oculos punctis utrinque 3 vel 4, inter oculos punctis 4 transversim positis, aequaliter distantibus. Oculi mediocres, parum prominuli. Thorax coleopterorum latitudine, latitudine summa vix brevior, basi et lateribus eodem arcu rotundatus, apice truncatus et utrinque subsinuatus, angulis anterioribus acutiusculis, leviter convexus, seriebus dorsalibus punctis tribus approximatis compositis, lateribus punctis praeter marginalia utrinque duobus impressus. Scutellum laevissimum. Elytra thorace paulo longiora, subtiliter obsoleteque punctulata, seriebus tribus punctorum maiorum impressis, prima suturali, reliquis duabus dorsalibus, lateribus et margine apicali pubescentibus, ceterum subglabra, margine laterali et apicali piceis. Abdomen apicem versus sensim angustatum, subtilius punctatum, longius pubescens, metallico-versicolor, segmentorum marginibus piceis. Pedes nigro-picei, tarsis piceis, anticis in utroque sexu sat fortiter dilatatis.

Obs. Affinis sequenti, at gracilior, elytris longioribus, piceo-marginatis et abdomine parcius punctato distinctus.

\* 10. *Q. capucinus*: *Niger, nitidus, palpis pedibusque piceis, elytris irregulariter triserialim fortius punctatis, fronte inter oculos punctis 4 transversalibus impressa. — Long. 3 lin.*

*Staph. capucinus Grav. Mon. 40. 2.*

*Philonth. capucinus Nordm. Symb. 79. 23.*

*Staph. inversus Say Transact. Am. Philosoph. Soc. Philadelph. Nov. Ser. II. 449. 5.*

Habitat in America septentrionali.

Statura omnino *Q. impressi*, at paulo minor, niger, nitidus, capite, thorace scutelloque pernitidis. Antennae capite sesqui longiores, articulo secundo tertio vix brevior, 4 — 10 sensim paulo brevioribus et crassioribus, penultimis crassitie paulo brevioribus, ultimo ovato, apice inferius abrupte acuminato, piceae, articulis tribus primis nigris. Palpi picei. Caput thorace paulo angustius, rotundatum, utrinque supra oculos punctis paucis, inter oculos punctis 4, transversim positis, aequaliter distantibus impressum. Thorax basi coleopteris vix angustior, latitudine summa vix brevior, antrorsum angustatus, basi fortius, lateribus leviter rotundatus, apice truncatus, angulis anterioribus subrectis, posterioribus subrotundatis, transversim convexus, seriebus dorsalibus punctis tribus modice approximatis compositis, lateribus punctis praeter marginalia utrinque nonnullis impressus, punctis subtilioribus. Elytra thoracis longitudine, nitida, parcius subtiliter obsoleteque punctulata, seriebus irregularibus tribus punctorum maiorum,

una suturali, reliquis duabus dorsalibus, lateribus et margine apicali parce pubescentia, ceterum glabra. Abdomen apicem versus sensim angustatum, crebrius punctulatum, nigro-pubescent, aeneo-nitidulum. Pedes picei, tarsi rufescentibus, anticis in utroque sexu dilatatis.

11. *Q. chlorophanus*: *Viridi-metallicus, nitidus, abdomine metallico-versicolore, ano ferrugineo.* — Long. 4 lin.

Habitat in Bengala, Mus. Dom. Hope.

Statura omnino *Q. fulgidi*, viridi-metallicus, nitidus. Antennae capite sesqui fere longiores, apice haud crassiores, articulo tertio secundo longiore, 4 — 10 subaequalibus, ultimo ovato, apice acuminato, nigrae. Mandibulae et palpi apice picei. Caput thorace dimidio angustius, subovatum, depressiusculum, supra oculos punctis utrinque tribus, inter oculos utrinque puncto singulo impressum. Oculi mediocres, parum prominuli. Thorax coleopterorum latitudine, latitudine summa tertia parte brevior, lateribus rotundatis, apicem versus leviter angustatus, basi leviter rotundatus, apice truncatus, angulis anterioribus subrectis, deflexis, posterioribus obtusis, leviter convexus, margine laterali subexplanato, seriebus dorsalibus punctis tribus, anterioribus duobus magis approximatis, compositis, lateribus punctis praeter marginalia utrinque 4 impressus. Scutellum laevissimum. Elytra thoracis longitudine, minus dense aequaliter punctata, parce griseo-pubescentia. Abdomen apicem versus sensim angustatum, parcius punctatum, tenuiter pubescens, fortiter metallico-versicolor, segmento quinto apice sextoque toto ferrugineis. Pedes tarsi picei, anticis in utroque sexu fortiter dilatatis.

## 2) Thorax seriebus dorsalibus 2-punctatis.

- \* 12. *Q. viridulus*: *Niger, nitidus, capite thoraceque obscure virescentibus, elytris laete viridibus, subtiliter punctatis, scutello punctulato.* — Long. 4 lin.

Habitat Bogotae in Columbia interiore, Dom. Buquet.

Statura omnino *Q. impressi*, niger, nitidus, capite thoraceque politis, nigro-virescentibus, elytris laete cyanescenti-viridibus, nitidis. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, penultimis subobconicis, crassitie vix longioribus, ultimo ovato, apice abrupte acuminato, nigrae. Palpi nigri, maxillares articulo quarto minore. Caput thorace paulo angustius, suborbiculatum, fronte planiuscula, punctis utrinque tribus oblique positis ad verticem, duobus approximatis supra oculos, singulo ad oculi marginem anteriorem impressum. Oculi parvi, rotundati, parum prominuli. Thorax coleopteris paulo angustior, basi et lateribus leviter rotundatis, his pone medium subsinuatis, antrorsum subangustatus, apice truncatus et utrinque subsinuatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, parum convexus, seriebus dorsalibus punctis duobus prope apicem sitis compositis, lateribus punctis praeter marginalia nullis, marginali maiusculo singulo utrinque ante medium impressus. Scutellum subtiliter punctulatum. Elytra thorace sesqui longiora, subtiliter minus dense punctata, tenuiter nigro-pubescentia. Abdomen breviusculum, coleopteris angustius, apicem versus sensim



attenuatum, nigro-subaeneum, nitidum, parce subtiliterque punctulatum, parce nigro-pubescent. Pedes nigri.

Tarsi antici feminae modice dilatati. Mas latet.

3) Thorax seriebus dorsalibus nullis.

\* 13. *Q. labiatus*: *Niger, nitidus, capite antice rufo, elytris punctis nonnullis profundis impressis. — Long. 3 lin.*

Habitat in Columbia, Dom. Moritz.

Statura omnino *Q. impressi*, at paulo minor, niger, capite, thorace scutelloque pernitidis. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 4—10 sensim brevioribus et crassioribus, penultimis crassitie paulo brevioribus, ultimo subovato, apice abrupte acuminato, rufae, medio fuscescentes. Palpi rufi. Caput thorace paulo angustius, oculis magnis, prominulis, infra oculos rufum, puncto singulo in margine oculi utrinque notatum. Thorax basi vix coleopteris paulo angustior, latitudine summa haud brevior, antrorsum subangustatus, basi cum angulis posterioribus fortius, lateribus leviter rotundatus, apice truncatus, angulis anterioribus subrectis, modice convexus, puncto singulo maiusculo utrinque ad marginem anteriorem notatus, ceterum praeter puncta nonnulla marginalia subtiliora laevissimus. Scutellum laevissimum. Elytra thoracis longitudine, laevigata, nitida, stria suturali profundius impressa, punctorum rariorem seriebus duabus, altera in stria suturali, altera dorsali, et praeterea punctis duobus vel tribus maioribus in medio disci, apice pilis parvis nigris ciliata. Abdomen longius parciusque nigro-pubescent, parce punctulatum, cupreo-chalybeo nitidulum, segmento quarto subtus medio late laevi. Pedes antici testacei, intermedii piceo-testacei, postici picci, tarsi testacei. Alae fusco-hyalinae.

Tarsi antici maris fortiter, feminae leviter dilatati.

Variat. 1, Thorace, elytris abdomineque concoloribus.

2, Thorace elytrisque concoloribus, abdomine segmentis ultimis duobus rufis.

3, Thorace concolore, elytris, abdominisque segmentis duobus ultimis rufis.

4, Elytris concoloribus, thorace abdominisque segmentis duobus ultimis rufis.

5, Thorace, elytris abdominisque segmentis ultimis duobus rufis.

14. *Q. frenatus*: *Testaceus, nitidus, thoracis pectorisque lateribus abdomineque nigris, elytris dorso triseriatim profunde punctatis. — Long. 3 lin.*

Habitat prope La-Guayram in Columbia, Dom. Moritz.

Praecedenti affinis. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, 4—10 sensim brevioribus et parum crassioribus, 4 et 5 cylindricis, reliquis subobconicis, penultimis crassitie paulo brevioribus, ultimo ovato, apice abrupte acuminato, totae testaceae. Caput thorace paulo angustius, oculis permagnis, prominulis, inter oculos puncto utrinque singulo, ad oculi marginem anteriorem sito, supra oculos punctis duobus notatum, totum cum ore testaceum, nitidum. Thorax basi coleo-

pteriorum latitudine, latitudine summa paulo brevior, antrorsum modice angustatus, lateribus, basi et angulis posterioribus rotundatis, apice truncatus, angulis anterioribus deflexis, subrotundatis, modice convexus, punctis 4 transversim positis ad marginem anteriorem impressus, ceterum praeter puncta plura marginalia subtiliora laevissimus, rufo-testaceus, nitidus, limbo laterali nigro. Scutellum laeve, nigro-piceum. Elytra thoracis longitudine, luteo-testacea, circa scutellum picea, nitida, disco glabra, dorso punctorum sat profundorum seriebus tribus, prima prope suturam, in stria suturali impressa sita, regulari, reliquis duabus discoidalibus, irregularibus, lateribus punctis pluribus profundis subseriatis impressis. Abdomen parce punctatum, nigro-pubescent, nigrum, segmento quinto margine apicali, sexto toto testaceis. Pectus obscure testaceum, lateribus nigrum. Pedes testacei.

Maris tarsi antici fortiter dilatati, abdominis segmenta quartum et quintum inferiora medio longitudinaliter leviter impressa, impressione dense fulvo-villosa. Femina latet.

15. *Q. buphthalmus*: *Niger, nitidus, capite, thorace elytrisque aeneis, his variolosis, femoribus basi albis, antennis clavatis, oculis maximis.* — *Long. 3½ lin.*

Habitat in Mexico, Mus. Dom. Chevrolat.

Insectum singulare, oculis magnis prominentibus et antennis tenuibus clavatis insigne, ab hoc genere autem rite haud distinguendum. Antennae capite prope breviores, tenues, articulis 5 primis elongatis, secundo primo plus duplo brevior, tertio secundo paulo longiore, 3 — 5 sensim paulo brevioribus, 6 — 10 brevibus, sensim paulo crassioribus, sexto subgloboso, 7 — 10 transversis, ultimo conico, acuminato, articulis 6 primis piceo-testaceis, sequentibus fuscis, ultimo luteo. Palpi piceo-testacei. Caput magnum, thorace parum angustius, oculis permagnis, fortiter prominentibus, fronte planiuscula, subtiliter vage punctata, utrinque longitudinaliter subimpressa, supra obscure viridi-aeneum, pernitidum, subtus nigrum. Thorax coleopterorum latitudine, latitudine dimidio fere brevior, antrorsum haud angustatus, basi et lateribus eodem arcu rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, fortiter convexus, ad marginem apicalem punctis 4 transversim positis notatus, dorso inter apicem et medium utrinque leviter oblique impressus, margine laterali explanato, inaequali, obscure viridi-aeneus, pernitidus. Scutellum punctato-variolosum, viridi-aeneum. Elytra thorace paulo longiora, inaequalia, fortiter inaequaliter punctata, aenea, nitida. Abdomen breviusculum, nigrum, nitidum, parce nigro-pubescent. Pectus nigrum, nitidum. Pedes graciliores, picei, femoribus basi ultra medium albidis.

Tarsi antici feminae modice dilatati. Mas latet.

16. *Q. curtus*: *Brevis, niger, nitidus, elytris rufis, parce subtiliterque punctatis.* — *Long. 4 lin.*

Habitat in Sardinia, Mus. Reg. Taurin.

*Q. fulgidi* maximis individuis latitudine aequalis, at multo brevior, statura curtata ab aliis huius generis speciebus discedens. Niger, nitidus, capite thorace scutelloque politis. Antennae capite parum longio-

res, tennes, articulo tertio secundo paulo longiore, ultimo apice oblique truncato, rufo-piceae. Palpi rufi. Labrum rufo-piceum. Caput thorace angustius, oblongum, basi vix constrictum, parce subtilissime punctulatum, fronte utrinque supra oculos et ad oculi marginem anteriorem puncto minusculo singulo impressa. Thorax coleopterorum latitudine, antrorsum subangustatus, latitudine tertia fere parte brevior, basi fortiter, lateribus leviter rotundatus, apice truncatus, utrinque subsinuatus, angulis anterioribus deflexis, antrorsum leviter prominentibus, rotundatis, transversim convexus, laevissimus, ad angulos posteriores utrinque subimpressus. Elytra thoracis longitudine, apice rotundata, parcius subtiliter punctata, rufa, nitida, parce subtiliterque griseo-pubescentia. Abdomen breve, triangulare, minus parce subtiliterque punctatum, nigrum, nitidum, segmentis singulis margine apicali obscure piceo, parce subtiliterque griseo-pubescentis. Pedes rufo-picei, femoribus nigris.

Tarsi antiqui maris modice, feminae leviter dilatati.

\* 17. *Q. brevis*: *Brevis, piceus, nitidus, antennis elytrisque rufis, his parcius punctatis.* — Long. 2 lin.

Habitat in Germania septentrionali.

*Q. fulgidus* minimis individuis latitudine aequalis at multo brevior. Piceus, nitidus, capite, thorace scutelloque politis. Antennae capite paulo longiores, articulo tertio secundo vix longiore, 4—7 sensim paulo brevioribus et crassioribus, 7—10 aequalibus, crassitie brevioribus, ultimo oblongo, apice oblique truncato, totae rufae. Os rufum. Caput thorace paulo angustius, basi profundius constrictum, supra punctis duobus fortibus baseos et puncto utrinque ad oculi marginem anteriorem impressum. Thorax coleopteris paulo latior, latitudine quarta fere parte brevior, antrorsum subangustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus subrectis, posterioribus obtusis, convexus, ante marginem anteriorem puncto utrinque singulo notatus, ceterum laevissimus, margine laterali, praecipue posterius, explanato, piceo-rufescens. Scutellum laevissimum. Elytra thorace paululum longiora, parcius et minus subtiliter punctata, tenuiter longius pubescentia, rufa, nitida, apice leviter rotundata. Abdomen breviusculum, apicem versus leniter angustatum, crebrius punctatum, pubescens, lateribus pilosellum, segmentis singulis apice rufo-marginatis, ano testaceo. Pedes rufi.

Tarsi antiqui feminae modice dilatati. Mas latet.

Obs. Affinis praecedenti, at multo minor et minus contractus.

\*\* Thorax processu pone coxas anticatas membranaceo.  
Series dorsales tripunctatae.

\* 18. *Q. molochinus*: *Niger, nitidus, antennis pedibusque rufo-piceis, elytris thorace brevioribus, castaneis, scutello punctulato.* — Long. 5 lin.

*Er. Col. March. I.* 489. 8.

*Staph. molochinus* Grav. Mon. 46. 6. — Gyll. Ins. Suec. II. 302. 20. — Sahlb. Ins.

Fenn. I. 314. 17. — Mannerh. Brachelytr. 25. 28. — Runde Brachelytr. Hal. 5. 21.

*Emus molochinus* Boisd. et Lacord. Faun. Ent. Paris. I. 377. 28.

*Philonth. molochinus* Nordm. Symb. 76. 6.



*Staph. laticollis* Grav. *Micr.* 173. 28. *Mon.* 47. 8.

*Staph. picipennis* Payk. *Faun. Suec.* III. 373. 8.

*Staph. Lapponicus* Zetterst. *Ins. Lappon.* 61. 12.

*Staph. laevicollis* Runde *Brachelytr. Hal.* 6. 24.

Habitat in omni Europa et in America septentrionali.

Niger, capite thoraceque pernitidis. Antennae capite duplo longiores, apicem versus haud incrassatae, tenuiores, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus, crassitie longioribus, ultimo subcompresso, apice oblique truncato et inferne acuminato, rufae, articulis 2 — 4 fusciscentibus. Palpi rufi. Labrum rotundatum, integrum. Caput thorace angustius, subovale, utrinque supra oculos punctis 3 vel 4, et in fronte utrinque ad oculi marginem interiorem puncto singulo notatum. Oculi mediocres, parum prominuli. Thorax coleopteris paulo latior, latitudine summa haud brevior, antrorsum subangustatus, basi modice, lateribus leviter rotundatus, apice subemarginatus, angulis anterioribus fortiter deflexis, subrectis, posterioribus subrotundatis, fortiter convexus, seriebus dorsalibus punctis tribus approximatis compositis, lateribus punctis praeter marginalia utrinque duobus, oblique positis, versus angulos anteriores situs impressus. Scutellum punctulatum. Elytra thorace breviora, subtiliter dense punctata, cinereo-pubescentia, opaca, aut castanea aut nigra. Abdomen elongatum, apicem versus vix attenuatum, densius subtiliter punctulatum, tenuiter griseo-pubescentia, cupreo-versicolor, nitidulum, subtus segmentis singulis castaneo-marginatis. Pedes picei.

Tarsi antichi maris fortiter, feminae modice dilatati.

\* 19. *Q. frontalis*: Niger, nitidus, antennis, palpis, tibiis tarsisque rufopiceis, elytris thorace longioribus, scutello punctulato, fronte inter oculos punctis sex transversalibus impressa. — Long. 5 lin.

*Philonth. frontalis* Nordm. *Symb.* 76. 5.

Habitat in Gallia, Italia, Dalmatia, Graecia.

Praecedenti affinis, at robustior, statura omnino sequentis, niger, capite thoraceque nitidis. Antennae capite paulo longiores, tenuiores, apicem versus haud incrassatae, articulo tertio secundo sesqui longiore, 4 — 10 sensim brevioribus, 4 — 6 cylindricis, 7 — 10 subobconicis, penultimis etiam crassitie longioribus, ultimo subovato, apice oblique truncato et inferne acuminato, totae rufae. Palpi rufi. Labrum rotundatum, integrum. Caput thorace angustius, subovale, utrinque supra oculos punctis nonnullis, fronte inter oculos punctis sex, in seriem transversalem medio late interruptam dispositis. Oculi maiusculi, parum prominuli. Thorax basi coleopterorum latitudine, latitudine summa haud brevior, antrorsum leviter angustatus, basi fortius, lateribus leviter rotundatus, apice subemarginatus, angulis anterioribus deflexis, subrectis, posterioribus subrotundatis, convexus, seriebus dorsalibus punctis tribus aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 5 impressus. Scutellum subtiliter punctulatum. Elytra thorace paululum longiora, subtiliter confertissime punctata, tenuiter nigro-pubescentia, opaca. Abdomen apicem versus parum angustatum, subtiliter crebre punctulatum, densius subtiliter nigro-pubescentia, plerumque laete

viridi-versicolor. Pedes nigri, tibiis tarsisque rufo-piceis, tarsis anticis in utroque sexu fortiter dilatatis.

Obs. Intermedius inter *Q. molochinum* et *fuliginosum*, illi scutello punctulato, huic fronte inter oculos sexpunctata affinis, abdomine laetis coloribus resplendente insignis, olim a Dahlio nomine *Staph. chloroventris* impositus.

\* 20. *Q. fuliginosus*: *Niger, nitidus, antennis, palpis tarsisque rufis, fronte inter oculos punctis sex transversalibus impressa. — Long. 5 lin.*

*Er. Col. March. I. 490. 9.*

*Staph. fuliginosus Grav. Micr. 34. 49. Mon. 48. 10. — Latr. Hist. nat. Crust. et Ins. IX. 319. 56.*

*Staph. tristis Grav. Micr. 34. 48. Mon. 48. 11. — Latr. Hist. nat. Crust. et Ins. IX. 319. 55. — Gyll. Ins. Suec. II. 301. 19. — Sahlb. Ins. Fenn. I. 314. 16. — Mannerh. Brachelytr. 25. 26. — Runde Brachelytr. Hal. 5. 20.*

*Emus tristis Boisd. et Lacord. Faun. Ent. Paris. I. 376. 27.*

*Philonth. tristis Nordm. Symb. 75. 4.*

*Staph. dilatatus Marsh. Ent. Britt. 504. 19.*

Habitat in Europa.

*Niger*, capite, thorace scutelloque pernitidis. Antennae capite sesqui longiores, tenuiores, apicem versus haud incrassatae, articulo tertio secundo sesqui prope longiore, 4—10 sensim paulo brevioribus, 4—6 cylindricis, 7—10 subobconicis, penultimis etiam crassitie paulo longioribus, ultimo oblongo-ovato, apice oblique truncato et inferne acuminato, rufae, articulis 4 primis dorso picescentibus. Palpi rufo-testacei. Labrum medio subsinuatum. Caput thorace paulo angustius, subovale, utrinque supra oculos punctis 2 vel 3, fronte inter oculos punctis 6, in seriem transversalem medio late interruptam dispositis impressum. Oculi maiusculi, modice prominuli. Thorax basi coleopterorum latitudine, iatitudine summa vix brevior, antrorsum subangustatus, basi et lateribus leviter rotundatus, apice truncatus et utrinque subsinuatus, angulis anterioribus subrectis, deflexis, posterioribus obtusis, convexus, seriebus dorsalibus punctis tribus, sat approximatis, aequaliter distantibus compositis, lateribus punctis praeter marginalia utrinque 3 vel 4 impressus. Scutellum laevissimum. Elytra thorace haud breviora, confertissime subtiliter punctata, pube tenui, depressa, nigra densius oblecta, opaca. Abdomen apicem versus parum angustatum, subtiliter densius punctulatum, tenuiter nigro-pubescent, obsolete cupreo-versicolor. Pedes picei, tarsis rufo-piceis, anticis in utroque sexu fortiter dilatatis.

\* 21 *Q. picipes*: *Niger, nitidus, antennis, pedibus elytrisque rufis, his crebre punctatis. — Long. 4 lin.*

*Er. Col. March. I. 491. 10.*

*Staph. picipes Mannerh. Brachelytr. 26. 34.*

*Philonth. picipes Nordm. Symb. 77. 11.*

*Philonth. varicolor Nordm. Symb. 76. 9.*

Habitat in Germania rarius, prope Berolinum, Weber, in Austria, Dom. Schüppel, in Corcyra frequens, Parreyss.

Habitu fere *Q. maurorufi*, at multo maior, *Q. fulgido* statura aequalis, niger, capite thoraceque pernitidis. Antennae capite sesqui longiores, tenuiores, apicem versus vix incrassatae, articulo tertio secundo sesqui lon-

giore, 4—10 sensim parum brevioribus et vix crassioribus, 4—6 cylindricis, 7—10 subobconicis, omnibus crassitie longioribus, ultimo oblongo, apice inferius sensim acuminato, totae rufae. Palpi rufi. Caput thorace parum angustius, orbiculatum, punctis utrinque supra oculos duobus, ad oculi marginem anteriorem singulo impressum. Thorax coleopterorum latitudine, latitudine summa vix brevior, antrorsum subangustatus, lateribus leviter, basi cum angulis posterioribus fortiter rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, convexus, seriebus dorsalibus punctis tribus aequaliter modice distantibus compositis, lateribus punctis praeter marginalia utrumque duobus vel tribus impressus. Scutellum laevissimum. Elytra thoracis longitudine, crebre subtilius punctata, rufa, nitida, subtiliter tenuiterque griseo-pubescentia. Abdomen apicem versus sensim sat fortiter attenuatum, minus dense punctatum, pube nigra longiore depresso minus subtili parcius vestitum, cupreo-versicolor. Pedes cum coxis rufo-testacei, tarsis anticis in utroque sexu leviter dilatatis.

Obs. Ph. varicolor Nordm. e Gallia meridionali quo differat a Q. picipede e descriptione satis accurata non patet.

22. *Q. plancus*: *Niger, capite thoraceque nigro-aeneis, nitidis, elytris crebre subtilius punctatis, abdominis segmentorum marginibus, antennarum basi pedibusque testaceis, tibiis fuscis.* — *Long. paulum ultra 3 lin.*

Habitat in Sardinia, Mus. Reg. Taurin.

Statura et summa affinitas Q. fimbriati, at paulo minor. Antennae capite duplo longiores, articulo tertio secundo sesqui longiore, ultimo apice leviter oblique emarginato, acuminato, nigrae, articulis tribus primis testaceis, secundo tertioque apice fuscis. Oris partes testaceae. Caput thoracis apice parum angustius, basi profundius constrictum, supra oculos utrinque crebre subtiliter punctatum, punctisque quatuor maioribus notatum, fronte subdepressa, inter oculos punctis quatuor, transversim positiss, aequaliter distantibus impressa, nigro-subaeneum, nitidum. Thorax basi coleopteris paulo angustior, antrorsum subangustatus, latitudini baseos longitudine aequalis, basi fortiter, lateribus leviter rotundatus, apice subemarginatus, angulis anterioribus acutiusculis, fortiter deflexis, transversim modice convexus, nigro-subaeneus, nitidus, lateribus piceus. Scutellum laevissimum, nigrum, nitidum. Elytra thorace tertia parte longiora, dense subtilius punctata, pube longiore depresso fulvescenti-grisea tenuiter vestita, fusco-testacea, subnitida. Abdomen crebre subtilius punctatum, pube longiore depresso fulvescenti-grisea vestitum, nigro-pilosellum, nigrum, nitidum, iridescens, segmentis singulis margine testaceo, anterioribus angustiore et saturatiore, ultimis duobus lato et dilutiore. Pectus nigrum. Pedes dilute testacei, coxis anterioribus concoloribus, tibiis omnibus fuscis, apice obscure testaceis, tarsis etiam posterioribus articulis omnibus depresso, ultimo s. unguiculari reliquis vix angustiore.

Mas latet. Femina, tarsis anticis fortiter dilatatis.

- \*23. *Q. ochropterus*: *Niger, nitidus, capite thoraceque nigro-subaeneis, antennis, elytris, abdominis segmentorum marginibus pedibusque rufis, coxis omnibus femoribusque posticis piceis.* — *Long. 4 lin.*

Habitat in Austria, Dom. Schüppel.



Habitu omnino sequentis, at sesqui maior, niger, capite thoraceque supra nigro-subaeneis, pernitidis. Antennae capite plus sesqui longiores, tenuiores, apicem versus haud incrassatae, articulo tertio secundo paulo longiore, 4—10 sensim brevioribus, cylindricis, penultimis etiam crassitie longioribus, subobconicis, ultimo oblongo-ovato, apice sensim acuminato, rufo-testaceae, apice summo flavescente. Palpi rufo-testacei. Caput thorace paulo angustius, orbiculatum, supra utrinque punctis duobus profundioribus, oblique positis supra oculos, et singulo ad oculi utriusque marginem interiorem impressum. Oculi magni, modice prominentes. Thorax basi coleopterorum latitudine, latitudine summa vix brevior, antrorsum subangustatus, basi et lateribus leviter rotundatus, apice truncatus, angulis anterioribus subrectis, leviter deflexis, posterioribus obtusis, subrotundatis, leviter convexus, seriebus dorsalibus punctis tribus aequaliter modice distantibus compositis, lateribus punctis praeter marginalia anteriores utrinque duobus, sat distantibus, oblique positis impressus. Scutellum laevissimum. Elytra thorace paululum longiora, crebre subtilius punctata, transversim subrugulosa, dilute rufo-brunnea, pube breviora depressa grisea tenuiter vestita. Abdomen apicem versus sensim et sat fortiter attenuatum, parcius subtiliter punctatum, pube subtiliore depressa fusco-grisea vestitum, metallico-versicolor, segmentis omnibus apice rufo-marginatis. Pedes graciliores, rufo-testacei, coxis omnibus nigro-piceis, femoribus posterioribus picescentibus.

Tarsi antici maris modice dilatati. Femina latet.

Obs. A Q. picipede thorace minus convexo, elytris paulo longioribus, coxis nigro-piceis, et abdominis segmentis apice rufo-marginatis facile distinguitur.

\* 24. *Q. fimbriatus*: *Niger, nitidus, antennis, pedibus elytrisque testaceis, his crebrius punctatis, macula magna communi quadrata nigricante.* — Long.  $3\frac{1}{2}$  lin.

Habitat in Austria.

Habitu omnino praecedentis, at minor, niger, capite thoraceque pernitidis. Antennae capite plus sesqui longiores, tenuiores, apicem versus vix incrassatae, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus et paululum crassioribus, cylindricis, penultimis subobconicis, crassitie vix longioribus, ultimo ovato, apice acuminato, totae rufo-testaceae. Palpi testacei. Caput thorace parum angustius, suborbiculatum, perparum convexum, utrinque supra oculos punctis duobus oblique positis, ad marginem oculi utriusque interiorem puncto singulo impressum. Oculi maiusculi, modice prominuli. Thorax basi coleopteris haud angustior, latitudine paulo brevior, basi lateribusque modice rotundatus, antrorsum parum angustatus, apice truncatus et utrinque subsinuatus, angulis anterioribus leviter deflexis, acutiusculis, posterioribus subrotundatis, modice convexus, seriebus dorsalibus punctis tribus sat approximatis, antico subtiliore, compositis, lateribus punctis praeter marginalia utrinque singulo ante medium prope marginem lateralem profundiore impressus. Scutellum laevissimum. Elytra thorace paulo longiora, crebrius profunde punctata, fortius griseo-pubescentia, dorso nigricantia, lateribus et margine apicali fusco-testaceis. Abdomen apicem versus sensim fortius attenuatum, crebrius subtiliter punctulatum, pube minus subtili longiore depressa lutescenti-grisea densius vestitum,

nitidulum, viridi-versicolor, segmentis omnibus testaceo-marginatis. Pedes cum coxis omnibus testacei, tarsis anticis in utroque sexu fortius dilatatis.

Obs. Proximus praecedenti, at distinctus pedibus totis cum coxis testaceis, abdomine elytrisque pube minus subtili, longiore, densiore, lutescente vestitis, his minus confertim at profundius punctatis, interstitiis punctorum laevibus, dorso praeter apicem nigris.

\* 25. *Q. peltatus*: *Brunneus, abdomine versicolore, antennis fusco-piceis, basi nigricantibus, capite orbiculato, nigro, thorace latiore, elytris densius subtiliter punctatis.* — Long. 3 lin.

*Qued. praecox* Er. Col. March. I. 492. 12.

Habitat in Germania, Hungaria.

*Q. maurorufus* antrorsum magis dilatatus, obscure brunneus, capite, thorace scutelloque nitidis. Antennae capite plus sesqui longiores, tenuiores, apicem versus sensim leniter incrassatae, articulo tertio secundo paulo longiore, 4—10 sensim brevioribus et crassioribus, subobconicis, penultimis etiam crassitie longioribus, ultimo oblongo, apice inferne sensim acuminato, pilosellae, piceae, articulis singulis, imprimis basalibus, medio nigricantibus. Caput depressiusculum, thorace multo angustius minusque, utrinque supra oculos punctis duobus, oblique positis, superiore minore, ad oculi utriusque marginem interiorem puncto singulo notatum, lateribus pone oculos sublaeve, nigrum, ore infra antennis piceo-testaceo. Oculi mediocres, prominentes. Thorax basi coleopterorum latitudine, basi fortius, lateribus cum angulis posterioribus leviter rotundatus, antrorsum leviter angustatus, apice subemarginatus, angulis anterioribus deflexis, leviter prominentibus, subrotundatis, leviter convexus, seriebus dorsalibus punctis tribus sat approximatis compositis, lateribus punctis praeter marginalia utrinque anterius duobus, altero prope seriem dorsalem, altero prope marginem lateralem, impressus, punctis minus subtilibus. Scutellum laevissimum. Coleoptera latiuscula, thorace parum longiora, subtilius dense punctata, opaca, pube depressa grisea tenuiter vestita, brunnea, dorso nigricantia, margine tenui apicali obscure testaceo. Abdomen coleopteris angustius, apicem versus sensim fortius angustatum, parcius punctulatum, tenuissime pubescens, segmentis singulis basi viridi-, apice cupreo-versicoloribus. Pedes rufo-picei, femoribus posticis medio viridi-micantibus. Alae fusco-hyalinae.

Tarsi antici maris fortiter, feminae modice dilatati.

Obs. Olim, cum *Q. praecox* specimen singulum a Gravenhorstio descriptum immaturum novissem, ab hoc mihi non distinctus visus est; speciminibus autem maturis a Prof. Gené communicatis, has duas species inter se simillimas distinxim. *Q. peltatus* thorace paulo latiore, capite orbiculato, planiusculo, antennis basi nigricantibus, elytris minus dense et minus profunde, abdomine paulo parcius et minus subtiliter punctatis praecipue discedit. A sequentibus thorace coleopterisque latioribus et elytris paulo longioribus differunt.

\* 26. *Q. praecox*: *Brunneus, abdomine versicolore, antennis basi pedibusque piceo-testaceis, capite ovato, nigro, elytris confertim subtilius punctatis.* — Long. 3 lin.

*Staph. praecox* Grav. Micr. 172. 27. Mon. 57. 21. — Latr. Hist. nat. Crust. et Ins. IX. 324. 68.

Habitat in Lusitania, Com. Hoffmannsegg, in Sardinia, Prof. Gené.

Praecedenti proximus. Obscure brunneus, capite thorace scutelloque nitidis. Antennae capite duplo longiores, tenuiores, apicem versus vix incrassatae, articulo tertio secundo sesqui longiore, 4—10 sensim paulo brevioribus, subcylindricis, penultimis etiam crassitie longioribus, leviter obconicis, ultimo oblongo, apice sensim acuminato, piceae, articulis tribus primis testaceis. Palpi testacei. Caput parvulum, thorace duplo angustius, ovatum, supra leviter convexum, punctis utrinque duobus oblique positus supra oculos, et singulo ad oculi utriusque marginem anteriorem impressum, lateribus supra oculos utrinque confertim subtilissime punctulatum, nigrum, nitidum, ore piceo-testaceo. Oculi mediocres, modice prominuli. Thorax basi coleopterorum prope latitudine, latitudine summa vix longior, basi fortiter, lateribus cum angulis posterioribus leviter rotundatus, antrorsum leviter angustatus, apice submarginatus, angulis anterioribus fortiter deflexis, acutiusculis, modice convexus, seriebus dorsalibus punctis tribus, posterioribus sat approximatis, antico paulo remotiore, compositis, lateribus punctis praeter marginalia antierius utrinque tribus, duobus prope seriem dorsalem, tertio prope marginem impressus, limbo undique brunneo, disco late nigropiceo. Scutellum laevissimum, piceum. Elytra thorace paulo longiora, dense subtilius punctata, pube longiore depressa grisea vestita, brunnea, dorso nigricantia, margine apicali obscure testaceo. Abdomen apicem versus sensim fortius angustatum, parcius subtiliter punctatum, pube depressa grisea tenuiter vestitum, piceum, metallico-versicolor, segmentis singulis margine apicali piceo-testaceo. Pectus nigrum. Pedes cum coxis omnibus piceo-testacei.

Tarsi antici maris fortiter, feminae leviter dilatati.

\* 27. *Q. umbrinus*: *Brunneus, capite nigro, abdomine versicolore, antennis totis fusco-piceis, elytris brevioribus, fortiter punctatis.* — Long. 3 lin.

*Er. Col. March. I.* 491. 11.

*Staph. maurorufus* Gyll. *Ins. Suec. II.* 309. 25. — *Sahlb. Ins. Fenn. I.* 317. 22, — *Zetterst. Ins. Lappon.* 61. 15. — *Mannerh. Brachelytr.* 26. 36.

*Philonth. maurorufus* Nordm. *Symb.* 77. 10.

Habitat in Suecia, Germania.

Obscure brunneus, nitidus, capite, thorace scutelloque pernitidis. Antennae capite duplo prope longiores, apicem versus vix crassiores, articulo tertio secundo duplo longiore, 4—10 subobconicis, sensim brevioribus et crassioribus, penultimis crassitie haud brevioribus, ultimo ovato, apice oblique truncato et inferne acuminato, pilosellae, piceae, articulis primis medio nigricantibus. Caput thorace paulo angustius, suborbiculatum, utrinque supra oculos punctis duobus, oblique positus, prope oculi utriusque marginem anteriorem puncto singulo impressum, nigrum, ore infra antennis piceo. Oculi maiusculi, modice prominentes. Thorax, coleopteris prope latior, latitudine summa haud brevior, basi modice, lateribus vix rotundatus, antrorsum subangustatus, apice submarginatus, angulis anterioribus deflexis, acutiusculis, posterioribus obtusis, modice convexus, seriebus dorsalibus punctis tribus, antico remotiore, posterioribus approximatis, compositis, lateribus punctis praeter marginalia antierius utrinque duobus, altero prope seriem dorsalem, altero prope marginem lateralem impressus, niger, margine omni sensim piceo-brunneo. Scutellum laevissimum, piceum. Elytra thorace paulo bre-



viora, confertim fortiter punctata, parum nitida, tenuiter griseo-pubescentia, nigro-brunnea, margine summo apicali obscure testaceo. Abdomen apicem versus sensim angustatum, parcius et minus subtiliter punctulatum, pube parca depressa minus subtili grisea vestitum, lucidum, segmentis singulis virescenti-apice cupreo-versicoloribus, infra apice testaceo-marginatis. Pedes rufo-brunnei.

Tarsi antici maris fortiter, feminae modice dilatati.

\* 28. *Q. maurorufus*: Testaceo-brunneus, capite nigro, abdomine versicolore, antennarum basi pedibusque testaceis, elytris dense subtilius punctulatis. — Long. 3 lin.

Er. Col. March. I. 492. 13.

*Staph. maurorufus* Grav. Mon. 56. 20.

*Staph. attenuatus* Grav. Mon. 61. 31.

*Staph. praecox* Gyll. Ins. Suec. II. 310. 26. — Sahlb. Ins. Fenn. I. 318. 24. — Zetterst. Ins. Lappon. 61. 16. — Mannerh. Brachelytr. 27. 37.

*Emus praecox* Boisd. et Lacord. Faun. Ent. Paris. I. 382. 37.

*Philonth. praecox* Nordm. Symb. 78. 12.

Var. *Thorace elytrisque testaceis*. — Long.  $3\frac{1}{2}$  lin.

*Staph. attenuatus* var. 3 Grav. Mon. 62.

Habitat in Germania, Suecia.

Plerumque testaceo-, rarius fusco- vel nigro-brunneus, nitidus, capite thorace scutelloque pernitidis. Antennae capite sesqui longiores, apicem versus haud incrassatae, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus, subcylindricis, penultimis crassitie haud brevioribus, subobconicis, articulo ultimo oblongo, apice abrupte acuminato, dilute rufo-piceae, articulo primo testaceo. Palpi testacei. Caput thorace paulo angustius, subovatum, supra oculos utrinque punctis duobus oblique positus, ad oculi utriusque marginem interiorem puncto singulo impressum, lateribus pone oculos crebre subtilissime punctulatum, nigrum, infra antennis dilute rufo-piceum. Oculi sat magni, prominuli. Thorax coleopteris paululum angustior, latitudine haud brevior, basi lateribusque modice rotundatus, antrosum subangustatus, apice truncatus, angulis anterioribus subrectis, posterioribus subrotundatis, modice convexus, seriebus dorsalibus punctis tribus sat approximatis compositis, lateribus punctis praeter marginalia anterior utrinque duobus, altero prope seriem dorsalem, altero prope marginem lateralem impressus, testaceo-brunneus, disco plerumque obscuriore. Scutellum laevisimum. Elytra thoracis longitudine, dense subtilius punctata, densius griseo-pubescentia, fusco-testacea, dorso plerumque nigricantia, subopaca. Abdomen apicem versus angustatum, subtiliter punctulatum, lucide irideo-versicolor, segmentis omnibus in individuis dilutioribus apice testaceo-marginatis. Pedes testacei, tarsis anticis maris fortius, feminae leviter dilatatis.

Obs. Var. citata, a Dahllo olim nomine *Staph. limbati signata*, colore thoracis elytrorumque testaceo, praecipue statura duplo maiore insignis, vix autem aliis notis rite distincta.

29. *Q. infuscatus*: *Nigro-piceus, capite nigro, antennarum basi, abdomine segmentorum marginibus pedibusque piceo-testaceis, scutello parce punctato, elytris fusco-testaceis, dorso nigricantibus, parce subtilius punctatis.* — Long.  $2\frac{1}{2}$  lin.

Habitat Lutetiae, coll. Dqm. Aubé.

Habitu omnino praecedentis, at plerumque dimidio minor. Antennae capite sesqui longiores, minus graciles, apicem versus leviter incrassatae, articulo tertio secundo paulo longiore, 4 — 10 subobconicis, sensim paulo brevioribus et crassioribus, penultimis etiam crassitie haud brevioribus, ultimo ovato, apice abrupte acuminato, piceae, basi piceo-testaceae. Palpi testacei. Caput thorace paulo angustius, suborbiculatum, supra leviter convexum, utrinque punctis duobus oblique positis supra oculos, et singulo ad oculi utriusque marginem interiorem impressum, nigrum, nitidum, ore infra antennis piceo-testaceo. Oculi mediocres, modice prominentes. Thorax coleopteris paulo angustior, latitudine haud brevior, antrorsum subangustatus, basi fortius, lateribus modice rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, modice convexus, seriebus dorsalibus punctis tribus sat approximatis compositis, lateribus punctis praeter marginalia antrorsum utrinque duobus, altero prope seriem dorsalem, altero prope marginem lateralem impressus, piceo-brunneus, nitidus. Scutellum nigrum, nitidum, parce obsoletius punctatum. Elytra thorace longiora, parcius subtiliusque punctata, parcius longius griseo-pubescentia, fusco-testacea, nitida, dorso nigricantia. Abdomen apicem versus sensim angustatum, parcius minus subtiliter punctatum, pube longiore depressa minus subtili grisea vestitum, nigro-pilosellum, nigrum, nitidum, segmentis omnibus margine apicali latiore piceo-testaceo. Pectus nigrum. Pedes testacei, coxis anterioribus concoloribus, postici coxis femorumque basi picescentibus.

Tarsi antici maris fortiter dilatati. Femina latet.

Obs. Praecedenti proximus, at statura minore, scutello parce punctato, elytris longioribus abdomineque parcius punctatis distinctus.

- X \* 30. *Q. rufipes*: *Niger, antennis pedibusque rufis, scutello elytrisque subtilissime confertissimeque punctatis, nigro-pubescentibus.* — Long.  $3\frac{1}{2}$  — 4 lin.

*Staph. rufipes* Grav. Micr. 171. 24. Mon. 67. 40. — Latr. Hist. nat. Crust. et Ins. IX. 324. 67.

*Staph. ruficornis* Grav. Mon. 50. 12. — Latr. Hist. nat. Crust. et Ins. IX. 326. 76.

*Emus attenuatus* Boisd. et Lacord. Faun. Ent. Paris. I. 383. 38.

Habitat in Europa australiori.

*Q. attenuato* maior et lator, niger, capite thoraceque politis, nitidissimis. Antennae capite sesqui longiores, graciliores, apicem versus leniter incrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, penultimis crassitie haud brevioribus, ultimo subovato, apice oblique truncato et inferne acuminato, totae rufo-testaceae. Palpi rufi. Caput minutum, thorace duplo angustius, ovatum, utrinque supra oculos punctis duobus, oblique positis, ad oculi utriusque marginem interiorem puncto singulo impressum, lateribus pone oculos subtilissime punctulatum. Oculi ovales, sat magni, modice prominuli. Thorax basi co-

leopterorum latitudine, latitudine baseos haud brevior, antrorsum angustatus, lateribus parum, basi cum angulis posterioribus fortius rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, transversim convexus, seriebus dorsalibus punctis tribus aequaliter distantibus compositis, lateribus punctis praeter marginalia anterieus utrinque duobus oblique positis impressus, punctis subtilioribus. Elytra thoracis longitudine et scutellum subtilissime confertissimeque punctulata, nitidula, pube densa depressa nigra fuscave oblecta. Abdomen apicem versus sensim angustatum, subtiliter punctulatum, subnitidum, cupreo-versicolor, pube densa depressa nigra fuscave vestitum. Pedes cum coxis anterioribus rufi.

Tarsi antici maris modice dilatati, feminae simplices.

\*31. *Q. monticola*: *Niger, nitidulus, antennis pedibusque rufo-testaceis, scutello punctulato, elytris subtiliter punctatis.* — Long.  $3\frac{1}{2}$  lin.

Habitat in montibus Tyrolensibus, Dom. Stentz.

Statura omnino *Q. maurorufi*, et maximis eius individuis aequalis, niger, subnitidus, capite thoraceque nitidissimis. Antennae capite sesqui longiores, graciliores, apicem versus vix incrassatae, articulo tertio secundo longiore, 4 — 10 sensim brevioribus at vix crassioribus, penultimis crassitie paulo longioribus, ultimo ovato, apice oblique truncato et inferne acuminato, totae rufo-testaceae. Palpi rufo-testacei. Caput thorace paulo angustius, suborbiculatum, parum convexus, utrinque punctis duobus oblique positis supra oculos, et puncto singulo ad oculi utriusque marginem interiorem impressum, lateribus pone oculos subtilissime punctulatum. Oculi sat magni et prominuli. Thorax coleopterorum latitudine, latitudine baseos haud brevior, antrorsum subangustatus, lateribus, basi et angulis posterioribus leviter rotundatis, apice truncatus, angulis anterioribus deflexis, subrectis, sat convexus, seriebus dorsalibus punctis tribus aequaliter distantibus compositis, lateribus punctis praeter marginalia anterieus utrinque binis oblique positis notatus. Scutellum punctulatum, fusco-pubescens. Elytra thoracis longitudine, densius subtiliter punctata, nitidula, pube depressa fusca densius vestita. Abdomen apicem versus sensim attenuatum, subtiliter punctulatum, fusco-pubescens, nitidulum, subcupreo-versicolor, subtus segmentorum marginibus tenuiter rufo-piceis. Pedes cum coxis anterioribus rufo-testacei, femoribus posticis medio viridi-micantibus.

Tarsi antici maris fortiter, feminae leviter dilatati.

Obs. A praecedente, cui proximus est, praecipue differt habitu, capite maiore, scutello elytrisque minus subtiliter denseque punctatis.

\*32. *Q. semiobscurus*: *Niger, pedibus cum coxis anticis antennisque testaceis, elytris thorace longioribus, fusco-subaeneis, subtilissime punctulatis, scutello punctulato.* — Long.  $2\frac{3}{4}$  —  $3\frac{1}{4}$  lin.

*Staph. semiobscurus* Marsh. Ent. Brit. 512. 41.

Habitat in Anglia, Dom. Spence, in Sardinia, Dom. Prof. Gené.

Habitu omnino praecedentis, at dimidio minor, *Q. attenuato* proximus, at plerumque paulo maior et robustior. Niger, capite thoraceque nitidissimis. Antennae capite sesqui longiores, graciliores, apicem versus haud incrassa-



tae, articulo secundo tertioque subaequalibus, 4—10 sensim paulo brevioribus, omnibus crassitie longioribus, ultimo oblongo-ovato, apice inferne sensim acuminato. Antennae totae dilute testaceae. Palpi testacei, articulo apicali piceo. Caput thorace paulo angustius, suborbiculatum, parum convexum, utrinque supra oculos punctis duobus oblique positis, ad oculi utriusque marginem interiorem puncto singulo impressum. Oculi magni, prominentes. Thorax basi coleopteris paululum angustior, latitudine baseos vix longior, antrorsum leviter angustatus, lateribus leviter, basi cum angulis posterioribus fortius rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, convexus, seriebus dorsalibus punctis tribus modice distantibus compositis, lateribus punctis praeter marginalia anterieus utrinque duobus, oblique positis, sat distantibus impressus. Scutellum dense subtiliterque punctulatum, subtiliter pubescens. Elytra thorace paulo longiora, confertim subtilissime punctulata, pube subtiliore longioreque depressa grisea densius vestita, virescenti-vel fusco-subaenea, subnitida, margine apicali testaceo. Abdomen apicem versus sensim angustatum, confertim subtilius punctatum, densius fusco-pubescens, subsericans, minus nitidum, supra parum, subtus fortius metallico-versicolor, segmentis ventralibus apice ferrugineo-marginatis. Pedes dilute flavo-testacei, coxis anticis concoloribus, intermediis piceis, posticis nigris, femoribus posticis picescentibus. Alae albo-hyalinae.

Tarsi antici maris modice dilatati, feminae simplices.

Obs. 1. Differt a *Q. attenuato* statura maiore, latiore, elytris longioribus, coxis intermediis piceis.

2. Descriptio Marshamiana manca ex auctoritate Spencei ad hanc speciem allata.

\* 33. *Q. collaris*: *Niger, antennis, pedibus cum coxis anterioribus, thoraceque rufo-testaceis, hoc medio piceo, elytris thoracis longitudine scutelloque subtiliter punctulatis. — Long. 4 lin.*

Habitat in Volhynia, Dom. Schüppel.

Habitu omnino *Q. attenuati*, at plus duplo maior. Antennae capite sesqui longiores, tenues, apicem versus haud incrassatae, articulo tertio secundo paululum longiore et tenuiore, 4—10 sensim paululum brevioribus, teretibus, ultimo oblongo, apice oblique truncato et inferne acuminato, totae rufo-testaceae. Palpi rufo-testacei, articulo ultimo concolore. Caput thorace paulo angustius, orbiculatum, parum convexum, utrinque punctis duobus oblique positis supra oculos, et puncto singulo ad oculi utriusque marginem interiorem notatum, lateribus pone oculos subtilissime punctulatum, nigrum, pernitidum. Oculi permagni, prominentes. Thorax basi coleopteris vix angustior, latitudine baseos haud longior, antrorsum leviter angustatus, basi modice, lateribus leviter rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, posterioribus obtusis, modice convexus, seriebus dorsalibus punctis tribus modice distantibus compositis, lateribus punctis praeter marginalia anterieus utrinque duobus, altero prope seriem dorsalem, altero prope marginem lateralem impressus, nitidus, rufo-testaceus, disco indeterminate nigricante. Scutellum punctulatum, nigrum, nitidum. Elytra thoracis longitudine, confertim subtiliter punctulata, pube densiore depressa nigra vestita, nigra, subnitida, margine summo apicali ferru-

gineo. Abdomen elongatum, apicem versus sensim attenuatum, confertim subtiliter punctulatum, pube subtiliore depressa fusca subsericans, minus nitidum, nigrum, segmentis singulis apice tenuiter rufo-marginatis. Pectus nigrum. Pedes dilute rufo-testacei, coxis anterioribus concoloribus, posticis piceo-testaceis. Alae hyalinae.

Tarsi antichi feminae leviter dilatati. Mas latet.

Obs. Q. attenuato proximus, praeter staturam maiorem autem et colorem thorace minus convexo, angulis posterioribus minus rotundatis diversus.

\* 34. *Q. attenuatus*: *Niger, pedibus cum coxis anterioribus antennisque testaceis, elytris thoracis longitudine, subaeneis, subtilissime punctatis, scutello punctulato.* — Long.  $2\frac{1}{4}$  lin.

*Er. Col. March. I. 493. 14.*

*Staph. attenuatus* Gyll. *Ins. Suec. II. 311. 27.* — Zetterst. *Faun. Lappon. I. 71. 12. Ins. Lappon. 62. 17.* — Sahlb. *Ins. Fenn. I. 318. 25.* — Mannerh. *Brachelytr. 27. 38.* — Runde *Brachelytr. Hal. 6. 15.*

*Staph. maurorufus* Runde *Brachelytr. Hal. 6. 26.*

*Philonth. attenuatus* Nordm. *Symb. 78. 13.*

*Emus scintillans* Boisd. et Lacord. *Faun. Ent. Paris. I. 384. 40.*

Habitat in Germania, Anglia, Suecia.

Niger, nitidus, capite thoraceque pernitidis, elytris nigro-subaeneis, nitidis. Antennae breviusculae, capite paulo longiores, tenuiores, apicem versus haud incrassatae, articulo secundo tertioque subaequalibus, sequentibus sensim paulo brevioribus, ultimo ovato, oblique truncato et inferne acuminato, torae testaceae. Palpi testacei, articulo apicali piceo. Caput thorace parum angustius, rotundatum, utrinque supra oculos punctis duobus oblique positis, ad oculi utriusque marginem interiorem puncto singulo impressum. Oculi permagni, prominentes. Thorax basi fere coleopterorum latitudine, latitudine baseos vix brevior, antrorsum modice angustatus, lateribus leviter, basi cum angulis posterioribus fortiter rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, sat convexus, seriebus dorsalibus punctis tribus modice distantibus compositis, lateribus punctis praeter marginalia utrinque duobus, oblique positis, distantibus impressus. Scutellum subtiliter punctulatum. Elytra confertim subtiliter punctulata, densius subtiliter griseo-pubescentia. Abdomen elongatum, apicem versus sensim angustatum, dense subtiliter punctulatum, pube brevi depressa tenuissima vestitum, parum nitidum, infra magis quam supra metallico-versicolor. Pedes cum coxis anterioribus laete testacei.

Tarsi antichi maris fortiter, feminae modice dilatati.

Variat elytris fusco-testaceis, magis minusve aeneo-micantibus.

Obs. 1. *Staph. maurorufus* Runde secundum collectionem Germanianam, ab auctore citatam, ab hac specie non distinctus.

2. *Staph. 4-punctatum* Zetterst. *Faun. Lappon. I. 76. 24. Ins. Lappon. 64. 34.* cuius specimen singulum in Lapponia a cel. auctore captum est, tantummodo huius speciei varietatem thoracis seriebus dorsalibus 4-punctatis habeo, neque auctori ipsi praeter magnitudinem minorem aliud discrimen observatum. Apud *Philonthus* punctorum in seriebus dorsalibus numerus saepius variabilis, neque in *Quediis* pluribus omnino constans repertus est.

- \* 35. *Q. hyperboreus*: *Niger, pedibus anterioribus cum coxis antennisque piceo-testaceis, elytris thorace brevioribus, nigro-subaeneis, subtiliter punctulatis. — Long. 3 lin.*

*Staph. Fellmanni* Zetterst. Ins. Lappon. 62. 18. (*forte*)

Habitat in Unalaschka, Dom. v. Chamisso.

Statura et summa affinitas praecedentis, niger, capite thoraceque pernitidis. Antennae capite sesqui longiores, tenues, apicem versus haud crassiores, articulo tertio secundo haud longiore et paulo tenuiore, 4 — 10 sensim brevioribus, penultimis longitudine haud crassioribus, ultimo paulo crassiore, subovato, apice oblique truncato et inferne acuminato, totae piceo-testaceae. Palpi testacei, articulo ultimo piceo. Caput thorace dimidio angustius, orbiculatum, parum convexum, utrinque supra oculos punctis duobus oblique positis, ad oculi utriusque marginem anteriorem puncto singulo impressum, lateribus pone oculos subtilissime punctulatum. Oculi permagni, prominentes. Thorax basi coleopterorum prope latitudine, latitudine summa haud longior, antrorsum leviter angustatus, lateribus leviter, basi cum angulis posterioribus fortius rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, sat convexus, seriebus dorsalibus punctis tribus modice distantibus compositis, lateribus punctis praeter marginalia anterieus utrinque duobus, oblique positis, sat distantibus impressus. Scutellum punctulatum. Elytra thorace paulo breviora, crebrius subtiliter punctata, pube subtiliore longiuscula depressa nigra densius vestita, nigro-subaenea, nitida. Abdomen elongatum, apicem versus sensim attenuatum, confertim subtilius punctatum, pube subtili depressa fusca densius vestitum, subsericans, segmentis inferioribus apice piceo-marginatis. Pedes cum coxis antici dilute testacei, intermedii piceo-testacei, postici picei.

Tarsi antici maris fortiter, feminae leviter dilatati.

Obs 1. Elytris brevioribus, rnfus subtiliter punctulatis et pedibus obscurioribus praecipue a praecedente distinctus.

2. *Staph. Fellmanni*, e Lapponia, num recte ad hanc speciem allatus sit, e descriptione manca non satis patet. Descriptio citata haec est: „Affinis et similis *St. attenuato*, sed forte paulo maior, antennis aperte paulo longioribus et pedibus tantum anterioribus testaceis distinctus. Elytra iam fusca iam nigra. Pedes postici fusci vel obscure testacei.”

- \* 36. *Q. virgulatus*: *Niger, nitidus, antennis pedibusque testaceis, scutello punctulato, elytris dense subtiliter punctatis, abdomine tristriato-cine-reo-pubescente. — Long. 2½ lin.*

Habitat in Sardinia, Dom. Prof. Gené.

Habitu et summa affinitas *Q. attenuati*, at minor, *Q. boope* parum maior. Niger, nitidus, capite thoraceque politis. Antennae capite sesqui longiores, articulo tertio secundo paululum longiore et tenuiore, 4—10 sensim paulo brevioribus, penultimis crassitie brevioribus, ultimo apice inferius acuminato, totae laete flavo-testaceae. Palpi testacei. Caput thorace angustius, suborbiculatum, utrinque supra oculos punctis duobus oblique positis, inter oculos ad oculi utriusque marginem anteriorem puncto singulo maiusculo impressum. Oculi permagni, prominuli. Thorax basi coleopteris paulo angustior, antrorsum leviter angustatus, latitudine postica vix longior, basi cum angulis posterioribus fortiter, lateribus parum rotundatus, apice trun-



catus, angulis anterioribus subrectis, deflexis, transversim sat convexus, seriebus dorsalibus punctis tribus modice distantibus compositis, lateribus punctis praeter marginalia anterieus utrinque duobus, altero prope seriem dorsalem, altero prope marginem lateralem. Scutellum punctulatum, pubescens. Elytra thorace paulo longiora, crebre subtiliter punctata, nigro-subaenea, pube subtilissima depressa vestita, immixtis pilis sparsis crassioribus, itidem depressis, fulvis, nitidis, apice fulvo-ciliata. Abdomen apicem versus sensim attenuatum, crebre subtilissime punctatum, nigrum, nitidum, subiridescens, supra pube subtilissima depressa nigra vestitum, lineis tribus cinereo-pubescentibus, in singuli segmenti apice ciliis fulvis interruptis, subtus magis iridescens, segmentis singulis margine posteriore rufo-piceo, tenuiter cinereo-pubescentibus, immixtis pilis rarioribus sparsis fulvis. Pedes laete flavo-testacei, anteriores coxis concoloribus, posteriores femoribus tibiisque fusciscentibus.

Tarsi antici maris fortiter, feminae leviter dilatati.

Obs. Elytrorum et abdominis pubescentia satis distinctus, ceterum *Q.* attenuato et boopi proximus.

\* 37. *Q. boops*: *Niger, nitidus, antennis pedibusque testaceis, elytris thoracis longitudine, subtilius punctatis, scutello punctulato.* — Long. 2 — 2½ lin.

*Er. Col. March. I. 494. 15.*

*Staph. boops* Grav. *Micr. 21. 26. Mon. 71. 54.* — *Latr. Hist. nat. Crust. et Ins. IX. 311. 31.* — *Gyll. Ins. Suec. II. 312. 28.* — *Zetterst. Faun. Lappon. I. 72. 13. Ins. Lappon. 62. 19.* — *Sahlb. Ins. Fenn. I. 318. 26.* — *Mannerh. Brachelytr. 27. 39.*  
*Emus boops* Boisd. et Lacord. *Faun. Ent. Paris. I. 385. 41.*  
*Philonth. boops* Nordm. *Symb. 78. 14.*

Habitat in Germania, Suecia.

*Q.* attenuato proximus, at duplo minor, niger, nitidus, capite thoraceque pernitidis, elytris nigro-piceis, nitidulis. Antennae capite sesqui longiores, articulo tertio secundo paulo minore, 4 — 10 sensim brevioribus, penultimis crassitie parum longioribus, subobconicis, ultimo ovato, apice acuminato; totae testaceae. Palpi testacei. Caput maiusculum, thorace vix angustius, rotundatum, utrinque supra oculos punctis duobus, ad marginem oculi utriusque anteriorem puncto singulo impressum, lateribus pone oculos subtilissime punctulatum. Oculi permagni, prominuli. Thorax coleopteris perparum angustior, latitudine vix longior, antrorsum subangustatus, basi cum angulis posterioribus fortiter, lateribus leviter rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, leviter convexus, seriebus dorsalibus punctis tribus, antico paulo remotiore, compositis, lateribus punctis praeter marginalia anterieus utrinque duobus, altero prope seriem dorsalem, altero prope marginem lateralem impressus. Scutellum subtilissime parce punctulatum. Elytra thoracis longitudine, confertim subtilius punctata, tenuiter griseo-pubescentia. Abdomen elongatum, parce subtiliter punctulatum, parceque tenuiter griseo-pubescentibus, apice nigro-pilosellum, subtus segmentis singulis piceo-marginatis. Pedes cum coxis laete testacei.

Tarsi antici maris fortiter, feminae modice dilatati.

- \* 38. *Q. obliteratus*: *Niger, nitidus, antennis, thoracis lateribus, abdominis segmentorum marginibus pedibusque testaceis, elytris dense punctatis, pallidis, plaga subsuturali nigra.* — Long. 3 lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura omnino *Q. attenuati*, niger, nitidus, capite thorace scutelloque politis. Antennae capite thoraceque paulo breviores, articulo tertio secundo tenuiore et sesqui longiore, 4 — 10 sensim paulo brevioribus at vix crassioribus, penultimis crassitie vix longioribus, ultimo apice distincte oblique emarginato et inferne acuminato, saturate rufo-testaceae, basi dilutiores. Palpi testacei. Caput thorace angustius, orbiculatum, utrinque supra oculos punctis duobus, ad oculi utriusque marginem interiorem puncto singulo impressum, lateribus supra oculos confertissime subtiliter punctulatum. Oculi magni. Thorax coleopteris paulo angustior, antrosum subangustatus, latitudine postica fere longior, basi cum angulis posterioribus fortiter, lateribus leviter rotundatus, apice truncatus, angulis anterioribus subrectis, deflexis, transversim sat convexus, seriebus dorsalibus punctis tribus, anteriore remotiore, compositis, lateribus punctis praeter marginalia antierius utrinque duobus, altero prope seriem dorsalem, altero prope marginem lateralem, nigro-piceus, lateribus testaceis. Scutellum laevissimum. Elytra thorace paululum longiora, subdepressa, dense minus fortiter punctata, pube longiore subdepressa grisea vestita, pallide lutea, plaga longitudinali prope suturam, parum distincta, nigra. Abdomen modice elongatum, apicem versus sensim angustatum, sat crebre subtilius punctatum, pube longiore depressa fulvescenti-grisea densius vestitum, nigrum, nitidulum, subiridescens, segmentis singulis testaceo-marginatis. Pedes laete testacei, coxis anterioribus totis, posticis apice concoloribus.

Tarsi antici maris fortiter, feminae leviter dilatati.

Obs. Intermedius inter plures species diffines: habitu omnino et capitis forma cum *Q. attenuato* et scintillante congruit, ab illo autem scutello laevissimo, ab hoc fronte inter oculos bipunctata differt; ab utroque antennarum structura discedit, qua *Q. maurorufus* appropinquatur.

- \* 39. *Q. scintillans*: *Nigro-aeneus, nitidus, antennarum basi pedibusque testaceis, elytris parcius fortiter punctatis, fronte inter oculos punctis 4 transversalibus impressa.* — Long. 2½ lin.

*Er. Col. March. I. 494. 16.*

*Staph. scintillans Grav. Mon. 70. 53.*

Habitat in Germania.

Statura omnino praecedentium, at minus elongatus, obscure aeneus, capite, thorace scutelloque nitidissimis. Antennae capite paulo longiores, apicem versus leviter incrassatae, articulo tertio secundo tenuiore at parum brevior, 4 — 10 sensim paulo brevioribus et crassioribus, ultimo oblongo-ovato, apice inferne acuminato, rufo-piceae, articulo primo testaceo. Palpi picei. Caput thorace angustius, rotundatum, utrinque supra oculos punctis duobus, inter oculos serie transversali e punctis quatuor, aequaliter distantibus, lateralibus in oculi ipso margine situs obsoletioribus, intermediis sat profundis notatum, lateribus pone oculos subtilissime punctulatis. Oculi magni, prominuli. Thorax basi coleopteris latitudine aequalis, latitudine

summa haud brevior, antrorsum leviter angustatus, basi et lateribus subaequaliter rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, posterioribus subrotundatis, leviter convexus, seriebus dorsalibus punctis tribus modice approximatis compositis, lateribus punctis praeter marginalia antius utrinque 4 impressus. Scutellum laevissimum. Elytra thorace paulo longiora, minus dense fortiter punctata, albido-pubescentia. Abdomen apicem versus sensim attenuatum, subtiliter parcius punctulatum, pube longiore depressa fusco-grisea densius vestitum, nigro-pilosellum, subtus lucide metallico-versicolor, segmentis singulis piceo-marginatis. Pedes cum coxis anticis testacei, tarsis anticis in utroque sexu fortiter dilatatis.

- \* 40. *Q. lucidulus*: *Viridi-aeneus, nitidus, antennarum basi, pedibus, elytris abdomineque testaceis, elytris dense subtilius punctatis, fronte inter oculos punctis 4 transversalibus impressis.* — Long.  $2\frac{1}{2}$  lin.

*Er. Col. March. I.* 495. 17.

Habitat in Germania.

Statura et summa affinitas praecedentis. Antennae capite paulo longiores, crassiusculae, articulo secundo tertioque subaequalibus, 4 — 10 sensim paulo crassioribus at vix brevioribus, subobconicis, ultimo ovato, apice inferius acuminato, rufo-piceae, basi testaceae. Os testaceum. Caput thorace angustius, rotundatum, lucide viridi-aeneum, politum, utrinque supra oculos punctis duobus, inter oculos serie transversali e punctis quatuor aequaliter distantibus impressum. Oculi magni, prominuli. Thorax basi coleopterorum latitudine, latitudine summa vix brevior, antrorsum angustatus, basi fortius, lateribus leviter rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, posterioribus subrotundatis, modice convexus, seriebus dorsalibus punctis tribus modice approximatis compositis, lateribus punctis praeter marginalia antius utrinque 4 impressus, lucide viridi-aeneus, politus. Scutellum laevissimum. Elytra thorace paulo longiora, fusco-testacea, supra aeneo-nitidula, confertim subtilius punctata, pube lutescenti-grisea minus subtili vestita. Abdomen apicem versus sensim attenuatum, confertim subtiliter punctatum, pube depressa, minus tenui, lutescenti-grisea dense vestitum, nigro-pilosellum, obscure testaceum, virescenti-nitidulum, linea longitudinali dorsali angusta viridi. Pectus nigricans. Pedes testacei, coxis femoribusque anticis flavis, tarsis anticis in utroque sexu fortiter dilatatis.

#### Species mihi invisae.

41. *Q. peregrinus*: *Nitidus, nigricans, antennis, palpis, elytris, pedibus, thoracis lateribus et segmentorum marginibus rufescentibus.* — Long. 3 lin.

*Staph. peregrinus Grav. Mon.* 53. 17.

Habitat in Carolina.

Habitu et thoracis forma *Q. fulgido* (nitido) affinis, capite autem ovato et colore satis differt. Caput ovatum, thoracis longitudine, sed eo angustius, nitidum, punctis perpaucis maioribus impressum, nigrum, antennis (quarum articulus ultimus deest) capite paulo longioribus palpisque rufis. Thorax punctis quibusdam impressus, niger, lateribus rufescentibus. Coleoptera



quadrata, subpilosa, piceo-rufescentia. Abdomen segmentorum marginibus rufis, supra nigricans, infra fuscum. Pedes testaceo-rufi, tarsis anticis dilatatis. (Grav.)

42. *Q. tenellus*: *Nigro-fuscus, abdomine pedibusque pallidioribus, elytris testaceis, thorace seriebus dorsalibus 4-punctis.* — Long.  $3\frac{1}{2}$  lin.

*Staph. tenellus* Grav. Mon. 54. 18.

*Staph. nitidi* var. 13. Grav. Micr. p. 32.

Habitat in Germania.

Statura graciliore, colore et thoracis punctura satis a *Q. fulgidus* (nitido) distinctus est. Caput suborbiculatum, latitudine paulo longius, thorace minus, nigrum, nitidissimum, in angulis baseos punctum, punctis 3 distinctioribus, antennis fuscis. Thorax orbiculatus, latitudine fere eadem ac in *Q. fulgidus*, nigro-fuscus, marginibus pallidioribus, puncto marginali uno, serie laterali 3-puncta, seriebus dorsalibus 4-punctis, punctis autem subconfluentibus. Praeterea punctum marginale in angulo baseos, ante humeros coleopterorum impressum est, distinctius quam in reliquis huius familiae Staphylinis. Scutellum nigro-fuscum. Coleoptera quadrata, dilute testacea, nitida, subtiliter puncta, pilis tenuissimis vestita. Abdomen coleopteris triplo longius, pilosellum fuscum, subtiliter punctum. Pedes fusi, tarsis anterioribus pallidioribus, anticis brevibus. (Grav.)

Obs. Forte huius generis.

43. *Q. rufo-cinctus*: *Niger, nitidus, elytris glabris, rufo-marginatis, haud striato-punctatis, abdomine subcoerulescente, segmentorum marginibus rufo-ferrugineis.*

*Staph. rufocinctus* Mannerh. Brachelytr. 26. 33.

*Philonth. rufocinctus* Nordm. Symb. 79. 21.

Habitat in Fennia meridionali.

Statura et magnitudo *Q. impressi*, a quo differt elytris haud striato-punctatis: an igitur species diversa? (Mannerh.)

44. *Q. maurus*: *Niger, nitidus, ore, antennis extrorsum pedibusque piceo-ferrugineis, elytris profunde punctatis, antennis thorace brevioribus, mediocribus.* — Long.  $2\frac{3}{4}$  lin.

*Staph. maurus* Sahlb. Ins. Fenn. I. 317. 23. — Mannerh. Brachelytr. 26. 35.

*Philonth. maurus* Nordm. Symb. 79. 22.

Habitat in Fennia meridionali.

Magnitudine et statura *Q. umbrinus* (maurorufus S.) similis, colore, antennis brevioribus et oculis minoribus ab illo distinctus. Caput thorace paulo angustius, orbiculatum, nigrum, nitidum, laeve, puncto pone singulum oculum magno et quibusdam minoribus in angulis verticis. Palpi et reliquae oris partes piceo-ferrugineae. Oculi mediocres, nigri. Antennae thorace breviores, extrorsum paulo crassiores, articulis tribus baseos nigris, nitidis, longioribus, ceteris piceo-ferrugineis, brevibus, ultimo longiore. Thorax antice submarginatus, angulis deflexis, lateribus et postice exacte orbiculato-rotundatus, supra convexus, niger, nitidus, laevis, punctura ut in

*Q. umbrino*. Scutellum triangulare, nigrum, nitidum, laevisimum. Elytra thoracis medio nonnihil angustiora, apice oblique truncata, supra depressa, nigra, nitida, profunde sed parcius punctata, tenue pubescentia. Abdomen pone elytra longitudine fere reliqui corporis, subnitidum, nigrum, punctulatum, pubescens. Corpus subtile nigrum, subnitidum, punctulatum, pubescens. Pedes piceo-ferruginei. (Sahlb.)

45. *Q. vicinus*: *Niger, nitidus, pedibus abdomineque testaceis, hoc cupreo-versicolore, elytris pallide flavis, fortiter punctatis.*

*Emus vicinus* Boisd. et Lacord. Faun. Ent. Paris. I. 383. 39.

Habitat Lutetiae.

*Q. attenuato* paulo brevior et latior. Caput nigrum, nitidum, in utroque sexu aequale, thorace angustius, oblongum, laevisimum, vertice punctis duobus, fronte ad oculi utriusque marginem interiorem singulo, et utrinque supra oculos tribus vel quatuor impressum. Thorax capiti concolor, amplius, suborbicularis, laevisimus, seriebus dorsalibus 3-punctatis, praeterea punctis duobus approximatis nonnunquam confluentibus prope marginem lateralem, nonnullis sparsis versus margines disci impressus, basi leviter punctata. Scutellum triangulare, nigrum, nitidum, laevisimum. Elytra pallide flava, elongata, apice oblique truncata, fortiter punctata, pubescentia. Abdomen modice elongatum, apice attenuatum, subtiliter rugulosum et pubescens, testaceum, laete aeneo cupreoque versicolor. Pedes obscurius testacei, tibiis intermediis perparum spinulosi. (Sec. Boisd. et Lacord.)

Obs. A *Q. lucidulo* fronte inter oculos utrinque 1-punctata discedit.

### **Astrapaenus Grav.**

*Grav. Mon. p. 199. — Latr. Gen. Crust. et Ins. I. p. 284. — Guér. Iconogr. Regn. An. Ins. t. 9. f. 2.*

*Palpi utrique articulo ultimo securiformi.*

*Ligula rotundata, medio incisa, paraglossis brevior.*

*Pedes intermediis approximatis.*

Corpus mediocriter, lineare, alatum. Caput thorace minus, ovatum, basi vix constrictum, declinatum, oculis haud prominulis. Labrum breve, transversum, bilobum, lobis rotundatis, limbo laterali membranaceo, apice setis rigidis obsitum. Mandibulae capite fere breviores, fortes, muticae, apice aduncae, acutae. Maxillae (t. IV. f. 11. b.) malis coriaceis, exteriore apice, interiore intus barbatis. Palpi maxillares mediocres, articulo primo minore, tenui, cylindrico, secundo tertioque crassiusculis, subobconicis, tertio paulo brevior, quarto maris fortiter, feminae leviter securiformi. Labium (t. IV. f. 11. a.) mento transversum, corneo, ligula brevi, rotundata, medio leviter incisa, membranaea, paraglossis longioribus, coriaceis, intus ciliatis. Antennae filiformes, articulo ultimo apice oblique submarginato. Thorax coleopteris fere latior, antrorsum leviter angustatus, latitudine brevior, transversim convexus, lateribus deflexis basiue rotundatus, basi utrinque subsinuatus, antice truncatus. Scutellum triangulare. Elytra thorace paulo longiora, truncata. Abdomen longiusculum, lineare, apice attenuatum. Pedes breviusculi, antici breves, femoribus compressis, subtile sinuatis, tibiis brevibus,

intus breviter parceque spinulosis, tarsis articulis 4 primis in utroque sexu dilatatis, subtus spongioso-tomentosis, pedes posteriores tibiis spinulosis, tarsis articulo primo ultimoque leviter elongatis. Coxae intermediae contiguae.

### Observationes.

1. Habitus omnino Quedii, palporum structura, et ligula medio incisa imprimis distinctus.

2. Mas palporum maxillarium articulo ultimo fortius securiformi, tarsis anticis fortius dilatatis, abdominisque segmento sexto subtus apice leviter emarginato distinctus.

\* 1. *A. Ulmi: Niger, nitidus, elytris abdominisque segmento quinto apice rufis.* — Long. 5 — 6 lin.

*Grav. Micr.* 199. 1. — *Panz. Faun. Germ.* 88. 4. — *Latr. Hist. nat. Crust. et Ins.* IX. p. 288. t. 79. f. 3. *Gen. Ins.* I. 284. 1. — *Mannerh. Brachelytr.* 19. 1. — *Nordm. Symb.* 16. 1.

*Staph. Ulmi Rossi Faun. Etr.* 1. 248. (*Ed. Hellw.* 307.) 611. t. 5, f. 6. — *Oliv. Ent.* III. 42. 17. t. 4. f. 37. — *Grav. Mon.* 47. 7.

*Staph. Ulmineus Fab. Syst. El.* II. 595. 28.

*Astrapaeus ulmineus Boisd. et Lacord. Faun. Ent. Paris.* I. 356.

Habitat in Gallia, Italia, Dalmatia, Hungaria („sub cortice Ulmi primo vere” Rossi).

Niger, capite thoraceque nitidissimis. Antennae capitis thoracisque longitudine, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus at non crassioribus, leviter obconicis, basi rufae, apice ferrugineae. Palpi rufi. Caput parce subtilissime obsolete punctulatum, supra oculos utrinque punctis duobus maioribus, ad oculi utriusque marginem interiorem puncto singulo impressum. Thorax parce omnium subtilissime punctulatus. Scutellum laeve. Elytra saturate rufa, nitida, subtiliter punctulata, stria suturali impressa, et in medio dorsi punctorum maiorum distantium serie longitudinali notata. Abdomen minus nitidum, crebre punctulatum, breviter nigro-pubescent, segmento quinto apice late rufo. Pedes nigri, tibiis tarsisque rufo-piceis.

### Euryporus Er.

*Er. Col. March.* I. p. 496.

*Pelecyphorus Nordm. Symb.* p. 13. t. 1. f. 5.

*Palpi maxillares filiformes, labiales articulo tertio securiformi.*

*Ligula rotundata, integra, paraglossis brevior.*

*Pedes intermediū approximati.*

Corpus mediocre, oblongum, alatum. Caput thorace multo minus, oblongo-ovatum, basi vix constrictum, deflexum, oculis vix prominulis. Labrum latum, transversum, medio profunde incisum, totum corneum, apice setulosum. Mandibulae fortiter arcuatae, acutae, basi unidentatae. Maxillae (t. IV. t. 10. b.) malis coriaceis, minutis, exteriore apice, interiore intus barbatis. Palpi maxillares mediocres, articulo primo minuto, brevi, tertio secundo paulo brevior, quarto secundo aequali, cylindrico. Labium (t. IV. f. 10. a.) mento transversum, corneo, ligula brevissima rotundata, integra,



membranea, paraglossis minutis, membranaceis, intus ciliatis, ligulam superantibus. Palpi labiales articulo primo secundoque subaequalibus, brevibus, subobconicis, tertio securiformi. Antennae filiformes, articulo ultimo apice oblique truncato. Thorax coleopteris paululum latior vel iisdem subaequalis, antrorsum angustatus, latitudine brevior, lateribus et basi rotundatus, apice truncatus, transversim convexus. Scutellum rotundato-triangulare. Elytra thoracis longitudine, truncata. Abdomen latiusculum, apice attenuatum. Pedes mediocres, graciliores, tibiis omnibus spinulosis, tarsis gracilibus, articulo primo ultimoque leviter elongatis, tarsis anticis etiam in utroque sexu simplicibus. Coxae intermediae contiguae.

#### O b s e r v a t i o n e s.

1. Habitus fere Quedii, at caput minus, praeterea palporum et tarsorum anteriorum structura distinctus. Paykull et Gyllenhal hoc genus minus apte ob palporum formam cum Oxyporo coniungunt: proxime accedit ad Astrapaeum, a quo praecipue palpis maxillaribus teretibus, mandibulis basi dentatis tarsisque anticis simplicibus discedit.

2. Mas palporum labialium articulo ultimo fortius securiformi abdominisque segmento sexto subtus apice leviter emarginato ab altero sexu differt.

3. Victus in silvis sub musco.

\* 1. *E. picipes*: *Niger, nitidus, thorace laevissimo, antice biserialim tripunctato*. — Long. 4 —  $4\frac{1}{2}$  lin.

*Er. Col. March. I.* 496. 1.

*Oxyporus picipes* Payk. *Faun. Suec. III.* 426. 2. — *Gyll. Ins. Suec. II.* 443, 2.

*Pelecyphorus picipes* Nordm. *Symb.* 14. 1.

Habitat in Suecia, Germania.

*Niger, nitidus*, capite thoraceque nitidissimis. Antennae capitis thoracisque prope longitudine, filiformes, apicem versus haud incrassatae, articulo tertio secundo sesqui longiore, 4 — 10 sensim brevioribus at haud crassioribus, 4 et 5 subcylindricis, sequentibus leviter obconicis, penultimo crassitie vix longiore, ultimo apice abrupte acuminato, nigrae, articulis duobus basalibus rufis. Os rufum. Caput laevissimum, supra oculos punctis utrinque duobus impressum. Thorax basi coleopteris paulo latior, latitudine baseos paulo brevior, antrorsum angustatus, basi cum angulis posterioribus et lateribus rotundatus, apice truncatus, angulis anterioribus leviter deflexis, subrectis, convexus, laevissimus, antice seriebus dorsalibus e punctis tribus sat approximatis, lateribus praeter marginalia punctis utrinque duobus, altero prope seriem dorsalem, altero prope marginem lateralem notatus. Scutellum laeve. Elytra thorace paulo breviora, crebre punctata, nigro-pubescentia. Abdomen punctulatum, breviter rigideque parce nigro-pubescentis, metallico-versicolor. Pedes nigri, tibiis piceis, tarsis rufescentibus.

\* 2. *E. puncticollis*: *Niger, nitidus, thorace crebre punctato, linea media laevi*. — Long.  $4\frac{1}{2}$  lin.

Habitat in America septentrionali.

Praecedente corpore medio paulo latior, niger, nitidus. Antennae capite thoraceque breviores, apicem versus sensim incrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim crassioribus at vix brevioribus, 6 — 10 cylindricis, penultimis crassitie paulo brevioribus, ultimo subgloboso.

apice leviter impresso et inferne acuminato, nigrae, apice summo ferrugineo. Os piceum. Caput minutum, crebre punctatum, fronte medio laevi. Thorax coleopteris paulo angustior, latitudine summa paulo brevior, apice truncatus, basi, lateribus et angulis omnibus, anterioribus leviter deflexis, rotundatis, leviter convexus, crebre punctatus, linea media longitudinali laevi subelevata. Scutellum punctulatum. Elytra thoracis longitudine, sat crebre punctata. Abdomen apicem versus sensim leniter angustatum, supra punctulatum, segmentis tribus primis basi utrinque striola abbreviata impressis, ultimis duobus subtilius punctulatis, parce tenuiterque griseo-pubescent, subtus segmentis singulis basi fortius densiusque punctatis, longius cinereo-pubescent. Pedes nigro-picei, tarsis rufescentibus.

Abdominis segmentum inferius quintum in mare ante apicem impressum, ante foveolam elevato-biplicatum. Femina latet.

### **Oxyporus Fab.**

*Fab. Syst. Ent. p. 267. — Grav. Micr. p. 151. — Latr. Gen. Crust. et Ins. I. p. 283.*

*Palpi maxillares filiformes, labiales articulo tertio lunato.*

*Ligula biloba, paraglossis nullis.*

*Pedes intermediï distantes.*

Corpus oblongum, infra modice convexum, alatum. Caput magnum, exsertum, basi parum constrictum, oculis parvis, rotundatis, parum prominulis. Labrum transversum, acute emarginatum. Mandibulae elongatae, porrectae, falcatae, decussatae, acutae, muticae. Maxillae (t. V. f. 6. b.) mala exteriore coriacea, apice rotundato, membraneo, margine barbato, interiore membranea, margine exteriore corneo, interiore pilis brevissimis dense ciliato. Palpi maxillares leviter elongati, articulo primo minuto, secundo elongato, tertio hoc paulo breviori, obconico, quarto hoc paulo breviori, apice subacuminato. Labium (t. V. f. 6. a.) mento corneo, transverso, utrinque antrorsum producto, portione intermedia inter mentum et ligulam coriacea, apice bicuspi, ligula biloba, lobis acuminatis. Palpi labiales articulo primo breviori, secundo sat elongato, tertio dilatato, apice emarginato, transversim triangulari vel semilunari. Antennae breviores, rectae, apicem versus sensim compressae dilatataeque, articulis penultimis 5 utrinque dilatatis, transversis, ultimo breviter ovato. Thorax coleopteris paulo angustior, basi rotundatus, apice truncatus, lateribus subtilissime marginatus, basin versus angustatus. Scutellum triangulare. Elytra apice truncata, angulo exteriore leviter rotundato. Abdomen corpore dimidio brevius, supra planum, lateribus late marginatum, subtus leviter fornicatum. Pectus amplum. Pedes leviter elongati, graciles, intermediï pectoris lateribus inserti, late distantes, tibiis omnibus muticis, tarsis breviusculis, articulo primo longiore, secundo leviter elongato, tertio quartoque decrescentibus, quinto his coniunctis sublongiore.

### **O b s e r v a t i o n e s.**

1. Caput et thorax laevia, hoc nonnunquam transversaliter impresso. Elytra stria suturali impressa. Abdominis segmenta 5 prima dorsalia prope marginem utrinque linea obliqua subarcuata insculpta.

2. Mas capite grandiore instructus esse solet: praeterea a femina differt abdominis segmento sexto apice leviter emarginato, septimo parvo conspicuo.

3. Victus in Agaricis et Boletis.

- \* 1. *O. rufus*: *Niger, thorace, abdomine pedibusque rufis, illo apice, his basi nigris, elytris macula magna humerali testacea.* — Long.  $2\frac{1}{2}$ —4 lin.

*Fab. Syst. Ent.* 267. 1. *Mant. Ins.* 222. 1. *Ent. Syst. I. II.* 531. 1. *Syst. El. II.* 604. 1. — *Rossi Faun. Etr.* 1. 252. 624. *Ed. Helv. I.* 313. 624. — *Payk. Faun. Suec. III.* 425. 1. — *Cederh. Faun. Ingr.* 120. 367. — *Walken. Faun. Paris. I.* 274. 1. — *Panz. Faun. Germ.* 16. 19. *Ent. Germ.* 359. 1. — *Oliv. Ent. III.* 43. 4. 1. t. 1. f. 1. *Encycl. meth. VIII.* 607. 2. — *Grav. Micr.* 151. 1. *Mon.* 235. 2. — *Latr. Hist. nat. Crust. et Ins. IX.* p. 358. 1. t. 80. f. 3. *Gen. Crust. et Ins. I.* 284. — *Gyll. Ins. Suec. II.* 442. 1. — *Sahlb. Ins. Fenn. I.* 405. 1. — *Guér. Iconogr. Regn. An. Ins. t. 9. f. 1.* — *Mannerh. Brachel.* 19. 1. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 354. 1. — *Runde Brachelytr. Hal.* 1. 1. — *Nordm. Symb.* 7. 1. — *Er. Col. March. I.* 498. 1.

*Staph. rufus* *Linn. Faun. Suec. n.* 844. *Syst. nat. I. II.* 684. 6. — *Degeer Ins. IV.* 24. 10. t. 1. f. 11—13. — *Schrank En. Ins. Austr.* 232. 438. — *Scop. Ent. Carn.* 101. 307. — *Müll. Prod.* 99. 1116. — *Goeze Ent. Beitr. I.* 717. 6. — *De Vill. Ent. I.* 412. 6. — *Fourcr. Ent. Paris. I.* 170. 22. — *Payk. Mon. Staph.* 18. 10. — *Marsh. Ent. Brit.* 502. 14.

*Staph. nigrofulvus* *Fourcr. Ent. Par. I.* 173. 32.

*Staph. octavus* *Schaeff. Icon t.* 85. f. 3.

*Le Staph. jaune à tête, étuis et anus noirs* *Geoffr. Hist. Ins. I.* 370. 22.

Habitat in Europae boletis.

Antennae et palpi rufo-testacei. Labrum testaceum. Caput maris thorace paulo latius, feminae eiusdem latitudine, parce subtilissime punctulatum, nigrum, nitidum, apice utrinque leviter impressum. Thorax coleopteris paulo angustior, latitudine dimidio brevior, leviter convexus, aequalis, laevis, rufus, nitidus. Scutellum nigrum. Elytra thorace paulo longiora, nigra, nitida, macula magna quadrata humerali testacea, disco punctorum seriebus duabus longitudinalibus minus regularibus, punctisque nonnullis sparsis impressa. Abdomen rufum, segmentis ultimis duobus totis, antepenultimo apice nigris. Pectus nigrum. Pedes rufo-testacei, femoribus basi, trochanteribus coxisque nigris.

Variat rarius pedibus totis cum coxis testaceis.

- \* 2. *O. maxillosus*: *Testaceus, capite, thorace pectoreque piceis, elytris angulo apicali exteriore nigro.* — Long. 3 — 4 lin.

*Fab. Ent. Syst. I. II.* 531. 2. *Syst. El. II.* 605. 2. — *Cederh. Faun. Ingr.* 120. 2. t. 3. f. 6. — *Walk. Faun. Paris. I.* 274. 2. — *Panz. Faun. Germ.* 16. 20. *Ent. Germ.* 359. 2. — *Grav. Micr.* 152. 2. *Mon.* 235. 5. — *Latr. Hist. nat. Crust. et Ins. IX.* 358. 2. — *Oliv. Encycl. meth. VIII.* 607. 3. — *Sahlb. Ins. Fenn. I.* 405. 2. — *Curt. Brit. Ent. IX.* t. 418. — *Mannerh. Brachelytr.* 19. 2. — *Nordm. Symb.* 7. 2. — *Er. Col. March. I.* 498. 2.

*Var. a: Abdomine nigro, pedibus basi interdum picescentibus.*

*Oxyp. angularis* *Gebler Ledebour It. App. III.* 69. 1.

*Var. b: Abdomine femoribusque nigris.*

*Oxyp. Schönherri* *Mannerh. Brachelytr.* 19. 3. — *Sahlb. Ins. Fenn. I.* 406. 3. — *Nordm. Symb.* 7. 3.

Habitat in Germaniae agaricis, passim.

Antennae et palpi rufo-testacei, labrum testaceum. Caput maris thorace maius, feminae ei aequale, parce subtilissime punctulatum, nitidum, apice utrinque obsolete impressum, nigro- vel brunneo- vel rufo-piceum, vel piceo-testaceum, ore dilutiore. Thorax coleopteris dimidio fere angustior,



latitudine tertia parte brevior, modice convexus, laevissimus, dorso ante medium utrinque transversim impressus, capitis colore, disco saepe dilutior. Scutellum testaceum. Elytra thorace sesqui longiora, pallide testacea, angulo apicis exteriori late nigro, stria suturali seriebusque duabus dorsalibus impressis sat regulariter punctatis, punctisque nonnullis prope striam dorsalem anteriorem impressis. Abdomen rufescenti-testaceum, supra ante apicem nonnunquam picescens, subtus segmentis penultimis duobus utrinque piceis, raro abdomen totum nigro-piceum. Pectus piceum. Pedes cum coxis flavo-testacei.

Obs. *O. angularis*, a Dom. Gebler in Sibiria detectus, etiam in Germania licet raro obvius, indubia *O. maxilloso* varietas. *O. Schönherrii* etiam in Fennia in agaricis lectus, huius speciei varietas videtur, ab *O. angulari* nonnisi femoribus nigris, et a *O. maxilloso* genuino, quantum e descriptione accuratissima Sahlbergiana patet, nonnisi abdominis pedumque colore distinctus, quam ob rem pro mera huius speciei varietate eum habere non dubito.

\* 3. *O. Mannerheimii*: *Niger, tarsi testaceis, antennis obscure testaceis, articulo primo nigro.* — Long.  $3\frac{2}{3}$  lin.

Gyll. Ins. Suec. IV. 495. 1 — 2. — Mannerh. Brachelytr. 19. 4. — Sahlb. Ins. Fenn. I. 407. 4. — Germ. Faun. Ins. Europ. XVI. 5. — Nordm. Symb. 19. 4.

Habitat in Lapponiae boletis, Dom. Sturm et Schüppel.

Statura omnino praecedentium, nigerrimus, nitidus. Antennae rufo-testaceae, articulo primo piceo. Labrum pallide testaceum. Palpi picei. Caput thorace parum latius, parce subtilissime punctulatum, apice utrinque impressum. Thorax basi coleopteris vix dimidio angustior, latitudine dimidio fere brevior, convexus, basi subtiliter canaliculatus, ceterum aequalis, laevissimus. Elytra thorace sesqui longiora, humeris prominulis, lateribus omnino rectis, dorso depressa, in medio disco minus regulariter biseriatim, introrsum vage obsolete punctata. Tarsi dilute testacei.

Variat thorace, elytris corporeque subtus piceis.

\* 4. *O. maior*: *Niger, tarsi testaceis, elytris flavis, margine, sutura vittae tenui nigris.* — Long.  $4\frac{1}{2}$  lin.

Grav. Mon. 234. 1. — Oliv. Encycl. meth. VIII. 607. 1.

Habitat in America septentrionali, e mus. Knoch.

Praecedentibus duplo maior, niger, nitidus. Antennae fusco-testaceae, articulo primo nigro. Palpi picei. Caput maximum, parce obsolete punctulatum, apice obsolete 3-impressum. Thorax basi coleopteris dimidio angustior et longitudine haud latior, lateribus parum rotundatus, antrorsum sensim incrassatus, dorso ante apicem et in medio leviter transversim impressus. Elytra thorace paulo longiora, humeris prominulis, pallide testacea, sutura antrorsum, margine laterali apice dilatatis, margineque tenui apicali nigris, dorso striis duabus obliquis punctatis approximatis, punctisque nonnullis sparsis utrinque impressa, striarum interstitio nigro elytri aream flavam discendente. Abdomen et pectus nigra, immaculata. Pedes nigri, tarsi dilute testaceis.

- \* 5. *O. femoralis*: *Niger, nitidus, elytris pallide flavis, sutura margineque laterali nigris, tibiis tarsisque testaceis.* — Long.  $3\frac{3}{4}$  lin.

*Grav. Micr.* 196. 3. *Mon.* 235. 7. — *Latr. Hist. nat. Crust. et Ins.* IX. 359. 5.  
— *Oliv. Encycl. meth.* VIII. 608. 6.

Habitat in America septentrionali.

Statura omnino *O. maxilloso* et minoribus eius individuis aequalis, niger, nitidus. Antennae fusco-piceae, articulo primo nigro, sequentibus rufescentibus. Palpi maxillares picei, labiales testacei. Caput maris thorace latius et maius, feminae eiusdem fere latitudine, basi subtiliter punctatum, ante apicem obsolete longitudinaliter impressum. Thorax basi coleopteris dimidio angustior, antrorsum leviter rotundato-incrassatus, latitudine paulo brevior, leviter convexus, ante apicem utrinque obsolete transversim impressus. Elytra humeris antrorsum prominulis, pallide flava, sutura basi, limboque laterali apice dilatato nigris, disco punctis minus regulariter biserialis nonnullisque sparsis subtilius impressa, angulo apicali exteriori late subtiliter aciculato-ruguloso. Abdomen et pectus nigra. Pedes nigri, tibiis tarsisque testaceis, illis basi apiceque summis nigris.

Variat 1, Pedibus flavis, coxis, trochanteribus, femorum tibiisque summa basi summoque apice nigris.

2, Pedibus totis flavis, coxis solis piceis.

- \* 6. *O. vittatus*: *Niger, nitidus, elytris pedibusque testaceis, illis sutura margineque laterali nigris, his immaculatis.* — Long. 3 lin.

*Grav. Micr.* 195. 2. *Mon.* 235. 6. — *Latr. Hist. nat. Crust. et Ins.* IX. 359. 4.  
— *Oliv. Encycl. meth.* VIII. 608. 5.

*Var.*: *Pectore ventrequ testaceis.*

*Oryp. cinctus* *Grav. Micr.* 196. 4. *Mon.* 235. 8. — *Latr. Hist. nat. Crust. et Ins.* IX. 359. 6. — *Oliv. Encycl. meth.* VIII. 608. 7.

Habitat in Pennsylvania, Knoch.

Niger, nitidus. Antennae parvae, fusco-piceae, basi testaceae. Palpi testacei. Labrum testaceum. Mandibulae mediocres. Caput thorace paulo maius, convexum, basi subtiliter obsoleteque punctatum, apice leviter biimpressum. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus fortius rotundatus, modice convexus, lateribus medio obsolete impressus. Elytra thorace sesqui longiora, humeris antrorsum prominulis, testacea, sutura basi leviter dilatata, limbo laterali apice dilatato margineque tenui apicali nigris, disco punctis biserialis aliisque sparsis impressa. Abdomen segmentis singulis margine summo testaceo. Pedes cum coxis toti testacei.

Variat 1, Thorace dorso medio rufo-piceo, pectore toto nigro, abdomine segmentis tribus primis piceis, limbo laterali et apicali late testaceis, quarto quintoque totis nigris, sexto testaceo.

2, Thorace supra nigro, basi apiceque nigro, metasterno testaceo, pectore reliquo nigro, abdomine testaceo, supra segmentis tribus primis summa basi medio piceis, quarto quintoque nigro-piceis, margine summo apicali nigro, infra segmentis ultimis tribus fusciscentibus.

3, Thorace nigro piceo, pectore toto flavo-testaceo, abdomine infra flavo-testaceo, unicolore, supra piceo, limbo laterali integro testaceo.

Obs. Exempla genuina et var. 1 mascula, var. 2 et 3 feminea. Oxy. cinctus Grav. ad var. 3 pertinet. — A praecedente statura minore, pedibus cum coxis testaceis et elytrorum angulo apicali exteriore laevissimo facile distinguendus.

\* 7. *O. lateralis*: *Rufo-testaceus, nitidus, capitis thoracisque lateribus elytrorumque angulo apicali exteriore nigris.* — Long.  $3\frac{2}{3}$  lin.

Grav. Micr. 195. 1. Mon. 235. 3. — Latr. Hist. nat. Crust. et Ins. IX. 358. 3. — Oliv. Encycl. meth. VIII. 607. 4.

Habitat in Pennsylvania, Knoch.

Rufo-testaceus, minus nitidus. Antennae longiusculae, minus compressae dilatataeque, articulis 5—8 obconicis, sensim paulo crassioribus, penultimis vix transversis, concolores. Palpi testacei. Mandibulae maris ante apicem obtuse dentatae. Caput maris thorace paulo maius, feminae ei aequale, apice utrinque leviter longitudinaliter impressum, ante medium foveola minuta oblonga magis minusve distincta notatum, laeve, utrinque vitta laterali nigro-picea maculaque frontali media fusca. Thorax basi coleopteris duplo fere angustior, lateribus modice rotundatus, latitudine dimidio brevior, leviter convexus, lateribus pone medium oblique impressus, utrinque vitta laterali nigro-picea. Elytra duplo prope longiora, testacea, angulo apicali exteriore late nigro, disco seriebus duobus impressis punctorum irregulariter dispositorum punctisque nonnullis sparsis notata. Abdomen aut immaculatum aut dorso segmento quoque utrinque macula fusca signato. Pedes toti flavo-testacei.

#### Species mihi invisae.

8. *O. Stygicus*: *Niger, tarsis rufis.* — Long. 3 lin.

Say. Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. pag. 459.

Habitat in Indiana Americae septentrionalis.

Niger, politus. Antennae articulis 2—5 rufis. Labrum rufum. Palpi testacei. Thorax margine laterali inflexo depresso. Elytra nigra, immaculata, stria subsuturali aliaque abbreviata media, et prope hanc stria obsoleta punctisque nonnullis maioribus impressa, margine laterali impunctato. Pedes nigro-picei, tarsis pallide rufis.

Obs. Thorax paulo longior, quam in *O. cincto* Gr., et elytrorum puncta numerosiora. (Say).

#### Observatio.

Speciem succino inclusam Gravenhorst Mon. p. 235 hoc modo descripsit nomine O. Blumenbachii: „Caput quadratum, angulis obtusis, thorace maius. Antennis capitis longitudine, pilosellis, articulo primo longiore, 7 sequentibus dimidia illius longitudine, ultimis haud rite dignoscendis, sed, uti videtur, materie heterogena obductis, qua speciem clavae praebent (antennae alterius pars dimidia deficit). — Clypeus exacte ut in Oxyporis hodiernis formatus, transversus, margine anteriore paululum emarginato. — Instrumenta cibaria maximam partem destructa sunt, ita ut forma eorum haud exacte dignosci possit. Thorax transversus, exacte ut in nostris Oxyporis formatus, margine anteriore recto, posteriore rotundato, capitis latitudine, sed dimidia eius longitudine. Scutellum haud dignoscendum. Coleoptera paululum e situ naturali remota, thoracis magnitudine, elytris quadratis. Abdomen coleopteris triplo longius, e segmentis 8 constat. Pedes numero, forma, longitudine et proportionem partium omnino respondentibus Oxyporum nostrorum; numerus autem articulorum in tarsis haud perspicue videri potest. Color pallide flavus est. Color naturalis verisimile omnino evanuit.”



## Tribus quinta.

### **P a e d e r i n i.**

*Stigmatica prothoracica occulta, spatio pone coxas anticas membraneo.*  
*Coxae posticae conicae.*

---

Labrum corneum, limbo laterali saepe membranaceo,  
bilobum, muticum, lobis elongatis, divergentibus Achenio;  
lobis brevibus, rotundatis Cryptobio, Scimbalio, Lathrobio;  
brevissimum, medio leviter emarginatum, Dolicaoni, Paedero;  
apice bidentatum, inter dentes incisum Scopaeo, Lithocharidi, Sti-  
lico, Ophiti, Sunio;  
apice 4-dentatum Echiasteri.

Mandibulae longae, tenues, falcatae, acutae, medio dentatae.

Maxillae malis brevioribus, interiore interne, exteriori apice barba-  
tis, huius barba introrsum vergente.

Palpi maxillares magis minusve elongati, articulo primo minuto,  
brevi, secundo tertioque elongatis, hoc apicem versus magis minusve acu-  
minato, quarto minimo;

acuminato, subulato, distincto Cryptobio, Scimbalio, Achenio, La-  
throbio, Scopaeo, Lithocharidi, Ophiti;

obtusio, distincto, Dolicaoni, Paedero;

acuto, at minutissimo, aegerrime conspiciendo Stilico, Echiasteri,  
Sunio.

Mentum corneum, transversum, breve.

Ligula membranaea, biloba,

lobis approximatis, Cryptobio, Dolicaoni, Scimbalio, Achenio,  
Lathrobio, Ophiti;

lobis distantibus Lithocharidi, Stilico, Echiasteri, Sunio, Paedero;  
cornea, apice tridentata Scopaeo.

Paraglossae lineares, membranaeae, interne ciliatae.

Palpi labiales triarticulati, articulo tertio tenuiore, acuminato.

Antennae infra oculos sub capitis margine laterali insertae, 11-arti-  
culatae, rectae plerisque;

geniculatae, articulo primo elongato Cryptobio, Ophiti.

Caput exsertum, collo quasi tenui thoraci affixum vel distinctius pedunculatum Scopaeo, Ophiti, Stilico, Echiasteri, Sunio.

Thorax immarginatus, basi truncatus, apice attenuatus Scopaeo, Ophiti, Stilico, Echiasteri, Sunio; spatio pone coxas anticas absque integumento corneo.

Scutellum distinctum, triangulare.

Elytra pectoris longitudine.

Abdomen elongatum, marginatum, segmento septimo saepius minuto, bivalvi;

valvulis styliformibus Cryptobio, Dolicaoni, Scimbalio, Ophiti, Paedero;

valvulis unciformibus Achenio.

Pedes coxis anticis conicis, exsertis, intermediis approximatis, posticis conicis, tibiis muticis, tarsis 5-articulatis, articulo quarto simplice, plerisque; bilobo Paedero;

lobo membraneo instructo Sunio.

tarsis posterioribus articulis 4 primis longitudine sensim decrescentibus plerisque;

longitudine aequalibus Achenio, Lathrobio;

tarsis anticis articulis 4 primis fortiter dilatatis Scimbalio, Achenio, Lathrobio;

leviter dilatatis, Dolicaoni, Paedero, Lithocharidi;

simplicibus Cryptobio, Scopaeo, Ophiti, Stilico, Echiasteri, Sunio.

#### O b s e r v a t i o .

Continetur haec tribus genere Fabriciano Paedero, quod Gravenhorst secundum palporum maxillarium haud satis exactam observationem in genera Lathrobium et Paederum dissolvit. Latreille secundum labri figuram Lathrobium Staphylinorum genuinorum sectioni associavit, Paederum Stenis apposuit. Primo intuitu Lathrobia habitu quodammodo divergunt a Paederis, imprimis pedibus robustioribus, tibiis apice oblique truncatis, palpis maxillaribus minus elongatis, genera autem Scopaeus et Lithocharis omnino intermedia, ut subtribu etiam distinguere queant.

#### Cryptobium Mannerh.

Mannerh. Brachelytr. p. 38. — Steph. Britt. Ent. V. p. 271. — Boisd. et Lacord.

Faun. Ent. Paris. I. p. 427. — Er. Col. March. I. p. 500.

Lathrobium Fam. III. Grav. Mon. p. 129.

Labrum bilobum.

Palpi maxillares articulo ultimo acuto.

Antennae fractae.

Tarsi articulo quarto simplice, posteriores articulis 4 primis decrescentibus.

Corpus elongatum, lineare, alatum. Caput plerumque oblongum, basi constrictum, collo quasi brevi thoraci immissum, exsertum, porrectum, lateribus pone oculos rotundatis, tumidulis, ante oculos abrupte paulo angustius, oculis minutis, rotundatis, parum prominulis. Labrum breve, transversum, bilobum, lobis divergentibus, margine setosum. Mandibulae longae,

falcatae, acutae, medio fortiter acuteque bi- vel tridentatae. Maxillae (t. IV. f. 28. b.) malis coriaceis, interiore interne, exteriori apice barbatis. Palpi maxillares modice elongati, articulo primo parvo, secundo tertioque aequalibus, hoc apicem versus sensim leviter incrassato, quarto minimo, subulato. Labium (t. IV. f. 28. a.) mento transversum, ligula membranacea, biloba, lobis rotundatis, paraglossis acuminatis, membranaceis, ciliatis, ligulam paulo superantibus. Palpi labiales articulis duobus primis cylindricis, secundo primo paulo longiore, tertio minuto, tenui, acuminato. Antennae filiformes, tenues, fractae, articulo primo elongato, interdum apice subclavato. Thorax pleurumque coleopteris angustior, subcylindricus, basi apiceque truncatus. Scutellum triangulare. Elytra truncata. Abdomen lineare, stylis analibus exsertis, pilosis. Pedes modice elongati, graciles, antici haud incrassati, tibiis omnibus pubescentibus pilosisque, spinis apicalibus longioribus, tarsis omnibus simplicibus, articulo primo elongato, reliquis sensim decrescentibus, ultimo praecedentibus ambobus subaequali.

#### O b s e r v a t i o n e s.

1. Genus valde distinctum tarsis anticis simplicibus et antennis geniculatis. Caput et thorax pube subtili erecta e punctis oriunda vestita. Thorax nonnunquam subseriatim punctatus.

2. Species Americanae a Com. Dejean in recentiore collectionis suae Catalogo sub genere proprio Stilico enumerantur, at neque hoc nomen iis vacat, neque omnino ab Europaeo Cryptobio fracticorni genere distingui possunt.

3. Mares alii abdominis segmento inferiore tertio apice in processum longum elongato, alii segmento inferiore sexto apice inciso, alii ambabus notis a femina distinguuntur.

4. Victus speciei indigenae in silvis palustribus, sub musco et sub foliis putrescentibus.

\* 1. *C. badium*: *Rufo-castaneum, nitidum, pedibus testaceis, capite coleopterorum latitudine, oblongo, crebrius punctato, piceo, thorace utrinque subseriatim punctato.* — *Long. 5 lin.*

*Latrob. badium* Grav. *Micr.* 53. 3. *Mon.* 135. 16. — *Latr. Hist. nat. Crust. et Ins.* IX. 338. 3.

Habitat in America septentrionali, Knoch.

Castaneo-rufum, nitidum, capite piceo, pedibus testaceis. Antennae capite thoraceque longe breviores, articulo tertio secundo sesqui prope longiore, sequentibus sensim brevioribus, ultimo apice abrupte acuminato, dilute rufo-piceae. Caput thorace sesqui fere latius, oblongum, subdepressum, crebrius punctatum, antice inter antennas laevius et utrinque subimpressum. Oculi infra capitis medium siti. Thorax coleopteris dimidio angustior, latitudine plus sesqui longior, subcylindricus, crebrius vage, passim subseriatim punctatus, plaga media longitudinali laevi. Scutellum laeve. Elytra thorace paulo longiora, crebre profundeque punctata, subtiliter fusco-pubescentia. Abdomen subtilius punctulatum, longius fusco-pubescentius. Pedes testacei, tibiis basi piceis.

Abdominis segmentum inferius tertium maris subtus medio processu elongato, segmentum quartum longe superante, margine nigro-piloso instructum, reliqua integra.

Obs. A Gravenhorstio errore quodam inter species Brunsvicensis descriptum.



- \* 2. *C. Carolinum*: Nigrum, nitidum, thorace, elytris anoque rufo-piceis, pedibus testaceis, capite coleopterorum latitudine, crebre punctato, thorace utrinque subseriatim punctato. — Long.  $4\frac{1}{2}$  lin.

Habitat in Carolina meridionali, Dom. Zimmermann.

Statura et summa similitudo praecedentis et sequentis. Antennae capite thoraceque parum breviores, tenues, articulo tertio secundo plus sesqui longiore, sequentibus sensim brevioribus, ultimo apice abrupte subacuminato, rufae, articulo secundo sequentibusque pluribus piceis. Palpi rufi. Labrum obscurius rufo-piceum. Caput thorace sesqui latius, sat magnum, oblongum, nigrum, nitidum, parce griseo-pubescent, fortius crebre punctatum, infra oculos laeve vel punctis paucis sparsis minus profundis impressum, antice inter antennas utrinque fortius impressum. Oculi infra capitis medium siti. Thorax coleopteris dimidio angustior, latitudine plus sesqui longior, subcylindricus, rufo-piceus, nitidus, utrinque crebrius subseriatim punctatus, plaga media longitudinali angusta laevi. Scutellum obsolete bipunctatum, piceum, Elytra thorace vix longiora, sat fortiter crebre punctata, tenuiter fusco-pubescentia, rufo-picea, nitida. Abdomen parcius punctulatum griseoque pubescens, nigrum, nitidum, segmentis ultimis duobus rufis. Pectus nigrum, mesosterno piceo. Pedes testacei, femoribus pallidioribus.

Abdominis segmentum inferius tertium maris subtus apice in processum, segmento quarto saepius brevior, latior, subsemiorbiculatum, margine nigro-pilosum productum, reliqua integra.

Variat minus maturus corpore rufo-testaceo, elytris abdominisque segmentis ultimis duobus dilutioribus, pedibus pallidis.

Obs. Distinctum a *C. bicolor* praecipue punctura fortiore, capite maiore, elytris brevioribus, ut et segmento quarto maris processu minus elongato instructo.

- \* 3. *C. bicolor*: Nigrum, nitidum, thorace, elytris anoque rufis, pedibus testaceis, capite coleopteris angustiore, oblongo, crebre punctato, thorace utrinque crebre punctato. — Long.  $4\frac{1}{2}$  lin.

*Luthrobium bicolor* Grav. *Micr.* 179. 1. *Mon.* 135. 17. — *Say Transact. Amer. Philosoph. Soc. Nov. Ser. IV.* 454. 1. — *Nordm. Symb.* 145. 15. — *Kirby Fauna Boreali-Amer.* 87. 4.

Habitat in America septentrionali, e mus. Knoch.

Antennae tenuiusculae, capitis thoracisque prope longitudine, articulo tertio secundo duplo prope longiore, sequentibus leviter elongatis, sensim brevioribus, ultimo apice truncato et abrupte subacuminato, rufae, medio fuscescentes. Palpi rufi. Labrum rufum. Mandibulae rufo-piceae. Caput thorace vix sesqui latius, coleopteris angustior, oblongo-subovatum, nigrum, parum nitidum, crebre punctatum, antice laeve, inter antennas utrinque impressum. Oculi infra capitis medium siti. Thorax coleopteris plus dimidio angustior, latitudine duplo prope longior, subcylindricus, utrinque crebrius vage punctatus, plaga media longitudinali angusta laevi, rufus, modice nitidus. Scutellum rufum, obsolete bipunctatum. Elytra thorace paulo longiora, fortius dense punctata, modice nitida, rufa. Abdomen distinctius punctulatum, griseo-pubescent, nigrum, segmentis ultimis duobus rufis. Pectus nigrum, mesosterno rufo. Pedes testacei, femoribus pallidioribus.

Abdominis segmentum inferius tertium maris apice processu elongato, segmentum quartum superante, margine longe piloso, apice rotundato, piceo instructo, reliqua integra.

- \* 4. *C. melanocephalum*: *Testaceum, pedibus pallidis, capite coleopterorum latitudine, oblongo, fortius crebre punctato, nigro, thorace utrinque fortius crebriusque punctato.* — Long.  $4\frac{1}{2}$  lin.

Habitat in America septentrionali, Dom. Zimmermann.

Statura et summa affinitas praecedentium, corpore pallidius testaceo, modice nitido, capite solo piceo vel nigro, elytris dorso infuscatis. Antennae capitis thoracisque prope longitudine, tenues, articulo tertio secundo vix sesqui longiore, 4 — 8 modice elongatis, aequalibus, 8 — 10 sensim brevioribus, ultimo subovato, apice abrupte subacuminato, basi pallide testaceae, medio fuscae, apice rufo-testaceae. Os rufo-testaceum. Caput thorace sesqui latius, oblongum, fortius sat crebre punctatum, antice infra oculos laeve, punctorum serie transversali in fronte antica notatum, inter antennis utrinque subimpressum, tenuiter nigro-pubescent. Oculi infra capitis medium siti. Thorax coleopteris dimidio angustior, latitudine sesqui fere longior, subcylindricus, rufo-testaceus, nitidus, subtiliter nigro-pubescent, utrinque fortius crebriusque subinaequaliter punctatus, linea longitudinali media laevi postice subelevata. Scutellum laeve. Elytra thorace paulo longiora, depressa, dense punctata, tenuiter cinereo-pubescentia, pallidius testacea, dorso communi infuscato. Abdomen punctulatum, parce supra tenuiter, subtus longius cinereo-pubescent, pallidius testaceum. Pectus pallide testaceum. Pedes pallide flavi.

Abdominis segmentum inferius tertium maris apice processu longiore, segmentum quartum longe superante, lato, apice rotundato, margine longe nigro-piloso instructo, reliqua integra.

Obs. Praecedenti proximum, colore autem, capite maiore, elytris amplioribus, et processu segmenti quarti abdominis in mare latiore etc, distinctum videtur.

- \* 5. *C. prolixum*: *Nigrum, subnitidum, pedibus testaceis, coxis nigropiceis, capite thorace latiore, oblongo, creberrime punctato, thorace angusto, crebre punctato, coleopteris thorace sesqui longioribus.* — Long.  $4\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Praecedentibus paululum gracilius, nigrum, subnitidum. Antennae capitis thoracisque fere longitudine, tenues, articulo tertio secundo duplo longiore, 4 — 8 modice elongatis, aequalibus, 8 — 10 sensim brevioribus, ultimo apice truncato et abrupte subacuminato, rufo-piceae, apice rufo-testaceae. Os rufo-piceum. Caput thorace sesqui fere latius, coleopteris plerumque paulo angustius, oblongum, subdepressum, creberrime sat fortiter punctatum, antice infra oculos laeviusculum, opacum, inter antennis utrinque impressum. Oculi infra capitis medium siti. Thorax coleopteris duplo fere angustior, latitudine plus sesqui longior, subcylindricus, sat crebre fortiterque subinaequaliter punctatus, linea media longitudinali subelevata laevi. Scutellum obsolete bipunctatum, opacum. Elytra thorace sesqui longiora, confertim punctata, subtiliter nigro-pubescentia. Abdomen parcius

subtiliusque punctulatum, griseo-pubescens, nigro-piceum, nitidum; subtus basi rufescenti-piceum, apice summo piceum. Pedes testacei, coxis piceis, femoribus flavo-testaceis.

Abdominis segmentum inferius tertium maris apice processu longiore, segmentum quartum superante, lineari, apice rotundato, margine longe nigro-piloso instructum, reliqua integra.

6. *C. longipenne*: Nigrum, subnitidum, pedibus testaceis, coxis piceis, capite thorace latiore, confertissime subtilius punctato, thorace angusto, creberrime punctato, elytris thorace plus sesqui longioribus. — Long. 5 lin.

Habitat in Mexico, coll. Dom. Chevrolat.

Habitus et summa affinitas praecedentis, at sesqui fere maius. Nigrum, subnitidum. Antennae capite thoraceque breviores, tenues, articulo tertio secundo duplo fere longiore, 4 — 6 modice elongatis, aequalibus, reliquis sensim brevioribus, articulo ultimo apice truncato et abrupte subacuminato, rufo-testaceae. Palpi rufo-testacei. Mandibulae piceae. Labrum rufo-testaceum. Caput thorace sesqui vel interdum fere duplo latius, oblongum, depressum, densius subtiliter nigro-pubescens, confertissime subtiliter punctatum, opacum, apice laeviusculum, utrinque modice impressum. Oculi paulo infra capitis medium siti. Thorax coleopteris duplo angustior, latitudine sesqui longior, subcylindricus, lateribus ante medium levissime rotundatus, creberrime punctatus, linea media longitudinali subelevata antice abbreviata laevi, subtiliter nigro-pubescens, lateribus nigro-pilosellus, nitidus. Scutellum laeve, opacum. Elytra thorace plus sesqui longiora, depressa, confertissime punctata, subrugulosa, subnitida, subtiliter densius nigro-pubescentia. Abdomen crebre subtiliter punctulatum, subnitidum, densius fusco-pubescens, apice piceum. Pedes testacei, coxis piceis.

Mas latet.

Obs. Proximum praecedenti, pube autem densiore, capite densius subtiliusque punctato, opaco, thorace crebrius punctato, elytris longioribus distinctum.

- \* 7. *C. fasciatum*: Rufo-piceum, subnitidum, pedibus testaceis, elytrorum fascia abdomineque ante apicem nigro-piceis, capite thorace latiore, oblongo, subtiliter crebrius punctato, opaco, nigro, thorace utrinque crebrius punctato. — Long. 4 lin.

Habitat in Columbia, Dom Moritz.

Antennae capite thoraceque breviores, tenues, articulo tertio secundo sesqui longiore, 4 — 7 modice elongatis, subaequalibus, sensim brevioribus, ultimo apice truncato et abrupte subacuminato, rufae, articulis 3 — 7 basi fusciscentibus, reliquis rufo-testaceis. Palpi rufo-testacei. Labrum rufum. Mandibulae rufo-piceae. Caput thorace vix sesqui latius, oblongum, subdepressum, nigrum, opacum, subtiliter nigro-pubescens, subtiliter crebrius punctatum, antice infra oculos laeve, inter antennas utrinque obsolete impressum. Oculi infra capitis medium siti. Thorax coleopteris dimidio fere angustior, latitudine plus sesqui longior, subcylindricus, dilute rufo-piceus, nitidus, crebre vageque sat fortiter punctatus, linea lata longitudinali media laevi. Scutellum laeve, nitidum, rufum. Elytra thorace paulo longiora, depressa, dense punctata, subtiliter cinereo-pubescentia, rufo-testacea, sub-



nitida, fascia lata nigricante, seu nigricantia, basi piceo-rufa, apice testacea. Abdomen subtiliter punctulatum, nitidum, rufo-piceum, ante apicem nigropiceum. Pedes testacei, coxis rufis, femoribus flavo-testaceis.

Abdominis segmentum inferius tertium maris apice processu longiore, segmentum quartum longe superante, elongato-triangulari, apice subrotundato, margine longe piloso instructum, reliqua integra.

- \* 8. *C. albipes*: *Nigro-piceum, subnitidum, pedibus pallide testaceis, capite coleopterorum latitudine, subquadrato, creberrime punctato, thorace utrinque crebre punctato. — Long. 3 lin.*

Habitat in Puerto Rico, Dom. Moritz.

Nigro-piceum, subnitidum. Antennae capite thoraceque paulo breviores, tenuiores, articulo tertio secundo sesqui longiore, 4—8 elongato-obconicis, aequalibus, ultimis sensim brevioribus, ultimo subovato, apice abrupte acuminato, testaceae, articulis 3—8 medio fuscis. Palpi testacei. Os rufo-piceum. Caput thorace sesqui latius, coleopterorum latitudine, oblongo-subquadratum, leviter convexum, tenuiter griseo-pubescentia, crebre punctatum, fronte antius infra oculos laevi, inter antennis utrinque distincte impressa. Oculi paulo infra medium capitis siti, modice prominuli. Thorax coleopteris dimidio angustior, latitudine sesqui longior, subcylindricus, fortius sat crebre punctatus, punctis versus medium subseriatis, linea longitudinali media laevi. Scutellum laeve, nitidum. Elytra thorace sesqui longiora, depressa, creberrime punctata, subtiliter cinereo-pubescentia, margine summo apicali ferrugineo. Abdomen densius sat distincte punctulatum, cinereo-pubescentia, segmentis ultimis duobus margine pallidis. Pedes pallide testacei, coxis piceo-testaceis.

Abdominis segmentum inferius tertium apice processu lato subsemicirculari, segmentum quartum vix superante, margine pilis longis nigris ciliato instructum, reliqua integra.

- \* 9. *C. fulvipes*: *Brunneo-piceum, nitidum, pedibus testaceis, capite coleopterorum latitudine, subquadrato, parcius profunde punctato, thorace utrinque crebre punctato, elytris thorace paulo longioribus. — Long. 3 lin.*

Habitat in ins. Puerto Rico, Dom. Moritz.

Statura omnino praecedentis. Castaneo-piceum, nitidum. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus, minus elongatis gracilibusque, ultimo truncato et apice abrupte subtiliter acuminato, rufo-piceae, apice testaceae. Palpi testacei. Mandibulae piceae. Labrum testaceum. Caput thorace sesqui latius, subquadratum, subdepressum, fortiter minus crebre punctatum, interstitiis parce sparsimque subtilius punctatis, fronte antice laeviore, inter antennis utrinque obsolete impressa. Oculi ad capitis medium siti, minus prominuli. Thorax coleopteris vix dimidio angustior, latitudine sesqui longior, subcylindricus, crebrius profunde vage punctatus, plaga latiore media longitudinali laevi. Scutellum laeve. Elytra thorace paulo longiora, creberrime punctata, minus nitida, densius subtiliter griseo-pubescentia. Abdomen subtiliter obsoleteque punctulatum, longius pubescens, apice segmen-

torumque singulorum marginibus piceo-testaceis. Pedes piceo-testacei, coxis concoloribus.

Abdominis segmentum inferius tertium maris apice processu lato, parallelo, apice truncato, ad segmenti quinti apicem prope elongato, distinctius punctato, longe nigro-piloso instructum, sextum apice acute emarginatum.

- \* 10. *C. antiquum*: Nigro-subaeneum, nitidum, pedibus piceis, femoribus supra testaceis, capite coleopterorum latitudine, depresso, subquadrato, parce obsolete punctato, opaco, thorace utrinque parce subseriatim punctato. — Long.  $4\frac{1}{2}$  lin.

Habitat in Brasilia, Dom. Virmond et Westermann.

Statura fere Cr. melanocephali, nigro-subaeneum, nitidum. Antennae capite sesqui longiores, tenues, articulo tertio secundo duplo prope longiore, 4 — 10 sensim brevioribus, ultimo apice abrupte acuminato, rufo-piceae, articulo primo crassiore, apice rufescente. Palpi et labrum rufa. Mandibulae piceae. Caput thorace vix sesqui latius, oblongum, lateribus pone oculos tumidulum, subdepressum, nigro-aeneum, opacum, obsolete parce subtiliter punctatum, anterieus sensim laeve, inter antennis leviter impressum. Oculi vix infra capitis dimidium siti. Thorax coleopteris paulo angustior, latitudine sesqui longior, subcylindricus, dorso fortius biserialim punctatus, lateribus punctis nonnullis subtilibus sparsus. Scutellum laeve. Elytra thoracis longitudine, depressa, fortius dense punctata, subtilissime griseo-pubescentia, subnitida. Abdomen parce subtiliterque punctulatum, subtiliter nigro-pubescent, apice nigro-pilosum, segmentis ultimis duobus margine apicali rufo-piceo. Pedes picei, femoribus superne piceo-testaceis.

Mas latet.

- \* 11. *C. femorale*: Nigro-piceum, nitidum, pedibus piceis, femoribus testaceis, capite coleopterorum latitudine, depresso, subquadrato, crebrius punctato, thorace utrinque crebre subseriatim punctato. — Long.  $4\frac{1}{2}$  lin.

Habitat in Brasilia, Dom. Virmond.

Subdepressum, nigro-piceum, nitidum. Antennae capitis thoracisque prope longitudine, tenues, articulo tertio secundo duplo longiore, 4 — 8 modice elongatis, aequalibus, sequentibus sensim brevioribus, ultimo apice abrupte acuminato, rufo-piceae. Os cum palpis rufo-piceum. Caput thorace duplo fere latius, oblongo-subquadratum, subdepressum, sat crebre profundius punctatum, anterieus parce obsoleteque punctatum, inter antennis utrinque subimpressum. Oculi infra capitis medium siti. Thorax coleopteris plus dimidio angustior, latitudine sesqui longior, subcylindricus, dorso depressiusculus, utrinque fortius sat crebre punctatus, punctis partim subseriatis, plaga media longitudinali laevi. Scutellum laeve, piceum. Elytra thorace paulo longiora, depressa, dense punctata, punctis maioribus nonnullis distantibus seriatis impressa. Abdomen parce obsoleteque subtilissime punctatum, margine laterali dilute, segmentorum marginibus apicalibus obscure rufo-piceis. Pectus piceum. Pedes rufo-picei, femoribus testaceis, coxis geniculisque piceis.

Abdominis segmentum superius sextum maris apice leviter emarginatum, inferius tertium apice processu magno, ad segmenti quinti apicem

usque prope elongato, parallelo, apice rotundato, distinctius punctato, margine longe nigro-piloso instructum, reliqua integra.

- \* 12. *C. dispar*: *Piceum, nitidum, pedibus rufis, capite thorace latiore, oblongo, parcius punctato, thorace dorso biserialim, lateribus disperse punctato.* — Long. 3 lin.

Habitat in Columbia, Dom. Moritz.

*C. albipedi* fere aequale, iam nigro-piceum, iam piceo-rufum, nitidum. Antennae capite thoraceque breviores, articulo secundo tertio vix brevior, sequentibus sensim paulo brevioribus, ultimo apice truncato et abrupte subacuminato, rufo-testaceae. Palpi rufo-testacei. Caput thorace paulo latius, oblongum, subdepressum, pone oculos minus crebre punctatum, ante oculos punctis parvis sparsum, inter antennis utrinque obsolete impressum. Oculi ad capitis medium siti. Thorax coleopteris dimidio angustior, latitudine sesqui longior, subcylindricus, dorso biserialim, lateribus minus crebre disperse punctatus. Scutellum laeve. Elytra thorace tertia parte longiora, dense subtiliter punctata, subtiliter cinereo-pubescentia, minus nitida. Abdomen parcius subtiliter obsoleteque punctulatum, piceum, segmentis singulis margine apicali late rufo-piceis. Pedes iam rufo-picei, iam rufo-testacei, coxis omnibus concoloribus.

Abdominis segmentum inferius sextum maris apice profundius triangulariter incisum, quintum apice late leviter emarginatum, tertium apice processu triangulari, acuto, fere conico, ad segmenti quarti medium fere porrecto instructum.

- \* 13. *C. pallipes*: *Nigro-piceum, nitidum, pedibus flavis, capite thorace latiore, oblongo, crebre punctato, thorace dorso utrinque crebre punctato.* — Long.  $4\frac{1}{2}$  lin.

Nordm. Symb. 149. 1.

*Lathrobium pallipes* Grav. Micr. 179. 2. Mon. 130. 4.

*Lathrobium Gravenhorstii* Kirby Faun. Boreali-Amer. 87. 3. t. 2. f. 2.

Habitat in Pennsylvania, e mus. Knoch.

Statura Cr. melanocephali, nigro-piceum, nitidulum. Antennae capite thoraceque paulo breviores, tenuiores, articulo tertio secundo paulo longiore, 4 — 7 modice elongatis, aequalibus, sequentibus sensim brevioribus, ultimo ovato, apice abrupte acuminato, rufo-testaceae. Palpi et labrum rufo-testacea. Caput thorace paulo latius, oblongum, crebre punctatum, anterieus laeve, inter antennis utrinque impressum, tenuiter subtiliterque griseo-pubescent. Oculi infra capitis medium siti. Thorax coleopteris paulo angustior, latitudine sesqui fere longior, subcylindricus, utrinque sat crebre vage punctatus, spatio medio longitudinali laevissimo. Scutellum punctis paucis obsoletis notatum. Elytra thorace paulo longiora, dense punctata, griseo-pubescentia. Abdomen crebrius subtiliter punctulatum, densius subtiliter griseo-pubescent, nigro-piceum, segmentis singulis margine apicali rufo-piceo. Pedes coxis omnibus rufo-testaceis, femoribus flavis, tibiis tarsisque dilute rufo-testaceis.

Abdominis segmentum inferius sextum maris apice triangulariter excisum.



- \* 14. *C. latebricola*: *Nigrum, nitidum, pedibus testaceis, capite thoracis latitudine, oblongo, crebrius fortiusque punctato, thorace dorso biseriatim, utrinque crebrius punctato.* — Long. 4 lin.

Nordm. Symb. 150. 2.

*Lathrob. cinctum* Say Transact. Amer. Philoph. Soc. Philadelph. Nov. Ser. IV. 454. 3. (forte)

Habitat in America septentrionali.

Praecedenti affine, at duplo minus, nigro-piceum, nitidum. Antennae capite thoraceque paulo breviores, tenues, apice subincrassatae, articulo tertio secundo sesqui longiore, 4 — 7 modice elongatis, subaequalibus, sequentibus sensim paulo brevioribus et crassioribus, ultimo ovato, apice abrupte acuminato, rufo-testaceae. Palpi et labrum testacea. Mandibulae rufo-testaceae. Caput thoracis latitudine, oblongum, pone oculos crebre sat fortiter, ante oculos parce vageque punctatum, antice inter antennis utrinque sat distincte impressum, parce nigro-pubescent. Oculi paulo infra capitis medium siti. Thorax coleopteris paulo angustior, latitudine sesqui longior, subcylindricus, utrinque crebrius vage punctatus, spatio dorsali longitudinali laevissimo, utrinque punctorum numerosorum serie minus regulari terminato. Scutellum laeve. Elytra thorace paulo longiora, dense fortius punctata, subtilius cinereo-pubescentia. Abdomen crebre distinctius punctulatum, supra nigro-, subtus cinereo-pubescent, infra segmentis singulis apice tenuissime piceo-marginatis. Pedes rufo-testacei, femoribus flavis.

Abdominis segmentum inferius sextum maris apice triangulariter incisum.

- \* 15. *C. geniculatum*: *Nigrum, pedibus testaceis, coxis geniculisque piceis, capite thorace latiore, subquadrato, subtilius creberrime punctato, thorace utrinque creberrime punctato.* — Long. 4 lin.

Habitat in Cayenna, Dom. Buquet.

Nigrum, parum nitidum, subtiliter griseo-pubescent. Antennae capite thoraceque prope longiores, tenues, articulo tertio secundo sesqui longiore, 4 — 8 modice elongatis, subaequalibus, sequentibus decrescentibus ultimo apice truncato et abrupte acuminato, rufo-piceae, articulis singulis apice fuscescentibus, ultimis immaculatis. Os cum palpis piceum, maxillaribus articulo terminali dilute testaceo. Caput thorace sesqui latius, subquadratum, angulis posterioribus rotundatis, ante oculos angustatum, depressum, supra creberrime punctatum, subopacum, nigro-pubescent, parce nigro-pilosum, antice inter antennis parce punctatum, utrinque impressum. Oculi infra capitis medium siti. Thorax coleopteris dimidio prope angustior, latitudine sesqui longior, parallelus, subdepressus, creberrime minus subtiliter subinaequaliter punctatus, linea longitudinali media latiore maculaque utrinque minuta basali laevissimis. Scutellum parce obsoleteque punctatum. Elytra thorace sesqui fere longiora, depressa, creberrime punctata, subtiliter cinereo-pubescentia. Abdomen parcius subtilissime punctulatum, pube subtili densiore griseo-sericans, segmentis ultimis duobus margine apicali rufo-piceo. Pedes obscure testacei, coxis nigro-piceis, femoribus flavis, geniculis fuscescentibus.

Mas latet.

16. *C. similipenne*: *Nigrum, nitidulum, pedibus testaceis, coxis nigro-piceis, capite thorace latiore, oblongo, dense punctato, elytris profunde punctatis, margine apicali piceo.* — Long. 4 lin.

*Lathrob. similipenne* Say *Transact. Amer. Philoph. Soc. Philadelph. Nov. Ser. IV.* 453. 1.

Habitat in Mexico, coll. Dom. Chevrolat.

Statura *C. sulphuripedis*, nigrum, nitidulum. Antennae capite sesqui longiores, tenues, articulo tertio secundo paulo longiore, 4 — 8 elongato-obconicis, subaequalibus, 9 et 10 sensim brevioribus, obconicis, ultimo ovato, apice abrupte obtuse acuminato, articulis duobus primis rufo-piceis, 3 — 7 nigro-piceis, reliquis 4 rufo-testaceis. Palpi piceo-testacei. Mandibulae rufae. Labrum testaceum. Caput thorace sesqui prope latius, oblongo-ovatum, supra parum convexum, sat crebre profundius punctatum, antice laevius et parce punctatum, subtiliter nigro-pubescent, minus nitidum. Oculi ante capitis medium siti. Thorax coleopteris dimidio angustior, latitudine sesqui longior, parallelus, subdepressus, sat crebre profundeque punctatus, linea media latiore subelevata laevis, parce nigro pubescens. Elytra thorace tertia parte longiora, subdepressa, dense fortius profunde punctata, nigro-pubescentia, nitidula, margine apicali tenui piceo-rufescente. Abdomen parcius subtiliusque punctatum, pube longiore depressa fusco-nigra vestitum, segmentis 4 primis apice anguste piceo-marginatis, quinto sextoque apice rufo-piceis. Pedes coxis piceis, femoribus pallide flavis, tibiis tarsisque rufo-piceis.

Mas latet.

Obs. A *C. sulphuripede* sat distinctum antennis obscuris, capite maiore, dense punctato, thorace paulo longiore, subdepresso, lateribus nullo modo rotundato, elytris paulo longioribus, abdomine distinctius punctulato, coxis piceis, etc., a praecedente magis discedit capite oblongo-ovato, elytris fortius profundiusque minus confertim punctatis, abdomine minus subtiliter obsoleteque punctulato, antennis minus elongatis, etc.

17. *C. assimile*: *Nigrum, subnitidum, pedibus piceis, femoribus flavis, elytris margine summo apicali testaceo, capite thorace paulo latiore, crebre punctato, opaco, thorace utrinque creberrime punctato.* — Long.  $3\frac{1}{2}$  lin.

Habitat in Mexico, coll. Dom. Chevrolat.

Corpus gracilius, *C. sulphuripedi* aequale, nigrum, subnitidum. Antennae capite sesqui longiores, tenuiores, articulo tertio secundo vix sesqui longiore, 4 — 8 elongato-obconicis, reliquis tribus brevioribus, ultimo subovato, apice abrupte acuminato, nigro-piceae, articulis ultimis tribus rufo-testaceis. Mandibulae elongatae, piceae. Caput thorace paulo angustius oblongum, supra leviter convexum, opacum, dense punctatum, densius subtiliter griseo-pubescent, fronte anterieus laevi, inter antennis impressa. Oculi ante capitis medium siti, sat prominuli. Thorax coleopteris paulo angustior, latitudine sesqui longior, subcylindricus, creberrime fortius profunde punctatus, margine posteriore lineaque media subelevata laevissimis, parce nigro-pubescent. Scutellum laeve. Elytra thorace tertia parte longiora, profunde hinc inde subseriatim creberrime punctata, subrugulosa, subtiliter dorso nigro-, lateribus griseo-pubescentia, margine tenui apicali rufo-testa-

ceo. Abdomen densius subtiliusque punctatum, longius griseo-pubescent, segmentis ultimis duobus margine apicali late rufo-testaceo, quatuor primis infra apice obscure piceo-marginatis. Pedes coxis piceis, trochanteribus rufo-testaceis, femoribus pallide flavis, tibiis nigro-piceis, apice cum tarsis saturate rufo-testaceis.

Mas latet.

- \* 18. *C. ochropus*: *Nigro-piceum, nitidum, antennis pedibusque cum coxis anticis testaceis, elytris margine apicali rufo-piceo, capite thorace latiore, parcius punctato, thorace utrinque crebre fortiter punctato.* — Long. 4 lin.

Habitat in Brasilia, Dom. Prof. Germar.

Statura et summa affinitas *C. sulphuripedis*, at paulo robustius, nigro-piceum, nitidum. Antennae breviusculae, capite paulo longiores, tenuiores, articulo tertio secundo sesqui longiore, 4 — 7 modice elongatis, aequalibus, sequentibus sensim brevioribus, ultimo cylindrico, apice truncato et abrupte acuminato, rufo-testaceae. Os testaceo-rufum, mandibulis apice piceis. Caput thorace sesqui latius, oblongum, supra subdepressum, pone oculos parcius subtiliusque punctatum, ante oculos laeve, inter antennis utrinque leviter impressum, tenuiter nigro-pubescent, lateribus parce nigro-pilosum. Oculi versus capitis medium siti. Thorax coleopteris dimidio angustior, latitudine sesqui longior, lateribus ante medium levissime rotundatus, parum convexus, fortiter profunde sat crebre punctatus, linea media longitudinali posterius subelevata laevi, parce nigro-pubescent. Scutellum laeve, opacum. Elytra thorace tertia fere parte longiora, subdepressa, subtiliter nigro-pubescentia, confertissime profunde punctata, minus nitida, brunneo-picea, humeris dilutioribus, margine apicali obscure rufescente. Abdomen parcius subtiliusque punctatum, tenuiter nigro-pubescent, nigrum, nitidum, segmento quinto sextoque apice testaceo-marginatis. Pedes testacei, coxis anticis concoloribus, posterioribus piceis, femoribus dilutioribus, anticis apicem versus picescentibus.

Mas latet.

- \* 19. *C. sulphuripes*: *Nigrum, nitidum, pedibus flavis, elytris piceis, margine apicali testaceo, capite thorace latiore, parcius punctato, thorace utrinque creberrime fortiter punctato.* — Long. 4 lin.

Habitat in Columbia, Dom. Moritz.

Praecedentibus paulo gracilius, supra nigrum, nitidum. Antennae capite thoraceque paulo breviores, tenuiores, articulo tertio secundo sesqui longiore, 4 — 8 modice elongatis, subaequalibus, sensim brevioribus, ultimo apice truncato, obscure rufo-testaceae. Os piceum. Caput thorace sesqui prope latius, oblongum, supra leviter convexum, minus crebre punctatum, medio et antrorsum laeve, punctis paucis sparsum, inter antennis utrinque profundius impressum, antice saepe dilutius piceum, nigro-pubescent pilosumque. Oculi paulo infra capitis medium siti. Thorax coleopteris dimidio angustior, latitudine sesqui longior, subcylindricus, lateribus ante medium leviter rotundatis, creberrime sat fortiter punctatus, linea media longitudinali subelevata laevissima, parce nigro-pubescent. Scutellum laeve. Elytra



thorace tertia parte longiora, subdepressa, dense sat fortiter punctata, subtiliter cinereo-pubescentia, nigro-picea, humeris dilutioribus, margine apicali testaceo. Abdomen crebrius subtiliusque punctulatum, nigrum, nitidum, segmentis ultimis duobus apice rufo-piceis, subtus interdum medio longitudinaliter dilute rufo-piceum. Caput et thorax subtus cum pectore magis minusve dilute picea. Pedes pallide flavi, coxis rufo-testaceis.

Mas latet.

20. *C. brevicorne*: Subcylindricum, rufo-testaceum, nitidum, elytris pedibusque testaceis, capite thorace latiore, oblongo, crebre punctato, thorace crassiore, lateribus crebre punctato, elytris thorace paulo longioribus, subseriatim punctatis. — Long. 3 lin.

Habitat Novae Valentiae in Columbia, Dom. Moritz.

Statura fere praecedentis, nitidum, subtiliter nigro-pubescent, elytris pedibusque dilutioribus. Antennae capite parum longiores, apicem versus subincrassatae, articulo secundo tertioque aequalibus, elongato-obconicis, 4 — 10 crassitie haud longioribus, fere moniliformibus, ultimo breviter ovato, apice abrupte acuminato. Caput thorace paulo latius, oblongum, supra leviter convexum, crebre profundius punctatum, fronte antice laeviore, inter antennis utrinque fortius impressa. Oculi ad capitis medium siti, parum prominuli. Thorax coleopteris parum angustior, latitudine vix longior, basin versus subangustatus, modice convexus, lateribus crebre punctatus, punctis passim subseriatis, margine posteriore et plaga longitudinali media laevissimis. Scutellum laeve. Elytra thorace paulo longiora, sat crebre fortiterque subseriatim punctata. Abdomen parce subtilissime punctulatum. Pedes breviusculi, tarsis posterioribus brevibus, articulo primo sequentibus perparum longiore.

Mas latet.

Obs. Specimen singulum, evidenter immaturum, capite thoraceque rufo-testaceis, nitidis, elytris abdomineque luteo-testaceis, pedibus flavis.

21. *C. curticorne*: Subcylindricum, nigrum, nitidum, antennis pedibusque rufo-testaceis, capite thorace latiore, oblongo, crebre punctato, thorace utrinque dense punctato, elytris thoracis longitudine, subseriatim punctatis. — Long.  $3\frac{1}{2}$  lin.

Habitat in Brasilia, coll. Dom. Chevrolat.

Statura et summa affinitas praecedentis. Niger, nitidulus, densius nigro-pubescent. Antennae capite vix longiores, apicem versus sensim subincrassatae, articulo tertio secundo paulo longiore, 4 — 10 subcylindricis, crassitie paululum longioribus, sensim paulo brevioribus, ultimo subovato, apice abrupte acuminato, totae rufo-testaceae. Palpi rufo-testacei. Mandibulae piceae. Caput maiusculum, thorace latius, supra leviter convexum, sat crebre fortius profunde punctatum, antice laeviusculum, inter antennis utrinque impressum. Oculi paulo ante capitis medium siti, vix prominuli. Thorax coleopteris vix angustior, latitudine apicis paulo longior, basin versus subangustatus, modice convexus, sat crebre minus fortiter punctatus, linea media longitudinali laevi margineque basali laevissimis. Scutellum laeve. Elytra thoracis longitudine, creberrime hinc inde subseriatim pro-

funde punctata, subrugulosa, basi picea. Abdomen crebrius subtiliter punctulatum, segmento quinto sextoque margine apicali rufo-testaceis. Caput et thorax subtus cum pectore rufescenti-picea. Pedes cum coxis saturate rufo-testacei, dense pubescentes, breviusculi, tarsis posterioribus brevibus, articulo primo sequentibus vix longiore.

Mas latet.

Obs. A praecedente distinctum videtur capite maiore, thorace longiore, crebrius et minus fortiter punctato, elytris thorace haud longioribus.

- \* 22. *C. curtum*: Breve, subdepressum, piceum, subnitidum, pedibus testaceis, capite coleopterorum latitudine, subquadrato, dense punctato, thorace subquadrato, crebre punctato. — Long. 3 lin.

Habitat in Brasilia, Dom. Virmond.

Praecedentibus parum angustius, at multo brevius, subdepressum, piceum, subnitidum. Antennae capite cum thorace paulo breviores, tenues, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus, ultimo ovato, apice abrupte acuminato, rufae. Labrum et palpi rufa. Mandibulae valde elongatae, rufo-piceae. Caput thorace plus dimidio latius, subquadratum, lateribus pone oculos rotundato-ampliatum, planum, punctis sat magnis at parum profundis, antrorsum evanescentibus, subrugulosum, antice subtilissime conferissimeque strigosum, tenuiter subtiliterque nigro-pubescent. Oculi versus capitis medium siti. Thorax coleopteris dimidio angustior, latitudine haud longior, lateribus subrotundatus, basin versus subangustatus, subdepressus, crebrius punctatus, linea longitudinali media laevi, basi subelevata, tenuiter subtiliterque nigro-pubescent. Scutellum punctulatum. Elytra thorace sesqui longiora, dense punctato-subrugosa, subtiliter cinereo pubescentia, margine laterali subcostato. Abdomen parcius punctulatum, nitidum, segmentis singulis margine apicali rufo-piceo. Caput et thorax subtus cum pectore rufo-picea. Pedes cum coxis testaceo-rufi. Alae albo-hyalinae.

Abdominis segmentum inferius quintum maris apice medio subsinuatum, sextum acute emarginatum.

- \* 23. *C. obscurum*: Nigrum, nitidum, pedibus rufo-testaceis, capite thorace paulo latiore, oblongo, subtiliter punctulato, opaco, thorace utrinque creberrime punctato. — Long. 4 lin.

Habitat in Cayenna, Dom. Buquet.

Nigrum. Antennae capitis thoracisque fere longitudine, tenues, articulo primo longiore, apice leviter incrassato, tertio secundo sesqui longiore, 4 — 8 modice elongatis, subaequalibus, reliquis sensim brevioribus, ultimo breviter ovato, apice abrupte obtuse acuminato, rufae. Palpi et labrum rufa. Mandibulae piceae. Caput thorace paulo latius, oblongum, basi fortiter rotundatum, angulis posterioribus obtusis, prominentibus, lateribus pone oculos rectis, ante oculos leviter constrictum, antrorsum superne leviter angustatum, inferne leviter dilatatum, margine laterali inferiore inde prominente, supra subdepressum, subtiliter confertim punctatum, opacum, subtiliter griseo-fusco-pubescent, antice inter antennas utrinque foveola minuta impressum. Oculi ad capitis medium siti. Thorax coleopteris tertia parte angustior,

latitudine sesqui longior, subcylindricus, modice nitidus, creberrime punctatus, linea angusta media longitudinali laevi subelevata. Scutellum opacum, laeve. Elytra thorace paulo longiora, dense punctata, densius subtiliter cinereo-pubescentia, opaca. Abdomen parum nitidum, subtiliter parcius punctulatum, griseo-pubescent, segmentis singulis margine tenui rufo-piceo terminatis. Pedes cum coxis testaceo-rufi.

Mas latet.

24. *C. scitum*: *Nigrum, subnitidum, pedibus antennisque testaceis, his articulo primo apice piceo, capite thoracis latitudine, oblongo, creberrime punctato, thorace elongato, utrinque crebre fortiterque punctato. — Long. 3 lin.*

Habitat valle Araguensi Columbiae, Dom. Moritz.

Statura omnino praecedentis, nigrum, subnitidum. Antennae capite thoraceque paulo breviores, tenuiores, articulis 2 — 7 modice elongatis, subaequalibus, sequentibus sensim paulo brevioribus, ultimo apice truncatulo, testaceae, articulo primo apice piceo, sequentibus nonnullis fusciscentibus. Palpi testacei. Os rufo-piceum. Caput thoracis latitudine, oblongum, creberrime punctatum, parum nitidum, subtilissime cinereo-pubescent. Oculi ad capitis medium siti, sat prominuli. Thorax coleopteris dimidio angustior, latitudine summa sesqui longior, lateribus ante medium cum angulis anterioribus rotundatus, basin versus subangustatus, modice convexus, lateribus crebre fortiterque punctatus, spatio medio longitudinali laevissimo, subelevato, utrinque stria leviter impressa terminato. Scutellum punctato-inaequale. Elytra thorace paulo longiora, confertissime punctata, subrugulosa, parce griseo-pubescentia. Abdomen confertim subtiliter punctulatum, segmentis singulis apice rufo-marginatis. Pedes flavi, coxis anterioribus rufo-testaceis.

Abdominis segmentum inferius maris sextum apice triangulariter excisum, quintum apice leviter emarginatum.

- \* 25. *C. pulchellum*: *Rufo-piceum, pedibus testaceis, capite oblongo, creberrime punctato thoraceque utrinque crebre punctato rufis, elytris nigro-piceis, margine apicali testaceo, abdomine supra nigro. — Long. 2½ lin.*

Habitat in Columbia, Dom. Moritz.

Cr. fracticorni vix maius. Antennae capitis thoracisque fere longitudine, articulo secundo tertio aequali, sequentibus sensim brevioribus, ultimo apice truncato et apice abrupte subtiliter acuminato, testaceae, articulo primo apice fusco. Palpi testacei. Caput thorace vix latius, oblongum, leviter convexum, rufum, opacum, subtiliter griseo-pubescent, creberrime punctatum, inter antennis utrinque leviter impressum. Oculi versus capitis medium siti, modice prominuli. Thorax coleopteris dimidio angustior, latitudine sesqui fere longior, lateribus ante medium leviter rotundatus, modice convexus, rufo-piceus, nitidus, creberrime sat fortiter punctatus, linea lata longitudinali media laevissima postice subelevata. Scutellum rufo-piceum, obsolete punctulatum. Elytra thorace paulo longiora, sat profunde confertissime punctata, subrugulosa, subtiliter cinereo-pubescentia, nigro-picea, margine



apicali testaceo. Abdomen parcius subtilissime punctulatum, cinereo-pubes-  
cens, nigrum, segmentis singulis margine apicali obscure piceis, ultimis  
duobus margine apicali latiore rufo-testaceo, subtus disco anteriore pecto-  
reque dilute rufo-piceis. Pedes pallide flavi, coxis rufo-testaceis.

- \* 26. *C. agile*: Nigrum, nitidum, pedibus rufo-testaceis, capite thorace la-  
tiore, oblongo, parcius punctato, thorace elongato, utrinque crebre pun-  
ctato. — Long. 3 lin.

Habitat in Brasilia, Dom. v. Olfers.

Cr. fracticorni sesqui maius, nigrum, nitidum. Antennae capite thorace-  
que longe breviores. Caput thorace paulo latius, oblongum, leviter conve-  
xum, parcius sat profunde punctatum, antice laevius, inter antennas utrinque  
leviter impressum, parce nigro-pubescentia. Oculi versus capitis medium  
siti, sat prominuli. Thorax coleopteris dimidio angustior, latitudine duplo  
fere longior, subcylindricus, lateribus crebre punctatus, spatio dorsali lon-  
gitudinali medio laevissimo, parce nigro-pubescentia. Scutellum punctulatum.  
Elytra thorace paulo longiora, densius minus fortiter punctata, subtiliter  
cinereo-pubescentia. Abdomen confertim subtilissime punctulatum, densius  
griseo-pubescentia, opacum. Pedes rufo-testacei.

Abdominis segmentum inferius sextum maris fissum, quintum apice  
profunde incisum.

Variat corpore rufo-piceo, pedibus pallide testaceis.

- \* 27. *C. fracticorne*: Nigrum, nitidum, pedibus testaceis, capite thorace  
angustiore, oblongo, parce punctato, thorace dorso biseriatim, utrinque  
crebrius punctato. — Long.  $2\frac{1}{2}$  lin.

Mannerh. Brachelytr. 38. 13. — Steph. Illustr. Britt. Ent. V. t. 27. f. 1. — Boisd.  
et Lacord. Faun. Ent. Paris. I. 427. 1. — Nordm. Symb. 151. 3. — Runde  
Brachelytr. Hal. 13. 1. — Er. Col. March. I. 500. 1.

*Paederus fracticornis* Payk. Faun. Suec. III. 430. 5.

*Lathrobium fracticorne* Grav. Micr. 54. 5. Mon. 130. 3. — Latr. Hist. nat.  
Crust. et Ins. IX. 339. 5. — Gyll. Ins. Suec. II. 369. 5. — Sahlb. Ins. Fenn. I.  
341. 7. — Germ. Faun. Ins. Europ. XII. 1.

*Paederus glaberrimus* Payk. Mon. Car. App. 136. 17 — 18.

Habitat in Europa.

Nigrum, nitidum. Antennae capite thoraceque breviores, articulo se-  
cundo tertioque subaequalibus, sequentibus sensim paululum brevioribus  
et crassioribus, oblongo-subobconicis, ultimo apice truncatulo, piceo-testa-  
ceae, articulo primo apice fusco. Palpi maxillares testacei, articulo tertio  
fuscescente. Labrum piceum. Mandibulae testaceae. Caput thorace paulo  
angustius, oblongum, disperse punctatum, fronte media laeviore, parce nigro-  
pubescentia. Oculi versus capitis medium siti, parum prominuli. Thorax  
coleopteris tertia parte angustior, latitudine sesqui longior, subcylindricus,  
lateribus crebrius punctatus, spatio longitudinali medio laevissimo, utrinque  
punctorum numerosorum serie sat regulari terminato, lateribus parce sub-  
tiliterque nigro-pubescentia. Scutellum punctulatum. Elytra thorace paulu-  
lum longiora, densius punctata, fusco-pubescentia. Abdomen confertim sub-

tilissime punctulatum, densius subtiliter fusco-pubescens. Pedes cum coxis anticis testacei, coxis posterioribus piceo-testaceis.

Abdominis segmentum sextum inferius maris apice profunde incisum, quintum leviter longitudinaliter impressum, apice medio subsinuatum.

### **Dolicaon Laporte.**

*Laporte Etud. Ent. I. p. 119.*

*Adelobium Nordm. Symb. p. 139.*

*Labrum brevissimum, bilobum.*

*Palpi maxillares articulo ultimo obtuso.*

*Antennae rectae.*

*Tarsi articulo quarto simplice, posteriores articulis 4 primis decrescensibus.*

Corpus elongatum, lineare, aliis apterum, aliis alatum. Caput thoracis latitudine vel eodem parum angustius, subovatum, basi modice constrictum, collo quasi brevi crassiusculo thoraci immissum, exsertum, porrectum, oculis parvis rotundatis, leviter prominulis, ante capitis medium sitis. Labrum transversum, brevissimum, medio incisum. Mandibulae validae, falcatae, medio obtuse dentatae. Maxillae (t. IV. f. 27. b.) malis membranaceis, interiore interne, exteriori subelongata apice barbatis. Palpi maxillares modice elongati, articulo primo brevi, minuto, secundo tertioque elongatis, hoc apicem versus sensim incrassato, ultimo minimo, globoso, obtuso. Labium (t. IV. f. 27. a.) mento transversum, apice truncato, ligula brevi, biloba, lobis rotundatis, membranea, paraglossis triangularibus, coriaceis, interne ciliatis, ligulam longe superantibus. Palpi labiales articulo primo obconico, secundo elongato, primo sesqui fere longiore, tertio minuto, tenui, subulato. Antennae filiformes, rectae, articulo primo reliquis paulo maiore, ultimo apice acuminato. Thorax oblongus, coleopterorum fere latitudine. Elytra nonnunquam breviora. Abdomen lineare, apice angustatum, stylis analibus exsertis. Pedes modice elongati, antici leviter incrassati, femoribus anticis subtus ante apicem obsolete dentatis, tarsis anticis articulis 4 primis leviter dilatatis, obcordatis, subtus spongioso-tomentosis, tarsis posterioribus longioribus, gracilibus, articulis 4 primis latitudine sensim decrescensibus.

### **O b s e r v a t i o n e s.**

1. Habitus omnino *Lathrobii*, genus autem palporum maxillarum articulo ultimo obtuso, labro brevissimo et tarsorum posteriorum articulis primis elongatis evidenter distinctum. Quibus notis species Europaeae cum Capensi gigantea congruunt.

2. Mas abdominis segmento inferiore sexto profunde inciso, nonnunquam etiam quinto apicem versus leviter impresso distinctus.

3. Victus latet.

\* 1. *D. Lathrobioides: Niger, subnitidus, antennis pedibusque rufis, elytris thorace duplo brevioribus.* — Long. 8 lin.

*Laporte Etud. Ent. I. p. 120. t. 4. f. 1.*

*Adelobium brachypterum Nordm. Symb. 140. 1.*

Habitat ad Promont. Bonae Spei, Dom. Prof. Lichtenstein.

Apterus, niger, subnitidus. Antennae capite paulo longiores, tenuiores, articulo tertio secundo duplo longiore, sequentibus decrescensibus, elongato.

subobconicis, ultimo fusiformi, praecedente paulo longiore, rubrae. Mandibulae piceae. Caput thoracis latitudine eodemque paulo brevius, minus depressum, crebre minus profunde punctatum, fronte anteriore et linea media longitudinali laevioribus. Thorax coleopteris prope latior, latitudine sesqui longior, basin versus vix angustatus, lateribus levissime rotundatus, basi apiceque truncatus, angulis anterioribus deflexis, subrectis, posterioribus rotundatis, convexus, sat crebre minus profunde punctatus, linea media longitudinali laevi. Scutellum piceum, laeve. Elytra thorace duplo breviora, subdepressa, crebre punctata, parce subtiliterque nigro-pubescentia, apice oblique truncata, angulo exteriori emarginato. Abdomen subtiliter minus confertim exasperato-punctatum, parce subtiliterque pubescens, subtus apice piceum. Pedes rufi, coxis piceis.

Abdominis segmentum inferius sextum maris profunde incisum, quintum integrum.

\* 2. *D. Illyricus*: Niger, nitidus, elytris thorace dimidio brevioribus, abdominis segmentis ultimis duobus, pectore antennisque rufis, pedibus testaceis. — Long. 3 lin.

Habitat in Illyria, Dom. Schüppel.

Habitu fere praecedentis, at longe minor, subcylindricus, apterus, niger, nitidus, capite thoraceque pernitidis. Antennae capite paulo longiores, apicem versus subincrassatae, articulo tertio secundo perparum longiore, sequentibus sensim paulo brevioribus et crassioribus, ultimo apice abrupte acuminato, rufae. Palpi maxillares articulo tertio fortiter incrassato. Mandibulae et labrum picea. Caput thorace paulo latius, minus depressum, parcius profundiusque punctatum, linea media longitudinali laevi, antice utrinque ad antennarum basin intra marginem incrassatum subimpressum. Thorax coleopteris paululum latior, latitudine sesqui longior, basin versus vix angustatus, lateribus subrectis, basi apiceque truncatus, angulis omnibus rotundatis, subcylindricus, crebrius profundiusque punctatus, linea latiore longitudinali laevi. Scutellum laeve. Elytra thorace dimidio breviora, dorso parce, lateribus crebrius punctata, tenuiter griseo-pubescentia, nitidissima, rufa. Abdomen parcius subtiliusque punctatum, tenuiter griseo-pubescentia, segmento quinto sextoque rufis, illo summa basi nigro. Pectus rufum. Pedes testacei, coxis posticis rufis.

Abdominis segmentum inferius sextum maris profunde incisum, quintum integrum.

Obs. Sequentibus statura magis cylindrica et elytris brevioribus diffinis. Specimen Musei olim nom. Xantholini Illyrici a Com. Dejean Domino Schüppel communicatum est. Exinde monente Dom. Schüppel inter Lathrobia ill. Comes hanc speciem aptius enumeravit.

\* 3. *D. haemorrhous*: Niger, nitidus, elytris thoracis longitudine anoque rufis, illis summa basi nigris, antennis pedibusque rufo-testaceis. — Long.  $2\frac{1}{2}$  lin.

Habitat in Sardinia, Dom. Prof. Gené, in Sicilia, Dahl.

Linearis, subdepressus, alatus, niger, nitidus. Antennae capite sesqui longiores, apicem versus nullo modo incrassatae, articulo tertio secundo



vix longiore, sequentibus sensim paulo brevioribus, ultimo apice abrupte acuminato, rufo-testaceae. Palpi rufo-testacei, maxillares articulo tertio fortius incrassato. Os rufum. Caput thorace paulo angustius, subdepressum, crebrius punctatum, spatio lato medio, inter oculos interrupto, laeviore, apice utrinque foveola minuta notatum, parce nigro-pilosellum. Thorax coleopterorum latitudine, latitudine sesqui longior, lateribus subrectis, basi apiceque truncatus, angulis omnibus obtusis, subrotundatis, perparum convexus, crebrius subtiliusque punctatus, linea media longitudinali laevi, tenuiter griseo-pubescentia. Scutellum laeve. Elytra thoracis longitudine, minus confertim subtilius punctata, tenuiter griseo-pubescentia, tota testaceo-rufa, nitida, basi summa magis minusve nigricante. Abdomen parcius subtiliter punctatum, minus subtiliter griseo-pubescentia, segmentis quinto sextoque rufis, illo summa basi nigricante. Pedes toti cum coxis rufo-testacei.

Abdominis segmentum inferius maris profunde incisum, quintum apice obsolete impressum.

Obs. Proximus sequenti, at distinctus videtur, elytris nonnisi summa basi nigricantibus, abdominis segmentis quatuor anterioribus margine concoloribus, thorace paulo longiore, crebrius punctato.

\* 4. *D. gracilis*: Niger, nitidus, elytris thorace longioribus, macula magna apicali anoque rufis, antennis pedibusque testaceis. — Long. 3 lin.

*Lathrobium gracile* Grav. Micr. 182. 5. — Latr. Hist. nat. Crust. et Ins. IX. 341. 11.

Habitat in Lusitania.

Statura omnino praecedentis, niger, nitidus, capite thoraceque pernitiidis. Antennae capite thoraceque breviores, tenues, apicem versus nullo modo incrassatae, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus, ultimo oblongo-ovato, apice sensim acuminato, rufo-testaceae. Os rufo-testaceum, palpis maxillaribus articulo tertio modice incrassato. Caput thorace paululum angustius, oblongum, parcius disperse punctatum. Thorax coleopterorum fere latitudine, latitudine tertia parte longior, lateribus subrectis, basi apiceque truncatus, angulis omnibus rotundatis, parum convexus, crebre subtilius punctatus, linea longitudinali media laevi, tenuiter subtiliterque griseo-pubescentia. Scutellum laeve. Elytra thorace paululum longiora, sat crebre punctata, tenuiter griseo-pubescentia, nitida, nigra, macula permagna apicali rufa, elytri dimidiam partem occupante. Abdomen parcius punctulatum, griseo-pubescentia, subnitidum, segmentis quatuor primis margine apicali tenui rufo, quinto rufo, basi dimidiatim nigro, sexto rufo. Pedes toti cum coxis testacei.

Abdominis segmentum inferius sextum maris profunde incisum, quintum apice obsolete longitudinaliter impressum.

\* 5. *D. biguttulus*: Niger, nitidus, punctatus, elytris thorace paulo longioribus, macula apicali rufa, antennis pedibusque testaceis. — Long.  $2\frac{1}{3}$  —  $2\frac{2}{3}$  lin.

*Lathrobium biguttulum* Boisd. et Lacord. Faun. Ent. Paris. I. 425. 7.

Habitat in Austria, Hungaria.

Statura praecedentis, at plerumque minor, niger, nitidus. Antennae

capite thoraceque breviores, tenues, articulo tertio secundo sesqui longiore. 4 — 10 sensim paulo brevioribus, ultimo ovato, apice abrupte obtuse acuminato, testaceae. Os rufo-testaceum, palpis maxillaribus articulo tertio fortiter incrassato. Caput thorace paulo angustius, oblongum, subdepressum, parcius disperse punctatum, punctis in media fronte rarioribus, apice utrinque leviter foveolatum. Thorax coleopteris paulo angustior, latitudine tertia parte longior, lateribus rectis, parallelis, basi apiceque truncatus, angulis anterioribus rectis, posterioribus rotundatis, parum convexus, sat crebre punctatus, linea media longitudinali laevi parum distincta, tenuiter subtiliterque griseo-pubescent. Scutellum laeve. Elytra thorace paululum longiora, crebre subtilius punctata, tenuiter griseo-pubescentia, nigra, magna apicali laete rufa, elytri quartam fere vel tertiam partem occupante, in marginem lateralem dilatata, sutura tamen nigra. Abdomen parcius punctulatum, griseo-pubescent, subtus apice piceum, segmentis intermediis tenuiter testaceo-marginatis. Pedes testacei, coxis omnibus picescentibus.

Abdominis segmentum inferius sextum maris profunde incisum, quintum apice obsolete impressum.

### **Scimbalium Er.**

*Labrum breve, bilobum.*

*Palpi maxillares articulo ultimo acuto.*

*Antennae rectae.*

*Tarsi articulo quarto simplice, posteriores articulis 4 primis sensim brevioribus.*

Corpus elongatum, valde depressum, alatum. Caput porrectum, exsertum, collo quasi angustiore thoraci immissum, antrorsum angustatum, oculis parvis, haud prominulis, ante capitis medium situs. Labrum breve, bilobum, lobis rotundatis, basi corneum, apice membranaceum. Mandibulae breviusculae, validiusculae, basi interne fimbria angusta membranacea instructae, medio fortiter 3-dentatae, apice fortiter falcatae, acutae. Maxillae malis parvis, coriaceis, interiore interne, exteriori apice barbatis. Palpi maxillares breviores, articulo primo minimo secundo tertioque longitudine aequalibus, hoc apicem versus sensim leniter incrassato, quarto minuto, brevi, subulato. Labium mento brevi, transversum, ligula membranacea, profunde biloba, lobis rotundatis, paraglossis ligulam superantibus, linearibus, coriaceis, interne barbatis. Palpi labiales articulo primo minore, leviter obconico, secundo maiore, incrassato, tertio tenui, subulato, secundo brevior. Antennae rectae, apicem versus subattenuatae, articulo primo crassiusculo, cylindrico, maiore, secundo duplo brevior, obconico, tertio secundo sesqui longiore, sequentibus sensim paulo brevioribus, elongato-subobconicis, ultimo elongato-fusiformi, apice sensim acuminato. Thorax coleopterorum fere latitudine, trapeziformis, basi apiceque truncatus, angulis anterioribus subrectis, posterioribus rotundatis. Scutellum apice rotundatum. Elytra apice introrsum oblique truncata, angulo exteriori rotundato. Abdomen lineare, apice attenuatum, segmento sexto conico, stylis analibus exsertis. Pedes breviores, antici modice incrassati, femoribus tibiisque interne leviter sinuatis, tibiis posterioribus parce spinulosis, spinis apicalibus tenuibus, distinctis, tarsis anticis in utroque sexu in patel-

lam ovalem, subtus tomentosam, dilatatis, posterioribus sat elongatis, tibiam longitudine aequantibus, articulis 4 primis sensim brevioribus, ultimo praecedentibus ambobus vix longiore.

### O b s e r v a t i o n e s.

1. Genus sat distinctum, licet ab omnibus hucusque auctoribus cum Achenio confusum, quocum corpore deplanato primo quidem intuitu congruit, oris autem, praecipue labii et tarsorum structura minime convenit.

2. Mas distinctus abdominis segmento inferiore sexto apice profunde triangulariter exciso, septimo inferiore conspicuo.

3. Victus latet.

\* 1. *Sc. anale: Piceum, nitidum, antennis, pedibus, ano elytrisque rufis, his thorace longioribus.* — Long.  $4\frac{1}{2}$  lin.

*Achenium anale Nordm. Symb. 153. 2.*

Habitat in Chersoneso Taurica et in Caucaso.

Achenio depresso paulo longius et minus depressum, nigro-piceum, nitidum. Antennae capite thoraceque longiores, rufae. Palpi rufi. Mandibulae rufo-piceae. Caput thorace angustius, antrorsum leviter angustatum, lateribus crebrius, medio parcius punctatum, parcius pube erecta nigra vestitum, lateribus pilosellum. Thorax coleopteris vix angustior, latitudine paululum longior, basin versus subangustatus, lateribus subrectis, subdepressus, minus fortiter aequaliter crebre punctatus, linea media longitudinali laevi, pube subtili erecta nigra parcius vestitus. Scutellum subtiliter punctulatum. Elytra thorace paulo longiora, subtiliter dense punctata, tenuiter griseo-pubescentia, rufa, parum nitida. Abdomen densius subtiliusque punctatum, densius griseo-pubescentia, piceum, segmentis 4 primis apice tenuiter rufo-marginatis, quinto rufo, basi piceo, sexto toto rufo. Pedes testaceo-rufi, coxis concoloribus.

Obs. 1. *Achenium Caucasicum* Laporte Etud. Ent. I. p. 119 „fortiter punctatum, leviter pubescens, rufo-brunneum, antennis, oris partibus pedibusque dilute castaneis,” a Dom. Godet in Caucaso lectum, forte idem, at descriptio nimis manca, ut equidem insectum illud pro alio quodam haberem, si ipse in Staphylinorum collectione sua, quam Dom. Godet examinandam mihi amice communicavit, aliud quoddam, cui magis conveniret, observassem.

2. Nomine *Lathrobii phoenicuri* hoc insectum Dom. Steven communicavit.

\* 2. *Sc. planicollis: Piceum, nitidum, antennis, pedibus, ano elytrisque rufis, his thoracis longitudine.* — Long.  $4\frac{1}{2}$  lin.

Habitat in Italia, Dom. Schüppel, in Sicilia, Dom. Dr. Schultze.

Statura et summa affinitas praecedentis. Nigro-piceum, capite thoraceque saepius supra nigris, nitidis. Antennae capite thoraceque paulo longiores, rufae. Palpi rufo-testacei. Caput basi thorace parum angustius, antrorsum modice angustatum, parcius profundiusque punctatum, fronte media late laeviore, pube brevi rara erecta nigra vestitum, lateribus nigro-pilosellum. Thorax coleopterorum latitudine, latitudine antica paulo longior, basin versus leviter angustatus, lateribus subrectis, supra subdepressus, crebrius minus subtiliter punctatus, linea media longitudinali laevi, parce subtilissime griseo-pubescentia. Scutellum punctulatum, nigrum. Elytra thoracis longi-



tudine, densius minus subtiliter punctata, rufa, nitida, subtiliter griseo-pubescentia. Abdomen minus subtiliter crebrius punctulatum, densius griseo-pubescentis, segmentis 4 primis apice piceo-marginatis, quinto apice, sexto toto rufis. Pedes rufo-testacei, coxis nigro-piceis.

Obs. Praecedenti proximum, elytris autem brevioribus, capite thoraceque minus subtiliter minusque dense punctatis distinctum videtur.

\* 3. *Sc. testaceum*: *Dilute testaceum, nitidum, capite thoraceque parce punctatis.* — Long.  $2\frac{1}{2}$  lin.

Habitat in Sicilia, Dom. Dr. Schultz.

Praecedentibus multo minus, fortiter depressum, testaceum, nitidulum. Antennae capite thoraceque paulo longiores. Caput thorace paulo latius, antrorsum leviter angustatum, parce punctatum, plaga angusta longitudinali media laevi, subtiliter griseo-pubescentis, lateribus longe nigro-pilosellum. Thorax coleopteris paulo angustior, latitudine longior, basin versus modice angustatus, utrinque parcius subtiliter punctatus, parcius subtiliter griseo-pubescentis. Scutellum punctulatum. Elytra thoracis longitudine, dense subtiliter punctata, pube depressa grisea dense vestita. Abdomen dense subtiliter punctulatum, pubescens, parce pilosellum, segmentis 4 primis nonnunquam fuscescentibus.

Obs. Forte apterum.

### **Achenium Leach.**

*Curt. Britt. Ent. III. t. 115. — Steph. Illustr. V. p. 265. — Boisd. et Lacord. Faun. Ent. Paris. I. p. 420. — Er. Col. March. I. p. 501.*

*Lathrobium Fam. II. Grav. Mon. p. 129.*

*Labrum bilobum, lobis divergentibus.*

*Palpi maxillares articulo ultimo acuto.*

*Antennae rectae.*

*Tarsi articulo quarto simplice, posteriores articulis 4 primis aequalibus, quinto his coniunctis aequali.*

Corpus elongatum, valde depressum vel planum, alatum. Caput subcordatum, porrectum, exsertum, collo quasi angusto thoraci immisum, basi leviter emarginatum, angulis posterioribus rotundatis, antrorsum triangulariter angustatum, ante oculos subito angustius, oculis versus capitis medium sitis, parvis, rotundatis, haud prominulis. Labrum angustius, profunde bilobum, lobis elongatis, leviter divergentibus. Mandibulae breviusculae, validiusculae, basi interne dilatatae, medio fortiter unidentatae, apice fortiter falcatae, acutae. Maxillae mala interiore membranacea interne, exteriore cornea apice membranaceo barbatis. Palpi maxillares minus elongatae, articulo primo minuto, secundo tertioque aequalibus, hoc apicem versus sensim subincrassato, quarto minimo, conico-acuminato, seu obtuse subulato. Labium mento brevi, transversum, ligula brevi, membranacea, biloba, lobis rotundatis, paraglossis coriaceis, ligulam haud superantibus. Palpi labiales articulis duobus primis subcylindricis, secundo paulo longiore et crassiore, tertio tenui, subulato, secundo brevior. Antennae rectae, apicem versus attenuatae, articulo primo crassiusculo, cylindrico, secundo hoc duplo brevior, obconico, tertio secundo sesqui longiore, sequentibus sensim minoribus,

elongato-subobconicis, ultimo elongato-fusiformi, apice sensim acuminato. Thorax coleopteris paulo angustior, trapeziformis, apice truncatus, angulis anterioribus lateraliter leviter prominulis, posterioribus rotundatis. Scutellum triangulare. Elytra truncata. Abdomen lineare, apice angustatum, segmento septimo prominulo, inferiore bivalvi, valvulis apice acuminato, sursum uncinato terminatis. Pedes breviusculi, validiusculi, antici incrassati, femoribus tibiisque interne sinuatis et subdentatis, tibiis posterioribus setulosis, apicem versus leviter dilatatis, apice oblique truncatis, spinis apicalibus brevissimis, tarsis anticis in utroque sexu articulis 4 primis in patellam orbicularem subtus tomentosam dilatatis, posterioribus tenuibus, articulo primo brevissimo, secundo parum elongato, tertio huic subaequali, quarto iterum breviori, ultimo his coniunctis longitudine aequali.

#### O b s e r v a t i o n e s.

1. Tarsis posterioribus brevibus *Lathrobio* affine, at tarsi tenuiores et articulus quintus longior, ceterum praecipue corpore deplanato, labro angusto, lobis magis elongatis, mandibulis brevioribus distinctum. Thorax saepius per plagas punctatus, plagis intermediis in medio dorso distinctioribus.

2. Mas abdominis segmento inferiore sexto apice inciso distinctus.

3. Victus palustris.

\* 1. *A. depressum*: *Nigrum, nitidum, pedibus, ano elytrisque rufis, his basi nigris, thorace paulo longioribus, capite parcius punctato, thorace per plagas punctato.* — Long. 4 lin.

*Nordm. Symb.* 152. 1.

*Lathrobium depressum* Grav. *Micr.* 182. 6. *Mon.* 129. 2. — *Latr. Hist. nat. Crust. et Ins.* IX. 341. 12. *Gen. Crust. et Ins.* I. 290. 2.

*Achenium cordatum* Boisd. et Lacord. *Faun. Ent. Paris.* I. 421. I.

Habitat in Lusitania, Italia.

Nigrum, nitidum. Antennae capite thoraceque paulo longiores, rufo-piceae. Palpi rufo-testacei. Labrum rufo-piceum. Caput basi thoracis latitudine vel paulo latius, leviter emarginatum, antrorsum angustatum, triangulare, parcius punctatum, interstitiis disperse subtiliter punctulatis, fronte media sublaevi, parce subtiliterque nigro-pubescens. Thorax coleopteris paulo angustior, latitudine vix longior, basin versus leviter angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, angulis anterioribus subrectis, depressus, utrinque plagis tribus punctatus, plagis intermediis densius punctatis, distinctioribus, posterius subimpressis, interstitio laevissimo. Scutellum subtiliter parce punctatum. Elytra thorace paulo longiora, minus dense basi subseriatim, apice vage punctata, rufa, nitida, basi tertia fere parte nigra. Abdomen subtiliter densius punctatum, subtiliter griseo-pubescens, nigrum, segmentis 4 primis apice tenuiter flavo-marginatis, quinto apice, sexto toto rufis. Pedes cum coxis anterioribus rufi, tibiis basi piceis.

Abdominis segmentum inferius sextum maris leviter triangulariter excisum.

Variat femoribus posticis picescentibus.

- \* 2. *A. aequatum*: Nigrum, nitidum, pedibus, ano elytrisque rufis, his thorace sesqui longioribus, capite thoraceque subtilius crebre punctatis. — Long. 4 lin.

Habitat in Aegypto, Dom. Prof. Ehrenberg.

Statura et summa affinitas praecedentis, at paulo latius, nigrum, nitidum. Antennae capite thoraceque vix longiores, rufo-piceae, densius pubescentes et pilosellae. Palpi rufo-testacei. Labrum rufo-piceum. Caput basi thoracis latitudine, leviter emarginatum, antrorsum angustatum, triangulare, supra subtilius licet profunde crebre fere aequaliter punctatum, fronte media vix laeviore, interstitiis parce subtilissime punctulatis, pube tenui subtilique erecta cinerea vestitum, lateribus parce pilosellum. Thorax coleopteris paulo angustior, latitudine antica vix longior, basin versus subangustatus, lateribus rectis, basi truncatus, angulis posterioribus late rotundatis, subtiliter sat crebre fere aequaliter punctatus, parce pube erecta subtili grisea vestitus, linea media longitudinali laevi, piceo-niger, angulis anterioribus dilutioribus. Scutellum parce punctatum, nigrum, opacum. Elytra thorace sesqui longiora, aequaliter sat crebre subtiliter punctata, tota testaceo-rufa, pube erecta grisea densius vestita. Abdomen densius subtiliter punctatum, pube longiore depresso grisea densius vestitum, parce nigropilosellum, minus nitidum, nigrum, segmentis 4 primis apice tenuiter flavo-marginatis, quinto margine apicali, sexto apice rufis, subtus magis piceum. Pedes cum coxis omnibus rufo-testacei, tibiis summa basi picescentibus.

Abdominis segmentum inferius quintum maris medio longitudinaliter impressum, sextum longitudinaliter impressum, apice triangulariter excisum.

Obs. A praecedente praecipue capitis, thoracis elytrorumque punctura crebriore, fere aequali, thorace basin versus vix angustato, elytris longioribus, totis rufis distinctum.

- \* 3. *A. planum*: Nigrum, nitidum, pedibus, ano elytrisque rufis, his basi late nigris, thorace sesqui longioribus, capite thoraceque confertim subtilius punctatis. — Long. 4 lin.

Habitat in Grusinia, Dom. Marschall v. Bieberstein.

Statura et summa affinitas praecedentis, nigrum, nitidum. Antennae capitis thoracisque longitudine, rufo-piceae. Palpi rufi. Labrum rufo-piceum. Caput basi thoracis latitudine, leviter emarginatum, antrorsum angustatum, triangulare, crebre fere aequaliter punctatum, fronte media vix laeviore, interstitiis hinc inde puncto subtilissimo notatis, pube rara subtili erecta grisea vestitum, subtus longius pubescens. Thorax coleopteris paulo angustior, latitudine parum longior, basin versus subangustatus, lateribus rectis, basi subtruncatus, angulis posterioribus late rotundatis, anterioribus subrectis, depressus, crebrius subtiliter punctatus, linea longitudinali media laevi, punctorum crebrorum plaga utrinque terminata, nigerrimus, glaber. Scutellum nigrum, parce subtiliter punctatum. Elytra thorace sesqui longiora, densius aequaliter subtiliter punctata, testaceo-rufa, nitida, basi ultra tertiam partem nigra. Abdomen subtiliter confertim punctatum, nigrum, parum nitidum, segmentis quatuor anterioribus apice tenuissime flavo-marginatis, quinto margine apicali, sexto apice rufis. Pedes testaceo-rufi, coxis anterioribus concoloribus, posticis piceis, tibiis basi piceis.



Abdominis segmentum inferius sextum maris apice leviter triangulariter excisum.

Obs. Differt a praecedente elytris basi nigris, thorace minus aequaliter punctato, glabro, capite minus crebre punctato; ab *A. depresso* et *A. ephippio* elytris longioribus, densius aequaliter punctatis, thorace basin versus vix angustato etc. magis distinctus.

- \* 4. *A. ephippium*: *Nigrum, nitidum, elytris apice oblique rufis, thorace paulo longioribus, capite parcius punctato, thorace, per plagas punctato.* — Long. 4 lin.

Habitat in Hungaria.

*A. depresso* aequale et simillimum at paulo fortius depressum, nigrum, nitidum. Antennae capite thoraceque paulo longiores, rufo-piceae. Palpi et labrum rufa. Caput, thorax et elytra pube parca brevi subtili erecta grisea vestita. Caput basi thoracis latitudine, leviter emarginatum, antrorsum angustatum, triangulare, supra parcius fortius punctatum, interstitiis parce subtilissime punctatis, fronte media laeviore, lateribus pilosellum. Thorax coleopteris tertia parte angustior, basin versus angustatus, lateribus rectis, basi truncatus, angulis posterioribus late rotundatis, anterioribus rectis, modice prominulis, supra parcius inaequaliter punctatus, spatio medio longitudinali laevissimo, utrinque plaga punctorum crebrorum terminata. Scutellum laeve. Elytra thorace paulo longiora, minus dense inaequaliter hinc inde subseriatim punctata, rufo-testacea, basi late a humeris usque ad suturae apicem prope triangulariter descendente nigra, seu nigra, angulo apicali exteriori late cum margine laterali et apicali rufo. Abdomen crebrius subtiliter punctatum, pube depressa grisea vestitum, segmentis 4 anterioribus apice tenuiter flavo-, quinto sextoque piceo-marginatis. Pedes cum coxis anterioribus iam rufi, iam rufo-, iam fusco-picei, tarsis testaceis.

Abdominis segmentum inferius sextum maris apice leviter triangulariter excisum, quintum medio late leviter sinuatum.

Obs. Distinctum ab *A. depresso* praecipue corpore magis deplanato, elytrorum spatio nigro ultra medium descendente, sat determinato, ano concolore, etc.

- \* 5. *A. basale*: *Nigrum, nitidum, elytris rufis, basi nigris, thorace paulo longioribus, capite parcius punctato, thorace per plagas punctato.* — Long. 4 lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura omnino *A. depressi*, et minoribus eius individuis aequale, nigerrimum, nitidum, capite thoraceque pube parciore erecta grisea nigraque vestitis. Antennae capite thoraceque paulo longiores, rufo-vel fusco-piceae, apice dilutiores. Palpi rufi. Labrum rufo-piceum. Caput basi thoracis latitudine, leviter emarginatum, antrorsum angustatum, triangulare, supra parcius, minus subtiliter punctatum, fronte media laeviore, interstitiis parce subtilissime punctatis, lateribus nigro-pilosellum. Thorax coleopteris paulo angustior, latitudine antica paulo longior, basin versus leviter angustatus, lateribus rectis, basi truncatus, angulis posterioribus late rotundatis, anterioribus acutiusculis, supra parcius punctatus, punctis utrinque in plagas tres longitudinales digestis, seriebus intermediis distinctioribus, crebrius pun-

etatis, posterius leviter impressis, interstitio intermedio laevisimo, interstitiis lateralibus interioribus ante medium punctis numerosis repletis. Scutellum laeve. Elytra thorace paulo longiora, densius hinc inde subseriatim punctata, tenuiter subtiliterque griseo-pubescentia, saturate rufa, nitida, basi et humeris arcuatim quarta vel tertia parte nigra. Abdomen nigrum, subnitidum, dense subtilius punctatum, tenuiter griseo-pubescentis, segmentis singulis apice tenuiter fusco-piceo-marginatis. Pedes nigro-picei, tarsis testaceis.

Abdominis segmentum inferius sextum maris profundius triangulariter incisum.

Variat pedibus rufo-piceis, coxis omnibus concoloribus.

Obs. Proximum A. depresso statura et elytrorum proportionem, distinctum autem capite thoraceque minus subtiliter, elytris crebrius punctatis, ano concolore, pedibus obscurioribus, etc.

\* 6. *A. debile*: Nigro-piceum, nitidum, pedibus testaceis, elytris thorace paulo longioribus, basi nigris anoque rufis, capite crebre punctato, thorace lateribus parce, medio biserialim punctato. — Long.  $3\frac{3}{4}$  lin.

Habitat in Mesopotamia, Dom. Dr. Helfer, in Aegypto Dom. Prof. Ehrenberg.

Statura et summa affinitas A. depressi, at paululum minus et gracilius, nigro-piceum, nitidum. Antennae capite thorace paulo longiores, testaceae. Caput thorace paulo latius, antrorsum angustatum, basi leviter emarginatum, triangulare, nitidum, crebre fere aequaliter punctatum, interstitiis disperse subtilissime punctatis, fronte media parum laeviore, subtilissime griseo-pubescentis. Thorax coleopteris paulo angustior, latitudine antica paulo longior, basin versus modice angustatus, lateribus rectis, basi truncatus, angulis posterioribus late rotundatis, anticis subrectis, nitidus, omnium subtilissime parce disperseque punctulatus, dorso biserialim, lateribus parce vageque subtiliter punctatus. Scutellum obsolete punctulatum. Elytra thorace paulo longiora, parcius, anterieus subseriatim, posterius vage punctata, rufa-testacea, nitida, basi quarta fere parte nigricantia, parce subtilius griseo-pubescentia. Abdomen crebre subtiliterque punctatum, subtilius griseo-pubescentis, nigrum, subnitidum, segmentis 4 primis apice subtiliter piceo-marginatis, quinto rufo, basi piceo, sexto toto rufo. Pectus piceum. Pedes cum coxis omnibus testacei, tibiis summa basi picescentibus.

Abdominis segmentum inferius sextum maris apice triangulariter excisum.

Variat immaturus colore corporis estaceo elytris anoque dilutioribus, pedibus flavis.

Obs. Ab A. depresso elytris parcius, capite crebre punctatis, ab A. striato elytris longioribus latioribusque, ab omnibus thorace lateribus parce punctato distinctum.

\* 7. *A. striatum*: Nigrum, nitidum, pedibus, abdominis segmentis ultimis duobus elytrisque rubris, his thorace parum longioribus, subseriatim punctatis, thorace per plagas punctato. — Long.  $3\frac{3}{4}$  lin.

*Lathrobium striatum* Latr. Hist. nat. Crust. et Ins. IX. 341. 13.

Habitat in Gallia, Italia.

Statura omnino *A. depressi*, at paulo angustius, nigrum, nitidum, capite, thorace elytrisque pube rara erecta grisea vestitis. Antennae capite thoraceque paulo longiores, rufo-piceae. Os rufo-piceum. Caput basi thorace paulo latius, antrorsum fortiter angustatum, basi leviter emarginatum, subcordatum, crebre punctatum, interstitiis subtilissime disperse punctulatis, fronte media laeviore. Thorax coleopteris haud angustior, latitudine antica parum longior, basin versus modice angustatus, lateribus rectis, basi truncatus, angulis posterioribus late rotundatis, anterioribus subrectis, subtilissime disperse punctulatus, utrinque triplagiatis subtilius parciusque punctatus, plagis angustis, intermediis distinctioribus, densius punctatis. Scutellum nigrum, parce subtiliterque punctatum. Elytra thorace parum longiora, parcius subseriatim punctata, tota rubra, nitida. Abdomen subtilius crebre punctatum, densius griseo-pubescent, nigro-piceum, subnitidum, segmentis 4 primis margine apicali tenuissimo, testaceo, ultimis duobus totis rubris. Pedes cum coxis omnibus rufo-testacei, tibiis basi summa picescentibus.

Abdominis segmentum inferius sextum maris apice profundius triangulariter excisum.

Obs. Distinctum a praecedentibus coleopteris brevioribus, thorace haud latioribus, abdominis segmentis duobus ultimis totis rufis. Caput latiusculum, lateribus magis rotundatum, ideoque magis cordatum. Abdomen ante apicem sensim subdilatum.

\* 8. *A. humile*: Nigrum, capite thoraceque piceis, antennis, pedibus, ano elytrisque testaceis, his summa basi fusciscentibus, thorace parum longioribus, vage punctatis. — Long. 3 lin.

*Er. Col. March. I. 501. 1.*

*Lathrobium humile Nicolai Col. Halens. 38. 1.*

*Achenium depressum Curt. Britt. Ent. III. t. 115. — Boisd. et Lacord. Faun. Ent. Paris. I. 421. 2.*

Habitat in Germania.

Habitu omnino praecedentium, at duplo minus. Antennae capite thoraceque longiores, rufae. Palpi rufo-testacei. Caput basi thorace paulo latius, vix emarginatum, subcordatum, antrorsum angustatum, parcius subtiliusque punctatum, fronte media laeviore, parce griseo-pubescent, lateribus longius pilosellum, plerumque dilute rufo-, rarius fusco- vel nigro-piceum, nitidum. Thorax coleopterorum latitudine, latitudine antica parum longior, basin versus leviter angustatus, lateribus rectis, basi truncatus, angulis posterioribus rotundatis, anterioribus acutiusculis, parcius subtiliter punctatus, punctis utrinque in plagas tres longitudinales digestis, plagis intermediis angustioribus, magis regularibus, distinctioribus, medio nonnunquam leviter impressis, plerumque dilute rufo-, rarius fusco- vel nigro-piceus, nitidus, pube parca erecta grisea vestitus. Scutellum nigro-piceum, laeve. Elytra thorace perparum longiora, subtilius parciusque aequaliter punctata, pube tenui erecta grisea vestita, nitida, testacea, basi indeterminate nigricantia. Abdomen crebrius subtiliter punctatum, opacum, densius griseo-pubescent, segmentis 5 primis apice subtiliter flavescenti-marginatis, sexto piceo-testaceo. Pectus nigrum. Pedes testacei, coxis concoloribus, tibiis basi piceis.

Abdominis segmentum inferius sextum maris apice triangulariter excisum, quintum medio subsinuatum.



9. *A.ieiunum*: *Nigrum, capite, thorace elytrisque rufo-piceis, capite crebrius punctato, elytris thoracis longitudine, inaequaliter punctatis.* — Long.  $3\frac{2}{3}$  lin.

Habitat in Sardinia, Mus. Reg. Taurin.

Statura et summa affinitas *A. humilis*, at paulo maius, *A. striato* aequale. Antennae capite thoraceque paulo longiores, rufae, apice dilutiores. Palpi rufo-testacei. Caput basi thorace paulo latius, cordatum, basi emarginatum, lateribus pone oculos leviter rotundatum, antice fortiter angustatum, depressum, crebrius punctatum, interstitiis subtiliter punctulatis, fronte media laeviore, parce pubescens pilosumque, saturate rufo-piceum, nitidum, ore dilutiore. Thorax apice coleopterorum latitudine vel iis paululum latior, basin versus leviter angustatus, latitudine antica parum longior, basi cum angulis posterioribus fortiter rotundatus, apice utrinque oblique truncatus, angulis anterioribus prominulis, depressus, dorso biseriatim, versus latera parce disperse punctatus, interstitiis subtilissime parce punctulatis, lateribus parce pilosellus, saturate rufo-piceus, nitidus. Scutellum saturate rufo-piceum, laeve. Elytra thoracis longitudine, crebre inaequaliter minus subtiliter punctata, subrugosa, longius griseo-pilosa, rufo-picea, nitida. Abdomen apicem versus sensim leviter dilatatum, confertim minus subtiliter punctatum, griseo-pubescent, minus nitidum, nigrum, segmentis 4 primis margine apicali tenui piceo-testaceo, quinto apice, sexto septimoque totis rufo-piceis. Pectus nigrum. Pedes dilute testacei, tibiis basi obscurioribus.

Obs. Ab *A. striato* differt colore, capite thoraceque minus crebre punctatis, ab *A. humili* statura maiore, capite basi profundius emarginato, abdomine minus subtiliter punctato, ab utroque elytris brevioribus, fortius et inaequaliter punctatis, subrugosis.

10. *A. tenellum*: *Nigro-piceum, nitidum, ore, antennis, pedibus elytrisque testaceis, his thoracis longitudine, parce subtiliterque punctatis, capite parcius punctato.* — Long. 2 lin.

Habitat in Sardinia, Mus. Reg. Taurin.

Minutum, *A. humili* plus duplo minus, angustulum, valde depressum, piceo-nigrum, nitidum. Antennae capite thoraceque paulo longiores, testaceae. Caput basi thorace paululum latius, subcordatum, basi subemarginatum, lateribus pone oculos leviter rotundatum, apice sat fortiter angustatum, planum, sat fortiter profundeque disperse punctatum, fronte media laeviore, interstitiis laevissimis, tenuiter pubescens, lateribus longius pilosellum, ore testaceo. Thorax oblongus, apice coleopterorum basi latitudine aequalis, basin versus leviter angustatus, latitudine longior, basi apiceque truncatus, angulis posterioribus fortiter rotundatis, anterioribus obtusiusculis, modice prominulis, dorso biseriatim, utrinque parce subtiliusque disperse punctatus, tenuiter pilosellus. Elytra thoracis longitudine, apicem versus sensim paulo latiora, depressa, parce subtiliterque aequaliter punctata, tenuiter pilosella, nitida, luteo-testacea, basi prope scutellum picescentia. Abdomen apicem versus parum dilatatum, confertim subtiliter punctatum, densius griseo-pubescent, minus nitidum, nigrum, segmentis 5 primis margine apicali tenui testaceo, sexto apice septimoque toto piceis, subtus fere totum rufo-piceum. Pedes luteo-testacei.

**Lathrobium Grav.**

*Mannerh. Brachelytr. p. 37. — Curt. Britt. Ent. XIV. t. 650. — Steph. Illustr. V. p. 265. — Boisd. et Lacord. Faun. Ent. Paris. I. p. 421.*

*Lathrobium Fam. IV. Grav. Mon. p. 130.*

*Labrum breve, bilobum.*

*Palpi maxillares articulo ultimo acuto.*

*Antennae rectae.*

*Tarsi articulo quarto simplice, posteriores articulis 4 primis aequalibus, ultimo his coniunctis brevior.*

Corpus elongatum, lineare, plerumque alatum. Caput subquadratum, vel suborbiculatum, thoracis circiter latitudine, basi profunde constrictum, collo quasi brevi angusto thoraci immissum, exsertum, porrectum, oculis parvis rotundatis, parum prominulis. Labrum transversum, bilobum, lobis divergentibus, totum corneum, apice setosum. Mandibulae falcatae, acutae, basi dilatatae, medio fortiter unidentatae. Maxillae (t. IV. f. 26. b.) malis coriaceis, interiore interne, exteriori apice dense barbatis. Palpi maxillares modice elongati, articulo primo minuto, secundo tertioque elongatis, hoc paulo longiore, apicem versus modice incrassato, quarto minuto, tenui, subulato. Labium (t. IV. f. 26. a.) mento brevi transversum, ligula membranea, biloba, lobis rotundatis, paraglossis acuminatis, membranaceis, ligulam paulum superantibus. Palpi labiales articulis duobus primis cylindricis, secundo paulo longiore, tertio secundo longitudine subaequali, at tenui, acuminato. Antennae filiformes, rectae, articulo primo reliquis paulo maiore, ultimo apice sensim acuminato. Thorax coleopterorum prope latitudine, saepius oblongus, basi apiceque truncatus, angulis omnibus rotundatis. Scutellum semicirculare. Elytra truncata, angulo exteriori rotundato. Abdomen lineare, apice angustatum, segmento septimo saepius conspicuo, stylis analibus valviformibus. Pedes breves, validi, antici incrassati, femoribus saepe subtus subdentatis, tibiis interne longitudinaliter excavatis, interne subdentatis, tibiis omnibus pubescentibus, posticis apice subcompressis, oblique truncatis, spinis apicalibus inaequalibus, altero sat distincto, altero minuto, tarsis anticis articulis 4 primis in patellam suborbicularem, subtus tomentosam dilatatis, posterioribus articulo primo secundo brevior, hoc subelongato, tertio quartoque brevibus, primo aequalibus, ultimo reliquis coniunctis brevior.

**Observationes.**

1. Corpus punctatum, punctis in capite et thorace saepius rarioribus, hoc nonnunquam dorso seriatis, pubescens, pube capitis thoracisque rariore, e punctis oriunda, erecta, brevissima, abdominis saepius densiore, depressa.

2. Mas abdominis segmento inferiore sexto inciso, praecedentibus saepius impressis, quinto nonnunquam apice sinuato, femina segmento inferiore sexto integro vel apice producto distinguuntur.

3. Victus sub musco, foliis deciduis et lapidibus in silvis paludosis.

\* 1. *L. brunripes*: Nigrum, nitidum, punctatum, thorace oblongo, elytris concoloribus, subtilius punctatis, antennis pedibusque rufis. — Long. 4 lin.

*Grav. Micr. 56. 10. Mon. 131. 6. — Latr. Hist. nat. Crust. et Ins. IX. 340. 10. — Gyll. Ins. Suec. II. 366. 3. — Sahlb. Ins. Fenn. I. 338. 4. — Zetterst. Faun. Lappon. I. 83. 3. Ins. Lappon. 67. 3. — Mannerh. Brachelytr. 37. 6. — Boisd.*

et Lacord. Faun. Ent. Paris. I. 422. 1. — Nordm. Symb. 141. 6. — Er. Col. March. I. 503. 1.

*Paeder. brunnipes* Fab. Ent. Syst. I. II. 537. 5. Syst. El. II. 609. 5. — Panz. Ent. Germ. 363. 6.

*Paeder. elongatus* var. c Payk. Faun. Suec. III. 429. 3.

*Paeder. elongatus* var. 1 Walken. Faun. Paris. I. 275. 3.

*Staph. dentatus* Marsh. Ent. Britt. I. 515. 53.

*Staph. punctatus* Fourcr. Ent. Par. I. 168. 14.

*Le Staph. à pattes fauves et à étuis pointillés* Geoffr. Hist. Ins. I. 366. 14.

Habitat in Europa.

Statura et summa affinitas *L. elongati*, at plerumque paululum maius et robustius, nigrum, capite thoraceque pernitidis, pube parca brevi subtile erecta grisea vestitis. Antennae capitis thoracisque longitudine, graciliores, articulo tertio secundo sesqui longiore, 4 — 10 subaequalibus, elongato-obconicis, rufae. Palpi et labrum rufa. Mandibulae rufo-piceae. Caput thoracis prope latitudine, suborbiculatum, leviter convexum, crebre punctatum, punctis in media fronte paulo rarioribus. Thorax coleopteris vix angustior, latitudine tertia parte longior, lateribus rectis, parallelis, angulis omnibus subrotundatis, leviter convexus, crebre punctatus, linea media longitudinali laevi. Scutellum obsolete punctatum, tenuiter griseo-pubescentis. Elytra thoracis longitudine, subtilius parcusque punctata, nitida, parcus subtiliter griseo-pubescentia. Abdomen omnium confertissime subtilissime punctatum, parum nitidum, pube subtili depressa densius vestitum, supra apice summo piceum, subtus apicem versus rufo-piceum. Pedes crassiusculi, rufo-testacei, coxis omnibus piceis.

Mas abdominis segmento inferiore quarto apicem versus obsolete impresso, quinto apice triangulariter impresso, sexto longitudinaliter impresso, apice utrinque sinuato, medio profunde inciso; femina eodem apice utrinque profundius sinuato, medio fortiter triangulariter producto.

\* 2. *L. elongatum*: Nigrum, nitidum, punctatum, capite suborbiculato, thorace oblongo, elytris thorace longioribus, crebre punctatis, rufis, basi nigris, antennis pedibusque rufis. — Long. 4 lin.

Grav. Micr. 55. 8. Mon. 132. 2. — Latr. Hist. nat. Crust. et Ins. IX. 340. 8. Gen. Crust. et Ins. I. 289. 1. — Guér. Iconogr. regn. an. Ins. t. 9. f. 4. — Nordm. Symb. 141. 1. — Er. Col. March. I. 504. 2.

*Staph. elongatus* Linn. Syst. Nat. I. II. 685. 14. — Goeze Ent. Beitr. I. 719. 14. — De Vill. Ent. I. 415. 14. — Schrank Enum. Ins. Austr. 232. 439. — Payk. Mon. Staph. 25. 17. — Marsh. Ent. Britt. 515. 52.

*Paeder. elongatus* Fab. Syst. Ent. 268. 2. Spec. Ins. I. 339. 3. Mant. Ins. I. 223. 3. Ent. Syst. I. II. 537. 3. Syst. El. II. 609. 3. — Walken. Faun. Paris. I. 275. 3. — Cederh. Faun. Ingr. 122. 374. — Payk. Faun. Suec. III. 428. 3. — Panz. Faun. Germ. 9. 12. Ent. Germ. 362. 4. — Rossi Faun. Etr. Mant. I. 101. 224. Ed. Helw. I. 446. 224. — Oliv. Encycl. meth. VIII. 628. 4.

*Staph. variegatus* Fourcr. Ent. Par. I. 172. 29.

*Lathrob. elongatum* Gyll. Ins. Suec. II. 363. 1. — Sahlb. Ins. Fenn. I. 337. 1. — Zetterst. Faun. Lappon. I. 81. 1. Ins. Lappon. 67. 1. — Munnerh. Brachelytr. 37. 1. — Runde Brachelytr. Hal. 12. 1. (forte)

Habitat in Europa.

Corpus elongatum, lineare, nigrum, nitidum. Antennae capite thora-



ceque breviores, crassiusculae, articulo secundo tertioque subaequalibus, 4—10 paulo minoribus, inter se iterum subaequalibus, obconicis, rufae. Palpi et labrum rufa. Caput thoracis prope latitudine, suborbiculatum, supra leviter convexum, crebre punctatum, punctis in media fronte rarioribus, parcius griseo-pubescentis. Thorax coleopteris paulo angustior, latitudine tertia parte longior, lateribus rectis, parallelis, angulis omnibus rotundatis, supra leviter convexus, crebre punctatus, linea longitudinali laevi, parce subtiliterque griseo-pubescentis. Scutellum obsoletius punctatum. Elytra thorace plerumque paulo longiora, dense punctata, tenuiter griseo-pubescentia, vel rufa, basi nigra, vel nigra, apice rufa. Abdomen subtiliter confertissime punctatum, dense subtiliter griseo-pubescentis, minus nitidum, apice piceum. Pedes testaceo-rufi, coxis piceis.

Mas abdominis segmento inferiore quinto leviter longitudinaliter impresso, sexto per totam longitudinem bicarinato, carinis parallelis, nigro-pubescentibus, interstitio angusto, impresso, apice triangulariter exciso; femina eodem apice rotundato, medio leviter emarginato.

Variat coleopteris thorace vel paulo longioribus, vel eidem longitudine aequalibus: specimina autem etiam coleopteris brevioribus alata occurrunt, in his autem, si mascula sunt, abdominis segmenti inferioris quinti structura minus quam in illis explicata, carinis et impressione longitudinali nec non sinibus apicalibus obsoletioribus, vel nonnunquam fere omnino obsoletis. Cave ne eiusmodi specimina pro specie peculiari habeas.

Obs. Abdominis segmentum sextum maris apud Gyllenh. 1. c. non bicarinatum sed „foveis duabus approximatis parallelis impressum” descriptum, nihilominus non ausim, insectum hocce pro alia quadam specie habere, cum fieri possit, errore quodam eius descriptionem turbatam esse, et iam minus quam olim (Col. March. 1. c.) hac de re dubito. Dom. Sahlberg et Zetterstedt forte bona fide diagnosi Gyllenhaliana usi sunt, et Dom. Runde eandem mutuavisse mihi persuasum, cum huncce auctorem maximam opusculi sui partem e Sahlbergio transscripsisse pateat.

\* 3. *L. fulvipenne*: *Nigrum, nitidum, punctatum, thorace oblongo, elytris thoracis longitudine, crebre punctatis, rufis, basi nigris, pedibus rufis.* — Long. 4 lin.

Gyll. Ins. Suec. II. 365. 2. — Sahlb. Ins. Fenn. I. 338. 3. — Zetterst. Faun. Lapp. I. 82. 2. Ins. Lappon. 67. 2. — Mannerh. Brachelytr. 37. 2. — Boisd. et Lacord. Faun. Ent. Paris. I. 424. 4. — Runde Brachelytr. Hal. 12. 3. — Nordm. Symb. 141. 2. — Er. Col. March. I. 505. 3.

*Staph. fulvipennis* Grav. Mon. 104. 106.

*Lathrob. elongatum* Boisd. et Lacord. Faun. Ent. Paris. I. 423. 3.

Habitat in Germania, Gallia.

*L. elongato* affine, angustius, nigrum, nitidum. Antennae vix capitis thoracisque longitudine, crassiusculae, articulo tertio secundo longiore, 4—10 sensim paulo brevioribus et paulo magis obconicis, rufae. Palpi et labrum rufo-picea. Mandibulae piceae. Caput thorace vix angustius, sub-ovale, lateribus crebrius, fronte parcius punctatum. Thorax coleopteris parum angustior, latitudine sesqui longior, lateribus rectis, parallelis, angulis omnibus rotundatis, parcius griseo-pubescentis, leviter convexus, sat fortiter crebre punctatus, linea media longitudinali laevi, subtiliter griseo-pubescentis. Scutellum obsoletius punctatum. Elytra thoracis vix longitudine, sat crebre aequaliter punctata, rufa, summa basi nigra. Abdomen subtiliter confertim

punctulatum, nitidulum, pube depressa grisea densius vestitum, summo apice piceo. Pedes crassi, rufo-testacei, coxis piceis.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso, femina eodem rotundato, integro.

Variat 1, Alatum, coleopteris thorace paululum longioribus, laete rubris.

2, Apterum, coleopteris thoracis longitudine vel eodem magis minusve brevioribus, rufo-piceis vel castaneis.

Obs. 1. Affine praecedenti, habitu graciliore, capite thoraceque magis oblongis saepius aegre distinguendum, constanter autem differt abdominis segmento inferiore in mare aequali, apice rotundato, medio profunde inciso, in femina apice rotundato, nullo modo emarginato. Utraque varietas abdominis structura convenit, neque praeter alarum defectum et elytrorum longitudinem, ab hoc dependentem, ullam differentiam video.

2. Nomen specificum fortuito idem huic speciei Gravenhorst et Gyllenhal imposuerunt. Paederum fulvipennem Fabr., a Gyllenhalio allatum, p. 295 descripsi.

\* 4. *L. rufipenne*: Nigrum, nitidum, punctatum, capite thoraceque oblongis, elytris thorace longioribus, crebre punctatis, rufis, basi nigris, antennis pedibusque rufis. — Long.  $3\frac{1}{2}$  lin.

Gyll. Ins. Succ. III. 704. 1 — 2. — Sahlb. Ins. Fenn. I. 337. 2. — Mannerh. Brachelytr. 37. 3. — Nordm. Symb. 141. 3. — Er. Col. March. I. 505. 4.

Habitat in Suecia et Germania septentrionali, Dom. Schüppel.

Praecedentibus affine, L. elongato habitu proximum, at dimidio minus, nigrum, nitidum. Antennae capite sesqui longiores, crassiusculae, articulo secundo tertioque subaequalibus, 4 — 10 obconicis, inter se aequalibus, rufae. Palpi rufi. Mandibulae et labrum rufo-picea. Caput thoracis latitudine, leviter oblongum, supra leviter convexum, creberrime sat fortiter profundeque punctatum, fronte media vix laeviore, minus nitidum, tenuiter griseo-pubescent. Thorax coleopteris paulo angustior, latitudine sesqui longior, lateribus rectis, parallelis, angulis anterioribus minus, posterioribus fortius rotundatis, leviter convexus, creberrime fortius punctatus, linea media longitudinali laevissima, tenuiter griseo-pubescent. Scutellum obsolete parum-punctatum. Coleoptera thorace paulo longiora, minus dense subtilius punctata, griseo-pubescentia, nitida, rufa, basi tertia parte nigra. Abdomen confertissime subtiliter punctatum, pube breviuscula depressa grisea densius vestitum, apice summo piceo. Pedes rufi, coxis omnibus nigris.

Mas abdominis segmento quinto sextoque leviter longitudinaliter impressis, hoc apice leviter triangulariter exciso, femina eodem acute rotundato.

Obs. L. punctulatum Mannerh. Brachelytr. 37. 4. Nordm. Symb. 141. 4. forte potius ad hanc speciem pertinet: descriptio enim optime quadrat, si supponere licet, pro L. rufipenni enumeratam quandam L. fulvipennis varietatem esse.

\* 5. *L. multipunctum*: Piceum, nitidum, capite thoraceque oblongo fortiter punctatis, elytris profunde serialim punctatis, rufo-testaceis, basi piceis, pedibus testaceis. — Long. 3 —  $3\frac{1}{2}$  lin.

Grav. Micr. 52. 2. Mon. 131. 11. — Latr. Hist. nat. Crust. et Ins. IX. 338. 2. — Gyll. Ins. Succ. III. 705. 3 — 4. — Mannerh. Brachelytr. 37. 5. — Boisd. et

*Lacord. Faun. Ent. Paris. I. 423. 2. — Nordm. Symb. 141. 5. — Er. Col. March. I. 506. 5.*

*Lathrob. lineare Grav. Micr. 54. 6. Mon. 132. 13. — Latr. Hist. nat. Crust. et Ins. IX. 340. 7.*

*Var. Rufo-testaceum, abdomine piceo, elytris anoque testaceis.*

*Paederus testaceus Oliv. Ent. III. 44. 5. 3. t. 1. f. 6. Encycl. meth. VIII. 628. 5.*

Habitat in Europa.

Statura omnino praecedentium, at minus et gracilius, vel nigro-piceum, vel castaneum, vel rufo-testaceum, abdomine obscuriore. Antennae capite thoraceque longe breviores, minus graciles, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus, rufae. Palpi rufi. Caput thorace paulo latius, subovatum, supra leviter convexum, crebre fortiter punctatum, punctis in fronte media parcioribus. Thorax coleopteris paulo angustior, latitudine plus sesqui longior, lateribus rectis, parallelis, angulis omnibus late rotundatis, leviter convexus, crebre profundeque punctatus, linea media longitudinali laevi, parce subtiliterque nigro-pubescent. Scutellum laeve. Elytra thorace vix longiora, anterieus ultra medium confertim profunde seriatim, dein parcius et minus fortiter punctata, rufa- vel rufo-testacea, basi saepe picea. Abdomen confertim subtilius punctatum, densius subtiliter griseo-pubescent, nigro-piceum, vel piceum, apice rufo-piceum, vel rufo-testaceum. Pedes testacei, coxis omnibus femoribusque anticis rufo-testaceis.

Mas abdominis segmento inferiore quinto fovea lata, antice tuberculis duobus rotundatis approximatis minutis at sat prominulis terminata impresso, sexto profunde inciso; femina eodem apice rotundato.

Variat colore dilutiore capite, thoraceque rufo-castaneis, abdomine piceo, ano elytris-que testaceis, vel totus dilute testaceus, elytris anoque pallidioribus.

\* 6. *L. spadiceum: Rufo-piceum, subtilius punctatum, thorace oblongo, basin versus subangustato, abdominis segmentorum marginibus- ano pedibusque rufo-testaceis. — Long. 4 lin.*

Habitat in Austriae alpinis, Dom. Schüppel, in Bavaria meridionali, Dom. Waltl.

L. elongato magnitudine aequale vel parum maius, paulo magis depressum, rufo-piceum, capite thoraceque nitidis. Antennae capitis thoracisque longitudine, minus tenues, articulo tertio secundo sesqui longiore, 4—10 sensim paulo brevioribus, elongato-obconicis, rufae. Palpi rufi. Caput thorace paulo latius, lateribus rotundatum, antice angustatum, parum convexum, subtilius crebre punctatum, magis fusco-piceum, subtiliter tenuiterque griseo-pubescent. Thorax coleopteris paulo angustior, latitudine sesqui longior, basin versus subangustatus, lateribus rectis, angulis anterioribus late, posterioribus leviter rotundatis, subdepressus, crebre minus fortiter punctatus, linea longitudinali media laevi, parce subtiliterque pube brevioris grisea obsitus. Scutellum parce obsoleteque punctatum. Elytra thorace vix longiora, subtilius aequaliter punctata, griseo-pubescentia, opaca. Abdomen dense subtilissime punctatum, pube brevi depressa grisea densiore obductum, opacum, rufo-piceum, segmentorum marginibus anoque rufo-testaceis. Pedes graciliores cum coxis anterioribus rufo-testacei.



Mas abdominis segmentis inferioribus 3 — 5 leviter at sensim paulo magis impressis, sexto obsoletius canalienlato, apice fortius triangulariter exciso: femina eodem obsolete longitudinaliter bi-impresso, apice obtuse rotundato.

\* 7. *L. angusticolle*: *Subdepressum, nigrum, nitidum, subtilius punctatum, thorace obsolete bisulcato, elytris rufis, basi nigris.* — Long.  $3\frac{1}{2}$  lin.

Boisd. et Lacord. Faun. Ent. Paris. I. 425. 6.

Habitat in Hungaria et Styriae alpinis, Dom. Schüppel, in Bavaria superiore, Dr. Waltl.

Habitu fere *L. quadrati*, at subdepressum, elongatum, gracile, nigrum, nitidum. Antennae capite thoraceque paulo longiores, graciles, articulo tertio secundo sesqui longiore, 4 — 10 aequalibus, sat elongatis, teretibus, rufo-piceae, articulo primo nigro-piceo. Palpi rufo-picei. Caput thorace paulo latius, suborbiculatum, subdepressum, creberrime subtilius punctatum, fronte media laeviore. Thorax coleopteris dimidio angustior, latitudine dimidio longior, basin versus subangustatus, lateribus subrectis, antice cum angulis anterioribus late rotundatis, angulis posterioribus leviter rotundatis, subdepressus, crebre subtilius punctatus, postice obsolete late bisulcatus, sulcis densius punctatis, linea longitudinali media subelevata laevi. Scutellum laeve. Elytra thorace tertia parte longiora, depressa, subtilius dense, basi subseriatim, praeterea aequaliter punctata, rufa, basi parte tertia nigra, subtiliter tenuiterque griseo-pubescentia. Abdomen dense subtiliter punctatum, pube subtili depressa grisea densius obductum. Pedes graciliores, nigri, tarsi rufo-picei.

Mas abdominis segmento inferiore quinto apice late leviter sinuato, sexto bicarinato, apice inter carinas leviter triangulariter exciso; femina eodem apice rotundato.

\* 8. *L. bicolor*: *Elongatum, depressum, subtilius punctatum, testaceum, nitidum, elytrorum basi abdomineque fusciscentibus, thorace angusto, obsolete bisulcato.* — Long. 3 lin.

Habitat in Carniolia (Dahl), Dom. Schüppel.

Habitu omnino praecedentis, at dimidio minus, gracile, depressum, rufo-testaceum, capite, thorace elytrisque nitidis. Antennae capite thoraceque multo longiores, minus tenues, articulo tertio secundo sesqui prope longiore, 4 — 10 sensim paulo brevioribus, elongato-subobconicis. Caput thorace sesqui prope latius, lateribus rotundatum, antrorsum angustatum, depressum, versus latera creberrime, disco inaequaliter parcius punctatum, subtiliter griseo-pubescent, lateribus parce nigro-pilosellum. Thorax coleopteris dimidio angustior, latitudine plus sesqui longior, basin versus subangustatus, lateribus subrectis, antice cum angulis anterioribus late rotundatis, angulis posterioribus leviter rotundatis, depressus, sat crebre subtiliter punctatus, linea media subelevata laevissima, sulco longitudinali obsoletissimo densius punctato utrinque terminata. Scutellum laeve. Elytra thorace vix longiora, depressa, dense subtilius, anterieus subseriatim, praeterea aequaliter punctata, densius subtiliter griseo-pubescentia, basi fusciscentia. Abdomen confertissime subtiliter punctatum, vix nitidum, pube brevi subtili de-

pressa grisea densiore obductum, fusco-testaceum, apice testaceum. Pectus fusco-testaceum. Pedes graciliores, cum coxis rufo-testacei.

Mas abdominis segmento inferiore quinto medio obsolete impresso, apice late leviter emarginato, sexto longitudinaliter impresso, apice leviter triangulariter exciso; femina eodem apice rotundato.

\* 9. *L. picipes*: *Depressum, piceum, parum nitidum, subtilius punctatum, thorace angusto, obsolete bisulcato.* — Long.  $2\frac{1}{2}$  lin.

Habitat in Austria, Dom. Ullrich, in Bavaria superiore, Dom. Walth, Tergesti ad littora maris (Dahl), Dom. Schüppel.

Statura omnino et summa affinitas praecedentis, at dimidio minus, depressum, nigro- vel brunneo-piceum, parum nitidum, subtiliter griseo-pubescent. Antennae capite thoraceque longiores, minus tenues, articulo tertio secundo parum longiore, 4—10 subaequalibus, elongato-subobconicis, dilute rufo-piceae. Palpi et labrum rufa. Caput thorace plus sesqui latius, suborbiculatum, ore angustato, depressum, confertim subtilius punctatum. Thorax coleopteris duplo fere angustior, latitudine plus sesqui longior, basin versus vix angustatus, lateribus subrectis, antice cum angulis anterioribus late rotundatis, angulis posterioribus modice rotundatis, depressus, creberrime subtilius punctatus, medio obsolete bisulcatus, linea media longitudinali subelevata laevissima. Scutellum parce punctulatum. Elytra thorace longiora, creberrime subtilius punctata, brunneo-picea, basi nigro-picea. Abdomen elytris angustius, subtiliter creberrime punctatum, nigrum, apice piceum. Pedes graciliores, picei, tarsis dilutioribus.

Mas abdominis segmento inferiore quarto leviter, quinto fortius longitudinaliter impressis, sexto bicarinato, inter carinas late canaliculato, et apice emarginato; femina eodem apice rotundato.

Variat elytris dilutius rufo-piceis, basi nigricantibus.

Obs. Praecedenti proximum, distinctum autem statura minore, colore, elytris paulo longioribus, capite crebrius punctato.

10. *L. labile*: *Elongatum, subdepressum, nigrum, antennis, tibiis tarsisque rufo-piceis, thorace oblongo, dorso biseriatis punctato, elytris striato-punctatis, apice rufo-testaceis.* — Long.  $1\frac{2}{3}$  lin.

Habitat in Sardinia, Mus. Reg. Taurin.

Habitu omnino praecedentis, at multo minus, L. longulo aequale, elongatum, subdepressum, nigrum, nitidulum. Antennae capitis thoracisque prope longitudine, articulis 3—10 subobconicis, sensim paulo brevioribus, ultimo praecedente sesqui longiore, subcylindrico, apice abrupte acuminato, rufo-piceae. Caput thorace paulo latius, oblongum, latitudine plus sesqui longius, antrorsum subangustatum, supra perparum convexum, sat crebre profundeque punctatum, linea tenui longitudinali media laevi, pube brevi erecta grisea tenuiter vestitum, oris partibus piceis. Thorax coleopteris dimidio prope angustior, latitudine duplo fere longior, lateribus parallelis, angulis anterioribus fortiter rotundatis, parum convexus, dorso biseriatis per totam longitudinem punctatis, seriebus regularibus, crebre punctatis, basin versus sulculi instar leviter impressis, utrinque minus regulariter plagiatis punctatus, interstitio inter series dorsales laevissimo, pube brevi erecta nigra parce

vestitus. Elytra thorace longiora, subdepressa, striis crebre fortiterque punctatis, apice evanescentibus, confertis obsita, pube brevi erecta nigra tenuiter vestita, apice laete rufo-testacea. Abdomen minus confertim subtiliter punctatum, pube depressa grisea tenuiore vestitum, apice piceum. Pedes picei, tibiis tarsisque rufo-piceis, tarsis posticis articulo primo leviter elongato.

\* 11. *L. quadratum*: Nigrum, nitidum, punctatum, thorace subquadrato.

Gyll. Ins. Suec. II. 367. 4. — Sahlb. Ins. Fenn. I. 339. 5. — Zetterst. Faun. Lappon. I. 83. 4. Ins. Lappon. 67. 4. — Er. Col. March. I. 507. 6.

Staph. quadratus Payk. Mon. Staph. 29. 21. — Marsh. Ent. Britt. 517. 58.

Paed. filiformis Payk. Faun. Suec. III. 429. 4.

Var. a: *Elytris immaculatis, antennis pedibusque piceis, his rarius testaceis*. — Long. 3 — 3½ lin.

Lathrob. pilosum Grav. Micr. 56. 9. Mon. 131. 7. — Latr. Hist. nat. Crust. et Ins. IX. 340. 9. — Boisd. et Lacord. Faun. Ent. Paris. I. 426. 9.

Paed. filiformis Fab. Ent. Syst. I. II. 538. 8. Syst. El. II. 609. 8. — Panz. Ent. Germ. 363. 9.

Lathr. quadratum Mannerh. Brachelytr. 38. 9. — Runde Brachelytr. Hal. 12. 5. — Nordm. Symb. 146. 16.

Var. b: *Elytris macula apicali rufo-testacea, antennarum basi apiceque pedibusque testaceis*. — Long. 3 lin.

Lathrob. terminatum Grav. Micr. 55. 7. Mon. 134. 17. — Latr. Hist. nat. Crust. et Ins. IX. 339. 6. — Mannerh. Brachelytr. 38. 10. — Curtis Britt. Ent. XIV t. 650. — Runde Brachelytr. Hal. 13. 6. — Nordm. Symb. 146. 17.

Habitat in Europa.

Corpus nigrum, nitidum, abdomine opaco. Antennae capitis thoracisque longitudine, articulo tertio secundo paulo longiore, 4 — 10 subaequalibus, elongato-subobconicis, piceae aut testaceae, medio obscuriores. Palpi et os rufo-picea. Caput thorace angustius et multo minus, suborbiculatum, subdepressum, confertim subtiliter punctatum, fronte media parcius punctata, subtiliter griseo-pubescent, lateribus nigro-pilosellum. Thorax coleopteris paulo angustior, latitudine haud longior, quadratus, angulis omnibus leviter rotundatis, parum convexus, confertim lateribus subtiliter, dorso fortius punctatus, linea media longitudinali laevi parum distincta, subtiliter tenuiterque griseo-pubescent. Scutellum subtiliter punctulatum. Elytra thorace paulo longiora, subtilius dense punctata, tenuiter pubescentia, in varietate altera immaculata, in altera macula apicali exteriori rotundata rufo-testacea notata. Abdomen breviusculum, coleopteris paulo angustius, apice attenuatum, stylis analibus longius exsertis, omnium subtilissime confertissimeque punctulatum, pube tenuissima grisea sericeum. Pedes crassiusculi, in varietate altera saepius piceo-testacei, in altera semper testacei, coxis in utraque piceis.

Mas abdominis segmento inferiore quarto quintoque obsolete canaliculatis, sexto profunde inciso; femina eodem apice fortiter triangulariter producto.

Obs. Paederum filiformem Fab. ad hanc speciem referendam esse, et ex citato Paykulo, et ex ipsa collectione auctoris patet, quamvis descriptio magis Cryptobio fracticorni congruere videtur, cui autem neque antennae neque pedes picei sunt. Nomen primum nihilominus praevaleat.



- \* 12. *L. scutellare*: *Nigrum, nitidum, punctatum, capite parcius punctato, thorace subquadrato, elytris rufis, circa scutellum nigricantibus, pedibus testaceis.* — Long.  $2\frac{2}{3}$  lin.

Nordm. Symb. 143. 10.

Habitat in Austria.

L. quadrato paulo minus et angustius, L. punctato aequale, nigrum, nitidum. Antennae capitis thoracisque fere longitudine, articulo tertio secundo sesqui prope longiore, 4—10 elongato-subobconicis, subaequalibus, rufo-piceae, medio obscuriores. Palpi rufo-testacei, maxillares articulo tertio piceo. Caput thorace angustius, suborbiculatum, parcius punctatum, media fronte sublaevi, subtiliter griseo-pubescentia, parce nigro-pilosellum. Thorax coleopteris parum angustior, latitudine paululum longior, lateribus rectis, parallelis, angulis anterioribus leviter, posterioribus fortius rotundatis, leviter convexus, crebre punctatus, linea media longitudinali laevi, tenuiter subtiliterque griseo-pubescentia. Scutellum sublaeve. Elytra thorace vix longiora, sat crebre punctata, laete rufa, circa scutellum nigricantia, subtiliter griseo-pubescentia. Abdomen subtilissime confertim punctatum, pube densiore subtili sericea grisea vestitum, opacum. Pedes testacei, coxis omnibus nigris.

Mas abdominis segmento inferiore sexto leviter longitudinaliter impresso, apice leviter emarginato; femina eodem apice acute rotundato.

13. *L. angustatum*: *Nigrum, nitidum, punctatum, capite confertissime subtiliter punctato, thorace subquadrato, elytris rufis, basi nigris, pedibus testaceis.* — Long.  $2\frac{2}{3}$  lin.

Boisd. et Lacord. Faun. Ent. Paris. I. 424. 5.

Habitat Lutetiae, Mus. Dom. Aubé.

Statura fere L. pilosi, at paululum angustius, et minoribus eius individuis aequale, nigrum, nitidum. Antennae capitis thoracisque longitudine, rufo-testaceae. Os rufo-testaceum. Caput thorace paululum angustius, subovatum, confertissime subtiliter punctatum, fronte media spatio angusto longitudinali laevi, tenuiter griseo-pubescentia. Thorax coleopterorum prope latitudine, latitudine paululum longior, leviter convexus, crebre punctatus, linea media longitudinali laevi. Scutellum sublaeve. Elytra thorace paululum longiora, sat crebre profundeque punctata, tenuiter griseo-pubescentia, rufa, nitida, basi (parte elytrorum quarta) nigra. Abdomen confertim subtilissime punctatum, minus nitidum, densius griseo-pubescentia. Pedes testacei, coxis anterioribus concoloribus, licet saturatioribus.

Obs. L. scutellari adeo simile, ut praeter capitis puncturam vix distinguatur.

14. *L. pectorale*: *Subdepressum, nigrum, ore, antennis, elytris, pectore anoque rufis, pedibus testaceis, capite parcius, thorace utrinque crebrius punctato.* — Long. 3 lin.

Habitat in Americae meridionalis ins. St. Thomae, Dom. Moritz.

Statura omnino L. filiformis, at paulo robustius et magis depressum. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus, leviter obconicis, rufo-testaceae.

Palpi rufo-testacei. Os rufum. Caput thoracis latitudine, subquadratum, depressiusculum, nigro-piceum, nitidum, lateribus crebrius punctatum, densius pilosum, fronte laevi, punctis raris sparsis notata, subtus rufo-piceum. Thorax coleopteris parum angustior, latitudine paulo longior, basin versus subangustatus, lateribus rectis, angulis anterioribus obtusiusculis, posterioribus rotundatis, subdepressus, lateribus crebre punctatus, plaga media longitudinali laevi, supra nigro-piceus, nitidus, subtiliter nigro-puberulus, subtus piceo-rufus. Scutellum laeve, piceum. Elytra thorace paulo longiora, crebrius minus fortiter, anterieus subseriatim punctata, rufa, subnitida, griseo-pubescentia. Abdomen subtilissime punctulatum, griseo-pubescentis, nigro-piceum, parum nitidum, segmento ultimo toto, reliquis margine rufis. Pectus rufum. Pedes breves, crassiusculi, testacei, coxis rufis.

Mas abdominis segmento inferiore quinto apice late leviter emarginato, sexto profunde triangulariter exciso. Femina latet.

\* 15. *L. Lusitanicum*: *Subdepressum, nigrum, capite thoraceque oblongo, dorso biserialim punctato, parce punctatis, pernitidis, elytris subseriatim punctatis anoque rufis, pedibus testaceis.* — Long.  $3\frac{1}{2}$  lin.

*Lathrob. longiusculum Lusitanicum* Grav. Micr. 181. 4.

Habitat in Lusitania. —

L. quadrato aequale, at magis depressum, nigrum, opacum, capite thoraceque pernitidis. Antennae capite thoraceque paulo longiores, apice subattenuatae, graciliores, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus, teretibus, penultimis leviter obconicis, rufo-testaceae. Palpi et labrum rufo-testacea. Caput thorace angustius et duplo fere minus, subovatum, depressiusculum, punctis raris sparsis impressum, lateribus longe nigro-pilosum. Thorax coleopteris paulo angustior, latitudine sesqui prope longior, basin versus vix angustatus, angulis anterioribus parum, posterioribus leviter rotundatis, subdepressus, dorso biserialim, lateribus disperse parceque punctatus. Scutellum laeve. Elytra thorace paulo longiora, minus dense subtilius, dorso biserialim, lateribus apiceque disperse punctata, rufa, nitida, parcius griseo-pubescentia. Abdomen minus elongatum, apice attenuatum, subtiliter confertissime punctatum, pube depressa grisea densius vestitum, opacum, segmentis quatuor primis margine summo, ultimis duobus totis rufis. Pedes cum coxis anterioribus rufo-testacei.

Mas abdominis segmento inferiore sexto profunde inciso. Femina latet.

Obs. A *L. longiusculo* Americano abunde distinctum.

\* 16. *L. longiusculum*: *Subdepressum, nigrum, nitidum, capite thoraceque oblongo, dorso biserialim punctato, disperse punctatis, elytris serialim punctatis pedibusque rufis.* — Long. 5 lin.

Grav. Micr. 181. 4. Mon. 131. 9. — Nordm. Symb. 144. 14.

Var. Corpore toto rufo-testaceo.

*Lathr. politum* Grav. Micr. 180. 3. Mon. 131. 10.

*Lathr. castaneum* Grav. Micr. 52. 1. Mon. 131. 10. — Latr. Hist. nat. Crust. et Ins. IX. 338. 1.

Habitat in America septentrionali.

*L. punctato* aequale, at paulo magis depressum, nigro-piceum, nitidum, capite thoraceque nitidissimis. Antennae vix capitibus thoracisque longitudine, articulo tertio secundo parum longiore, 4—10 sensim paulo brevioribus, rufo-testaceae. Palpi rufo-testacei. Os rufo-piceum. Caput thoracis prope latitudine, suborbiculatum, subdepressum, punctis raris sparsis impressum, lateribus parce pilosellum. Thorax coleopteris parum angustior, latitudine paulo longior, lateribus rectis, parallelis, angulis anterioribus leviter, posterioribus fortiter rotundatis, subdepressus, dorso irregulariter biseriatim, lateribus disperse parcius punctatus, glaber. Scutellum piceum, laeve. Elytra thoracis longitudine, parcius leviterque dorso seriatim, lateribus et apice disperse punctata, parce griseo-pubescentia, nitida, rufa, basi picescentia. Abdomen minus dense subtiliter punctulatum, parce subtiliterque griseo-pubescentia, nitidum, segmentis omnibus margine apicali rufo-piceo. Pedes cum coxis rufo-testacei.

Mas abdominis segmento inferiore sexto apicem versus triangulariter impresso, apice intra impressionem triangulariter exciso; femina eodem apice triangulariter producto.

Obs. *L. politum* Grav. nonnisi colore corporis rufo-testaceo differt, forte immaturum. — *L. castaneum* haud distinctum videtur, forte errore quodam inter *Brunsvicensia* cum *L. badio* allatum.

17. *L. discolor*: *Supra nigro-piceum, nitidum, corpore subtus, antennis pedibusque rufis, capite thoraceque oblongo, dorso biseriatim punctato, disperse punctatis, elytris subseriatim punctatis.* — Long. 3 lin.

Habitat in Americae meridionalis ins. St. Thomae, Dom. Moritz.

Statura omnino *L. longiusculi*. Antennae capite thoraceque paulo breviores, crassiusculae, articulo tertio secundo paulo longiore, 4—10 sensim paulo brevioribus, leviter obconicis, rufo-testaceae. Palpi rufo-testacei. Os rufo-piceum. Caput thoracis latitudine, subdepressum, disperse sat profunde punctatum, punctis lateribus et basi crebrioribus, in fronte raris, supra nigro-piceum, nitidum, parce nigro-pilosellum, subtus piceo-rufum. Thorax coleopteris paululum angustior, latitudine paulo longior, basin versus subangustatus, lateribus rectis, angulis anterioribus obtusis, posterioribus leviter rotundatis, parum convexus, crebrius punctatus, punctis lateribus vagis, dorso in series duas longitudinales irregulares, saepius duplicatas digestis, spatio interiecto laevissimo, supra nigro-piceus, nitidus, glaber, subtus rufo-piceus. Scutellum piceum, laeve. Elytra thorace vix longiora, minus dense, dorso subseriatim, lateribus et apice vage punctata, parce griseo-pubescentia, nigro-picea, nitida, margine apicali obsolete rufescente. Abdomen subtiliter punctulatum, pube densiore depressa grisea vestitum, minus nitidum, supra piceum, apice segmentorumque singulorum marginibus, rufis, subtus cum pectore rufo-piceum. Pedes rufo-testacei.

Mas abdominis segmento inferiore sexto profunde inciso, quinto foveola oblonga, ovata, marginibus carinatis inclusa, notato. Femina latet.

- \* 18. *L. dimidiatum*: *Lineare, nigrum, nitidum, elytris apice thoraceque rufis, antennis pedibusque testaceis, capite subovato, parce punctato, thorace oblongo, dorso biseriatim punctato.* — Long. 2 lin.

Say. *Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV.* 455. 5.



Habitat in Carolina meridionali, Dom. Zimmermann.

Statura et summa affinitas sequentis. Antennae capite thoraceque paulo breviores, articulo tertio secundo vix longiore, 4 — 10 sensim paulo brevioribus, rufo-testaceae, medio picescentes. Os rufo-testaceum. Caput thoracis latitudine, subovatum, depressiusculum, parce punctatum, fronte media sublaevi, parce pilosum, nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine paulo longior, basin versus vix angustatus, lateribus subrectis, angulis posterioribus rotundatis, anterioribus obtusiusculis, subdepressus, dorso punctorum seriebus duabus, e punctis 7 vel 8 compositis, posterius subimpressis, lateribus punctis parvis minus subtilibus notatus, rufus, nitidus. Scutellum laeve, nigro-piceum. Elytra thorace paulo longiora, stria suturali impressa et dorso seriebus punctorum duabus sat regularibus, postice evanescentibus notata, lateribus parce subtiliter punctata, nitida, nigra, apice dimidiato rufa. Abdomen subtiliter punctulatum et subtiliter griseo-pubescent, nigrum, nitidum, segmentis 4 primis apice tenuiter testaceo-marginatis, quinto apice, sexto toto testaceo. Pectus nigrum. Pedes pallide testacei.

Abdominis segmentum inferius sextum feminae apice rotundatum. Mas latet.

Obs. Differt a *L. nitido* capite minus quadrato, thoracis seriebus dorsalibus e punctis paucis compositis, punctis quoque lateralibus parrioribus.

\* 19. *L. nitidum*: Lineare, nigrum, nitidum, thorace, elytris apice, antennis pedibusque rufo-testaceis, capite subquadrato, parce punctato, thorace oblongo, dorso biserialim punctato. — Long.  $2\frac{1}{3}$  lin.

Habitat in Columbia, Dom. Moritz.

Habitu fere *Lithocharidis melanocephalae*, at duplo maius et huius generis. Antennae capitis thoracisque prope longitudine, crassiusculae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus, leviter obconicis, rufo-testaceae. Os rufo-testaceum, mandibulis piceis. Caput thoracis latitudine, subquadratum, supra depressiusculum, subtilius parce punctatum, fronte sublaevi, lateribus subtiliter pubescens, nigrum, nitidum. Thorax coleopteris paulo angustior, latitudine paulo longior, basin versus subangustatus, lateribus rectis, angulis anterioribus leviter, posterioribus fortius rotundatis, dorso punctorum seriebus duabus minus regularibus multipunctatis, utrinque punctorum plaga irregulari notatus, rufus, nitidus. Scutellum laeve. Elytra thorace paulo longiora, stria suturali impressa, seriebus nonnullis punctorum subtilium obsoletorumque, ultra medium evanescentibus notata, glabra, nitida, nigra, apice late laete rufo-testacea. Abdomen dense subtiliter punctatum, densius subtiliter cinereo-pubescent, nigrum, segmentorum marginibus apicalibus rufo-piceis. Pectus rufo-piceum. Pedes dilute testacei.

Mas abdominis segmento inferiore quinto apice leviter emarginato, sexto leviter triangulariter exciso; femina eodem apice rotundato.

Variat thorace piceo.

\* 20. *L. collare*: Lineare, nigrum, nitidum, antennis, pedibus anoque rufo-testaceis, thorace rufo, elytrisque subserialim punctatis. — Long.  $2\frac{2}{3}$  lin.

Habitat in America septentrionali, Dom. Prof. Germar.

Habitus et summa affinitas praecedentis, at sesqui maius. Antennae capitis thoracisque prope longitudine, minus tenues, articulo tertio secundo paulo longiore, sequentibus sensim paulo brevioribus, leviter obconicis, rufo-testaceae. Os rufo-testaceum. Caput thorace paulo angustius, suborbiculatum, perparum convexum, parce sparsim punctatum, lateribus nigro-pilosellum, nigerrimum, pernitidum. Thorax coleopteris paulo angustior, latitudine paulo longior, lateribus rectis, angulis anterioribus leviter, posterioribus fortius rotundatis, leviter convexus, lateribus parcius disperse, dorso crebrius subseriatim punctatis, spatio medio longitudinali laevi, utrinque punctorum numerosorum serie sat regulari terminato, laete rufus, pernitidus, pilis parvis minutis erectis e punctis oriundis vestitus. Scutellum nigro-piceum, laeve. Elytra thorace tertia parte longiora, dorso sat regulariter seriatim punctata, seriebus 5, ante apicem evanescentibus, interstitiis laevissimis, lateribus et apice parce subtiliterque disperse punctata, nigra, nitida, callo humerali et margine apicali picescentibus. Abdomen subtilissime punctulatum, pube brevi depressa grisea densius vestitum, nigrum, segmentis 4 primis margine, quinto apice, sexto toto rufo testaceis. Pectus nigrum. Pedes cum coxis testacei.

Mas latet. Femina abdominis segmento inferiore sexto apice leviter producto, acute rotundato.

Variat elytris rufo-piceis, disco picescentibus.

\* 21. *L. punctatum*: *Lineare, nigrum, nitidum, antennis pedibusque rufo-piceis, capite parce punctato, thorace subquadrato elytrisque thoracis longitudine crebrius punctatis. — Long. 3 lin.*

*Zetterst. Faun. Lappon. I. 84. 5. Ins. Lappon. 68. 5. — Sahlb. Ins. Fenn. I. 340. 6. — Nordm. Symb. 141. 7. — Er. Col. March. I. 508. 7.*

*Lathrob. lineare* Boisd. et Lacord. *Faun. Ent. Paris. I. 426. 8.*

Habitat in Germania.

*L. filiformi* affine, at robustius, nigrum, capite, thorace elytrisque nitidis. Antennae capitis thoracisque fere longitudine, articulo tertio secundo paulo longiore, 4—10 sensim paululum brevioribus et distinctius obconicis, rufo-testaceae. Palpi testacei. Os rufo-piceum. Caput thoracis prope latitudine, suborbiculatum, supra parum convexum, parce disperse punctatum, fronte media laeviore, parce griseo-puberulum, lateribus nigro-pilosellum. Thorax coleopterorum fere latitudine, latitudine parum (quadrante) longior, subquadratus, lateribus rectis, parallelis, angulis anterioribus leviter, posterioribus fortius rotundatis, parum convexus, crebrius punctatus, spatio angusto dorsali longitudinali laevi, tenuiter subtiliterque griseo-pubescentis. Scutellum laeve. Elytra thoracis longitudine, crebrius punctata, tenuiter griseo-pubescentia. Abdomen subtiliter confertissime punctatum, densius griseo-pubescentis, apice piceum, segmentis omnibus tenuiter piceo-marginatis. Pedes crassiusculi, rufo-picei.

Mas abdominis segmento inferiore sexto leviter canaliculato, apice leviter exciso; femina eodem apice producto, rotundato.

- \* 22. *L. filiforme*: *Lineare, nigrum, nitidum, antennis pedibusque piceis, capite, thorace oblongo elytrisque crebre punctatis, his thorace brevioribus.* — Long.  $2\frac{2}{3}$  lin.

Grav. Mon. 134. — Nordm. Symb. 144. 13. — Er. Col. March. I. 508. 8.

*Lathrob. lineare* Gyll. Ins. Suec. II. 370. 6. — Sahlb. Ins. Fenn. I. 341. 8. — Zetterst. Faun. Lappon. I. 85. 6. Ins. Lappon. 68. 6. — Mannerh. Brachelytr. 37. 7. — Nordm. Symb. 142. 9.

Habitat in Europa.

Elongatum, lineare, tenue, nigrum, nitidum. Antennae capite thoraceque breviores, articulo secundo tertioque aequalibus, 4 — 10 sensim paulo brevioribus, subobconicis, piceo-testaceae. Palpi dilute piceo-testacei. Os piceum. Caput thoracis latitudine, subovatum, supra leviter convexum, sat crebre punctatum, fronte media laeviore, subtiliter pubescens. Thorax coleopteris vix angustior, latitudine plus sesqui fere longior, lateribus rectis, parallelis, angulis omnibus late rotundatis, sat fortiter crebre punctatus, linea media longitudinali laevi, subtiliter pubescens. Scutellum laeve. Elytra thorace paulo breviora, subtiliter crebre punctata, punctis passim subseriatis, tenuiter griseo-pubescens. Abdomen subtiliter dense punctatum, pube subtiliore depressa grisea densius vestitum. Pedes picei vel nigri, tarsi ferrugineis.

Mas abdominis segmento inferiore quinto obsolete longitudinaliter impresso, sexto apice anguste triangulariter impresso et exciso; femina eodem apice producto, rotundato.

- \* 23. *L. pallidum*: *Lineare, testaceum, capite thoraceque oblongo parcius, elytris densius punctatis, nitidis, abdomine supra fusciscente.* — Long.  $2\frac{1}{2}$  lin.

Nordm. Symb. 142. 8. — Er. Col. March. I. 509. 9.

Habitat in Germania, Berolini, ipse, in Austria et Illyria, Dom. Schüppel.

Statura omnino *L. filiformis*, at minus elongatum, fulvo-testaceum, capite, thorace elytrisque nitidis. Antennae capite thoraceque paulo breviores, articulo tertio secundo parum brevior, 4 — 10 sensim minoribus, leviter obconicis. Caput thoracis latitudine, suborbiculatum, ore angustato, parce disperse punctatum, lateribus parce pilosellum. Thorax coleopterorum fere latitudine, latitudine tertia parte longior, parum convexus, crebrius punctatus, spatio medio longitudinali angusto laevi, glaber. Scutellum laeve. Elytra thorace paululum breviora, crebre minus profunde punctata, tenuiter pubescens. Abdomen subtilissime confertissimeque punctatum, pube densa subtili grisea sericeum, vix nitidum, segmentis 5 anterioribus dorso nigricantibus. Pedes breves, validi, femoribus anticis fortius incrassatis.

Mas abdominis segmento inferiore quarto quintoque longitudinaliter leviter impressis, sexto profunde inciso; femina eodem apice producto, apice summo subtruncato.

- \* 24. *L. dividuum*: *Lineare, depressum, piceum, capite thoraceque oblongis, dense punctatis, nitidis, elytris testaceis, basi piceis.* — Long. 2 lin.

Habitat in Sardinia, Dom. Prof. Gené.

*L. longulo* paululum maius, at multo magis depressum. Antennae capite thoraceque vix breviores, crassiusculae, articulo tertio secundo paulo



longiore, 4 — 10 sensim paulo brevioribus, rufo-testaceae, articulo ultimo apice pallidiore. Palpi testacei. Caput thoracis latitudine, oblongum, antrosum leviter angustatum, depressum, profunde sat crebre punctatum, linea media longitudinali laevi, subtiliter griseo-pubescens, testaceo-piceum, subnitidum. Thorax coleopteris dimidio angustior, latitudine antica plus sesqui longior, basin versus leviter angustatus, lateribus rectis, angulis anterioribus leviter, posterioribus fortius rotundatis, subdepressus, sat crebre punctatus, linea media longitudinali laevi, subtiliter tenuiterque griseo-pubescens, testaceo-piceus, nitidus. Scutellum piceum, laeve. Elytra thorace paulo longiora, depressa, parce subtilissime obsoleteque punctata, parce pubescentia, dilute testacea, nitida, basi picea. Abdomen confertissime subtilissimeque punctatum, pube brevissima subtili grisea dense obductum, opacum, nigrum, segmentis 4 primis apice subtiliter piceo-marginatis, quinto apice, sexto toto testaceo-piceis. Pectus piceum. Pedes breves, crassiusculi, picei, tarsis dilutioribus.

Mas abdominis segmento inferiore quinto longitudinaliter impresso, apicem versus bicarinato, interstitio apice carinula brevissima notato, carinulis omnibus apice prominulis, apice tridentato; sexto apice profunde triangulariter exciso; femina eodem apice rotundato.

- \* 25. *L. dilutum*: Lineare, piceo-testaceum, nitidum, capite thoraceque oblongo crebre punctatis, elytris thorace brevioribus, distincte punctatis. — Long. 2 lin.

*Er. Col. March. I. 509. 10.*

Habitat in Germania septentrionali.

*L.* longulo affine, at duplo maius, piceo-testaceum, nitidum. Antennae capite thoraceque paulo breviores, crassiusculae, articulo secundo tertioque subaequalibus, 4 — 10 sensim paulo brevioribus, subobconicis. Caput thoracis latitudine, suborbiculatum, ore angustato, producto, parum convexum, crebre punctatum, media fronte laeviore. Thorax coleopteris paulo angustior, latitudine sesqui longior, lateribus rectis, parallelis, angulis anterioribus late, posterioribus modice rotundatis, subdepressus, crebre punctatus, spatio angusto longitudinali medio laevi. Scutellum laeve. Elytra thorace paulo breviora, crebre fortius punctata. Abdomen subtiliter confertissime punctatum, subtiliter densius griseo-pubescens. Pedes breves, crassiusculi, pallide testacei.

Mas latet. Femina abdominis segmento inferiore sexto apice producto, rotundato.

- \* 26. *L. longulum*: Lineare, nigro-piceum, nitidum, capite thoraceque oblongo distincte, elytris obsolete punctatis, antennis pedibusque rufis. — Long.  $1\frac{2}{3}$  lin.

*Grav. Micr. 53. 4. Mon. 134. 15. — Latr. Hist. nat. Crust. et Ins. IX. 339. 4. — Gyll. Ins. Suec. II. 371. 7. IV. 483. 7. — Nordm. Symb. 143. 12. — Er. Col. March. I. 510. 11.*

*Lathrobium minutum* Sahlb. *Ins. Fenn. I. 341. 9. — Mannerh. Brachelytr. 38. 8. — Runde Brachelytr. Hal. 12. 4. — Nordm. Symb. 143. 11.*

Habitat in Europa.

Minutum, lineare, tenue, apterum, nigrum, vel nigro-piceum, nitidum. Antennae crassiusculae, capitis thoracisque longitudine, articulo tertio secundo aequali, 4 — 10 sensim paulo brevioribus, rufae. Os rufum. Caput thorace paulo latius, subovatum, ore angustato, producto, subdepressum, parcius profundiusque punctatum, subtiliter griseo-pubescent. Thorax coleopteris parum angustior, latitudine plus sesqui longior, lateribus rectis, parallelis, angulis omnibus late rotundatis, crebre punctatus, spatio angusto longitudinali medio laevi, subtiliter griseo-pubescent. Scutellum laeve. Elytra thorace tertia parte breviora, minus dense obsolete punctata, subtiliter griseo-pubescentia. Abdomen subtiliter dense punctatum, pube subtili depressa grisea densius vestitum. Pedes breviores, crassiusculi, cum coxis anterioribus rufo-testacei vel rufi.

Mas abdominis segmento inferiore sexto obsolete canaliculato, bisinuato; femina eodem apice producto, rotundato.

Obs. Nomine *L. minuti* ill. Com. Dejean *L. longulum* Gyll. a Gravenhorstiano distinxit, equidem neque e descriptione Gyllenhaliana, neque in specimine Suecico a Dom. Prof. Zetterstedt mihi communicato ullam differentiam video.

\* 27. *L. scabricolle*: *Nigrum, antennis pedibusque rufis, capite orbiculato thoraceque suborbiculato confertissime punctato-subrugulosis. — Long. 3¼ lin.*

Habitat in Hercynia, Austria.

Corpus apterum, habitu Stilicium simulans, nigrum, opacum. Antennae capite thoraceque paulo breviores, articulo tertio secundo sesqui longiore, sequentibus sensim paulo brevioribus, subobconicis, rufo-ferrugineae. Palpi rufi, maxillares articulo tertio fusciscent. Caput thorace paulo latius, orbiculatum, leviter convexum, confertissime punctato-subrugosum, subtiliter griseo-pubescent. Thorax coleopteris paulo latior, latitudine paulo longior, antice posticeque rotundatus, subovatus, leviter convexus, confertissime subtilius punctato-subrugulosus, subtiliter griseo-pubescent. Elytra thorace haud longiora, creberrime punctata, cinereo-pubescentia, obscure piceo-subaenea, nitidula. Abdomen subclavatum, confertim subtiliter punctulatum, cinereo-pubescent, subtile segmentis singulis piceo-marginatis. Pedes longiores, crassiusculi, rufi, coxis piceis.

Mas abdominis segmento inferiore quinto apice late subemarginato, sexto late obsolete canaliculato, apice leviter triangulariter exciso; femina eodem apice producto, acute rotundato.

Species mihi invisae, forte huius generis.

28. *L. armatum*: *Nigrum, punctatum, thorace oblongo-quadrato, elytris antennisque piceis, pedibus rufis. — Long. 3½ lin.*

Say Transact. Amer. Philoph. Soc. Philadelph. Nov. Ser. IV. 453. 1.

Habitat in Indiana Americae septentrionalis.

Corpus nigrum, punctatum. Antennae piceae, articulo primo obconico, secundo tertioque coniunctis haud longiore, his aequalibus, 4 — 10 moniliformibus, ultimo ovato, acuto. Palpi dilute picei. Thorax oblongo-quadratus, angulis rotundatis, absque linea media longitudinali laevi. Elytra nigro-

picea, minus profunde punctata, stria suturali impressa. Pedes rufi, femoribus robustis, praecipue anticis, his infra prope apicem angulatis, tibiis leviter dilatatis compressisque, apice oblique truncatis, anticis intus obtuse sinuatis, tarsis articulo primo secundo haud longiore, quarto subtus sublobato.

Obs. *L. brunnipedi* affine videtur.

29. *L. puncticolle*: *Nigrum, thorace fere toto punctato, elytris thorace longioribus, antennis (?) pedibusque castaneis. — Long. 5 lin.*

*Kirby Faun. Boreali-Amer.* 86. 2.

Habitat in America boreali sub lat. 54°.

Corpus nigrum, subnitidum, thorace excepto pubescens. Caput obovatum, subtiliter dense punctatum, antennis mutilatis basi, mandibulis palpisque castaneis. Thorax oblongo-quadrangulus, angulis rotundatis, minus confertim punctatus, spatio ullo longitudinali laevi vix relicto. Elytra thorace longiora, dense punctata, obscure castanea. Pedes brunnei. — *L. brunnipedi* affine, at elytris longioribus, pedibus obscurioribus, femoribus anticis minus fortibus, muticis. (Kirby).

Obs. Tibiis anticis basi dentatis et tarsis anticis dilatatis indubie ad *Lathrobia* genuina referendum.

30. *L. brevicorne*: *Depressum, castaneum, capite, thorace elytrisque subtiliter confertissime punctatis, antennis brevibus, apicem versus incrassatis. — Long. 3 lin.*

*Latr. Hist. nat. Crust. et Ins.* IX. 342, 14.

Habitat in Gallia.

Depressum, obscure castaneum, subopacum, exceptis thorace elytrisque. Antennae rufescentes, articulis brevibus, apicem versus sensim crassioribus. Palpi rufescentes. Caput elongatum, subtilissime punctatum, subrugulosum, linea media longitudinali (laevi?). Thorax elongatus, punctatissimus, aequalis. Elytra subtilissime punctata, leviter pubescentia. Pedes breves, rufopicei, femoribus obscurioribus. (Latr.)

Obs. De genere dubito, forte *Lithocharidi* adscribendum.

## Scopaenus Er.

*Labrum denticulatum.*

*Palpi maxillares articulo ultimo acuto, distincto.*

*Ligula cornea, tricuspidata.*

*Antennae rectae.*

*Tarsi articulo quarto simplice.*

Corpus parvulum, elongatum, alatum. Caput a thorace remotum, pedunculo brevi tenui thoracis apici affixum, porrectum, thoracis latitudine, oculis parvis, rotundatis, modice prominulis. Labrum transversum, breve, apice denticulatum, denticulis intermediis distinctioribus, interstitio profundius triangulariter inciso. Mandibulae leviter falcatae, acutae, medio fortiter acuteque tridentatae. Maxillae malis breviusculis, coriaceis, inferiore interne, exteriore apice introrsum vergente barbatis. Palpi maxillares articulo primo brevi, secundo tertioque longitudine subaequalibus, illo leviter clavato, hoc



fortiter incrassato, quarto minuto, tenui, subulato. Labium mento transversum, ligula cornea, apice tricuspidata, dentibus aequalibus, acutis, paraglossis linearibus, membranaceis, interne ciliatis. Palpi labiales articulo primo minuto, cylindrico, secundo maiusculo, crassiusculo, subovali, tertio minuto, tenui, acuminato. Antennae filiformes, rectae, articulo primo modice elongato, secundo tertioque obconicis, ultimo ovato, apice subacuminato. Thorax oblongus, apice magis minusve attenuatus. Scutellum subsemiorbiculare. Elytra apice truncata. Abdomen lineare, segmento septimo vix prominulo, stylis analibus haud distinctis. Pedes breviores, validiusculi, antici leviter incrassati, tibiis anticis interne subsinuatis et ante medium obtuse dentatis, posterioribus apice oblique truncatis, tarsis omnibus simplicibus, posterioribus articulis 4 primis sensim paulo brevioribus, primo subelongato, quinto his coniunctis multo brevioribus.

#### Observationes.

1. Genus intermedium inter *Lathrobium*, *Lithocharidem* et *Stilicum*, corporis forma, imprimis capite remoto et thorace apice attenuato cum *Stilico* convenit, discedit autem praeter labrum minutum et palporum maxillarium articulum quartum distinctum corporis punctura subtilissima et pedibus breviusculis, minus gracilibus. A *Lathrobio*, quocum pedum structura omnino congruit, discedit tarsis anticis simplicibus et labro denticulato. Ad *Lithocharidem* proxime aggreditur, pedibus autem praecipue anticis validioribus, labro minore, lateribus haud membranaceo differt. Praeterea ab omnibus huius familiae generibus ligula cornea, tricuspidata distinctum.
2. Sexus differentia in abdominis segmentis inferioribus ultimis quaerenda.
3. Victus in locis humidis.

\* 1. *Sc. laevigatus*: Subtilissime punctulatus, piceus, subnitidus, antennis pedibusque testaceis, capite orbiculato, thorace subovato, apice attenuato, basi foveola gemina impresso, elytris thorace longioribus. — Long.  $1\frac{1}{2}$  lin.

*Paederus laevigatus* Gyll. Ins. Suec. IV. 483. 4 — 5.

*Rugilus laevigatus* Mannerh. Brachelytr. 40. 2.

*Lathrobium laevigatum* Er. Col. March. I. 510. 12.

Habitat in Europa.

Minutus, nigro-piceus, nitidulus, pube subtilissima brevissimaque cinerea sericans. Antennae capite thoraceque longe breviores, crassiusculae, articulo tertio secundo paulo minore, 3 — 10 sensim paulo brevioribus, penultimis crassitiae vix longioribus, ultimo subovato, apice subito acuminato, rufo-testaceae. Palpi testacei, maxillares articulo tertio crasso, fusco. Os rufo-testaceum. Labrum apice 4-denticulatum, denticulis exterioribus obtusis, intermediis acutis, interstitio triangulariter exciso. Caput thorace paulo latius, suborbiculatum, modice convexum, confertissime subtilissimeque punctatum. Thorax coleopteris dimidio angustior, latitudine sesqui longior, subovatus, basi rotundatus, apice attenuatus, lateribus cum angulis anterioribus leviter rotundatus, basin versus subangustatus, modice convexus, confertissime subtilissimeque punctatus, basi foveolis duabus approximatis rotundatis impressus. Elytra thorace tertia parte longiora, confertim subtilissime punctata, nigro-picea, margine apicali dilutiore. Abdomen subtilissime confertissimeque punctulatum, apice piceum, infra apice rufum, segmentis 4 an-

terioribus apice testaceo-marginatis. Caput et thorax infra rufescunt. Pedes rufo-testacei.

Mas abdominis segmento inferiore sexto apice profunde exciso, excisura utrinque dente duplici, altero deorsum, altero longiore retrorsum spectante, terminata; femina eodem leviter triangulariter producto.

Variat immaturus colore testaceo.

- \* 2. *Sc. didymus*: *Subtilissime punctulatus, niger, subnitidus, antennis pedibusque piceis, thorace oblongo, apice attenuato, basi foveola gemina impresso, elytris thorace longioribus.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Sardinia, Dom. Prof. Gené.

Præcedente magis, sequente minus gracilis, niger, subnitidus, pube brevissima subtilissimaque grisea sericans. Antennae capite sesqui longiores, articulo tertio secundo paulo minore, 3 — 10 sensim brevioribus, ultimo apice subito acuminato, ferrugineae, basi piceae. Palpi rufi, maxillares articulo tertio nigro. Os piceum. Caput thorace paulo latius, oblongum, basi truncatum, angulis posterioribus rotundatis, antrorsum subangustatum, supra leviter convexum, omnium subtilissime confertissime punctatum. Thorax coleopteris dimidio prope angustior, latitudine tertia parte longior, basin versus subangustatus, apice attenuatus, angulis anterioribus obtusis, lateribus subrectis, angulis posterioribus rotundatis, minus convexus, confertissime subtilissimeque punctatus, basi foveolis duabus, carinula brevi distinctis, obsolete impressus, apice linea subtilissima longitudinali medium versus evanescente exaratus. Elytra thorace paulo longiora, confertissime subtilissimeque punctata. Abdomen confertissime subtilissimeque punctatum, densius pubescens, infra segmentis 4 primis margine apicali tenui testaceo. Pedes picei.

Mas abdominis segmento inferiore quinto apice late levissime emarginato, sexto fortiter triangulariter exciso.

- \* 3. *Sc. minutus*: *Elongatus, subdepressus, subtilissime punctatus, piceus, antennis pedibusque testaceis, capite oblongo, thorace oblongo, apice attenuato, elytris thoracis longitudine.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Germania, Gallia.

L. minimo simillimus, at paululum maior, elongatus, linearis, subdepressus, nigro-piceus, subnitidus, pube subtili brevissima grisea sericans. Antennae capite sesqui longiores, articulo tertio secundo subaequali, 3 — 10 sensim brevioribus, ultimo ovato, apice acuminato, testaceae. Palpi testacei. Labrum apice sexdenticulatum, dentibus lateralibus obtusis, obsolete, intermediis acutissimis, interstitio leviter triangulariter exciso. Caput oblongum thorace latius, basi truncatum, angulis posterioribus leviter rotundatis, apicem versus sensim angustatum, parum convexum, omnium subtilissime punctulatum. Thorax coleopteris haud angustior, latitudine sesqui prope longior, basin versus leviter angustatus, apice attenuatus, lateribus subrectis, angulis anterioribus obtusis, posterioribus rotundatis, subdepressus, omnium subtilissime punctulatus, basi obsolete bifoveolatus, apice linea longitudinali abbreviata subtili exaratus. Elytra thoracis longitudine, confertissime sub-

tiliter punctata. Abdomen subtilissime punctulatum, densius pubescens, segmentis singulis apice subtilissime testaceo-marginatis. Pedes testacei.

Mas abdominis segmento inferiore quinto apice obsoletissime bisinuato, sexto triangulariter exciso; femina eodem leviter triangulariter producto.

Variat colore dilutiore piceo-testaceo.

Obs. L. minimo perquam affine, at paulo maior, antennis, palpis pedibusque testaceis, capite basi latiore, elytris distinctius punctatis etc. distinguendus.

- \* 4. *Sc. minimus*: *Omnium subtilissime punctatus, niger, antennis pedibusque piceis, capite oblongo, thorace oblongo, apice attenuato, elytris thoracis longitudine.* — Long.  $1\frac{1}{4}$  lin.

*Lathrobium minimum* Er. Col. March. I. 511. 13.

Habitat in Germania.

Praecedenti affinis, at dimidio minor, angustus, linearis, niger, pube subtilissima cinerea obductus. Antennae capite plus sesqui longiores, articulo tertio secundo subaequali, 4 — 10 sensim vix brevioribus at paululum crassioribus, ultimo maiore, ovato, apice acuminato, rufo-piceae. Palpi rufo-picei, maxillares articulo tertio crasso, fusco. Caput basi thorace paulo latius, oblongum, basi truncatum, angulis posterioribus rotundatis, antrorsum angustatum, parum convexum, omnium confertissime subtilissimeque punctatum. Thorax coleopteris paulo angustior, latitudine sesqui longior, basin versus subangustatus, apice attenuatus, lateribus rectis, angulis anterioribus obtusis, subrotundatis, posterioribus rotundatis, subdepressus, subtilissime confertissimeque punctatus, basi nonnunquam obsolete bifoveolatus, apice saepius lineola longitudinali brevi subtilissima impressus. Elytra thoracis longitudine, confertim subtilissime punctata, apice picescentia. Abdomen omnium subtilissime confertissimeque punctulatum, apice piceum, segmentis 4 primis subtiliter testaceo-marginatis. Pedes picei testacei.

Mas abdominis segmento inferiore quinto apice late distinctius emarginato, sexto triangulariter exciso; femina eodem leviter triangulariter producto.

- \* 5. *Sc. infirmus*: *Omnium subtilissime punctulatus, opacus, rufo-piceus, abdomine nigricante, antennis pedibusque testaceis, thorace oblongo, elytris thorace longioribus.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Aegypto, Dom. Prof. Ehrenberg.

Habitu fere Lithocharidis ochraceae, at huius generis, et duplo illa minor, subdepressus, dilute rufo-piceus, opacus, totus omnium subtilissime confertissimeque punctatus, pube subtilissima brevissimaque cinerea densius obductus, sericans. Antennae capite plus sesqui longiores, articulo secundo tertioque subaequalibus, 4 — 10 sensim paulo brevioribus, penultimis prope transversis, ultimo subgloboso, apice subito subacuminato, dilute testaceae. Caput thoracis latitudine, oblongum, basi cum angulis posterioribus leviter rotundatum, antrorsum vix angustatum, depressiusculum. Thorax coleopteris dimidio prope angustior, latitudine paulo longior, basin versus subangustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, subrotundatis, posterioribus rotundatis, subdepressus, aequalis.



Elytra thorace tertia parte longiora. Abdomen nigricans, segmentis 4 primis margine apicali, quinto apice, sexto toto dilute rufo-piceis. Pedes flavo-testacei.

Mas abdominis segmento inferiore sexto emarginato; femina eodem rotundato.

- \* 6. *Sc. exiguus*: *Omnium subtilissime punctulatus, opacus, niger, antennis pedibusque testaceis, capite, thorace elytrorumque apice rufo-piceis, elytris thorace oblongo parum longioribus.* — Long. 1 lin.

Habitat in Pennsylvania, Dom. Schüppel.

Habitu omnino praecedentis, at multo minor, linearis, subdepressus, omnium subtilissime confertissime punctatus, opacus, pube subtilissima brevissimaque cinerea densius obductus, sericans. Antennae capite paulo longiores, articulo tertio secundo minore, 3 — 10 sensim paulo brevioribus, ultimo ovato, apice acuminato, laete testaceae. Palpi dilute luteo-testacei. Caput thorace paulo latius, oblongum, basi levissime, angulis posterioribus fortius rotundatis, antrorsum vix angustatum, supra parum convexum, rufo-piceum, ore dilutius. Thorax coleopteris paulo angustior, latitudine summa sesqui longior, basin versus leviter angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, posterioribus rotundatis, parum convexus, basi linea longitudinali, mox evanescente, subtilissima obsolete carinatus, dilute rufo-piceus. Scutellum rufo-piceum. Elytra thorace paululum longiora, nigricantia, apice indeterminate rufo-picea. Abdomen nigrum, segmentis 5 anterioribus margine summo apicali, sexto toto testaceis. Pectus piceum. Pedes laete flavo-testacei.

Mas abdominis segmento inferiore sexto apice triangulariter exciso. Femina latet.

7. *Sc. pygmaeus*: *Omnium subtilissime confertissimeque punctatus, opacus, niger, antennis pedibusque testaceis, capite thoraceque rufo-piceis, elytris thorace oblongo longioribus.* — Long. 1½ lin.

Habitat in ins. Puerto-Rico, Dom. Moritz.

Praecedenti proximus, at paulo maior et elytris longioribus distinctus, linearis, subdepressus, omnium subtilissime confertissimeque punctatus, opacus, pube subtilissima brevissimaque cinerea densius obductus, sericans. Antennae capite sesqui fere longiores, articulo secundo tertio paulo maiore, 4 — 10 sensim vix brevioribus at paulo crassioribus, ultimo apice acuminato, testaceae. Palpi dilute testacei. Caput thorace paulo latius, oblongum, basi truncatum, angulis posterioribus vix rotundatis, ante antennas subito angustatum, supra minus convexum, rufo-piceum. Thorax coleopteris dimidio prope angustior, latitudine plus sesqui longior, basin versus subangustatus, lateribus rectis, apice utrinque oblique truncatus, fere attenuatus, angulis anterioribus fortius rotundatis, posterioribus obtusis, subrotundatis, subdepressus, basi utrinque obsolete impressus, impressionibus carinula brevissima obsolete distinctis, dilutius rufo-piceus. Scutellum rufo-piceum. Elytra thorace tertia fere parte longiora, nigricantia, humeris, sutura margineque apicali indeterminate anguste rufo-piceis. Abdomen nigrum, seg-

mentis 4 primis margine summo apicali, quinto apice, sexto toto piceo-testaceis. Pectus nigro-piceum. Pedes dilute flavo-testacei.

8. *Sc. fasciatellus*: *Omnium subtilissime confertissimeque punctatus, opacus, niger, antennis, pedibus elytrisque testaceis, his fascia lata inaequali, capite piceo, subquadrato.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Puerto-Rico, Dom. Moritz.

Statura omnino praecedentium, omnium subtilissime confertissimeque punctatus, opacus, pube brevissima subtilissimaque cinerea densius obductus, sericans. Antennae capite paulo longiores, crassiusculae, articulo tertio secundo paulo minore, 3 — 10 sensim paulo brevioribus et crassioribus, penultimis fere transversis, ultimo apice subito acuminato, laete testaceae. Palpi dilute testacei. Caput thorace paulo latius, basi truncatum, angulis posterioribus leviter rotundatis, pone oculos haud, ante oculos subito angustatum, supra leviter convexum, rufo-piceum, ore dilutiore. Thorax coleopteris angustior, latitudine summa tertia parte longior, basin versus subangustatus, lateribus rectis, apice utrinque oblique truncatus, angulis omnibus obtusis, subrotundatis, anterioribus leviter prominentibus, subdepressus, aequalis, nigro-piceus. Elytra thorace paulo longiora, basi summa rufo-picea, dein nigra, apice late inaequaliter cum lateribus inflexis testacea. Abdomen segmentis 4 anterioribus margine tenui apicali testaceo, infra quinto apice sextoque toto rufo-piceis. Pectus nigro-piceum. Pedes flavi, coxis anterioribus concoloribus, posticis testaceis.

Mas latet. Femina abdominis segmento inferiore sexto leviter triangulariter producto.

- \* 9. *Sc. pulchellus*: *Laevissimus, nitidus, niger, antennis abdominisque basi testaceis, elytrorum puncto apicali pedibusque flavis, capite orbiculato thoraceque oblongo-ovato, convexo rufis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Sc. laevigato paulo minor et gracilior, nitidus, capite, thorace elytrisque laevissimis. Antennae capite thoraceque paulo breviores, apicem versus leviter incrassatae, articulo secundo tertioque subaequalibus, 4 — 10 sensim brevioribus et crassioribus, ultimo apice abrupte subacuminato, rufo-testaceae. Palpi rufo-testacei. Caput thorace paulo latius, suborbiculatum, supra leviter convexum, piceo-rufum, parce nigro-pilosum. Thorax coleopteris angustior, latitudine summa sesqui longior, utrinque, licet antrorsum minus sensim quam basin versus angustatus, oblongo-ovatus, convexus, piceo-rufus, parce nigro-pilosus. Scutellum piceum. Elytra thorace paulo longiora, nigerrima, apice exterius puncto flavo notata. Abdomen omnium subtilissime punctulatum, nigrum, nitidum, segmentis anterioribus tribus totis, reliquis summo margine apicali testaceis, illis plerumque basi piceis. Pectus nigro-piceum. Pedes elongati, flavi.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso. Femina latet.

Variat capite thoraceque piceis.

**Lithocharis Dej.**

*Boisd et Lacord. Faun. Ent. Paris. I. p. 431. — Er. Col. March. I. p. 511.*  
*Medon Steph. Illustr. V. p. 273. t. 127. f. 2. Sunius ibid p. 274.*  
*Paederus Fam. I. Grav. Mon. p. 138.*

*Labrum apice bidenticulatum.*

*Palpi maxillares articulo quarto distincto, acuto.*

*Ligula membranacea, biloba.*

*Antennae rectae.*

*Tarsi articulo quarto simplice.*

Corpus minutum, elongatum, subdepressum, alatum. Caput plerumque thoracis latitudine, subquadratum, basi profunde constrictum, collo quasi tenui brevi thoraci immissum, exsertum, porrectum, oculis parvis rotundatis, parum prominulis. Labrum latum, transversum, lateribus explanatis rotundatisque membranaceis magis minusve dilatatis, apice medio plerumque bidenticulatum, inter dentes magis minusve emarginatum vel excisum. Mandibulae falcatae, acutae, medio fortiter acuteque 3- vel 4-dentatae. Maxillae (t. IV. f. 29. b.) malis coriaceis, interiore interne, exteriori apice barbatis. Palpi maxillares leviter elongati, articulo primo minuto, secundo tertioque elongatis, longitudine aequalibus, hoc incrassato, quarto minimo, tenuissimo, acuto. Labium (t. IV. f. 29. a.) mento transversum, apice leviter late emarginato, ligula biloba, lobis angustioribus, apice subrotundatis, membraneis, valde distantibus, interstitio dense setoso, paraglossis membraneis, linearibus, ligulae laciniis vix superantibus. Palpi labiales articulo primo minuto, cylindrico, secundo subovato, maiusculo, tertio minuto, tenui, acuto. Antennae filiformes, rectae, articulo primo subelongato, ultimo apice acuminato. Thorax magis minusve distincte quadrangulus. Scutellum triangulare. Elytra truncata. Abdomen lineare, apice angustatum, segmento septimo parum conspicuo, stylis analibus haud distinctis. Pedes leviter elongati, antici nonnunquam subincrassati, tarsis anticis saepius leviter dilatatis, articulis primis feminae triangularibus, maris obcordatis, posterioribus simplicibus, magis minusve elongatis, articulis 4 primis longitudine sensim decreascentibus.

## O b s e r v a t i o n e s.

1. Oris fabrica ceterisque characteribus Stilico nimis forte affinis, a quo tamen habitu facillime distinguitur. Color plerisque testaceus vel ferrugineus, parum nitidus, punctura subtilis. Ceterum a Stilico praecipue antennis longioribus gracilioribusque, labro minus amplo, palporum maxillarium articulo quarto magis conspicuo, paraglossis ligulam haud superantibus, tarsorumque anticorum structura differt. Species nonnullae Lathrobiis similes, a quibus labro denticulato, tarsis anticis, paucis masculis exceptis, perparum vel haud dilatatis, posterioribus articulo primo magis minusve elongato discedunt.

2. Medon Steph. a reliquis huius generis speciebus (Sunius Steph.) labri bidentati causa distinctum, quod in his falso muticum exhibetur.

3. Mas abdominis segmento sexto inferiore exciso, quinto etiam nonnunquam emarginato, femina segmento inferiore sexto apice aut rotundato, aut triangulariter producto. Tarsi antici maris saepius leviter dilatati.

4. Victus sub lapidibus, sub foliis deciduis.

\* 1. *L. castanea*: *Rufo-picea, capite oblongo thoraceque creberrime punctatis, elytris thorace sesqui longioribus. — Long. 3 lin.*

*Er. Col. March. I. 512. 1.*



*Paederus castaneus* Grav. Micr. 60. 3. — Latr. Hist. nat. Crust. et Ins. IX. 348. 8. — Oliv. Encycl. meth. VIII. 628. 6.

*Rugilus castaneus* Mannerh. Brachelytr. 40. 5.

*Paederus quadratus* Beck Beitr. 25. 39. t. 7. f. 39.

*Medon Ruddii* Steph. Illustr. V. t. 127. f. 2.

Habitat in Germania, Gallia.

Sequentibus maior, rufo-picea, subnitida, capite abdomineque obscurioribus. Antennae capite thoraceque paulo breviores, tenuiusculae, articulo tertio secundo sesqui longiore, 4—10 sensim brevioribus et sensim magis obconicis, ultimo fusiformi, apicem versus sensim acuminato. Labrum amplum, lateribus limbo membranaceo dilatatum, medio acute bidentatum. Caput thorace paulo latius, eodemque sesqui longius, oblongum, parum convexum, creberrime punctatum, fronte anteriori medio laeviore, subtiliter griseo-pubescent, lateribus parce nigro-pilosellum. Thorax coleopteris paulo angustior, latitudine antica vix brevior, basin versus angustatus, apice utrinque oblique truncatus, lateribus rectis, angulis omnibus subrotundatis, anterioribus lateraliter prominentibus, leviter convexus, subtiliter confertissime punctatus, subtiliter griseo-pubescent, lateribus parce nigro-pilosellus. Elytra thorace sesqui longiora, subtiliter confertissime punctata, densius subtiliter pubescentia, piceo-ferruginea. Abdomen subtilissime crebre punctulatum, pube depressa subtili fulvescente densius vestitum, piceum, apice et subtus ferrugineum. Pedes longiusculi, graciles, ferruginei.

Mas abdominis segmento inferiore quinto apice medio late emarginato, utrinque subsinuato, sexto acute triangulariter exciso; femina eodem apice leviter triangulariter producta.

\* 2. *L. fuscus*: Ferruginea, capite oblongo, subtiliter punctato, subruguloso, fusco, thorace lateribus confertissime subtilissimeque, dorso confertim subtiliterque punctato, elytris thorace sesqui longioribus. — Long. 2 lin.

Boisd. Lacord. Faun. Ent. Paris. I. 431. 1. — Er. Col. March. I. 513. 2.

*Rugilus fuscus* Mannerh. Brachelytr. 40. 3.

Habitat in Germania, Gallia, Helvetia.

Praecedente plus duplo minor, ferruginea, nitida, capite obscuriore. Antennae capite thoraceque breviores, crassiusculae, articulo tertio secundo parum longiore, 4—10 sensim brevioribus, ultimo apice subito acuminato, Labrum amplum, limbo laterali membranaceo auctum, medio bidentatum. Caput thorace vix latius, at paulo longius, oblongum, confertissime minus subtiliter punctato-subrugulosum, subopacum, subtiliter griseo-pubescent, lateribus pilosellum. Thorax coleopteris parum angustior, latitudine antica vix brevior, basin versus angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, posterioribus rotundatis, parum convexus, lateribus confertissime subtiliter punctatus, opacus, dorso confertim minus subtiliter punctatus, nitidulus, linea media longitudinali laevi basi subelevata, antrorsum obsoleta, subtiliter griseo-pubescent, lateribus parce pilosellus. Elytra thorace sesqui longiora, subtilius confertim punctata, subtiliter griseo-pubescentia. Abdomen subtiliter confertim punctatum, fulvescenti-griseo-dense pubescens. Pedes graciliores, dilute testacei.

Mas abdominis segmento quinto inferiore apice late et profunde sinuato, utrinque in dentem fortem producto, sexto acute emarginato; femina eodem apice leviter triangulariter producto.

- \* 3. *L. brunnea*: *Dilute rufo-picea, capite oblongo, confertim punctato, thorace creberrime, lateribus subtilius punctato, elytris thorace parum longioribus.* — *Long. 2 lin.*

*Er. Col. March. I. 513. 3.*

Habitat in Germania.

Statura et summa affinitas praecedentis, dilute rufo-picea, nitida, capite abdomineque obscurioribus. Antennae capite sesqui longiores, crassiusculae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus, penultimis crassitie vix longioribus, ultimo apice subito acuminato. Labrum latiusculum, lateribus limbo membranaceo dilatatum, apice obsolete bidenticulatum. Caput thorace paulo latius, oblongo-subquadratum, confertissime minus subtiliter punctatum. Thorax coleopteris parum angustior, latitudine paulo brevior, basin versus subangustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, subrotundatis, posterioribus fortius rotundatis, parum convexus, creberrime dorso minus subtiliter, lateribus subtilius punctatus, linea longitudinali media laevi subelevata, antice obsoleta, subtilissime griseo-pubescentia, lateribus parce nigro-pilosellus. Elytra thorace paulo longiora, minus confertim subtilius punctata, minus subtiliter griseo-pubescentia. Abdomen subtilissime punctulatum, densius griseo-pubescentia, apice nigro-pilosellum. Pedes dilute rufo-testacei.

Mas abdominis segmento inferiore quinto apice levissime emarginato, sexto triangulariter exciso; femina eodem apice leviter triangulariter producto.

Obs. Praecedenti simillima, praecipue elytris brevioribus, minus confertim punctatis distinguenda.

- \* 4. *L. diluta*: *Rufo-testacea, capite oblongo, antrorsum angustato thoraceque subtiliter punctatis, elytris thorace sesqui longioribus.* — *Long. 2 lin.*

*Er. Col. March. I. 514. 4.*

Habitat in Germania.

Statura et summa affinitas praecedentium, rufo-testacea, nitida, abdomine piceo. Antennae capite plus sesqui longiores, articulo tertio secundo paulo longiore, 4 — 8 sensim brevioribus, ultimis tribus sensim paulo maioribus, ultimo maiusculo, apice acuminato. Labrum latiusculum, lateribus limbo angustiore membranaceo auctum, apice obsoletius bidenticulatum. Caput thorace parum latius, at tertia parte longius, oblongum, antrorsum angustatum, subtiliter dense punctatum, antice linea tenui longitudinali media laevi subelevata, subtilissime griseo-pubescentia, lateribus nigro-pilosellum. Thorax coleopteris paulo angustior, latitudine paulo brevior, basin versus modice angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, posterioribus rotundatis, parum convexus, aequaliter confertissime subtiliter punctatus, subtilissime griseo-pubescentia, lateribus parce nigro-pilosellus. Elytra thorace sesqui longiora, subtiliter confertis-

sime punctata, subtiliterque griseo-pubescentia. Abdomen confertissime subtiliter punctulatum, densius griseo-pubescent, piceum, apicem versus rufo-testaceum. Pedes dilute testacei.

Mas latet. Femina abdominis segmento inferiore sexto apice leviter triangulariter producto.

- \* 5. *L. rufiventris*: *Rufo-picea, capite thoraceque nigro-piceis, capite subquadrato, fortius punctato, medio laevi, thorace subtiliter punctato, medio subtiliter canaliculato, elytris thorace sesqui longioribus. — Long. 2 lin.*

*Er. Col. March. 1. 514. 5.*

*Lathrobium rufiventre Nordm. Symb. 147. 19.*

Habitat in Germania.

Statura et affinitas praecedentium, picea, nitida. Antennae capite sesqui longiores, apice leviter incrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus, penultimis crassitie vix longioribus, ultimo apice abrupte acuminato, rufae. Labrum apice obsoletius bidenticulatum. Os rufo-piceum. Caput basi thorace paululum latius, subquadratum, antrosum subangustatum, punctatum, fronte media laeviore, subtiliter pubescens, lateribus pilosellum, nigrum, nitidum. Thorax coleopteris paulo angustior, longitudine antica paulo brevior, basin versus subangustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus leviter, posterioribus fortius rotundatis, subtilius et minus confertim punctatus, linea longitudinali media laevi parum distincta, medio subtiliter canaliculata, subtiliter pubescens, nigro-pilosellus. Elytra thorace sesqui longiora, minus confertim subtilius punctata, griseo-pubescentia, lateribus parce nigro-pilosella. Abdomen subtiliter dense punctulatum subtiliterque griseo-pubescent, parce nigro-pilosellum, apice rufo-piceum. Pedes dilute rufo-piceis.

Femina abdominis segmento inferiore sexto apice leviter triangulariter producto; mas eodem „paulo emarginato” sec. Nordm.

Variat corpore rufo, capite, thorace abdomineque supra praeter apicem piceis.

Obs. Capitis punctura minus conferta praecipue ab affinis distincta.

- \* 6. *L. ferruginea*: *Ferruginea, capite subquadrato, confertissime fortiter punctato, thorace crebrius profundiusque punctato, subtiliter canaliculato, elytris thorace paulo longioribus. — Long. 2 lin.*

Habitat in Austria, Dom. Schüppel.

Statura et affinitas praecedentium. Corpus ferrugineum, nitidum. Antennae capite plus sesqui longiores, apice subincrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, ultimo apice subito acuminato. Labrum amplum, lateribus limbo membranaceo auctum, apice obsoletius bidentatum. Caput thorace vix latius, subquadratum; supra confertissime sat fortiter punctatum, fronte media vix laeviore, antius subrugulosa, subtiliter pubescens, lateribus parce pilosellum. Thorax coleopteris paulo angustior, latitudine apicis vix brevior, basin versus modice angustatus, apice utrinque oblique truncatus, angulis anterioribus leviter, posterioribus fortius rotundatis, leviter convexus, minus



confertim licet sat crebre fortius et sat profunde punctatus, linea media longitudinali laevi subtiliter canaliculata, subtiliter griseo-pubescentis, lateribus parce nigro-pilosellus. Elytra thorace paulo longiora, minus confertim aequaliter subtilius punctata, tenuiter pubescentia. Abdomen parcius subtiliusque punctatum, tenuiter pubescens, segmento quinto basi piceo. Pedes dilute testacei.

Mas abdominis segmento inferiore quinto apice leviter bisinuato, sexto leviter triangulariter exciso. Femina latet.

Obs. A praecedentibus praecipue capitis punctura forti confertissima distincta.

\* 7. *L. melanocephala*: Nigro-picea, nitida, capite suborbiculato, parce punctato, elytris piceo-testaceis, thorace paulo longioribus. — Long.  $1\frac{1}{2}$  — 2 lin.

Er. Col. March. I. 515. 6.

*Paederus melanocephalus* Fab. Ent. Syst. I. II. 538. 10. Syst. El. II. 610. 10. — Panz. Ent. Germ. 363. 11. Faun. Germ. 36. 22.

*Staph. tricolor* Marsh. Ent. Britt. 516. 55.

*Paederus bicolor* Oliv. Ent. III. 44. 7. t. I. f. 4. — Grav. Micr. 59. 2. Mon. 138 3. — Latr. Hist. nat. Crust. et Ins. IX. 348. 7. — Oliv. Encycl. meth. VIII. 630. 12. — Panz. Faun. Germ. 104. 15.

*Rugilus bicolor* Mannerh. Brachelytr. 40. 4. — Runde Brachelytr. Hal. 14. 4.

*Lithocharis bicolor* Boisd. et Lacord. Faun. Ent. Paris. I. 433. 3.

*Lathrobium ruficorne* Latr. Gen. Crust. et Ins. I. 290. 3.

Habitat in Europa temperata.

Antennae capite thoraceque breviores, apicem versus vix crassiores, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus, ultimo acuminato, rufo-testaceae, medio nonnunquam fuscescentes. Os rufo-piceum. Labrum margine obsolete denticulatum. Caput thoracis latitudine, suborbiculatum, depressiusculum, parce subtilius punctulatum, fronte media laeve, lateribus parce nigro-pilosum, piceo-nigrum, nitidum. Thorax coleopteris parum angustior, latitudine haud brevior, basin versus subangustus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, posterioribus rotundatis, parum convexus, crebre punctatus, linea media longitudinali laevi, testaceo-rufus, nitidus. Scutellum rufum. Elytra thorace parum longiora, densius subtiliter punctata, tenuiter griseo-pubescentia, piceo-testacea, nitida. Abdomen confertim subtiliter punctulatum, pube subtili depressa grisea densius vestitum, parum nitidum, nigrum, segmentorum margine posteriore apiceque piceis. Pectus nigro-piceum. Pedes testacei, crassiusculi, tarsi posterioribus sat elongatis, anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore sexto longitudinaliter leviter impresso, apice leviter triangulariter exciso; femina eodem apice leviter triangulariter producta.

\* 8. *L. angularis*: Rufo-testacea, nitida, thorace rufo, capite, ano elytrisque nigris, his angulis exterioribus testaceis, capite thoraceque convexis elytrisque parce punctatis. — Long.  $2\frac{1}{2}$  lin.

Habitat Novae Valentiae in Columbia, Dom. Moritz, in Mexico, coll. Dom. Chevrolat.

Primo intuitu Paederium simulat. Antennae capite thoraceque paulo breviores, tenuiusculae, articulo tertio secundo sesqui longiore, quarto iterum dimidio breviorē; 4 — 11 sensim paulo longioribus, teretibus, ultimo apice acuminato, dilute testaceae. Palpi testacei. Labrum transversum, apice acute bidenticulatum, inter dentes profundius sinuatum. Mandibulae validiores, rufo-piceae. Caput thoracis latitudine, transversim subquadratum, leviter convexum, ad latera et inter oculos parce vage punctatum (et parce nigro-pilosum, nigrum, pernitidum. Thorax coleopteris parum angustior, latitudine tertia parte brevior, lateribus rectis, apice truncatus, angulis omnibus subrotundatis, supra convexus, punctis raris obsoletisque versus latera notatus, parce nigro-pilosus, ceterum laevis glaberrimusque, testaceo-rufus, pernitidus. Scutellum nigrum. Elytra thorace paulo longiora, leviter convexa, stria forti suturali punctisque subtilioribus parvis subseriatis impressa, parce griseo-pilosa, pernitida, nigerrima, angulo exteriorē late ad suturam usque testaceo. Abdomen parce subtilissime punctulatum, tenuiter subtiliterque cinereo-pubescent, nitidum, segmentis 4 anterioribus rufo-testaceis, apicalibus duobus nigris. Pectus rufum. Pedes graciliores, rufo-testacei, femoribus pallide flavescens, tarsis posterioribus sat elongatis, antice simplicibus.

Mas abdominis segmento inferiore quinto apice subsinuato, sexto leviter longitudinaliter impresso, apice emarginato. Femina latet.

9. *L. macularis*: *Rufa, nitida, elytrorum macula magna transversali anoque nigris, thorace fortiter convexo, elytris parce serialim punctatis.* — Long. 3 lin.

Habitat in Guyana Venezuelensi, Dom. Moritz.

Affinis praecedenti. Antennae capite thoraceque paulo breviores, tenuiores, articulo tertio secundo vix breviorē, 3 — 10 sensim paulo brevioribus et crassioribus, levissime obconicis, ultimo oblongo, apice subito acuminato, testaceae. Palpi testacei. Labrum apice acute bidenticulatum, inter dentes profundius sinuatum. Mandibulae validiores. Caput thorace prope latius, basi truncatum, medio subsinuatum, antrorsum subangustatum, supra leviter convexum, punctis raris sparsisque minus subtilibus impressum, parce nigro-pilosum, saturate rufum, pernitidum. Thorax coleopteris vix angustior, latitudine haud brevior, basin versus leviter angustatus, lateribus subrectis, apice truncatus, angulis anterioribus subrectis, posterioribus rotundatis, supra valde convexus, laevis, parce nigro-pilosus, saturate rufus, nitidus. Scutellum concolor. Elytra thorace parum longiora, stria lata suturali punctisque parvis seriatis minus subtilibus impressa, nitida, rufa, macula magna transversali a medio margine laterali ad suturae apicem deflexa summoque margine suturali nigris, apice testaceo. Abdomen subtilissime punctulatum, tenuiter subtiliterque cinereo-pubescent, rufum, segmentis apicalibus duobus nigris. Pectus rufum. Pedes longiores, graciliores, rufi, femoribus pallide flavis, tarsis posterioribus modice elongatis, anticis leviter dilatatis.

- \* 10. *L. confluens*: *Subdepressa, nigra, ore thoracisque limbo rufo-piceis, elytris apice testaceis, capite thoraceque transversim subquadratis, creberrime punctato-rugulosis.* — Long. 1 $\frac{3}{4}$  lin.

*Lathrobium confluentum* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 456. 8. (forte)

Habitat in Carolina meridionali, Dom. Zimmermann.

Reliquis brevior, depressa. Antennae capite thoraceque breviores, crassiusculae, articulo secundo tertioque subaequalibus, 4—10 sensim paulo brevioribus et crassioribus, ultimo apice subito acuminato, medio fuscae, basi apiceque rufo-piceae. Os rufo-piceum, palpis testaceis. Caput thorace paulo latius, subquadratum, depressum, confertissime minus subtiliter punctato-rugosum, nigrum, opacum, subtilissime cinereo-pubescent. Thorax coleopteris paulo angustior, latitudine antica paulo longior, basin versus angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus prominulis posterioribusque subrotundatis, lateribus distinctius marginatus, subdepressus, confertissime minus subtiliter punctato-subrugosus, opacus, rufo-piceus, disco nigro, subtiliter cinereo-pubescent. Scutellum piceum. Elytra thorace sesqui longiora, crebre punctata, tenuiter pubescentia, nitida, nigro-picea, apice late testacea. Abdomen subtilius confertim punctatum, densius pubescens, subnitidum, supra nigrum, apice summo rufo-piceo, subtus medio rufo-piceum, apice testaceum. Pectus piceum. Pedes breviusculi, cum coxis rufo-testacei, tarsis posterioribus breviusculis, anticis in utroque sexu levissime dilatatis.

Mas abdominis segmento inferiore sexto apice leviter triangulariter inciso; femina eodem leviter triangulariter producto.

11. *L. dorsalis*: Subdepressa, confertissime subtiliter punctata, opaca, piceo-rufa, antennis pedibusque rufo-testaceis, elytris dorso nigris, thorace subtilissime canaliculato. — Long. 2 lin.

Habitat in Puerto-Rico, Dom. Moritz.

Statura omnino praecedentis, subdepressa, obscure rufo-ferruginea, vix nitida, tota confertissime subtiliter punctata, pube brevi subtili grisea densius vestita. Antennae capite paulo longiores, apicem versus sensim subincrassatae, articulo tertio secundo parum longiore, 4—10 sensim paulo brevioribus et crassioribus, penultimis vix transversis, ultimo ovato, apice leviter acuminato, saturate rufo-testaceae. Palpi rufo-testacei. Labrum apice medio bidenticulatum. Caput thorace latius, suborbiculatum, depressiusculum. Thorax coleopteris paulo angustior, latitudine antica paulo brevior, basin versus leviter angustatus, apice utrinque oblique truncatus, angulis anterioribus obtusis, modice prominulis, posterioribus rotundatis, pone medium lineola subtilissima vix perspicue canaliculatus. Elytra thorace sesqui longiora, subdepressa, dorso late nigra, margine tenui apicali et limbo lato laterali ferrugineis. Abdomen apice summo testaceo. Pedes rufo-testacei, crassiusculi, breviusculi, femoribus anterioribus infra subdentatis, tarsis posterioribus minus elongatis, anticis leviter dilatatis.

Mas abdominis segmento inferiore sexto apice late triangulariter exciso, septimo conspicuo. Femina latet.

- \* 12. *L. depressa*: Subdepressa, confertim subtiliter punctata, parum nitida, rufo-ferruginea, capite transversim subquadrato, subruguloso, thorace basin versus obsolete subtilissime carinato, elytris thorace paulo longioribus. — Long. 1½ lin.

Habitat in Columbia.



Statura breviuscula depressa praecedentium, rufo-ferruginea, nitida, pube subtilissima grisea vestita, antennis pedibusque ferrugineo-testaceis. Antennae capite vix sesqui longiores, apicem versus leniter incrassatae, articulo tertio secundo haud longiore, 3 — 10 sensim paulo brevioribus et crassioribus, ultimo apice abrupte acuminato, rufo-testaceae. Palpi testacei. Labrum apice 4-denticulatum. Caput thorace paulo latius, basi truncatum, angulis posterioribus et lateribus leviter rotundatum, depressum, confertissime subtiliter punctatum, subrugulosum. Thorax coleopteris angustior, latitudine paulo brevior, basin versus modice angustatus, apice utrinque oblique truncatus, angulis anterioribus subrectis, modice prominentibus, posterioribus subrotundatis, supra depressus, confertim subtiliter punctatus, linea media longitudinali subelevata obsoleta, ante medium evanescente, nonnunquam subtilissime canaliculata, basi utrinque prope lineam elevatam obsolete longitudinaliter impressus, basi summa obsolete 4-punctatus. Elytra thorace paulo longiora, subtiliter dense punctata, stria suturali, pone medium abbreviata, punctata impressa. Abdomen subtilissime punctulatum, piceum, segmentorum marginibus apiceque summo dilutioribus, subtus cum reliquo corpore piceo-rufescens. Pedes rufo-testacei, breves, tarsis brevioribus, articulo quarto subcordato, quinto brevi, anticis in utroque sexu levissime dilatatis.

Mas abdominis segmento inferiore sexto apice triangulariter exciso; femina eodem acute rotundato.

\* 13. *L. biseriata*: *Depressa, confertissime subtiliter punctata, opaca, ferruginea, capite subquadrato, subruguloso, thorace basi 4-punctato, elytris thorace sesqui longioribus, dorso obsolete biserialim punctatis.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Columbia, Dom. Moritz.

Statura omnino et summa affinitas praecedentis, fusca ferrugineave, opaca, pube subtilissima grisea tenuiter vestita, omnium confertissime subtiliter punctata. Antennae capite paulo longiores, apice subincrassatae, articulo tertio secundo paulo longiore, articulis 4 — 10 sensim paulo brevioribus et crassioribus, ultimo ovato, apice subito acuminato, testaceo-rufae. Palpi rufo-testacei. Labrum apice obsoletius bidenticulatum, inter denticulos sinuatum. Caput thorace paulo latius, transversim subquadratum, basi truncatum, angulis posterioribus leviter rotundatis, lateribus rectis, ante oculos subito angustatum, depressum, confertissime longitudinaliter subrugulosum. Thorax coleopteris paulo angustior, latitudine brevior, basin versus modice angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus subrectis, prominulis, posterioribus obtusis, depressus, linea media longitudinali elevata vix perspicua subtilissime carinatus, basi punctis 4 sat distinctis, transversim positis, intermediis subtilioribus, notatus. Elytra thorace sesqui longiora, depressa, stria suturali duabusque dorsalibus paulo fortius punctatis, apice obsolescentibus notata. Abdomen apice testaceum. Pedes testacei, breviusculi, tarsis brevioribus, depressiusculis, articulo quarto subcordato.

Mas abdominis segmento inferiore sexto apice emarginato; femina eodem rotundato.

Obs. Proxima praecedenti, at elytris paulo longioribus, dorso bistriatis thorace basi evidenter 4-punctato distinguenda.

- \* 14. *L. oblita*: *Subdepressa, picea, capite subquadrato, creberrime punctato, thorace creberrime subtiliter punctato, elytris thorace sesqui longioribus anoque rufo-piceis, pedibus rufo-testaceis.* — Long.  $1\frac{2}{3}$  lin.

Habitat in Columbia, Dom. Moritz.

Statura brevis lataque praecedentis, nigro-picea, nitidula, ore elytris anoque rufo-piceis. Antennae capite sesqui longiores, apice subincrassatae, articulo secundo tertioque subaequalibus, 4 — 10 sensim brevioribus et crassioribus, ultimo apice subito acuminato, densius pilosellae, rufo-testaceae. Labrum apice obsolete bidenticulatum, interstitio inter dentes leviter triangulariter exciso. Caput thoracis latitudine, subquadratum, angulis posterioribus rotundatis, subdepressum, creberrime minus subtiliter punctatum, fronte media vix laeviore, subtiliter nigro-pubescent, lateribus nigro-pilosum. Thorax coleopteris parum angustior, latitudine paulo brevior, basin versus subangustatus, lateribus rectis, angulis anterioribus obtusis, modice prominulis, posterioribus rotundatis, subdepressus, creberrime subtilius punctatus, linea media longitudinali laevi, parum distincta, subtiliter nigro-pubescent, lateribus parce nigro pilosus. Elytra thorace sesqui longiora, subtilius dense punctata, griseo-pubescentia. Abdomen omnium subtilissime punctulatum, densius griseo-pubescent, segmentorum marginibus summis apiceque rufo-piceis. Corpus subtile rufo-piceum. Pedes breviusculi, rufo-testacei, tarsis posterioribus minus elongatis, anticis leviter dilatatis.

Mas abdominis segmento inferiore sexto apice triangulariter exciso; femina eodem rotundato.

- \* 15. *L. curtula*: *Subdepressa, nigro-picea, capite transversim subquadrato, parcius fortiusque punctato, medio laevi, thorace punctato, elytris thorace sesqui longioribus, apice anoque rufo-piceis, pedibus rufo-testaceis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Statura omnino et summa similitudo praecedentis, nigro-picea, nitida. Antennae capite sesqui longiores, apicem versus haud crassiores, articulo secundo tertioque aequalibus, 4 — 10 sensim paulo brevioribus, penultimis crassitie vix longioribus, ultimo subito fortius acuminato, rufo-testaceae. Palpi rufo-testacei. Labrum apice obsolete bidentatum, inter dentes profundius incisum. Os rufo-piceum. Caput thorace vix latius, transversim subquadratum, angulis posterioribus subrectis, depressum, minus dense fortius punctatum, fronte media laeviore, subtiliter nigro-pubescent, parce nigro-pilosum, nigrum, nitidum. Thorax coleopteris angustior, latitudine tertia parte brevior, basin versus subangustatus, lateribus rectis, angulis anterioribus obtusis, leviter prominentibus, posterioribus subrotundatis, parum convexus, densius fortiusque punctatus, linea media longitudinali laevi, subtiliter tenuiterque nigro-pubescent, lateribus nigro-pilosus, niger, nitidus. Elytra subtilius minus dense punctata, parcius griseo-pubescentia, fusco-picea, nitida, apice dilute rufo-picea. Abdomen breviusculum, obsolete punctulatum, longius et minus dense griseo-pubescent, nigro-piceum,

ano rufo-piceo. Corpus subtus rufo-piceum. Pedes testacei, tarsi posterioribus leviter elongatis, anticis fortius dilatatis.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso; femina eodem rotundato.

16. *L. posticata*: *Subdepressa, picca, nitidula, capite thoraceque parcius punctatis, elytris apice nigris, pedibus flavis.* — Long. 1 lin.

Habitat in Puerto-Rico, Dom. Moritz.

Habitu omnino praecedentium, at duplo minor. Antennae capite sesqui longiores, articulo tertio secundo paulo minore, 3 — 10 sensim paulo brevioribus et crassioribus, penultimis fere transversis, ultimo maiusculo, subovato, pilosellae, totae testaceae. Os testaceum. Labrum apice medio bidenticulatum, inter dentes leviter excisum. Caput thorace paulo latius, basi, lateribus et angulis posterioribus leviter rotundatis, antrorsum subangustatum, depressiusculum, crebrius punctatum, fronte media laeviore, subtiliter parceque puberulum, piceum, nitidum. Thorax coleopteris paulo angustior, latitudine antica prope brevior, basin versus modice angustatus, apice utrinque oblique truncatus, angulis anterioribus subrectis, leviter prominentibus, posterioribus subrotundatis, leviter convexus, sat crebre minus subtiliter punctatus, linea media longitudinali laevi, pube erecta brevi subtilique parce vestitus, piceus, nitidus. Elytra thorace tertia parte longiora, densius subtiliusque punctata, subtiliter pubescentia, laete testacea, apice tertia parte nigerrima. Abdomen subtilissime punctatum, longius tenuiter pubescens, supra piceum, apice testaceum, infra cum reliquo corpore rufo-testaceum. Pedes flavi, breviusculi, femoribus anticis posticisque leviter incrassatis, tarsi posterioribus minus elongatis, anticis leviter dilatatis.

Mas abdominis segmento inferiore sexto apice triangulariter exciso. Femina latet.

- \* 17. *L. corticina*: *Brevis, subdepressa, subtilissime confertissimeque punctata, opaca, rufo-picea, capite subquadrato, basi emarginato, thorace leniter carinato.* — Long. 3 lin.

*Paederus corticinus* Grav. Micr. 184. 1. — Oliv. Encycl. meth. VIII. 629. 9.

*Lathrob. rufescens* Nordm. Symb. 148. 20.

*Lathrob. millepunctatum* Say Transact. Amer. Philosoph. Transact. Philadelph. Nov. Ser. IV. 456. 7.

Habitat in America septentrionali.

Corpus breviusculum, latum, subdepressum, rufo-piceum, opacum, pube subtilissima depressa grisea densius vestitum, capite abdomineque fuscis. Antennae capitis thoracisque prope longitudine, graciliores, apicem versus haud crassiores, articulo tertio secundo sesqui longiore, 4 — 10 sensim brevioribus, ultimo subovato, apice acuminato, rufo-testaceae. Labrum margine laterali dilatato, medio unidentatum. Caput maiusculum, thoracis latitudine, subquadratum, basi profundius emarginatum, depressiusculum, subtilissime confertissimeque punctatum, linea tenuissima longitudinali laevi. Thorax coleopteris paulo angustior, latitudine antica haud brevior, basin versus leviter angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus modice prominentibus, obtusis, posterioribus leviter ro-



tundatis, parum convexus, subtilissime confertissimeque punctatus, linea media longitudinali tenui carinata laevi. Elytra thorace paulo longiora, subtilissime confertissimeque punctata, disco fuscесcentia. Abdomen confertissime subtilissimeque punctatum, fuscum, segmentorum marginibus apiceque rufopiceis. Pedes testacei, tarsi posterioribus modice elongatis, anticis leviter dilatatis.

Mas abdominis segmento inferiore sexto apice late triangulariter emarginato; femina eodem leviter triangulariter producto.

Variat dilutior corpore testaceo, elytris plaga dorsali determinata fusca.

- \* 18. *L. hepatica*: Subtilissime confertissimeque punctata, opaca, fusca, antennis, abdominis segmentorum marginibus, elytrisque ferrugineis, his plaga fusca, pedibus testaceis, capite quadrato, thorace transversim subquadrato, leviter carinato. — Long. 3 lin.

Habitat in Columbia, Dom. Moritz.

*L. corticinae* aequalis, opaca, pube subtilissima grisea densius vestita. Antennae capite thoraceque vix breviores, apicem versus leviter attenuatae, articulo tertio secundo sesqui prope longiore, 4 — 10 sensim paulo brevioribus et paulo magis transversis, ultimo apice fortiter acuminato, piceo-rufae. Labrum margine laterali membranaceo dilatato, apice medio denticulo prominulo instructum. Caput thorace paulo latius, subquadratum, lateribus rectis, basi late leviter emarginatum, angulis posterioribus subrectis, depressiusculum, subtilissime confertissime punctatum, linea tenuissima obsoleta longitudinali laevi, lateribus parce pilosum, fuscum. Thorax coleopteris paulo angustior, latitudine antica paulo brevior, basin versus leviter angustatus, apice utrinque oblique truncatus, angulis anterioribus subrectis, modice prominentibus, posterioribus rotundatis, subdepressus, subtilissime confertissimeque punctatus, linea angusta media longitudinali laevi, fortius elevata carinatus, lateribus parce pilosus, rufescenti-fuscus. Elytra thorace paulo longiora, subtilissime confertissimeque punctata, ferruginea testaceave, plaga dorsali longitudinali indeterminata fusca. Abdomen omnium subtilissime punctulatum, fuscum, ano testaceo, segmentorum marginibus ferrugineis. Pedes longiusculi, dilute testacei, tarsi posterioribus minus elongatis, anticis in utroque sexu leviter dilatatis.

Mas abdominis segmento inferiore quinto apice trisinuato, sexto late triangulariter exciso, septimo conspicuo; femina segmento sexto apice rotundato, septimo occulto.

Obs. Differt ab *L. corticina* praecipue statura graciliore, capite basi minus profunde emarginato et elytris brevioribus.

- \* 19. *L. cinnamomea*: Subtilissime confertissimeque punctata, opaca, rufotestacea, capite, elytrorum plaga obsoleta abdomineque fuscis, pedibus testaceis, capite subquadrato, thorace obsolete carinato. — Long. 1½ lin.

Habitat in Columbia, Dom. Moritz.

Habitu omnino praecedentis, at dimidio minor, opaca, pube subtilissima depressa cinerascens obducta. Antennae capitis thoracisque prope longitudine, apicem versus haud attenuatae, articulo tertio secundo sesqui longiore, 4 — 10 sensim brevioribus, cylindricis, ultimo subovato, apice acuminato,

rufo-testaceae. Labrum transversum, limbo laterali membranaceo, fortiter dilatato, apice medio denticulo obsoleto armatum. Caput thorace vix latius, subquadratum, lateribus rectis, basi late leviter emarginatum, angulis posterioribus subrectis, supra leviter convexum, aequaliter confertissime subtilissime punctatum, lateribus pilosellum, rufo-piceum, ore dilutiore. Thorax coleopteris paulo angustior, latitudine antica vix brevior, basin versus modice angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusiusculis, posterioribus subrotundatis, parum convexus, subtilissime confertissimeque punctatus, linea media longitudinali tenui subelevata laevi obsoleto carinatus, lateribus parce pilosus, saturate rufo-testaceus vel rufus. Elytra thorace tertia fere parte longiora, subtilissime confertissimeque punctata, testacea, plaga indeterminata obsoletaque prope suturam fusca. Abdomen confertissime subtilissimeque punctatum, fuscum, segmento sexto toto, reliquis margine rufo-testaceis ferrugineisve. Pectus rufo-testaceum. Pedes dilutius testacei, longiusculi, tarsis posterioribus minus elongatis, anticis in utroque sexu leviter dilatatis.

Mas abdominis segmento inferiore sexto apice late triangulariter exciso; femina eodem apice leviter producto, acute rotundato.

Obs. Proxima praecedenti, at statura minore, graciliore, thorace obsoleto carinato, elytris paulo longioribus, etc. distincta.

20. *L. limbata*: Subtilissime confertissimeque punctata, opaca, fusca, antennis testaceis, elytrorum limbo laterali et apicali pedibusque flavis, capite suborbiculato, thorace subquadrato, subtiliter carinato. — Long. 2 lin.

Habitat in Columbia, Dom. Moritz.

Habitus omnino *L. ochraceae*, at duplo fere maior, opaca, pube subtilissima cinerascenti densius vestita. Antennae capite thoraceque breviores, validiusculae, apicem versus neque attenuatae, neque incrassatae, articulo tertio secundo sesqui longiore, 4 — 10 sensim paululum brevioribus, cylindricis, ultimo oblongo-subovato, apice fortius acuminato, rufo-testaceae, ante medium fuscescentes. Os testaceum. Labrum transversum, limbo laterali membranaceo fortiter dilatato, apice truncatulum et obsoleto denticulatum. Caput thoracis prope latitudine, basi, lateribus et angulis posterioribus leviter rotundatis, supra leviter convexum, omnium, subtilissime confertissime punctatum, utrinque nigro-pilosellum, nigro-fuscum. Thorax coleopteris tertia parte angustior, latitudine haud brevior, basin versus haud angustatus, apice utrinque oblique truncatus, angulis anterioribus obtusis, posterioribus subrotundatis, parum convexus, omnium subtilissime confertissime punctatus, linea media longitudinali tenui, posterius fortius elevata, subtiliter carinatus, lateribus parce nigro-pilosus, fuscus. Elytra thorace paulo longiora, depressa, confertissime subtilissimeque, licet distinctius quam caput et thorax punctata, dorso fusca, limbo lato laterali apicalique determinate pallide flavo-testaceis. Abdomen confertissime subtilissimeque punctatum, supra nigro-fuscum, segmentis singulis testaceo-marginatis, infra rufo-testaceum, segmentis singulis basi lateraliter fuscis. Pectus fuscum. Pedes dilute testacei, femoribus flavis, tarsis posterioribus modice elongatis, anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore sexto apice truncato, septimo exserto; femina sexto apice rotundato, septimo occulto.

- \* 21. *L. fumata*: Subtilissime confertissimeque punctata, brunnea, pedibus testaceis, capite suborbiculato, nigro, thorace subquadrato, subtiliter carinato. — Long.  $2\frac{1}{4}$  lin.

Habitat in Brasilia, Dom. Melly.

Habitu omnino *L. ochraceae*, at plus sesqui maior. Antennae capite thoraceque paulo breviores, crassiusculae, apicem versus neque incrassatae, neque attenuatae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus, ultimo subovato, apice abrupte acuminato, piceae, apice rufo-ferrugineae. Labrum transversum, limbo laterali membranaceo leviter dilatato, apice truncatum, obsolete denticulatum, rufum. Palpi rufo-picei. Caput thoracis latitudine, basi et lateribus leviter, angulis posterioribus fortius rotundatum, ante oculos subito angustatum, supra parum convexum, anterius depressum, subtilissime confertissimeque punctatum, lateribus parce nigropilosum, nigrum, subnitidum. Thorax coleopteris paulo angustior, latitudine haud brevior, basin versus vix angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, posterioribus leviter rotundatis, subdepressus, omnium subtilissime confertissime punctatus, linea media longitudinali tenui leviter elevata, subtiliter at sat distincte carinatus, subtilissime griseo-pubescent, lateribus parce nigro-pilosus, brunneus, parum nitidus. Elytra thorace paulo longiora, confertim subtiliter pubescentia, rufescenti-brunnea, apicem versus sensim nigra, nitidula. Abdomen subtilissime punctulatum, subtiliter pubescens, supra obscure brunneum, subtus ferrugineum, segmentis basi obscurioribus. Pectus brunneum. Pedes testacei, tarsis posterioribus modice elongatis.

Mas latet. Femina abdominis segmento inferiore sexto apice leviter triangulariter producto, tarsis anticis simplicibus.

- \* 22. *L. infuscata*: Subtilissime confertissimeque punctata, fusca, antennis pedibusque testaceis, elytris ferrugineis, disco fuscis, capite suborbiculato, nigro, thorace quadrato, obsolete carinato. — Long.  $1\frac{1}{4}$  lin.

Habitat in Columbia, Dom. Moritz.

Habitus omnino et summa similitudo *L. ochraceae*, at dimidio minor, pube subtilissima grisea obducta, capite thoraceque lateribus parce nigropilosis. Antennae capitis thoracisque prope longitudine, crassiusculae, apicem versus subincrassatae, articulo secundo tertioque subaequalibus, 4 — 10 sensim paulo brevioribus et crassioribus, ultimo apice subito subacuminato, totae rufo-testaceae. Palpi rufo-testacei. Os rufo-piceum. Labrum apice subsinuaturn, medio obsolete bidenticulatum. Caput thoracis latitudine, basi, lateribus et angulis posterioribus leviter rotundatis, ante oculos subito angustatum, supra posterius leviter convexum, anterius subdepressum, omnium subtilissime confertissimeque punctatum, nigrum, subnitidum. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus rectis, parallelis, apice utrinque oblique truncatus, angulis anterioribus obtusis, posterioribus subrotundatis, parum convexus, omnium subtilissime confertissimeque punctatus, linea media longitudinali tenui, subelevata, obso-



lete carinatus, fusco-piceus, subnitidus. Elytra thorace tertia parte longiora, subdepressa, confertissime subtilissimeque, licet distinctius quam caput et thorax, punctata, ferruginea vel fusco-testacea, plaga magna lata communi fusca, seu fusca, limbo lato laterali apicalique fusco-testaceis. Abdomen confertissime subtilissimeque punctatum, longius pubescens, fuscum, segmentis 4 primis summo margine apicali, quinto apice lato, sexto toto testaceis. Pectus fuscum. Pedes flavi, femoribus anticis posticisque leviter incrassatis, tarsis posterioribus minus elongatis, anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem rotundato.

\* 23. *L. ochracea*: Subtilissime confertissimeque punctata, opaca, ferruginea, capite subtriangulari nigro, thorace quadrato, obsolete carinato, — Long.  $1\frac{2}{3}$  lin.

Boisd. et Lacord. Faun. Ent. Paris. I. 432. 1. — Er. Col. March. I. 516. 7.

*Paederus ochraceus* Grav. Micr. 59. 1. Mon. 138. 1. — Latr. Hist. nat. Crust. et Ins. IX. 347. 6. — Panz. Faun. Germ. 104. 14. — Oliv. Encycl. meth. VIII. 630. 14.

*Paederus rubricollis* Grav. Mon. 138. 2. — Oliv. Encycl. meth. VIII. 630. 15. — Gyll. Ins. Suec. II. 376. 5. — Sahlb. Ins. Fenn. I. 344. 5.

*Rugilus rubricollis* Mannerh. Brachelytr. 40. 6. — Runde Brachelytr. Hat. 14. 3. *Lithoch. testacea* Boisd. et Lacord. Faun. Ent. Paris. I. 432. 2. (forte)

Habitat in Europa, in Americae ins. St. Johannis, Dom. Moritz, in Brasilia, Dom. Lhotzki.

Corpus ferrugineum vel rufo-testaceum, opacum, pube subtilissima griseo-sericea densius vestitum. Antennae capitis thoracisque fere longitudine, apicem versus neque attenuatae, neque incrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim brevioribus, ultimo apice fortius acuminato, testaceae. Palpi et os testacea. Labrum apice medio obsoletius bidenticulatum, inter dentes leviter triangulariter excisum. Caput thoracis latitudine, basi truncatum, angulis posterioribus rotundatis, antrorsum angustatum, supra posterius leviter convexum, anterius depressiusculum, aequaliter subtilissime confertissimeque punctatum, lateribus parce pilosum, nigrum. Thorax coleopteris tertia parte angustior, latitudine haud brevior, basin versus haud angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, posterioribus subrotundatis, subdepressus, subtilissime confertissimeque punctatus, linea tenni subelevata obsolete carinatus. Elytra thorace tertia parte longiora, confertissime subtilissimeque punctata. Abdomen confertissime subtilissimeque punctatum. Pedes dilute flavo-testacei, tarsis posterioribus modice elongatis, anticis in utroque sexu leviter dilatatis.

Mas abdominis segmento inferiore quinto apice late subsinuato, sexto fortiter triangulariter exciso; femina eodem apice leviter producto rotundatoque.

\* 24. *L. obsoleta*: Omnium subtilissime confertissimeque punctata, opaca, nigro-picea, capite suborbiculato, thorace quadrato, aequali, pedibus antennisque rufo-piceis, his basi fuscis. — Long.  $1\frac{2}{3}$  lin.

Er. Col. March. I. 516. 8.

*Lathrobium obsoletum* Nordm. Symb. 146. 18.

Habitat in Germania.

Statura et summa similitudo praecedentis, nigro-picea, opaca, pube subtili griseo-sericea densius vestita, capite thoraceque lateribus parce nigro-pilosellis. Antennae capitis thoracisque longitudine, apicem versus neque attenuatae, neque incrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et paulo magis obconicis, ultimo ovato, apice acuminato, rufo-piceae, basi fuscae. Palpi rufo-picei, articulo tertio fusco. Labrum apice obsolete denticulatum. Caput thorace paulo angustius, basi, lateribus et angulis posterioribus leviter rotundatis, ante oculos subito angustatum, leviter convexum, subtilissime confertissimeque punctatum, ore rufo-piceo. Thorax coleopteris paulo angustior, latitudine haud longior, lateribus rectis, parallelis, basi apiceque truncatus, angulis anterioribus leviter, posterioribus fortius rotundatis, subdepressus, aequalis, confertissime subtilissimeque punctatus. Elytra thorace tertia parte longiora, subtilissime confertissimeque punctata. Abdomen omnium subtilissime punctulatum. Pedes rufo-picei vel rufo-testacei, tarsis posterioribus modice elongatis.

Mas abdominis segmento inferiore sexto apice leviter emarginato, tarsi anticis fortiter dilatatis; femina abdominis segmento inferiore sexto apice rotundato, tarsi anticis leviter dilatatis.

Variat thorace, elytris anoque rufo-piceis, rarius antennis palpisque totis rufo-testaceis.

Obs. A praecedente, cui varietates dilutiores colore etiam similes sunt, differt thorace latiore, angulis anticis minus obtusis, supra ne obsolete quidem carinato, capiteque subtilius punctatis, antennarum basi, palporumque max. articulo tertio plerumque in dilutioribus etiam speciminibus piceis, tarsi anticis maris fortiter dilatatis, patellam ovalem formantibus.

\* 25. *L. obscurella*: Omnium subtilissime confertissimeque punctata, opaca, fusco-picea, antennis pedibusque testaceis, capite subtriangulari, thorace oblongo-subquadrato, aequali. — Long.  $1\frac{1}{2}$  lin.

Habitat in Sardinia, Dom. Prof. Gené.

Praecedenti proxima, opaca, tota omnium subtilissime confertissimeque punctata, pube subtilissima fusco-grisea densiore sericans. Antennae capite thoraceque paulo breviores, validiusculae, apicem versus neque incrassatae, neque attenuatae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus, ultimo subovato, apice subito acuminato, totae testaceae. Os testaceum. Labrum apice medio acute bidenticulatum. Caput basi thoracis latitudine, basi et angulis posterioribus leviter rotundatis, antrorsum sensim angustatum, supra parum convexum, nigrum. Thorax coleopteris paulo angustior, latitudine paulo longior, basin versus subangustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, posterioribus leviter rotundatis, parum convexus, aequalis, fusco-piceus. Elytra thorace paulo longiora, subdepressa, fusco-picea. Abdomen nigrum, segmentorum ventralium margine apicali rufo-piceo. Pedes dilute testacei, tarsis posterioribus leviter elongatis, anticis leviter dilatatis.

Obs. Distinguitur a praecedente praecipue capite antrorsum magis angustato, thorace latitudine longiore, elytris contra paulo brevioribus.

- \* 26. *L. nigritula*: *Linearis, subtilius punctata, nigra, antennis pedibusque piceis, capite suborbiculato, thorace subquadrato.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Sicilia, Dom. Dr. Helfer.

Statura fere praecedentium, at magis linearis, nigra, subnitida, subtiliter cinereo-pubescent. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo minore, 3 — 10 sensim paulo brevioribus, ultimo apice subito acuminato, rufo-piceae. Os rufo-piceum. Labrum apice medio obsolete bidenticulatum. Caput thoracis latitudine, basi, lateribus et angulis posterioribus leviter rotundatis, parum convexum, confertim subtilius punctatum, linea media longitudinali laevi parum distincta. Thorax coleopteris vix angustior, latitudine paululum longior, lateribus rectis, parallelis, apice utrinque oblique truncatus, angulis anterioribus obtusis, posterioribus subrotundatis, parum convexus, subtilius confertim punctatus, linea media longitudinali laevi, posterius subelevata. Elytra thorace paulo longiora, subtilius confertim punctata, nigro-picea. Abdomen confertim subtiliter punctatum, segmentorum margine apicali anoque piceis. Pedes saturate rufo-testacei vel rufo-picei, tarsi posterioribus minus elongatis, anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem rotundato.

Obs. A praecedentibus praecipue punctura multo minus subtili distinguenda.

- \* 27. *L. debilis*: *Subdepressa, rufo-testacea, subtilissime confertissimeque punctata, capite thoraceque subquadratis, elytris fuscis, apice pedibusque flavo-testaceis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Linearis, subdepressa, rufo-testacea, opaca, pube subtilissima cinerea obducta. Antennae capite sesqui longiores, apicem versus neque attenuatae, neque incrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus, ultimo oblongo, apice abrupte acuminato, flavo-testaceae. Palpi testacei. Labrum apice subsinuatum, medio obsoletius bidenticulatum. Caput thoracis latitudine, lateribus rectis, basi truncatum, angulis posterioribus subrectis, subquadratum, supra posterius leviter convexum, anterius subdepressum, omnium subtilissime confertissimeque punctatum, lateribus parce pilosellum. Thorax coleopterorum latitudine, latitudine haud brevior, lateribus rectis, parallelis, apice utrinque oblique truncatus, angulis anterioribus obtusis, posterioribus rotundatis, subdepressus, subtilissime confertissimeque punctatus, linea media longitudinali, basin versus obsoletissima, vix observanda. Elytra thorace sesqui longiora, subtilissime confertissimeque punctata, nigro-fusca, apice determinate testacea. Abdomen coleopteris angustius, dense punctulatum, segmentis singulis nonnunquam basi fusciscentibus. Pedes flavi, breviusculi, femoribus posticis leviter incrassatis, tarsi posterioribus gracilioribus, leviter elongatis, anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore quinto apice levissime, sexto leviter emarginato; femina eodem rotundato.



- \* 28. *L. humilis*: *Brevis, subdepressa, subtilissime confertissimeque punctata, rufo-picea, capite transversim subquadrato, basi emarginato, thorace obsolete carinato, basi subtilissime canaliculato, antennis tenuibus, articulis duobus primis crassis.* — Long.  $1\frac{3}{4}$  lin.

Habitat in Columbia, Dom. Moritz.

Statura brevis, latiuscula *L. corticinae*, at quadruplo minor, rufo-picea, pube subtilissima depressa grisea vestita, capite thoraceque nigricantibus, lateribus nigro-pilosellis. Antennae capite paulo longiores, apicem versus attenuatae, articulis duobus primis crassiusculis, sequentibus tenuibus, tertio secundo vix brevior, 4 — 10 sensim paulo brevioribus, ultimo oblongo, acuminato, pilosellae, rufo-testaceae. Labrum margine dilatatum, medio bidenticulatum. Caput thorace latius, transversim subquadratum, basi late emarginatum, angulis posterioribus vix rotundatis, depressiusculum, subtilissime aequaliter punctatum. Thorax coleopteris angustior, latitudine antica tertia parte brevior, basin versus modice angustatus, lateribus rectis, apice utrinque oblique truncatus, angulis anterioribus obtusis, modice prominentibus, posterioribus rotundatis, subdepressus, confertissime subtilissimeque punctatus, linea media longitudinali, subelevata, tenui, antrorsum subobsoleta, basi subtiliter canaliculata. Elytra thorace tertia parte longiora, subtiliter dense punctata. Abdomen confertissime subtilissimeque punctatum. Corpus subtile dilute rufo-piceum. Pedes breviusculi, testacei, tarsis posterioribus leviter elongatis, anticis parum dilatatis.

Mas latet. Femina abdominis segmento inferiore sexto apice leviter producto, emarginato.

Obs. Antennarum structura cum sequentibus duabus a praecedentibus facile distinguenda.

- \* 29. *L. sordida*: *Subtilissime confertissimeque punctata, rufo-testacea, subnitida, subdepressa, capite subquadrato, thorace subtilissime canaliculato, antennis tenuibus, articulis duobus primis crassis.* — Long.  $1\frac{1}{4}$  lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

*L. ochracea* minor, praecipue brevior, praecedente plus duplo minor, subdepressa, rufo-testacea, subnitida, pube subtilissima grisea vestita, capite thoraceque lateribus parce nigro-pilosellis. Antennae capite sesqui longiores, articulis duobus primis crassis et sat magnis, sequentibus tenuibus, 3 — 10 sensim paulo brevioribus, ultimo oblongo, apice acuminato, pilosellae, testaceae. Palpi testacei. Labrum apice medio obsolete bidenticulatum. Caput thorace prope latius, basi truncatum, angulis posterioribus et lateribus rectis, ante oculos subito angustatum, subdepressum, aequale. Thorax coleopteris paulo angustior, latitudine antica tertia parte brevior, basin versus angustatus, apice utrinque oblique truncatus, angulis anterioribus obtusis, leviter prominentibus, posterioribus rotundatis, parum convexus, linea media longitudinali exarata subtilissima, apice evanescente canaliculatus. Elytra thorace tertia parte longiora, angulo-apicali exteriori dilutiori, dorso-plaga obsoleta fuscescente. Abdomen densius pubescens. Pedes flavo-testacei, breviores, tarsis posterioribus brevioribus, anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore sexto apice profundius triangulariter exciso; femina eodem rotundato.

- \* 30. *L. exilis*: *Omnium subtilissime confertissimeque punctata, rufo-testacea, capite suborbiculato, nigro thoraceque subquadratis, pedibus flavis, antennis tenuibus, articulis duobus primis crassis. — Long. 1 lin.*

Habitat in Columbia, Dom. Moritz.

*L. ochracea* habitu similis at quadruplo fere minor, rufo-testacea, parum nitida, tota subtilissime confertissimeque punctulata, pube subtilissima cinerea vestita, capite solo nigro-piceo. Antennae capite thoraceque paulo breviores, articulis duobus primis crassis ac reliquis multo maioribus, his tenuibus, 3 — 10 subaequalibus, ultimo oblongo, apice abrupte acuminato, pilosellae, flavo-testaceae. Palpi testacei. Os rufo-testaceum. Labrum apice leviter sinuatum, medio acute bidentatum. Caput thoracis latitudine, suborbiculatum. Thorax coleopteris paulo angustior, longitudine haud brevior, basin versus vix angustatus, lateribus rectis, angulis omnibus rotundatis, parum convexus, aequalis. Elytra thorace paulo longiora, dilutius testacea, plaga dorsali indeterminata obsoletiore fusca. Abdomen apice pilosellum. Pedes flavi, breviores, femoribus anticis posticisque leviter incrassatis, tarsis posterioribus tenuibus, minus elongatis, anticis in utroque sexu simplicibus.

Mas abdominis segmento inferiore sexto apice late triangulariter exciso, septimo exserto; femina sexto apice rotundato, septimo occulto.

### **Ophites Er.**

*Labrum breve, apice bidenticulatum.*

*Palpi maxillares articulo quarto distincto, acuto.*

*Antennae fractae.*

*Tarsi articulo quarto simplice.*

Corpus elongatum, lineare, alatum. Caput oblongum, postice attenuatum, collo quasi longo tenui thoraci affixum, exsertum, porrectum, oculis parvis, rotundatis, prominulis. Labrum transversum, breve, medio sinuatum et bidenticulatum. Mandibulae falcatae, acutae, medio fortiter acuteque dentatae. Maxillae (t. IV. f. 32. b.) malis corneis, interiore interne apiceque, exteriore subelongata apice dense barbatis. Palpi maxillares elongati, articulo primo brevi, secundo valde elongato, tenui, tertio huic aequali, apicem versus sensim subincrassato, quarto parvo, subulato, acuto. Labium (t. IV. f. 32. a.) mento transversum, ligula lata, biloba, lobis rotundatis, approximatis, membranacea, paraglossis angustis, acuminatis, coriaceis, pubescentibus, ligulam haud superantibus. Palpi labiales filiformes, articulo primo secundo paulo brevior, tertio huic aequali, acuminato. Antennae tenues, elongatae, fractae, articulo primo valde elongato, recto, secundo reliquis tenuiore et brevior, sequentibus sensim paulo brevioribus, ultimo apice truncatulo. Thorax angustus, fortiter elongatus, cylindricus, a medio apicem versus sensim attenuatus. Scutellum triangulare. Elytra truncata. Abdomen lineare, segmento superiore sexto apice trisinuato, septimo retractili, stylis analibus distinctis, longioribus, acuminatis. Pedes elongati, graciles, coxis anticis elongatis, tarsis omnibus simplicibus, articulis 4 primis longitudine sensim decrescentibus.

### **Observationes.**

1. Habitus in hac familia singularis, capitis thoracisque forma Casnoniam referens, neque etiam Raphidia dissimilis, unde a Dejeanio nomen Rugili

*Raphidioidis* speciei cuiusdam impositum est. Genus a *Stilico* oris fabrica, quam *Paedero* magis convenit, satis distinctum, a *Paedero* tarsorum articulo quarto simplice, ab omnibus huius familiae generibus, excepto *Cryptobio*, antennarum forma differt, ab hoc autem habitu, labri, palporum pedumque structura satis discedit.

2. Sexus differentia in abdominis segmento inferiore, in maribus inciso, in femina integro, rotundato, quaerenda.

\* 1. *O. versatilis*: *Nigro-piceus, nitidus, elytris thoracis nitidi longitudine, basi subseriatim punctatis, margine apicali pedibusque albidis, his medio piceis.* — Long.  $4\frac{1}{2}$  lin

T. 2. f. 1.

Habitat in Columbia, Dom. Moritz.

Niger, nitidus. Antennae nigro-piceae, articulo primo basi rufescente, ultimis ferrugineis. Palpi maxillares rufo-picei, articulo tertio medio nigricante. Os rufo-piceum. Caput nitidum, parcius subtiliter punctulatum, pube parciore minus depressa nigra tenuiter vestitum, posterius attenuatum, pedunculo piceo. Thorax coleopteris duplo angustior, elongatus, cylindricus, postice utrinque compressus, a medio inde antrorsum sensim attenuatus, utrinque parce subtiliterque subseriatim punctatus, nitidus, pube parca erecta instructus. Scutellum piceum, laeve. Elytra thoracis longitudine, basi interioris fortiter densius seriatim, exterius apiceque parcius vage punctata, parce griseo-pubescentia et nigro-pilosa, nigro picea, nitida, margine apicali pallide flavo. Abdomen parcius subtilissime punctulatum, pube depressa cinerea tenuiter vestitum, apice piceo-testaceum, segmento superiore sexto apice acute bidentato. Pedes albidus, coxis nigro-piceis, antici femoribus subtus tibiisque totis piceis, posteriores femoribus apice late nigro-piceis, tibiis etiam plerumque basi piceis.

Mas abdominis segmento inferiore sexto profunde triangulariter exciso; femina eodem rotundato.

\* 2. *O. Raphidioides*: *Niger, nitidus, elytris thorace paulo longioribus, basi seriatim punctatis, margine apicali pedibusque albidis, femoribus apice tibiisque anticis piceis.* — Long.  $4\frac{1}{2}$  lin.

Habitat Carthagenae in Columbia, Dom. Gory.

Praecedenti proximus, at paulo robustior. Antennae ferrugineae, articulo primo apice piceo. Palpi maxillares rufo-testacei, articulo tertio medio piceo. Os nigro-piceum. Caput confertim subtilissime punctatum, pube subtili depressa cinerea dense obductum, posterius attenuatum, pedunculo piceo. Thorax coleopteris duplo angustior, elongatus, cylindricus, postice subcompressus, medio leviter incrassatus, antrorsum sensim attenuatus, utrinque parcius subtiliter subseriatim punctatus, nitidus, pilis parvis erectis nigris instructus. Scutellum nigrum, laeve. Elytra thorace paulo longiora, basi densius seriatim punctata, seriebus interioribus longioribus, apice parce vage punctata, parce subtiliterque cinereo-pubescentia et nigro-pilosa, nigra, nitida, margine apicali pallide flavo. Abdomen confertissime subtilissimeque punctatum, pube subtilissima cinerea densius vestitum, apice piceo-testaceum, segmento sexto dorsali apice acute bidentato. Pedes albi, coxis omnibus nigris, femoribus apice late nigricantibus, tibiis anticis nigro-



piceis, posterioribus albis, apice subtestaceis, tarsis testaceis, posterioribus basi albis.

Mas abdominis segmento inferiore sexto apice emarginato; femina latet.

Obs. Distinguendus a praecedente antennis brevioribus, dilutioribus, capite latiore, pube densa cinerea vestito, thorace medio leviter incrassato, elytris longioribus, abdomine confertissime punctulato, densius pubescente, tibiis posterioribus basi haud nigricantibus.

3. *O. velitaris*: *Niger, capite thoraceque opacis, elytris thorace longioribus, aequaliter punctatis, margine apicali pedibusque albidis, his medio piceis.* — Long.  $4\frac{1}{4}$  lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Statura omnino et summa affinitas praecedentium, at vere distinctus. Corpus nigrum. Antennae rufo-piceae, articulo primo apice fusco. Palpi maxillares rufo-picei, articulo secundo tertioque medio fuscescentibus. Os piceum. Caput opacum, omnium confertissime subtilissimeque punctato-alutaceum, pube subtili depressa grisea obductum. Thorax coleopteris duplo angustior, elongatus, cylindricus, pone medium lateribus compressus, a medio inde apicem versus sensim leniter angustatus, glaber laevisque, opacus, linea media longitudinali, ante medium latiore, pone medium interrupta lateribusque pone medium nitidis. Scutellum laeve, opacum. Elytra thorace longiora, aequaliter densius, profundius et minus fortiter, basi etiam vix seriatim punctata, pube parca depressa grisea instructa, nigro-picea, nitida, margine apicali albido-testaceo. Abdomen confertim subtiliter punctatum, pube densiore subtili depressa grisea vestitum, subnitidum, segmento quinto sextoque apice testaceis, sexto superiore apice obtuse bidentato. Pedes coxis omnibus piceis, trochanteribus testaceis, femoribus albis, apice late nigro-piceis, posterioribus apice late testaceis, tarsis omnibus testaceis.

Mas latet. Femina abdominis segmento inferiore sexto apice rotundato.

### **Stilicis Latr.**

*Latr. Regn. Anim. IV. p. 436. — Encycl. meth. X. p. 494. — Er. Col. March. I. p. 519. Rugilus Curt. Britt. Ent. IV. t. 168. — Steph. Illustr. Britt. Ent. V. p. 277. — Boisd. et Lacord. Faun. Ent. Paris. I. p. 433.*

*Labrum amplum, apice bispinosum.*

*Palpi maxillares articulo quarto vix distincto.*

*Antennae rectae.*

*Tarsi articulo quarto simplice.*

Corpus elongatum, alatum. Caput sat magnum, orbiculatum, pedunculo tenui brevi thoraci affixum, exsertum, porrectum, oculis parvis, rotundatis, prominulis. Labrum amplum, mandibulas obtegens, lateribus rotundatum, limbo laterali membranaceo, medio subsinuaturn, acute bidenticulatum. Mandibulae falcatae, acutae, medio acute 3- vel 4-dentatae. Maxillae (t. IV. f. 31. b.) malis corneis, interiore interne, exteriori apice dense barbatis. Palpi maxillares modice elongati, articulo primo parvo, secundo tertioque elongatis, longitudine subaequalibus, hoc incrassato, quarto subulato, minimo, aegre conspicuo. Labrum (t. IV. f. 31. a.) mento transversum, ligula mem-

branacea biloba, lobis rotundatis, brevissimis, distantibus, interstitio spisse setoso, paraglossis linearibus, membraneis, ligulam longe superantibus. Palpi labiales articulo primo cylindrico, secundo subovato, praecedente paulo longiore crassioreque, tertio minuto, tenui, acuto. Antennae filiformes, rectae, brevisculae crassiusculaeque, articulo primo maiusculo, longiore, secundo tertioque sequentibus paulo longioribus. Thorax subovatus, apice attenuatus. Scutellum parvum, triangulare. Elytra truncata. Abdomen lineare, segmento septimo retractili, stylis analibus haud distinctis. Pedes sat longi, graciles, coxis anticis elongatis, tarsis omnibus simplicibus, anticis articulis omnibus in utroque sexu linearibus.

### Observationes.

1. Corpus obscurum, subtilissime pubescens, capite thoraceque confertissime punctatis, subrugulosis, hoc magis minusve distincte carinato. Capite lato, rotundato, a thorace remoto, hoc angustiore, apice attenuato, pedibus longioribus gracilioribusque habitum peculiarem prae se fert.

2. Nomen a Latreillio huic generi impositum, a Com. Dejean Cryptobiis Americanis tributum, restitui, qua de re conf. quod l. c. monui.

3. Sexus differentia abdominis segmentis inferioribus ultimis cognoscenda, segmento sexto in femina integro, rotundato, in mare apice exciso, nonnunquam etiam quinto fortiter impresso vel excavato.

4. Victus sub lapidibus et sub foliis deciduis.

✕ \* 1. *St. fragilis*: Niger, thorace oblongo, obsolete carinato, scutello pedibusque anticis sanguineis, elytris dense subtiliter punctatis, apice pallidis. — Long. 3 lin.

*Enc. meth. X. p. 495. — Er. Col. March. I. 520. 1.*

*Paederus fragilis* Grav. Mon. 110. 7. — Latr. Hist. nat. Crust. et Ins. IX. 347.

4. *Gen. Crust. et Ins. I. 292. 2. — Oliv. Encycl. meth. VIII. 629. 8.*

*Rugilus fragilis* Curtis Britt. Ent. IV. t. 168. — Boisd. et Lacord. Faun. Ent. Paris. 1. 434. 2. — Germ. Faun. Ins. Europ. 18. 5. — Runde Brachelytr. Hal. 14. 2.

*Staph. angustatus* Fourc. Ent. Par. I. 172. 28.

Habitat in Gerania.

*St. rufipede* paululum maior et gracilior, niger, parum nitidus. Antennae capite thoraceque longe breviores, articulo tertio secundo sesqui longiore, sequentibus sensim paulo brevioribus et crassioribus, ultimo ovato, apice acuminato, basi nigro-piceae, extrorsum obscure testaceae. Palpi maxillares rufo-piceae, articulo tertio nigro, apice pallido. Labrum latum, amplum, nigrum, margine piceum, medio acute bidenticulatum. Caput coleopterorum latitudine, orbiculatum, confertissime subtiliter punctato-rugulosum, tenuiter subtilissime griseo-pubescens; opacum, nigrum. Thorax capitis prope longitudine, coleopteris duplo angustior, a medio antorsum attenuatus, basin versus angustatus, convexus, rufo-sanguineus, opacus, subtilius crebre punctatus, tenuiter subtilissime griseo-pubescens, linea longitudinali media angusta laevi, haud elevata, subtiliter canaliculata. Scutellum punctulatum, rufum. Elytra thorace paulo longiora, subtiliter crebre punctata, densius subtiliter cinereo-pubescentia, subnitida, nigro-picea, extrorsum summo apice pallide testaceo. Abdomen subtilissime confertissimeque punctulatum, dense subtiliter cinereo-pubescens. Pectus nigrum, mesosterno rufo.

Pedes antici cum coxis rufi, intermediis coxis femoribusque rufis, tibiis piceis, tarsis testaceis, postici picei, tarsis testaceis.

Mas abdominis segmento inferiore quarto apice medio tuberculo minuto prominulo rufo, ciliarum longarum arcu cineto munito, quinto medio profunde excavato, fovea laevis, marginibus eius lateralibus acutis, carnatis, nigro-ciliatis, apice prominulis, sexto inciso.

\* 2. *St. rufipes*: *Niger*, thorace obsolete carinato, antennis pedibusque rufo-piceis, elytris subtiliter punctatis, apice piceis. — Long.  $2\frac{3}{4}$  lin.

Er. Col. March. I. 521. 2.

*Rugilus rufipes* Germ. Faun. Ins. Europ. 18. 4.

*Paed. orbiculatus* Fab. Ent. Syst. I. II. 538. 9. Syst. El. II. 603. 91

*Staph. orbiculatus* Marsh. Ent. Brit. 528. 85.

*Rugilus orbiculatus* Boisd. et Lacord. Faun. Ent. Paris. I. 434. 1.

Habitat in Germania, Gallia.

*Niger*, parum nitidus, capite thoraceque opacis, subtiliter breviterque pubescentibus. Antennae capite sesqui longiores, apicem versus haud incrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim brevioribus, non autem crassioribus, rufae, apice magis testaceae. Palpi rufo-picei. Labrum rufescenti-piceum, apice medio acute bidentatum. Mandibulae piceae. Caput coleopterorum latitudine, orbiculatum, confertissime minus subtiliter punctatum, longitudinaliter subrugulosum. Thorax coleopteris dimidio angustior, latitudine parum longior, basin versus leviter angustatus, modice convexus, confertissime minus subtiliter punctatus, subrugosus, linea longitudinali media tenui laevi, subtiliter canaliculata, minus distincta, antrosum obsoleta. Elytra thorace paulo longiora, subtiliter parcius punctata, tenuiter cinereo-pubescentia, fusco-picea, nitidula, apice sensim subdilutiora. Abdomen subtilissime dense punctatum, pube subtili cinerea subsericans, nigrum, subnitidum, apice piceum. Pedes piceo-rufo-testacei, coxis saturatius piceis.

Mas abdominis segmento inferiore sexto apice inciso.

\* 3. *St. subtilis*: *Niger*, thorace subcarinato, antennis pedibusque rufo-testaceis, femoribus posterioribus apice fuscescentibus, elytris apice concoloribus. — Long.  $2\frac{3}{4}$  lin.

Habitat in Austria, Dom. Schüppel, in Saxonia montana, Dom. Märkel.

Statura omnino praecedentis, niger, capite thoraceque opacis, subtiliter densius griseo-pubescentibus. Antennae capite paulo longiores, apicem versus sensim leviter incrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, rufae. Mandibulae et palpi piceo-rufi. Labrum piceum, medio acute bidentatum. Caput coleopterorum latitudine, subovatum, omnium confertissime subtilius punctato-subrugosum. Thorax coleopteris dimidio angustior, latitudine longior, lateribus basin versus modice angustatus, leviter convexus, omnium confertissime subtilius punctato-rugulosus, linea media longitudinali laevi subelevata, subtilissime canaliculata, apicem versus evanescente. Elytra thorace paulo longiora, densius subtiliusque punctata, piceo-nitidula, apice concolora. Abdomen confertim subtiliter punctatum, segmentis ultimis duobus apice piceis. Pedes



rufo-testacei, coxis omnibus nigris, femoribus posterioribus apice fuscescentibus.

Mas abdominis segmento inferiore sexto apice triangulariter exciso.

Obs. 1. Differt a praecedente praecipue staturā graciliore, capite thoraceque magis oblongis, multo subtilius punctato-rugulosis, thoracis linea laevi distinctiore, geniculis posterioribus infuscatis, a sequentibus thoracis linea media longitudinali minus elevata, elytris apice concoloribus.

2. Nomine *Paederi subtilis* a Dahlio Dom. Schüppel missus.

\* 4. *St. similis*: *Niger, thorace carinato, antennis pedibusque rufo-testaceis, elytris crebre punctatis, apice testaceis.* — Long.  $2\frac{1}{2}$  lin.

*Er. Col. March. I.* 521. 4.

*Paederus orbiculatus* Grav. *Micr.* 63. 6. *Mon.* 141. 8.

*Rugilus orbiculatus* Runde *Brachelytr.* Hal. 14. 1.

Habitat in Germania.

Praecedentibus paulo minor, niger, capite thoraceque opacis, subtiliter breviterque pubescentibus. Antennae capite parum longiores, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, rufae. Palpi testacei. Labrum amplum, acute bidenticulatum, piceum, margine rufescens. Mandibulae piceae. Caput coleopterorum latitudine, orbiculatum, confertissime punctatum, longitudinaliter subrugulosum. Thorax coleopteris dimidio fere angustior, latitudine vix longior, basin versus leviter angustatus, modice convexus, confertissime punctatus, subrugulosus, linea longitudinali media angusta subelevata laevi, per totam longitudinem canaliculata. Elytra thorace paulo longiora, profundius fortius crebre punctata, griseo-pubescentia, nitidula, saturate nigro-picea, margine apicali testaceo. Abdomen nigrum, subnitidum, confertim subtilissime punctatum, pube subtili depressa grisea subsericans. Pedes rufo-testacei, geniculis concoloribus, coxis omnibus nigro-piceis.

Mas abdominis segmento inferiore quinto semicirculariter impresso, impressione apice utrinque denticulo minuto prominulo terminata, sexto medio leviter impresso, apice acute emarginato.

Obs. Nomine *Paederi analis* a Dahlio Dom. Schüppel communicatus.

\* 5. *St. geniculatus*: *Niger, thorace carinato, antennis pedibusque testaceis, femoribus posterioribus apice nigrificantibus, elytris crebre punctatis, apice testaceis.* — Long.  $2\frac{3}{4}$  lin.

*Er. Col. March. I.* 522. 4.

Habitat in Germania.

Statura et summa affinitas praecedentis, niger, capite thoraceque opacis, breviter subtiliterque pubescentibus. Antennae capite parum longiores, apicem versus leviter incrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim brevioribus et crassioribus, ultimo breviusculo, apice acuminato, rufo-testaceae. Palpi rufo-testacei. Labrum amplum, apice acute bidenticulatum, piceum, margine rufescens. Mandibulae rufo-piceae. Caput coleopterorum latitudine, orbiculatum, confertissime punctatum, longitudinaliter subrugulosum. Thorax coleopteris vix dimidio angustior, latitudine

perparum longior, basin versus leviter angustatus, modice convexus, confertissime punctatus, subrugulosus, linea longitudinali media tenui laevi, subtiliter per totam fere longitudinem canaliculata. Elytra thorace paulo longiora, fortius crebre punctata, griseo-pubescentia, nigro-picea, nitidula, margine apicali piceo-testaceo. Abdomen confertim subtilissime punctatum, nigrum, nitidulum, pube subtili depressa grisea sericans. Pedes dilute testacei, coxis omnibus piceis, femoribus posterioribus apice fuscis.

Mas abdominis segmento inferiore quinto late profundeque impresso, impressione apice utrinque spina dentiformi, acuta, retrorsum prominula terminata, sexto leviter impresso, apice acute emarginato.

Obs. Proximus praecedenti et femina nisi elytris paulo longioribus et geniculis pedum posteriorum infuscatis vix distinguenda, mas praeterea abdominis segmento inferiore quinto apice spinis duabus longioribus armato diversus.

\* 6. *St. affinis*: *Niger, thorace carinato, pedibus testaceis, femoribus posterioribus apice tibiisque fuscis, elytris subtiliter punctatis, apice testaceis.* — Long.  $2\frac{1}{4}$  lin.

*Er. Col. March. I. 522. 5.*

Habitat in Germania, Gallia.

Habitus et summa affinitas praecedentium, at paulo minor, niger, capite thoraceque opacis, breviter subtiliterque pubescentibus. Antennae capite paulo longiores, apicem versus subincrassatae, articulo tertio secundo sesqui prope longiore, 4 — 10 sensim brevioribus et crassioribus, rufo-piceae. Palpi maxillares rufo-picei, articulo tertio fusciscente. Labrum amplum, apice acute bidenticulatum, piceum, margine rufescente. Caput vix coleopterorum latitudine, orbiculatum, confertissime fortius punctatum, longitudinaliter rugulosum. Thorax coleopteris duplo fere angustior, latitudine paulo longior, basin versus leviter angustatus, modice convexus, confertissime fortius punctatus, subrugulosus, linea longitudinali media latiore laevi, subelevata, subtiliter canaliculata, canalicula antrorsum evanescente. Elytra thorace paulo longiora, parcius subtiliter punctata, parce subtiliterque cinereo-pubescentia, piceo-subaenescencia, nitida, margine apicali testaceo. Abdomen confertim subtiliter punctatum, densius subtiliter griseo-pubescent, nigro-subaeneum, nitidulum. Pedes testacei, coxis omnibus piceis, femoribus apice tibiisque magis minusve saturate fuscis.

Mas abdominis segmento inferiore sexto apice inciso.

Obs. Caput et thorax fortius, elytra parcius et subtilius quam in praecedentibus punctata.

\* 7. *St. fuscipes*: *Niger, thorace carinato, pedibus fusco-piceis, tarsis testaceis, elytris parce subtilissime punctatis, apice testaceis.* — Long.  $2\frac{1}{4}$  lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura praecedentis, niger, parum nitidus, capite thoraceque opacis. Antennae capite sesqui longiores, apicem versus sensim subincrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim brevioribus et paulo crassioribus, rufo-piceae, articulo primo nigro. Palpi maxillares nigro-picei. Labrum nigro-piceum, apice subtiliter bidenticulatum. Caput coleopterorum

latitudine, orbiculatum, confertissime fortius punctatum, fronte spatio minuto medio laeviore, tenuiter subtiliterque griseo-pubescentia, parce nigro-pilosum. Thorax coleopteris duplo angustior, latitudine longior, apicem versus magis quam in praecedentibus attenuatus, modice convexus, confertissime fortius punctatus, subtilissime tenuiter griseo-pubescentia, lateribus parce nigro-pilosus, linea media longitudinali lata laevis subelevata, summa basi obsolete canaliculata. Elytra thorace parum longiora, parce subtiliter obsolete punctata, tenuiter griseo-pubescentia, piceo-nigra, nitida, margine apicali testaceo. Abdomen nitidulum, subtiliter punctulatum, densius subtiliter griseo-pubescentia. Pedes crassiusculi, fusco-picei, trochanteribus, summa tibiatarum basi apiceque tarsisque rufescenti-testaceis.

Mas abdominis segmento inferiore sexto apice triangulariter exciso.

\* 8. *St. orbiculatus*: *Niger, thorace carinato, antennis pedibusque cum coxis rufo-testaceis, elytris subtiliter punctatis, apice piceis.* — Long. 2 lin.

*Encycl. meth. X. p. 495. — Er. Col. March. I. 523. 6.*

*Staph. orbiculatus Payk. Mon. Staph. 35. 26.*

*Paed. orbiculatus Payk. Faun. Suec. III. 431. 6. — Walken. Faun. Paris. I. 275.*

*2. — Latr. Hist. nat. Crust. et Ins. IX. 346. 3. Genera Crust. et Ins. I. 291. 1.*

*— Oliv. Encycl. meth. VIII. 628. 9. — Gyll. Ins. Suec. II. 374. 3. — Zetterst.*

*Faun. Lappon. I. 85. 1. Ins. Lappon. 69. 3. — Sahlb. Ins. Fenn. I. 343. 3.*

*Rugilus orbiculatus Mannerh. Brachelytr. 40. 1.*

Habitat in Germania, Suecia.

Praecedentibus minor, niger, capite thoraceque opacis, subtiliter pubescentibus. Antennae capite vix longiores, apicem versus sensim subincrassatae, articulo tertio secundo haud longiore et paulo tenuiore, 4 — 10 sensim brevioribus et crassioribus, rufae. Palpi rufo-testacei. Labrum acute bidentatum, fere bispinosum. Caput coleopteris latius, suborbiculatum, basi truncatum, confertissime punctatum, vix rugulosum, linea media longitudinali laevi obsoleta. Thorax coleopteris dimidio angustior, latitudine paulo longior, basin versus leviter angustatus, modice convexus, confertissime punctatus, subrugulosus, linea longitudinali media latiore laevi, subelevata, per totam longitudinem subtiliter canaliculata. Elytra thorace paulo longiora, subtiliter parcius punctata, subtilius cinereo-pubescentia, obscure fusco-picea, margine apicali testaceo. Abdomen confertim subtiliter punctatum, subtiliter griseo-pubescentia, subnitidum. Pedes toti rufo-testacei, coxis omnibus concoloribus.

Mas abdominis segmento inferiore sexto apice triangulariter exciso.

\* 9. *St. angularis*: *Piceus, capite thoraceque carinato rufescentibus, elytris angulis posterioribus pedibusque cum coxis testaceis.* — Long. 2 lin.

Habitat in Pennsylvania et in Carolina, Dom. Zimmermann.

Statura fere *St. affinis*, at paulo minor. Antennae capite paulo longiores, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus et crassioribus, piceo-rufae. Palpi rufo-testacei. Labrum piceum, apice medio subtiliter licet acute bidenticulatum. Caput coleopterorum latitudine, suborbiculatum, supra confertissime subtilius punctato-subrugulosum, densius subtiliter griseo-pubescentia, opacum, piceo-subrufescentia, subtus



parce punctatum, nitidum, rufo-piceum. Thorax coleopteris plus dimidio angustior, latitudine longior, basin versus leviter angustatus, apicem versus fortius attenuatus, modice convexus, crebre subtilius punctatus, linea media longitudinali laevis, rufo-piceus, subnitidus, tenuiter subtiliterque griseo-pubescent. Scutellum piceum. Elytra thorace paulo longiora, minus dense subtilius licet profunde punctata, tenuiter griseo-pubescentia, nitida, nigro-picea, margine lato apicali humerisque testaceis. Abdomen confertim subtiliter punctatum, dense subtiliter griseo-pubescent, nitidulum, nigro-piceum, supra segmentis ultimis duobus, subtus omnibus margine apicali testaceo. Pectus nigro-piceum. Pedes toti cum coxis omnibus testacei.

Mas abdominis segmento inferiore quinto apice emarginato, sexto triangulariter exciso.

10. *St. iugalis: Niger, thorace fortius carinato, antennis pedibusque cum coxis testaceis, elytris fortiter punctatis, apice testaceis. — Long. 2½ lin.*

Habitat prope Caracas Columbiae, Dom. Moritz.

Statura omnino praecedentium, niger, capite thoraceque opacis. Antennae capite paulo longiores, apice subincrassatae, articulo tertio secundo paulo longiore, 4 — 10 sensim brevioribus et crassioribus, testaceae. Palpi testacei, maxillares articulo ultimo apice fusciscent. Labrum amplum, apice medio obsoletius bidenticulatum, nigro-piceum. Caput coleopterorum latitudine, orbiculatum, minus subtiliter confertissime punctata-rugulosum, tenuiter subtiliterque griseo-pubescent, parce nigro-pilosellum. Thorax coleopteris duplo prope angustior, basin versus modice angustatus, apicem versus fortius attenuatus, convexus, confertissime minus subtiliter punctatus, subtilissime griseo-pubescent, parce nigro-pilosellus, linea media longitudinali modice elevata laevis. Elytra thorace vix longiora, leviter inaequalia, fortius profunde punctata, tenuiter nigro-pubescentia, nitidula, margine apicali dilute testaceo. Abdomen confertissime subtilissimeque punctulatum, subtiliter nigro-pubescent. Pedes graciliores, toti cum coxis omnibus testacei.

Mas abdominis segmento inferiore quinto apice trisinuato, sinu intermedio maiore, medio denticulo acuto rufescente instructo, sexto apice inciso.

\* 11. *St. carinatus: Niger, thorace fortius carinato, antennis testaceis, pedibus flavis, elytris fortiter profundeque punctatis. — Long. 2¼ lin.*

Habitat in Columbia, Dom. Moritz.

Statura omnino praecedentium, niger, capite thoraceque opacis. Antennae capite paulo longiores, apicem versus leniter incrassatae, articulo tertio secundo vix longiore, 4 — 10 sensim brevioribus et crassioribus, testaceae. Palpi testacei. Labrum amplum, apice medio subtilissime bidenticulatum. Caput coleopterorum fere latitudine, suborbiculatum, confertissime fortius punctatum, subtilissime nigro-pubescent, parce nigro-pilosellum. Thorax coleopteris dimidio angustior, latitudinis longior, basin versus modice angustatus, apice fortius attenuatus, modice convexus, confertissime fortius punctatus, parce griseo-pubescent, linea longitudinali media laevi, elevata. Elytra thorace vix longiora, fortiter profundeque punctata, parce pubescentia, nigro-subaenea, nitida. Abdomen nigro-subaeneum, nitidulum, subtiliter griseo-pubescent. Pedes flavi, coxis concoloribus.

Mas abdominis segmento inferiore quinto apice sinuato, sinu medio acute dentato, sexto profunde inciso.

Species mihi invisae.

12. *St. dentatus*: Niger, pedibus pallidis, elytris obscure rufis. — Long. ultra  $1\frac{1}{2}$  lin.

*Rugilus dentatus* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. p. 457.

Habitat in America septentrionali.

Nigro-piceus. Caput longitudinaliter rugulosum. Antennae et palpi picei. Labrum amplum, medio bidentatum, piceum, lateribus rotundatum. Thorax capite minor, longitudinaliter rugulosus, convexus, linea longitudinali laevi. Elytra confertim subtiliter punctata, subtiliter pubescentia, stria suturali impressa, margine apicali obscure rufo testaceove. Pedes pallide flavescens.

Variat margine elytrorum apicali suturam versus nigro, thoracis linea laevi fere obsoleta. St. orbiculato minor, capite thoraceque fortius distinctiusque rugulosus. (Say.)

### Echiaster Er.

*Labrum apice 4-dentatum.*

*Palpi maxillares articulo quarto vix distincto.*

*Antennae rectae.*

*Tarsi articulo quarto simplice.*

Corpus elongatum, subdepressum, alatum. Caput exsertum, pedunculo tenui brevissimo thoraci affixum, porrectum, suborbiculatum, oculis maiusculis, lateribus modice prominulis, infra fortius convexus. Labrum (t. II. f. 2. c.) transversum, mandibulas haud obtegens, apice fortiter acuteque 4-dentatum. Mandibulae longae, tennes, arcuatae, acutae, medio dente longo arcuato, interne subtilius dentato armatae. Maxillae (t. II. f. 2. b.) malis membranaceis, interiore interne, exterior apice barbatis. Palpi maxillares articulo primo brevi, secundo modice elongato, tertio hoc paulo longiore, incrassato, ultimo subulato, acuto, minutissimo, vix conspicuo. Labium (t. II. f. 2. a.) mento transversum, apice emarginato, angulis anterioribus in E. longicollis haud prominulis, in E. melanuro fortiter productis, ligula membranae, lobis brevissimis, leviter rotundatis, distantibus, interstitio setoso, paraglossis multo longioribus, linearibus, acuminatis, interne ciliatis. Antennae breves, capite vix longiores, articulis duobus primis crassiusculis, primo perparum elongato, secundo breviusculo, subgloboso, sequentibus sensim paulo brevioribus et crassioribus, ultimo subovato, apice subacuminato. Thorax elongatus, coleopteris angustior. Scutellum triangulare. Elytra depressa. Abdomen apicem versus sensim attenuatum, segmento quinto sextoque angustatis, elongatis, tubuliformibus, septimo retractili, stylis analibus haud exsertis. Pedes graciliores, breviusculi, tarsi articulis 4 primis longitudine decrescentibus, anticis simplicibus.

### Observationes.

1. Corpus punctura conferta scabrusculum, habitu fere Sunii, at tarsi arti-

culo quarto simplice, a Lithocharide palporum maxillarium articulo quarto minimo, a Stilico labro brevior, 4-dentato, ab omnibus abdominis segmentis ultimis duobus elongatis et attenuatis discedens.

2. Sexus differentia latet.

- \* 1. *E. longicollis*: Testaceus, opacus, punctato-scabriusculus, thorace elongato, apicem versus attenuato, abdomine segmentis ultimis duobus fuscis. — Long.  $2\frac{1}{4}$  lin.

T. II. f. 2.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Testaceus, opacus, dense fortius punctatus, interstitiis scabriusculis, pilis brevissimis reclinatis rigidis vestitus. Antennae capite paululum longiores. Caput latitudine longius, pone oculos leviter angustatum et rotundatum, supra depressiusculum. Thorax coleopteris plus duplo angustior, elongatus, latitudine summa duplo longior, a medio basin versus subangustatus, apicem versus attenuatus, subdepressus. Elytra thoracis longitudine, depressa, posterius infusca. Abdomen transversim subrugosum, segmentis ultimis duobus angustis, elongatis, fuscis, apice summo pilosum. Pedes dilute rufi-testacei.

- \* 2. *E. melanurus*: Testaceus, opacus, confertissime punctatus, thorace basin versus angustato, abdomine segmentis ultimis duobus nigris. — Long.  $1\frac{1}{2}$  lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Praecedente duplo minor et minus elongatus, testaceus, opacus, confertissime subtilius punctatus, pube brevissima depressa grisea fulvo-micante tenuiter vestitus. Antennae capite haud longiores. Caput suborbiculatum, thorace duplo latius, depressiusculum, antice utrinque leviter longitudinaliter impressum. Thorax coleopteris duplo angustior, latitudine summa sesqui longior, ante medium leviter rotundatus, basin versus leviter angustatus, apicem versus attenuatus, depressiusculus, obsolete carinatus. Elytra thorace paulo longiora, depressa, fusciscentia. Abdomen segmentis ultimis duobus nigris. Pedes pallide testacei.

Obs. Oris partes huius speciei T. II. f. 2. a. b. c. dilineatae sunt.

### **Sunius Leach.**

Er. Col. March. I. p. 523.

*Astenus* Dej. Boisd. et Lacord. Faun. Ent. Paris. I. p. 435. — Steph. Illustr. Brit. Ent. V. p. 275.

*Labrum medio bidenticulatum.*

*Palpi maxillares articulo quarto vix distincto.*

*Antennae rectae.*

*Tarsi articulo quarto lobo membraneo instructo.*

Corpus elongatum, lineare, plerumque alatum. Caput magnum, basi fortiter constrictum, collo quasi angusto brevi thoraci affixum, exsertum, porrectum, oculis parvis, rotundatis, leviter prominulis. Labrum latum, transversum, medio bidenticulatum. Mandibulae longae, tenues, arcuatae, acutae, medio dente longo acuto et pone dentem denticulis 2 vel 3 armatae. Ma-



xillae (t. IV. f. 33. b.) malis breviusculis, corneis, interiore interne, exteriore apice barbatis. Palpi maxillares leviter elongati, articulo primo minuto, brevi, secundo tertioque elongatis, longitudine subaequalibus, hoc incrassato, quarto subulato, subtilissimo, vix nisi oculo fortiter armato conspicuo. Labium (t. IV. f. 33. a.) mento brevi, transverso, ligula membranea, biloba, lobis rotundatis, sinu profundo distantibus, paraglossis angustis, acuminatis, membraneis interne ciliatis, ligulam parum superantibus. Palpi labiales articulo primo brevior, cylindrico, secundo maiore ovato, tertio minuto, tenui, subulato. Antennae filiformes, rectae. Thorax subcordatus, licet antice haud emarginatus, sed subito angustatus. Scutellum parvum, triangulare. Elytra truncata. Abdomen lineare, apice obtusum, segmento septimo retractili, stylis analibus haud distinctis. Pedes aliis breviusculi, aliis elongati, tarsis omnibus articulis tribus primis elongatis, longitudine decreascentibus, quarto brevissimo, subtus lobo membraneo aucto, quinto minuto, quantum parum excedente.

#### O b s e r v a t i o n e s.

1. Corpus plerisque opacum, punctura densiore fere asperatum, pube brevissima parca inspersum, a reliquis huius tribus generibus imprimis tarsorum structura distinctum.

2. Abdominis segmentum inferius sextum in mare triangulariter incisum, in femina integrum, rotundatum.

3. Victus sub lapidibus et foliis deciduis.

4. Sectiones in hoc genere praecipue secundum corporis habitum distinguendae videntur:

\* (Genuini.) Corpore valde elongato, lineari, subdepresso, pedibus breviusculis, labro transverso, capite oblongo, apice leviter emarginato, thorace depressiusculo, plerumque oblongo, coleopteris latitudine subaequali, abdomine elongato, saepius basin versus leviter angustato.

\*\* (Spurii.) Capite suborbiculato, apice truncato, labro plerumque medio leviter producto, thorace coleopteris angustiore, subgloboso, pedibus longioribus, habitu omnino Stilici, oris autem tarsorumque structura omnino cum genuinis congruentes.

#### \* (G e n u i n i).

\* 1. *S. filiformis*: *Elongatus, linearis, niger, antennis pedibusque testaceis, capite thorace latiore, thorace basin versus angustato, elytris thorace paulo longioribus, apice latius testaceis.* — Long.  $2\frac{1}{4}$  lin.

*Er. Col. March. I. 525. 3.*

*Paederus filiformis* Latr. *Gen. Crust. et Ins. I. 293. 4.* — Oliv. *Encycl. meth. VIII. 629. 10.*

*Paederus procerus* Grav. *Mon. p. 141.*

*Astenus procerus* Boisd. et Lacord. *Faun. Ent. Paris. I. 436. 1.*

*Paederus extensus* Mannerh. *Brachelytr. 39. 6.*

Habitat in Europa.

*S. angustato* parum latior, at multo longior, niger, opacus, pube brevi depressa fulvo-grisea vestitus. Antennae capite sesqui longiores palpique testacei. Mandibulae rufo-testaceae. Caput thorace paulo latius, oblongo-subquadratum, basi truncatum, lateribus pone oculos et angulis posteriori-

bus rotundatis, supra et infra subtiliter confertissime punctato-rugosum. Thorax coleopteris angustior, latitudine summa paulo longior, apice cum angulis anterioribus rotundatus, basin versus angustatus, subdepressus, subtiliter confertissime punctato rugulosus. Elytra thorace paulo longiora, dense minus fortiter at profunde punctata, apice late subsinuatum testacea. Abdomen elongatum, parallelum, crebrius punctulatum, subnitidum, subtus segmentis singulis margine apicali piceo. Pedes cum coxis anterioribus testacei.

Mas abdominis segmento inferiore sexto apice inciso.

\* 2. *S. prolixus*: *Elongatus, linearis, niger, antennis pedibusque testaceis, capite thorace latiore, thorace basin versus subangustato, elytris apice testaceis.* — Long.  $2\frac{1}{3}$  lin.

*Paederus cinctus* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV 457. 2. (*forte*).

Habitat in America septentrionali, Dom. Zimmermann.

Statura et summa affinitas praecedentis, niger, opacus, subtiliter cinereo-pubescent. Antennae tenues, capite plus sesqui longiores palpique flavi. Mandibulae et labrum rufo-testacea. Caput thorace parum latius, oblongo-subquadratum, basin versus ab oculis subangustatum, basi cum angulis posterioribus rotundatum, supra confertissime subtilius punctato-rugosum, opacum, subtus crebre fortiterque punctatum, nitidum. Thorax coleopteris vix angustior, latitudine paulo longior, lateribus ante medium subangulatus, basin versus leviter angustatus, subdepressus, confertissime rugoso-punctatus. Coleoptera thorace parum longiora, apicem versus leviter angustata, minus fortiter profunde dense punctata, interstitiis subrugulosis, apice testacea. Abdomen elongatum, basin versus subangustatum, profundius punctatum, nitidulum, segmentis omnibus margine apicali rufo-piceo. Pedes cum coxis anterioribus flavo-testacei.

Mas abdominis segmento inferiore sexto apice profunde inciso.

Variat elytrorum margine apicali anguste lateve testaceo, nonnunquam etiam elytris rufo-piceis, macula magna discoidali nigra, apice late testaceis, vel thorace etiam et abdominis segmentis 4 anterioribus rufo-piceis.

Obs. A praecedente differt praecipue pube longiore, antennis tenuioribus, thorace basin versus minus angustato, et capite subtus minus confertim fortiter punctato, coleopteris apicem, abdomine basin versus angustatis, etc.

\* 3. *S. linearis*: *Elongatus, niger, antennis pedibusque testaceis, capite thoracis latitudine, thorace basin versus subangustato, elytris thorace paulo longioribus, apice late testaceis.* — Long.  $1\frac{3}{4}$  lin.

Habitat in America septentrionali, Dom. Zimmermann.

Statura elongata et summa affinitas praecedentium, at minor, S. angustato aequalis, niger, opacus, pube subtili depressa cinerea vestitus. Antennae tenuiores, capite plus sesqui longiores, et palpi testacei. Mandibulae et labrum rufo-picea. Caput thoracis latitudine, oblongo-subquadratum, basi truncatum, angulis posterioribus fortiter rotundatis, supra confertissime subtiliter punctato-rugosum, opacum, subtus parcius profundiusque punctatum. Thorax coleopteris paulo angustior, latitudine paulo longior, lateribus

ante medium subangulato-rotundatus, basin versus subangustatus, subdepressus, subtiliter confertissime punctato-rugulosus, margine basali apicali-que piceo. Coleoptera apicem versus subangustata, thorace paulo longiora, crebrius fortiusque punctata, interstitiis subrugosis, subnitida, apice latius sinuatim testacea. Abdomen elongatum, basin versus sensim subangustatum, minus dense punctatum, nitidulum, segmentis singulis margine summo rufo-piceis. Pedes cum coxis anterioribus flavo-testacei.

Mas abdominis segmento inferiore sexto apice acute triangulariter exciso.

- ✕ \* 4. *S. intermedius*: Niger, antennis pedibusque testaceis, capite thorace latiore, thorace ovato, elytris thorace paulo longioribus, fortiter punctatis, margine tenui apicali testaceo. — Long. 2 lin.

Er. Col. March. I. 524. 2.

Habitat in Germaniae silvis.

Statura omnino *S. angustatus*, at paululum maior et abunde distinctus, nitidulus, capite thoraceque opacis, subtiliter cinereo-pubescens. Antennae capite vix duplo longiores et palpi testacei. Mandibulae piceo-testaceae. Caput thorace latius, oblongo-subquadratum, basi truncatum, lateribus pone oculos cum angulis posterioribus rotundatum, supra leviter convexum, fortius confertim punctato-rugosum, subtus crebre fortiter profundeque punctatum. Thorax ante medium coleopteris paulo angustior, latitudine summa vix longior, lateribus ante medium subangulato-rotundatus, basin versus fortiter angustatus, leviter convexus, fortius confertim punctato-rugosus. Elytra thorace paulo longiora, crebre fortiter profunde punctata, nigra, margine summo apicali pallide testaceo. Abdomen basin versus subangustatum, crebrius fortiusque punctatum, segmentis singulis apice tenuiter testaceo-marginatis. Pedes cum coxis anterioribus flavo-testacei.

Mas abdominis segmento inferiore quinto longitudinaliter obsolete impresso, sexto acute triangulariter exciso.

- ✕ \* 5. *S. angustatus*: Niger, antennis pedibusque testaceis, capite thorace latiore, thorace basin versus angustato, elytris thorace sesqui longioribus, apice late testaceis. — Long.  $1\frac{2}{3}$  lin.

Er. Col. March. I. 524. 1.

*Staph. angustatus* Payk. Mon. Staph. 36. 27. — Fab. Ent. Syst. I. II. 528. 41. Syst. El. II. 599. 50. — Panz. Ent. Germ. 356. 31. Faun. Germ. 11. 18. — Oliv. Ent. III. 42. 21. 24. t. 2. f. 81. — Marsh. Ent. Britt. 527. 83.

*Paederus angustatus* Payk. Faun. Suec. III. 431. 7. — Grav. Micr. 63. 7. Mon. 141. 9. — Latr. Hist. nat. Crust. et Ins. IX. 347. 5. Genera Crust. et Ins. I. 292. 3. — Oliv. Encycl. meth. VIII. 630. 13. — Gyll. Ins. Suec. II. 375. 4. — Sahlb. Ins. Fenn. I. 343. 4. — Mannerh. Brachelytr. 40. 7. — Runde Brachelytr. Hal. 13. 4.

*Astenus angustatus* Boisd. et Lacord. Faun. Ent. Paris. I. 436. 2.

*Staph. gracilis* Payk. Mon. Staph. 38. 28. — Fab. Ent. Syst. I. II. 528. 44. Syst. El. II. 599. 54. — Panz. Ent. Germ. 357. 34.

Habitat in Europa.

Niger, nitidus, capite thoraceque opacis, subtiliter cinereo-pubescens. Antennae capite vix duplo longiores, apice subincrassatae, testaceae. Palpi testacei. Mandibulae piceo-testaceae. Caput thorace paulo latius, oblongo-



subquadratum, basi truncatum, lateribus pone oculos et angulis posterioribus rotundatis, supra depressiusculum, confertim punctato-subrugosum, infra sat crebre fortius profundiusque punctatum. Thorax coleopteris plus dimidio angustior, latitudine summa vix longior, lateribus ante medium fortiter rotundato-angulatis, basin versus valde angustatus, depressiusculus, confertim punctato-subrugosus. Elytra thorace sesqui prope longiora, crebre fortiter punctata, nigra, apice late sinuatim testacea. Abdomen basin versus subangustatum, parcius minus subtiliter punctulatum, longius pubescens, segmentis 4 primis margine tenui apicali testaceo. Pedes cum coxis anterioribus dilute flavo-testacei.

Mas abdominis segmento inferiore sexto apice acute triangulariter exciso.

- \* 6. *S. bimaculatus*: *Niger, capite thorace latiore, thorace subovato, rufo-piceo, antennis, elytris pedibusque testaceis, his macula media nigra.* — *Long.*  $1\frac{3}{4}$  lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura fere praecedentis, at paululum gracilior. Corpus totum pubes minus brevi depressa cinerea parcius vestitum. Antennae capite duplo prope longiores, apice vix crassiores, flavo-testaceae. Palpi flavi. Os rufo-piceum. Caput thorace paululum latius, oblongum, basi et lateribus pone oculos leviter, angulis posterioribus fortius rotundatis, supra perparum convexum, confertissime punctato-rugulosum, opacum, nigro-piceum, subtus creberrime profundius punctatum, nigrum, subnitidum. Thorax coleopteris plus dimidio angustior, latitudine summa paulo longior, lateribus ante medium rotundatis, basin versus angustatus, subdepressus, confertim punctato-rugulosus, opacus, picescenti-rufus. Scutellum nigro-piceum. Elytra thorace paulo longiora, creberrime fortius profunde punctata, testacea, macula rotundata media, interdum sat minuta, nigra notata. Abdomen basin versus subangustatum, crebre fortius punctatum, nitidulum, nigrum, segmentis singulis piceo-testaceo-marginatis. Pectus nigro-piceum. Pedes pallide flavo-testacei.

Mas abdominis segmento inferiore sexto apice inciso.

Variat aut thorace etiam nigro, abdominisque segmentis 4 primis supra concoloribus, aut dilutior, capite abdomineque rufis, illo fronte media nigricante, hoc segmento quinto solo praeter marginem apicalem nigro, nunquam etiam elytrorum macula evanescente.

- \* 7. *S. limbatus*: *Niger, antennis pedibusque testaceis, capite thorace latiore, thorace basin versus angustato, elytris thoracis longitudine, sutura dimidia apiceque testaceis.* — *Long.*  $1\frac{3}{4}$  lin.

Habitat in Columbia, Dom. Moritz.

S. angustato parum minor, niger, minus nitidus, pube depressa cinerea, in capite thoraceque subtiliore, in elytris abdomineque magis conspicua vestitus. Antennae capite thoraceque parum breviores, tenues, dilute testaceae. Palpi testacei. Mandibulae rufae. Caput thorace plus sesqui latius, coleopterorum prope latitudine, oblongum, pone oculos rotundatum, supra inter oculos leviter convexum, ceterum depressum, subtilissime punctato-rugulo-

sum. Thorax coleopteris duplo angustior, latitudine sesqui longior, lateribus ante medium leviter rotundatus, apice attenuatus, basin versus leviter angustatus, subdepressus, confertissime punctato-rugulosus. Elytra thoracis longitudine, dense profundeque punctata, limbo apicali cum sutura ad elytri medium fere adscendente testaceo. Abdomen basin versus sensim subangustatum, crebrius minus subtiliter punctatum, segmentis singulis margine tenui testaceo terminatis. Pedes cum coxis dilute testacei.

Mas abdominis segmento quinto longitudinaliter impresso, apice subsinuato, sexto apice inciso.

- \* 8. *S. lateralis*: *Linearis, niger, antennis pedibusque flavis, capite thoraceque piceis, hoc basin versus angustato, elytris thorace paulo longioribus, testaceis, macula magna laterali nigra.* — Long.  $1\frac{1}{4}$  lin.

Habitat Novae Valenciae in Columbia, Dom. Moritz.

Habitu omnino *S. filiformis*, at paulo brevior. Antennae capite thoraceque paulo breviores, tenues, albido-flavae. Palpi pallide flavi. Caput thorace vix latius, oblongum, basi truncatum, angulis posterioribus subrotundatis, supra perparum convexum, subtiliter punctato-rugulosum, subtilissime parce griseo-pubescent, opacum, rufo-piceum. Thorax coleopteris parum angustior, latitudine summa perparum longior, lateribus ante medium cum angulis anterioribus fortius rotundatus, basin versus leviter angustatus, subdepressus, confertissime punctato-rugulosus, densius subtilissime griseo-pubescent, lateribus parce nigro-pilosus, fusco-piceus, opacus. Scutellum nigro-piceum. Elytra thorace paululum longiora, confertim fortius punctata, subtiliter tenuiterque griseo-pubescentia, testacea, macula permagna laterali suturaque tenui nigris. Abdomen parallelum, minus dense subtiliterque punctatum, pube longiore depressa fusco-grisea densius vestitum, nigrum, nitidum, segmentis singulis tenuiter testaceo-marginatis. Pectus nigrum. Pedes cum coxis pallide flavi.

Mas latet.

- \* 9. *S. bisignatus*: *Elongatus, testaceus, elytris macula magna laterali, abdomine segmento quarto quintoque nigris.* — Long.  $1\frac{2}{3}$  lin.

Habitat prope Caracas Columbiae, Dom. Moritz.

Statura *S. longiusculi*, testaceus, vix nitidus, tenuiter subtiliterque griseo-pubescent, capite thoraceque lateribus parce nigro-pilosis. Antennae capite thoraceque paulo breviores, tenues, flavae. Palpi flavi. Caput thorace vix latius, basi truncatum, angulis posterioribus rotundatis, latitudine paulo longius, subdepressum, subtiliter punctato-rugulosum. Thorax coleopterorum prope latitudine, latitudine paulo longior, lateribus ante medium subangulatis, depressus, subtilissime et vix perspicue punctato-rugulosus. Coleoptera apicem versus subangustata, thorace parum longiora, crebrius sat profunde punctata, interstitiis aequalibus, utroque macula magna rotundata laterali pone medium signato. Abdomen basin versus sensim leviter angustatum, anterior fortius, posterior subtiliter punctatum, segmento quarto piceo, quinto nigro. Pedes albido-testacei.

Mas abdominis segmento inferiore sexto apice profundius inciso.

- \* 10. *S. attenuatus*: *Elongatus, testaceus, elytris puncto posteriore, abdomine segmento quinto nigris*. — Long. 2 lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Praecedenti affinis, at magis elongatus, testaceus, capite thoraceque rufo-testaceis, vix nitidus, pube subtili depressa cinerea tenuiter vestitus. Antennae capite paulo longiores, tenues, flavo-testaceae. Palpi testacei. Caput thoracis latitudine, oblongum, pone oculos cum basi rotundatum, subdepressum, subtiliter punctato-rugulosum. Thorax coleopteris vix angustior, latitudine longior, lateribus ante medium subangulatus, basin versus modice angustatus, subdepressus, sat evidenter punctato-rugulosus. Coleoptera apicem versus modice angustata, thorace tertia parte longiora, dense profundius punctata, interstitiis transversim subrugulosis, lateribus ante apicem macula minuta rotundata nigra notata. Abdomen elongatum, basin versus sensim angustatum, crebre minus subtiliter punctatum, segmento quinto nigro, margine apicali testaceo. Pedes dilute testacei.

Mas abdominis segmento inferiore quinto longitudinaliter impresso, sexto apice profundius inciso.

Obs. Differt a praecedente statura magis elongata, abdomine basin versus fortius angustato, elytris longioribus, capite basi haud truncato, thorace perspicue punctato-ruguloso, etc.

- \* 11. *S. longiusculus*: *Elongatus, linearis, testaceus, abdomine segmentis ultimis duobus nigris*. — Long.  $1\frac{2}{3}$  lin.

*Paederus longiusculus* Mannerh. *Brachelytr.* 39. 5.

*Paederus discopunctatus* Say *Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV.* 457. 1.

Habitat in America septentrionali.

Statura omnino praecedentium, rufo-testaceus, opacus, tenuiter subtiliterque griseo-pubescent, capitis thoracisque lateribus abdominisque apice nigro-pilosellis. Antennae capite thoraceque paulo breviores, tenues, flavae. Palpi flavi. Caput thorace vix latius, oblongum, basi cum angulis posterioribus rotundatum, supra confertissime subtiliter punctato-rugulosum, subtus crebre fortius punctatum. Thorax coleopteris angustior, latitudine longior, lateribus ante medium rotundatis, basin versus leviter angustatus, parum convexus, confertissime punctato-rugosus. Elytra thorace tertia parte longiora, crebre punctata. Abdomen elongatum, basin versus sensim subangustatum, crebrius minus subtiliter punctulatum, segmento quinto nigro, sexto nigro-piceo. Pedes flavi.

Mas abdominis segmento inferiore sexto profunde inciso.

- \* 12. *S. testaceus*: *Elongatus, linearis, testaceus, immaculatus, antennis pedibusque pallidioribus*. — Long.  $2\frac{1}{3}$  lin.

Habitat in Columbia, Dom. Moritz.

Habitu fere praecedentium, elongatus, linearis, subdepressus, totus testaceus, capite thoraceque rufo-testaceis, vix nitidus, pube subtili depressa cinerea tenuiter vestitus, capitis thoracisque lateribus abdominisque apice parce nigro-pilosellis. Antennae capite thoraceque paulo breviores, tenues, pallide flavae. Palpi pallide testacei. Caput thorace haud latius, oblongum,



basi truncatum, lateribus pone oculos et angulis posterioribus leviter rotundatis, supra anterior depressiusculum, subtiliter punctato-rugulosum, subtus crebre punctatum. Thorax coleopteris haud angustior, latitudine longior, lateribus ante medium rotundatis, basin versus leviter angustatus, subdepressus, subtiliter punctato-rugulosus, lateribus ante medium utrinque subimpressus. Elytra thorace paululum longiora, dense minus subtiliter sat profunde punctata. Abdomen parallelum, subtilius punctulatum, segmento quinto basi fuscescente. Pedes pallide flavi.

Mas abdominis segmento inferiore quinto apice subsinuato, sexto profunde triangulariter exciso.

\* 13. *S. coarctatus*: *Elongatus, testaceus, capite nigro thoraceque ovato rugulosis, elytris parce punctatis fascia media, abdomine fascia ante apicem nigris.* — Long. 2 lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Reliquis fere gracilior, elongatus, subdepressus. Antennae capite thoraceque paulo breviores, tenues, articulo tertio secundo sesqui prope longiore, sequentibus sensim paulo brevioribus, ultimo elongato, acuminato, pallide testaceae, articulis 3 — 5 medio fuscis. Palpi testacei. Caput thorace coleopterisque sesqui prope latius, basi cum angulis posterioribus rotundatum, supra posterius leviter convexum, subtiliter minus confertim longitudinaliter rugulosum, tenuiter griseo-pubescent, nigrum, minus nitidum, ore dilutius, rufo-piceo. Thorax latitudine summa paulo longior, ante medium coleopteris prope latior, basin apicemque versus fortiter attenuatus, depressiusculus, longitudinaliter dense rugosus, parce pubescens, minus nitidus, rufus, medio piceus. Elytra thoracis latitudine, basi attenuata, parce profundius subseriatim punctata, parce pubescentia, nitida, fascia media nigra, ante fasciam rufo-, pone fasciam flavo-testacea. Abdomen apicem versus leviter incrassatum, parcius punctatum, tenuiter griseo-pubescent, subnitidum, testaceum, segmento quinto nigro-piceo. Pectus nigro-piceum. Pedes subelongati, pallide flavi, femoribus apice nigris, tibiis basi fuscis.

Mas abdominis segmento inferiore quinto medio leviter impresso, sexto apice utrinque leviter sinuato, medio profunde inciso.

14. *S. tristis*: *Niger, antennis pedibusque rufo-testaceis, capite thorace latiore, elytris thoracis latitudine, concoloribus.* — Long. 1½ lin.

Habitat in Sardinia, Mus. Reg. Taurin.

S. filiformi latitudine aequalis, at brevior et minus gracilis, niger, opacus, pube depressa griseo-fusca densius vestitus. Antennae capite sesqui longiores, rufo-testaceae. Palpi testacei. Mandibulae et reliquae oris partes, labro nigro excepto, rufo-piceae. Caput basi thorace latius, latitudine baseos haud longius, basi emarginatum, angulis posterioribus retrorsum prominentibus, rotundatis, lateribus pone oculos leviter rotundatis, ante oculos paulum angustatis, apice leviter emarginatum, supra postice leviter convexum, antrorsum planatum, confertissime punctato-rugulosum, subtus confertissime punctatum. Thorax basin versus angustatus, latitudine summa haud longior, basi apiceque utrinque oblique truncatus, depressus, confertissime punctato-rugulosus, posterius linea longitudinali subtilissima impressa tenuissime ca-

*naliculatus*. Coleoptera thorace haud longiora, neque eius basi latiora, subdepressa, apice oblique truncata, dense minus fortiter punctata, nigra, apice concolore. Abdomen minus confertim punctatum, punctis transversim rugulosis, longius pubescens, segmentis singulis margine summo apicali piceo-testaceis. Pedes cum coxis anterioribus rufo-testacei.

\* 15. *S. curtulus*: *Niger, opacus, antennis pedibusque rufo-testaceis, thorace transversim subquadrato, elytris thorace brevioribus.* — Long. 2 lin.

Habitat in Sardinia, Dom. Prof. Gené.

Brevis, latiusculus, subdepressus, niger, opacus, pube depressa grisea densius vestitus. Antennae capite paulo longiores, tenuiores, rufo-testaceae. Palpi testacei. Caput thoracis latitudine, latitudine summa haud longior, basi leviter emarginatum, angulis posterioribus subrotundatis, lateribus ante oculos angustatum, supra parum convexum, subtilissime punctato-rugulosum, infra confertissime minus subtiliter punctatum, rufo-piceum. Thorax coleopterorum latitudine, latitudine paulo brevior, basi apiceque utrinque oblique truncatus, angulis omnibus obtusis, lateribus rectis, basin versus vix angustatus, subdepressus, densius subtiliter punctatus. Elytra thorace paulo breviora, apice utrinque oblique truncata, dense minus subtiliter profunde punctata, margine summo apicali rufo-piceo. Abdomen coleopterorum latitudine, parallelum, minus subtiliter punctulatum, segmentis singulis margine apicali rufo-piceo. Pedes breviusculi, cum coxis testaceo-rufi.

Mas abdominis segmento inferiore sexto apice acute triangulariter exciso.

#### Species mihi invisae.

16. *S. binotatus*: *Testaceus, capite, elytrorum parte anoque nigris, pedibus pallidis.* — Long. 2 lin.

*Paederus binotatus* Say Journ. Acad. Nat. Scienc. Philadelph. III. p. 154.

Habitat in America septentrionali.

Pallide testaceus, leviter pubescens, punctatus. Caput thorace maius, nigrum. Antennae et oris partes pallidae. Thorax longitudinaliter subovatus, dense punctatus. Elytra ante apicem macula magna laterali nigra. Abdomen segmento ultimo anoque nigris. Pedes albid.

#### \*\* (S p u r i i).

\* 17. *S. bipunctatus*: *Elongatus, subdepressus, testaceus, pedibus pallidis, capite thoraceque confertim subtiliter punctato-subrugulosis, elytris puncto medio nigro notatis.* — Long. 2½ lin.

Habitat in Columbia, Dom. Moritz.

Corpus elongatum, subdepressum, testaceum, parum nitidum, pube depressa grisea minus dense vestitum. Antennae capitis thoracisque prope longitudine, tenues, articulo primo sequentibus ambobus longitudine aequali, tertio secundo plus sesqui longiore, 4 — 10 sensim paulo brevioribus, ultimo reliquis paululum crassiore, apice abrupte acuminato, flavae. Palpi flavi. Labrum transversum, medio leviter productum et acute bidentatum

vel bispinosum. Caput thorace paulo latius, latitudine paulo longius, pone oculos sensim rotundato-angustatum, supra depressiusculum, subtiliter punctatum et concinne reticulato-rugulosum, subtus parce obsoletius punctatum, lateribus parce nigro-pilosellum. Thorax coleopteris dimidio angustior, latitudine summa vix longior, lateribus paulo ante medium subangulatus, apicem versus sensim angustatus, lateribus rectis, basin versus vix angustatus, angulis posterioribus rotundatis, depressiusculus, eodem modo ac caput subtiliter reticulato-rugulosus, versus latera utrinque obsolete impressus. Elytra thorace plus sesqui longiora, subdepressa, minus subtiliter sat profunde densius punctata, puncto medio nigro. Abdomen coleopteris angustius, apicem versus subangustatum, subtiliter punctulatum, apice densius pilosum. Pedes pallide flavi.

Mas abdominis segmento inferiore sexto apice profunde exciso.

- \* 18. *S. hybridus*: *Niger, obscurus, elytris apice pedibusque flavis, capite thoraceque confertissime punctato-subrugulosus, hoc oblongo, lateribus obsolete impresso.* — Long.  $1\frac{2}{3}$  lin.

Habitat in Columbia, Dom. Moritz.

Niger vel nigro-piceus, parum nitidus. Antennae capitis thoracisque longitudine, testaceae, articulo primo sequentibus ambobus paulo longiore. Palpi testacei. Labrum rufo-piceum, transversum, apice acute bidentatum. Caput thorace paulo latius, suborbiculatum, supra leviter convexum, subtiliter confertissime punctato-subrugulosum, opacum, parce subtiliter griseo-pubescent, subtus parcius profundiusque punctatum, subnitidum. Thorax coleopteris duplo fere angustior, latitudine paulo longior, lateribus paulo ante medium rotundatus, basin versus angustatus, apicem versus subsinuatum attenuatus, modice convexus, subtiliter confertissime punctato-rugulosus, parce subtiliter cinereo-pubescent, utrinque lateribus obsolete impressus. Elytra thorace paulo longiora, fortius sat profunde crebreque punctata, subtiliter cinereo-pubescentia, apice late sinuatum flava, sutura nigra. Abdomen subtiliter punctulatum, densius cinereo-pubescent, segmentis singulis apice piceo-marginatis. Pedes pallide flavi.

Mas abdominis segmento inferiore sexto late profundeque sinuato.

- \* 19. *S. stigma*: *Rufo-piceus, opacus, capite thoraceque oblongo confertissime subtiliter punctato-rugulosus, nigris, elytris thorace longioribus, testaceis, macula laterali obliqua nigra.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Statura omnino et summa affinitas praecedentis. Antennae capitis thoracisque prope longitudine, articulo primo sequentibus ambobus longiore, tertio secundo parum longiore, 4 — 10 sensim paululum brevioribus et crassioribus, ultimo ovato, apice subcompresso, testaceae. Palpi testacei. Os obscure rufo-testaceum. Labrum transversum, apice medio leviter productum et acute bidentatum. Caput thorace latius, suborbiculatum, basi truncatum, lateribus pone oculos cum angulis posterioribus rotundatis, supra levissime convexum, confertim subtiliter punctato-rugulosum, opacum, tenuiter griseo-pubescent, subtus crebre profundeque punctatum, nigrum vel nigro-piceum. Thorax coleopteris plus dimidio angustior, latitudine summa



paulo longior, lateribus ante medium angulatis, dein subrotundatis, basin versus leviter angustatus, apicem versus sinuato-attenuatus, supra leviter convexus, confertissime subtilius punctato-rugulosus, opacus, tenuiter griseo-pubescent, niger vel nigro-piceus. Scutellum rufo-piceum. Elytra thorace paulo longiora, dense profundius punctata, testacea, macula obliqua in medio latere, et sutura infra medium nigris. Abdomen dilute rufo-piceum, segmentorum marginibus apiceque testaceis, subnitidum, subtiliter punctulatum, pube longiore depressa grisea densius vestitum. Pectus rufo-piceum. Pedes pallide flavo-testacei.

Mas abdominis segmento inferiore sexto apice late profundeque sinuato.

- \* 20. *S. exiguus*: *Rufo-piceus, opacus, capite nigricante thoraceque confertissime punctato-rugulosus, elytris thoracis longitudine antennisque testaceis, pedibus pallidis.* — Long. 1 lin.

Habitat in Puerto-Rico, Dom. Moritz.

Statura fere praecedentium, at longe minor. Antennae capitis thoracisque prope longitudine, articulo primo sequentibus tribus coniunctis longitudine aequali, tertio secundo paulo longiore, 4 — 10 sensim paululum brevioribus et crassioribus, ultimo maiusculo, ovato, apice acuminato, totae testaceae. Palpi flavi. Labrum transversum, apice medio leviter productum et acute dentatum. Caput thorace paulo latius, suborbiculatum, supra leviter convexus, confertim punctato-rugulosus, opacus, tenuiter griseo-pubescent, nigro-piceum, os versus rufescens, subtus concolor, subnitidum, crebre profundius punctatum. Thorax coleopteris paulo angustior, latitudine summa vix longior, lateribus ante medium angulatis, basin versus angustatus, supra modice convexus, dense minus subtiliter punctato-rugulosus, opacus, tenuiter griseo-pubescent, rufo-piceus. Elytra thoracis vix longitudine, basin versus sensim angustata, depressa, parce minus subtiliter punctata, parce cinereo-pubescentia. Abdomen subtiliter punctulatum, longius cinereo-pubescent, piceo-testaceum, segmentorum margine apicali anoque dilutioribus. Pectus piceo-testaceum. Pedes albid, femoribus anticis subtus leviter dilatatis.

- \* 21. *S. naevius*: *Rufo-piceus, opacus, capite thoraceque transversim subquadrato subtiliter punctato-rugulosus, elytris thoracis longioribus testaceis, macula laterali transversa nigra.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Columbia, Dom. Moritz.

Praecedentibus affinis, at corpus breviusculum crassiusculumque, rufo-piceum, opacus, pube depressa grisea densius vestitum. Antennae capitis thoracisque prope longitudine, articulo primo sequentibus ambobus paulo longiore, tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus at vix crassioribus, ultimo crassiore, ovato, apice acuminato, flavo-testaceae. Palpi testacei. Labrum breve, apice medio leviter productum et bidenticulatum. Caput thorace paulo latius, latitudine baseos prope brevius, basi leviter emarginatum, angulis posterioribus subrectis, ante oculos angustatum, supra parum convexus, subtiliter punctato-rugulosus, subtus parcius punctatum, nitidulum. Thorax coleopteris paulo angustior, latitudine paulo brevior, basin versus leviter angustatus, apice utrinque oblique truncatus, an-

gulis anterioribus rectis, posterioribus obtusis, subrotundatis, supra parum convexus, confertissime subtiliter punctato-rugulosus. Elytra thorace tertia parte longiora, dense profundius punctata, testacea, lateribus paulo ante medium macula transversa nigra notata. Abdomen breviusculum, subtiliter punctulatum, apice testaceum. Pedes flavo-testacei.

Mas abdominis segmento inferiore sexto late profundeque sinuato.

\* 22. *S. labeo*: *Testaceus, thorace utrinque impresso, labro maximo, trilobo, femoribus anticis basi dilatatis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Americae ins. St. Johannis, Dom. Moritz.

Corpus totum dilute testaceum, nitidum, elytris abdomineque pallidioribus. Antennae tenuiores, capite thoraceque longiores, articulis omnibus elongatis, primo sequentibus ambobus subaequali, tertio secundo sesqui longiore, sequentibus sensim paulo brevioribus at haud crassioribus, ultimo apice abrupte acuminato. Labrum maximum, bisinuatum, lobis lateralibus brevissimis, rotundatis, intermedio producto. Caput magnum, thorace paulo latius, postice rotundatum, antice elongatum, posterius leviter convexum, inter oculos utrinque profunde foveolatum, antierius utrinque leviter longitudinaliter impressum, margine laterali ab oculis inde elevato, parcius subtiliter punctulatum. Thorax coleopteris angustior, latitudine summa paulo longior, lateribus rotundato-amplius, basin versus angustatus, apicem versus attenuatus, ante apicem leviter constrictus, crebrius subtiliusque punctatus, dorso obsolete carinatus, pone medium secundum marginem lateralem utrinque longitudinaliter impressus. Scutellum punctulatum. Elytra thorace vix longiora, densius punctata. Abdomen subtilissime punctulatum. Pedes pallide flavi, femoribus anticis fortiter compressis, basi dilatatis, subtus parce nigro-setulosis.

\* 23. *S. troglodytes*: *Rufo-castaneus, nitidus, subtiliter punctatus, capite aequali, thorace subgloboso elytrisqne nigro-piceis, his apice testaceis.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Columbia, Dom. Moritz.

Habitus, statura et summa affinitas sequentis. Caput et thorax pube subtili erecta grisea, elytra et abdomen pube depressa cinerea vestita. Antennae capite thoraceque paulo breviores, crassiusculae, articulo primo sequentibus ambobus paulo brevioribus, his obconicis, subaequalibus, 4 — 10 sensim paulo brevioribus et crassioribus, ultimo breviter ovato, pilosellae, nigro-fuscae, articulis tribus primis testaceis, ultimo ferrugineo. Palpi testacei. Labrum medio leviter sinuatum. Caput thoracis latitudine, orbiculatum, supra leviter convexum, subtilius sat crebre punctatum, nigro-piceum, nitidum. Thorax coleopteris paulo angustior, latitudine summa haud longior, basi, apice et lateribus, his praecipue ante medium, rotundatis, basin versus angustatus, convexus, subtiliter obsoleteque confertim punctatus, nigro-piceus, nitidus. Elytra thorace sesqui longiora, leviter convexa, parcius subtiliter obsoleteque punctata, nigro-picea, nitida, apice tertia parte dilute testacea. Abdomen laevigatum, dilute rufo-castaneum, nitidum. Pectus piceum. Pedes pallide flavi.

Mas abdominis segmento inferiore sexto late profundeque triangulariter exciso.

Variat capite, thorace elytrisque dilute rufo-castaneis, his apice testaceis.

Obs. A sequente praecipue distinctus thorace latiore, subgloboso, capite in utroque sexu aequali, fortius punctatis, antennis medio fusco-nigris, in individuis etiam iis, quae colore dilutiore cum illo congruunt.

\* 24. *S. pusto*: *Rufo-castaneus, nitidus, subtilissime punctatus, capite foveolato, thorace subovato, elytris apice testaceis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Dilute rufo-castaneus, nitidus, capite thoraceque pube brevi subtili erecta fusca tenuiter vestitis. Antennae capite thoraceque breviores, crassiusculae, articulo primo sequentibus ambobus paulo brevior, tertio secundo paulo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, ultimo apice abrupte obtuse acuminato, pilosellae, rufo-testaceae. Palpi testacei. Labrum transversum, apice rotundatum, medio sinuatum. Caput thorace paulo latius, suborbiculatum, convexum, subtilissime punctatum, fronte media in mare tri-impressa, foveis anterioribus duabus maioribus, laevigatis, interstitio elevato, compresso, piloso, feminae inter oculos utrinque late impressa. Thorax coleopteris duplo fere angustior, latitudine paulo longior, subovalis, fortiter convexus, subtiliter parcius punctatus. Elytra thorace paulo longiora, leviter convexa, parcius subtiliter punctata, apice testacea. Abdomen subtilissime parcius punctulatum. Pedes flavo-testacei.

Mas abdominis segmento inferiore sexto apice late profundeque triangulariter excisis.

Obs. Habitus omnino reliquorum, at corporis superficie laevi in hoc genere alienus videtur.

### **Paederus Grav.**

*Curt. Britt. Ent. III. t. 108. — Steph. Illustr. V. p. 297. — Boisd. et Lacord. Faun. Ent. Paris. I. p. 428. — Er. Col. March. I. p. 517.*

*Paederus* Fam. III. Grav. Mon. p. 142.

*Labrum apice emarginatum.*

*Palpi maxillares articulo ultimo obtuso, distincto.*

*Antennae rectae.*

*Tarsi articulo quarto bilobo.*

Corpus elongatum, lineare, aliis alatum, aliis apterum. Caput thoracis plerumque latitudine, suborbiculatum, basi profunde constrictum, pedunculo brevissimo tenui thoraci immisum, exsertum, porrectum, oculis parvis, modice prominulis. Labrum transversum, medio anguste leviter emarginatum. Mandiculae falcatae, acutae, medio acute bidentatae. Maxillae (t. IV. f. 30. b.) malis membranaceis, interiore interne, exteriori apice barbatis. Palpi maxillares modice elongati, articulo primo brevi, secundo tertioque elongatis, subaequalibus, hoc apicem versus sensim incrassato, quarto minimo, obtuso. Labium (t. IV. f. 30. a.) mento transversum, ligula lata, biloba, lobis rotundatis, membranacea, paraglossis subacuminatis, membranaceis, pu-



bescentibus, interne ciliatis, ligulam paulo superantibus. Palpi labiales articulis duobus primis subaequalibus, cylindricis, tertio minimo, obtusiusculo. Antennae filiformes, rectae, articulo primo tertioque subelongatis, ultimo acuminato. Thorax ovatus, convexus, interdum globosus. Scutellum rotundato-triangulari. Elytra truncata. Abdomen lineare, apice conico-attenuatum, segmento septimo eminulo, stylis analibus distinctis. Pedes elongati, graciles, tarsis anticis in utroque sexu articulis 4 primis modice dilatatis, obcordatis, aequalibus, subtus dense tomentosis, posterioribus articulis 3 primis longitudine sensim decrescentibus, quarto profunde bilobo, subtus subtilius pubescentibus.

### Observationes.

1. Genus characteribus valde distinctum, corpore plerumque laetis coloribus insigni, nitido, capite thoraceque parce subtiliter punctatis, parce subtiliterque pilosellis.

2. Abdominis segmentum inferius sextum in mare per totam prope longitudinem incisum, feminae integrum.

3. Victus imprimis in ripis. Nonnullae species sociales occurrunt.

X \* 1. *P. littoralis*: *Apterus, niger, elytris cyaneis, thorace, abdominis segmentis 4 primis pedibusque rufis, femoribus omnibus apice late nigris, antennis testaceis, medio fuscis*. — Long. 4 lin.

*Grav. Micr.* 61. 4. — *Mannerh. Brachelytr.* 39. 2. — *Sahlb. Ins. Fenn.* I. 342.

1. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 429. 2. — *Runde Brachelytr. Hal.* 13. 1. — *Er. Col. March.* I. 519. 3.

*Paed. riparius* var. *Latr. Hist. nat. Crust. et Ins.* I. 346. 2.

*Paed. riparius* Oliv. *Ent.* III. 44. 4. 2. t. 1. f. 2.

*Staph. riparius* *Schrunk Enum. Ins. Austr.* 233. 441.

Habitat in Germania, Gallia, Italia, Lusitania.

*P. ripario* paulo maior, praesertim robustior. Antennae capite thoraceque paulo breviores, articulo tertio secundo sesqui longiore, 4—10 sensim paulo brevioribus, testaceae, articulis 5—9 fuscis. Palpi maxillares testacei, articulo tertio apice fusciscenti, labiales fuscis. Mandibulae nigrae. Caput thoracis latitudine vel eodem paulo latius, orbiculatum, nigrum, nitidum, supra parum convexus, sparsim punctatum, utrinque nigro-pilosellum, ad antennarum basin utrinque leviter impressum. Thorax coleopterorum latitudine, vel iisdem paulo latior, globosus, convexus, rufus, nitidus, dorso biseriatim, lateribus sparsim parce subtiliterque punctatus, parce subtiliterque pilosus. Scutellum punctulatum, obscure rufum. Elytra thorace haud longiora, fortiter punctata, tenuiter cinereo-pubescentia, cyanea, nitida. Abdomen parce subtiliterque punctulatum, rufum, nitidum, segmentis ultimis duobus nigris. Pectus totum nigrum. Pedes cum coxis anterioribus rufo-testacei, femoribus omnibus apice late nigris, tarsis apice fusciscentibus.

Obs. Specimina ex Europa meridionali: Lusitania, Sicilia, Sardinia, Carinthia, paulo maiora, validiora, capite thoraceque maioribus, specie autem nullo modo a genuinis distincta. Individua Lusitanica in collectione Hoffmannseggiana nomine *P. Finisterrae* exstiterunt, Carinthiaca misit Dahl nom. *P. speculatoris* et diversicollis signata.

- \* 2. *P. brevipennis*: *Apterus, niger, elytris coeruleis, thorace, abdominis segmentis 4 primis pedibusque rufis, femoribus posterioribus apice nigricantibus, antennis testaceis.* — Long. 3 lin.

*Boisd. et Lacord. Faun. Ent. Paris. I. 430. 3.*

Habitat in Austria et Carinthia, Dahl, in Saxonia, Dom. Märkel, in Gallia, Dom. Aubé.

Statura et summa affinitas *P. littoralis*, at duplo minor. Antennae capite thoraceque parum breviores, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus, testaceae, articulis 5 — 9 nonnunquam fuscescentibus. Palpi testacei, maxillares articulo tertio apice summo subfuscescente. Mandibulae nigrae. Caput thoracis fere latitudine, orbiculatum, nigrum, nitidum, supra parum convexum, sparsim punctatum, fronte media late laevi, inter oculos et inter antennarum basin utrinque leviter impressa, parce nigro-pilosum. Thorax coleoptera latitudine fere excedens, globosus, convexus, rufus, nitidus, dorso biseriatim, lateribus disperse parce punctatus, parce subtiliterque pilosus. Scutellum punctulatum, rufum. Elytra thorace paulo breviora, fortiter punctata, coerulea, nitida, nigro-pubescentia pilosellaque. Abdomen parce punctulatum, nigro-pilosellum, nitidum, rufum, segmentis ultimis duobus nigris. Pectus nigrum. Pedes toti rufo-testacei, femoribus posterioribus apice summo perparum fuscescentibus, coxis omnibus testaceis, posticis basi nigricantibus.

Obs. *P. brachypteri* et *tenuipennis* nominibus a Dahlio missus.

- \* 3. *P. Capensis*: *Apterus, niger, elytris coeruleis, thorace, abdominis segmentis 4 primis, mesosterno pedibusque rufis, femoribus apice tibiisque basi fuscis, antennis testaceis, medio fuscis.* — Long. 3½ lin.

Habitat in Africa australi, Dom. Drège.

*P. littoralis* paulo minor et gracilior. Antennae capitis thoracisque longitudine articulo tertio secundo duplo longiore, 4 — 10 sensim paulo brevioribus et apicem versus magis incrassatis, testaceae, articulis 4 — 9 obscure fuscis. Palpi testacei. Mandibulae rufae. Caput thoracis fere latitudine, suborbiculatum, nigrum, nitidum, supra perparum convexum, utrinque crebrius subtiliusque punctatum, medio late longitudinaliter laeve, utrinque inter oculos obsolete, ad antennarum insertionem distinctius impressum. Thorax coleopterorum latitudine, ovatus, lateribus ante medium rotundatis, basin versus modice angustatus, modice convexus, rufus, nitidus, dorso biseriatim, lateribus disperse parce obsoleteque punctatus. Scutellum piceum. Elytra thorace haud longiora, profunde fortiterque punctata, griseo-pubescentia, coerulea, nitida. Abdomen parce punctatum, nitidum, rufum, segmentis duobus ultimis nigris. Pectus nigrum, mesosterno rufo. Pedes coxis omnibus rufis, femoribus testaceis, apice late fuscis, tibiis fuscis, apice tarsisque obscure testaceis.

- \* 4. *P. longipennis*: *Alatus, niger, elytris coeruleis, mandibulis, thorace oblongo, abdominis segmentis 4 primis pedibusque rufis, his geniculis nigris, antennis nigris, basi testaceis.* — Long. 3 lin.

*Er. Col. March. I. 517. 1.*

*Paed. riparius Grav. Micr. 65. 5. Mon. 143. 11. — Zetterst. Ins. Lappon. 68. 1.*

*Var. Tarsis tibiisque fuscis, his basi nigricantibus.*

*Paederus fuscipes* Curtis Britt. Ent. III. t. 108.

Habitat in Europa.

Antennae capitis thoracisque longitudine, articulo tertio secundo duplo prope longiore, nigrae, articulis 3 vel 4 primis testaceis. Palpi maxillares testacei, articulo tertio apice nigro, labiales fuscii. Mandibulae testaceae. Caput thoracis latitudine, suborbiculatum, nigrum, nitidum, supra parum convexum, utrinque parce punctatum, fronte media laevi, antice utrinque longitudinaliter impressum, parce pilosum. Thorax oblongus, coleopteris duplo fere angustior, latitudine sesqui longior, basin versus vix angustatus, lateribus ante medium subrotundatis, leviter convexus, dilute rufus, nitidus, dorso biseriatim, lateribus disperse parce punctatus, parce pilosellus. Scutellum sublaeve, nigrum. Elytra thorace sesqui longiora, dense punctata, tenuiter albido-pubescentia, parce nigro-pilosa, coerulea, nitida. Abdomen parce obsoleteque punctulatum, nitidum, rufum, segmentis ultimis duobus nigris. Pectus nigrum. Pedes cum coxis omnibus rufo-testacei, femoribus apice nigris, tibiis basi, tarsis apice nigricantibus.

Variat pedum geniculis minus magisque late nigris, vel interdum tibiis tarsisque fuscis, illis basi, his apice nigris, vel totis nigris.

\* 5. *P. Madagascariensis*: *Alatus, niger, elytris coeruleis, mandibulis, thorace oblongo, abdominisque segmentis 4 primis rufis, antennarum, palporum pedumque basi testaceis.* — Long. 3 lin.

Habitat in Madagascar, Dom. Goudot.

Statura et summa affinitas praecedentis. Antennae capitis thoracisque prope longitudine, articulo tertio secundo duplo prope longiore, 4 — 10 sensim paulo brevioribus et apicem versus paulo magis incrassatis, articulis 3 primis rufo-testaceis, secundo et tertio apice fuscescentibus, quarto basi rufo-testaceo. Palpi maxillares articulis duobus primis testaceis, tertio nigro, basi testaceo, labiales fuscii. Mandibulae rufo-testaceae. Caput thorace paululum latius, orbiculatum, nigrum, nitidum, supra perparum convexum, utrinque crebrius punctatum, fronte media longitudinaliter laevi, utrinque inter oculos et ad antennarum insertionem subimpressum, parce pilosum. Thorax coleopteris dimidio angustior, latitudine tertia parte longior, lateribus anterieus leviter rotundatus, basin versus leviter angustatus, modice convexus, rufus, nitidus, dorso biseriatim, lateribus disperse parce punctatus, parce subtiliterque pilosus. Scutellum sublaeve, nigrum. Elytra thorace sesqui longiora, dense fortiter profundeque punctata, tenuiter albido-pubescentia, coerulea vel virescenti-coerulea, nitida. Abdomen parce punctulatum, tenuiter pubescens, parce nigro-pilosellum, nitidum, segmentis 4 anterioribus rufis, posticis duobus nigris. Pectus nigrum. Pedes coxis trochanteribusque omnibus rufis, coxis posticis basi nigris, femoribus omnibus basi vix ultra medium testaceis, apice tibiisque totis nigris, tibiis nonnunquam apicem versus picescentibus.

\* 6. *P. caligatus*: *Alatus, niger, elytris coeruleis, mandibulis, thorace subovato abdominisque segmentis 4 primis rufis, palpis, antennarum pedumque basi testaceis.* — Long. 3 lin.

Habitat Lutetiae, Dom. Aubé.



Statura *P. longipenni* aequalis, at angustior, habitu omnino *P. riparii*. Antennae capitis thoracisque prope longitudine, articulo tertio secundo sesqui prope longiore, 4 — 10 sensim paulo brevioribus et crassioribus, ultimo breviter subovato, apice acuminato, nigrae, articulis tribus primis dilute testaceis. Palpi maxillares toti testacei. Mandibulae rufo-testaceae. Caput thoracis fere latitudine, orbiculatum, nigrum, nitidum, supra parum convexum, lateribus parce subtiliterque punctatum, utrinque inter oculos obsolete, ad antennarum basin distinctius impressum, parce nigro-pilosum. Thorax coleopteris paululum angustior, latitudine summa sesqui prope longior, ovatus, basin versus leviter angustatus, convexus, rufus, nitidus, punctis raris subtilissimis piligeris notatus. Scutellum punctulatum, nigrum. Elytra thorace vix longiora, parcius sat fortiter punctata, coerulea, nitida, pube tenui depressa albida pilisque parvis erectis nigris vestita. Abdomen segmentis anterioribus 4 parce, ultimis duobus crebrius subtiliter punctulatis, illis rufis, his nigris. Pectus nigrum, mesosterno concolore. Pedes coxis omnibus trochanteribusque et femoribus ultra medium testaceis, dein nigri.

- \* 7. *P. limnophilus*: *Alatus, niger, elytris coeruleis, thorace ovato abdominisque segmentis 4 primis rufis, antennis pedibusque basi testaceis, palpis testaceis, articulo tertio nigro.* — Long. 3 lin.

Habitat in Saxonia, Dom. Märkel, in Gallia, Dom. Aubé.

Statura omnino praecedentis, at paululum minor. Antennae capitis thoracisque longitudine, articulo tertio secundo duplo prope longiore, 4 — 10 sensim paulo brevioribus at vix crassioribus, ultimo oblongo-ovato, acuminato, nigrae, articulis 4 primis testaceis, quarto apice piceo. Palpi maxillares testacei, articulo tertio toto nigro. Mandibulae nigrae, apice piceae. Caput thoracis latitudine, suborbiculatum, nigrum, nitidum, supra parum convexum, utrinque crebrius punctatum, fronte media longitudinaliter late laevi, anteriis longitudinaliter utrinque leviter impressa, parce nigro-pilosum. Thorax coleopteris vix angustior, latitudine summa tertia parte longior, basin versus modice angustatus, ovatus, convexus, dilute rufus, nitidus, lateribus parce, dorso biserialiter subtiliter punctatus. Scutellum sublaeve, nigrum. Elytra thorace paululum longiora, crebre fortius punctata, coerulea, nitida, pube parca depressa albida pilisque raris erectis nigris vestita. Abdomen subtilius punctatum, tenuiter albedo-pubescens, rufo-testaceum, nitidum, segmentis ultimis duobus nigris. Pectus nigrum, mesosterno concolore. Pedes coxis trochanteribusque et femoribus ad medium usque testaceis, dein nigri.

Variat tibiis apice tarsis basi fuscis.

Obs. Differt a praecedente praeter palporum mandibularumque colorem elytris paulo longioribus, crebrius minus profunde punctatis. A *P. longipenni* habitu, et palporum articulo tertio toto nigro facile distinguitur, a sequentibus mesosterno nigro discedit.

- \* 8. *P. riparius*: *Alatus, niger, elytris coeruleis, thorace, abdominis segmentis 4 primis, mesosterno pedibusque rufis, femoribus omnibus apice nigris, antennis nigris, basi testaceis.* — Long.  $3\frac{2}{3}$  lin.

*Fab. Syst. Ent.* 268. 1. *Spec. Ins.* 1. 339. 1. *Mant. Ins.* 1. 223. 1. *Ent. Syst.* 1. II. 536. 1. *Syst. El.* II. 608. 1. — *Payk. Faun. Suec.* III. 427 2. — *Walken. Faun. Paris.* I. 275. 1. — *Cederh. Faun. Ingr.* 121. 373. t. 3. f. d. — *Panz. Faun. Germ.* 9. 11. *Ent. Germ.* 362. 1. — *Rossi Faun. Etr.* 253. 626. (Ed. Hellw.

- I.* 314. 626.) — *Latr. Hist. nat. Crust. et Ins.* IX. 345. 2. t. 79. f. 8. *Gen. Crust. et Ins.* I. 293. 5. — *Gyll. Ins. Suec.* II. 372. 1. — *Mannerh. Brachelytr.* 39. 3. — *Sahlb. Ins. Fenn.* I. 343. 2. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 428. 1. — *Rande Brachelytr. Hal.* 13. 1. — *Er. Col. March.* I. 518. 1.
- Staph. riparius* Linn. *Faun. Suec.* n. 846. *Iter Gotth.* p. 273. *Syst. Nat.* I. II. 684. 8. — *Degeer Ins.* IV. 28. 14. t. I. f. 18. — *Müll. Prodr.* 99. 1125. — *Goeze Ent. Beitr.* I. 217. 8. — *De Vill. Ent.* I. 413. 8. — *Harr. Besch. d. Ins.* I. 253. 417. — *Fourcr. Ent. Par.* I. 170. 21. — *Payk. Mon. Staph.* 27. 19. — *Marsh. Ent. Britt.* 503. 17.
- Paederus confinis* Zetterst. *Ins. Lappon.* 69. 2.
- Staph. gregarius* Scop. *Ent. Carn.* 102. 308.
- Le Staph. à tête noire et à étuis bleus* Geoffr. *Ins.* I. 369. 21.
- Staph. septimus* Schaeff. *Icon.* t. 71. f. 3.

Habitat in Europa.

*P. longipenni* paulo maior, statura formaque intermedius inter hunc et *P. littoralem*, ab utroque mesosterno rufo distinctus. Antennae capitis thoracisque longitudine, articulo tertio secundo plus sesqui longiore, 4 — 10 sensim paulo brevioribus et crassioribus, ultimo ovato, apice acuminato, nigrae, articulis 4 primis testaceis. Palpi maxillares testacei, articulo tertio apice nigro. Mandibulae rufo-testaceae. Caput thoracis latitudine, orbiculo latum, nigrum, nitidum, supra perparum convexum, utrinque subtiliter parcius punctatum, fronte media late laevi, utrinque inter oculos et ad antennarum insertionem leviter impressa, parce nigro-pilosum. Thorax coleopteris angustior, basin versus sensim angustatus, convexus, rufus, nitidus, dorso biseriatim, lateribus parcissime subtilius punctatus. Scutellum punctulatum, rufo-piceum. Elytra thorace paululum longiora, fortiter punctata, tenuiter griseo-pubescentia, coerulea, nitida. Abdomen parcius punctulatum, tenuiter pubescens, nitidum, rufum, segmentis ultimis duobus nigris. Pectus nigrum, mesosterno rufo. Pedes cum coxis omnibus rufo-testacei, femoribus apice nigris, tarsis apice fusciscentibus.

Obs. Paed. riparius F. a Kirbio in opere splendido *Fauna Boreali-Americana* p. 86. n. 1. inter insecta in America boreali (sub. Lat. 54°) enumeratus, num huic nostro, an potius speciei cuidem confini subiungendus sit mihi haud perspectum.

9. *P. Javanus*: *Alatus, niger, elytris cyaneis, thorace, abdominis segmentis 4 primis mesosternoque rufis, pedibus antennisque testaceis, his medio fuscis, femoribus posterioribus, tibiisque tarsisque omnibus nigris.* — *Long.* 4 lin.

Habitat in Java, collect. Dom. Chevrolat.

Statura et summa affinitas *P. riparii*, at paululum maior. Antennae capitis thoracisque longitudine, articulo tertio secundo duplo longiore, 4 — 10 sensim paulo brevioribus, ultimo oblongo-ovato, apice fortiter acuminato, testaceae, articulis 5 — 9 fuscis. Palpi toti flavo-testacei. Mandibulae rufae. Caput thoracis latitudine, orbiculatum, nigrum, nitidum, supra perparum convexum, fronte utrinque parce inaequaliter punctata, inter oculos utrinque obsolete, ad antennarum insertionem leviter impressa, parce pilosellum. Thorax coleopteris paulo angustior, latitudine paulo longior, basin versus leviter angustatus, lateribus subrectis, convexus, utrinque parce subtiliter punctatus, rufus, nitidus. Scutellum rufum. Elytra thorace paulo

longiora, crebre sat profunde punctata, cyanea, nitida, pilis depressis albidis aliisque erectis nigris tenuiter vestita. Abdomen parce punctatum, nigro-pilosellum, rufum, segmentis ultimis duobus nigris. Pectus nigrum, mesosterno rufo. Pedes testacei, tibiis tarsisque omnibus, femorumque posticorum apice lato nigris.

Obs. A *P. ripario* differt praecipue pedum colore, a *P. peregrino* statura multo maiore, elytris fortius punctatis, nigro-pilosellis, palpis immaculatis.

- \* 10. *P. aestuans*: *Alatus, niger, elytris coeruleis, thorace abdominis segmentis 4 primis, mesosterno pedibusque rufis, femoribus posticis apice nigris, antennis nigris, basi testaceis.* — Long. 3 lin.

Habitat in Senegalia, Dom. Buquet, in Aegypto, Dom. Prof. Ehrenberg.

Statura et summa affinitas *P. longipennis*. Antennae capitis thoracisque longitudine, articulo tertio secundo duplo longiore, 4 — 10 sensim paulo brevioribus et apicem versus magis incrassatis, nigrae, articulis 4 primis flavo-testaceis. Palpi maxillares testacei, articulo tertio apice summo fuscescente. Mandibulae testaceae. Caput thoracis latitudine, suborbiculatum, nigrum, nitidum, parum convexum, utrinque crebrius subtiliusque punctulatum, fronte media late laevi, anterieus utrinque leviter longitudinaliter impressa. Thorax oblongus, coleopteris duplo fere angustior, lateribus ante medium subrotundatis, basin versus vix angustatus, modice convexus, dilute testaceo-rufus, nitidus, dorso biserialim, lateribus disperse subtiliter punctatus. Scutellum rufum. Elytra thorace sesqui fere longiora, crebre punctata, coerulea, nitida, pube brevi depressa albida parcius vestita. Abdomen subtiliter punctulatum, tenuiter cinereo-pubescent, nitidum, testaceo-rufum, segmentis ultimis duobus nigris. Pectus nigrum, mesosterno rufo. Pedes cum coxis omnibus rufo-testacei, femoribus posticis apice nigris, tarsis articulis singulis apice summo nigricantibus.

- \* 11. *P. Sabaeus*: *Alatus, niger, elytris coeruleis, thorace, abdominis segmentis 4 primis, mesosterno pedibusque anterioribus totis, posticis basi rufis, antennis totis testaceis.* — Long. 3 lin.

Habitat in Abessynia et in Arabia felice, Dom. Prof. Ehrenberg, in Guinea, Mus. Dom. Westermann.

Statura et summa similitudo praecedentis. Antennae capite thoraceque paulo longiores, graciles, articulo tertio secundo duplo longiore, 4 — 10 sensim brevioribus et apicem versus paulo magis incrassatis, ultimo oblongo, apice fortiter acuminato, totae testaceae. Caput thorace paululum latius, orbiculatum, nigrum, nitidum, supra perparum convexum, utrinque subtilius crebriusque punctatum, fronte media late laevi, anterieus utrinque leviter longitudinaliter impressa. Thorax oblongus, coleopteris dimidio angustior, lateribus anterieus leviter rotundatis, basin versus angustatus, modice convexus, dilute rufus, nitidus, dorso biserialim, lateribus disperse parce subtiliusque punctatus. Scutellum rufum. Elytra thorace plus sesqui longiora, crebre fortiterque punctata, tenuiter albido-pubescentia, cyanea, nitida. Abdomen parcius minus subtiliter punctulatum, nitidum, testaceo-rufum, abdominis segmentis ultimis duobus nigris. Pectus nigrum, mesosterno rufo.



Pedes anteriores toti rufo-testacei, posteriores coxis femoribusque testaceis, his apice, tibiis tarsisque nigris.

Obs. A praecedente praecipue pedum colore, a sequente antennis longioribus, tenuioribus, totis testaceis, thorace basin versus magis angustato, elytris longioribus et abdomine minus subtiliter punctato satis distinctus.

\* 12. *P. peregrinus*: *Alatus, niger, elytris coeruleis, thorace, abdominis segmentis 4 primis, mesosterno pedumque basi rufis, antennis nigris, basi testaceis.* — Long. 3 lin.

Habitat in Java, Dom. Com. Hoffmannsegg, in Bengala et in Ins. St. Mauriti, Mus. Dom. Westermann.

Statura et summa affinitas praecedentium. Antennae capitis thoracisque fere longitudine, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus at haud crassioribus, ultimo oblongo, apice acuminato, nigrae, articulis tribus primis testaceis, tertio apice nigro. Palpi maxillares testacei, articulo tertio apice summo fuscescente. Mandibulae testaceae. Caput thoracis latitudine, orbiculatum, nigrum, nitidum, supra perparum convexum, utrinque subtilius punctatum, fronte media late laevi, inter oculos et ad antennarum insertionem leviter impressa. Thorax coleopteris dimidio angustior, latitudine tertia parte longior, lateribus ante medium leviter rotundatis, basin versus subangustatus, modice convexus, testaceo-rufus, nitidus, dorso biseriatim, lateribus disperse subtiliter punctatus. Scutellum piceo-nigrum. Elytra thorace sesqui fere longiora, crebre profundeque punctata, tenuiter albido-pubescentia, coerulea, nitida. Abdomen parce subtiliterque punctulatum, subtiliter tenuiterque griseo-pubescent, nitidum, testaceo-rufum, segmentis ultimis duobus nigris. Pectus nigrum, mesosterno rufo. Pedes antici toti rufo-testacei, posteriores coxis femoribusque testaceis, his apice late, tibiis tarsisque nigro-fuscis.

\* 13. *P. littorarius*: *Alatus, niger, elytris cyaneis, thorace, abdominis segmentis 4 primis, mesosterno pedibusque rufis, femoribus posticis apice fuscis, antennis testaceis, medio fuscis.* — Long. 2½ lin.

Grav. Mon. 142. 10.

*Paederus littoralis* Oliv. *Encycl. meth.* VIII. 627. 2.

Habitat in America septentrionali.

Habitu *P. riparii* at triplo minor. Antennae capitis thoracisque longitudine, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus et crassioribus, ultimo oblongo, testaceae, articulis 5 — 9 fuscis. Palpi testacei, immaculati. Mandibulae piceae. Caput thoracis latitudine, orbiculatum, nigrum, nitidum, supra parum convexum, utrinque sparsim punctatum, medio late longitudinaliter laeve, fronte utrinque prope oculos leviter impressa. Thorax coleopteris parum angustior, latitudine tertia parte longior, lateribus ante medium leviter rotundatus, basin versus leviter angustatus, convexus, rufus, nitidus, dorso biseriatim, lateribus disperse parce punctatus. Scutellum rufo-piceum. Elytra thorace parum longiora, fortiter profundeque punctata, parce griseo-pubescentia et nigro-pilosella, nigro-cyanea, nitida. Abdomen parce obsoleteque punctulatum, parce griseo-pubescent, nigro-pilosellum, nitidum, rufum, segmentis ultimis duobus nigris. Pectus nigrum, mesosterno rufo. Pedes cum coxis omnibus rufi, femoribus posticis apice saepius fuscescentibus.

- \* 14. *P. Mexicanus*: *Alatus, niger, elytris cyaneis, thorace, abdominis segmentis 4 primis, mesosterno pedibusque rufis, femoribus posterioribus apice nigris, antennis longioribus, testaceis.* — Long.  $3\frac{2}{3}$  lin.

Habitat in Mexico, Dom. Deppe.

Statura omnino *P. aequinoctialis*. Antennae capite thoraceque paulo longiores, articulo tertio secundo plus duplo longiore, 4 — 10 sensim brevioribus at vix crassioribus, ultimo apice elongato, acuminato, totae testaceae. Palpi testacei, immaculati. Mandibulae nigro-piceae. Caput thorace paulo latius, orbiculatum, nigrum, nitidum, subdepressiusculum, crebrius minus subtiliter punctatum, spatio minuto medio et antico laevibus, utrinque ad antennarum basin impressum. Thorax coleopteris duplo prope angustior, latitudine summa sesqui longior, lateribus anterieus rotundatis, basin versus angustatus, ovatus, convexus, saturate rufus, nitidus, dorso utrinque subseriatim, lateribus disperse parcius subtiliusque punctatus. Scutellum rufum. Elytra thorace sesqui longiora, crebre profundeque punctata, tenuiter cinereo-pubescentia, parce nigro-pilosella, cyanea, nitida. Abdomen coleopteris angustius, apicem versus sensim attenuatum, parce obsoletius punctatum, nigro-pilosellum, nitidum, saturate rufum, segmentis ultimis duobus nigris. Pectus nigrum, mesosterno rufo. Pedes cum coxis omnibus rufi, femoribus posterioribus apice fuscis, tarsis articulis singulis summa apice fuscescentibus.

- \* 15. *P. aequinoctialis*: *Alatus, niger, elytris thorace paulo longioribus cyaneis, thorace, abdominis segmentis 4 primis mesosternoque rufis, pedum basi antennisque longioribus testaceis, his medio nigris.* — Long. 4 lin.

*Paed. columbinus* Laporte Etud. Ent. I. 123. 3. (forte)

Habitat in Columbia, Cayenna.

*P. ripario* nonnunquam paulo maior. Antennae capite thoraceque longiores, articulo tertio secundo duplo longiore, 4 — 10 sensim paulo brevioribus at non crassioribus, ultimo apice abrupte acuminato, testaceae, articulis 4 — 8 piceis, ultimis tribus ferrugineis. Palpi testacei, immaculati. Mandibulae piceae. Caput thoracis latitudine, orbiculatum, nigro-piceum, nitidum, supra depressiusculum, parce subtiliter punctatum, fronte media laevi, nigro-pilosellum. Thorax coleopteris dimidio angustior, latitudine sesqui longior, lateribus anterieus leviter rotundatis, basin versus angustatus, subovatus, convexus, saturate rufus, nitidus, dorso biserialiter, utrinque disperse parcius subtilissime punctatus, subtiliter parceque nigro-pilosus. Scutellum rufum. Elytra thorace paulo longiora, crebre profundeque punctata, tenuiter cinereo-pubescentia, virescenti-cyanea. Abdomen coleopteris paulo angustius, parce punctulatum, omnium subtilissime alutaceum, subtiliter longius nigro-pilosum, subnitidum, saturate rufum, segmentis ultimis duobus nigris. Pectus nigrum, mesosterno rufo. Pedes coxis omnibus rufis, femoribus flavo-testaceis, posterioribus apice late nigris, tibiis tarsisque omnibus nigris.

Variat capite rufo-piceo.

- \* 16. *P. Brasiliensis*: *Alatus, niger, elytris thorace sesqui longioribus, cyaneis, thorace, abdominis segmentis 4 primis mesosternoque rufis, pedibus basi flavo-testaceis, antennis testaceis, medio fuscis.* — Long. 4 lin.

Habitat in Brasilia.

Statura et magnitudo praecedentis. Antennae capite thoraceque paulo longiores, articulo tertio secundo duplo longiore, 4 — 10 sensim paulo brevioribus at vix crassioribus, ultimo apice abrupte acuminato, nigro-piceae, articulis tribus primis flavo-testaceis, ultimis tribus ferrugineis. Palpi maxillares testacei, articulo tertio apice nigricante. Mandibulae piceae. Caput thoracis latitudine, nigrum, nitidum, lateribus longe nigro-pilosum, supra depressiusculum, parce subtiliusque punctatum, fronte media late laevi, anterieus utrinque longitudinaliter leviter impressa. Thorax coleopteris dimidio angustior, latitudine summa sesqui longior, lateribus anterieus leviter rotundatis, basin versus angustatus, ovatus, convexus, rufus, nitidus, dorso biseriatim, lateribus disperse subtiliter punctatus, utrinque nigro-pilosellus. Scutellum rufum. Elytra thorace sesqui longiora, crebre profundeque fortiter punctata, parce cinereo-pubescentia et nigro-pilosella, cyanea, nitida. Abdomen parce subtilius punctulatum, tenuiter nigro-pilosellum, nitidum, rufum, segmentis ultimis duobus nigris. Pectus nigrum, mesosterno rufo. Pedes omnes coxis femoribusque flavo-testaceis, his apice late nigris, tibiis tarsisque piceis, illis basi late nigris.

Obs. Differt a praecedente imprimis statura paulo robustiore, antennis minus elongatis, elytris paulo longioribus et fortius punctatis, tibiis apice et tarsis basi piceis.

- \* 17. *P. tempestivus*: *Alatus, niger, elytris thoracis longitudine, cyaneis, thorace, abdominis segmentis 4 primis mesosternoque rufis, antennis testaceis, medio fuscis, pedibus anticis nigris, basi testaceis, posterioribus testaceis, femoribus apice tibiis basi nigris.* — Long. 4 lin.

Habitat in Columbia, Dom. Moritz, in Brasilia, Dom. Thorey.

Praecedentibus aequalis. Antennae capitis thoracisque longitudine, articulo tertio secundo plus duplo longiore, 4 — 10 sensim paululum brevioribus at vix crassioribus, ultimo oblongo-fusiformi, nigro-fuscae, articulis 3 primis flavo-testaceis, ultimis 3 vel 4 ferrugineis. Palpi rufo-testacei, immaculati. Mandibulae validiores, piceae. Caput thoracis latitudine, orbiculatum, nigrum, nitidum, supra depressiusculum, vage subtilius punctatum, fronte ad antennarum insertionem utrinque obsolete impressa. Thorax coleopterorum latitudine, latitudine sesqui fere longior, lateribus anterieus modice rotundatus, basin versus angustatus, ovatus, sat convexus, saturate rufus, nitidus, vage subtiliter punctatus, linea media longitudinali laevissima. Scutellum obscure rufum. Elytra thoracis longitudine, crebre profundeque punctata, virescenti-cyanea, nitida, parce cinereo-pubescentia et nigro-pilosa. Abdomen coleopterorum latitudine, parcius punctatum, cinereo-nigroque pilosum, saturate rufum, segmentis ultimis duobus nigris. Pectus nigrum, mesosterno rufo. Pedes posteriores dilute testacei, femoribus apice tibiis basi late nigris, antici nigri, coxis obscure rufo-testaceis, femoribus basi flavo-testaceis, tarsis nonnunquam piceis.



Obs. 1. A praecedente praecipue elytris brevioribus et pedibus posterioribus apice dilute testaceis distinctus.

2. Specimen reliquis duplo maius, capite thoraceque grandioribus, ceterum autem haud distinctum, in Mexico in monte Orizaba lectum, nomine P. ruge-punctati Dom. Chevrolat examinandum communicavit.

\* 18. *P. mandibularis*: *Alatus, niger, elytris cyaneis, thorace, abdominis segmentis 4 primis mesosternoque rufis, pedibus antennisque testaceis, medio nigris.* — Long.  $4\frac{1}{2}$  lin.

Habitat in Brasilia meridionali, Sellow.

Praecedentibus paulo maior, robustior. Antennae capitis thoracisque longitudine, articulo tertio secundo plus duplo longiore, 4 — 10 subaequalibus, ultimo oblongo-fusiformi, testaceae, articulis 4 — 8 fuscis. Palpi maxillares testacei, immaculati. Mandibulae validiores, piceae, feminae utraque simplici, maris sinistra incrassata, valida, apice oblique truncata, ante apicem obtuse dentata. Caput thoracis prope latitudine, orbiculato-subquadratum, nigrum, nitidum, supra depressiusculum, parce disperse punctatum, fronte inter oculos obsolete, inter antennis distinctius impressa. Thorax coleopteris parum angustior, latitudine antica haud longior, basin versus modice angustatus, lateribus vix rotundatus, convexus, saturate rufus, dorso biseriatim, lateribus disperse parceque subtiliter punctatus. Scutellum rufo-piceum. Elytra thorace paulo longiora, fortiter profundeque punctata, parce cinereo-pubescentia, cyanea, nitida. Abdomen parce punctulatum, omnium subtilissime alutaceum, parce nigro-pilosum, subnitidum, saturate rufum, segmentis ultimis duobus nigris. Pectus nigrum, mesosterno rufo. Pedes omnes coxis femoribusque testaceis, tibiis apice late nigris, tibiis nigro-piceis, apice tarsisque obscure testaceis.

\* 19. *P. ferus*: *Alatus, niger, elytris cyaneis, thorace paulo longioribus, thorace, abdominis segmentis 4 primis mesosternoque rufis, pedibus nigris, basi flavis, tarsis antennisque testaceis.* — Long. 5 lin.

Habitat in Brasilia, Dom. Virmond.

Statura omnino P. conspicui, at paulo brevior. Antennae capitis thoracisque longitudine, articulo secundo tertio dimidio brevior, 4 — 10 leviter decrescentibus, ultimo oblongo-fusiformi, testaceae, medio fusciscentes, basi flavescens. Palpi testacei, immaculati. Mandibulae piceae, feminae simplices, maris sinistra incrassata, apice dilatata et oblique truncata. Caput thoracis latitudine, orbiculatum, nigrum, nitidum, supra depressiusculum, disperse punctatum, fronte utrinque inter oculos leviter, inter antennis distinctius impressa. Thorax antice coleopteris paulo latior, latitudine summa paulo longior, basin versus fortius angustatus, subovatus, leniter convexus, dorso biseriatim, lateribus disperse subtilius punctatus, rufus, nitidus, parce nigro-pilosellus. Scutellum piceo-rufum. Elytra thorace quadrante fere longiora, fortiter profundeque punctata, cyanea, nitida, parce cinereo-pilosella. Abdomen parce punctatum, nigro-pilosum, nitidum, segmentis 4 primis rufis, reliquis nigris. Pectus nigrum, mesosterno rufo. Pedes coxis testaceis, femoribus basi flavis, apice nigris, tibiis nigris, tarsis testaceis.

Obs. Proximus P. mandibulari, distinctus autem statura paulo maiore, capite haud quadrato, thorace anterieus coleopteris latiore, posterius fortius angustato, tibiis

totis nigris, a praecedentibus statura validiore, coleopteris thorace angustioribus et mandibula sinistra maris dilatata.

- \* 20. *P. conspicuus*: *Alatus, niger, elytris cyaneis, capite, thorace, abdominis segmentis 4 anterioribus mesosternoque rufis, pedibus basi testaceis, antennis testaceis, medio nigris.* — Long.  $5\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Praecedentibus maior. Antennae capitis thoracisque longitudine, articulo tertio secundo plus duplo longiore, 4 — 10 sensim paulo brevioribus at haud crassioribus, testaceae, articulis 4 — 8 nigris. Palpi testacei. Mandibulae rufae, feminae simplices, maris sinistra valida. Caput thoracis latitudine, orbiculatum, rufum, nitidum, versus os piceum, supra depressiusculum, subtiliter parceque punctatum, nigro-pilosellum, fronte media laevi, utrinque inter oculos obsolete, ad antennarum basin distinctius impressa. Thorax antice coleopterorum fere latitudine, latitudineque apicis vix longior, basin versus fortiter angustatus, lateribus subrotundatis, modice convexus, saturate rufus, nitidus, dorso biseriatim, lateribus disperse parceque subtiliter punctatus, nigro-pilosellus. Scutellum rufum. Elytra thorace paulo longiora, fortiter profundeque punctata, nigro-pilosa, cyanea, nitida. Abdomen parce punctulatum, subnitidum, rufum, segmentis ultimis duobus nigris, subtilius nigro-pilosellum. Pectus nigrum, mesosterno rufo. Pedes coxis testaceis, femoribus flavis, apice late nigris, tibiis nigris, tarsis piceis, articulis ultimis duobus testaceis.

Variat fronte antierius nigra.

- \* 21. *P. laetus*: *Apterus, niger, elytris coeruleis, thorace, abdomine toto, mesosterno pedibusque rufis, his geniculis posterioribus nigris, antennis longioribus, totis testaceis.* — Long.  $3\frac{2}{3}$  lin.

Habitat in Mexico, Dom. Deppe.

Statura omnino *P. littoralis*, at elytris longioribus. Antennae capite thoraceque longiores, articulo tertio secundo duplo longiore, 4 — 10 sensim paulo brevioribus at haud crassioribus, ultimo elongato, apice abrupte acuminato, totae testaceae. Palpi testacei, immaculati. Mandibulae rufo-piceae. Caput thoracis latitudine, orbiculatum, nigrum, nitidum, supra depressiusculum, parce disperseque punctatum, utrinque inter oculos obsolete, ad antennarum basin distinctius impressum, nigro-pilosellum. Thorax coleopterorum latitudine, latitudine summa vix longior, basin versus leviter angustatus, lateribus rotundatus, subglobosus, convexus, testaceo-rufus, nitidus, dorso biseriatim, lateribus disperse parce subtilissime punctatus, parce subtiliterque nigro-pilosus. Scutellum rufum. Elytra thorace paulo longiora, fortiter punctata, parce cinereo-pubescentia, nigro-pilosella, coerulea, nitida. Abdomen parce obsoletiusque punctulatum, nigro-pilosellum, subnitidum, totum testaceo-rufum. Pectus nigrum, mesosterno rufo. Pedes cum coxis omnibus rufo-testacei, posteriores femoribus apice, tibiis basi nigris.

22. *P. melampus*: *Alatus, niger, elytris coeruleis, thorace abdominisque segmentis 4 primis rufis, palpis pedibusque nigris, basi mesosternoque piceis.* — Long. 3 lin.

Habitat in India orientali, Mus. Dom. Hope.

Statura *P. ruficollis* at dimidio minor. Antennae articulis duobus primis obscure testaceis (reliqui desunt). Palpi nigri, articulo primo piceo. Caput thoracis latitudine, orbiculatum, subcoerulescenti-nigrum, nitidum, supra parum convexum, vage subtiliter punctulatum, anterius utrinque late impressum. Thorax coleopteris paulo angustior, latitudine summa sesqui longior, lateribus ante medium leviter rotundatis, basin versus angustatus, ovatus, convexus, subtiliter punctatus, linea media longitudinali laevi, rufus, nitidus. Scutellum nigrum. Elytra thorace paulo longiora, dense minus fortiter punctata, tenuiter nigro-pubescentia. Abdomen segmentis anterioribus 4 rufo-testaceis, ultimis duobus nigris. Pectus nigrum, mesosterno piceo. Pedes nigri, coxis, trochanteribus femorumque summa basi piceis.

23. *P. extraneus*: *Niger, elytris cyaneis, thorace abdominisque segmentis 4 primis rufis, antennis palpisque basi piceis.* — Long.  $3\frac{2}{3}$  lin.

Wiedem. Zool. Mag. II. I. 133. 200.

Habitat in Bengala, Mus. Dom. Westermann.

Statura omnino *P. riparii*, at paululum maior. Antennae capitis thoracisque prope longitudine, articulo tertio secundo duplo prope longiore, 4—10 sensim brevioribus at vix crassioribus, ultimo elongato, nigrae, articulis duobus primis obscure rufo-testaceis, superne picescentibus. Palpi piceo-testacei, articulo tertio nigro. Mandibulae nigrae. Caput thoracis latitudine, orbiculatum, supra perparum convexum, crebrius obsolete subtiliter punctulatum, nigrum, nitidum. Thorax breviter ovatus, coleopterorum prope latitudine, latitudine summa parum longior, lateribus leviter rotundatus, basin versus angustatus, convexus, crebre obsoletius subtiliter punctatus, linea media longitudinali laevi, rufus, nitidus. Scutellum nigrum. Elytra thorace vix quarta parte longiora, crebre minus fortiter punctata, albo-pubescentia, cyanea, nitidula. Abdomen supra subtiliter punctatum, longius albido-pubescentis, segmentis 4 prioribus rufis, reliquis nigris. Pectus nigrum. Pedes totum cum coxis nigri.

24. *P. tamulus*: *Alatus, niger, capite elytrisque chalybeis, thorace abdominisque segmentis 4 primis rufis.* — Long. 3 lin.

Habitat Leontopoli (Singhapur) in India orientali, Mus. Dom. Hope.

Habitu *P. riparii*, at paululum minor. Antennae capitis thoracisque longitudine, articulo tertio secundo duplo longiore, 4—10 sensim paulo brevioribus et paululum crassioribus, ultimo oblongo-fusiformi, nigrae, cinereo-pubescentes, articulis duobus primis subtestaceis. Palpi nigri. Caput thoracis latitudine, orbiculatum, chalybeum, nitidum, supra depressiusculum, subtiliter crebre punctulatum, fronte media laeviore, parcius cinereo-pubescentis parceque nigro-pilosum. Thorax coleopteris dimidio angustior, latitudine summa parum longior, lateribus rotundatus, basin versus angustatus, breviter ovatus, convexus, crebre subtiliter punctatus, linea longitudinali media laevi, rufus, nitidus, subtilissime albido-pubescentis. Scutellum nigrum. Elytra thorace paulo longiora, dense subtilius punctata, nigro-coerulea, nitida, densius albido-pubescentia. Abdomen tenuiter cinereo-pubescentis, parce nigro-pilosum, segmentis 4 primis rufis, reliquis



nigris. Pectus nigrum, albedo-pubescent. Pedes toti cum coxis omnibus nigris, albedo-pubescentes.

Obs. Affinis praecedenti, at duplo fere minor, caput distinctius punctulatum, cyaneum, thorax magis oblongus, etc.

\* 25. *P. cyanocephalus*: *Alatus, niger, capite elytrisque cyaneis, thorace abdominisque segmentis 4 primis rufis, palpis pedibusque nigris, basi testaceis, mesosterno piceo.* — Long. 3 lin.

Habitat in Siam, Dom. Kaden.

Statura omnino praecedentium. Antennae capite thoraceque longiores, graciles, articulo tertio secundo plus duplo longiore, 4 — 10 sensim paulo brevioribus at vix crassioribus, ultimo elongato, nigrae, articulis duobus primis rufo-piceis. Palpi piceo-testacei, articulo tertio nigro. Mandibulae rufo-piceae, apice rufescentes. Caput thorace paulo latius, orbiculatum, supra perparum convexum, disperse punctatum, punctis medio rarioribus, anterius utrinque leviter longitudinaliter impressum, parce albedo-pilosum, cyaneum, nitidum. Thorax coleopteris duplo prope angustior, latitudine summa plus duplo longior, lateribus leviter rotundatus, basin versus angustatus, oblongo-ovatus, convexus, disperse subtiliter punctatus, rufus, nitidus. Scutellum piceum. Elytra thorace paulo longiora, parce obsoleteque punctata, sublaevigata, parcius albedo-pubescentia, cyanea vel chalybea, nitida. Abdomen parce subtiliter obsoleteque punctulatum, longius albedo-pubescent, segmentis 4 prioribus rufo-testaceis, reliquis nigris. Pectus nigrum, mesosterno rufo-piceo. Pedes densius albedo-pubescentes, coxis, trochanteribus femoribusque rufo-testaceis, his apice, tibiis tarsisque nigris.

\* 26. *P. ruficollis*: *Alatus, nigro-cyaneus, elytris cyaneis, thorace rufo, antennis, palpis pedibusque nigris.* — Long.  $3\frac{1}{2}$  —  $3\frac{3}{4}$  lin.

*Fab. Spec. Ins. I. 339. 2. Mont. Ins. 223. 2. Ent. Syst. I. II. 537. 2. Syst. El. II. 608 2. — Payk. Faun. Suec. III. 427. 1. — Oliv. Ent. III. 44. 4. 1. t. I. f. 1. Encycl. meth. VIII. 627. 3. — Panz. Faun. Germ. 27. 22. Ent. Germ. 362. 2. — Grav. Micr. 185. 2. Mon. 143. 13. — Latr. Hist. nat. Crust. et Ins. IX. 345. 1. — Gyll. Ins. Suec. II. 373. 2. — Mann. Brachelytr. 39. 4. — Boisd. et Lacord. Faun. Ent. Paris. I. 430. 4. — Guér. Iconogr. Regn. An. Ins. t. 9. f. 5. — Runde Brachelytr. Hal. 13. 3.*

*Staph. ruficollis* Payk. Mon. Staph. 26. 18.

*Staph. thoracicus* Fourcr. Ent. Par. I. 170. 23.

*Le Staph. noir à corcelet rouge* Geoffr. Ins. I. 370. 23.

Habitat in Europa et in Asia occidentali (Persia).

*P. ripario* fere aequalis, nigro-cyaneus, nitidulus, pube subtilissima albida sericans, antennis, palpis pedibusque nigris, thorace solo rufo. Antennae capite thoraceque paulo longiores, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus, ultimo elongato-fusiformi. Caput thoracis latitudine, orbiculatum, supra parum convexum, subtiliter punctatum, inter oculos et inter antennis utrinque leviter impressum, breviter nigro-pilosum. Thorax coleopteris plus dimidio angustior, latitudine summa sesqui prope longior, lateribus ante medium rotundatis, basin versus angustatus, ovatus, convexus, subtiliter vage punctatus, linea media longitudinali laevi. Scutellum nigrum. Elytra thorace tertia parte longiora, crebrius subtiliusque punctata, cyanea. Abdomen subtiliter punctulatum.

- \* 27. *P. tricolor*: *Alatus, niger, elytris cyaneis, thorace mesosternoque rufis, antennis pedibusque basi testaceis.* — Long.  $4\frac{1}{2}$  lin.

Habitat in ins. St. Vincentii Americae, Dom. Hope.

*P. ruficollis* paulo maior et gracilior. Antennae capite thoraceque longiores, articulo tertio secundo duplo longiore, 4 — 10 sensim paulo brevioribus, ultimo oblongo-fusiformi, nigrae, articulis tribus primis flavo-testaceis, ultimo ferrugineo. Palpi maxillares rufi, articulo tertio apice late nigro. Mandibulae piceae. Caput thoracis latitudine, orbiculatum, nigrum, nitidum, supra depressiusculum, subtiliter parceque punctulatum, fronte anteriori utrinque leviter longitudinaliter impressa, parce nigro-pilosellum. Thorax antice coleopteris vix angustior, latitudine antica vix longior, basin versus modice angustatus, lateribus leviter rotundatis, ovatus, convexus, testaceo-rufus, nitidus, utrinque parce subtilissime punctatus. Scutellum obscure rufum. Elytra thorace paulo longiora, crebre profundius punctata, subrugosa, cyanea, nitida, tenuissime griseo-pubescentia, parce nigro-pilosella. Abdomen nigrum, subnitidum, parce punctulatum, nigro-pilosellum. Pectus nigrum, mesosterno rufo. Pedes coxis femoribusque testaceis, his apice late nigris, tibiis tarsisque nigris.

28. *P. erythroderus*: *Apterus, niger, elytris cyaneis, thorace mesosternoque rufis, pedibus antennisque testaceis, his medio, femoribus omnibus apice nigris.* — Long.  $4\frac{1}{2}$  lin.

Habitat in monte Orizaba in Mexico, collect. Dom. Chevrolat.

Antennae capitis thoracisque longitudine, articulo tertio secundo duplo prope longiore, 4 — 10 sensim paululum brevioribus, ultimo apice abrupte acuminato, articulis tribus primis flavo-testaceis, 4 — 9 fuscis, ultimis duobus rufo-testaceis. Palpi testacei, immaculati. Mandibulae piceae. Caput thoracis latitudine, orbiculatum, nigrum, nitidum, supra depressiusculum, vage obsolete punctatum, fronte media laevi, anteriori utrinque leviter longitudinaliter impressa, parce nigro-pilosellum. Thorax coleopteris latior, latitudine antica vix longior, basin versus subangustatus, lateribus leviter rotundatus, valde convexus, dilute rufus, nitidus, lateribus parce vageque obsolete punctatus. Scutellum piceum. Coleoptera antrorsum leviter angustata, parce obsoletius exasperato-punctata, viridi-cyanea; nitida, nigro-pilosella. Abdomen parce punctatum, nigrum, subnitidum, nigro-pilosellum. Pectus nigrum, mesosterno rufo. Pedes antici rufo-testacei, coxis concoloribus, femoribus apice nigris, postici flavo-testacei, femoribus apice coxisque nigris.

- \* 29. *P. thoracicus*: *Apterus, niger, elytris cyaneis, thorace rufo, antennis pedibusque testaceis.* — Long.  $4\frac{1}{2}$  — 5 lin.

Habitat in Mexico.

Habitu omnino *P. littoralis* at duplo fere maior. Antennae capitis thoracisque fere longitudine, articulo tertio secundo duplo longiore, 4 — 10 sensim paulo brevioribus, ultimo oblongo-ovato, totae testaceae. Palpi rufo-testacei. Mandibulae piceae. Caput thoracis latitudine, orbiculatum, supra leviter convexum, crebrius punctatum, anteriori utrinque leviter longitudinaliter impressum, nigrum, nitidum, nigro-pilosellum. Thorax coleopteris

paulo latior, latitudine antica haud longior, basin versus leviter angustatus, lateribus subrotundatus, valde convexus, subglososus, vage subtiliter punctatus, linea media longitudinali laevi, rufus, nitidus. Scutellum rufo-piceum. Coleoptera thorace fere breviora, basin versus sensim leviter angustata, apice coniunctim emarginata, crebrius fortiusque exasperato-punctata, cyanea vel viridi-cyanea, nitida, nigro-pilosella. Abdomen apicem versus modice incrassatum, sat crebre minus fortiter punctatum, nigrum, subnitidum, parcius nigro-pubescent. Pectus nigrum. Pedes nigri, coxis anticis rufis, tarsis omnibus piceo-testaceis, articulis ultimis duobus dilute testaceis.

- \* 30. *P. chalybeus*: *Alatus, nigro-cyaneus, abdomine, pedibus antennisque nigris, his articulo secundo basi piceo.* — Long. 4 lin.

Habitat in Madagascar, Dom. Goudot.

Statura omnino *P. ruficollis* et maximis eius individuis parum maior, niger, capite, thorace elytrisque obscure cyaneis, subnitidulus. Antennae capitibus thoracisque latitudine, articulo secundo subtus rufo-piceo, tertio secundo plus duplo longiore, 4 — 10 sensim paulo brevioribus, ultimo elongato-fusiformi. Palpi nigri, articulo secundo basi piceo. Caput thoracis latitudine, orbiculatum, supra leviter convexum, crebrius subtiliusque vage punctatum, fronte media vix laeviore, anterieus utrinque longitudinaliter leviter impressa, parce nigro-pilosellum. Thorax coleopteris duplo fere angustior, latitudine summa sesqui longior, lateribus ante medium rotundatis, basin versus angustatus, ovatus, valde convexus, subtiliter vage punctatus, linea media longitudinali laevi, parce breviter subtiliterque nigro-pilosellus. Scutellum nigrum. Elytra thorace longiora, crebrius subtiliter punctata, tenuiter cinereo-pubescentia. Abdomen subtiliter punctulatum, subtiliter cinereo-nigroque pubescens.

- \* 31. *P. coerulescens*: *Alatus, nigro-coeruleus, abdomine, pedibus antennisque nigris, his articulis duobus primis subtus testaceis.* — Long. 2½ lin.

Habitat in Madagascar, Dom. Goudot.

Praecedente plus duplo minor, niger, capite, thorace elytrisque nigro-coeruleis, subnitidus, pube subtili brevique albida subsericans. Antennae capite thoraceque paulo longiores, articulo tertio secundo duplo prope longiore, 4 — 10 sensim brevioribus, ultimo oblongo, apice acuminato, nigrae, articulis duobus primis subtus testaceis. Palpi maxillares nigri, articulo secundo toto testaceo. Caput thoracis latitudine, orbiculatum, supra convexum, crebre subtilius punctatum, punctis in media fronte rarioribus, antice utrinque longitudinaliter impressum, parce nigro-pilosum. Thorax coleopteris plus dimidio angustior, latitudine summa tertia parte longior, lateribus ante medium rotundatis, basin versus angustatus, ovatus, convexus, aequaliter densius subtiliusque punctatus. Scutellum nigrum. Elytra thorace sesqui longiora, densius subtiliusque punctata, transversim subrugosa. Abdomen subtiliter punctulatum.

- \* 32. *P. memnonius*: *Alatus, niger, albido-pubescent, elytris cyaneis, antennis, palpis pedibusque nigris, immaculatis.* — Long. 3 lin.

Habitat in Aegypto, Dom. Prof. Ehrenberg.



Statura *P. ruficollis*, et minimis eius individuis aequalis, niger, subnitidus, capite thoraceque nitidis, elytris cyaneis, nitidis, pube crassiore depressa albida nitidula densius vestitus. Antennae capitis thoracisque longitudine, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus et magis obconicis, ultimo fusiformi, totae nigrae, albido-sericantes. Palpi nigri, immaculati, albido-sericantes. Caput thoracis latitudine, orbiculatum, supra leviter convexum, crebre punctatum, fronte media sublaevi, anterius utrinque leviter longitudinaliter impressa, parce nigro-pilosum. Thorax coleopteris duplo prope angustior, latitudine summa tertia parte longior, lateribus ante medium rotundatis, basin versus modice angustatus, ovatus, convexus, crebrius subtiliusque punctatus, linea media longitudinali laevi parum distincta. Scutellum nigrum. Elytra thorace paulo longiora, crebre fortius punctata. Abdomen subtiliter punctulatum.

\* 33. *P. fastuosus*: *Alatus, niger, elytris cyaneis, antennis palpisque testaceis.* — Long.  $3\frac{3}{4}$  lin.

*Klug Col. Madag.* 52. 45.

Habitat in Madagascar, Dom. Goudot.

Statura fere *P. littoralis*, at elytris longioribus, niger, nitidus, elytris vel virescenti- vel violaceo-cyaneis. Antennae capitis thoracisque longitudine, articulo tertio secundo duplo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, ultimo fusiformi, totae testaceae, apice saepius obscuriores. Palpi testacei. Mandibulae rufae. Caput thoracis latitudine, orbiculatum, supra depressiusculum, utrinque crebrius punctatum, medio longitudinaliter laeve, anterius utrinque longitudinaliter impressum. Thorax coleopteris parum angustior, latitudine antica vix longior, basin versus leviter angustatus, lateribus vix rotundatus, modice convexus, utrinque crebrius disperse subtilius punctatus, linea media longitudinali laevi. Scutellum nigrum. Elytra thorace paulo longiora, fortiter profundeque crebre punctata, parce nigro-pilosella. Abdomen parcius subtiliusque punctulatum, nigropilosellum. Tarsi picei.

\* 34. *P. fulvicornis*: *Apterus, niger, elytris brevibus, nigro-cyaneis, tibiis piceis apice, tarsis, antennis palpisque testaceis.* — Long. 4 lin.

Habitat in Java, Dom. Dr. De Haen.

Statura fere *P. littoralis*, at sesqui maior, niger, nitidus, elytris nigro-coeruleis. Antennae capite thoraceque paulo longiores, articulo tertio secundo duplo prope longiore, 4 — 10 sensim paulo brevioribus, ultimo elongato-fusiformi, fortiter acuminato, rufo-testaceae. Palpi rufo-testacei. Mandibulae piceae. Caput thorace paulo latius, orbiculatum, supra depressiusculum, parce subtiliterque punctatum, anterius utrinque fortius impressum. Thorax coleopterorum latitudine, latitudine summa paulo longior, lateribus anterius rotundatis, basin versus angustatus, fortiter convexus, subglobosus, utrinque parce disperseque subtilius punctatus, medio tenuissime obsolete canaliculatus. Scutellum nigrum. Coleoptera vix thoracis longitudine, basin versus angustata, parum exasperato-punctata, tenuiter griseo-pubescentia, parce nigro-pilosella. Abdomen apicem versus sensim leniter incrassatum,

minus nitidum, parce obsoletius punctatum, fusco-pilosellum. Pedes nigri, tibiis piceis apice tarsisque fulvis.

\* 35. *P. rutilicornis*: *Apterus, niger, elytris longioribus, nigro-subviolaceis, antennis palpisque testaceis.* — Long.  $5\frac{1}{2}$  lin.

Habitat Bogotae in Columbia interiore, Dom. Buquet.

Habitu *P. littoralis*, at magnus, niger, nitidus, elytris nigro-subcaneis, aut nigro-violaceis, aut nigris, parce nigro-pilosus. Antennae capitis thoracisque longitudine, articulo tertio secundo duplo prope longiore, 4 — 10 sensim brevioribus, ultimo fusiformi, apice acuminato, testaceae. Palpi maxillares testacei, labiales picei. Mandibulae nigro-piceae. Caput thoracis latitudine, subovatum, supra depressiusculum, parce subtiliterque punctatum, fronte media laeviore, antice utrinque leviter longitudinaliter impressa. Thorax coleopteris parum angustior, latitudine summa tertia parte longior, lateribus leviter rotundatus, basin versus angustatus, ovatus, convexus, utrinque disperse parceque subtiliter punctatus, plaga angusta longitudinali media laevi. Scutellum nigrum. Elytra thoracis longitudine, crebre fortiter profundeque punctata, subrugosa. Abdomen apicem versus sensim incrassatum, parcius subtiliusque punctatum, tenuiter nigro-pilosellum. Pedes nigri, tibiis tarsisque piceis, subtus fulvescenti-tomentosis.

\* 36. *P. coarctatus*: *Apterus, niger, elytris brevibus, viridibus, abdomine apice obscure sanguineo, antennis palpisque testaceis.* — Long. 5 lin.

Habitat in Brasilia, Dom. Prof. Germar.

Praecedente paulo minor. Antennae capite thoraceque longiores, articulo tertio secundo duplo longiore, sequentibus sensim paulo brevioribus, ultimo apice abrupte acuminato, testaceae, medio obscuriores, nigro-pilosellae. Palpi testacei. Mandibulae piceae. Caput thoracis latitudine, orbiculatum, nigrum, nitidum, nigro-pilosellum, supra depressiusculum, parce subtiliterque punctatum, fronte media late laevi, antice utrinque leviter impressa. Thorax coleopteris multo latior, latitudine summa paulo longior, lateribus leviter rotundatus, basin versus angustatus, valde convexus, niger, nitidus, dorso biseriatim, lateribus disperse parceque subtiliter punctatus, nigro-pilosellus. Scutellum nigrum, laeve. Coleoptera angusta, thorace haud longiora, fortiter profundeque punctata, viridia, nitida, nigro-pilosella. Abdomen apicem versus leniter incrassatum, parcius punctatum, nigro-pilosellum, nigrum, nitidum, segmentis duobus ultimis obscure sanguineis. Pectus nigrum. Pedes picei, femoribus apicem versus sensim nigris.

### Species mihi invisae.

37. *P. australis*: *Rufus, capite abdominisque apice pedibusque nigris, elytris coeruleis.*

*Guér. Voyag. de la Coquille t. 1. f. 23. — Boisd. Faun. Ent. Ocean. p. 57.*

Habitat ad Portum Jacksoni Novae Hollandiae.

*P. ripario* similis.

38. *P. morio*: *Ater, opacus, elytris brevissimis, coerulescentibus, thorace obovato, postice truncato, antennis ferrugineis.*

*Mannerh. Bruchelytr.* 39. 1.

Habitat in ins. St. Dominici.

*P. ripario* duplo maior. Caput thorace multo latius, subdepressum, disperse punctatum, fronte obsoletius bi-impressa. Antennae tenues, capite cum thorace longiores. Thorax latitudine longior, ovatus, basi truncatus, convexus, disperse punctatus. Elytra thorace angustiora, latitudine breviora, apice introrsum oblique truncata, punctis leviter insculptis pilosis. Abdomen reliquo corpore fere longius, inflatum. Pedes praesertim postici, elongati. (Mannerh.)



## Tribus quinta.

### **P i n o p h i l i n i .**

*Stigmata prothoracica occulta, spatio pone coxas anticus corneo.*

*Coxae posticae conicae.*

*Antennae sub frontis margine laterali insertae.*

---

Labrum latum, transversum vel brevissimum,  
integrum Pinophilo, Taenodemati;  
medio sinuatum, Palamino, Oedichiro, Procirro.

Mandibulae longae, tenues, falcatae, acutae, medio dentatae.

Maxillae malis brevioribus, interiore interne, exteriore apice barbatis.

Palpi maxillares articulo primo minuto, secundo elongato, tertio  
hoc plerumque brevior, quarto magno,

transverso, lineari Pinophilo;

transverso, securiformi Taenodemati, Palamino, Oedichiro;  
elongato-fusiformi, acuminato Procirro.

Mentum corneum, transversum, breve.

Ligula brevis, biloba, plerisque;  
apice subsinuata Procirro.

Paraglossae distinctae,  
ligulam superantes, lineares Pinophilo, Taenodemati, Procirro;  
ligulae aequales Oedichiro;  
ligula breviores Palamino.

Palpi labiales articulo tertio minuto, fere subulato.

Antennae rectae, filiformes, 11-articulatae, infra oculos sub capitis  
margine laterali insertae.

Caput exsertum, collo quasi tenui thoraci affixum,  
basi truncatum Pinophilo, Palamino;  
basi rotundatum Oedichiro;  
basi attenuatum Procirro.

Thorax basi ab elytris remotus, subtus spatio pone coxas anticas corneo.

Elytra pectoris longitudine.

Abdomen elongatum, segmento septimo saepius occulto,

marginatum Pinophilo, Taenodemati;  
 immarginatum Palamino, Oedichiro, Procirro.

Pedes graciliores, mutici, coxis anticis conicis, exsertis, intermediis approximatis, posticis conicis, tarsis 5-articulatis, posterioribus articulo primo leviter elongato, anticis dilatatis,

articulis 4 primis patellam formantibus, singulis transversis, subtus membrana instructis Pinophilo, Palamino;

tomentoso-spongiosis Taenodemati;

articulis 4 primis singulis dilatatis, subquadratis, fortiter convexis Oedichiro, Procirro.

#### O b s e r v a t i o.

Affinis tribui praecedenti et sequenti, cum utraque mandibulis longis, tenuibus, falciformibus et palpis maxillaribus elongatis, cum illa palporum maxillarum articulo primo brevi et coxis anticis oblongis, conicis, cum hac spatio pone coxas corneo convenit, ab utraque palporum maxillarum articulo ultimo magno, plerumque magis minusve securiformi, discedit.

### Pinophilus Grav.

*Grav. Micr. p. 201. — Laporte Etud. Ent. I. p. 121.*

*Pinophilus Nordm. Symb. p. 154. t. 2. f. 6. et Araecerus Nordm. ibid. p. 157.*

*Pityophilus Brullé Hist. nat. Ins. III. p. 75.*

*Lathrobium Fam. I. Grav. Mon. p. 129.*

*Palpi maxillares articulo quarto transverso, lineari.*

*Abdomen marginatum.*

*Tarsi antici articulis 4 primis dilatatis, coniunctis patellam formantibus.*

Corpus mediae magnitudinis, elongatum, lineare, alatum. Caput thoracis prope latitudine, subquadratum, basi profunde constrictum, collo quasi brevi thoraci immissum, exsertum, porrectum, subdepressum, oculis medio-cribus, rotundatis, prominulis. Labrum latum, brevissimum, integrum. Mandibulae sat longae, falcatae, acutae, medio dente valido truncato armatae. Maxillae (t. IV. f. 34. b.) malis coriaceis, interiore interne, exteriori apice barbatis. Palpi maxillares elongati, articulo primo brevi, minuto, secundo tertioque subaequalibus, quarto his magnitudine aequali, at transverso, in trorsum directo, facie palpitante per totam articuli latitudinem extensa. Labium (t. IV. f. 34. a.) mento transverso, ligula brevi, biloba, membranacea, paraglossis acuminatis, membranaceis, ligulam longe superantibus. Palpi labiales articulo primo maiore secundoque leviter elongatis, tertio parvo, subacuminato. Antennae filiformes, tenues, articulis singulis apice incrassatis, primo fortiore, secundo saepius sequentibus paulo crassiore, ultimo a medio inde abrupte acuminato. Thorax quadrangulus, plerumque basin versus angustatus, basi apiceque truncatus, angulis anterioribus subrectis, posterioribus obtusis vel rotundatis. Scutellum rotundatum. Elytra truncata. Abdomen lineare, coleopterorum latitudine, teretiusculum, marginatum, stylis analibus haud exsertis. Pedes breviores, antici incrassati, femoribus tibiisque interne sinuatis, his apice oblique truncatis, tarsis articulis quatuor primis fortiter dilatatis, subtus articulum singulo lobo transversali semicirculari membraneo, saepius ultra articulum ipsum dilatato, instructo, posteriores graciliores, tibiis apice oblique truncatis, posticis apicem versus subdilatatis,

compressis, tarsis articulo primo modice elongato, sequentibus duobus coniunctis aequali, quarto parvo, lobato.

### Observationes.

1. Corpus punctulatum, plerumque tenuiter pubescens, capite thoraceque nonnunquam glabris, thorace dorso in nonnullis biserialim punctato, fronte saepius posterius et ante oculos utrinque fortius punctata.

2. Genus *Araeocerus*, corporis subdepressi, thoracis parum convexi, elytrorum planorum causa a Nordmanno distinctum, omnibus characteribus cum hoc genere congruit, neque habitu etiam satis dissimile videtur.

3. Dom. Brullé e latino vocabulo *Pinus* generis huius nomen a Gravenhorstio compositum esse opinatus, ut vocem hybridam e systemate elideret, *Pinophilum* in *Pityophilum* mutavit. Cum autem eruditum huius generis inventorem nomen illud e graeco *πινος* (squalor) composuisse patere videatur, non est quod istam a Dom. Brullé adhibitam emendationem recipiamus.

4. Postea cel. Gravenhorst ipse hoc genus *Lathrobio* subiunxit (Mon. p. 129.), quo forte evenit, quod exinde a plurimis auctoribus sit oblitum. Dom. Leach nomen novum, *Lycidius*, proposuit, a plurimis Entomologis receptum, in Catalogo collectionis Dejeaniana lapsu typographico *Lyeidius* dictum.

5. Mas pene prominulo, corneo, acuto, sursum curvato saepius cognoscendus, plerumque quoque abdominis segmenti inferioris sexti, interdum etiam quinti structura distinctus.

6. De victu nihil constat. Genus Europae omnino alienum, in America, Africa et Asia obvium.

1. *P. tenebrosus*: *Niger, subopacus, antennis, palpis tarsisque fusco-testaceis, thorace oblongo-quadrato, dorso biserialim punctato.* — *Long. 11 lin.*

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Subcylindricus, niger, subopacus, densius griseo-pubescens, capite thoraceque glabris, parum nitidis. Antennae capite sesqui longiores, fusco-testaceae, articulo tertio nigricante, secundo plus sesqui longiore, sequentibus sensim decrescentibus. Palpi fusco-testacei. Caput thorace paulo angustius, lateribus pone oculos crebre punctatum, fronte punctis paucis dispersis, secundum latera crebrioribus et confertioribus, impressa. Thorax coleopterorum latitudine, latitudine paulo longior, basin versus subangustatus, lateribus rectis, basi apiceque truncatus, angulis praecipue posterioribus rotundatis, leviter transversim convexus, punctis parvis, dorso subbiserialis, ad latera dispersis, impressus, interstitiis omnium subtilissime punctulatis. Scutellum punctulatum. Elytra thorace longiora, dense minus fortiter punctata. Abdomen confertim subtilius punctatum, apice summo piceum. Pedes nigri, griseo-pubescentes, tarsis ferrugineis, anticis minoribus, tibia sesquilatioribus, orbiculatis.

Abdominis segmentum sextum inferius maris longitudinaliter impressum, apice profundius emarginatum, quantum apicem versus leviter longitudinaliter impressum.

- \* 2. *P. torosus*: *Niger, nitidus, antennis tarsisque fusco-testaceis, thorace subquadrato, vage punctato.* — *Long. 10 lin.*

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Praecedenti aequalis, licet paulum minus elongatus, subcylindricus, ni-



ger, subnitidus, capite thoraceque pernitidis, glabris. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, sequentibus sensim decrescentibus, fusco-testaceae, articulo primo nigro-piceo. Palpi fusco-picei. Caput thoracis prope latitudine, fronte disperse sat fortiter punctata. Thorax coleopterorum latitudine, latitudine haud longior, basin versus subangustatus, lateribus subrectis, basi apiceque truncatus, angulis praecipue posterioribus rotundatis, leviter convexus, sparsim sat profunde punctatus, plagis tribus longitudinalibus laevioribus. Scutellum obsoletius parum-punctatum. Elytra thorace paulo longiora, crebre profunde punctata, parcius nigro-pubescentia. Abdomen densius nigro-pubescentia, creberrime minus subtiliter punctatum, segmentis ultimis duobus punctato-rugosis, ultimo apice acute emarginato. Pedes nigri, tarsis piceis, anticis tibia parum latioribus, orbiculatis.

Abdominis segmentum sextum inferius maris leviter longitudinaliter impressum, apice profunde incisum, quintum apicem versus leviter canaliculatum, feminae integra, sextum crebre fortius punctatum, longitudinaliter subrugosum.

3. *P. caliginosus*: *Niger, nitidus, antennis, palpis tarsisque fusco-testaceis, thorace oblongo, crebre punctato, linea media longitudinali laevi.* — *Long. 7 lin.*

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Praecedentibus minor et angustior, subcylindricus, niger, subnitidus, griseo-pubescentia, capite thoraceque glabris, pernitidis. Antennae capite thoraceque multo breviores, articulo tertio secundo parum longiore, 3 — 5 subaequalibus, sequentibus sensim decrescentibus, fusco-testaceae, articulis duobus primis piceis. Palpi fusco-testacei. Caput thoracis paulo angustius, fronte disperse sat profunde punctata. Thorax coleopterorum latitudine, latitudine longior, basin versus subangustatus, lateribus rectis, basi apiceque truncatus, angulis praecipue posterioribus subrotundatis, leviter transversim convexus, crebre punctatus, punctis prope marginem lateralem et prope lineam mediam longitudinalem laevem magis confertis. Scutellum parce punctulatum. Elytra thoracis latitudine, sat dense punctata, griseo-pubescentia. Abdomen confertim punctatum, griseo-pubescentia. Pedes nigri, tarsis fusco-testaceis, anticis tibia sesqui latioribus, orbiculatis.

Mas latet.

- \* 4. *P. picipes*: *Niger, nitidus, pedibus rufo-piceis, thorace oblongo, dorso biseriatim punctato.* — *Long. 7 lin.*

Habitat Novi Aureliani in Georgia Americae, Dom. Philippi.

Subcylindricus, niger, subnitidus, capite thoraceque glaberrimis, pernitidis. Antennae articulis 2 primis rufis, secundo tertio paulo brevioribus, 3 — 4 aequalibus, quintoque iterum paulo brevioribus (reliqui desunt). Palpi rufi. Caput thoracis paulo angustius, vertice et oculorum margine interno crebrius punctatis, fronte punctis paucis dispersis impressa. Thorax coleopteris paululum latior, latitudine paulo longior, basin versus vix angustatus, lateribus rectis, basi apiceque truncatus, angulis omnibus leviter rotundatis, transversim convexus, omnium subtilissime punctulatus, dorso

serie utrinque longitudinali sat regulari e punctis 6 — 7, lateribus punctis paucis dispersis impressus. Scutellum laeve. Elytra thoracis longitudine, fortiter profundeque punctata, parce pubescentia. Abdomen crebre subtilius punctatum, pube longiore depressa parce vestitum. Pedes crassiusculi, rufo-picei, coxis nigris, femoribus posterioribus basi nigricantibus, tarsis anticis tibia sesqui latioribus, subovatis.

Abdominis segmentum inferius sextum maris longitudinaliter obsolete impressum, apice emarginatum, emarginatura bisinuata.

\* 5. *P. complanatus*: *Niger, pedibus piceis, thorace dense subtiliter punctato. — Long. 6 lin.*

Habitat in Siam, Dom. Kaden.

Depressus, niger, nitidus, subtiliter pubescens. Antennae articulis duobus primis rufo-piceis (reliqui desunt). Mandibulae piceae. Caput thorace paulo angustius, supra parce disperseque minus fortiter punctatum, puncto quoque pilo brevi erecto instructo, interstitiis omnium subtilissime punctulatis. Thorax apice coleopterorum latitudine, basin versus angustatus, latitudine apicis haud brevior, lateribus leviter rotundatus, basi truncatus, apice utrinque subsinuatus, angulis anterioribus acutiusculis, posterioribus leviter rotundatis, subdepressus, dense subtiliter punctatus, subtiliter tenuiterque griseo-pubescent. Scutellum punctulatum. Elytra thorace paulo longiora, dense punctata, pube longiore grisea tenuiter vestita. Abdomen crebre punctatum, griseo-pubescent, apice summo picescente. Pedes picei, tarsis fulvis, anticis tibia sesqui latioribus, oblongis.

Mas latet.

\* 6. *P. niger*: *Niger, nitidus, tarsis rufo-piceis, thorace basin versus angustato, vage punctato, basi obsolete bi-impresso. — Long. 5½ lin.*

*Araecerus niger* Nordm. *Symb.* 157. 1.

Habitat in Brasiliae meridionalis prov. Montevideo, Sellow.

Subdepressus, niger, nitidus, capite thoraceque glabris. Antennae capite thoraceque paulo breviores, tenues, articulo tertio secundo paulo longiore, nigrae, apice piceae. Palpi nigri. Mandibulae piceae. Caput thoracis latitudine, latitudine brevius, subtilissime parce punctulatum, utrinque parce vageque fortiter punctatum. Thorax coleopteris paulo angustior, latitudine antica haud brevior, basin versus leviter angustatus, lateribus rectis, basi apiceque truncatus, angulis anterioribus subrectis, posterioribus rotundatis, leviter transversim convexus, basi obsolete longitudinaliter bi-impressus, dorso biplagiatus crebre, lateribus parce sparsim punctatus, interstitiis subtilissime disperse punctatis. Scutellum subtiliter punctulatum. Elytra thorace paulo longiora, fortiter profunde punctata, parce subtiliterque nigro-pubescentia. Abdomen crebre punctatum, tenuiter nigro-pubescent. Pedes graciliores, nigri, tarsis dilute rufo-piceis, anticis membranis dilatatis tibia duplo prope latioribus.

\* 7. *P. Javanus*: *Niger, griseo-pubescent, pedibus rufo-testaceis, thorace subquadrato, aequaliter dense punctato. — Long. 6 lin.*

Habitat in Java, Dom. Westermann.

Statura omnino *P. latipedis*, niger, subopacus, griseo-pubescent. Antennae capite thoraceque paulo breviores, tenues, articulo tertio secundo sesqui longiore, sequentibus sensim decrescentibus, rufo-testaceae. Palpi testacei. Caput thoracis fere latitudine, postice creberrime rugoso-punctatum, fronte anterieus laevigata, nitida, parce subtiliter punctulata, ante oculos utrinque vage fortiter punctata. Thorax coleopteris paulo angustior, latitudine apicis paululum longior, basin versus subangustatus, lateribus rectis, basi apiceque truncatus, angulis anterioribus acutis, posterioribus obtusis, subrotundatis, leviter transversim convexus, aequaliter dense minus profunde punctatus, basi lineola tenuissima longitudinali laevi subelevata. Scutellum punctulatum. Elytra thorace paulo longiora, dense punctata, interstitiis transversim subrugosis. Abdomen crebre punctatum, metallico-versicolor, segmento quinto sextoque margine, sexto inferiore etiam summa basi rufo-piceis. Pedes rufo-testacei, coxis piceis, tarsis anticis membranis valde dilatatis tibia plus duplo latoribus.

- \* 8. *P. Aegyptius*: *Nigro-piceus, dense griseo-pubescent, ano pedibusque rufis, thorace basin versus angustato, aequaliter dense subtiliterque punctato. — Long. 6 lin.*

Habitat in Aegypto, Dom. Dr. Waltl.

Statura et summa affinitas praecedentis, at paulo magis depressus, nigro-piceus, parum nitidus, pube depressa fusco-grisea densius vestitus. Antennae capite thoraceque breviores, tenues, articulo tertio secundo sesqui longiore, sequentibus sensim decrescentibus, testaceae. Palpi testacei. Caput subtriangulare, thorace paulo angustius, supra postice utrinque late creberrime punctatum, ceterum laevigatum, nitidum, omnium subtilissime parce punctulatum, fronte inter oculos utrinque vage fortius punctata. Thorax apice coleopteris paulo lator, latitudine apice vix longior, basin versus distincte angustatus, subdepressus, aequaliter dense subtiliter punctatus. Scutellum punctulatum. Elytra thorace paulo longiora, subtilius crebre punctata. Abdomen crebrius punctatum, supra segmento singulo medio puncto maiore utrinque unico notato, subtus pluribus sparso, segmento quinto margine apicali, sexto toto rufo-piceis. Pectus piceum. Pedes rufi, coxis omnibus nigris, tarsis anticis membranis fortiter dilatatis tibia plus duplo latoribus.

- \* 9. *P. Capensis*: *Niger, nigro-pubescent, pedibus rufis, abdomine aeneo-nitidulo, thorace subquadrato, aequaliter dense punctato. — Long. 6 lin.*

Habitat in Africa australi, Dom. Drège.

Statura et summa affinitas praecedentium, niger, opacus, nigro-pubescent, abdomine aeneo-versicolore. Antennae capite thoraceque paulo breviores, tenues, articulo tertio secundo paulo brevior, sequentibus decrescentibus, rufo-testaceae. Palpi rufo-testacei. Caput thorace vix angustius, supra postice utrinque triangulariter creberrime fortiter intricato-punctatum, anterieus laevigatum, nitidum, omnium subtilissime parce punctulatum, ante oculos utrinque vage fortiter punctatum. Thorax apice coleopterorum latitudine, latitudine haud longior, basin versus vix angustatus, lateribus rectis, basi apiceque truncatus, angulis anterioribus acutiusculis, posterioribus rotundatis, supra leviter transversim convexus, aequaliter confertissime punctatus, dense nigro-pubescent. Scutellum punctulatum. Elytra thorace



tertia parte longiora, confertissime punctata, dense nigro-pubescentia. Abdomen crebre punctatum, tenuius pubescens, fortiter aeneo-relucens, subtus segmentis ultimis duobus punctis maioribus sparsis, apice piceis. Pedes rufi, coxis piceis, tarsis anticis membranis fortiter dilatatis tibia plus duplo latioribus.

\* 10. *P. latipes*: Niger, griseo-pubescent, pedibus flavis, abdomine fortius punctato, thorace subquadrato, crebrius vage punctato. — Long.  $6\frac{1}{2}$  lin.

Grav. Micr. 202. 1. — Latr. Hist. nat. Crust. et Ins. IX. p. 336. — Laporte Etud. Ent. I. 122. 1. — Nordm. Symb. 154. 1.

*Lathrobium latipes* Grav. Mon. 129. 1.

Habitat in America septentrionali.

Niger, parum nitidus, griseo-pubescent. Antennae capite thoraceque paulo breviores, tenues, articulo tertio secundo sesqui prope longiore, sequentibus decrecentibus, flavo-testaceae, articulis omnibus inde a tertio apice fusciscentibus. Os dilute rufo-piceum, palpis pallidis. Caput thoracis fere latitudine, dense subtilissime punctulatum, vage subtiliter punctatum, postice et in media fronte grossius et parum profunde punctatum, ante oculos utrinque impressum. Thorax coleopteris paulo angustior, latitudine parum longior, basin versus leviter angustatus, lateribus subrectis, basi apiceque truncatus, angulis anterioribus subrectis, posterioribus obtusis, subrotundatis, leviter convexus, minus dense fortiter punctatus, linea tenui longitudinali media calloque utrinque antico laevioribus, undique omnium subtilissime punctulatus. Scutellum punctulatum. Elytra thorace tertia parte longiora, creberrime fortiter profundeque punctata. Abdomen crebrius fortiusque punctatum, segmentis omnibus margine rufo-piceo. Pedes flavi, coxis piceo testaceis, tarsis anticis membranis fortiter dilatatis tibia plus duplo latioribus, suborbiculatis.

Abdominis segmentum inferius quintum maris apice leviter emarginatum, sextum integrum, praeter basin laevissimum.

\* 11. *P. flavipes*: Niger, griseo-pubescent, pedibus flavis, abdomine subtilius punctato, thorace basin versus angustato, vage punctato. — Long.  $5\frac{1}{2}$  lin.

Habitat in Puerto-Rico, Dom. Moritz.

Proximus praecedenti, at minor, praesertim angustior, niger, parum nitidus, griseo-pubescent. Antennae capite thoraceque paulo breviores, tenues, articulo tertio secundo paulo longiore, sequentibus decrecentibus, flavo-testaceae, articulis singulis a tertio inde apice fuscis. Os rufo-testaceum, palpis flavis. Caput thorace parum angustius, confertissime subtilissimeque punctulatum, postice utrinque crebrius fortiter punctatum, antice utrinque punctis nonnullis sparsis fortibus impressum, margine antico rufo-testaceo. Thorax apice coleopteris vix angustior, latitudine antica paulo longior, basin versus modice angustatus, lateribus subrectis, basi apiceque truncatus, angulis omnibus leviter rotundatis, supra parum convexus, subtilissimeque punctulatus, minus fortiter vage punctatus, linea angusta longitudinali subelevata calloque utrinque antico laevibus. Scutellum parce punctulatum. Elytra thorace tertia parte longiora, crebre profunde punctata, opaca. Abdomen crebre subtilius punctatum, subtus, praecipue versus api-

cem, punctis maioribus sparsum, apice summo et infra segmentorum omnium margine summo apicali rufo-piceis. Pedes flavi, coxis nigro-piceis, tarsis anticis membranis fortiter dilatatis tibia plus duplo latioribus, suborbiculatis.

Abdominis segmentum inferius sextum maris integrum, praeter basin laevigatum.

12. *P. albipes*: *Niger, griseo-pubescent, pedibus flavis, abdomine subtiliter punctulato, thorace basin versus angustato, crebrius punctato.* — Long.  $5\frac{1}{2}$  lin.

Habitat in Mexico, Mus. Dom. Chevrolat.

Statura et summa affinitas praecedentis *Niger*, opacus, griseo-pubescent. Antennae capite thoraceque paulo breviores, tenues, articulo tertio secundo sesqui longiore, sequentibus sensim decreascentibus, testaceae, articulis singulis a quarto inde fusciscentibus. Palpi flavi. Mandibulae rufo-piceae. Caput thorace vix angustius, supra densius subtiliter punctulatum, posterius utrinque obsolete punctatum, fronte anterieus nitidiore, infra oculos utrinque punctis nonnullis maiusculis sparsa, ad antennarum basin utrinque leviter impressa, margine apicali angusto saturate rufo-testaceo. Thorax apice coleopterorum latitudine, latitudine antica paulo longior, basin versus leviter angustatus, lateribus subrectis, basi apiceque truncatus, angulis anterioribus et posterioribus subrotundatis, parum convexus, crebrius minus subtiliter punctatus, basi, linea angusta media longitudinali et callo utrinque pone apicem sublaevibus, ante basin et pone apicem obsolete bi-impressus. Elytra thorace paulo longiora, dense fortiter profundeque punctata, dorso nigro-, lateribus et margine apicali griseo-pubescentia. Abdomen densius subtiliter punctatum, densius griseo-pubescent, segmentis ultimis margine tenui apicali pallide testaceo. Pedes pallide flavi, coxis nigris, tarsis anticis membranis fortiter dilatatis tibia plus duplo latioribus, suborbiculatis.

Abdominis segmentum inferius sextum maris apice leviter emarginatum.

Obs. Differt a praecedente thorace paulo longiore, crebrius punctato, fronte distinctius punctulata, elytris fortius, abdomine subtilius punctatis, abdominis segmento inferiore sexto maris apice emarginato.

13. *P. sulphuripes*: *Niger, griseo-pubescent, pedibus flavis, thorace basin versus subangustato, crebre punctato, bilineatim impresso.* — Long.  $5\frac{2}{3}$  lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Statura omnino et summa similitudo *P. flavipedis*. *Niger*, opacus, griseo-pubescent. Antennarum articuli 2 primi (reliqui desunt) palpique flavi. Mandibulae piceae. Caput thoracis prope latitudine, supra parcius sparsim subtiliter punctatum, fronte antice utrinque punctis nonnullis fortibus impressa, margine antico summo piceo. Thorax apice coleopteris latitudine aequalis, latitudine summa parum longior, basin versus leviter angustatus, lateribus subrectis, basi apiceque truncatus, angulis anterioribus subrectis, posterioribus obtusis, subrotundatis, parum convexus, sat crebre fortius punctatus, interstitiis, margine postico, callisque duobus minutis prope apicem laevibus, dorso per totam longitudinem lineis duabus parallelis leniter impressis, postice latioribus bi-canaliculatus. Scutellum punctulatum. Elytra thorace paulo

longiora, dense profundeque punctata, apice pube densiore longioreque ciliata. Abdomen crebre minus fortiter punctatum, apice summo segmentorumque penultimorum margine apicali ferrugineo. Pedes pallide flavi, coxis nigris, tarsis anticis membranis dilatatis tibia plus duplo latioribus, suborbiculatis.

Abdominis segmentum inferius quintum maris apice subemarginatum, sextum longitudinaliter impressum, apice incisum.

Obs. Differt a praecedentibus duobus corpore paulo crassiore, thorace fortius punctato, distinctius longitudinaliter bi-impresso et abdominis segmento inferiore sexto maris impresso, apice profundius inciso.

\* 14. *P. fulvipes*: *Niger, griseo-pubescent, pedibus testaceis, thorace crebrius subtiliter punctato, basi obsolete bifoveolato. — Long. 4½ lin.*

Habitat in Columbia, Dom. Moritz.

Praecedentibus simillimus, at paulo minor, praecipue brevior, niger, opacus, griseo-pubescent. Antennae capite thoraceque breviores, tenues, articulo tertio secundo parum longiore, 3 et 4 aequalibus, sequentibus decrescentibus, testaceae, articulis a tertio inde singulis apice fuscis. Palpi testacei. Mandibulae rufo-piceae. Caput thorace parum angustius, fronte laevi, pone oculos et ante oculos utrinque et apice punctis nonnullis maioribus sparsa, ad antennarum basin utrinque leviter impressa, margine apicali saturate rufo-testaceo, longius griseo-pilosum. Thorax coleopterorum prope latitudine, latitudine vix longior, basin versus subangustatus, lateribus subrectis, angulis anterioribus subrectis, posterioribus rotundatis, parum convexus, aequaliter minus dense subtiliter punctatus, basi foveolis duabus obsoletiusculis impressus. Scutellum crebre punctatum. Elytra thorace tertia parte longiora, creberrime profundius punctata, secundum suturam leviter impressa, longius pubescentia. Abdomen crebre minus fortiter punctatum, longius pubescens, segmentis ultimis duobus margine apicali piceo-testaceo. Pedes testacei, coxis piceis, tarsis anticis membranis dilatatis tibia duplo latioribus, suborbiculatis.

Obs. *P. Cayennensis* Laport. Etud. Ent. I. 122. 1.: „Simillimus *P. latipedi*, statura autem multo minore, capite minore, thorace obsolete punctato, sublaevi, corpore minus pubescente, long. 4 lin, e Cayenna”, forte a specie modo descripta haud distinctus, hac descriptione autem nequaquam cognoscendus. *Lathrobium* maius Brull. D' Orbign. It. Am. mer. Ins. t. 5. f. 9. ad hoc genus pertinere et huic speciei similis videtur, icon autem minus exacta, ut ad quamnam tot tantopere affinium specierum adhibenda sit discernatur.

15. *P. suffusus*: *Niger, griseo-pubescent, palpis pedibusque flavis, fronte elytrisque obscure castaneis, thorace crebre subtilius punctato. — Long. 5 lin.*

Habitat in Venezuela Guyanensi, Dom. Moritz.

Habitu omnino praecedentium, *P. latipede* paulo minor, niger, opacus, griseo-pubescent, capite thoraceque subnitidis, glabris. Antennae capitis thoracisque prope longitudine, tenues, articulo tertio secundo paulo longiore, sequentibus sensim decrescentibus, flavae, articulis singulis a quarto inde apice summo fuscis. Palpi flavi. Caput thorace parum latius, supra confertissime subtilissimeque punctulatum, posterius grossius parum profunde



punctatum, fronte ante oculos utrinque punctis nonnullis sat fortibus impressa, castaneum, vertice nigricante, subtus nigrum, nitidum, ore testaceo. Thorax coleopterorum prope latitudine, latitudine haud longior, lateribus rectis, basin versus vix angustatus, basi apiceque truncatus, angulis anterioribus subrotundatis, posterioribus late rotundatis, leviter transversim convexus, confertissime subtilissimeque alutaceus, crebre minus profunde punctatus, margine postico, linea media longitudinali, calloque utrinque prope apicem parum distincto laevioribus, basi obsolete bifoveolatus. Scutellum sublaeve, opacum. Elytra thorace paulo longiora, confertissime profunde punctata, longius pubescentia, obscure castanea, opaca, basi nigricantia. Abdomen supra creberrime punctatum, subtus parcius profundiusque longius pubescens, segmentis singulis apice castaneo-marginatis. Pedes flavi, coxis nigris.

Sexus differentia latet.

\* 16. *P. palmatus*: *Niger, nitidus, glaber, pedibus flavis, geniculis nigricantibus, abdomine obsolete punctato, thorace basin versus angustato, crebre punctato.* — Long.  $5\frac{1}{2}$  lin.

Nordm. Symb. 155. 2.

Habitat in Parà interiore, Sieber.

Niger, nitidus, glaber. Antennae capite thoraceque paulo longiores, tenues, articulo tertio secundo sesqui longiore, sequentibus sensim decrescentibus, flavo-testaceae. Mandibulae piceae. Caput thoracis fere latitudine, supra postice et in media fronte transversim fortiter punctatum, apice punctorum serie transversali impressum, interstitiis laevissimis. Thorax coleopteris vix angustior, latitudine anteriore haud brevior, basin versus angustatus, lateribus anterioribus leviter, posterioribus rotundatis, basi apiceque truncatus, angulis anterioribus subrectis, posterioribus obtusis, subdepressus, crebre fortiterque punctatus, linea media longitudinali parum distincta calloque utrinque anteriore laevibus. Scutellum laeve. Elytra thorace sesqui longiora, fortiter profundeque punctata. Abdomen coleopteris angustius, parcius obsolete punctatum. Pedes graciliores, flavi, coxis nigris, geniculis piceis, tarsis anticis membranis fortiter dilatatis tibia plus duplo latioribus, subovatis.

Mas latet.

\* 17. *P. terminatus*: *Rufo-castaneus, nitidus, capite, thorace, elytrorum abdominisque apice pedibusque posterioribus nigris, his basi pallidis, thorace basin versus angustato, vage punctato.* — Long.  $5\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Antennae capite thoraceque paulo breviores, tenues, articulo tertio secundo paulo longiore, sequentibus decrescentibus, rufo-testaceae, articulis singulis a secundo inde medio fusciscentibus. Palpi rufo-testacei. Mandibulae rufo-castaneae. Caput thoracis prope latitudine, nigerrimum, glaberrimum, pernitidum, supra disperse subtiliter punctulatum, postice et ante oculos utrinque fortiter vage punctatum. Thorax coleopteris paulo angustior, latitudine haud brevior, basin versus angustatus, lateribus leviter rotundatis, basi apiceque truncatus, angulis anterioribus subrectis, posterioribus

late rotundatis, subdepressus, vage sat profunde punctatus, linea media longitudinali subelevata calloque utrinque antico laevibus, glaberrimis, niger, pernitidus. Scutellum basi punctatum, castaneum. Elytra thorace sesqui longiora, minus crebre sat profunde fortiterque punctata, rufo-castanea, apice sinuatum nigra, nitida, parce cinereo-pubescentia. Abdomen subtilius minus crebre punctatum, pube parca longiore depressa grisea vestitum, dilute rufo-castaneum, segmento quinto nigro, basi rufo-castaneo, sexto toto nigro. Pectus castaneum. Pedes antichi testacei, posteriores nigri, femoribus basi ultra medium pallide testaceis.

Mas latet.

18. *P. brunneus*: *Niger, nitidus, pubescens, thorace, elytris pectoreque brunneo-rufis, antennis, palpis pedibusque flavis, thorace basin versus angustato, disperse punctato.* — Long.  $3\frac{2}{3}$  lin.

Laporte Etud. Ent. I. 122. 2.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Praecedentibus minor. Antennae capitis thoracisque prope longitudine, tenues, articulo tertio secundo paulo longiore, sequentibus sensim decrescentibus, flavo-testaceae. Palpi flavo-testacei. Os rufo-testaceum. Caput thoracis latitudine, nigrum, nitidum, longius parce pilosum, supra leviter convexum, punctis parvis dispersis fortioribus posterius et inter oculos impressum, interstitiis laevissimis. Thorax vix coleopterorum latitudine, latitudine antica paulo longior, basin versus angustatus, lateribus leviter rotundatis, basi apiceque truncatus, angulis anterioribus subrectis, posterioribus late rotundatis, convexus, disperse fortius profundiusque punctatus, parce pilosus, brunneo-rufus, nitidus. Scutellum brunneo-rufum, laeve. Elytra thorace sesqui longiora, crebre fortiter profundeque punctata, parce pilosa, brunneo-rufa, nitidula. Abdomen parcius minus subtiliter punctatum, longius pubescens, nigrum, nitidum, segmentis singulis margine apicali tenui rufo-piceo. Pectus brunneo-rufum, coxis concoloribus, pedibus flavo-testaceis, tarsis anticis membranis fortiter dilatatis tibia plus duplo longioribus, suborbiculatis.

Mas latet.

- \* 19. *P. crassicollis*: *Niger, nitidus, antennis pedibusque flavis, elytris rufo-castaneis, rude punctatis, thorace lato, biseriatim punctato.* — Long. 4 lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Habitu fere *P. brevicollis* at sesqui maior, niger, capite thoraceque pernitidis. Antennae capite sesqui longiores, tenues, articulo primo magis elongato, secundo sequentibus paulo longiore et crassiore, 3 — 5 aequalibus, sequentibus decrescentibus, flavae. Palpi flavi. Mandibulae rufo-piceae. Caput thorace multo angustius, laevissimum, fronte punctis paucis sat magnis impressa. Thorax coleopterorum latitudine, latitudine paulo brevior, lateribus subrectis, basi apiceque truncatis, angulis anterioribus subrectis, posterioribus fortiter rotundatis, convexus, laevissimus, dorso utrinque serie minus regulari, circiter 6-punctata, lateribus utrinque punctis tribus quatuorve singulis sat profundis impressus. Scutellum parum-punctatum. Ely-

tra thorace paulo longiora, dense punctis magnis profundis impressa, parce subtiliter pubescentia, castaneo-rufa, minus nitida. Abdomen crebrius punctatum, subtus parcius longiusque pubescens, supra pube densa depressa grisea micante vestitum, segmentis ultimis duobus margine apicali pallido. Pedes flavi, coxis piceis, tarsis anticis membranis sat dilatatis tibia duplo latioribus, suborbiculatis.

Sexus differentia latet.

\* 20. *P. brevicollis*: Nigro-piceus, griseo-pubescent, pedibus rufis, thorace lato, transverso, aequaliter crebre punctato. — Long.  $3\frac{1}{4}$  lin.

Habitat in Aegypto, Dom. Prof. Ehrenberg.

Reliquis plerisque minor et praecipue brevior, nigro-piceus, parum nitidus, griseo-pubescent. Antennae capite parum longiores, minus tenues, articulo secundo tertioque subaequalibus, sequentibus subobconicis, sensim brevioribus, ultimis leviter transversis, ultimo apice abrupte acuminato, rufo-testaceae. Palpi rufo-testacei. Caput thorace angustius, crebre fortius punctatum, fronte ante medium laeviore. Thorax coleoptera latitudine fere excedens, latitudine dimidio fere brevior, basin versus haud angustatus, lateribus subrectis, basi leviter rotundatus, apice truncatus, angulis omnibus rotundatis, leviter transversim convexus, aequaliter crebreque punctatus, basi medio obsolete bi-impressus. Scutellum subtiliter punctulatum. Elytra thorace sesqui longiora, crebre punctata, margine summo apicali rufo. Abdomen breviusculum, crebre punctatum, apice segmentorumque singulorum margine apicali rufo-piceis. Pedes rufi, coxis omnibus piceis, tarsis anticis tibia sesqui latioribus, suborbiculatis.

Sexus differentia latet.

### **Taenodema Laporte.**

Lap. Etud. Ent. 1. p. 120.

*Gymnurus Nordm. Symb. p. 158. t. 2. f. 7.*

*Palpi maxillares articulo quarto securiformi.*

*Abdomen marginatum.*

*Tarsi antici articulis 4 primis dilatatis coniunctis patellam formantibus.*

Corpus mediae magnitudinis, elongatum, lineare, alatum. Caput thorace minus, subdepressum, subtriangulare, basi truncatum, collo quasi tenuiore brevi thoraci immissum, exsertum, porrectum, oculis ad angulos capitis posteriores sitis, mediocribus, rotundatis, prominulis. Labrum latum, brevissimum, integrum. Mandibulae falcatae, acutae, medio dente longiore bicuspi armatae. Maxillae (t. IV. f. 35. b.) malis coriaceis, interiore interne, exteriore apice barbatis. Palpi maxillares leviter elongati, articulo primo minimo, secundo elongato, subcylindrico, tertio hoc dimidio breviori, obconico, quarto triangulariter introrsum producto, securiformi. Labium (t. IV. f. 35. a.) mento transversum, ligula brevi, membranacea, biloba, lobis rotundatis, apice pilorum fasciculo instructis, paraglossis acuminatis, membranaceis, ligulam longe superantibus. Palpi labiales articulo primo minuto, secundo leviter elongato, tertio parvo, subacuminato. Antennae filiformes, articulo ultimo ovato. Thorax coleopterorum prope latitudine, depressus, quadran-



gulus, angulis anterioribus subrectis, posterioribus rotundatis. Scutellum triangulare. Elytra truncata. Abdomen lineare, coleopteris angustius, tere-tiusculum, marginatum, stylis analibus haud exsertis. Pedes breviores, antici incrassati, femoribus tibiisque intus sinuatis, tarsis articulis 4 primis fortiter dilatatis, transversis, subtus spongioso-tomentosis, posteriores graciliores, tibiis posticis compressis, apice oblique truncatis, tarsis articulo primo leviter elongato, sequentibus duobus coniunctis aequali, quarto parvo, lobato.

#### Observationes.

1. Corpus punctulatum, pubescens, habitu generi praecedenti assimilabile, distinctum antennis pedibusque minus gracilibus, abdomine angustiore, tarsis anticis subtus tomentosis, palpis maxillaribus articulo securiformi terminatis.

2. Sexus differentia latet.

3. De victu nihil constat. Species nonnisi ex America aequinoctiali cognitae.

\* 1. *T. cyanescens*: *Nigra, nitida, capite thoraceque laevibus, cyaneis, nitidis, elytris punctatis, abdomine apice obscure rufo.* — Long. 6 lin.

*Gymnurus cyanescens* Nordm. Symb. 159. 1. t. 2. f. 7.

Habitat in Parà Brasiliae, Sieber.

Nigra, subnitida, capite thoraceque supra cyaneis, nitidis, elytris nigro-subcyanescentibus. Antennae vix capitis thoracisque longitudine, articulo primo tertio parum maiore, hoc secundo sesqui longiore, sequentibus sensim paulo brevioribus nec crassioribus, leviter obconicis, penultimo crassitie haud longiore, ultimo ovato. Caput et thorax depressa, laevissima, glaberrima, polita, punctis raris hinc inde impressa. Caput thorace angustius, latitudine brevius. Thorax coleopterorum latitudine, latitudine paulo brevior, basin versus haud angustatus, basi leviter rotundatus, apice truncatus, angulis anterioribus levius, posterioribus fortius rotundatis. Scutellum obsolete subtiliter punctatum. Elytra thorace paulo longiora, crebrius punctata, parce nigro-pubescentia. Abdomen coleopteris dimidio angustius, punctulatum, nigro-pubescens, segmento sexto apice rufo-castaneo. Pedes breviusculi, crassiusculi. Alae fuscae.

\* 2. *T. vestita*: *Nigra, nitida, plagiatis cinereo-tomentosa, elytris cyaneis, pedibus piceis.* — Long. 6 lin.

*Gymnurus vestitus* Nordm. Symb. 160. 2.

Habitat in Parà Brasiliae, Sieber.

*T. cyanescens* gracilior, nigra, nitida. Antennae nigro-piceae, articulo primo tertio haud longiore, licet paulo crassiore, subtus testaceo, secundo tertio plus duplo brevior, 3—6 elongatis, teretibus sensim paululum brevioribus (reliqui desunt). Caput thoracis prope latitudine, latitudine postica haud brevior, fronte antice depressa et utrinque subimpressa, creberrime punctatum, cinereo-pubescens, macula frontali media subelevata laevissima, glabra. Thorax coleopteris parum angustior, latitudine tertia parte longior, basin versus leviter angustatus, depressus, basi truncatus, apice subbisinuatus, angulis anterioribus subrectis, posterioribus rotundatis, creberrime punctatus, postice carinatus, cinereo-tomentosus, margine basali, lateribus usque ad medium adscendente, parce subtiliterque punctato, calloque utrinque ad

apicem sito laevisimo glabris, lateribus parce nigro-pilosellis. Scutellum cinereo-tomentosum. Elytra thoracis longitudine, crebre punctata, nigro-cyanea, nitida, dense cinereo-tomentosa, margine basali angustiore apicalique latiore laevigatis, glaberrimis. Abdomen elongatum, coleopteris dimidio angustius, segmentis singulis basi punctulatis, cinereo-tomentosis, apice laevigatis, glabris, lateribus nigro-pilosis, segmento septimo prominulo, bicuspi. Pedes cinereo-pubescentes, picei, posteriores elongati, femoribus dilutioribus.

### Species mihi invisae.

3. *T. semicyanea*: *Ruguloso-punctata, nigra, capite, thorace elytrisque cyaneis.* — Long. 6 lin.

Laporte *Etud. Ent.* I. 121. 1.

*Paederus semicyaneus* Perty *Delect. an. art.* p. 33. t. 7. f. 8.

Habitat in Brasilia.

Caput, thorax et elytra ruguloso-punctata. Abdomen subcylindricum. Antennae capitis fere et thoracis longitudine. Pedes nigri, tarsorum articulo quarto bifido. (Perty).

Obs. Secundum iconem citatam caput thoracis longitudine, thorax latitudine multo longior, basin versus vix angustatus, elytra thoracis longitudine. Secundum Laporte abdominis segmenta apice testaceo-marginata.

4. *T. aenea*: *Nigra, capite, thorace elytrisque aeneis, abdomine segmentis duobus margine ultimoque toto rubris.*

*Staph. aeneus* Oliv. *Ent.* III. 42. 10. 6. t. 2. f. 11.

Habitat Surinami.

Antennae nigrae, longitudine thoracis. Caput, thorax elytraque aenea. Abdomen nigrum, segmento quarto quinto margine ultimoque toto rubris. Pedes nigri. (Oliv.)

Obs. Forte huius generis. Secundum iconem abdominis segmentum tertium et quartum apice rubra, quintum totum nigrum.

### **Palaminus Er.**

*Palpi maxillares articulo quarto securiformi.*

*Abdomen immarginatum.*

*Tarsi antici articulis 4 primis coniunctis patellam formantibus.*

Corpus minutum, elongatum, lineare, alatum. Caput thoracis fere latitudine, subtriangulare, basi truncatum, collo quasi angusto perbrevis thoraci immissum, exsertum, porrectum, oculis sat magnis, rotundatis, prominulis. Labrum transversum, breve, medio sinuatum, sinu utrinque denticulo terminato. Mandibulae falcatae, acutae, medio dente valido simplice acuto armatae. Maxillae (t. IV. f. 36. b.) malis membranaceis, interiore intus, exteriori apice introrsum vergente barbata. Palpi maxillares modice elongati, articulo primo brevi, minuto, secundo leviter elongato, tertio hoc paulo brevior, obconico, quarto triangulariter introrsum producto, securiformi. Labium (t. IV. f. 36. a.) mento transversum, apice profundius emarginato, ligula membranaea, sinu lato biloba, lobis interae ciliatis, paraglossis ligula paulo brevioribus. Palpi labiales articulis duobus primis subaequalibus.

tertio minuto acuminato. Antennae breviusculae, tenues, filiformes, articulis duobus primis reliquis paulo crassioribus, ultimo saepius praecedentibus maiore, obconico, truncato, apice abrupte acuminato. Thorax subcordatus, latitudine antica brevior, antice truncatus. Coleoptera apice coniunctim emarginata. Abdomen coleopteris angustius, elongatum, lineare, immarginatum, subdepressum, stylis analibus exsertis, acuminatis. Pedes parum elongati, antici subincrassati, tibiis apicem versus triangulariter sensim modice dilatatis, tarsis articulis 4 primis dilatatis, 3 primis triangularibus, transversis, subtus lobatis, quarto minuto, aegre distinguendo, quinto sat tenui, posteriores graciles, tibiis intermediis simplicibus, posticis ante apicem dilatatis, dein emarginatis, unde ante apicem externe dente triangulari armatae videntur, tarsis articulo primo modice elongato, quarto brevi, lobato.

#### O b s e r v a t i o n e s .

1. Corpus punctatum, pilosellum, habitu omnino generum praecedentium, at minutum, et abdomine immarginato ab illis, depressiusculo a sequentibus distinctum. Capitis figura et oculorum situs ut in praecedentibus duobus generibus. Abdomen segmentorum 4 anteriorum sculptura e lineis impressis transversim reticulatis imbricatim texta insigne.

2. Sexus differentia abdominis segmenti inferioris sexti structura cognoscenda.

3. De victu nihil compertum. Species omnes ex America meridionali allatae.

\* 1. *P. pilosus*: *Niger, griseo-pilosus, antennis pedibusque albidis, thorace lateribus leviter rotundato, basin versus angustato, elytris thorace plus sesqui longioribus.* — Long. 3 lin.

Habitat in Columbia, Dom. Moritz.

*P. variabilis* duplo maior, niger, subnitidus, antennis, palpis pedibusque albidis-flavescentibus. Antennae elongatae, capite thoraceque longiores, tenues, articulo ultimo reliquis aequali. Caput thoracis latitudine, latitudine baseos haud brevius, subtriangulare, fronte depressiuscula, parce obsoleteque inaequaliter punctatum, longius griseo-pilosum. Thorax coleopteris dimidio angustior, latitudine antica vix brevior, basin versus leviter angustatus, lateribus subrotundatis, basi apiceque truncatus, angulis anterioribus rectis, posterioribus rotundatis, leviter convexus, parcius inaequaliter et fere obsolete punctatus, griseo-pilosus. Elytra thorace plus sesqui longiora, obsolete punctata, pilis longis depressis griseis vestita. Abdomen segmentis 4 primis fortius dense imbricato-granulatis, quinto parce obsolete granulato, sexto laevi, pilis parvis depressis griseis vestitum.

Abdominis segmentum inferius sextum maris apice emarginatum. Femina latet.

\* 2. *P. piceus*: *Piceus, parcius albidis-pubescentibus, antennis pedibusque albidis, thorace basin versus angustato, elytris thorace sesqui longioribus.* — Long. 2 lin.

Habitat in Columbia, Dom. Moritz.

*Piceus*, nitidus, pube longiore minus depressa albida parcius vestitus, antennis, palpis pedibusque albidis. Antennae capitis thoracisque longitudine, tenues, articulo ultimo reliquis aequali. Caput thorace paulo latius, latitudine baseos paulo brevius, supra modice convexus, sat dense fortiter profundeque punctatum. Thorax coleopteris plus dimidio angustior, latitudine



baseos parum brevior, basin versus fortius angustatus, lateribus rectis, basi apiceque truncatus, angulis anterioribus acutiusculis, posterioribus obtusis, parum convexus, crebrius sat fortiter profundeque punctatus, linea media longitudinali subelevata calloque utrinque pone basin subelevato laevibus. Elytra thorace sesqui longiora, crebre fortiter profundeque punctata, margine apicali laevi, dilutiora. Abdomen segmentis 4 primis dense fortiter imbricato-granulatis, quinto basi obsolete tuberculato, apice sextoque toto laevibus.

Obs. Specimen singulum, abdominis segmento inferiore sexto integro, subrotundato, sexu ambiguum. A praecedente statura multo minore, pilis brevioribus, punctis crebrioribus, profundis, a sequente colore et antennarum articulo ultimo haud crassiore, ab utroque thorace angustiore, lateribus haud rotundato et elytris brevioribus distinctus.

\* 3. *P. variabilis*: Testaceus, parcius pubescens, thorace lateribus rotundato, basin versus angustato, elytris thorace plus sesqui longioribus. — Long. 2 lin.

T. 1. f. 8.

Habitat in Columbia et in Americae meridionalis ins. Puerto-Rico et St. Thomae, Dom. Moritz.

Testaceus, nitidus, albedo-pubescens pilosusque, antennis pedibusque pallide flavis. Antennae capite thoraceque paulo breviores, tenues, articulo ultimo abrupte maiore. Caput thorace parum latius, transversum, antrorsum angustatum, basi truncatum, supra leviter convexum, grosse punctatum, parce breviterque pilosellum. Thorax coleopteris paulo angustior, latitudine antica paulo brevior, basin versus angustatus, lateribus rotundatus, basi truncatus, apice late leviter emarginatus, angulis anterioribus acutiusculis, posterioribus obtusis, parum convexus, parcius grosse punctatus, tenuiter pubescens, lateribus parce nigro-pilosellus. Scutellum laeve. Elytra thorace duplo prope longiora, minus dense fortiter punctata, pube depressa parcius vestita, apice summo pallidiora laevi. Abdomen coleopteris angustius, subcylindricum, segmentis quatuor anterioribus confertim grosse imbricato-granulatis, ultimis duobus laevibus.

Abdominis segmentum inferius sextum maris apice subemarginatum; feminae acuminatum, interstitio inter acumen et latera membranaceo.

Variat: 1, Coleopteris macula media communi fusca notatis.

2, Coleopteris plaga lata communi suturali apice abbreviata nigra.

3, Coleopteris plaga lata suturali aliaque laterali, ante apicem confluentibus, fuscis.

4, Coleopteris cruce dorsali vittaque laterali nigris, abdomine nigro.

5, Duplo vel triplo minor (long.  $1\frac{1}{2}$  lin.), pallide flavo-testaceus, immaculatus, antennis pedibusque albidis.

\* 4 *P. testaceus*: Testaceus, subtilius griseo-pubescens, thorace lateribus rotundato, elytris thorace duplo longioribus. — Long.  $1\frac{1}{2}$  lin.

Habitat in Pennsylvania. Specimen a Dom. Zimmermann lectum dedit Dom. Schüppel.

Rufescenti-testaceus, nitidus, parcius subtiliusque griseo-pubescens, abdomine saturatiore, longius pubescente, antennis pedibusque dilutioribus. Antennae capitis thoracisque prope longitudine, apicem versus sensim paululum crassiores, ultimo reliquis paulo maiore. Caput thoracis latitudine, basi truncatum, apice rotundatum, modice convexum, crebrius fortiusque punctatum. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus rotundatus, basin versus vix magis quam apicem versus angustatus, basi truncatus, apice leviter emarginatus, angulis anterioribus subrectis, posterioribus obtusis, convexus, disco crebrius profundiusque punctatus, limbo sublaevi. Elytra thorace duplo longiora, sat crebre profundeque punctata, margine apicali pallidiore laevi. Abdomen segmentis 4 primis dense grosseque imbricato-granulatis, quinto sextoque laevibus.

Abdominis segmentum inferius sextum maris apice profunde incisum. Femina latet.

### **Oedichirus Er.**

*Palpi maxillares articulo quarto securiformi.*

*Abdomen immarginatum.*

*Tarsi antici articulis 4 primis singulis dilatatis, incrassatis.*

Corpus elongatum, teretiusculum, apterum. Caput thoracis fere latitudine, rotundatum, basi fortiter constrictum, collo quasi angustulo thoraci immissum, exsertum, porrectum, oculis mediocribus, rotundatis, leviter prominulis. Labrum brevissimum, medio sinuatum, margine subtiliter denticulato. Mandibulae falcatae, acutae, medio dente simplice armatae. Maxillae (t. IV. f. 37. b.) malis tenuiter corneis, interiore minuta interne, exteriore latiore extus rotundata, apice introrsum verso barbatis. Palpi maxillares elongati, articulo primo minuto, secundo valde elongato, apicem versus parum incrassato, tertio hoc duplo prope brevior, obconico, quarto triangulariter introrsum producto, fortiter securiformi. Labium (t. IV. f. 37. a.) mento transversum, ligula biloba, lobis divaricatis, subrotundatis, membranea, paraglossis acuminatis, membraneis, pubescentibus, ligulam haud superantibus, una cum ligulae lobis palporum basi eorumque articulo primo occultis, difficulter observandis. Palpi labiales breviores, articulis duobus primis crassiusculis, primo brevior, tertio parvo, tenuiore. Antennae filiformes, tenuiores, articulo primo parum elongato, ultimo apice abrupte acuminato. Thorax oblongo-cordatus, apice truncatus, convexus. Scutellum elongatum, triangulare. Coleoptera thorace breviora, apice coniunctim emarginata. Abdomen elongatum, apicem versus sensim subincrassatum, immarginatum, subcylindricum, segmento septimo prominulo, bicorniculato, stylis analibus nullis. Pedes leviter elongati, graciles, tibiis posticis apice compressis, subdilatatis, oblique emarginatis, tarsis anticis articulis quatuor primis in utroque sexu fortiter dilatatis, singulis subquadratis, tumidis, subtus membrana semicirculari lobatis, quinto minuto, posterioribus articulo primo elongato, quarto brevi, lobato.

### **O b s e r v a t i o n e s.**

1. Genus insigne, habitu omnino Paederi, abdomine segmento septimo spinis duabus porrectis armato singulare, capitis orbiculati thoracisque figura, abdomine apicem versus incrassato, pedibus magis elongatis, tarsis anticis articulis quatuor primis singulis fortiter dilatatis et incrassatis a praecedentibus satis discedens.

2. Mas abdominis segmento inferiore sexto apice emarginato, sinu membrana repleto distinctus.

3. De victu nihil constat.

\* 1. *O. Paederinus*: *Rufo-testaceus, nitidus, capite, elytris anoque nigris.* — Long. 3 lin.

T. 1. f. 9.

Habitat in Sicilia, Dahl.

Paedero littorali duplo minor et gracilior, apterus, nitidus, rufus, capite, elytris abdominisque segmentis ultimis duobus nigris, undique parcius longiusque pilosus. Antennae capite duplo longiores, articulo tertio secundo paulo longiore, 3 — 10 sensim paulo brevioribus, subobconicis, ultimo oblongo-ovato, apice abrupte acuminato, breviter pilosellae, testaceae. Palpi testacei. Os rufo-piceum. Caput thoracis latitudine, orbiculatum, vertice et inter oculos transversim fortiter parcius punctatum. Thorax apice coleopterorum fere latitudine, latitudine antica paulo longior, basin versus angustatus, basi apiceque truncatus, angulis anterioribus rotundatis, sat convexus, punctis magnis profundis, lateribus dispersis, dorso irregulariter biseriatis, impressus. Scutellum laeve, piceum. Elytra brevia, apice oblique truncata, fortiter profundeque punctata, parce nigro-pilosa. Abdomen cylindricum, segmento singulo punctorum profundorum seriebus tribus transversalibus cincto, griseo-pilosum, segmento septimo prominulo, corniculis duobus acuminatis, leviter divergentibus armato. Pedes testacei.

### **Procirrus Latr.**

*Latr. Regn. An. IV. p. 436. — Laporte Etud. Ent. I. p. 123.*

*Palpi maxillares articulo quarto elongato-fusiformi, acuminato.*

*Abdomen immarginatum.*

*Tarsi antici articulis 4 primis singulis dilatatis, incrassatis.*

Corpus minoris magnitudinis, valde elongatum, teres, alatum. Caput rotundatum, basi attenuatum, collo quasi tenui thoraci affixum, exsertum, porrectum, oculis mediocribus, rotundatis, leviter prominulis. Labrum transversum, breve, medio sinuatum. Mandibulae falcatae, acutae, medio dente magno bicuspi armatae. Maxillae (t. IV. f. 38. b.) malis membraneis, interiore interne, exteriori brevi apice barbatis. Palpi maxillares elongati, articulo primo minore obconico, reliquis tribus valde elongatis, longitudine aequalibus, ultimo fusiformi, apice acuminato. Labium (t. IV. f. 38. a.) mento transversum, ligula brevi, apice subsinuata, membranea, paraglossis acuminatis, membraneis, pubescentibus, ciliatis, ligulam longe superantibus. Palpi labiales breviores, articulis duobus primis cylindricis, secundo primo paulo longiore, tertio minuto, tenuiore. Antennae filiformes, tenues, articulis duobus primis reliquis paulo crassioribus, ultimo cylindrico, apice obtuse acuminato. Thorax elongatus, subcylindricus. Scutellum rotundatum. Elytra truncata. Abdomen elongatum, lineare, immarginatum, segmento sexto conico-acuminato, stylis analibus haud exsertis. Pedes elongati, antici leviter incrassati, tarsi articulis 4 primis in utroque sexu fortiter dilatatis, singulis subquadratis, tumidis, posteriores graciles, tibiis posticis apice com-



pressis, oblique emarginatis, tarsis posterioribus articulo primo valde elongato, quarto minuto, lobato.

#### O b s e r v a t i o n e s.

1. Habitus fere Paederi, at multo gracilior, a reliquis huius tribus generibus palporum maxillarium articulo ultimo acuminato valde distinctus, capitis basin versus attenuati figura, oculis ante eius medium sitis, thorace valde elongato habitu ab illis divergens, tarsorum anticorum structura cum genere praecedente congruens.

2. Mas abdominis segmento inferiore sexto apice summo emarginato distinctus.

3. Victus latet.

×

\* 1. *P. Lefebvrei*: Fusco-ferrugineus, opacus, confertim punctatus, subtiliter fulvo-pubescent. — Long. 4 lin.

Latr. Regn. An. IV. p. 436. not. — Guér. Iconogr. Ins. t. 9. f. 6. — Laporte Etud. Ent. I. p. 124.

Habitat in Sicilia.

Totus fusco-ferrugineus, abdominis apice, antennis, palpis pedibusque dilutioribus, ferrugineo-testaceis, confertim punctatus, opacus, subtiliter fulvo-pubescent. Antennae capite sesqui longiores, articulo primo modice elongato, cylindrico secundoque hoc dimidio brevior, elongato, obconico crassioribus, sequentibus tenuibus, tertio secundo paulo brevior, 3 et 4 aequalibus, 4 — 10 sensim paulo brevioribus, penultimis paulo crassioribus, leviter obconicis, ultimo praecedentibus ambobus paulo longior, cylindrico, apice obtuse acuminato. Caput thoracis latitudine, supra modice convexum, basi subcarinatum. Thorax coleopteris angustior, latitudine duplo longior, basin versus sensim leviterque angustatus, basi apiceque truncatus, angulis anterioribus fortiter rotundatis, dorso depressiusculus. Coleoptera thoraci longitudine aequalia.

## Tribus sexta.

### S t e n i n i.

*Stigmata prothoracica occulta, spatio pone coxas anticas corneo.*

*Coxae posticae conicae.*

*Antennae in fronte insertae*

---

Labrum corneum, apice rotundatum,  
integerrimum Dianoo, Steno;  
denticulatum Euaestheto.

Mandibulae longae, tenues, falcatae, acutae, ante apicem fortiter unidentatae.

Maxillae malis brevibus, interiore interne, exteriore apice membranaceis barbatisque.

Palpi maxillares elongati, graciles, articulo primo elongato, secundo tertioque singulis paulo longioribus, hoc apice magis minusve leviter incrassato, quarto minimo, vix nisi oculo fortissime armato conspiciendo.

Mentum corneum.

Ligula membranea.

Paraglossae membranae, ligulae connatae eamque paulo superantes Dianoo, Euaestheto;

liberae, ovatae Steno.

Palpi labiales breves articulo secundo incrassato, tertio subulato.

Antennae breviores, tenues, II-articulatae, articulis ultimis tribus magis minusve incrassatis clavatae subclavatae, in fronte inter oculos insertae Dianoo, Steno;

ad capitis marginem anteriorem insertae Euaestheto.

Caput maiusculum, porrectum, collo quasi crasso thoraci immissum, oculis magnis, magis minusve prominentibus.

Thorax immarginatus, saepius cylindricus, basi apiceque truncatus, subtus integumento corneo integro.

Scutellum saepius vix distinctum.

Elytra pectore haud longiora.

Abdomen segmentis sex distinctis, septimo saepius prominulo.

Pedes graciles, coxis anticis parvis, subconicis, leviter prominentibus, intermediis tenui spatio distantibus, posticis conicis, trochanteribus posticis fulcrantibus, tibiis inermibus, tarsis

5-articulatis Dianoo, Steno;

4-articulatis Euaestheto.

### O b s e r v a t i o.

Sat distincta videtur haec tribus a praecedentibus duabus antennarum insertione et structura, palporum maxillarium articulo primo elongato, coxis anticis minutis. Ceterum palporum structura magis cum Paederinis convenit.

### Dianous Leach.

*Curt. Brit. Ent. III. t. 107. — Mannerh. Bruchelytr. p. 41. — Steph. Illustr. V. p. 304. — Boisd. et Lacord. Faun. Ent. Paris. I. p. 440. — Er. Col. March. I. p. 527.*

*Mentum transversum, quadratum.*

*Tarsi 5-articulati.*

Corpus elongatum, subcylindricum, alatum. Caput thorace latius, basi constrictum, collo quasi cylindrico thoraci immissum, exsertum, oculis mediocribus, rotundatis, leviter prominulis. Labrum latum, transversum, antice truncatum, margine integerrimo. Mandibulae falcatae, acutae, ante apicem unidentatae. Maxillae (t. V. f. 1. b.) malis corneis, interiore margine interno membranaceo, recto, barbato, exteriori apice introrsum verso membranaceo, barbato. Palpi maxillares elongati, articulis tribus primis elongatis, sensim paulo longioribus, teretibus, quarto subulato, subtilissimo, vix observando. Labium (t. V. f. 1. a.) mento transversim quadrangulare, spatio membranaceo inter mentum et palporum basin brevi, transverso, ligula biloba, lobis distantibus, oblongis, membraneis, paraglossis membraneis, subtiliter ciliatis, ligulae lobis aequalibus iisque valde approximatis, fere connatis et aegre distinguendis. Palpi labiales breves, articulo primo tenui, leviter arcuato, secundo hoc paulo longiore, ovato, tertio minuto, subtili, subulato. Antennae in fronte anteriore inter oculos insertae, graciles, articulis duobus primis reliquis vix crassioribus, primo secundo brevior, tertio elongato, quarto duplo prope longiore, sequentibus 5 (4—8) longitudine sensim decrescentibus, ultimis tribus paululum crassioribus, penultimis duobus obconicis, ultimo ovato, apice acuminato. Thorax coleopteris multo angustior, cylindricus. Scutellum minutum, triangulare. Elytra apice truncata. Abdomen marginatum, apice sensim angustatum, setis duabus exsertis terminatum. Pedes graciles, modice elongati, tarsis filiformibus, articulo primo leviter elongato.

### O b s e r v a t i o n e s.

1. Corpus subtilissime pubescens, subtilius punctatum.
2. Habitus omni Steni, at oculi minus grandes prominentesque, antennae longiores, corpus subtiliter punctatum differentiam quandam habitualement praebent.
3. Mas abdominis segmento sexto ventrali apice emarginato, feminae eodem acute rotundato noscuntur.
4. Dom. Brullé hoc genus tarsorum articulo quarto bilobo distinxit a Steno (Hist. nat. Ins. VI. p. 85.), fugit autem huncce auctorem, permultas Stenorum species articulo tarsorum quarto fortiter bilobo instructas esse, Dianoum autem a reliquis Stenis tarsorum structura nullo modo differre.
5. Victus in rivorum ripis humidis.



\* 1. *D. coerulescens*: *Nigro-coeruleus, nitidulus, subtiliter albido-pubescent, elytris macula rotundata rubra.* — Long.  $2\frac{1}{2}$  lin.

Curt. Britt. Ent. III. t. 107. — Mannerh. Brachelytr. 41. 1. — Boisd. et Lacord. Faun. Ent. Paris. I. 440. — Er. Col. March. I. 528. 1.

*Stenus coerulescens* Gyll. Ins. Suec. II. 463. 1. — Sahlb. Ins. Fenn. I. 421. 1.

*Stenus biguttatus* Ljungh Web. et Mohr Arch. I. 62. 5.

Habitat in Europa.

Corpus nigro-coeruleum, nitidulum, pube brevissima albida vestitum. Antennae capitis thoracisque prope longitudine, nigrae, apice obscure ferrugineae. Palpi nigri. Mandibulae rufae. Caput thorace sesqui latius, transversim suborbiculatum, basi profunde constrictum, dense subtilius punctatum, fronte subdepressa, leviter bisulcata, interstitio lato, leviter convexo, aequae ac frontis latera elevato. Thorax coleopteris dimidio angustior, lateribus ante medium modice rotundatus, posterius magis angustatus, basi coleopteris duplo angustior, latitudine summa longitudini aequali, subcylindricus, subtilius punctatus, punctis in medio dorso rarioribus. Elytra thorace sesqui fere longiora, sutura longitudinaliter depressa et basi impressa, dense subtilius punctata, pone medium macula rotunda rubra vel fulva, colore violaceo cincta, notata. Abdomen subtilissimo punctulatum. Pedes tibiis tarsisque nigris, densius albido-pubescentibus.

### **Stenus Latr.**

Latr. Précis des caract. p. 77. Gen. Crust. et Ins. I. p. 294. — Fab. Syst. El. II. p. 602. — Grav. Micr. p. 153. — Curt. Britt. Ent. IV. t. 164.

*Mentum triangulare.*

*Tarsi 5-articulati.*

Corpus elongatum, subcylindricum, aliis alatum, aliis apterum. Caput thorace paulo latius, basi leviter constrictum, collo quasi crasso thoraci immisum, oculis permagnis, rotundatis, prominulis. Labrum sat magnum, transversim semicirculare, margine integerrimo. Mandibulae falcatae, acutae, ante apicem fortiter acuteque unidentatae (seu apice acute bidentatae). Maxillae (t. V. f. 2. b.) malis corneis, interiore margine interno membranaceo, subrotundato, barbato, exteriori apice introrsum verso membranaceo, barbato. Palpi maxillares elongati, articulis primis elongatis, sensim longioribus, duobus primis terebibus, tertio apicem versus leviter incrassato, subclavato, quarto subulato, subtilissimo, oculo etiam fortiter armato aegre observando. Labium (t. V. f. 2. a.) mento triangulari, ligula biloba, lobis minimis, ovatis, approximatis, membraneis, paraglossis distantibus, basi coarctatis, apice rotundato-ampliat, ligula multo maioribus, membraneis. Palpi labiales breves, articulo primo tenui, leviter arcuato, secundo hoc brevior, subovato, tertio minuto, subtili, subulato. Antennae in fronte anteriore insertae, tenuiusculae, articulis duobus primis crassioribus, sequentibus magis minusve elongatis, dein sensim brevioribus, ultimis tribus reliquis distincte crassioribus, clavam oblongam formantibus. Thorax coleopteris angustior, magis minusve cylindricus. Scutellum vix conspicuum. Coleoptera coniuncta apice emarginata. Abdomen lineare, aliis marginatum, aliis immarginatum. Pedes magis minusve elongati gracilesque.

## O b s e r v a t i o n e s.

1. Habitus huius generis peculiaris. Corpus omnibus magis minusve rude punctatum, colore nigro, rarius ad cupreum, saepius ad plumbeum vergente, pilis brevibus albidis, saepius subargenteis rarioribus densioribusve cinerascens, ore plerumque densius pubescente.

2. Mas plerumque abdominis segmento sexto inferiore apice emarginato, saepe etiam segmentis ventralibus pluribus emarginatis impressisve distinctus.

3. Singulariter in hoc genere ligula levissime mento affixa est, quo fit, ut agone mortis vi antiperistaltica oesophagus cum ligula, paraglossis palpisque labialibus prolabatur et promuscidis porrectae speciem prae se ferat. Hanc rem primus observasse videtur Olivier, bene eam recensuit Gyllenhal (Ins. Suec. II. p. 476). Dein Dom. Thion peculiarem de Stenorum instrumentis cibariis commentationem (Annal. Soc. Ent. Gall. IV. p. 153.) edidit, qua huius partis structuram exacte descripsit, quam partem autem pro vera promuscide habuit. Ipse specimibus recentibus dissectis non possum quin Gyllenhalianae opinioni accedam.

4. Victus palustris vel riparius. Species nonnullae gregariae in arena, riparia humida agilissimi cursitant.

## I. Tarsi articulo quarto simplice.

## A. Abdomen marginatum.

## \* Elytra maculata.

\* 1. *St. biguttatus*: Nigro-subaeneus, dense profundeque punctatus, subtiliter albedo-pubescent, palpis basi testaceis, elytris macula fulva. — Long.  $2\frac{1}{3}$  lin.

*Fab. Syst. El. II.* 602. 1. — *Grav. Micr.* 154. 2. *Mon.* 225. 2. — *Latr. Hist. nat. Crust. et Ins. IX.* 352. 2. *t.* 80. *f.* 1. *Gen. Crust. et Ins. I.* 294. 1. — *Gyll. Ins. Suec. II.* 464. 2. — *Sahlb. Ins. Fenn. I.* 421. 2. — *Zetterst. Faun. Lappon. I.* 86. 1. *Ins. Lappon.* 69. 1. — *Guér. Iconogr. Regn. An. Ins. t.* 9. *f.* 8. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 443. 3. — *Er. Col. March. I.* 529. 1.

*Staph. biguttatus* Linn. *Faun. Suec. n.* 851. *Syst. Nat. I. II.* 685. 15. — *Fab. Gen. Ins.* 241. 11—12. *Spec. Ins. I.* 336. 13. *Mant. Ins. I.* 221. 20. *Ent. Syst. I. II.* 527. 36. — *Müll. Prodr.* 90. 1124. — *Schrunk Enum. Ins. Aust.* 235. 446. *Fauna Boica I.* 643. 871. — *Goeze Ent. Beitr. I.* 719. 15. — *De Vill. Ent. I.* 415. 15. — *Panz. Faun. Germ. II.* 17. *Ent. Germ.* 355. 26. — *Rossi Faun. Etr. ed. Helv. I.* 312. 2. — *Walk. Faun. Paris. I.* 276. 2.

*Paed. biguttatus* Oliv. *Ent. III.* 44. 5. 4. *t.* 1. *f.* 3. *a. b.*

*Sten. bipustulatus* Ljungh *Web. u. Mohr Archiv. I.* 1. 63. 4. — *Mannerh. Brachelytr.* 41. 1. — *Runde Brachelytr. Hal.* 14. 1.

*Staph. bipustulatus* Marsh. *Ent. Britt.* 527. 82.

Habitat in Europa.

Nigro-subaeneus, nitidulus, pube brevi depressiuscula albida cinerascens. Antennae tenniusculae, medio saepe piceae, articulo tertio quarto duplo longiore. Palpi nigri, articulo primo toto et secundo basi dilute testaceis. Mandibulae nigrae, apice piceae. Caput cum oculis thorace sesqui latius, crebre punctatum, fronte profunde excavata, carinula minuta antice abbreviata longitudinali inter oculos leviter elevata. Thorax basi coleopteris duplo angustior, latitudine baseos duplo prope longior, subcylindricus, ante medium leviter rotundato-ampliat, confertissime punctatus, pone medium canalicula dorsali utrinque abbreviata sat profunde impressus. Elytra thorace paulo longiora, ad medium baseos torulosa, confertissime rugoso-punctata, macula rotundata paululum infra medium fulva, annulo nigro cincta. Abdomen coleopteris paulo angustius, apicem versus sensim

leviter attenuatum, subtiliter punctulatum, pilis brevibus depressis albidis argenteo-subsericans, praecipue versus latera. Pedes elongati, graciles, nigri, trochanteribus testaceis, tarsis elongatis, apice picescentibus.

Mas abdominis segmento inferiore sexto apice profunde emarginato, quinto apice medio leviter sinuato, intra sinum oblique retuso, excavatione laevi, margine utrinque prominulo obtuso terminata, quarto apice eodem modo at obsolete impresso; femina segmento sexto leviter triangulariter producto, apice subrotundato.

\* 2. *St. bipunctatus: Nigro-subaenens, dense profundeque punctatus, densius albido-pubescent, palpis articulo primo testaceo, elytris macula fulva.* — Long.  $2\frac{1}{2}$  lin.

Er. Col. March. I. 530. 2.

Habitat in Europa, in Mesopotamia, Dr. Helfer.

Praecedenti proximus, at distinctus, minus gracilis, nigro-subaeneus, nitidulus, pube brevi depressiuscula albida cinerascens. Antennae nigrae, articulo tertio quarto duplo longiore. Palpi nigri, articulo primo solo obscure testaceo. Mandibulae rufo-piceae, basi nigrae. Caput thorace paulo latius, crebre punctatum, fronte modice excavata, excavatione medio longitudinaliter subelevata, carinula tenui longitudinali, antice sensim evanescente notata. Thorax basi coleopteris duplo angustior, latitudine baseos duplo prope longior, lateribus ante medium leviter rotundatus, subcylindricus, confertissime punctatus, interstitiis subrugosis, pone medium canalicula utrinque abbreviata profunde impressus. Elytra thorace paulo longiora, ad baseos medium torulosa, confertissime rugoso-punctata, paulo infra medium macula rotundata fulva, annulo nigro cincta. Abdomen apicem versus sensim angustatum, crebre subtiliter punctatum, pilis brevibus depressis albidis, versus latera condensatis, argenteo-subsericans. Pedes nigri, trochanteribus nigro-piceis vel concoloribus, tarsis elongatis, apice picescentibus.

Mas abdominis segmento inferiore sexto longitudinaliter impresso, apice profunde emarginato, quinto inde ab apice ad medium usque leviter late impresso, impressione laevi, lateribus margine utrinque leviter prominulo obtuso terminata, apice leviter sinuato, quarto integro; femina segmento sexto apice triangulariter producto.

Obs. 1. Differt a praecedente statura paulo maiore; praecipue robustiore, antennis pedibusque minus elongatis, fronte minus profunde excavata, elytrorum macula paulo magis a sutura remota, aequali spatio a sutura et a margine exteriori distante, trochanteribus obscuris. Facilius distinguitur palporum articulo primo solo testaceo, secundo toto nigro.

2. Specimen in Lusitania a Com. Hoffmannsegg lectum, elytrorum macula dilatata insigne, in collectione Hoffmannseggiana nom. *St. bimaculati* distinctum fuit.

\* 3. *St. guttula: Niger, dense profundeque punctatus, densius albido-pubescent, pedibus testaceis, femoribus apice late nigris, elytris macula fulva.* — Long. 2 lin.

Müll. Germ. Mag. IV. 225. 23. — Er. Col. March. I. 531. 3.

Sten. Kirbyi Gyll. Ins. Suec. IV. 499. 2—3. — Curt. Britt. Ent. IV. t. 164.

Sten. biguttatus var. Grav. Mon. p. 226.

Sten. biguttatus var. b Gyll. Ins. Suec. II. 464. 2.



Habitat in Europa.

Habitu omnino praecedentis, at duplo minor, niger, nitidulus, pube brevi depressiuscula albida obsitus. Antennae piceae, articulo primo nigro, tertio quarto sesqui longiore. Palpi pallide flavo-testacei, articulo tertio apice fusciscente. Mandibulae rufo-piceae, basi nigrae. Caput cum oculis thorace sesqui latius, creberrime sat fortiter punctatum, fronte profunde excavata, carinula minuta inter oculos parum elevata. Thorax basi coleopteris duplo angustior, latitudine baseos duplo longior, lateribus ante medium leviter rotundato-ampliatus, subcylindricus, creberrime sat fortiter punctatus, subinaequalis. Elytra thorace perparum longiora, dorso leviter convexa, confertissime sat fortiter profundeque punctata, macula paulo infra medium rotundata sat magna fulvo-rufa, annulo nigriore cincta. Abdomen coleopteris paulo angustius, apicem versus sensim leviterque attenuatum, sat crebre minus subtiliter punctulatum. Pedes testacei, coxis nigris, femoribus apice late nigricantibus, tarsis elongatis, picescentibus.

Mas abdominis segmentis inferioribus ultimis duobus apice medio subsinuatis, quinto medio utrinque longitudinaliter longius ciliato; femina segmento sexto apice modice producto, rotundato.

Variat palpis articulo secundo testaceo, apice nigricante, tertio nigricante, basi testaceo, tibiis nigro-piceis, medio dilutioribus.

\* 4. *St. bimaculatus*: Niger, confertim profunde punctatus, parcius albedo-pubescent, palpis pedibusque testaceis, illis apice fuscis, his geniculis nigris, elytris macula fulva. — Long.  $2\frac{2}{3}$  lin.

Gyll. Ins. Suec. II. 466. 3. — Sahlb. Ins. Fenn. I. 422. 3. — Zetterst. Faun. Lappon. I. 87. 2. Ins. Lappon. 69. 2. — Runde Brachelytr. Hal. 14. 2. — Er. Col. March. I. 532. 4.

Sten. Juno Grav. Micr. 154. 1. Mon. 225. 1. — Latr. Hist. nat. Crust. et Ins. IX. 352. 1. — Boisd. et Lacord. Faun. Ent. Paris, I. 441. 1.

Staph. biguttatus Marsh. Ent. Britt. 526. 81.

Habitat in Europa.

St. Junone sesqui maior, robustior, niger, opacus, pube brevissima depressiuscula albida subcinerascens. Antennae tenues, piceae, articulo primo nigro, tertio quarto sesqui longiore. Palpi dilute testacei, articulo tertio apice fusco. Maxillae rufo-testaceae, basi nigrae. Caput cum oculis thorace paulo latius, confertissime profunde punctatum, fronte haud excavata, at inter oculos utrinque longitudinaliter sulcato-impressa, interstitio medio sat elevato, antrorsum angustato. Thorax basi coleopteris dimidio angustior, lateribus praecipue ante medium rotundatus, latitudine summa vix longior, parum convexus, confertissime fortiter profundeque punctatus, subinaequalis, medio obsolete canaliculatus. Elytra thorace parum longiora, subdepressa, profunde fortiterque dense punctata, interstitiis subrugosis, macula minuta rotundata testacea paululum infra medium. Abdomen sat crebre at minus fortiter profundeque punctatum, subnitidulum, apicem versus leviter angustatum. Pedes testacei, coxis nigris, femoribus summo apice nigricantibus, tarsis elongatis, picescentibus.

Mas abdominis segmento inferiore sexto apice profunde emarginato, medio laevigato, basi denticulo compresso prominulo armato, quinto medio

leviter impresso, impressione laevigata, utrinque carinula compressa, sat prominula, postice dentes instar producta terminata, apice inter carinas emarginato, medio denticulo prominulo minuto armato, quarto apice leviter impresso, impressione laevi, medio itidem denticulo minuto prominulo instructo; femina segmento sexto modice producto, apice summo emarginato.

- \* 5. *St. stigmula*: *Niger, opacus, confertim punctatus, densius albido-pubescens, palpis pedibusque testaceis, his geniculis fuscescentibus, elytris macula fulva.* — Long. 2 lin.

*Stenus Kirbyi* Boisd. et Lacord. Faun. Ent. Paris. I. 442. 2.

Habitat in Austria, Dom. Ullrich, in Gallia, Dom. Aubé.

Habitu omnino *St. speculatoris*, at paulo minor, niger, opacus, pube brevissima depressa densiore albida cinereus. Antennae piceae, articulo primo nigro, tertio quarto sesqui longiore. Palpi pallide testacei, articulo tertio apice vix fuscescente. Mandibulae rufae, basi nigrae. Caput cum oculis thorace paulo latius, dense fortius punctatum, fronte modice impressa, inter oculos longitudinaliter bisulcata, interstitio leviter elevato, subcarinato. Thorax basi coleopteris duplo fere angustior, lateribus praecipue ante medium leviter rotundatus, latitudine summa paulo longior, subcylindricus, confertissime punctato-subrugosus, medio obsolete canaliculatus. Elytra thorace vix longiora, subdepressa, confertissime fortius at minus profunde punctato-subrugosa, macula rotundata minuta rufo-testacea paulo infra medium. Abdomen apice attenuatum, confertim subtilius punctatum. Pedes rufo-testacei, coxis piceis, femoribus apice summo tibiisque basi fuscescentibus.

Mas abdominis segmento inferiore quarto quintoque late profundeque impressis, sexto apice profunde triangulariter exciso; femina sexto apice leviter producto, rotundato.

- \* 6. *St. colon*: *Niger, opacus, confertim profunde punctatus, parce albido-pubescens, antennarum articulo primo, palpis pedibusque testaceis, elytris macula minuta fulva.* — Long. 2 lin.

*Say Transact. Amer. Philoph. Soc. Philadelph. Nov. Ser. IV. 458. 2.*

Habitat in Carolina meridionali, Dom. Prof. Germar.

Statura omnino praecedentis, niger, opacus, pube brevissima parca albida cinerascens. Antennae piceae, articulo primo testaceo, tertio quarto sesqui longiore. Palpi pallide flavo-testacei. Caput thorace sesqui latius, confertissime minus subtiliter punctatum, fronte inter oculos late at parum profunde excavata, late leviter longitudinaliter bisulcata, interstitio parum elevato. Thorax basi coleopteris duplo angustior, latitudine baseos duplo prope longior, lateribus ante medium rotundato-amplius, subcylindricus, confertissime fortius punctatus, interstitiis subrugosis, pone medium canalicula, dorso utrinque abbreviata, profundiore impressus. Elytra thorace vix longiora, subdepressa, fortiter profundeque dense punctata, interstitiis subrugulosis, macula minuta rufo-testacea paulo infra medium notata. Abdomen apicem versus sensim subangustatum, sat crebre minus subtiliter punctatum, nitidulum, dorso segmentis singulis basi subtiliter longitudinaliter carinatis. Pedes testacei, coxis concoloribus, femoribus apicem versus obscurioribus, tarsis elongatis.

Mas abdominis segmento inferiore quinto sextoque apice profundius emarginatis, hoc obsolete longitudinaliter impresso, illo apice intra sinum oblique retuso; femina segmento sexto apice leviter producto et rotundato.

**\*\* Elytra immaculata. Pedes concolores.**

\* 7. *St. Juno*: Niger, parum nitidus, parcius albido-pubescent, palpis testaceis, apice fuscis, thorace obsolete canaliculato, abdomine dense fortius punctato. — Long.  $2\frac{1}{2}$  lin.

*Fab. Syst. El. II.* 602. 2. — *Ljungh Web. et Mohr Arch. I.* 1. 60. 1. — *Gyll. Ins. Suec. II.* 467. 4. — *Sahlb. Ins. Fenn. I.* 423. 4. — *Zetterst. Faun. Lappon. I.* 87. 3. *Ins. Lappon. 69.* 3. — *Mannerh. Brachelytr. 41.* 3. — *Runde Brachel. Hal. 15.* 3.

*Staph. Juno Walken. Faun. Paris. I.* 276. 1.

*Staph. clavicornis Fab. Gen. Ins. 242.* 11 — 12. *Spec. Ins. I.* 336. 14. *Mant. Ins. I.* 221. 21. *Ent. Syst. I. II.* 527. 37. — *Rossi Faun. Etr. Mant. 98.* 220. *Ed. Hellv. I.* 443. 220.

*Sten. buphthalmus Latr. Hist. nat. Crust. et Ins. IX.* 353. 6. t. 80. f. 2. *Gen. Crust. et Ins. I.* 295. t. 9. f. 2.

*Sten. boops Grav. Mon. 226.* 4. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 447. 10.

Habitat in Europa et in America septentrionali.

Niger, opacus, abdomine nitidulo, pube brevissima rariore albida parum cinerascens. Antennae breviusculae, nigrae, articulo tertio quarto vix sesqui longiore. Palpi flavi, articulo secundo apice fusco, tertio fusco, basi summa testaceo. Mandibulae nigrae, apice piceae. Caput cum oculis thorace paulo latius, sat fortiter dense punctatum, fronte modice impressa, inter oculos longitudinaliter profunde bisulcata, interstitio leviter elevato. Thorax basi coleopteris duplo fere angustior; lateribus antè medium modice rotundatus, latitudine summa paulo longior, subcylindricus, sat fortiter profundeque dense punctatus, dorso canalicula obsoleta, neque apicem, neque basin attingente, impressus, basi depressiusculus. Elytra thorace paulo longiora, dorso a medio baseos longitudinaliter subtorosa, confertissime fortiter profundeque punctata, interstitiis praecipue medio intricato-rugosis. Abdomen coleopteris paulo angustius, dense minus subtiliter punctatum, dorso segmentis singulis summa basi longitudinaliter carinatis. Pedes nigri.

Mas pectore impresso, cinereo-villoso, abdominis segmento inferiore sexto inciso, quinto medio late profundeque excavato, excavatione laevi, utrinque margine acute carinato terminata, antice carinula abbreviata sat prominula armato, apice medio profunde emarginato, reliquis quatuor medio leviter impressis, subtiliter acute carinatis, femoribus posterioribus leviter incrassatis et incurvis, tibiis posterioribus ante apicem subsinuatis, apice unco introrsum vergente terminatis; femina abdominis segmento inferiore sexto leviter triangulariter producto, apice summo subemarginato, pedibus simplicibus.

Variat femoribus basi piceo-testaceis.

\* 8. *St. intricatus*: Ater, parum nitidus, confertim punctatus, palpis articulo primo flavo, thorace canaliculato elytrisque inaequalibus, his intricato-rugosis, abdomine dense subtilius punctato. — Long.  $2\frac{1}{3}$  lin.

Habitat in Hispania meridionali, Dom. Schüppel, in Sardinia, Dom. Prof. Gené.



Habitu omnino praecedentis et minoribus eius individuís aequalis, ater, fere opacus, pube brevissima subtilissimaque albida vix cinereo-versicolor. Antennae graciliores, articulo tertio quarto sesqui longiore. Palpi nigri, articulo primo toto secundoque summa basi flavo-testaceis. Caput thorace paulo latius, creberrime profundius punctato-rugulosum, fronte modice excavata, bisulcata, sulcis latoribus minus profundis, interstitio convexo, leviter elevato. Thorax basi coleopteris triplo prope angustior, lateribus ante medium leviter rotundatus, latitudine summa tertia parte longior, subcylindricus, inaequalis, dorso obsoletius canaliculato, basin versus depressiusculo, lateribus longitudinaliter subimpressis, creberrime fortiter profundeque punctatus, interstitiis inaequalibus et praecipue lateribus et pone medium intricato-rugulosis. Elytra thorace paulo longiora, medio longitudinaliter torosa, in medio disco et secundum suturam subimpressa, dense fortius profunde punctata, interstitiis inaequalibus et praecipue posterius fortiter intricato-rugosis. Abdomen coleopteris angustius, apicem versus sensim angustatum, crebre minus fortiter punctatum, segmentis dorsalibus singulis basi impressis et medio longitudinaliter carinulatis. Pedes nigri, tarsi elongatis.

Mas abdominis segmento inferiore sexto apice triangulariter exciso, basi bidentulato, quinto longitudinaliter excavato, excavatione laevigata, utrinque carina acuta bidentata terminata, apice inter carinas emarginato, quarto longitudinaliter impresso, impressione angustiore, parum profunda, laevigata, carina acuta subtiliore simplice terminata; femina segmento sexto leviter producto, apice subemarginato.

9. *St. asphaltinus*: Niger, nitidus, dense fortiter punctatus, parce subtiliter albido-pubescent, palpi's basi testaceis, thorace obsolete canaliculato, elytris intricato-rugosis, abdomine punctato. — Long.  $2\frac{1}{2}$  lin.

Habitat in Tyrol, Mus. Dr. Aubé.

Statura gracilior omnino *St. atri*, niger, nitidulus, pube minus brevi at rara vix cinereo-micans. Antennae mediocres, articulo tertio quarto paulo longiore. Palpi nigri, articulo primo toto et secundo basi ad medium usque dilute flavo-testaceis. Caput thorace sesqui latius, profunde fortiterque minus dense punctatum, fronte leviter impressa, late bisulcata, interstitio modice convexo elevatoque. Thorax basi coleopteris duplo angustior, lateribus ante medium fortiter rotundatus ibique coleopteris vix dimidio angustior, latitudine summa paulo longior, subcylindricus, medio canalicula brevi impressus, densius fortiter profundeque punctatus, interstitiis passim rugulosis. Elytra thorace parum longiora, subaequalia, crebre fortiter profundeque punctata, interstitiis intricato-rugosis. Abdomen basi coleopteris vix angustius, apicem versus sensim angustatum, parce minus subtiliter punctatum, nitidum. Pedes graciliores, nigri, tarsi modice elongatis.

Mas latet. Femina abdominis segmento inferiore sexto apice leviter producto, rotundato.

Obs. A praecedentibus elytris brevioribus, a *St. atro* praeterea punctura multo fortiore, abdominis parca, a sequentibus reliquis tarsis elongatis distinctus.

- \* 10. *St. ater*: *Niger, confertim profunde punctatus, densius albido-pubes-  
cens, palpis articulo primo flavo, thorace elongato, subtiliter canalicu-  
lato, abdomine dense fortius punctato.* — Long.  $2\frac{3}{4}$  —  $2\frac{1}{2}$  lin.

*Mannerh. Brachelytr.* 42. 4. — *Boisd. et Lacord. Faun. Ent. Paris.* 1. 447. 11. —  
*Er. Col. March.* 1. 534. 6.

*Sten. maurus* *Mannerh. Brachelytr.* 41. 2. — *Runde Brachelytr. Hal.* 15. 4.

Habitat in Germania.

*St. Junoni* affinis, at saepe dimidio minor, gracilior, statura fere *St. biguttati*, ater, pube brevi depressiuscula densa alba cinereus. Antennae nigrae, articulo tertio quarto sesqui longiore. Palpi nigri, articulo primo toto et secundo summa basi pallide flavis. Mandibulae piceae, apice rufescentes. Caput cum oculis thorace vix sesqui latius, dense profundeque punctatum, fronte profundius excavata, inter oculos obsoletus bisulcata, interstitio parum elevato. Thorax basi coleopteris duplo angustior, lateribus ante medium leviter rotundatis, latitudine summa sesqui fere longior, subcylindricus, dense profundeque punctatus, interstitiis subrugulosis, dorso distinctius canaliculatus, canalicula neque basin neque apicem attingente. Elytra thorace paulo longiora, antice versus suturam leviter longitudinaliter torosa, confertim profunde fortiterque punctata, subrugulosa. Abdomen coleopteris angustius, apicem versus sensim leviterque attenuatum, sat dense et minus subtiliter punctatum, dorso segmentis singulis summa basi carinatis. Pedes nigri, graciles, tarsis posticis elongatis.

Mas abdominis segmento inferiore sexto apice profunde et acute emarginato, longitudinaliter laevigato, basi subtiliter bidenticulato, quinto late longitudinaliter excavato, excavatione laevigata, utrinque carina elevata, postice denticuli instar prominula, apice inter carinas emarginato, quarto tertioque pariter longitudinaliter impressis, impressione antrorsum sensim laeviore, illo apice sinuato, impressione carina prominula, licet minus acuta terminata, carinis segmenti quarti quintique pilis introrsum vergentibus ciliatis, tibiis posticis ante apicem interne subdentatis; femina abdominis segmento inferiore sexto apice leviter producto, rotundato, tibiis posticis simplicibus.

- \* 11. *St. carbonarius*: *Niger, opacus, confertim profunde punctatus, spissius albido-pubescent, palpis concoloribus, thorace obsolete canaliculato, elytris thoracis longitudine, abdomine parcius fortiusque punctato.* — Long. 2 lin.

*Gyll. Ins. Succ. IV.* 505. 13 — 14. — *Suhlb. Ins. Fenn.* 1. 432. 19. — *Mannerh. Brachelytr.* 44. 21. — *Er. Col. March.* 1. 535. 7.

Habitat in Germania.

Praecedente minor, praesertim brevior, *St. buphalmo* valde affinis, at robustior, niger, opacus, pube brevi depressa crassiuscula densiore albida argenteo-versicolor. Antennae breviusculae, articulo tertio quarto sesqui longiore. Palpi nigri, articulo etiam primo concolore. Caput thorace paulo latius, coleopteris dimidio fere angustius, fortiter profundeque dense punctatum, fronte subimpressa, late bisulcata, interstitio leniter convexo, parum elevato. Thorax basi coleopteris duplo angustior, anterior lateribus modice rotundatus, latitudine baseos duplo longior, subcylindricus,

profunde creberrime punctatus, subrugosus, subinaequalis, medio obsolete canaliculatus, canalicula utrinque abbreviata. Elytra thoracis longitudine, dense profundeque punctata, subrugosa, antarius versus suturam longitudinaliter torulosa. Abdomen parcius minus subtiliter punctatum. Pedes nigri, tarsis posticis leviter elongatis.

Mas abdominis segmento inferiore sexto quintoque longitudinaliter impressis, apice emarginatis, quarto leviter longitudinaliter impresso, tertio apice obsolete impresso; femina segmento sexto apice rotundato.

- \* 12. *St. labilis: Plumbeo-niger, subnitidus, confertim punctatus, densius albido-pubescent, palpis articulo primo piceo, fronte profunde bisulcata, thorace elongato, subtiliter obsolete canaliculato, abdomine densius minus subtiliter punctato. — Long. 2 lin.*

*Sten. canaliculatus Sahlb. Ins. Fenn. I. 428. 12.*

Habitat in Fennia, Dom. Prof. Sahlberg.

Statura omnino *St. bupthalmi*, at gracilior, plumbeo-niger, nitidulus, pube densiore albida incanus. Antennae breviusculae, articulo tertio quarto parum longiore. Palpi nigri, articulo primo piceo. Caput thorace paulo latius, coleopteris plus dimidio angustius, dense profundeque punctatum, fronte haud excavata, late profundeque longitudinaliter bisulcata, interstitio convexo, aequae ac frontis latera elevato. Thorax elongatus, coleopteris duplo angustior, lateribus ante medium levissime rotundatus, cylindricus, dense profundeque punctatus, interstitiis passim subrugulosis, dorso subtiliter obsoletius per totam longitudinem canaliculatus. Elytra thorace parum longiora, aequalia, densius profunde punctata, interstitiis planis. Abdomen coleopteris paulo angustius, basin versus sensim angustatum, crebrius minus fortiter punctatum, nitidum. Pedes nigri, tarsis breviusculis.

Mas abdominis segmento inferiore sexto apice leviter emarginato, quinto quartoque medio longitudinaliter late depressis; femina latet.

Obs. Fronte haud impressa at fortiter bisulcata, thorace elongato, lateribus vix rotundato, canaliculato ab affinibus distinctus.

- \* 13. *St. ruralis: Niger, opacus, confertissime punctatus, subtilius densiusque albido-pubescent, palpis concoloribus, fronte late leviter bisulcata, thorace oblongo, obsolete subtiliter canaliculato, elytris thorace longioribus, abdomine parcius subtiliusque punctato. — Long. 1½ — 2 lin.*

Habitat in Austria, Dom. Ullrich.

Statura et summa affinitas *St. bupthalmi*. Niger, parum nitidus, pube brevi depressa albida cinerascens. Antennae breviusculae, articulo tertio quarto paulo longiore, nigrae. Palpi nigri, articulo primo concolore. Caput thorace paulo latius, dense fortius profunde punctatum, fronte depressiuscula, late at parum profunde bisulcata, interstitio leviter convexo, parum elevato. Thorax basi coleopteris duplo angustior, lateribus ante medium leviter rotundatis, latitudine summa tertia parte longior, subcylindricus, dense profundeque punctatus, interstitiis leviter rugulosis, subinaequalis, obsolete canaliculatus, dorso pone medium depressiusculo. Elytra thorace paulo longiora, basin versus medio torulosa, dense profundeque punctata,



interstitiis subrugosis. Abdomen apice angustatum, parcius subtiliusque punctatum. Pedes nigri, tarsis breviusculis.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem apice leviter producto, rotundato.

Obs Proximus et simillimus *St. bupthalmio*, distinctus autem palpis basi concoloribus, elytris longioribus et abdomine parcius subtiliusque punctato. A praecedente thorace minus elongato et fronte minus profunde bisulcata, nec non abdominis in mare structura differt.

- \* 14. *St. stygicus*: Plumbeo-niger, opacus, dense profundeque punctatus, densius albido-pubescent, palpis concoloribus, fronte profunde bisulcata, thorace elongato, utrinque subimpresso, abdomine densius fortiusque punctato. — Long.  $1\frac{1}{2}$  lin.

Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 458. 3.

Habitat in America septentrionali, Dom. Zimmermann.

Statura omnino *St. bupthalmi* at paulo gracilior, plumbeo-niger, pube breviori subdepressa albida densius aequaliter canus. Antennae breviusculae, nigrae, articulo tertio quarto paulo longiore. Palpi nigri, articulo primo apice piceo. Caput thorace paulo latius, coleopteris dimidio angustius, dense sat profunde punctatum, fronte leviter excavata, late profundius bisulcata, interstitio convexo et modice elevato. Thorax coleopteris duplo angustior, anterieus lateribus parum rotundatus, latitudine summa sesqui fere longior, subcylindricus, dense sat fortiter punctatus, interstitiis subrugulosis, medio utrinque leviter impressus. Elytra thorace vix longiora, sat fortiter confertim punctata, anterieus versus suturam leviter torulosa. Abdomen crebre minus subtiliter punctatum. Pedes nigri, tarsis breviusculis.

Mas latet. Femina abdominis segmento inferiore sexto apice rotundato.

Obs. Differt a *St. bupthalmio* fronte angustiore, profundius bisulcata, thorace elongato, lateribus parum rotundato, haud canaliculato, palpis basi concoloribus.

- \* 15. *St. egeus*: Nigro-subaeneus, parum nitidus, subtilissime albido-pubescent, palpis concoloribus, thorace oblongo, dorso obsolete longitudinaliter bi-impresso, elytris thorace longioribus, abdomine dense fortius punctato. — Long.  $1\frac{1}{2}$  lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Habitu omnino *St. bupthalmi* at dimidio minor, nigro-subaeneus, obscure nitidulus, pube brevi subtilissima albida parce vestitus et vix cinerascens-micans. Antennae breviusculae, articulo tertio quarto vix longiore, nigrae. Palpi nigri, basi concolores. Caput thorace vix latius, coleopteris duplo prope angustius, dense profundeque punctatum, fronte haud impressa, late obsolete bisulcata, interstitio vix convexo, haud elevato. Thorax ante medium modice rotundatus et coleopteris dimidio, basi iis duplo angustior, latitudine summa paulo longior, minus fortiter at profunde dense punctatus, interstitiis subrugulosis, subinaequalibus, dorso obsolete secundum totam longitudinem bi-impressus, impressionibus basi apiceque distinctioribus. Elytra thorace paulo longiora, dorso medio basin versus torulosa, confertim minus fortiter profunde punctata, interstitiis subtiliter rugulosis. Abdomen crebre minus subtiliter punctatum. Pedes nigri, tarsis breviusculis.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso, quinto quartoque leviter longitudinaliter impressis, impressione utrinque carina obsoleta terminata; femina segmento sexto apice leviter producto, rotundato.

- \* 16. *St. buphthalmus*: *Niger, opacus, confertissime profunde punctatus, subtilius parciusque albido-pubescent, palpis articulo primo testaceo, thorace oblongo, obsolete canaliculato, elytris thoracis longitudine, abdomine densius fortiusque punctato.* — Long.  $1\frac{3}{4}$  — 2 lin.

*Grav. Micr.* 156. 6. *Mon.* 230. 9. — *Gyll. Ins. Suec. II.* 475. 10. — *Sahlb. Ins. Fenn. I.* 428. 11. — *Mannerh. Brachelytr.* 43. 11. — *Runde Brachelytr. Hal.* 16. 9. — *Er. Col. March. I.* 536. 8.

*Sten. boops Ljungh Web. Beitr. II.* 158. 12.

*Sten. clavicornis Panz. Faun. Germ. 27. 11. Ent. Germ.* 356. 27.

*Sten. canaliculatus Boisd. et Lacord. Faun. Ent. Paris. I.* 449. 14.

*Sten. palposus Zetterst. Ins. Lappon. 70. 6. (forte)*

Habitat in Europa.

*Niger*, pube brevi subtili subdepressa rariore albida subcinerascens. Antennae breviusculae, nigrae, articulo tertio quarto paulo longiore. Palpi nigri, articulo primo testaceo. Caput thorace paulo latius, coleopteris parum angustius, creberrime sat profunde punctatum, fronte vix impressa, late bisulcata, interstitio leviter convexo, subelevato. Thorax basi coleopteris duplo angustior, anterieus lateribus leviter rotundatus, latitudine summa parum longior, subcylindricus, dense profunde punctatus, interstitiis subrugulosis, medio plerumque obsolete canaliculatus, dorso pone medium depressiusculus. Elytra thorace vix longiora, confertissime fortius sat profunde punctata, interstitiis subrugulosis, anterieus versus basin longitudinaliter torulosa. Abdomen apicem versus sensim angustatum, densius minus subtiliter punctatum, parum nitidum. Pedes nigri, breviusculi, tarsi breviusculi.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso; femina eodem apice rotundato.

Obs. *St. palposus* Zett. „palpis elongatis, articulo ultimo magno inflexo” forte specimen monstrosum vel fortuito deformatum.

- \* 17. *St. colonus*: *Niger, nitidulus, confertissime profunde punctatus, subtiliter parcius albido-pubescent, palpis articulo primo testaceo, thorace oblongo, elytris thorace vix longioribus, abdomine parce subtiliterque punctato.* — Long.  $1\frac{3}{4}$  lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Statura et summa affinitas praecedentis at minoribus eius individuis paululum minor; niger, pube brevi depressa subtili albida parca subcinerascens. Antennae breviores, articulo tertio quarto paulo longiore, nigropiceae. Palpi nigri, articulo primo testaceo. Caput thorace paulo latius, coleopteris vix dimidio angustius, sat fortiter profundeque dense punctatum, fronte haud impressa, late parum profunde bisulcata, interstitio aequae ac frontis latera elevato, subcarinato. Thorax basi coleopteris duplo prope angustior, ante medium lateribus parum rotundatus, latitudine summa paulo longior, subcylindricus, creberrime profunde punctatus, interstitiis subrugulo-

sis, dorso pone medium depressiusculus. Elytra thorace parum longiora, sat fortiter profunde punctata, interstitiis transversim rugulosis, antierius longitudinaliter torulosa. Abdomen apicem versus sensim subangustatum, parce subtiliusque punctatum, nitidum. Pedes nigri, tarsis breviusculis.

Mas abdominis segmento inferiore quinto apice obsolete emarginato, sexto leviter triangulariter exciso; femina eodem rotundato.

Obs. A praecedente thorace haud canaliculato et imprimis abdomine parce punctato, a *St. cinerascens* colore magis nigro, punctura thoracis et elytrorum fortiore, interstitiis punctorum rugulosis distinctus.

\* 18. *St. morio*: *Plumbeo-niger, nitidulus, albido-pubescent, densius punctatus, capite coleopteris dimidio angustiore, fronte leviter excavata, obsolete bisulcata, thorace oblongo, aequali, elytris thorace longioribus, abdomine densius subtiliusque punctato.* — Long.  $1\frac{1}{2}$  lin.

Grav. Mon. 230. 10. — Er. Col. March. I. 537. 9.

*Sten. buphthalmus* Zetterst. Faun. Lappon. 88. 5. Ins. Lappon. 70. 5.

Habitat in Germania.

*Plumbeo-niger, nitidulus, pube densiore brevi depressa albida cinerascens. Antennae mediocres, nigrae, articulo tertio quarto longiore. Palpi nigri, articulo primo testaceo. Caput thorace sesqui latius, coleopteris dimidio angustius, crebre profunde punctatum, fronte leviter excavata, excavatione subaequali, sulcis longitudinalibus obsoletissimis, interstitio vix elevato. Thorax basi coleopteris duplo angustior, antierius lateribus leviter rotundatus, latitudine summa haud longior, subcylindricus, aequalis, crebre profundeque punctatus, interstitiis planis. Elytra thorace sesqui fere longiora, leviter convexa, dense profundeque punctata, interstitiis aequalibus. Abdomen coleopteris paulo angustius, apicem versus sensim subangustatum, crebrius subtiliusque punctatum, nitidum. Pedes corpori concolores, tarsis breviusculis.*

Mas abdominis segmento inferiore quinto apice leviter emarginato, sexto leviter triangulariter exciso; femina sexto apice leviter producto, rotundato.

\* 19. *St. incanus*: *Plumbeo-niger, nitidulus, densius albido-pubescent, subtilius punctatus, capite coleopteris dimidio angustiore, fronte profunde bisulcata, thorace oblongo, aequali, elytris thorace longioribus, abdomine parcius subtiliter punctato.* — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 538. 10.

Habitat in Germania.

Statura omnino et summa affinitas praecedentis. *Plumbeo-niger, nitidulus, pube densiore albida cinereus. Antennae breviusculae, articulo tertio quarto paulo longiore, nigrae. Palpi nigri, articulo primo piceo. Caput thorace sesqui prope latius, coleopteris dimidio angustius, minus dense punctatum, fronte haud impressa, late profundeque bisulcata, interstitio sulcorum convexo et aequae ac frontis latera elevato. Thorax angustior, lateribus ante medium leviter rotundatus, posterius angustatus, basi coleopteris plus duplo angustior, latitudine summa paulo longior, aequalis, densius subtiliusque punctatus, interstitiis planis. Elytra thorace paulo longiora, dorso lon-*



gitudinaliter torosa, fortius profundiusque minus dense punctata, interstitiis planis. Abdomen parcius subtiliter punctatum. Pedes nigri, tarsis breviusculis.

Mas abdominis segmento inferiore sexto apice leviter emarginato. Femina latet.

Obs. A praecedente fronte haud impressa, abdomine parce subtiliterque punctato, a sequente punctura subtiliore, ab utroque fronte profunde bisulcata distinctus.

\* 20. *St. cinerascens*: Plumbeo-niger, nitidulus, albido-pubescent, densius punctatus, capite coleopteris plus dimidio angustiore, fronte late leviter bisulcata, interstitio convexiusculo, thorace oblongo, postice depressiusculo, elytris thorace longioribus, abdomine parcius subtiliter punctato. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 539. 11.

Habitat in Germania.

Statura et summa affinitas *St. morionis*. Plumbeo-niger, nitidulus, pube densiore brevi depressa albida cinerascens. Antennae breviusculae, articulo tertio quarto vix longiore, nigrae. Palpi nigri, articulo primo piceo-testaceo. Caput thorace paulo latius, coleopteris dimidio prope angustius, dense fortius profundiusque punctatum, fronte haud excavata, leviter late bisulcata, interstitio latiusculo, leviter convexo, aequae ac frontis latera elevato. Thorax lateribus ante medium leviter rotundatus, posterius angustatus, basi coleopteris duplo angustior, latitudine summa paulo longior, dense profundeque punctatus, interstitiis angustis, planis, dorso pone medium depressiusculo vel late subimpresso. Elytra thorace paulo longiora, dorso antico medio levissime toroso, densius profunde punctata, interstitiis planis. Abdomen parce subtiliter punctatum. Pedes nigri, tarsis breviusculis.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem apice rotundato.

Obs. Facillime cum *St. morione* confundendus, at distinctus videtur, capite paulo angustiore, fronte haud excavata, sulcis longitudinalibus distinctioribus, interstitio distincte elevato, abdomine parce subtiliterque punctato, antennarum articulo tertio quarto subaequali, tarsorum posticorum articulo secundo sequentibus haud longiore, nec non abdominis segmento inferiore quinto in mare apice haud emarginato.

\* 21. *St. atratulus*: Plumbeo-niger, nitidulus, albido-pubescent, fortiter profunde punctatus, capite coleopteris plus dimidio angustiore, fronte leviter bisulcata, interstitio subcarinato, thorace oblongo, aequali, elytris thorace longioribus, abdomine densius punctato. — Long.  $1\frac{1}{4}$ — $1\frac{1}{2}$  lin.

Er. Col. March. I. 540. 12.

Habitat in Germania.

Affinis praecedentibus, at saepius paulo minor, plumbeo-niger, pube brevi depressa albida cinerascens. Antennae breves, articulo tertio quarto haud longiore, nigrae. Palpi nigri, articulo primo testaceo. Caput thorace paulo latius, coleopteris plus dimidio angustius, fortiter profundeque densius punctatum, interstitiis aequalibus, fronte haud excavata, at distincte late bisulcata, interstitio sulcorum leviter elevato, convexo, subcarinato. Thorax lateribus ante medium modice rotundatus, posterius angustatus, basi coleo-

pteris duplo prope angustior, latitudine summa haud longior, aequalis, fortiter profundeque densius punctatus, interstitiis planis, pone medium nunquam obsolete canaliculatus. Elytra thorace tertia parte longiora, dorso anterieus medio toruloso, fortiter profundeque densius punctata, interstitiis planis. Abdomen coleopteris paulo angustius, apicem versus sensim angustatum, fortius crebriusque punctatum. Pedes nigri, tarsis breviusculis.

Mas abdominis segmento inferiore quinto sextoque apice leviter emarginatis; femina sexto apice leviter producto et rotundato.

Obs. A praecedentibus punctura fortiore, elytris paulo longioribus, antennis pedibusque brevioribus, a St. incrassato statura minore, fronte distinctius sulcata et abdomine parcius punctato distinctus. Corporis habitu et statura, punctura et frontis figura eximie cum St. nigritulo congruit, at abdomen distincte marginatum.

- \* 22. *St. mendicus: Plumbeo-niger, nitidulus, griseo-pubescent, densius subtiliusque punctatus, capite coleopteris dimidio angustiore, parce punctato, fronte profunde bisulcata, elytris thorace paulo longioribus, abdomine parcius punctato.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Lusitania, Com. Hoffmannsegg.

Corpus breviusculum, crassiusculum, statura fere St. morionis, plumbeo-nigrum, nitidulum, pube subtili grisea tenuiter vestitum. Antennae breves, articulo tertio quarto paulo longiore, piceae, articulo primo nigro. Palpi nigro-picei, articulo primo testaceo. Caput thorace paulo latius, coleopteris dimidio angustius, parcius minus subtiliter punctatum, fronte depressa vel late subimpressa, sulcis minus latis at distinctioribus, profundis, interstitio haud convexo at aequae fere ac frontis latera elevato. Thorax lateribus ante medium modice rotundatus, coleopteris dimidio angustior, postice angustatus, basi coleopteris duplo angustior, latitudine summa haud longior, subcylindricus, densius subtiliusque licet sat profunde punctatus, interstitiis planis. Elytra thorace paulo longiora, dorso basin versus parum toroso, conferim minus fortiter profunde punctata, interstitiis aequalibus. Abdomen crassiusculum, minus conferim subtiliter punctatum. Pedes nigri, tarsis breviusculis.

Mas latet. Femina abdominis segmento inferiore sexto apice rotundato.

Obs. Capitis sculptura praecipue ab affinibus distinctus.

- \* 23. *St. incrassatus: Plumbeo-niger, parcius albido-pubescent, confertissime profunde punctatus, capite coleopteris duplo angustiore, fronte obsolete bisulcata, thorace oblongo, aequali, elytris thorace longioribus, abdomine nitidulo, parce subtiliterque punctato.* — Long.  $1\frac{2}{3}$  lin.

Er. Col. March. I. 541. 13.

Habitat in Germania, Suecia.

St. cinerascens affinis, plumbeo-niger, opacus, pube brevi depressa albida subcinerascens. Antennae breves, nigrae, articulo tertio quarto paulo longiore. Palpi nigri, articulo primo testaceo. Caput thorace parum latius, coleopteris duplo angustius, dense fortiter profundeque punctatum, fronte haud impressa, late obsoleteque bisulcata, interstitio parum convexo. Thorax basi coleopteris plus duplo angustior, anterieus lateribus leviter rotun-

datum, latitudine summa paulo longior, subcylindricus, fortiter profundeque confertissime punctatus, interstitiis subrugulosis, dorso ante basin late levissime subimpressus. Elytra lata, thorace paulo longiora, dense fortius sat profunde punctata, interstitiis praecipue posterius subrugulosis, anterius versus suturam leviter torosa. Abdomen maris apicem versus sensim leviterque angustatum, crassiusculum, feminae crassum, late marginatum, parce subtiliterque punctatum, nitidulum. Pedes corpori concolores, tarsis brevibus.

Mas abdominis segmento inferiore longitudinaliter subimpresso, apice leviter emarginato, sexto apice leviter triangulariter exciso; femina eodem apice leviter producto, rotundato.

Obs. Praecedentibus magnitudine et colore similis, differt autem capite minore, coleopteris abdomineque crasso latioribus, elytris rugulosis, etc. Feminas tres prope Berolinum legi, maris exemplum e Suecia nomine St buphthalmi mecum communicavit Dom. Prof. Zetterstedt.

- \* 24. *St. foraminosus*: Plumbeo-niger, nitidulus, dense fortiter profundeque punctatus, subtiliter parceque albido-pubescent, capite coleopteris dimidio angustiore, fronte obsolete bisulcata, elytris thorace sesqui longioribus, abdomine fortius punctato. — Long.  $1\frac{1}{2}$  lin.

Habitat in Germania.

Statura omnino praecedentium, plumbeo-niger, nitidulus, pube brevi depressiuscula albida parca vix cinerascens. Antennae breves, articulo tertio quarto paulo longiore, nigrae, pubescentes. Palpi toti nigri. Caput thorace paulo latius, coleopteris dimidio angustius, crebrius fortiter profundeque punctatum, fronte haud impressa, leviter bisulcata, interstitio lato, vix convexo. Thorax coleopteris paulo angustior, anterius lateribus sat fortiter rotundatus, postice angustatus, basi coleopteris duplo prope angustior, latitudine summa haud longior, leviter convexus, aequalis, fortiter profundeque crebre punctatus, interstitiis subrugulosis. Elytra thorace sesqui longiora, fortiter profunde dense punctata, interstitiis subrugulosis, anterius versus suturam leviter torulosa. Abdomen crebre fortius profunde punctatum. Pedes corpori concolores, tarsis mediocribus.

Mas latet. Femina abdominis segmento inferiore sexto apice leviter producto, rotundato.

Obs. A praecedentibus thorace brevior, anterius fortius rotundato, praecipue autem punctura multo fortiori distinctus.

- \* 25. *St. nitidus*: Plumbeo-niger, nitidus, subtiliter albido-pubescent, parcius profunde punctatus, capite coleopteris plus dimidio angustiore, fronte profundius bisulcata, thorace pone medium canaliculato, elytris thorace paulo longioribus, abdomine parcius subtiliusque punctato. — Long.  $1\frac{1}{2}$  lin.

Boisd, et Lacord. Faun. Ent. Paris. I. 450. 16.

Habitat in Sardinia, Dom. Prof. Gené, in Hispania meridionali, Dom. Schüppel, in Gallia, coll. Dom. Aubé.

Statura omnino St. cinerascens, plumbeo-niger, nitidus, pube depressa albida parcius vestitus. Antennae breviusculae, articulo tertio quarto sesqui longiore, nigrae. Palpi nigri, articulo primo testaceo. Caput thorace parum latius, crebrius minus fortiter profunde punctatum, fronte haud im-



pressa, profunde bisulcata, sulcis convergentibus, interstitiis aequae ac frontis latera elevato, subcarinato. Thorax ante medium fortius rotundatus, coleopteris dimidio angustior, postice angustatus, basi coleopteris duplo angustior, latitudine summa paululum longior, subcylindricus, minus confertim et minus fortiter profunde punctatus, interstitiis planis, posterius profundius canaliculatus. Elytra thorace paulo longiora, dorso longitudinaliter torulosa, parcius minus fortiter punctata, interstitiis planis. Abdomen parcius subtiliusque punctatum. Pedes nigri, tarsi breviusculi.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem rotundato.

Obs. Punctura parciore et thorace posterius profunde canaliculato imprimis distinctus.

\* 26. *St. aemulus*: Plumbeo-niger, fortius profundeque punctatus, subtiliter albido-pubescent, fronte planiuscula, thorace elongato, subtiliter canaliculato, elytris thoracis longitudine, abdomine parcius subtiliter punctato. — Long. 2 lin.

Er. Col. March. I. 541. 14.

Habitat in Germania et in Lapponia, Dom. Schüppel.

Plumbeo-niger, nitidulus, pube subtili albida tenuiter adpersus. Antennae breviusculae, articulo tertio quarto sesqui longiore, nigrae. Palpi nigri, articulo primo toto et secundo summa basi dilute testaceis. Caput thorace sesqui latius, crebre fortiter profundeque punctatum, fronte planiuscula, utrinque obsolete sulcata, linea tenui longitudinali media subelevata laevi. Thorax elongatus, ante medium leviter rotundatus, posterius angustatus, basi coleopteris duplo angustior, latitudine summa sesqui longior, fortiter profundeque densius punctatus, interstitiis planis, dorso per totam longitudinem distincte tenuiter canaliculatus. Elytra thorace vix longiora, fortiter profundeque densius punctata, interstitiis planis. Abdomen subtiliter minus profunde parcius punctatum. Pedes nigri, tarsi breviusculi.

Mas latet. Femina abdominis segmento inferiore sexto apice leviter producto, rotundato.

Obs. Sequenti fronte planiuscula et thorace distincte subtiliter canaliculato affinis, corpore autem nitido, statura magis elongata, capite thorace elytrisque fortius et minus confertim, abdomine parcius subtiliusque punctatis distinctus.

X \* 27. *St. canaliculatus*: Plumbeo-niger, dense profundius punctatus, subtiliter albido-pubescent, fronte plana, thorace oblongo, subtiliter canaliculato, elytris thorace paulo longioribus, abdomine crebrius subtiliter punctato. — Long.  $1\frac{2}{3}$  lin.

Gyll. Ins. Suec. IV. 501. 10 — 11. — Mannerh. Brachelytr. 43. 12. — Er. Col. March. I. 542. 15.

*Sten. buphthalmus* var. (canaliculatus Kn.) Grav. Mon. p. 230.

*Sten. buphthalmus* var. b Zetterst. Faun. Lappon. I. 88. 5. Ins. Lappon. 70. 5.

Habitat in Europa.

Statura omnino *St. morionis*, plumbeo-niger, parum nitidus, pube brevi, depressiuscula densiore cinerascens. Antennae perbreves, articulo tertio quarto subaequali, nigrae. Palpi nigri, articulo primo toto, secundo summa

basi testaceo. Mandibulae rufae. Caput thorace paulo latius, coleopteris plus dimidio angustius, creberrime minus fortiter at profunde punctatum, fronte plana, aequali, sulcis frontalibus omnino fere obsoletis. Thorax antèrius lateribus leviter rotundatus, posterius angustatus, basi coleopteris duplo angustior, latitudine summa paulo longior, subcylindricus, aequaliter dense minus fortiter profunde punctatus, interstitiis subrugulosis, per totam longitudinem linea tenui at distinctiore canaliculatus. Elytra thorace paulo longiora, minus fortiter at sat profunde dense punctata, interstitiis planis. Abdomen nitidulum, minus subtiliter crebre punctatum. Pedes corpori concolores, tarsis brevibus, apice piceis.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem leviter producto, rotundato.

\* 28. *St. opacus: Subdepressus, niger, opacus, dense punctatus, subtilissime albido-pubescens, palpis fuscis, basi testaceis, fronte lata, obsolete bisulcata, thorace subrotundato, abdomine densius punctato. — Long. 1½ lin.*

*Er. Col. March. I. 543. 16.*

Habitat Berolini, Dom. Schüppel.

Habitu fere *St. latifrontis*, niger, opacus, pube albida subtilissima vix cinerascens. Antennae breves, articulo tertio quarto haud longiore, piceae, articulo primo nigro. Palpi fuscii, articulo primo toto, secundo basi testaceo. Caput breviusculum, coleopteris latitudine subaequale, confertissime punctatum, fronte plana, late obsolete bisulcata, interstitio lato, levissime convexo. Thorax lateribus fortiter rotundatus, basin versus magis quam antèrius angustatus, basi coleopteris duplo prope angustior, latitudine summa haud longior, parum convexus, aequalis, subtilius licet profunde confertim punctatus, interstitiis angustis, planis. Elytra thorace vix longiora, depressa, apice coniuncta late emarginata, minus fortiter sat profunde confertim punctata, interstitiis angustis, planiusculis. Abdomen subtiliter densius punctatum, subnitidum. Pedes nigri, tarsis brevibus.

Mas latet. Femina abdominis segmento inferiore sexto apice rotundato.

\* 29. *St. pusillus: Niger, dense punctatus, parce albido-pubescens, palpis articulo primo testaceo, fronte profunde bisulcata, thorace rotundato, elytris thorace sesqui longioribus, abdomine parcius punctato. — Long. 1 — 1½ lin.*

*Er. Col. March. I. 544. 17.*

Habitat in Germania, Anglia.

Statura omnino *St. circularis*, niger, pube brevi depressa albida parciore vix cinerascens. Antennae crassiusculae, articulo tertio quarto paulo longiore, nigrae. Palpi nigri, articulo primo testaceo. Caput thorace paulo latius, coleopteris dimidio fere angustius, confertissime punctatum, fronte sulcis duobus latis profundisque antice conniventibus impressa, interstitio aequè ac frontis latera elevato, subcarinato. Thorax lateribus fortiter rotundatus, coleopteris dimidio angustior, basin versus coarctatus, basi coleopte-

ris plus duplo angustior, latitudine summa paulo brevior, dense punctatus, subinaequalis, dorso pone medium utrinque distinctius impressus. Elytra thorace sesqui fere longiora, sat fortiter profundeque dense punctata, interstitiis subrugulosis, subdepressa, prope suturam longitudinaliter subelevata. Abdomen breviusculum, apicem versus sensim angustatum, subtilius minus crebre punctatum. Pedes nigri, tarsis brevibus.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem rotundato.

\* 30. *St. exiguus*: *Niger, dense punctatus, parce albido-pubescent, palpis articulo primo testaceo, fronte profunde bisulcata, thorace rotundato, elytris thoracis longitudine, abdomine parcius punctato.* — Long. 1 —  $1\frac{1}{4}$  lin.

Habitat in Anglia, Dom. Hope.

Statura et summa affinitas praecedentis, niger, nitidulus, pube brevi depressa albida parca vix cinerascens. Antennae breves, crassiusculae, articulo tertio quarto paulo longiore. Palpi nigri, articulo primo testaceo. Caput thorace sesqui latius, coleopteris paulo angustius, confertim fortius profunde punctatum, fronte haud impressa, at sulcis duobus fortibus profundisque antice conniventibus exarata, interstitio aequae ac frontis latera elevato, subcarinato. Thorax antierius modice rotundatus, coleopteris tertia parte angustior, postice coarctatus, basi coleopteris duplo angustior, latitudine summa vix brevior, minus confertim profunde punctatus, interstitiis vix rugosis, dorso pone medium utrinque oblique obsolete impressus, ceterum aequalis. Elytra thorace vix longiora, depressa, dorso antierius obsolete torulosa, dense profundeque punctata, interstitiis subrugulosis. Abdomen apicem versus sensim angustatum, minus confertim subtilius punctatum. Pedes nigri, tarsis breviusculis.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem rotundato.

Obs. Differt a praecedente praeter elytra breviora punctura fortiore et thorace convexiore, minus inaequali.

\*\*\* Elytra immaculata. Pedes testacei.

\* 31. *St. speculator*: *Niger, opacus, creberrime punctatus, antennis testaceis, fronte excavata, bisulcata, pedibus testaceis, genibus nigris, tarsis fuscis.* — Long.  $2\frac{1}{2}$  lin.

Boisd. et Lacord. Faun. Ent. Paris. I. 445. 6. — Er. Col. March. I. 545. 18.

Staph. clavicornis Scop. Ent. Carn. 100. 303. — Goetze Ent. Beitr. I. 725. 13.

Staph. buphthalmus Schrank Beitr. zur Naturgesch. 72. 21. Enum. Ins. Austr. 235. 445. Faun. Boica I. 613. 870.

Sten. boops Gyll. Ins. Suec. II. 469. 5. — Sahlb. Ins. Fenn. I. 424 5. — Zetterst. Faun. Lappon. I. 87. 4. Ins. Lappon. 70. 4. — Mannerh. Brachelytr. 42. 5. — Runde Brachelytr. Hal. 15. 5.

Sten. boops var. 1 Grav. Mon. p. 227.

Sten. cicindeloides Ljungh Weber u. Mohr Archiv. I. 1. 62. 1.

Habitat in Europa.

Statura omnino St. Junonis, niger, opacus, pube brevissima alba in



thorace elytrisque parca, in abdomine pectoreque densiore, infra antennas densa. Antennae breviores, tenues, articulo tertio quarto paulo longiore, fusco-testaceae, clava fusca, articulis duobus primis nigris. Palpi dilute testacei, immaculati. Caput cum oculis thorace vix sesqui latius, creberrime sat fortiter profundeque punctatum, fronte leviter excavata, profundius bisulcata, interstitio leviter convexo, vix elevato. Thorax coleopteris vix dimidio angustior, lateribus modice rotundatus, basin versus paulo magis quam apicem versus angustatus, basi coleopteris duplo angustior, latitudine summa paulo longior, subinaequalis, creberrime sat fortiter profundeque punctatus, interstitiis subrugulosis, posterius obsolete canaliculatus. Elytra thoracis longitudine, creberrime fortiter profundeque punctata, interstitiis rugulosis, humeris et antierius prope suturam torulosa. Abdomen fortius dense punctatum, nitidulum, distinctius cinereo-micans, dorso segmentis 4 anterioribus singulis summa basi carinatis. Pedes testacei, coxis geniculisque anguste nigris, tarsis elongatis, fusciscentibus.

Mas pectore impresso, longius densiusque griseo-pubescente, abdominis segmento inferiore quinto medio late longitudinaliter impresso, impressione lateribus posterius carina terminata, apice inter carinas utrinque denticuli instar prominulas leviter emarginato, sexto apice triangulariter exciso, femoribus posterioribus subincrassatis, interne densius pubescentibus. Femina abdominis segmento inferiore sexto apice summo emarginato.

\* 32. *St. providus*: *Niger, creberrime punctatus, parce pubescens, palpis testaceis, apice piceis, fronte leniter excavata, bisulcata, thorace medio canaliculato, pedibus testaceis, genibus, tibiis apice tarsisque fuscis.* — Long.  $2\frac{1}{2}$  lin.

*Er. Col. March. I. 546. 19.*

Habitat in Germania.

Statura omnino et summa similitudo praecedentis. *Niger*, opacus, pube parca brevissima albida vix cinerascens. Antennae mediocres, articulo tertio quarto sesqui longiore, nigrae. Palpi testacei, articulo tertio apice piceo. Caput thorace paulo latius, creberrime fortiter profundeque punctatum, fronte leniter excavata, profundius bisulcata, interstitio leviter convexo, parum elevato. Thorax coleopteris vix dimidio angustior, lateribus ante medium rotundatis, posterius angustatus, basi coleopteris duplo angustior, latitudine summa paulo longior, leviter convexus, dense fortius profunde punctatus, interstitiis passim subrugulosis, medio canalicula brevi obsolete notatus. Elytra thoracis longitudine, subdepressa, dense fortiter profundeque punctata, interstitiis passim subrugulosis, dorso longitudinaliter, humeris leviter torosa. Abdomen minus fortiter crebre punctatum, nitidulum, distinctius cinereo-micans, dorso segmentis 4 anterioribus summa basi subtiliter carinatis. Pedes saturate testacei, coxis trochanteribusque nigris, femoribus apice, tibiis basi apiceque, tarsis totis fusco-piceis, his sat elongatis.

Mas pectore impresso, densius longiusque griseo-pubescente, abdomine infra segmentis 5 anterioribus longitudinaliter leviter impressis, impressione densius griseo-pubescente, lateribus a segmento secundo inde ciliis sensim longioribus, introrsum vergentibus, in segmento quinto carina, sensim magis prominente, apice in denticulum producta terminata, hoc apice

inter denticulos acutos late emarginato, segmento sexto profunde acuteque triangulariter exciso, femoribus posterioribus subincrassatis, posticis interne longius pubescentibus. Femina abdominis segmento inferiore sexto apice summo leviter emarginato.

Obs. Praecedenti simillimus, mas autem abdominis structura facile distinguendus, femina imprimis fronte paulo minus excavata, thorace dorso distinctius canaliculato, antennis nigris, palpis apice fuscis et pedibus obscurioribus distincta.

\* 33. *St. scrutator*: Niger, creberrime punctatus, parce pubescens, palpis basi pallide testaceis, fronte impressa, bisulcata, thorace obsolete canaliculato, pedibus piceis, femoribus basi testaceis. — Long.  $2\frac{1}{4}$  lin.

*Sten. femoralis* Er. Col. March. I. 547. 20.

Habitat in Germania.

Statura et summa affinitas praecedentium et parum minor, niger, opacus, pube brevissima depressa albida parciore vix cinerascens. Antennae tenues, articulo tertio quarto sesqui longiore, nigrae. Palpi picei, articulo primo toto, secundo basi, tertio summa basi testaceis. Caput thorace paulo latius, confertim fortius profunde punctatum, fronte medio excavata, profunde late bisulcata, interstitio convexo, modice elevato. Thorax antrorsum leviter rotundatus, coleopteris dimidio angustior, posterius angustatus, basi coleopteris plus duplo angustior, latitudine summa paulo longior, subcylindricus, creberrime fortius profunde punctatus, interstitiis subrugulosis, medio obsolete canaliculatus. Elytra thorace tertia parte longiora, parum convexa, fortiter profundeque dense punctata, interstitiis subrugulosis, anterieus versus suturam longitudinaliter leviter torosa. Abdomen supra apicem versus sensim subtilius dense punctatum, segmentis singulis basi subtiliter carinatis, minus dense subtilius punctatum, nitidum. Pedes picei, coxis nigris, femoribus ultra medium piceo-testaceis, tarsis elongatis.

Mas pectore leviter impresso, haud villosus, abdominis segmento inferiore sexto apice inciso, quinto medio late impresso, impressione laevi, utrinque carina elevata, apice prominula terminata, apice late emarginata, quinto medio leviter impresso; femina segmento inferiore sexto apice summo subemarginato.

Obs. 1. A praecedentibus praeter palpos pedesque obscuriores et abdominis in maribus structuram praecipue elytris longioribus distinctus.

2. *St. femorati* (n. 43.) causa nomen mutavi.

\* 34. *St. sylvester*: Niger, creberrime punctatus, parce pubescens, palpis testaceis, articulo tertio fusco, fronte leviter excavata bisulcataque, thorace obsolete canaliculato, pedibus piceis, femoribus testaceis. — Long. 2 lin.

Er. Col. March. I. 547. 21.

Habitat in Germania.

Habitus et summa affinitas *St. speculatoris*, at duplo fere minor, niger, opacus, pube brevissima depressa albida parca vix cinerascens. Antennae mediocres, articulo tertio quarto sesqui longiore, nigro-piceae, articulis primis duobus nigris. Palpi testacei, articulo ultimo fusco. Caput thorace paulo latius, confertissime fortius punctatum, punctis confluentibus subrugulosum, fronte leviter excavata, medio bisulcata, interstitio leviter convexo,

parum elevato. Thorax coleopteris vix dimidio angustior, medio modice rotundatus, posterius fortius angustatus, basi coleopteris plus duplo angustior, latitudine summa paulo longior, subcylindricus, confertissime fortiter profundeque punctatus, interstitiis rugulosis, dorso medio obsolete canaliculatus. Elytra thorace parum longiora, dense fortiter profundeque punctata, interstitiis subrugulosis, antierius versus suturam longitudinaliter torulosa. Abdomen coleopteris paulo angustius, apicem versus sensim subtilius crebre punctatum, supra segmentis singulis basi subtiliter carinatis. Pedes piceo-testacei, coxis nigris, femoribus apice tarsisque picescentibus.

Mas abdominis segmento inferiore sexto apice triangulariter exciso, quinto medio late impresso, impressione laevissima, utrinque carinula elevata, postice denticuli instar prominula terminata, apice late emarginata, quarto tertio secundoque apicem versus triangulariter impressis, impressionibus sensim obsoletioribus, laevibus; femina segmento sexto apice acute rotundato.

- \* 35. *St. augur*: *Nigro-subaeneus, parum nitidus, creberrime fortiterque punctatus, palpis albidis, antennis pedibusque testaceis, femoribus apice picescentibus, fronte excavata, medio subcarinata, abdomine crebre subtilius punctato.* — Long.  $2\frac{3}{4}$  lin.

Habitat in Columbia, Dom. Moritz.

St. speculatore paulo maior, nigro-subaeneus, nitidulus, pube brevi depressa albidā passim densiore praecipue in abdomine variegatus. Antennae testaceae, clava angusta, fusca, articulo tertio quarto sesqui longiore. Palpi albi. Caput thorace paulo latius, confertissime profunde punctato-rugulosum, fronte late excavata, in fundo obsolete bisulcata, interstitio angusto, parum elevato, subcarinato. Thorax lateribus rotundatus, coleopteris paulo angustior, basin versus paulo magis quam apicem versus angustatus, basi coleopteris dimidio angustior, latitudine summa haud longior, convexus, crebre fortiter punctatus, transversim rugosus, medio obsolete canaliculatus. Elytra thorace paulo longiora, dense profunde fortiterque punctata, interstitiis subrugosis, prope suturam longitudinaliter torosa. Abdomen coleopteris angustius, apicem versus sensim angustatum, dense subtilius punctatum, dorso segmentis singulis basi carinatis. Pedes testacei, coxis nigris, femoribus apice late piceis, maris leviter incrassatis.

Mas abdominis segmento inferiore sexto per totam fere longitudinem inciso; femina eodem integro, apice rotundato.

- \* 36. *St. metallicus*: *Niger, supra viridi-cupreus, profunde creberrime rugoso-punctatus, palpis albidis, antennis pedibusque testaceis, thorace obsolete canaliculato.* — Long. 2 lin.

Habitat in montibus prope Puerto Cabello Columbiae, Dom. Moritz.

Statura omnino *St. speculatoris*, at paulo minor, subtus niger, nitidus, supra obscure viridi-cupreus, nitidus, parce subtiliterque albidopubescent. Antennae graciliores, articulo tertio quarto sesqui longiore, testaceae, articulo primo flavo, clava fuscescente. Palpi albi, articulo tertio pallide flavo. Caput thorace paulo latius, profunde rugoso-punctatum, fronte late leniter excavata, in medio profundius bisulcata, interstitio angusto, haud



elevato, subcarinato, anterieus leviter tricarinata. Thorax lateribus ante medium leviter rotundatis, coleopteris plus dimidio angustior, posterius angustatus, basi coleopteris plus duplo angustior, latitudine summa paulo longior, dorso subdepressus, creberrime profunde rugoso-punctatus, obsolete canaliculatus. Elytra thorace vix longiora, profunde creberrime punctata, interstitiis rugulosis, versus suturam longitudinaliter obsolete torosa. Abdomen apicem versus sensim leviter angustatum, segmentis dorsalibus anterioribus crebre, posterioribus parce punctatis, inferioribus omnibus parce profundeque punctatis. Pedes testacei, coxis concoloribus, maris modice incrassati.

Mas abdomine infra per totam longitudinem medio planato, segmento sexto profunde inciso; femina eodem apice acuminato.

- \* 37. *St. croceipes*: *Niger, parum nitidus, subtiliter albido-pubescent, dense profundeque punctatus, palpis pallide testaceis, antennis pedibusque testaceis, fronte leniter excavata, abdomine subtiliter punctato.* — Long.  $1\frac{2}{3}$  lin.

Habitat in montibus prope Puerto Cabello Columbiae, Dom. Moritz.

Niger, subnitidus, pube subtili densiore cinerascens. Antennae graciliores, articulo tertio quarto paulo longiore, testaceae, clava fuscescente. Palpi pallide testacei. Caput thorace sesqui latius, creberrime profundius punctatum, fronte late leniter excavata, medio leviter bisulcata, interstitio angusto, leviter elevato, subcarinato, antice leviter tricarinulata. Thorax lateribus ante medium leviter rotundatis, coleopteris duplo prope angustior, posterius angustatus, basi coleopteris triplo fere angustior, latitudine summa paulo longior, subcylindricus, profunde creberrime punctatus, interstitiis transversim subrugulosis, aequalis. Elytra thorace paulo longiora, densius profunde punctata, interstitiis vix rugulosis, dorso prope suturam longitudinaliter leviter torosa. Abdomen thorace paulo angustius, apicem versus sensim angustatum, confertim subtiliter punctatum, densius pubescens, segmentis dorsalibus anterioribus basi subtiliter carinatis. Pedes in utroque sexu graciles, testacei, coxis nigris.

Mas abdominis segmento inferiore tertio apice medio late leviter impresso, quinto apice triangulariter impresso, impressione utrinque carinula apice prominula terminata, sexto profunde inciso; femina segmento sexto apice testaceo, rotundato.

38. *St. palpalis*: *Niger, fortiter punctatus, densius albido-pubescent, palpis albidis, antennis pedibusque testaceis, fronte excavata, abdomine subtiliter punctato.* — Long.  $1\frac{2}{3}$  lin.

Habitat in montibus prope Puerto Cabello Columbiae, Dom. Moritz.

Statura fere praecedentis, niger, nitidus, pube depressa cinerascens. Antennae elongatae, tenues, articulo quarto prope tertio aequali, testaceae, clava angusta, fuscescente. Palpi toti albi. Caput thorace multo latius, oculis sat magnis, prominentibus, crebre minus fortiter punctatum, fronte profunde excavata, medio bisulcata, sulcis angustis, approximatis, interstitio angusto, vix elevato, subcarinato, laeviore, antice leviter tricarinata. Thorax medio rotundatus et coleopteris dimidio angustior, basi coleopteris duplo fere angustior, latitudine summa paulo longior, convexus, crebre sat fortiter

profundeque punctatus, interstitiis transversim rugulosis. Elytra thorace parum longiora, subconvexa, crebre fortiter profundeque punctata, interstitiis vix rugosis, versus suturam longitudinaliter subelevata. Abdomen coleopteris angustius, apicem versus sensim attenuatum, confertim subtiliter punctatum, dense griseo-pubescent, segmentis dorsalibus anterioribus basi subtiliter carinatis. Pedes testacei, coxis omnibus nigro-piceis, maris modice incrassati.

Mas abdomine infra secundum longitudinem ad segmentum quintum usque late impressum, segmento sexto apice profunde inciso. Femina latet.

39. *St. agilis*: *Niger, nitidus, crebre profundeque punctatus, palpis albidis, antennis pedibusque testaceis*. — Long.  $1\frac{1}{2}$  lin.

Habitat prope La Guayra Columbae, Dom. Moritz.

Niger, nitidus, parce tenuissime pubescens. Antennae obscure testaceae, clava fuscescente. Palpi albidi, articulo tertio pallide flavescente. Caput thorace latius, crebre fortiter profundeque punctatum, fronte late impressa, fundo carina longitudinali laevi leviter elevata. Thorax basi coleopteris duplo fere angustior, lateribus ante medium rotundato-ampliatus, latitudine summa longior, subcylindricus, creberrime profunde fortiterque punctatus, aequalis. Elytra thorace parum longiora, punctis magnis profundis dense impressa, medio longitudinaliter subelevata. Abdomen coleopteris angustius, apicem versus sensim leviter angustatum, crebre minus subtiliter punctatum, tenuiter pubescens. Pedes flavo-testacei, coxis nigris.

Mas latet.

- \* 40. *St. fossulatus*: *Plumbeo-niger, nitidulus, dense punctatus, densius albido-pubescent, palpis testaceis, pedibus testaceis, tibiis apice geniculisque fuscis, coleopteris pone scutellum impressis*. — Long.  $2\frac{1}{2}$  lin.

Habitat in Saxonia, Dom. Märkel.

Statura omnino *St. lustratoris*, plumbeo-niger, subnitidus, pube minus brevi subtilique albida densiore cinerascens. Antennae mediocres, articulo tertio quarto sesqui longiore, piceae, articulis duobus primis nigerrimis. Palpi testacei, articulo tertio ante apicem picescente. Caput thorace paulo latius, dense profundius punctatum, fronte leniter excavata, medio bisulcata, sulcorum interstitio convexo, leviter elevato, subtiliter carinato. Thorax coleopteris dimidio angustior, lateribus ante medium rotundatis, posterius angustatus, basi coleopteris plus duplo angustior, latitudine summa paulo longior, confertim fortius profunde punctatus, subrugulosus, dorso medio obsolete canaliculatus, lateribus medio utrinque leviter oblique impressis. Elytra thorace vix longiora, dense fortiter profundeque punctata, interstitiis subrugulosis, humeris et dorso ante medium longitudinaliter torosis, spatio communi infra scutellum orbiculari depressiore et leviter impresso. Abdomen coleopteris paulo angustius, apicem versus sensim angustatum, dense minus subtiliter punctatum. Pedes testacei, coxis nigris, femoribus apice latiore, tibiis basi et apice, tarsorum elongatorum articulis singulis summo apice nigricantibus.

Mas abdominis segmento inferiore sexto apice triangulariter exciso, quinto medio late longitudinaliter impresso, impressione lateribus carina ob-

tusa, pilis longioribus griseis ciliata terminata, apice inter carinas sinuato, quarto apice obsoletius longitudinaliter impresso, apice vix sinuato; femina segmento quinto apice late leviter emarginato, sexto apice rotundato.

- \* 41. *St. lustrator*: *Plumbeo-niger, subnitidus, dense punctatus, subtiliter albedo pubescens, palpis basi albidis, pedibus testaceis, femoribus apice, tibiis tarsisque piceis.* — Long.  $2\frac{1}{4}$  —  $2\frac{1}{2}$  lin.

*Er. Col. March. I. 548. 22.*

Habitat in Germania, in Gallia Mus. Dom. Aubé.

Plumbeo-niger, nitidulus, pube breviuscula depressa albida parciore subcinereus. Antennae longiores, tenues, articulo tertio quarto sesqui longiore, nigrae. Palpi nigri, articulo primo toto, secundo basi testaceo-albidis. Caput cum oculis thorace sesqui latius, dense fortius punctatum, fronte late leniter excavata, medio profundius bisulcata, interstitio leviter convexo, parum elevato. Oculi perinagni. Thorax coleopteris vix dimidio angustior, lateribus paulo ante medium fortius rotundatis, posterius fortius angustatus, basi coleopteris plus duplo angustior, latitudine summa vix longior, convexus, aequalis, dense fortiter profundeque punctatus, interstitiis vix rugulosis, nonnunquam obsolete canaliculatus. Elytra thorace paulo longiora, dense fortiter profundeque punctata, interstitiis subrugulosis, leviter convexa, anterius versus suturam longitudinaliter torulosa. Abdomen minus subtiliter crebre punctatum, dorso convexum, segmentis singulis basi subtiliter carinatis. Pedes picei, coxis nigris, femoribus basi ultra medium testaceis, tarsis elongatis.

Mas abdominis segmento inferiore sexto apice triangulariter exciso, quinto utrinque carinato, carinis fortiter compressis, acutis, medio valde prominulis, spatio inter carinas vix impresso, polito, medio obsolete carinato, apice emarginato, segmentis antecedentibus tribus longitudinaliter late impressis, impressionis margine pilis longis flavescentibus dense ciliato; femina segmento sexto apice summo emarginato.

- X \* 42. *St. aterrimus*: *Subdepressus, aterrimus, crebre punctatus, antennis fusco-testaceis, basi nigris, palpis fuscis, basi testaceis, pedibus piceo-testaceis.* — Long.  $2\frac{1}{2}$  lin.

*Er. Col. March. I. 549. 23.*

Habitat in Germania, Gallia: Berolini, Dom. Schüppel, in Saxonia Dom. Märkel, Lutetiae, Dom. Aubé.

Habitus et statura *St. tempestivi*, aterrimus, vix nitidus, pube albida brevissima, subtilissima, parca, vix observanda. Antennae longiores, tenues, articulo tertio quarto paulo longiore, fusco-testaceae, articulis duobus primis nigris. Palpi picei, articulo primo toto, secundo basi testaceo. Caput transversum, thorace sesqui latius, nitidum, minus dense punctatum, fronte vix impressa, late obsoletius bisulcata, sulcorum interstitio parum convexo. Thorax lateribus ante medium modice rotundatis, statim pone medium angustatus, basi coleopteris plus duplo angustior, latitudine summa paulo longior, dense minus fortiter punctatus, punctorum interstitiis planis, dorso medio foveola minuta oblonga, pone medium utrinque obsolete oblique impressus. Coleoptera lata, thorace paulo longiora, parum convexa, minus profunde fortius densiusque punctata, interstitiis planis. Abdomen coleopte-



ris angustius, apicem versus sensim angustatum, parcius subtiliusque punctatum, nitidum. Pedes picei vel rufo-picei, femoribus omnibus basi, anticis cum coxis trochanteribusque dilutioribus, tarsis valde elongatis.

Mas abdominis segmento inferiore quinto apice late leviter emarginato, sexto triangulariter exciso; femina eodem apice leviter producto, rotundato.

Obs. A D. Aubé et Märkel in Formicae rufae societatibus lectus.

- \* 43. *St. femoratus*: Niger, dense punctatus, parce pubescens, palpis articulo primo testaceo, fronte leviter bisulcata, thorace oblongo, medio obsolete canaliculato, elytris thorace longioribus, pedibus ferrugineis. — Long.  $1\frac{1}{2}$  lin.

Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 459. 4.

Habitat in America septentrionali Knoch, e Pennsylvania quoque frequens a Dom. Zimmermann missus.

Statura omnino sequentium, niger, parum nitidus, pube brevissima albida rariore vix cinerascens. Antennae mediocres, articulo tertio quarto paulo longiore, piceae, articulis duobus primis nigris. Palpi fusci, articulo primo toto, secundo basi pallide testaceo. Caput thorace sesqui prope latius, dense fortius profundiusque punctatum, fronte levissime excavata, late bisulcata, interstitio leviter convexo, parum elevato. Thorax coleopteris dimidio angustior, lateribus anterieus rotundatus, posterius angustatus, basi coleopteris plus dimidio angustior, latitudine paulo longior, subcylindricus, fortius crebre punctatus, interstitiis subrugulosis, medio obsolete canaliculatus. Elytra thorace paulo longiora, subdepressa, fortiter profundeque dense punctata, interstitiis subrugulosis, anterieus versus suturam longitudinaliter leviter torosa. Abdomen apicem versus sensim leviter angustatum, late emarginatum, confertim minus subtiliter punctatum. Pedes saturate rufo-testacei, coxis piceis, femoribus apice picescentibus, tarsis modice elongatis.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso, quinto late emarginato; femina segmento sexto apice rotundato.

- \* 44. *St. proditor*: Niger, crebre punctatus, parce pubescens, palpis articulis duobus primis testaceis, fronte fortius bisulcata, thorace oblongo, obsolete canaliculato, elytris thoracis longitudine, femoribus basi rufo-piceis. — Long.  $1\frac{2}{3}$  lin.

Er. Col. March. I. 550. 24.

Habitat in Germania septentrionali.

Statura St. Argi. Niger, parum nitidus, pube brevi albida parca vix cinerascens. Antennae mediocres, articulo tertio quarto sesqui longiore, piceae, articulo primo nigro. Palpi testacei, articulo tertio fusco, secundo nonnunquam apice fusciscente. Caput thorace latius, crebre minus fortiter at profunde punctatum, fronte leviter excavata, fortius bisulcata, interstitio convexo, sat elevato, subcarinato. Thorax coleopteris vix dimidio angustior, ante medium rotundatus, posterius angustatus, basi coleopteris duplo angustior, latitudine summa vix longior, crebre fortius punctatus, interstitiis vix rugulosis, obsolete canaliculatus. Elytra thoracis longitudine, dense profundeque sat fortiter punctata, interstitiis vix rugulosis, anterieus versus suturam longitudi-

dinaliter leviter torosa. Abdomen minus confertim subtilius punctatum. Pedes nigro-picei, femoribus basi ultra medium rufo-piceis, tarsis modice elongatis.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso, quinto longitudinaliter impresso, impressione apicem versus latiore, utrinque carina levi, antrorsum obsolescente terminata, apice inter carinas profundius sinuato; femina segmento sexto apice leviter producto, rotundato.

Variat pedibus totis nigro-piceis.

Obs. A *St. Argo* praeter abdominis in mare structuram differt colore magis nigro, elytris brevioribus, fronte profunde bisulcata, interstitio sat elevato. Forte ob frontis structuram merus *St. Argus* Grav. videatur, quem secundum specimina typica ad speciem mox describendam retuli.

\* 45. *St. excubitor*: Plumbeo-niger, nitidulus, albido-pubescent, crebre fortiterque punctatus, palpis articulo primo testaceo, fronte subimpressa, obsolete bisulcata, thorace oblongo, aequali, elytris thoracis longitudine, pedibus rufo-brunneis. — Long.  $1\frac{3}{4}$  lin.

*Er. Col. March. I.* 551. 25.

Habitat Berolini, Dom. Schüppel.

*St. Argo* proximus, plumbeo-niger, nitidulus, pube subtili albida densiore cinerascens. Antennae mediocres, articulo tertio quarto paulo longiore, obscure piceae, articulo primo nigro. Palpi nigro-picei, articulo primo pallide testaceo. Caput thorace sesqui latius, dense fortius profunde punctatum, fronte levissime excavata, medio late obsoletius bisulcata, interstitio latiore, perparum convexo, vix elevato. Thorax coleopteris paulo angustior, ante medium leviter rotundatus, posterius angustatus, basi coleopteris duplo prope angustior, latitudine summa vix longior, subcylindricus, aequalis, densius fortiter profundeque punctatus, interstitiis punctorum planis. Elytra thoracis longitudine, subdepressa, anterieus versus suturam levissime torosa, densius fortiter profundeque punctata, interstitiis punctorum planis. Abdomen parcius subtiliter punctatum, late marginatum. Pedes rufo-brunnei, tarsis leviter elongatis.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso, quinto late leviter emarginato; femina segmento sexto apice rotundato.

Obs. A praecedente differt colore et frontis structura, a sequente fronte leviter excavata, elytris brevioribus et abdomine minus anguste marginato distinguendus.

\* 46. *St. Argus*: Plumbeo-niger, subnitidulus, albido-pubescent, crebre profundeque punctatus, palpis articulo primo testaceo, fronte obsoletius bisulcata, thorace oblongo, aequali, elytris thorace longioribus, femoribus basi rufo-piceis. — Long.  $1\frac{1}{2}$  —  $1\frac{3}{4}$  lin.

*Grav. Mon.* 231. 12. — *Er. Col. March. I.* 352. 26.

*Sten. opticus* Gyll. *Ins. Suec. IV.* 504. 13 — 14. — *Sahlb. Ins. Fenn. I.* 431. 18. — *Mannerh. Brachelytr.* 44. 20. — *Runde Brachelytr. Hal.* 17. 14.

Habitat in Germania.

Plumbeo-niger, nitidulus, pube brevi depressa albida densiore cinera-

scens. Antennae mediocres, articulo tertio quarto paulo longiore, nigropiceae, articulis duobus primis nigris. Palpi nigri, articulo primo testaceo. Caput thorace paulo latius, dense fortiter profundeque punctatum, fronte planiuscula, obsolete late bisulcata, interstitio perparum convexo. Thorax antérieur leviter rotundatus, coleopteris dimidio angustior, postérius angustatus, basi coleopteris plus duplo angustior, latitudine summa vix longior, subcylindricus, aequalis, dense aequaliter fortiter profundeque punctatus, interstitiis planis. Elytra thorace paulo longiora, leviter convexa, dense aequaliter sat fortiter profundeque punctata, antérieur versus basin obsolete torosa. Abdomen subcylindricum, minus late marginatum, subtilius minus dense punctatum. Pedes picei vel rufo-picei, femoribus basi ultra medium dilutioribus, tarsis leviter elongatis, articulo quarto subcordato.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso, quinto leviter emarginato; femina segmento sexto apice rotundato.

Obs. A praecedentibus abdomine subcylindrico et tarsorum articulo quarto subcordato discedens, ceterum primo intuitu *St. morioni* et cinerascens similis.

\* 47. *St. cautus*: Plumbeo-niger, nitidulus, albido-pubescent, crebre profundeque punctatus, capite elytris dimidio angustiore, fronte fortius bisulcata, thorace oblongo, aequali, elytris thorace longioribus, pedibus palpis antennisque brunneis, his articulo primo nigro. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 553. 27.

Habitat Berolini.

Praecedente paulo minor, statura omnino *St. atratuli*, plumbeo-niger, pube brevi subtili albida densiore cinerascens. Antennae mediocres, articulo tertio quarto paulo longiore, piceae, articulo primo nigro. Palpi picei, articulo primo testaceo. Caput thorace paulo latius, coleopteris plus dimidio angustius, densius fortiter profundeque punctatum, fronte haud excavata, late at parum profunde bisulcata, interstitio convexo, elevato. Thorax coleopteris plus dimidio angustior, lateribus ante medium modice rotundatus, postérius angustatus, basi coleopteris plus duplo angustior, aequalis, densius fortiter profundeque punctatus, interstitiis punctorum planis, vix rugulosis. Elytra thorace tertia parte longiora, subdepressa, densius fortiter profundeque punctata, interstitiis vix rugulosis. Abdomen coleopteris angustius, apicem versus sensim leviter angustatum, late marginatum, densius subtilius punctatum. Pedes rufo-brunnei, tarsis breviusculis.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem rotundato.

Obs. Statura intermedius inter *St. Argum* et *fuscipidem*, ab utroque fronte profundius bisulcata, elytris longioribus et abdomine late marginato distinctus.

\* 48. *St. vafellus*: Niger, nitidulus, albido-pubescent, fortiter punctatus, capite elytrorum latitudine, fronte fortius bisulcata, thorace oblongo, aequali, elytris thorace longioribus, palpis fuscis, articulo primo testaceo, pedibus rufo-brunneis. — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 554. 28.

Habitat Berolini.

Statura omnino *St. fuscipedis*, subplumbeo-niger, nitidulus, pube albida



parca leviter cinerascens. Antennae mediocres, articulo tertio quarto paulo longiore, obscure piceae, articulo primo nigro. Palpi fusci, articulo primo testaceo. Caput thorace paulo latius, coleopteris latitudine aequale, densius fortiter profundeque punctatum, fronte haud excavata, late et parum profunde bisulcata, sulcorum interstitio sat convexo. Thorax coleopteris paulo angustior, lateribus anterieus modice rotundatus, posterius angustatus, basi coleopteris duplo prope angustior, latitudine summa parum longior, subcylindricus, aequalis, densius fortiter profundeque punctatus, punctorum interstitiis planis. Elytra thorace paulo longiora, depressiuscula, aequalia, densius fortiter profundeque punctata, interstitiis planis. Abdomen coleopterorum latitudine, late marginatum, parce subtiliterque punctatum, pube densiore grisescente vestitum. Pedes rufo-brunnei, coxis nigris, tarsis breviusculis.

Mas latet. Femina abdominis segmento inferiore sexto apice subacuminato.

Obs. Affinis praecedenti, at minor, et coleopteris angustioribus distinctus. A *St. fuscipede* punctura subtiliore et abdomine late marginato, a *St. Argo* statura minore, frontis interstitio sulcorum magis elevato et coleopteris angustioribus depressis aequalibus differt.

\* 49. *St. fuscipes*: *Niger, subnitidus, griseo-pubescens, palpis fuscis, articulo primo flavo, pedibus ferrugineis, fronte planiuscula, obsolete bisulcata, thorace oblongo, aequali, elytris thoracis longitudine, abdomine tenuiter marginato.* — Long.  $1\frac{1}{3}$  —  $1\frac{1}{2}$  lin.

*Grav. Micr.* 157. 8. *Mon.* 232. 13. — *Latr. Hist. nat. Crust. et Ins.* IX. 354. 8. — *Gyll. Ins. Suec.* II. 478. 13. — *Sahlb. Ins. Fenn.* I. 431. 17. — *Zetterst. Faun. Lappon.* I. 89. 8. *Ins. Lappon.* 70. 9. — *Mannerh. Brachelytr.* 44. 19. — *Runde Brachelytr. Hal.* 17. 13. — *Er. Col. March.* I. 555. 30.

*Sten. fulvipes* *Boisd. et Lacord. Faun. Ent. Paris.* I. 446. 9.

Habitat in Europa.

*Niger*, subnitidus, pube brevi crassiuscula albida minus conferta subcinerascens. Antennae breviusculae, piceae, articulis duobus primis nigris, tertio quarto sesqui longiore. Palpi nigro-picei, articulo primo testaceo. Caput thorace vix sesqui latius, dense sat fortiter profundeque punctatum, fronte planiuscula, late at parum profunde bisulcata, interstitio lato, perparum elevato, antice subcarinato. Thorax coleopteris parum latior, medio et anterieus satis fortiter rotundatus, posterius angustatus, basi coleopteris dimidio angustior, latitudine paulo longior, subcylindricus, aequalis, densius fortiter profundeque punctatus, interstitiis planis. Elytra thorace vix longiora, apice oblique truncata, aequaliter profunde fortiterque punctata, interstitiis planis, anterieus versus suturam obsolete torulosa. Alae coleopteris vix duplo longiores. Abdomen subcylindricum, tennius marginatum, parcius subtiliusque punctatum, nitidum. Pedes saturate rufo-testacei vel dilutius rufo-brunnei, coxis nigris, geniculis picescentibus, tarsis breviusculis.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem apice leviter producto, acute rotundato.

\* 50. *St. humilis*: *Apterus, niger, densius albido-pubescens, palpis testaceis, articulo tertio fusco, pedibus ferrugineis, fronte planiuscula, bisulcata, elytris depressis, thorace subbrevioribus.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March.* I. 551. 29.

*Sten. fuscipes* Ljungh Web. Beitr. II. 159. 13.

*Sten. Argus* Gyll. Ins. Suec. IV. 503. 12—13. — *Sahlb. Ins. Fenn. I.* 430. 19. — *Mannerh. Brachelytr.* 43. 18.

*Sten. carbonarius* Boisd. et Lacord. Faun. Ent. Paris. I. 450. 17.

Habitat in Germania.

*St. fuscipede* parum maior, depressiusculus, apterus, niger, opacus, pube brevissima albida parciore subcinerascens. Antennae breves, articulo tertio quarto paulo longiore, piceae, articulis duobus primis nigris. Palpi testacei, articulo tertio fusco. Caput thorace paulo latius, dense profundeque punctatum, fronte vix impressa, obsolete late bisulcata, interstitio lato vix convexo elevatoque. Thorax medio coleopteris dimidio prope angustior, lateribus rotundatus, basi coleopteris duplo angustior, latitudine haud longior, subdepressus, creberrime profunde punctatus, interstitiis rugulosis, dorso utrinque leviter oblique impressus. Elytra thoracis vix longitudine, apice oblique truncata, subdepressa, confertissime profunde punctata, interstitiis subrugulosis. Abdomen latius marginatum, dense profunde punctatum, subnitidum. Pedes rufescenti-testacei, coxis nigris, femoribus apice picescentibus, tarsis breviusculis.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso, quinto late longitudinaliter subimpresso, apice emarginato; femina segmento sexto apice rotundato.

\* 51. *St. circularis*: Niger, parum nitidus, thorace rotundato, palpis totis, antennis pedibusque testaceis. — Long. 1 — 1½ lin.

*Grav. Micr.* 157. 8. — *Latr. Hist. nat. Crúst. et Ins. IX.* 354. 9. — *Er. Col. March. I.* 556. 31.

Habitat in Europa.

Minutus, depressiusculus, niger, opacus, pube brevi subtilique albida parcius adpersus. Antennae breviores, articulo tertio quarto parum longiore, piceo-testaceae, articulis duobus primis dilutioribus, apice piceae. Palpi toti flavo-testacei. Caput thorace paulo latius, creberrime punctatum, fronte planiuscula, late et parum profunde bisulcata, interstitio lato, leviter convexo, parum elevato, antice subcarinato. Thorax coleopteris plus dimidio angustior, lateribus sat fortiter rotundatus, posterius fortius angustatus, basi coleopteris triplo angustior, latitudine brevior, parum convexus, confertissime profundius punctatus, interstitiis rugulosis. Elytra thorace sesqui longiora, creberrime profunde punctata, interstitiis rugulosis, anterieus versus suturam obsolete torosa. Abdomen breviusculum, apicem versus sensim attenuatum, late marginatum, dorso subdepressum, subtilius parciusque punctatum, nitidulum. Pedes saturate rufo-testacei, coxis piceis, geniculis vix fuscescentibus, tarsis breviusculis.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem rotundato.

\* 52. *St. declaratus*: Niger, nitidulus, fronte plana, bisulcata, thorace rotundato, palpis pedibusque piceo-testaceis, illis articulo tertio, his geniculis fuscis. — Long. 1½ lin.

*Er. Col. March. I.* 557. 32.

*Sten. circularis* Grav. Mon. 233. 15. — *Gyll. Ins. Suec. II.* 479. 14. — *Sahlb.*

*Ins. Fenn. I. 432. 20. — Zetterst. Ins. Lappon. 71. 13. — Mannerh. Brachelytr. 44. 22. — Boisd. et Lacord. Faun. Ent. Paris. I. 451. 18. — Runde Brachelytr. Hal. 17. 16.*

Habitat in Europa.

Statura minuta et summa similitudo praecedentis. Niger, subnitidus, pube brevissima albida subcinerascens. Antennae breves, articulo tertio quarto paulo longiore, nigro-piceae, articulis duobus primis nigris. Palpi piceo-testacei, articulo tertio nigro-piceo. Caput thorace paulo latius, crebre punctatum, fronte vix impressa, profunde bisulcata, interstitio sat elevato, subcarinato. Thorax coleopteris plus dimidio angustior, lateribus fortiter rotundatus, posterius magis angustatus, coleopteris triplo fere angustior, latitudine summa paulo brevior, convexiusculus, aequalis, creberrime fortius punctatus, interstitiis vix rugulosis. Elytra thorace sesqui longiora, confertim profunde punctata, interstitiis subrugulosis, anterieus versus basin leviter torosa. Abdomen breviusculum, apicem versus sensim angustatum, late marginatum, subtilius parciusque punctatum, nitidulum. Pedes rufo-picei, geniculis obscurioribus, coxis nigris, tarsis breviusculis.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem apice rotundato.

Obs. Differt a praecedente corpore paulo magis depresso, magis cinerascens, fronte profundius bisulcata, interstitio sulcorum magis elevato, thorace elytris minus rugulosis, pedibus obscurioribus, palpis apice piceis, antennis piceis, basi nigris.

\* 53. *St. pumilio*: Niger, parum nitidus, fronte impressa, thorace rotundato, palpis nigris, articulo primo testaceo, pedibus testaceo-piceis. — Long.  $\frac{5}{8}$  lin.

*Er. Col. March. I. 558. 33.*

Habitat in Germania.

Proximus praecedentibus duobus, at dimidio prope minor, niger, vix nitidus, pube albida subtilissima parca haud cinerascens. Antennae breviusculae, articulo tertio quarto subaequali, nigro-piceae, articulis duobus primis nigris. Palpi nigro-picei, articulo primo testaceo. Caput thorace paulo latius, densius profunde fortiter punctatum, fronte levissime excavata, late bisulcata, interstitio angusto, leviter convexo, subcarinato. Thorax coleopteris plus duplo angustior, lateribus rotundatus, posterius vix magis quam anterieus angustatus, basi coleopteris plus duplo angustior, latitudine summa prope brevior, leviter convexus, dense fortius profunde punctatus, interstitiis rugulosis. Elytra thorace sesqui longiora, leviter convexa, fortiter profundeque confertim punctata, interstitiis subrugulosis. Abdomen coleopteris angustius, apicem versus sensim angustatum, crebre profundeque punctatum, angustius marginatum. Pedes obscure testacei vel rufo-picei, geniculis vix obscurioribus, tarsis breviusculis, nigricantibus.

Mas abdominis segmento inferiore sexto apice triangulariter exciso, quinto late subemarginato; femina segmento sexto apice rotundato.

Obs. Praecedentibus minor et minus depressus, a *St. circulari* antennis palpisque nigricantibus et pedibus obscurioribus, a *St. declarato* colore haud cinerascens, frontis structura et palporum articulo secundo piceo distinctus.



## B. Abdomen immarginatum.

- \* 54. *St. nigritulus*: Plumbeo-niger, densius profunde punctatus, albido-pubescent, palpis articulo primo testaceo, abdomine parcius subtiliter punctato. — Long.  $1\frac{1}{4}$  —  $1\frac{3}{4}$  lin.

Gyll. Ins. Suec. IV. 502. 10 — 11. — Sahlb. Ins. Fenn. I. 429. 13. — Zetterst. Faun. Lapp. I. 91. 11. Ins. Lappon. 71. 14. — Mannerh. Brachelytr. 43. 14. — Runde Brachelytr. Hal. 16. 11. — Er. Col. March. I. 559. 34.

Habitat in Europa.

Plumbeo-niger, nitidulus, pube breviuscula depressa albida cinerascens. Antennae breviusculae, articulo tertio quarto paulo longiore; nigrae. Palpi nigri, articulo primo piceo-testaceo. Caput thorace vix latius, dense fortiter profundeque punctatum, fronte planiuscula, obsoletius late bisulcata, interstitio leviter elevato, subcarinato. Thorax coleopteris dimidio angustior, lateribus anterieus rotundatus, posterius angustius, basi coleopteris duplo angustior, latitudine summa paulo longior, subcylindricus, aequalis, densius aequaliterque fortiter profundeque punctatus, interstitiis planis. Elytra thorace paulo longiora, densius aequaliterque fortiter profundeque punctata, interstitiis planis, anterieus versus suturam longitudinaliter leviter torosa. Abdomen breviusculum, apicem versus sensim leviterque angustatum, parcius subtiliusque punctatum. Pedes nigri, tarsi breviusculi.

Mas abdominis segmento inferiore sexto apice levissime emarginato; femina eodem rotundato.

Variat pedibus piceis.

Obs. Magnitudo variat, maiores St. lophthalmus, minores St. fuscipedi aequales, plerique statura St. morionis, ceterum mares plerumque minores, feminae maiores. In Lapponia se nonnisi specimina minimae staturae observasse Dom. Prof. Zetterstedt affert. — Habitu et colore similis St. morioni, cinerascens et imprimis atratulo, abdomine autem immarginato facile distinguendus.

- \* 55. *St. campestris*: Niger, confertissime fortiter punctatus, parcius albido-pubescent, palpis articulo primo testaceo, abdomine dense fortius punctato. — Long. 2 lin.

Er. Col. March. I. 559. 35.

Habitat in Germania.

Affinis praecedenti, at paulo maior, niger, pube depressa albida subcinerascens. Antennae breves, articulo tertio quarto sesqui longiore, nigropiceae, articulo primo nigro. Palpi nigro-picei, articulo primo pallide testaceo. Caput thorace paulo latius, dense fortiter profundeque punctatum, fronte haud impressa, late at parum profunde bisulcata, interstitio lato, levissime convexo, parum elevato. Thorax basi coleopteris dimidio angustior, lateribus anterieus leviter rotundatus, posterius fortius angustatus, basi coleopteris duplo prope angustior, latitudine summa paulo longior, subcylindricus, aequalis, confertissime sat fortiter profundeque punctatus, interstitiis subrugulosis. Elytra thorace vix longiora, profunde fortiterque confertim punctata, interstitiis subrugulosis, anterieus versus suturam obsolete torosa. Abdomen elongatum, cylindricum, segmentis quatuor primis basi leviter constrictis, dense sat fortiter profundeque punctatum. Pedes nigri, tarsi breviusculi, picescentibus.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem leviter producto, subacuminato.

Obs. A praecedente colore magis opaco, statura paulo maiore et magis elongata, abdomine cylindrico, fortius crebriusque punctato distinctus.

\* 56. *St. unicolor*: *Niger, dense fortiterque punctatus, albido-pubescent, palpis piceis, articulo primo testaceo, pedibus antennisque rufo-piceis, his basi nigris, abdomine crebre fortius punctato.* — Long.  $1\frac{2}{3}$  lin.

Habitat in Anglia, Dom. Hope, in Lusitania, Com. Hoffmannsegg, in Hercynia, Dom. Saxesen, in Austria, Dom. Schüppel.

Statura omnino praecedentis, at paulo minor, niger, subnitidus, pube brevi depressa albida cinerascens. Antennae breves, articulo tertio quarto paulo longiore, piceae, articulis duobus primis nigris. Palpi nigro-picei, articulo primo toto, secundo basi pallide testaceo. Caput thorace paulo latius, fortiter profundeque crebre punctatum, fronte haud impressa, at late profundius bisulcata, interstitio convexo, aequae ac frontis latera elevato. Thorax coleopteris parum angustior, lateribus anterieus rotundatus, posterius magis angustatus, basi coleopteris dimidio angustior, latitudine paulo longior, subcylindricus, dense sat fortiter profundeque punctatus, interstitiis subrugulosis. Elytra thoracis longitudine, apice oblique truncata, dense fortiter profundeque punctata, interstitiis planis, aequalia. Alae coleopteris vix duplo longiores. Abdomen cylindricum, segmentis primis basi parum constrictis, crebrius fortiusque punctatum, subnitidum. Pedes rufo-brunnei, coxis geniculisque piceis.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem leviter producto, subacuminato.

Obs. 1. *St. latifronti* similis, at tarsorum articulo quarto integro sat distinctus.

2. Specimina nom. *St. nigri* a Dahlio missa dedit Dom. Schüppel, at quem Com. Mannerheim sub nomine eodem (*Brachelytr.* 43. 13.) descripserit, mihi dubius est, cum abdomine marginato distinctus videatur. *St. nigriritulus* Runde *Brachelytr.* Hal. 16. 11. forte magis ad hanc speciem vel ad *St. latifrontem*, quam ad *Sten. nigriritulum* verum referendus.

\* 57. *St. opticus*: *Niger, confertissime punctatus, parce albido-pubescent, palpis piceis, articulo primo testaceo, antennis pedibusque rufo-piceis, abdomine parcius subtiliter punctato.* — Long.  $1\frac{1}{4}$  lin.

*Grav. Mon.* 231. 11. — *Er. Col. March.* I. 560. 36.

*Sten. femorellus* Zetterst. *Ins. Lappon.* 71. 13.

Habitat in Europa.

Praecedentibus minor, *St. circulari* aequalis, niger, opacus, pube brevissima albida parca vix cinerascens. Antennae breviusculae, articulo tertio quarto paulo longiore, piceae. Palpi picei, articulo primo testaceo. Caput thorace latius, creberrime punctatum, subrugulosum, fronte depressa, sulcis interstitioque elevato omnino fere obsoletis. Thorax coleopteris dimidio prope angustior, lateribus modice rotundatus, posterius magis angustatus, basi coleopteris duplo angustior, latitudine vix longior, subcylindricus, aequalis, confertissime minus fortiter at sat profunde punctatus, interstitiis subrugulosis. Elytra thorace paulo longiora, profunde fortius confertissime

punctata, interstitiis subrugulosis, anterieus versus suturam vix torosa. Abdomen coleopteris angustius, apicem versus sensim attenuatum, parcius subtiliter punctatum, nitidulum. Pedes rufo-picei, coxis nigris.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso, quinto late leviter emarginato; femina segmento sexto apice acuminato.

- \* 58. *St. chalcites*: *Nigro-subaeneus, nitidulus, fortiter punctatus, palpis pallidis, antennis pedibusque flavis, femoribus apice fuscis, abdomine subtiliter punctulato.* — Long.  $1\frac{3}{4}$  lin.

Habitat in Columbia, Dom. Moritz.

Nigro-subaeneus, nitidulus, lateribus parce albido-pubescent. Antennae mediocres, articulo tertio quarto paulo longiore, testaceae, apice obscuriores. Palpi pallide flavescentes. Caput thorace sesqui latius, crebre punctatum, fronte late excavata, obsolete bisulcata, interstitio vix elevato, angusto, evidenter carinato. Thorax coleopteris dimidio angustior, lateribus sat fortiter rotundatus, anterieus et posterius aequaliter angustatus, basi coleopteris duplo angustior, latitudine haud longior, convexiusculus, crebre profundeque punctatus, transversim rugulosus. Elytra thorace paulo longiora, dense sat fortiter profundeque punctata, interstitiis subrugulosis, leviter convexa, anterieus versus suturam leviter torosa. Abdomen coleopteris angustius, apicem versus sensim attenuatum, subtiliter punctulatum, supra segmentis singulis basi subtiliter carinatis, densius subtiliter dorso griseo-, lateribus albido-pubescent. Pedes flavo-testacei, coxis nigris, femoribus apice fuscescentibus, maris leviter incrassatis.

Mas abdominis segmento inferiore sexto profunde inciso; femina eodem apice acute rotundato.

## II. Tarsi articulo quarto bilobo.

### A. Abdomen marginatum.

- \* 59. *St. binotatus*: *Plumbeo-niger, confertim punctatus, densius albido-pubescent, antennis palpisque piceis, his basi testaceis, illis articulo primo nigro.* — Long.  $2\frac{1}{2}$  lin.

*Ljungh Weber u. Mohr Archiv. I. 1. 66. 6. — Grav. Mon. 229. 9. — Gyll. Ins. Suec. II. 474. 9. — Sahlb. Ins. Fenn. I. 427. 9. — Zetterst. Faun. Lappon. I. 89. 7. Ins. Lappon. 70. 8. — Mannrh. Brachelytr. 42. 9. — Boisd. et Lacord. Faun. Ent. Paris. I. 448. 12. — Runde Brachelytr. Hal. 16. 8. — Er. Col. March. I. 561. 37.*

Habitat in Europa.

Elongatus, subdepressus, plumbeo-niger, parum nitidus, pube depressa albida aequaliter densius vestitus, incanus. Antennae breviusculae, articulo tertio quarto sesqui longiore, rufo-piceae, clava fusca, articulo primo nigro. Palpi nigro-picei, articulo primo testaceo. Caput thorace paulo latius, crebrius minus fortiter punctatum, fronte planiuscula, inter oculos utrinque leviter impressa. Thorax coleopteris dimidio angustior, lateribus leviter rotundatus, latitudine summa paulo longior, depresso-cylindricus, crebre punctatus, interstitiis planis, dorso pone medium utrinque oblique impressus. Elytra thorace tertia parte longiora, dense minus fortiter profunde punctata, interstitiis pla-



nis, aequalia. Abdomen crebre minus subtiliter punctatum. Pedes nigri, tarsis nigro-fuscis, brevibus, articulo tertio subcordato, quarto profunde bilobo.

Mas abdominis segmento inferiore sexto apice triangulariter exciso, quarto apice obsoletius impresso; femina segmento sexto apice triangulariter producto.

- \* 60. *St. subimpressus: Plumbeo-niger, subtilius punctatus, densius albido-pubescentis, antennis testaceis, articulo primo nigro, tarsis palpsque piceis, his basi pallidis.* — Long.  $2\frac{3}{4}$  lin.

*Er. Col. March. I. 561. 3.*

Habitat in Germania, Anglia.

St. binotato proximus, at duplo maior, subdepressus, plumbeo-niger, subnitidus, pube depressa albida densiore incanus. Antennae breviores, articulo tertio quarto sesqui longiore, rufo-testaceae, apice vix obscuriores, articulo primo nigro. Palpi nigro-picei, articulo primo toto, secundo basi pallide testaceo. Caput thorace parum latius, crebre minus fortiter at profunde punctatum, fronte utrinque leviter impressa, interstitio leviter longitudinaliter convexo. Thorax coleopteris plus dimidio angustior, lateribus leviter rotundatus, latitudine summa parum longior, subdepressus, subtilius minus confertim punctatus, interstitiis planis, dorso pone medium utrinque oblique impressus. Elytra thorace sesqui prope longiora, crebrius subtilius at profunde punctata, interstitiis planis, anterieus versus suturam obsolete torosa. Abdomen crebrius subtiliter punctatum. Pedes nigri, tarsis fuscescentibus, brevibus, articulo tertio cordato, quarto profunde bilobo.

Mas abdominis segmento inferiore sexto apice profunde semicirculariter exciso, quarto tertioque late semicirculariter impressis, impressione carinula leviter prominente utrinque terminata; femina segmento sexto apice medio triangulariter producto, acuminato.

- \* 61. *St. plantaris: Plumbeo-niger, subtilius punctatus, densius albido-pubescentis, tarsis, palpis antennisque rufo-testaceis, his articulo primo nigro.* — Long.  $2\frac{1}{2}$  lin.

*Er. Col. March. I. 562. 39.*

*Sten. binotatus var. b Gyll. Ins. Suec. IV. 500. 9.*

Habitat in Germania, Suecia.

Statura et summa affinitas St. binotati, at certe distinctus, elongatus, subdepressus, plumbeo-niger, subnitidus, pube depressa densiore incanus. Antennae breviores, articulo tertio quarto paulo longiore, testaceae, articulo primo nigro, clava ferruginea. Palpi toti testacei. Caput thorace paulo latius, subtilius punctatum, inter oculos utrinque leviter impressum, interstitio leviter convexo. Thorax coleopteris duplo prope angustior, lateribus leviter rotundatus, latitudine paulo longior, depresso-subcylindricus, subtilius at sat profunde crebrius punctatus, interstitiis planis, dorso pone medium utrinque oblique impressus. Elytra thorace sesqui fere longiora, densius minus fortiter profunde punctata, interstitiis planis, aequalia. Abdomen subtilius at sat profunde crebrius punctatum. Pedes nigri, tarsis testaceis, brevibus, articulo tertio quartoque bilobis.

Mas abdominis segmento inferiore sexto profunde inciso, quarto tertioque apice leviter impressis; femina segmento sexto apice medio fortiter triangulariter producto, acuminato.

- \* 62. *St. plancus*: *Niger, subtilius punctatus, parcius pubescens, antennis testaceis, articulo primo nigro, palpis testaceis, articulo tertio fusco, femoribus obscure piceis.* — Long. 2 lin.

Habitat in Anglia, Dom. Hope, in Saxonia, Dom. Märkel.

Statura *St. binotati*, at minus elongatus depressusque, niger, nitidulus, parcius cinereo-pubescens. Antennae breviores, articulo tertio quarto paulo longiore, rufo-testaceae, clava fuscescente, articulo primo nigro. Palpi testacei, articulo tertio fusco. Caput thorace paulo latius, crebrius punctatum, fronte minus profunde late bisulcata, interstitio lato, leviter convexo elevatoque. Thorax coleopteris dimidio prope angustior, lateribus anterieus rotundatus, posterius magis angustatus, basi coleopteris duplo prope angustior, latitudine summa vix longior, subcylindricus, minus dense subtilius at sat profunde punctatus, dorso pone medium utrinque obsolete oblique impressus. Elytra thorace paulo longiora, densius fortiusque profunde punctata, interstitiis planis, parum inaequalia, versus suturam longitudinaliter torulosa, anterieus callosa. Abdomen subcylindricum, licet distincte marginatum, crebrius subtiliter punctatum. Pedes nigri, femoribus picescentibus, tarsis brevibus, dilatatis, articulo primo triangulari, secundo tertioque transversis, integris, quarto bilobo.

Mas abdominis segmento inferiore sexto apice triangulariter exciso, quinto quartoque longitudinaliter impressis, impressione utrinque carinula obtusa terminata, illo medio leviter excavato; femina segmento sexto apice acute rotundato.

- \* 63. *St. bifoveolatus*: *Niger, nitidulus, dense punctatus, parcius albido-pubescens, palpis antennisque testaceis, his articulo primo piceo, pedibus piceo-testaceis, geniculis nigris, elytris thoracis longitudine.* — Long. 1½ lin.

*Gyll. Ins. Suec. IV. 500. 9 — 10. — Sahlb. Ins. Fenn. I. 427 10. — Mannerh. Brachelytr. 43. 10.*

Habitat in Suecia, Fennia, Dom. Schüppel.

Statura subcylindrica praecedentis, at paulo minor, niger, nitidulus, pube brevi depressa albida parcius aequaliter adpersus, subcinerascens. Antennae breviusculae, articulo tertio quarto paulo longiore, testaceae, articulo primo piceo, clava fuscescente. Palpi testacei. Caput thorace paulo latius, crebre profundius punctatum, fronte depressiuscula, obsolete late bisulcata, interstitio latiusculo, leviter convexo, paululum magis quam frontis latera elevato. Thorax coleopteris dimidio angustior, lateribus anterieus leviter rotundatus, posterius magis angustatus, basi coleopteris duplo angustior, subcylindricus, dense fortius profunde punctatus, interstitiis planis, utrinque obsolete oblique impressus, ceterum aequalis. Elytra thoracis longitudine, aequalia, densius fortiter profunde punctata, punctorum interstitiis planis. Abdomen breviusculum, subcylindricum, licet distincte marginatum, dense profundeque punctatum. Pedes piceo-testacei, coxis nigris, femoribus apice

nigricantibus, tarsis brevibus, articulis duobus primis elongato-triangularibus, tertio triangulari, quarto bilobo.

Mas abdominis segmento inferiore sexto triangulariter exciso (Gyll); femina eodem apice triangulariter producto, leviter acuminato.

\* 64. *St. rusticus*: Plumbeo-niger, subnitidus, dense punctatus, albido-pubescent, palpis antennisque testaceis, his articulo primo nigro, pedibus piceo-testaceis, geniculis nigris, elytris thorace longioribus, — Long. 2 lin.

Habitat in Germania meridionali, Dom. Schüppel, Lutetiae, Dom. Aubé, in Anglia, Dom. Rudd.

Habitu fere *St. tarsalis*, et minoribus eius individuis aequalis, plumbeo-niger, nitidulus, pube brevi depressa albida aequali minus densa cinerascens. Antennae mediocres, articulo tertio quarto paulo longiore, rufo-testaceae, articulo primo nigro, clava fuscescens. Palpi toti rufo-testacei. Caput thorace paulo latius, dense profundius punctatum, fronte depressiuscula, late bisulcata, interstitio sulcorum latiusculo, leviter convexo, aequae ac frontis latera elevato. Thorax coleopteris plus dimidio angustior, lateribus anterieus leviter rotundatus, posterius magis angustatus, basi coleopteris plus duplo angustior, dense profundeque punctatus, punctorum interstitiis planis, pone medium utrinque obsolete oblique impressus, ceterum aequalis. Elytra thorace tertia parte longiora, dorso longitudinaliter leviter toroso, dense profundeque punctata, interstitiis planis. Abdomen coleopteris paulo angustius, apicem versus leviter angustatum, subcylindricum, licet lateribus distincte marginatum, sat dense profundeque punctatum. Pedes piceo-testacei, coxis nigris, femoribus apice nigricantibus, tarsis brevibus, articulis duobus primis linearibus, parum elongatis, tertio subtriangulari, quarto bilobo.

Mas abdominis segmento inferiore sexto triangulariter exciso; femina eodem medio subacuminato.

\* 65. *St. tempestivus*: Apterus, niger, nitidus, subtilius punctatus, parce pubescens, palpis antennisque testaceis, his articulo primo nigro, pedibus testaceis, geniculis nigris. — Long.  $2\frac{1}{3}$  lin.

*Er. Col. March. I. 563. 40.*

Habitat in Germania, Gallia, Anglia.

*St. binotati* speciminibus maioribus haud angustior, at brevior, apterus, subdepressus, niger, nitidus, pube brevissima parca albida haud cinerascens. Antennae mediocres, articulo tertio quarto paulo longiore, testaceae, articulo primo nigro, clava fuscescens. Palpi testacei. Caput thorace sesqui latius, crebrius minus fortiter punctatum, fronte late profundeque bisulcata, interstitio angustiore, aequae ac frontis latera elevato, subcarinato. Thorax coleopteris medio dimidio, basi duplo angustior, lateribus leviter rotundatus, latitudine summa vix longior, subdepresso-cylindricus, minus confertim subtilius punctatus, interstitiis planis, subinaequalis, dorso pone medium utrinque oblique impressus. Elytra thorace parum longiora, minus confertim subtilius punctata, interstitiis punctorum planis, inaequalia, apice oblique truncata. Abdomen parcius subtiliusque punctatum. Pedes testacei, coxis



nigris, femoribus apice tibiisque basi piceis, tarsis breviusculis, articulo primo elongato-triangulari, secundo triangulari, tertio obcordato, quarto bilobo.

Mas abdominis segmento inferiore sexto apice triangulariter exciso; femina eodem apice utrinque subsinuato, medio triangulariter producto, acuminato.

\* 66. *St. picipennis*: *Niger, nitidulus, subtilius punctatus, parce pubescens, antennis palpisque testaceis, pedibus testaceis, medio fuscescentibus, elytris piceis.* — Long. 2 lin.

Habitat in Saxonia, Dom. Märkel.

Praecedenti affinis, at brevior et minus depressus, niger, nitidulus, pube brevi depressa albida parce vestitus. Antennae breviusculae, articulo tertio quarto paulo longiore, dilute testaceae, articulo primo concolore, clava fuscescente. Palpi dilute testacei. Caput thorace latius, transversum, sat crebre profundeque minus fortiter punctatum, fronte late minus profunde bisulcata, interstitio convexo et paulo magis quam frontis latera elevato, subcarinato. Thorax coleopteris dimidio angustior, lateribus ante medium fortius rotundatus, pone medium fortius angustatus, basi coleopteris duplo angustior, latitudine summa haud longior, minus dense fortiterque licet profundius punctatus, interstitiis planis, dorso medio utrinque oblique leviter impressus, ceterum aequalis. Elytra thorace paulo longiora, densius fortiusque profunde punctata, interstitiis planis, inaequalia, picea. Abdomen breviusculum, parcius subtiliter punctatum, segmentis dorsalibus 4 primis margine apicali subelevato, polito. Pedes testacei, coxis, femoribus apice, tibiis basi piceis, tarsis brevibus, articulis duobus primis subtriangularibus, tertio transversim obcordato, quarto profunde bilobo.

Mas latet. Femina abdominis segmento inferiore sexto apice testaceo-marginato, medio leviter triangulariter producto.

\* 67. *St. languidus*: *Niger, nitidulus, subtilius punctatus, parcius pubescens, palpis antennisque obscure testaceis, his articulo primo nigro, pedibus piceis, basi apiceque obscure testaceis.* — Long. 2½ lin.

Habitat in Sicilia, Dom. Dr. Schultz.

Habitu fere *St. cordati*, at duplo minor, niger, nitidulus, pube brevissima depressa albida minus dense vestitus, vix cinerascens. Antennae mediocres, articulo tertio quarto sesqui longiore, obscure rufo-testaceae, clava fuscescente, articulo primo nigro. Palpi testacei, articulo tertio picescente. Caput thorace paulo latius, crebrius subtiliusque punctatum, fronte late profundeque bisulcata, interstitio latiusculo, leviter convexo, aequae ac frontis latera elevato, laeviore. Thorax basi coleopteris duplo angustior, lateribus ante medium leviter rotundatis, latitudine summa haud longior, parum convexus, parcius subtiliter punctatus, dorso pone medium utrinque oblique fortius impressus, ceterum aequalis. Elytra thorace sesqui prope longiora, dense crebre subtilius punctata, inaequalia. Abdomen basi coleopteris paulo angustius, apicem versus sensim attenuatum, parcius subtiliter punctatum. Pedes picei, femoribus basi ultra medium, tibiis apice tarsisque obscure rufo-testaceis, his breviusculis, articulis duobus primis linearibus, tertio subtriangulari, integro, quarto profunde bilobo.

Mas abdominis segmento inferiore sexto apice triangulariter emarginato; femina eodem apice utrinque subsinuato, medio triangulariter producto, acuminato.

- \* 68. *St. cordatus*: *Niger, nitidus, fortius crebre punctatus, parce pubescens, palpis antennisque elongatis testaceis, his articulo primo piceo, pedibus testaceis, femoribus apice, tibiis basi nigris.* — Long. 3 lin.

Grav. Micr. 198. 1. Mon. 226. 3.

Habitat in Lusitania, Com. Hoffmannsegg, in Sardinia, Prof. Gené.

Subdepressus, niger, nitidus, pube brevior depressa albida parcius vestitus. Antennae elongatae, capitis thoracisque fere longitudine, articulo tertio quarto paulo longiore, clava parum incrassata, rufo-piceae, articulo primo piceo. Palpi elongati, articulo tertio minus incrassato, testacei. Caput thorace vix latius, fortiter crebrius punctatum, fronte late et parum profunde bisulcata, interstitio leviter convexo, aequae ac frontis latera elevato. Thorax coleopteris duplo prope angustior, lateribus ante medium fortiter rotundatus, posterius fortius angustatus, basi coleopteris triplo fere angustior, latitudine summa haud longior, parum convexus, crebre fortius punctatus, dorso medio late canaliculatus, pone medium utrinque leviter impressus. Elytra lata, thorace paulo longiora, fortius profunde dense punctata, inaequalia, anterieus versus suturam distinctius callosa. Abdomen basi coleopterorum latitudine, apicem versus sensim fortius attenuatum, parce subtiliter punctatum, segmentis singulis basi e pube densiore cinereo-versicoloribus. Pedes testacei, femoribus apice late nigris, tibiis basi nigris, tarsis longioribus, articulo primo sat elongato, quarto angusto, bilobo.

Mas abdominis segmento inferiore sexto apice emarginato; femina eodem apice medio leviter producto, rotundato.

- \* 69. *St. hospes*: *Niger, subnitidus, dense fortiter punctatus, parce albido-pubescens, antennis elongatis, piceis, palpis pedibusque testaceis, his geniculis nigris.* — Long. 2 $\frac{3}{4}$  lin.

Habitat in Coreyra, Dóm. Schüppel,

Habitu fere praecedentis, at gracilior et duplo minor, niger, nitidus, pube longiore depressa albida parcius vestitus. Antennae elongatae, graciles, articulo tertio quarto paulo longiore, testaceo-piceae, clava fuscescente. Palpi elongati, dilute testacei, articulo tertio obscuriore, minus incrassato. Caput thorace paulo latius, crebre fortiter profundeque punctatum, fronte utrinque fortius late longitudinaliter excavata, interstitio convexo, aequae fere ac frontis latera elevato, parcius punctato. Thorax coleopteris plus dimidio angustior, lateribus medio modice rotundatus, basin et apicem versus aequaliter angustatus et ibidem coleopteris duplo angustior, latitudine summa paulo longior, subcylindricus, dense fortiter profundeque punctatus, interstitiis subrugulosis, dorso leviter canaliculatus et utrinque obsolete bi-impressus. Elytra thorace tertia parte longiora, leviter inaequalia, dorso distinctius longitudinaliter toroso, dense fortiter profundeque punctata, interstitiis subaequalibus. Abdomen elongatum, basi coleopteris paulo angustius, basin versus sensim attenuatum, apicem versus sensim subtilius punctatum, nitidum. Pedes testacei, coxis nigris, femoribus apice late nigricantibus,

tibiis basi piceis, tarsis longioribus, articulis tribus primis linearibus, primo sat elongato, quarto leviter bilobo.

Mas abdominis segmento inferiore sexto apice leviter emarginato. Femina latet.

\* 70. *St. subaeneus*: *Nigro-subaeneus, nitidulus, dense fortius punctatus, densius albido-pubescent, antennis medio, palpis basi testaceis, pedibus testaceis, medio piceis.* — Long. 2 lin.

*Sten. geniculatus* Mannerh. *Brachelytr.* 43. 15.

Habitat Lutetiae, Dom. Aubé, in Gallia meridionali, Dom. Schüppel, in Sardinia, Dom. Prof. Gené, in Anglia, Dom. Rudd.

Statura *St. impressi*, nigro-subaeneus, nitidulus, pube depressa albida parciore subcinerascens. Antennae longiusculae, graciles, articulo tertio quarto paulo longiore, obscure testaceae, articulis duobus primis nigris, clava picea. Palpi articulo primo dilute, secundo obscure testaceo, tertio piceo. Caput thorace paulo latius, fortius dense profundeque punctatum, fronte utrinque leviter longitudinaliter excavata, interstitio leviter convexo, aequae fere ac frontis latera elevato, medio paulo laeviore. Thorax coleopteris plus dimidio angustior, lateribus medio modice rotundatus, posterius paulo magis angustatus, basi coleopteris duplo angustior, latitudine summa parum longior, subcylindricus, confertissime fortiter profundeque punctatus, interstitiis rugulosis, dorso leviter canaliculatus et utrinque obsoletius bi-impressus. Elytra thorace tertia parte longiora, dense fortiter profundeque punctata, interstitiis vix rugulosis, dorso longitudinaliter leviter toruloso. Abdomen coleopteris paulo angustius, apicem versus sensim subangustatum, dense sat fortiter, licet apicem versus sensim subtilius punctatum. Pedes saturate testacei, coxis nigris, femoribus apice, tibiis basi late piceis, tarsis longioribus, articulis tribus primis linearibus, primo sat elongato, quarto profunde bilobo.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem rotundato.

Obs. Minus accurate a Mannerheimio descriptus. Specimen a Com. Dejean ipso sub eodem nomine communicatum dedit Dom. Schüppel.

\* 71. *St. aerosus*: *Nigro-aeneus, nitidus, dense punctatus, tenuiter albido-pubescent, antennis palpisque testaceis, illis articulo primo nigro, his apice fusciscentibus, pedibus flavo-testaceis, geniculis nigricantibus.* — Long. 2 lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura omnino praecedentis, laete nigro-aeneus, nitidus, pube subtili brevissima depressa albida parciore haud cinerascens. Antennae sat elongatae, tenues, articulo quarto tertio vix brevior, testaceae, articulo primo nigro, clava fusciscente. Palpi pallide testacei, articulo tertio apice fusciscente. Caput thorace sesqui fere latius, sat dense profunde punctatum, fronte late bisulcata, interstitio leviter convexo, aequae ac frontis latera elevato, medio longitudinaliter laeviore. Thorax coleopteris dimidio angustior, lateribus ante medium modice rotundatis, posterius magis angustatus, basi coleopteris duplo prope angustior, latitudine summa vix longior, dense for-



tius profunde punctatus, interstitiis subrugulosis, leviter canaliculatus et dorso utrinque obsoletius bi-impressus. Elytra thorace tertia parte longiora, dense profundeque fortius punctata, interstitiis subaequalibus, leviter inaequalia, dorso longitudinaliter distinctius torosa. Abdomen apicem versus sensim subangustatum, crebre minus subtiliter punctatum. Pedes flavo-testacei, coxis, femoribus apice, tibiis basi piceis, tarsis longioribus, articulis tribus primis linearibus, primo sat elongato, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto apice leviter emarginato; femina eodem rotundato.

Obs. A praecedente colore, punctura minus forti et capite latiore, a St. impresso punctura densiore, elytris longioribus, antennarum articulo primo nigro, palpis apice geniculisque piceis distinctus.

X \* 72. *St. impressus*: Nigro-subaeneus, nitidus, fortius punctatus, parce albido-pubescentis, antennis palpisque testaceis, pedibus flavis. — Long. 2 lin.

Germ. Spec. Ins. 36. 59. — Er. Col. March. I. 564. 41.

*Stenus proboscideus* Germ. Faun. Ins. Europ. XV. 1.

*Stenus Aceris* Boisd. et Lacord. Faun. Ent. Paris. I. 445. 7.

*Stenus pallipes* var. Grav. Mon. p. 233. (forte)

Habitat in Germania, Gallia, Anglia.

Subdepressus, nigro-subaeneus, nitidulus, pube brevissima albida parcius adpersus. Antennae mediocres, graciliores, articulo tertio quarto sesqui longiore, totae testaceae. Palpi testacei. Caput thorace sesqui latius, crebre punctatum, fronte haud excavata, at late profundeque bisulcata, interstitio convexo, aequae ac frontis latera elevato, linea media longitudinali laevi. Thorax coleopteris dimidio angustior, lateribus ante medium modice rotundatus, posterius magis angustatus, basi coleopteris duplo angustior, latitudine summa vix longior, parum convexus, densius punctatus, interstitiis subinaequalibus, dorso medio obsoletius canaliculatus, utrinque et ante et pone medium obsolete impressus. Elytra thorace vix longiora, densius fortiusque punctata, subinaequalia. Alae elytris duplo longiores. Abdomen apicem versus sensim angustatum, crebre minus subtiliter punctatum, nigrum, nitidum. Pedes flavo-testacei, coxis piceis, tarsis longiusculis, articulo primo sat elongato, quarto profunde bilobo.

Mas abdominis segmento inferiore sexto apice acute emarginato; femina rotundato.

\* 73. *St. geniculatus*: Niger, fortiter punctatus, leviter albido-pubescentis, antennis palpisque testaceis, apice fusciscentibus, pedibus testaceis, geniculis fuscis. — Long. 2 lin.

Grav. Mon. p. 228. — Er. Col. March. I. 564. 42.

Habitat in Germania.

Elongatus, niger, minus nitidus; pube brevi depressa albida cinerascens. Antennae graciliores, articulo tertio quarto sesqui longiore, testaceae, clava fusciscente, articulo primo piceo. Palpi testacei, articulo tertio apice fusciscente. Caput thorace paulo latius, crebre sat profunde punctatum, fronte haud excavata, at late et parum profunde bisulcata, interstitio lato,

parum elevato, linea tenui subelevata laevi subcarinato. Thorax coleopteris dimidio angustior, lateribus medio modice rotundatus, posterius paulo magis quam anterie angustatus, basi coleopteris duplo fere angustior, latitudine summa paulo longior, subcylindricus, creberrime sat fortiter punctatus, interstitiis subrugulosis, dorso obsoletissime canaliculatus. Elytra thorace vix longiora, creberrime fortiter punctata, interstitiis subrugulosis, anterie versus suturam longitudinaliter leviter torulosa. Alae integrae. Abdomen elongatum, fortius crebre punctatum. Pedes testacei, coxis piceis, geniculis fuscis, tarsis elongatis, gracilibus, articulis 1—3 linearibus, primo sat elongato, quarto angusto, bilobo.

Mas abdominis segmento inferiore sexto apice leviter sinuato; femina eodem rotundato.

\* 74. *St. flavipes*: *Niger, nitidulus, fortius punctatus, parcius albido-pubescentis, antennis, palpis pedibusque flavis, immaculatis, thorace medio obsoletius canaliculato, abdomine late marginato.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I.* 566. 44.

Habitat in Germania.

Habitu omnino *St. impressi*, at paulo minor, subdepressus, niger, nitidus, pube parca subtili depressa albida adpersus. Antennae longiusculae, articulo tertio quarto parum longiore, flavo-testaceae, articulo primo concolore, clava fusciscente. Palpi flavo-testacei. Caput coleopterorum latitudine, densius fortiusque profunde punctatum, fronte late parum profunde bisulcata, interstitio leviter convexo, aequae ac frontis latera elevato, linea longitudinali laevi subcarinato. Thorax coleopteris paulo angustior, lateribus ante medium rotundatus, posterius fortius angustatus, basi coleopteris duplo angustior, latitudine summa vix brevior, parum convexus, densius fortiusque profunde punctatus, interstitiis subaequalibus, dorso leviter canaliculatus, canalicula basi apiceque obsolescente, medio utrinque obsolete oblique impressus. Elytra thorace paululum longiora, depressiuscula, vix inaequalia, densius fortiter profundeque punctata, interstitiis planis. Alarum rudimenta elytris breviora. Abdomen apicem versus sensim attenuatum, late marginatum, minus dense et apicem versus sensim subtilius punctatum. Pedes flavo-testacei, coxis anticis rufo-testaceis, posterioribus rufo-piceis, tarsis longiusculis, articulis tribus primis linearibus, primo sat elongato, quarto bilobo.

Mas abdominis segmento inferiore sexto apice emarginato; femina eodem rotundato.

\* 75. *St. palustris*: *Niger, nitidulus, fortius punctatus, parcius albido-pubescentis, antennis, palpis pedibusque flavis, antennis articulo primo, palpis articulo tertio, femoribus apice, tibiis basi late fuscis, thorace medio obsoletius canaliculato.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I.* 565. 43.

*Stenus proboscideus* Gyll. *Ins. Suec. II.* 476. 11. — *Sahlb. Ins. Fenn. I.* 429. 14. — *Zetterst. Faun. Lappon. I.* 88. 6. *Ins. Lappon. 70.* 7. — *Mannerh. Brachelytr. 43.* 16.

Habitat in Germania.

Statura et summa affinitas praecedentis. Niger, nitidulus, pube brevi depressa parciore albida parum cinerascens. Antennae longiusculae, articulo tertio quarto vix sesqui longiore, rufo-testaceae, extrorsum fuscae, articulo primo piceo. Palpi testacei, articulo tertio picescente. Caput thorace sesqui latius, crebre dense punctatum, fronte late parum profunde bisulcata, interstitio lato, leviter convexo, aequae ac frontis latera elevato, medio laeviore, subcarinato. Thorax coleopteris paulo angustior, lateribus ante medium lateribus sat fortiter rotundatus, posterius magis angustatus, basi coleopteris dimidio angustior, latitudine summa vix brevior, parum convexus, crebrius fortiusque punctatus, interstitiis subrugulosis, medio leviter canaliculatus, pone medium utrinque obsolete oblique impressus. Elytra thorace parum longiora, densius fortiusque profunde punctata, subinaequalia, anterieus versus suturam subcallosa. Alarum rudimenta coleopteris breviora. Abdomen late marginatum, dense anterioribus fortius, posterius subtilius punctatum. Pedes saturate testacei, coxis nigris, femoribus apice late et tibiis basi fusco-piceis, tarsis longiusculis, articulis tribus primis linearibus, primo sat elongato, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso; femina eodem rotundato.

Obs. Distinctus a praecedente praeter antennarum, palporum pedumque constantem colorem thorace paulo longiore, posterius paulo minus angustato, tarsis minus elongatis, articulo quarto fortius bilobo, abdominis segmento inferiore sexto in mare fortius emarginato, etc.

\* 76. *St. fuscicornis*: Niger, nitidulus, fortiter punctatus, antennis palpisque piceis, his basi testaceis, pedibus piceo-testaceis, medio late nigricantibus. — Long.  $1\frac{1}{2}$  lin.

Habitat Lutetiae, Dom. Aubé, in Helvetia, Dom. Godet.

Statura et summa affinitas praecedentis, niger, nitidulus, pube brevi depressa parca albida vix cinerascens. Antennae longiusculae, articulo tertio quarto vix longiore, fusco-piceae, basi apiceque nigricantes. Palpi fusco-picei, articulo primo pallide testaceo. Caput thorace sesqui latius, fortius profundeque dense punctatum, fronte late bisulcata, interstitio convexo, aequae ac frontis latera elevato. Thorax coleopteris paulo angustior, lateribus medio modice rotundatus, basin apicemque versus fere aequaliter angustatus et ibidem coleopteris dimidio angustior, latitudine summa haud longior, leviter convexus, dense sat fortiter profundeque punctatus, interstitiis praecipue latera versus subrugulosis, medio utrinque obsolete impressus. Elytra thorace paululum longiora, dense fortiter profundeque punctata, interstitiis subaequalibus, perparum convexa, sutura longitudinaliter depressiuscula. Abdomen apicem versus sensim subangustatum, late marginatum, dense, segmentis anterioribus 4 fortius, posterioribus 2 subtilius punctatum. Pedes rufo-picei, coxis nigris, femoribus apice, tibiis basi late nigricantibus, tarsis longioribus, articulis tribus primis linearibus, primo sat elongato, quarto fortiter bilobo.

Mas latet. Femina abdominis segmento inferiore sexto apice acute rotundato.

Obs. Proximus praecedenti, distinctus autem antennis, palpis pedibusque obscurioribus, punctura crebriore et fortiore, thorace haud canaliculato, posterius paulo minus angustato.



- \* 77. *St. pallipes*: *Niger, subnitidus, dense fortius punctatus, densius albido-pubescent, antennis, palpis pedibusque testaceis, abdomine subcylindrico, subtiliter marginato.* — Long.  $1\frac{2}{3}$  lin.

*Grav. Micr.* 157. 7. *Mon.* 233. 14. — *Ljungh Web. et Mohr Beitr.* II. 156. 10. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 416. 8. — *Er. Col. March.* I. 567. 45.

Habitat in Germania.

Statura et summa affinitas *St. flavipedis*, *niger*, *subnitidus*, pube brevissima depressa albida subcinerascens. Antennae longiusculae, tenues, articulo tertio quarto sesqui longiore, testaceae, articulo primo concolore, clava fuscescente. Palpi flavo-testacei. Caput thorace paulo latius, crebre punctato-subrugulosum, fronte levissime excavata, late obsolete bisulcata, interstitio lato, perparum convexo elevatoque. Thorax coleopteris paulo angustior, lateribus sat fortiter rotundatus, posterius paulo magis quam anterieus angustatus, basi coleopteris plus dimidio angustior, latitudine summa haud longior, parum convexus, dense fortius punctatus, interstitiis subaequalibus, obsolete canaliculatus. Elytra thorace vix longiora, dense fortius sat profunde punctata, interstitiis subinaequalibus, anterieus versus suturam leviter torosa. Abdomen coleopteris paulo angustius, apicem versus sensim attenuatum, tenuiter marginatum, subcylindricum, dense anterieus fortius, posterius subtilius punctatum. Pedes flavo-testacei, coxis posterioribus piceis, tarsis longiusculis, articulis tribus primis linearibus, primo sat elongato, quarto bilobo.

Mas abdominis segmento inferiore sexto apice leviter triangulariter exciso; femina eodem apice medio leviter triangulariter producto.

Obs. A *St. flavipede* corpore angustiore, magis cinerascens, thorace medio ipso neque ante medium latiore, abdomine subtiliter marginato.

- \* 78. *St. filum*: *Apterus, elongatus, subdepressus, niger, nitidulus, parce subtiliter punctatus, parcius albido-pubescent, pedibus, palpis antennisque flavis, his articulo primo nigro.* — Long.  $1\frac{2}{3}$  lin.

*Er. Col. March.* I. 568. 46.

Habitat in Germania, Gallia, Anglia.

*St. flavipede* haud brevior, at dimidio angustior et magis linearis, subdepressus, *niger*, *nitidulus*, pube brevi depressa parca albida subcinerascens. Antennae breves, articulo tertio quarto sesqui longiore, flavae, clava apice fuscescente, articulo primo nigro. Palpi flavi. Caput thorace sesqui latius, parce punctatum, fronte depressa, late minus profunde bisulcata, interstitio lato, perparum elevato, laeviore. Thorax coleopteris paulo angustior, medio leviter rotundatus, posterius paulo magis angustatus, basi coleopteris vix dimidio angustior, latitudine summa paulo longior, depresso-subcylindricus, parce subtilius punctatus, aequalis. Elytra thorace parum longiora, parcius subtiliusque punctata, aequalia. Alarum rudimenta coleopteris breviora. Abdomen elongatum, lineare, late marginatum, parcius subtiliusque punctatum. Pedes breviusculi, flavi, coxis nigris, tarsis brevibus, dilatatis, articulo tertio subcordato, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto apice triangulariter emarginato, quinto longitudinaliter subimpresso; femina segmento sexto apice triangulariter producto, subacuminato.

## B. Abdomen immarginatum.

- \*79. *St. tarsalis*: Plumbeo-niger, densius albido-pubescent, crebre punctatus, tarsis, palpis antennisque rufo-testaceis, his articulo primo nigro, abdomine crebrius punctato. — Long. 2 —  $2\frac{1}{3}$  lin.

Ljungh Web. u. Mohr Beitr. II. 157. 11. — Gyll. Ins. Suec. II. 472. 8. — Sahlb. Ins. Fenn. I. 426. 8. — Zetterst. Faun. Lappon. I. 90. 10. Ins. Lappon. 71. 12. — Mannerh. Brachelytr. 42. 8. — Er. Col. March. I. 569. 47.

*Sten. buphthalmus* Ljungh Web. et Mohr Archiv. I. 1. 67. 7.

*Sten. clavicornis* Grav. Micr. 156. 5. Mon. 229. 7. — Boisd. et Lacord. Faun. Ent. Paris. I. 448. 13.

*Staph. clavicornis* Rossi Faun. Etr. ed. Hellw. I. p. 312. not. n. 1.

*Stenus riparius* Runde Brachelytr. Hal. 16. 10.

Habitat in Europa.

Plumbeo-niger, nitidulus, pube brevi depressa albida densiore cinerascens. Antennae mediocres, articulo tertio quartoque subaequalibus, rufo-testaceae, clava fusciscente, articulo primo minuto, nigro. Palpi rufo-testacei. Caput thorace paulo latius, crebre subtilius punctatum, fronte depressiuscula utrinque leviter impressa. Thorax coleopteris dimidio angustior, lateribus medio leviter rotundatus, posterius paulo magis angustatus, basi coleopteris duplo angustior, latitudine summa haud longior, profunde crebrius punctatus, interstitiis planis, dorso utrinque pone medium oblique leviter impressus. Elytra thorace paulo longiora, densius fortiusque profunde punctata, anteriùs versus basin longitudinaliter leviter torosa. Abdomen apicem versus sensim attenuatum, minus subtiliter densius punctatum. Pedes corpori concolores, tarsis rufo-testaceis, breviusculis, dilatatis, articulo tertio obcordato, quarto bilobo.

Mas abdominis segmento inferiore sexto apice triangulariter emarginato; femina eodem apice leviter bisinuato, medio leviter triangulariter producto.

Obs. *Steni riparii* Rund. specimen inspiciendum benevole communicavit Dom. Prof. Germar.

- \*80. *St. vestigialis*: Plumbeo-niger, albido-pubescent, crebre punctatus, tarsis, palpis antennisque testaceis, abdomine crebrius punctato. — Long.  $2\frac{1}{3}$  lin.

Habitat in Mexico.

*St. tarsali* gracilior, plumbeo-niger, subnitidus, pube subtili depressa albida subcinerascens. Antennae mediocres, articulo tertio quarto paulo longiore, totae testaceae. Palpi testacei. Caput thorace sesqui latius, profunde crebre punctatum, fronte depressiuscula, utrinque leviter impressa. Thorax elongatus, cylindricus, basi coleopteris duplo angustior, medio perparum rotundato-incrassatus, latitudine summa plus sesqui longior, profunde confertissime punctatus, interstitiis subrugulosis, dorso pone medium oblique obsolete utrinque impressus. Elytra thoracis longitudine, dense profundeque punctata, basi medio leviter torulosa. Abdomen coleopteris paulo angustius, cylindricum, fortius profunde dense punctatum. Pedes corpori concolores, tarsis testaceis, longioribus, articulis duobus primis linearibus, primo elongato, tertio obcordato, quarto bilobo.

Mas latet. Femina abdominis segmento inferiore sexto apice leviter triangulariter producto, subacuminato.

Obs. St. tarsali affinis, antennarum articulo primo testaceo thoraceque elongato, cylindrico facile distinguendus.

\* 81. *St. oculatus*: Plumbeo-niger, crebre subtilius punctatus, albido-pubescent, palpis antennisque testaceis, his articulo primo nigro, pedibus flavis, geniculis nigricantibus. — Long.  $2\frac{1}{2}$  lin.

Grav. Micr. 155. 3. Mon. 227. 5. — Gyll. Ins. Suec. II. 471 7. — Sahlb. Ins. Fenn. I. 425. 7. — Zetterst. Faun. Lappon. I. 90. 9. Ins. Lappon. 71. 11. — Mannerh. Brachelytr. 42. 7. — Boisd. et Lacord. Faun. Ent. Paris. I. 444. 5. — Runde Brachelytr. Hal. 15. 7. — Er. Col. March. I. 569. 48.

*Staph. similis* Herbst Archiv. 151. 15.

*Sten. similis* Ljungh Web. u. Mohr Archiv. I. 1. 65. 5.

Habitat in Europa.

Plumbeo-niger, subnitidus, pube depressa albida densiore cinerascens. Antennae longiores, articulo tertio quarto sesqui longiore, testaceae, articulo primo nigro. Palpi testacei. Caput thorace paulo latius, crebre punctatum fronte utrinque late at parum profunde longitudinaliter impressa, interstitio lato, levissime convexo, vix elevato. Thorax coleopteris dimidio angustior, lateribus medio modice rotundatus, posterius vix magis quam anterie angustatus, basi coleopteris duplo prope angustior, latitudine summa haud longior, subcylindricus, dense punctatus, interstitiis subaequalibus, dorso pone medium utrinque oblique obsolete impressus. Elytra thorace paulo longiora, dense profundeque punctata, anterie versus suturam obsolete torosa. Abdomen crebre profundeque punctatum. Pedes flavi, coxis nigris, femoribus apice, tibiis basi nigris, tarsis longiusculis, articulis tribus primis linearibus, apice summo nigricantibus, primo elongato, quarto bilobo.

Mas abdominis segmento inferiore sexto profunde exciso, quarto apice medio sinuato et leviter impresso, sinu denticulo obtusiusculo utrinque terminato; femina segmento sexto apice rotundato, medio leviter acuminato.

\* 82. *St. flavicornis*: Plumbeo-niger, crebre fortius punctatus, albido-pubescent, antennis palpisque testaceis, pedibus flavis, femoribus posticis apice fuscis, abdomine subtilissime punctato. — Long.  $2\frac{1}{2}$  lin.

*Sten. geniculatus* Say Transact. Amer. Philoph. Soc. Philadelph. Nov. Ser. IV. 458. 1.

Habitat in Pensylvania, Dom. Zimmermann.

Statura et summa affinitas praecedentis. Plumbeo-niger, nitidulus, pube depressa albida cinerascens. Antennae longiores, articulo tertio quarto sesqui longiore, totae flavo-testaceae. Palpi flavo-testacei. Caput thorace sesqui latius, crebrius minus fortiter punctatum, fronte utrinque late leviter parum profunde longitudinaliter impressa, interstitio lato, perparum convexo, minus fere quam frontis latera elevato. Thorax basi coleopteris dimidio angustior, lateribus medio leviter rotundatus, latitudine summa vix longior, subcylindricus, dense fortius profunde punctatus, interstitiis vix rugulosis, pone medium utrinque obsolete impressus. Elytra thorace paulo longiora,



minus dense profunde fortiterque punctata, leviter convexa, antcrius versus suturam leviter torosa. Abdomen crebre subtilissime punctatum, cylindricum, segmentis quatuor primis basi leviter constrictis, summa basi paulo fortius punctatis. Pedes flavi, coxis piceis, femoribus posticis solis apice fuscis, tarsis brevioribus, articulis duobus primis linearibus, primo leviter elongato, tertio triangulari, quarto fortius bilobo.

Mas abdominis segmento inferiore sexto apice profunde triangulariter exciso; quarto apice vix impresso; femina segmento sexto apice rotundato.

Obs. Nomine *St. flavicornis* Dej. signatam hanc speciem in collectione Dom. Chevrolat vidi.

\* 83. *St. solutus: Plumbeo-niger, dense subtilius punctatus, albedo-pubescentis, antennis, palpis pedibusque flavis, femoribus anticis ante apicem geniculisque posterioribus nigris, abdomine subtilissime punctato. — Long. 2½ lin.*

Habitat Lutetiae, Dom. Dr. Aubé.

Statura et summa affinitas praecedentis. Plumbeo-niger, nitidulus, pube depressa albida cinerascens. Antennae longiusculae, articulo tertio quarto sesqui longiore, totae flavo-testaceae. Palpi flavo-testacei. Caput thorace sesqui prope latius, sat crebre minus fortiter punctatum, fronte depressiuscula, parum profunde bisulcata, interstitio lato, perparum convexo, medio laeviore. Thorax coleopteris paulo angustior, lateribus medio modice rotundatus, posterius paulo magis angustatus, basi coleopteris dimidio angustior, latitudine summa paulo longior, subcylindricus, sat dense subtilius profundus punctatus, interstitiis planiusculis, utrinque pone medium leviter oblique impressus. Elytra thorace paulo longiora, crebre minus fortiter profundius punctata, interstitiis subaequalibus, dorso antcrius longitudinaliter leviter toroso. Abdomen cylindricum, segmentis 4 anterioribus basi leviter constrictis, subtilissime crebre punctatum. Pedes flavo-testacei, coxis nigris, femoribus anticis ante apicem superne nigricantibus, posterioribus apice, tibiis posterioribus basi late nigris, tarsis brevioribus, articulis duobus primis linearibus, primo leviter elongato, tertio triangulari, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto apice profunde triangulariter exciso, quarto apice medio leviter sinuato, et intra sinum leviter exciso; femina segmento sexto apice leviter producto, subacuminato.

Obs. A *St. oculato* facile distinctus antennarum articulo primo concolore et abdomine subtilissime punctato, a *St. flavicorni* punctura thoracis elytrorumque subtiliore et densiore, tibiis posterioribus basi nigris etc.

\* 84. *St. cicindeloides: Niger, nitidus, grosse profunde aequaliter punctatus, parce pubescens, antennis palpisque testaceis, pedibus flavis, femoribus apice tibiisque basi late nigris, abdomine parcius punctato. — Long. 2½ lin.*

Grav. Micr. 155. 4. Mon. 229. 6. — Gyll. Ins. Suec. II. 470. 6. — Sahlb. Ins. Fenn. I. 424. 6. — Mannerh. Brachelytr. 42. 6. — Boisd. et Lucord. Faun. Ent. Paris. I. 444. 4. — Runde Brachelytr. Hal. 15. 6. — Er. Col. March. I. 570. 49. *Staph. buphthalmos* Rossi Faun. Etr. I. 252. 623. ed. Hellw. I. 311. 623 et not. p. 312. n. 3.

*Staph. clavicornis* Rossi Favn. Etr. Mant. I. 98. 22. ed. Helv. I. 443. 220.

*Paed. biguttatus* var. Oliv. Ent. III. 44. t. 1. f. 3. d.

*Sten. similis* var.  $\beta$  Ljungh Web. u. Mohr Archiv. I. 1. p. 66.

Habitat in Europa.

Niger, nitidulus, pube subtili depressiuscula albida rara vix cinerascens. Antennae mediocres, articulo tertio quarto paulo longiore, testaceae, clava fuscescens. Palpi flavo-testacei. Caput thorace parum latius, fortius crebriusque punctatum, fronte depressiuscula, late parum profunde bisulcata, interstitio leviter elevato. Thorax coleopteris dimidio angustior, lateribus medio leviter rotundatus, latitudine summa parum longior, subcylindricus, profunde fortiterque crebre punctatus, interstitiis subrugosis. Elytra thorace paulo longiora, grosse profundeque dense punctata, interstitiis rugulosis, leviter convexa, anterieus versus suturam longitudinaliter vix torosa. Abdomen crassiusculum, segmentis singulis basi crebrius fortiusque, dein parce subtiliusque punctatis. Pedes coxis nigris, femoribus flavo-testaceis, basi summa apiceque lato nigro-piceis, tibiis piceis, apice tarsisque testaceis, his brevioribus, articulis tribus primis linearibus, primo leviter elongato, quarto bilobo.

Mas abdominis segmento sexto apice acute triangulariter exciso; femina eodem acute rotundato.

Obs. Mas multo rarior obvius.

\* 85. *St. annularis*: Niger, nitidus, grosse profundeque punctatus, parce pubescens, antennis palpis pedibusque piceis, femoribus basi annulo flavo, abdomine parce punctato. — Long.  $2\frac{2}{3}$  lin.

Habitat in Pensylvānia, Dom. Zimmermann.

Statura et summa affinitas praecedentis, niger, nitidulus, pube depressa albida parca vix cinerascens. Antennae mediocres, articulo tertio quarto sesqui longiore, nigro-piceae, medio rufo-piceae. Palpi rufo-piceae, articulo tertio nigro-piceo. Caput thorace paulo latius, minus crebre fortius punctatum, fronte depressiuscula, late parum profunde bisulcata, interstitio lato, parum elevato, medio callo laevissimo instructo. Thorax coleopteris plus dimidio angustior, lateribus medio leviter rotundatus, posterius paulo magis angustatus, basi coleopteris duplo angustior, latitudine summa haud longior, subcylindricus, fortiter profundeque vage punctatus, interstitiis inaequalibus, dorso pone medium utrinque obsolete impressus, posterius rugis tribus longitudinalibus subelevatis laevissimis, callosis. Elytra thorace paulo longiora, grosse profundeque minus dense inaequaliter punctata, leviter convexa, versus suturam vix longitudinaliter torosa. Abdomen coleopteris paulo angustius, cylindricum, segmentis singulis parcius, basi fortius, apice subtilius punctatis. Pedes nigri, femoribus ante medium annulo lato flavo, tibiis apice tarsisque piceis, his brevioribus, articulis tribus primis linearibus, primo modice elongato, quarto fortiter bilobo.

Mas abdominis segmento sexto apice profunde exciso; femina eodem acute rotundato.

- \* 86. *St. operosus*: *Niger, nitidus, grosse profundeque inaequaliter punctatus, glaber, antennis, palpis pedibusque testaceis, callis frontalibus carinulaque media thoracis laevissimis, abdomine crebrius punctato.* — *Long.*  $2\frac{1}{2}$  lin.

Habitat in Brasilia, Sellow.

Statura fere *St. oculati*, niger, nitidus, glaber. Antennae mediocres, articulo tertio quarto sesqui longiore, testaceae, clava fusca. Palpi flavo-testacei. Caput thorace paulo latius, subtilius punctatum, fronte leviter late impressa, profundius bisulcata, interstitio lato, leviter elevato rugulisque duabus elevatis minoribus lateralibus prope oculos laevissimis, politis, facie infra antennis dense albo-pubescente. Thorax coleopteris plus dimidio angustior, lateribus modice rotundatus, posterius aequaliter atque anterieus angustatus, basi coleopteris duplo angustior, latitudine summa haud longior, subcylindricus, minus confertim inaequaliter fortiter profundeque punctatus, linea media longitudinali latiore, utrinque abbreviata, laevi, subelevata. Coleoptera thorace paulo longiora, fortiter profundeque minus dense inaequaliter punctata, modice convexa, anterieus versus suturam torosa. Abdomen coleopteris angustius, apicem versus sensim leviter attenuatum, crebrius basi fortius, apice subtilius punctatum. Pedes rufo-testacei, coxis anterioribus concoloribus, posticis piceis, femoribus apice fusciscentibus, tarsis brevioribus, articulis duobus primis linearibus, primo leviter elongato, tertio triangulari, quarto fortiter bilobo.

Mas latet. Femina abdominis segmento inferiore sexto apice leviter triangulariter producto.

87. *St. ingalis*: *Niger, nitidus, antennarum basi pedibusque flavis, his geniculis fuscis, capite, thorace elytrisque crebre profundeque punctatis, callis frontalibus laevissimis, abdomine parce subtiliter punctato.* — *Long.* 2 lin.

Habitat in Columbia, Dom. Moritz.

Statura fere praecedentis, antennis autem pedibusque magis elongatis. Corpus nigrum, nitidum, infra abdomine parce griseo-pubescente. Antennae elongatae, articulo tertio quarto paululum longiore, fuscae, basi flavo-testaceae. Palpi articulis duobus primis flavis (tertius deest). Caput thorace paulo latius, profunde punctatum, fronte bisulcata, interstitio leviter elevato, antrorsum angustato, callo minuto rotundato prope oculum utrinque et carinulis duabus supra antennarum basin laevissimis, politis, infra antennis dense albedo-pubescentis. Thorax coleopteris dimidio angustior, medio leviter rotundatus, latitudine summa paulo longior, subcylindricus, crebre sat profunde punctatus, interstitiis leviter rugulosis, praecipue versus latera. Elytra thorace paulo longiora, leviter convexa, medio longitudinaliter late at parum distincte torosa, dense fortiter profundeque punctata, interstitiis subinaequalibus. Abdomen coleopteris paulo angustius, apicem versus sensim leviter angustatum, segmentis singulis basi leviter constrictis, anterioribus subtilius punctatis, ultimis duobus laevibus. Pedes flavi, coxis anterioribus testaceis, femoribus apice, tibiis basi, tarsis medio fuscis, his longioribus, articulis tribus primis linearibus, primo modice elongato, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto apice truncato. Femina latet.



- \* 88. *St. callosus*: *Niger, nitidus, fortiter profundeque punctatus, parce pubescens, antennis, palpis pedibusque testaceis, femoribus apice nigris, fronte callis tribus laevissimis, abdomine parce punctato.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Pensylvania, Dom. Zimmermann.

Habitu omnino *St. cicindeloidis*, at quadruplo minor, niger, nitidus, pube brevi parca albida vix cinerascens. Antennae longiores, articulo tertio quarto sesqui longiore, testaceae, apice fuscrescentes. Palpi testacei. Caput thorace paulo latius, subtilius confertim punctatum, densius albidopubescens, fronte depressa, obsolete bisulcata, interstitio latiusculo, leviter elevato, callo utrinque prope oculum minore carinulisque duabus minutis supra antennarum basin laevissimis, politis. Thorax coleopteris duplo fere angustior, lateribus leviter rotundatus, latitudine paulo longior, subcylindricus, fortius profunde crebreque inaequaliter punctatus, carinula media longitudinali subelevata, utrinque abbreviata, laevi. Elytra thorace paulo longiora, fortiter profundeque minus dense punctata, leviter convexa, anterius versus basin torosa. Abdomen coleopteris angustius, apicem versus sensim leviterque angustatum, parce subtiliter punctatum, segmentis quatuor primis basi modice constrictis. Pedes testacei, femoribus summa apice coxisque nigro-piceis, tarsis brevioribus, articulis duobus primis linearibus, primo leviter elongato, tertio triangulari, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto apice emarginato, quinto late subsinuato; femina segmento sexto apice rotundato.

89. *St. frontalis*: *Niger, nitidus, fortiter profundeque punctatus, antennis, palpis pedibusque testaceis, fronte callis tribus elevatis laevissimis, abdomine parcius punctato.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Affinis praecedenti, niger, nitidus, parce subtiliterque albidopubescens. Antennae breviores, articulo tertio quartoque subaequalibus, testaceae, clava fuscrescente. Palpi testacei. Caput thorace parum latius, creberrime sat profunde punctatum, fronte depressa, leviter bisulcata, interstitio lato, subelevato, callo prope utrumque oculum minore carinulisque duabus supra antennarum basin laevissimis, politis. Thorax coleopteris duplo angustior, lateribus ante medium leviter rotundatis, latitudine summa paulo longior, cylindricus, lateribus crebre, dorso parcius fortiter profundeque punctatus, carinula media subelevata obsoleta laevi. Elytra thorace paulo longiora, leviter convexa, medio oblique leviter torosa, densius fortiter profundeque punctata, interstitiis subaequalibus. Abdomen coleopteris angustius, apicem versus sensim attenuatum, segmentis singulis medio leviter constrictis, parcius obsoletiusque punctatum. Pedes obscure testacei, coxis nigris, tarsis brevioribus, articulo primo leviter elongato, tertio triangulari, quarto bilobo.

Mas abdominis segmentis inferioribus longitudinaliter late impressis, impressione densius pubescente, quinto apice subsinuato, sexto profunde inciso. Femina latet.

- \* 90. *St. colubrinus*: *Elongatus, niger, nitidus, glaber, antennis, palpis pedibusque elongatis flavis, fronte excavata, laevissima, abdomine parce subtiliterque punctato.* — Long. 3 lin.

Habitat in Madagascar, Dom. Goudot.

Valde elongatus, tenuis, cylindricus, proxime sequentibus gracilior, niger, nitidus, glaber. Antennae elongatae, tenues, articulo tertio quarto paulo longiore, flavo-testaceae. Palpi elongati, flavo-testacei. Labrum margine testaceo. Caput thorace duplo latius, laevissimum, antice albido-pubescent, fronte inter oculos fortiter excavata. Oculi magni, valde prominuli. Thorax basi coleopteris vix duplo angustior, medio perparum incrassatus, latitudine plus sesqui longior, cylindricus, crebrius fortiusque subinaequaliter punctatus, linea media longitudinali abbreviata laevi. Elytra thoracis latitudine, parcius profunde punctata, humeris obtusis, vix prominulis. Abdomen elongatum, subtiliter parce punctatum, segmentis tribus primis basi modice constrictis. Pedes elongati, flavo-testacei, coxis rufo-testaceis.

Mas abdominis segmento inferiore sexto apice triangulariter emarginato, quinto, quarto tertioque late profundius canaliculatis, canalicula pubescente, utrinque ciliata. Femina latet.

\* 91. *St. attenuatus*: *Elongatus, niger, nitidus, glaber, antennis, palpis pedibusque testaceis, fronte modice excavata, parce punctata, abdomine crebrius punctato.* — Long.  $2\frac{1}{2}$  lin.

Habitat in Madagascar, Dom. Goudot.

Elongatus, gracilis, niger, nitidus, glaber. Antennae elongatae, tenues, articulo tertio quarto paulo longiore, testaceae. Palpi elongati, testacei. Labrum margine summo piceo. Caput thorace duplo latius, antice albido-pubescent, fronte medio excavata, obsolete subtiliterque bisulcata, interstitio lato, nullo modo elevato, vage parceque punctata. Thorax coleopteris duplo angustior, medio levissime rotundato-incrassatus, latitudine duplo fere longior, cylindricus, dorso crebre, lateribus creberrime punctatus, interstitiis lateribus rugulosis, dorso linea media longitudinali, utrinque abbreviata, minus distincta, maculisque nonnullis minutis inaequalibus sparsis subelevatis laevibus. Elytra thorace paulo longiora, parcius profunde punctata, interstitiis aequalibus, leviter convexa, anterius versus suturam longitudinaliter leviter torosa, humeris fere rectangulis, prominulis. Abdomen coleopteris dimidio angustius, elongatum, cylindricum, basi fortius, apice subtilius crebrius punctatum, segmentis quatuor primis basi leviter constrictis. Pedes elongati, tenues, rufescenti-testacei, coxis anticis concoloribus, tarsis longioribus, articulo primo elongato, tertio triangulari, apice emarginato, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto apice triangulariter exciso. Femina latet.

\* 82. *St. prolixus*: *Elongatus, niger, nitidus, glaber, antennis, palpis pedibusque flavis, fronte modice excavata, utrinque punctata, abdomine parce punctato.* — Long.  $2\frac{3}{4}$  lin.

Habitat in Madagascar, Dom. Goudot.

Statura elongata gracili omnino praecedentis, at paulo maior, niger, nitidus, glaber. Antennae elongatae, tenues, articulo tertio quarto sesqui longiore, flavae, clava testacea. Palpi flavi. Caput thorace plus dimidio latius, antice albido-pubescent, fronte leniter excavata, subtiliter bisulcata, interstitio lato, depresso, parcius subtiliusque punctata, linea media longitu-

dinali laevi. Thorax coleopteris duplo prope angustior, medio haud incrassatus, latitudine plus sesqui longior, cylindricus, dorso parcius, lateribus crebre punctatus, interstitiis lateribus subrugulosis, dorso planis, linea media longitudinali laevi. Elytra thorace paululum longiora, minus dense sat fortiter profundeque punctata, leviter convexa, antice medio levissime torosa, humeris fere rectangulis. Abdomen coleopteris dimidio angustius, elongatum, cylindricum, segmentis 4 primis basi leviter constrictis, parce subtiliterque, segmentis 4 primis lateribus paulo fortius constrictis, pedes graciles, pallide flavi, coxis dilute testaceis, tarsis longiusculis, articulo primo elongato, tertio triangulari, apice acute emarginato, quarto fortiter bilobo.

Mas latet. Femina abdominis segmento inferiore sexto apice rotundato.

Obs. Proximus praecedenti, at statura maiore, antennis, palpis pedibusque pallidioribus, capite angustiore, thorace medio haud crassiore, minus confertim punctatis, abdomine parcius punctato distinctus videtur.

\* 93. *St. gracilis: Elongatus, niger, nitidus, glaber, antennis, palpis pedibusque testaceis, fronte leviter excavata, utrinque punctata, abdomine laevissimo.* — Long.  $2\frac{2}{3}$  lin.

Habitat in Madagascar, Dom. Goudot.

Elongatus, gracilis, cylindricus, niger, nitidus, glaber. Antennae modice elongatae, tenues, articulo tertio quarto paulo longiore, testaceae. Palpi elongati, flavi. Labrum margine tenui testaceo. Caput thorace sesqui latius, antice subtiliter albido-pubescent, fronte modice excavata, utrinque crebre subtilius punctata, linea media longitudinali laevi. Thorax coleopteris duplo fere angustior, medio vix incrassatus, latitudine duplo longior, cylindricus, lateribus longitudinaliter late subimpressus, dorso parcius, lateribus dense punctatus. Elytra thorace parum longiora, parcius minus profunde punctata, leviter convexa, antice versus basin obsolete torosa, humeris fere rectangulis, prominulis. Abdomen coleopteris angustius, elongatum, cylindricum, laevissimum, segmentis quatuor primis basi leviter constrictis. Pedes modice elongati, testacei, coxis anticis concoloribus, tarsis longioribus, articulo primo sat elongato, tertio triangulari, quarto fortiter bilobo.

Mas abdominis segmento inferiore quarto quintoque medio longitudinaliter breviter subtiliterque cinereo-pubescentibus, sexto apice leviter emarginato. Femina latet.

\* 94. *St. lubricus: Elongatus, niger, nitidus, glaber, antennis, palpis pedibusque flavo-testaceis, fronte leviter excavata, crebre punctata, elytris thorace paulo longioribus, inaequalibus, abdomine crebrius punctato.* — Long.  $2\frac{1}{2}$  lin.

Habitat in Brasilia.

Proximus praecedentibus, at minus cylindricus, niger, nitidus, glaber. Antennae elongatae, tenues, articulo tertio quarto paulo longiore, flavae. Palpi elongati, pallide flavi. Caput thorace sesqui latius, antice densius albido-pubescent, fronte late levissime excavata, crebre fortius profunde punctata, breviter subtiliterque albido-pubescente. Thorax coleopteris plus dimidio angustior, medio haud incrassatus, latitudine duplo fere longior,



cylindricus, dense fortiter profundeque aequaliter punctatus, interstitiis subrugulosus. Elytra thorace paulo longiora, minus confertim sat fortiter profundeque punctata, dorso prope suturam longitudinaliter anguste torosa, toro antice humeros versus vergente, extra torum late subimpressa, sutura depressiuscula, humeris fere rectangulis, leviter prominulis. Abdomen coleopteris dimidio angustius, cylindricum, segmentis anterioribus 4 crebre minus subtiliter punctatis, basi leviter constrictis, reliquis duobus parce subtiliterque punctatis, subtus segmentis intermediis longitudinaliter subtiliter cinereopubescentibus. Pedes flavo-testacei, coxis omnibus piceis, tarsis longioribus, articulo primo elongato, tertio triangulari, quarto fortiter bilobo.

Mas latet. Femina abdominis segmento inferiore sexto apice rotundato.

Obs. Distinctus a praecedente corpore minus gracili, elytris paulo latioribus et longioribus, inaequalibus.

\* 95. *St. anguinus*: *Elongatus, cylindricus, niger, nitidus, parce albido-pubescentis, antennis, palpis pedibusque flavis, fronte leviter excavata, parcius punctata, elytris thoracis longitudine, abdomine segmentis quatuor anterioribus profundius punctatis, basi dense pubescentibus.* — Long. 2 lin.

Habitat in Guyana Venezuelensi, Dom. Moritz.

Statura et summa affinitate *St. elongati*, at paulo minor, niger, nitidus, pube brevissima albida parca vestitus. Antennae elongatae, articulo tertio quarto paulo longiore, albido-flavae, apice fuscae. Palpi elongati, pallide flavi. Labrum margine testaceo. Caput thorace sesqui latius, fronte late levissime excavata, parcius et minus profunde punctata, parcius subtiliterque albido-pubescente, infra antennis dense albido-villosum. Thorax coleopteris dimidio angustior, lateribus medio levissime rotundato-angustatus, latitudine duplo prope longior, cylindricus, aequaliter dense fortius profundeque punctatus, posterius linea media longitudinali tenui subelevata laevi. Elytra thoracis longitudine, leviter convexa, minus dense fortiter profundeque punctata, interstitiis aequalibus, humeris fere rectangulis, prominulis. Abdomen coleopteris dimidio angustius, elongatum, segmentis quatuor primis crebre fortius profunde punctatis, basi constrictis et densius pubescentibus, ultimis parce subtiliterque punctatis. Pedes graciliores, pallide flavi, tarsis articulis 3 primis linearibus, primo elongato, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto profunde triangulariter exciso, quinto fortius longitudinaliter impresso, impressione lateribus ciliis densis longioribus introrsum vergentibus terminata, apice sinuato. Femina latet.

Obs. A praecedentibus corpore pubescente, a sequentibus fronte leviter excavata et elytris brevioribus distinctus.

\* 96. *St. teres*: *Elongatus, cylindricus, niger, nitidus, glaber, antennis, palpis pedibusque flavis, fronte leviter excavata, crebrius punctata, abdominis segmentis quatuor anterioribus profundius parciusque punctatis, elytris thorace brevioribus.* — Long. 2½ lin.

Habitat in Brasilia, Dom. Melly.

Statura elongata, gracilis, cylindrica praecedentium. Niger, nitidus, gla-

ber. Antennae elongatae, tenues, articulo tertio quarto parum longiore, testaceae, basi flavae. Palpi elongati, pallide flavi. Labrum rufo-testaceum, basi nigrum. Caput thorace sesqui latius, antice albedo-pubescent, fronte modice excavata, crebre fortius subinaequaliter punctata, posterius carinula media longiore, anterieus carinulis duabus supra antennarum basin brevioribus, angustis, elevatis, laevibus. Thorax coleopteris paulo angustior, medio haud incrassatus, latitudine duplo fere longior, cylindricus, creberrime fortiter profundeque punctatus, interstitiis subrugulosis. Elytra thorace paulo breviora, fortiter profundeque densius punctata, interstitiis subinaequalibus, humeris obtusis, haud prominulis. Abdomen coleopteris parum angustius, elongatum, cylindricum, segmentis quatuor primis vage profundius punctatis, basi fortius constrictis, ultimis duobus parce subtilissime punctatis. Pedes modice elongati, flavi, coxis anticis testaceis, tarsis brevioribus, articulo primo leviter elongato, tertio obcordato, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto profunde triangulariter inciso, quinto longitudinaliter late impresso, apice sinuato, quarto tertioque longitudinaliter excavatis, excavatione margine ciliata. Femina latet.

Obs. Coleopteris thorace brevioribus et perparum latoribus ab affinis distinctus.

\* 97. *St. elongatus*: *Elongatus, cylindricus, niger, nitidus, glaber, antennis, palpis pedibusque flavis, fronte leviter excavata, crebre punctata, abdomine segmentis quatuor anterioribus profundius punctatis, elytris thoracis longitudine*. — Long.  $2\frac{1}{4}$  lin.

Habitat in Brasilia.

Statura praecedentium, niger, nitidus, glaber. Antennae elongatae, articulo tertio quartoque subaequalibus, pallide flavae. Palpi elongati, flavi. Labrum nigrum, margine exteriore flavescente. Caput thorace sesqui latius, antice albedo-pubescent, fronte late levissime excavata, crebrius profunde punctata. Thorax coleopteris duplo prope angustior, medio vix incrassatus, latitudine duplo fere longior, cylindricus, fortiter profunde dense aequaliter punctatus, interstitiis subaequalibus. Elytra thoracis longitudine, fortiter profundeque minus dense punctata, leviter convexa, aequalia, humeris fere rectangulis, prominentibus. Abdomen coleopteris dimidio fere angustius, elongatum, cylindricum, segmentis quatuor primis profundius punctatis, basi leviter constrictis, ultimis duobus parce subtilissime punctatis, subtus medio longitudinaliter pubescens. Pedes modice elongati, flavi, coxis anticis testaceis, posterioribus piceis, tarsis longioribus, articulo primo elongato, tertio triangulari, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto profunde inciso, quinto apice late profundius sinuato; femina segmento sexto apice acute rotundato.

\* 98. *St. iunceus*: *Elongatus, cylindricus, nigro-subvirescent, nitidulus, parce subtiliterque albedo-pubescent, antennis, palpis pedibusque pallide flavis, fronte depressa, elytris thorace haud longioribus, abdomine crebre fortius punctato*. — Long.  $2\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Habitu fere praecedentium, St. oculato fere aequalis, at gracilior, plum-

beo-niger, virescenti-nitidulus, pube brevi subtili albida aequali parcaque vix cinerascens. Antennae elongatae, tenues, articulo tertio quarto paulo brevior, pallide flavae, clava testacea. Palpi elongati, flavo-testacei. Labrum testaceum, summa basi nigricante. Caput thorace paulo latius, fronte depressa, minus dense sat profunde punctata, densius subtiliter albido-pubescente, infra antennis albo-villosum. Thorax coleopteris dimidio angustior, medio vix incrassatus, latitudine duplo longior, cylindricus, aequaliter dense profundeque punctatus, interstitiis parum rugulosi. Elytra thorace haud longiora, leviter convexa, minus confertim fortiter profundeque punctata, interstitiis aequalibus. Abdomen coleopteris dimidio angustius, modice elongatum, segmentis 4 primis basi leviter constrictis, fortiter profundiusque crebre punctatum, segmento sexto parce subtiliterque punctato. Pedes graciliores, flavi, coxis testaceis, tarsis articulo primo elongato, secundo tertioque triangularibus, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto profunde longitudinaliter inciso, quinto longitudinaliter, quarto apice triangulariter impresso, impressionibus densius pubescentibus; femina segmento sexto apice rotundato.

\* 99. *St. ariolus*: Plumbeo-niger, subnitidus, dense punctatus, densius albido-pubescentis, antennis, palpis pedibusque flavis, fronte depressa, elytris thorace paulo longioribus, his et abdominis profundius punctulati segmentis singulis basi densius pubescentibus. — Long.  $2\frac{1}{2}$  lin.

Habitat in Columbia, Dom. Moritz.

Statura fere *St. oculati*, et parum gracilior, plumbeo-niger, subnitidus. Antennae elongatae, tenues, articulo tertio quartoque subaequalibus, flavae, clava fusciscente. Palpi elongati, flavi. Labrum testaceum, basi nigricans. Caput thorace sesqui prope latius, subtilius parciusque punctatum, fronte depressa, anterie obsolete bisulcata, densius subtiliter breviterque albido-pubescentis, infra antennis dense albido-villosum. Thorax coleopteris duplo angustior, medio leviter rotundato-incrassatus, latitudine sesqui fere longior, subcylindricus, creberrime minus fortiter punctatus, interstitiis subrugulosis, densius breviter subtiliterque albido-pubescentis. Elytra thorace paululum longiora, fortiter profundeque dense punctata, leviter convexa, humeris obtusiusculis, prominentibus, basi dense, ceterum parce albido-pubescentia. Abdomen coleopteris dimidio angustius, modice elongatum, crebrius fortius profundiusque punctatum, segmentis quatuor primis basi fortius constrictis, parce albido-pubescentis, segmentis 5 primis basi pube dense depressa albida obductis. Pedes graciliores, albido-flavi, tarsis articulo primo elongato, tertio triangulari, apice emarginato, quarto fortiter bilobo.

Mas latet. Femina abdominis segmento inferiore sexto apice rotundato.

Obs. Differt a praecedente praecipue statura minus gracili, thorace densius, elytris et abdominis segmentis basi dense sericeo-pubescentibus.

\* 100. *St. paganus*: Plumbeo-niger, nitidulus, crebre profundeque punctatus, subtiliter albido-pubescentis, antennis palpisque testaceis, his apice, illis basi apiceque fuscis, pedibus rufo-brunneis, tarsis testaceis. — Long.  $1\frac{2}{3}$  lin.

Er. Col March. I. 571. 50.

Habitat in Germania, Anglia.



Habitu et statura omnino *St. unicoloris*, plumbeo-niger, nitidulus, pube brevi depressa albida parciore cinerascens. Antennae breves, articulo tertio quarto paulo longiore, testaceae, articulis duobus primis piceis, clava fusca. Palpi testacei, articulo secundo apice fusco, tertio fusco, basi testaceo. Caput breviusculum, transversum, thorace paulo latius, dense fortius profunde punctatum, fronte depressa, late leviter bisulcata, interstitio lato, parum convexo, aequae ac frontis latera elevato. Thorax coleopteris dimidio angustior, lateribus leviter rotundatus, posterius magis angustatus, basi coleopteris duplo angustior, latitudine summa vix longior, subdepressus, dense profundeque aequaliter punctatus, interstitiis latera versus subrugulosus. Coleoptera thorace paululum longiora, fortius dense profundeque punctata, interstitiis subrugulosus. Abdomen coleopteris perparum angustius, apice leviter attenuatum, dense fortius profundeque punctatum. Pedes breviusculi, rufo-brunnei, tarsis testaceis, brevibus, articulo primo vix elongato, secundo tertioque subtriangularibus, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto profunde inciso, quinto longitudinaliter late impresso, impressione carinula obtusiuscula terminata, apice trisinuato, quarto levius longitudinaliter impresso; femina segmento sexto apice rotundato.

\* 101. *St. latifrons*: Subdepressus, plumbeo-niger, subnitidus, crebre punctatus, subtiliter albido-pubescent, antennis medio palpisque piceis, his basi testaceis, pedibus rufo-piceis, abdomine fortius crebre punctato. — Long.  $1\frac{2}{3}$  lin.

*Er. Col. March. I.* 572. 51.

*Sten. morio* var. *Grav. Mon. p.* 231.

Habitat in Europa.

Praecedenti proximus, linearis, subdepressus, plumbeo-niger, subnitidus, pube brevi albida cinerascens. Antennae breviusculae, articulo tertio quarto paulo longiore, piceae, basi apiceque obscuriores. Palpi picei, articulo primo toto, secundo basi testaceo. Caput breve, thorace paulo latius, crebre profunde punctatum, fronte obsoletius late bisulcata, interstitio lato, parum convexo, aequae ac frontis latera elevato. Thorax coleopteris paulo angustior, lateribus leviter rotundatus, posterius paulo magis quam antierius angustatus, basi coleopteris dimidio angustior, latitudine summa vix longior, subdepressus, confertissime aequaliter profunde punctatus, interstitiis latera versus subrugulosus. Elytra thorace paulo longiora, dense profundeque punctata, interstitiis aequalibus, antierius versus suturam obsolete torosa. Abdomen elongatum, lineare, dense profundiusque punctatum. Pedes breviusculi, rufo-picei, coxis nigris, tarsis brevibus, articulo primo parum elongato, secundo tertioque triangularibus, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto profunde inciso, quinto longitudinaliter impresso, impressione apicem versus utrinque carinula obtusa terminato, apice sinuato, quarto apicem versus leviter impresso; femina segmento sexto apice acute rotundato.

Obs. Differt a praecedente corpore paululum magis depresso, punctura paulo minus forti, coleopteris thorace parum latioribus, palpis, antennis tarsisque dilutioribus. Simillimus quoque *St. unicolori*, sulcis frontalibus autem minus profundis et tarsorum articulo quarto bilobo facile distinguendus.

- \* 102. *St. punctatus*: *Plumbeo-niger, subnitidus, crebre punctatus, albido-pubescent, antennis palpisque testaceis, apice fuscis, pedibus obscure rufo-testaceis, abdomine fortius crebre punctato.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Habitu omnino *St. cicindeloidis*, at plus quadruplo minor, plumbeo-niger, subnitidus, pube brevi albida cinerascens. Antennae breviores, articulo tertio quarto sesqui longiore, obscure testaceae, clava fusciscente. Palpi testacei, basi pallidi, apice fusciscentes. Caput thorace paulo latius, crebre fortiter profundeque punctatum, fronte depressa, obsolete bisulcata. Thorax coleopteris duplo angustior, lateribus medio parum rotundatis, latitudine vix longior, cylindricus, creberrime profunde punctatus, interstitiis rugulosis. Elytra thorace paulo longiora, fortiter profundeque dense punctata, interstitiis rugulosis, leviter convexa, anterius versus suturam longitudinaliter leviter torosa. Abdomen coleopteris angustius, apicem versus sensim leviter attenuatum, crebre profundeque punctatum, segmentis quatuor primis basi leviter constrictis. Pedes obscure rufo-testacei, coxis posterioribus piceis, femoribus apice fusciscentibus, tarsis breviusculis, articulo primo leviter elongato, tertio triangulari, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto profunde triangulariter inciso; femina eodem apice rotundato.

Obs. In collectione Dom. Chevrolat hunc Stenum vidi nomine *St. punctati* Dej.

- \* 103. *St. arculus*: *Plumbeo-niger, nitidulus, dense fortiter punctatus, parce albido-pubescent, antennis, palpis pedibusque testaceis, pedibus rufo-piceis, tibiis basi testaceis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Habitu omnino *St. callosi*, at sesqui maior, plumbeo-niger, nitidulus, pube brevi depressa albida rariore vix cinerascens. Antennae articulis duobus primis testaceis (reliqui desunt). Palpi rufescenti-testacei, basi dilutiores. Caput thorace parum latius, crebre fortiter profundeque punctatum, fronte planiuscula, obsolete late bisulcata, interstitio parum convexo, vix magis quam capitis latera elevato, subcarinato. Thorax coleopteris duplo angustior, lateribus ante medium leviter rotundatus, latitudine summa vix longior, subcylindricus, dense fortiter profundeque punctatus, interstitiis leviter rugosis, dorso pone medium callo minuto ovali laevissimo. Elytra thorace paulo longiora, leviter convexa, dense fortiter profundeque punctata, interstitiis leviter rugosis. Abdomen coleopteris angustius, apicem versus sensim leviter attenuatum, segmentis 4 anterioribus basi constrictis, minus dense profundius punctatum. Pedes dilute rufo-picei, coxis, trochanteribus, femorum apice tibiisque obscurioribus, his basi determinate testaceis, tarsis breviusculis, articulo primo leviter elongato, tertio triangulari, quarto fortiter bilobo.

Mas latet. Femina abdominis segmento inferiore sexto apice rotundato.

- \* 104. *St. contractus*: *Niger, nitidus, fortiter punctatus, parce albido-pubescent, palpis nigris, basi testaceis, tibiis basi albidis, coleopteris amplis, convexis.* — Long. 1 lin.

Er. Col. March. I. 573. 52.

Habitat in Germania, Gallia, Italia.

Magnitudine *St. circulari* aequalis, niger, nitidus, pube brevi albida parcius vestitus. Antennae breviusculae, articulo tertio quarto parum longiore, piceae, apice nigrae. Palpi nigro-picei, articulo primo pallide testaceo. Caput parvulum, thorace parum latius, inaequaliter profundius punctatum, fronte obsolete bisulcata, interstitio parum convexo, non magis quam frontis latera elevato, laeviore. Thorax coleopteris plus duplo angustior, lateribus sat fortiter rotundatus, latitudine haud longior, convexiusculus, fortiter profundeque crebre punctatus, interstitiis rugulosis. Coleoptera ampla, capite plus duplo latiora, thorace sesqui longiora, convexiuscula, densius fortiter profundeque punctata, interstitiis subrugosis. Abdomen coleopteris multo angustius, apicem versus sensim attenuatum, breviusculum, parce profundius punctatum. Pedes nigri, tibiis basi albis, tarsis piceis, breviusculis, articulo primo leviter elongato, tertio triangulari, quarto fortiter bilobo.

Mas abdominis segmento inferiore sexto apice triangulariter inciso; femina eodem acute rotundato.

#### Species mihi invisae.

105. *St. rotundatus*: Nigro-aeneus, antennis longitudine capitis, elytris postice rotundatis, thorace globoso, abdomine subconico.

*Ljungh Web. u. Mohr Archiv. I. 1. 67. 8.*

Habitat in pratis Sjärsjöensibus (Sueciae).

Caput nigro-aeneum, breve, thorace latius, intra oculos porca tridente bis canaliculatum, punctatum. Oculi magni prominuli. Antennae nigrae, longitudine capitis, apice albo-pilosae, clavatae. Palpi nigri. Thorax aeneo-niger, convexus, immarginatus, orbiculatus, temere punctatus, capite et elytris angustior, capiteque vix duplo longior. Scutellum nullum. Elytra aeneo-nigra, thorace dimidio latiora, pauloque longiora, postice rotundata, punctata. Alae — — —. Abdomen apice attenuatum, vix marginatum, nigrum, nitidum, longitudine elytra vix superans. Subtus totus niger, tibiis tarsisque solis piceis. Statura *Steni buphthalmi*, sed vix magnitudine *Staphylini pygmaei* Payk. (*Ljungh*).

106. *St. 4-punctatus*: Niger, punctatus, thorace hexagonali, inaequali. — Long. ultra 1 lin.

*Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 459. 5.*

Habitat in Missuria.

Corpus nigerrimum, immaculatum, profunde dense punctatum. Os testaceum. Palpi nigri. Thorax hexagonalis, margine laterali medio angulato, disco foveolis quatuor dilatatis, quincunce dispositis, linea media longitudinali leviter elevata distinctis, foveola vel foveolis duobus obsoletis ad marginem lateralem impressus. Abdomen et pectus impunctata, illo breviter piloso. Pedes picei. (*Say*).

Obs. An rite huic generi adscriptus?



**Euaesthetus Grav.**

Grav. Mon. p. 201.

*Eristhetus Mannerh. Brachelytr. p. 41. — Boisd. et Lacord. Faun. Ent. Paris. I. p. 438.*

*Mentum transversum, apice sinuato-emarginatum.*

*Tarsi 4-articulati.*

Corpus minutum, oblongum, teretiusculum, alatum. Caput sat magnum, thoracis fere latitudine, basi leviter constrictum, thoraci immissum, exsertum, oculis parvis, modice prominulis. Labrum latum, transversum, apice dense acuteque denticulatum seu acute serratum. Mandibulae sat longae, falcatae, acutae, medio dente valido acuto armatae, pone dentem subtiliter denticulatae. Maxillae (t. V. f. 3. b.) malis corneis, interiore interne, exteriori apice membraneo barbatis. Palpi maxillares elongati, articulis tribus primis longitudine increscentibus, duobus primis teretibus, tertio incrassato, clavato, quarto minimo, subtilissimo, subulato, nonnisi oculo fortiter armato conspicuo. Labium (t. V. f. 3. a.) mento transversum, apice late emarginato, angulis anterioribus acute prominulis, spatio membranaceo inter mentum et ligulam brevi, transversum, ligula lata, medio leviter sinuata, membranacea, paraglossis ligulae connatis, eamque superantibus, acuminatis, membranaceis, subtiliter ciliatis. Palpi labiales breves, articulo primo minore, subovato, secundo crassiusculo, subovato, basi breviter pedunculato, tertio minuto, subtili, subulato. Antennae in frontis margine anteriore intra mandibularum basin insertae, basi distantes, breves, articulis duobus primis sequentibus paulo crassioribus maioribusque, articulis 3—5 subaequalibus, 6—8 sensim paulo brevioribus, ultimis tribus sensim maioribus, clavulam oblongam formantibus. Thorax lateribus anterioribus rotundatus, posterioribus angustatus, basi apiceque truncatus. Scutellum minimum, triangulare. Coleoptera apice late conjunctim emarginata. Abdomen plerumque marginatum, apice obtusum. Pedes breviusculi, tarsis 4-articulatis, articulo primo ultimoque intermediis longioribus.

**Observationes.**

1. *Euaesthetus* cum *Steno* summa affinitate conjunctus, licet aliquomodo *Oxyteli* habitum simulans, genus constituit tarsis 4-articulatis singulare et omnino optime distinctum, unde nomen (ab *eu* et *αἰσθητικός*), quare emendatio Mannerheimii, qui illud errore quodam pro *Eristheto* exortum putat, mihi haud suscipienda videtur.

2. Mas abdominis segmento inferiore sexto apice inciso, praecedentibus impressis tuberculativis distinguendus.

3. Victus in silvis sub foliis deciduis et in pratis.

\* 1. *E. scaber*: *Nigro-piceus, opacus, capite pedibusque rufis, capite, thorace elytrisque confertissime punctatis, thorace lineolis duabus rectis impresso. — Long. 1 lin.*

Grav. Mon. 202. 1. — Gyll. Ins. Suec. II. 462. 1. — Germ. Faun. Ins. Europ. VII. 13. — Er. Col. March. I. 574. 1.

*Eristhetus scaber* Mannerh. Brachelytr. 41. 1. — Sahlb. Ins. Fenn. I. 420. 1. — Boisd. et Lacord. Faun. Ent. Paris. I. 439. 1.

*Stenus bipunctatus* Ljungh Web. et Mohr Arch. I. 1. 68. 9.

Habitat in Germania, Suecia.

Nigro-piceus, opacus, densius breviter subtilissime fusco-pubescent. Caput cum antennis palpisque saturate rufum, confertissime punctatum. Thorax coleopteris parum angustior, lateribus anterior rotundatus, posterius angustatus, basi coleopteris duplo angustior, perparum convexus, confertissime punctatus, dorso posterior lineolis duabus, ante medium abbreviatis, distantibus, parallelis, rectis impressus. Elytra thorace paulo breviora, depressa, sat crebre punctata. Abdomen subtilissime punctulatum. Pedes rufi.

Mas abdominis segmento inferiore sexto apice profunde exciso, quinto aequali, quarto prope apicem tuberculis duobus minutis compressis approximatis instructo; femina segmento sexto apice rotundato.

\* 2. *E. ruficapillus*: Piceus, nitidulus, capite pedibusque rufis, capite thoraceque confertim punctatis, hoc lineolis duabus arcuatis impresso, elytris subtilissime punctulatis. — Long.  $\frac{3}{4}$  lin.

Er. Col. March. I. 575. 2.

*Eristhetus ruficapillus* Boisd. et Lacord. Faun. Ent. Paris. I. 439. 2.

Habitat in Germania, Gallia.

Habitu et summa similitudine praecedentis, at duplo fere minor, piceus, nitidulus, densius breviter subtilissime fusco-pubescent. Caput cum antennis palpisque rufum, confertim punctatum. Thorax coleopteris paulo angustior, lateribus anterior fortiter rotundatus, posterius angustatus, basi coleopteris duplo angustior, parum convexus, confertim punctatus, dorso posterior lineolis duabus leviter arcuatis, (introrsum convexis,) ante medium abbreviatis, modice distantibus, impressus. Elytra thoracis longitudine, subdepressa, parcius subtilissime punctata, prope suturam subtilissime unistriata. Abdomen subtilissime punctulatum, segmentis singulis margine posteriore rufescente. Pedes rufi.

Mas abdominis segmento inferiore sexto apice exciso, quinto medio longitudinaliter canaliculato, quarto prope apicem tuberculis duobus minutis compressis, approximatis instructo; femina segmento sexto apice rotundato.

Obs. Differt a praecedente statura minore, colore dilutiore, minus opaco, elytris subtilissime punctatis, thoracis foveolis subarcuatis, minus distantibus. Primus hanc speciem nomine distinxit Dom. v. Winthem.

\* 3. *E. Americanus*: Rufo-piceus, nitidulus, capite pedibusque rufis, capite thoraceque aequali confertim, elytris subtilissime punctatis. — Long.  $\frac{3}{4}$  lin.

Habitat in Pensylvania, Dom. Zimmermann.

Magnitudine praecedentis, at crassior, rufo-piceus, nitidulus, omnium subtilissime brevissimeque parcius fusco-pubescent. Caput confertim punctatum, rufum, antennis palpisque rufo-testaceis. Thorax coleopteris dimidio angustior, lateribus anterior modice rotundatus, posterius angustatus, basi coleopteris plus duplo angustior, convexus, confertim punctatus, aequalis. Elytra thoracis longitudine, convexiuscula, subtilissime punctulata, prope suturam distincte unistriata. Abdomen vix subtilissime punctulatum. Pectus piceum. Pedes rufo-testacei.

Mas abdominis segmento sexto profunde inciso, quinto apice late emarginato; femina segmento sexto apice rotundato.

4. *E. immarginatus*: *Rufo-testaceus, confertissime punctatus, abdomine immarginato.* — Long. 2 lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Rufo-testaceus, parum nitidus, subtilissime griseo-pubescens. Antennae tenues, elongatae, capite sesqui longiores, pallide flavae, clava parum incrassata, fusciscente. Palpi pallide flavi. Caput thorace paulo latius, confertim subtiliter punctatum. Thorax coleopteris dimidio angustior, elongatus, latitudine longior, ante medium leviter rotundatus, posterius leviter angustatus, basi coleopteris duplo prope angustior, subcylindricus, confertissime punctatus, aequalis. Elytra thorace paulo breviora, dense punctata. Abdomen immarginatum, dense subtiliter punctatum. Pedes flavo-testacei.

Mas latet. Femina abdominis segmento inferiore sexto acute rotundato.



## Tribus septima.

### **O x y t e l i n i.**

*Stigmata prothoracica occulta.*

*Coxae posticae transversae, anticae conicae, prominentes.*

*Trochanteres postici simplices.*

---

Labrum transversum, corneum, apice utrinque lacinia membranacea auctum plerisque;

vel apice limbo membranaceo instructum Bledio, Syntomio;  
bispinum Megalopi.

Mandibulae validiores, saepius prominentes,  
dentatae Genuinis;  
muticae Megalopi, Osorinis, Coprophilinis.

Maxillae mala interiore interne, exteriore apice barbatis, hac haud dilatata,

corneis Megalopi, Osorinis;  
membranaceis Genuinis, Coprophilinis.

Palpi maxillares filiformes Megalopi, Osorinis, Coprophilinis;  
articulo ultimo subulato Genuinis.

Mentum corneum,  
oblongum Megalopi;  
latitudine haud brevius Osorinis;  
transversum Genuinis, Coprophilinis.

Ligula cornea Megalopi, Osorinis;  
membranacea Genuinis, Coprophilinis.

Paraglossae distinctae, membranaceae, plerisque,  
liberae Megalopi, Osorinis;  
ligulae connatae Genuinis plerisque, Coprophilo;  
nullae Bledio.

Palpi labiales breves, filiformes.

Antennae sub frontis margine laterali insertae,

11-articulatae plerisque;

magis minusve refractae Osorio, Genuinis, Acrognatho;

10-articulatae, clavatae Megalopi.

Caput porrectum, thoraci immissum, basi parum constrictum plerisque; exsertum, thoraci pedunculo tenui affixum Apocello.

Thorax ab elytris magis minusve remotus, spatio pone coxas antic as absque integumento corneo.

Scutellum distinctum Megalopi, Osorinis, Coprophilinis et Trogophloeo \*;

indistinctum Genuinis plerisque.

Elytra pectoris longitudine.

Abdomen segmentis septem distinctis, septimo minuto at haud retractili, marginatum plerisque;

immarginatum Osorinis.

Pedes coxis anticis conicis, exsertis, posticis transversis, trochanteribus posticis simplicibus, tibiis

omnibus muticis Megalopi, Holotrocho, Trogophloeo, Apocello, Deleasteri, Micralymmati, Syntomio;

anticis vel anterioribus spinulosis Platystheto, Oxytelo, Phloeonaeo, Coprophilo;

anticis muricatis Osorio, Bledio, Acrognatho;

tarsis 5-articulatis Megalopi, Osorinis, Coprophilinis;

3-articulatis Genuinis.

#### O b s e r v a t i o.

Distinguitur haec tribus a Piestinis et Proteininis coxis anticis exsertis, conicis, spatio pone eas absque integumento corneo, a Phloeocharinis et Omalinis trochanteribus posticis haud fulcrantibus. Praeterea abdominis segmento septimo distincto haud retractili insignis.

### S u b t r i b u s   p r i m a.

## M e g a l o p i n i.

*Antennae 10-articulatae, clavatae.*

*Ligula cornea.*

*Tarsi 5-articulati.*

Labrum bispinum.

Mandibulae breves, exsertae.

Maxillae malis angustis, leviter elongatis, corneis.

Palpi maxillares filiformes.

Mentum oblongum, corneum.

Ligula cornea.

Paraglossae membranaceae.

Palpi labiales articulo ultimo subovato.

Antennae breves, 10-articulatae, rectae, apice distincte clavatae.

Oculi magni, prominentes.

Thorax subcylindricus, basi truncatus, ab elytris parum remotus.

Scutellum distinctum.

Abdomen marginatum.

Pedes coxis anticis minutis, intermediis approximatis, tibiis muticis, tarsis 5-articulatis.

## Megalops Dej.

Corpus oblongum, crassiusculum, alatum. Caput sat magnum, thorace latius, porrectum, oculis lateralibus, permagnis, convexis. Labrum occultum, at nonnunquam labri loco spinæ duæ acutæ corneæ prominent. Mandibulae exsertæ, falcatae, decussatae, medio denticulo minuto armatae. Maxillae (t. II. f. 3. b.) malis corneis, interiore brevior, interne membranea ciliataque, apice unco corneo armata, exterior longior, apice barbatula. Palpi maxillares mala exterior paulo longiores, articulo primo parvo, tertio secundo paulo brevior, quarto hoc sesqui longior, apice acuminato. Labium (t. II. f. 3. a.) mento latitudine prope longior, apicem versus leviter angustato, apice truncato, ligula cornea, apice obtuse bidentata, paraglossis membranaceis, ligulam parum superantibus. Palpi labiales articulo primo leviter elongato, secundo hoc paulo brevior, tertio oblongo-ovato, apice subacuminato. Antennae breves, 10-articulatae, articulo primo crassiusculo, secundo elongato, ultimo maximo, cum praecedentibus duobus clavam sat distinctam, oblongo-ovalem, pubescentem formantibus. Thorax coleopteris angustior, subcylindricus. Scutellum sat magnum, semiorbiculare. Elytra pectus haud superantia, truncata, angulo exterior rotundato. Abdomen parallelum, marginatum. Pedes graciliores, minus elongati, coxis intermediis levi spatio distantibus, tibiis omnibus simplicibus, tarsis articulis 4 primis sensim decrescentibus, quinto primo paulo longior.

### Observationes.

1. Nescio qua similitudine deceptus cel. Gravenhorst speciem huius generis sibi cognitam inter *Oxyporos* collocaverit, quocum neque forma, neque insertione antennarum, neque palporum structura singulari convenit. Habitus potius quodammodo *Steno* similis, praecipue oculis magnis, protuberantibus, antennis breviusculis, tenuibus, clavatis, et coxis anticis minutis: antennarum autem insertionem et oris structuram longe distat. Maxillarum et labri structura proxime ad *Osorinos* accedit.

2. Labri structura in hoc genere singularis videtur, mihi autem nondam satis perspecta. Labrum reliquorum plerorumque insectorum merum non vidi, loco eius spinam utrinque acutam prominere in nonnullis individuis observo.

3. Sexus differentia latet.

4. De victu nihil compertum. Species omnes hucusque cognitae Americanae.

\* 1. *M. caelatus*: *Niger, nitidus, cicatricosus, elytris pedibusque testaceis, nigro-variegatis*. — Long.  $2\frac{1}{2}$  lin.

*Oxyporus caelatus* Grav. Micr. 197. 5. Mon. 235. 4. — Latr. Hist. nat. Crust. et Ins. IX. 359. 7. — Oliv. Encycl. VIII. 608. 8.

Habitat in America septentrionali, Knoch.

*Niger, nitidus*. Antennae capite breviores, rufo-testaceae, articulo ultimo maximo, fusco-piceo. Mandibulae rufo-piceae. Caput inaequale, fortiter profundeque punctatum. Thorax coleopteris paulo angustior, latitudine paulo brevior, basin versus leviter angustatus, lateribus leviter rotundatus, margine laterali pluries denticulato, convexus, dense fortiter profundeque punctatus, callis irregularibus elevatis laevigatis inaequalis. Elytra inaequaliter fortiter profundeque punctata, callis elevatis laevigatis inaequalia, secundum suturam impressa, rufescenti-testacea, nitida, macula magna anguli



apicis exterioris, maculisque nonnullis sparsis nigris. Abdomen laeve, segmentis singulis basi utrinque bifoveolatis, anterioribus quatuor singulis margine laterali macula testacea notatis. Corpus subtus capite thorace abdomineque punctatis, pectore laevisimo. Pedes testacei, femoribus ante apicem piceis.

- \* 2. *M. punctatus*: Niger, nitidus, capite thoraceque utrinque transversim plicato parcius punctatis, elytris disco punctis nonnullis approximatis impressis, maculis duabus magnis testaceis. — Long.  $1\frac{1}{2}$  lin.

Var. *Elytris nigris, macula minima apicali anteriore testacea.*

Habitat in Columbia, Dom. Moritz.

Praecedente plus duplo minor, niger, nitidus. Antennae vix capitis longitudine, rufo-testaceae, clava magna, ovata, picea. Palpi flavi. Mandibulae rufo-piceae. Caput thorace sesqui fere latius, parcius grosse inaequaliter punctatum, fronte utrinque prope oculum longitudinaliter, antice transversim leviter impressa. Thorax coleopteris dimidio angustior, latitudine haud brevior, basin versus leviter angustatus, lateribus ante medium modice rotundatus, margine laterali integro, subcylindricus, plicis tribus transversalibus, fundo punctatis, medio late interruptis impressus, interstitiis laevissimis. Elytra in disco anteriore punctis nonnullis sat profundis fortibusque approximatis biseriatis notata, ceterum laevissima, secundum striam suturalem impressa, nigra, nitida, in altero specimine macula magna subquadrata infra humerum, aliaque ad angulum apicalem anteriorem, in altero punctulo minuto ad suturae apicem testaceo signata. Abdomen laeve. Pedes testacei.

- \* 3. *M. cephalotes*: Niger, nitidus, pedibus testaceis, thorace utrinque transversim plicato punctatoque, elytris disco anteriore striola impressa. — Long.  $1\frac{1}{2}$  lin.

T. 2. f. 3.

Habitat in Columbia, Dom. Moritz.

Statura omnino praecedentis, niger, nitidus. Antennae capitis fere longitudine, rufo-piceae, basi testaceae, apice fuscae. Mandibulae rufo-piceae. Caput sat magnum, thorace plus sesqui latius, fronte punctis grossiusculis inaequaliter notata, prope oculum utrumque longitudinaliter, antice transversim leviter impressa. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus ante medium subangulato-rotundatis, basin versus non magis quam apicem versus angustatus, basi coleopteris dimidio angustior, transversim convexus, plicis 4 transversalibus fundo punctatis, medio interruptis impressus. Elytra stria suturali et in disco anteriore striola utrinque abbreviata longitudinali impressa, ceterum laevissima. Abdomen laeve. Pedes testacei.

## Subtribus secunda.

### O s o r i n i.

*Antennae 11-articulatae.*

*Ligula cornea.*

*Tarsi 5-articulati.*

Labrum transversum, apice utrinque lacinia parva membranacea instructum.

Mandibulae validae.

Maxillae malis corneis, angustis, subelongatis.

Palpi maxillares filiformes, articulo primo minuto.

Mentum corneum,

apice emarginatum Osorio;

apice rotundatum Holotrocho.

Ligula cornea.

Paraglossae membranaceae.

Palpi labiales breves, filiformes.

Antennae 11-articulatae, filiformes aut apicem versus sensim subincrassatae.

Caput magnum, crassum, basi haud constrictum.

Oculi parvi, haud prominentes.

Thorax basi truncatus, ab elytris minus remotus.

Scutellum distinctum.

Abdomen immarginatum.

Pedes coxis intermediis approximatis, tibiis

anticis spinuloso-muricatis Osorio;

omnibus muticis Holotrocho;

tarsis 5-articulatis.

### Osorius Leach.

*Latr. Regn. Anim. IV. p. 438. Nov. Annal. Mus. Hist. Nat. I. p. 83.*

*Molosoma Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. p. 462.*

*Tibiae anteriores spinulosae.*

Corpus elongatum, cylindricum, alatum. Caput sat magnum, crassum, oculis parvis, rotundatis, haud prominulis. Labrum transversum, apice emarginatum, laciniiis membraneis, elongato-triangularibus, interne fortiter ciliatis. Mandibulae validae, acutae, decussatae. Maxillae (t. V. f. 13. b.) malis leviter elongatis, corneis, interiore interne ciliata, exteriori apice barbata, utraque unco corneo introrsum verso armata. Palpi maxillares malam exteriorem parum excedentes, articulo primo minimo, secundo elongato, tertio hoc plus duplo brevior, quarto secundo subaequali, apice acuminato. Labium (t. V. f. 13. a.) mento latitudine sua haud brevior, apicem versus sensim angustato, apice emarginato, ligula cornea, integra, apice subacuminata, paraglossis linearibus, ligulam haud excedentibus, membranaceis, interne longe ciliatis. Palpi labiales articulo primo brevior crassiorque, reliquis duobus longitudine aequalibus, tertio apice subacuminato. Antennae brevior.

res, apicem versus haud incrassatae, fractae, articulo primo elongato, sub capitis margine laterali recondito, tertio subobconico, parum elongato, ultimo ovato, subacuminato. Thorax basin versus sensim angustatus, basi coleopteris angustior, lateribus marginatus, basi truncatus, subtus antice tuberculo sat prominulo instructus. Scutellum apice rotundatum. Elytra lateribus tenuiter marginata, apice truncata. Abdomen immarginatum, cylindricum, segmento sexto conico, apice truncato, segmento septimo parvo, conspicuo. Pedes breviusculi, intermedii basi approximati, tibiis margine exteriori angulato, supra angulum spinulis, infra angulum dentibus sat longis spiniformibus serrato, anterioribus facie posteriore muricatis, tarsis gracilibus, articulis 4 primis subaequalibus, quinto his coniunctis longitudine aequali.

#### O b s e r v a t i o n e s.

1. Corpus parce pubescens, capite thoraceque glabris.
2. Sexus differentia in plerisque latet.
3. Victus sub arborum putrescentium cortice, in cuniculis longitudinaliter erosis. Nympham quoque in simili cuniculo reperit Dom. Lacordaire (Nov. Annal. Mus. Hist. nat. II. p. 65.).

X \* 1. *O. ater: Niger, nitidus, thorace parce subtilissime punctato, capite subtiliter punctulato, apice bispinoso. — Long. 5½ — 6½ lin.*

*Perty Delect. An. Art. p. 30. t. 7. f. 1.*

Habitat in Brasilia.

Corpus nigerrimum, nitidum. Antennae capite parum longiores, piceae. Palpi picei. Mandibulae muticae. Caput magnum, subtiliter punctulatum, lateribus supra oculos subtilissime longitudinaliter rugulosum, apice utrinque in spinam sat validam, licet parum elongatam, porrectam productum. Thorax latitudine antica brevior, basin versus sensim angustatus, basi coleopteris paulo angustior, lateribus anterie tenuiter, posterius fortius reflexo-marginatus, angulis posterioribus subprominulis, transversim convexus, parce omnium subtilissime punctulatus. Elytra thorace paulo longiora, subtilissime confertissime longitudinaliter ruguloso-strigosa, stria suturali impressa. Abdomen supra et lateribus laeve, subtus et segmento septimo parcius punctatum. Pedes picei.

\* 2. *O. intermedius: Niger, nitidus, thorace omnium subtilissime punctulato, capite subtiliter punctulato, apice emarginato. — Long. 6 lin.*

Habitat in Columbia interiore, Dom. Buquet.

Statura et summa affinitas praecedentis. Niger, nitidus. Antennae capite paulo longiores, nigro-piceae. Palpi rufo-picei. Mandibulae muticae. Caput sat magnum, thoracis latitudine, posterius laeve, anterie subtiliter punctulatum, lateribus supra oculos nonnihil longitudinaliter rugulosum, apice late emarginatum, angulis anterioribus leviter prominulis. Thorax latitudine antica paulo brevior, basin versus a medio inde sensim angustatus, basi coleopteris paulo angustior, apice usdem paulo latior, lateribus anterie subtiliter, posterius fortius reflexo-marginatis, angulis posterioribus subrectis, haud prominulis, transversim convexus, laevis, vel omnium subtilissime obsolete punctatus. Elytra thorace paulo longiora, laevia, stria suturali



impressa. Abdomen laeve, segmento sexto infero punctulato, septimo laevi. Pedes nigri, tarsis fulvo-piceis.

Obs. Praecedenti simillimus, differt autem capitis angulis anticis haud spinosis, thorace paulo longiore, angulis posterioribus haud prominulis, elytris laevibus.

\* 3. *O. Brasiliensis*: *Niger, nitidus, thorace capiteque sat crebre distinctius punctatis, hoc apice emarginato.* — Long. 5 lin.

Guér. Iconogr. Ins. t. 9. f. 11. — Latr. Nov. Annal. Mus. Hist. nat. I. 86. 2.

Habitat Bahiae in Brasilia, Dom. Sellow.

Statura et summa affinitas praecedentium. Niger, nitidus. Antennae capite vix longiores, piceae. Palpi rufo-picei. Mandibulae muticae. Caput magnum, subtilius at sat distincte crebre punctatum, ante basin laevius, utrinque prope oculos longitudinaliter rugulosum, apice late emarginatum, angulis anterioribus prominentibus. Thorax latitudine antica tertia parte brevior, basin versus sensim angustatus, basi coleopteris paulo angustior, lateribus anterieus subtiliter, posterius fortius reflexo-marginatis, angulis posterioribus subrectis, haud prominulis, transversim convexus, crebre subtiliter, licet sat distincte punctatus. Elytra thorace paulo longiora, sparsim subtiliter punctata, subtiliter longitudinaliter rugulosa. Abdomen laeve, lateribus subtilissime longitudinaliter rugulosa, segmento ultimo inferiore crebre fortiterque punctato-ruguloso, segmento septimo laevi. Pedes nigri, tarsis fulvis.

Obs. Praecedentibus simillimus, differt autem ab *O. atro* capite mutico, a praecedente thorace brevior, ab utroque capite thoraceque distincte punctatis.

\* 4. *O. piceus*: *Niger, nitidus, antennis pedibusque rufis, thorace longiore capiteque parce subtilissime punctatis, hoc apice emarginato.* — Long. 3½ lin.

Habitat in Brasilia, Dom. Westermann.

Habitu omnino praecedentium, at multo minor, nigro-piceus, nitidus. Antennae capite parum longiores, rufae. Palpi rufo-testacei. Mandibulae muticae. Caput thoracis apice paululum angustius, subtiliter obsoleteque parce punctatum, ante basin transversim laeve, circa oculos vix rugulosum, apice emarginatum, angulis anterioribus modice prominentibus. Thorax latitudine antica perparum brevior, basin versus sensim leviter angustatus, basi coleopteris paululum angustior, lateribus medio levissime rotundatis, anterieus leviter, posterius fortius reflexo-marginatis, angulis posterioribus rectis, anterioribus acutis, leviter productis, modice transversim convexus, subtiliter obsoleteque parce punctatus, infra tuberculo iugulari rufo. Elytra thorace paulo longiora, laevia, stria suturali tenui impressa, brunneo-picea. Abdomen supra laeve glabrumque, infra parce punctatum et parce lutescenti-pilosum, supra nigro-piceum, apice et infra piceo-rufum. Pedes rufi.

\* 5. *O. incisicrurus*: *Niger, nitidus, thorace parce punctulato, longitudinaliter bi-impresso, capite maris apice tridentato, feminae fortiter emarginato, tibiis anterioribus profunde excisis.* — Long. 6 lin.

Latr. Nov. Annal. Mus. Hist. nat. I. 85. 1. — Klug Col. Madag. 51. 43. t. 1. f. 13.

Habitat in Madagascar, Dom. Goudot.

Statura omnino *O. atri*, nigerrimus, nitidus. Antennae capite paulo longiores, nigrae. Palpi picei. Mandibulae ante apicem dentatae. Labrum emarginatum. Caput thoracis latitudine, maris parce, feminae crebrius punctatum, postice laeve, lateribus prope oculos longitudinaliter rugulosum, fronte praecipue apicem versus depressiuscula, apice in mare tridentata, dente intermedio longiore, in femina medio triangulariter erosa. Thorax latitudine antica brevior, basi coarctatus, lateribus subtiliter reflexo-marginatis, angulis posterioribus rectis, leviter prominulis, disperse subtilius punctatus, dorso sulculis duobus longitudinalibus, lateribus pone medium secundum marginem lateralem impressus. Elytra thorace paulo longiora, vage obsolete punctata, stria suturali impressa. Abdomen laeve, segmento sexto inferiore maris parce punctato. Pedes nigri, tarsis rufo-piceis, tibiis minus dilatatis, anterioribus extus prope apicem profunde excisis, postice haud muricatis.

- \* 6. *O. rugifrons*: *Nigro-piceus, nitidus, punctatus, parce fulvo-pubescent, fronte dense longitudinaliter rugosa.* — Long.  $4\frac{2}{3}$  lin.

Habitat in Java, Dom. Dr. De Haen.

Habitu praecedentium at minor, nigro-piceus, nitidus, capite thoraceque parcius, elytris abdomineque densius fulvo-pubescentibus. Antennae capite paulo longiores, piceae. Palpi rufo-picei. Caput thoracis latitudine, postice laeve, nitidum, spatio transversali inter oculos dense longitudinaliter strigoso, opaco, antice subtiliter longitudinaliter rugulosum, latera versus disperse exasperato-punctatum, apice emarginatum, angulis anterioribus haud prominentibus. Thorax latitudine antica brevior, basin versus sensim angustatus, basi coleopteris parum angustior, lateribus medio levissime rotundatis, posterius leviter reflexo-marginatis, angulis posterioribus obtusiusculis, transversim convexus, crebrius subtiliusque punctatus, linea media longitudinali laevi. Elytra thorace paulo longiora, sat crebre minus profunde punctata, stria suturali impressa. Abdomen apicem versus sensim modice incrassatum, crebre punctatum, segmentis singulis margine postico tenui rufo-piceo. Pedes rufo-picei.

- \* 7. *O. cylindricus*: *Niger, nitidus, elytris, antennis pedibusque piceis, capite, thorace elytrisque parce, abdomine parcius punctato.* — Long.  $3\frac{1}{2}$  lin.

*Latr. Nov. Annal. Mus. Hist. nat. I. 86. 3.*

Habitat in Mexico (Orizaba), Dom. Deppe.

Corpus elongatum, nigrum, nitidum. Antennae capite vix longiores, ab articulo septimo inde paulo crassiores, piceae, basi apiceque magis rufescentes. Palpi testacei. Labrum rufo-piceum. Caput thoracis latitudine, lateribus passim exasperato-punctatum, fronte inter oculos utrinque transversim longitudinaliter rugulosa, apice emarginata, angulis anterioribus acutiusculis, haud prominentibus. Thorax latitudine antica vix brevior, basin versus sensim angustatus, basi coleopteris paulo angustior, angulis posterioribus obtusis, sat convexus, punctis oblongis sat profundis at raris impressus, dorso subseriatis. Elytra thorace parum longiora, punctis raris subseriatis impressa, picea. Abdomen subtilissime alutaceum, parcius minus

profunde punctatum, punctis in segmento quinto crebrioribus. Pedes dilute picei.

Variat minus maturus colore testaceo; tale specimen a Latreillio descriptum.

Obs. Affinis certe speciei sequenti, at magis elongatus, parcius punctatus.

- \* 8. *O. latipes*: Nigro-piceus, nitidus, elytris piceis, antennis pedibusque testaceis, capite, thorace elytrisque parce, abdomine crebre punctatis.  
— Long.  $2\frac{1}{2}$  lin.

*Oxytelus latipes* Grav. Mon. 198. 13. — Oliv. Encycl. meth. VIII. 616. 15.

*Molosoma latipes* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. p. 462.

Habitat in America septentrionali, e coll. Knochiana.

Nigro-piceus, nitidus, breviter subtiliter capite et thorace parce, elytris et abdomine densius fulvescenti-pubescent. Antennae testaceae, capite parum longiores, articulis ultimis 5 distincte crassioribus. Palpi rufo-testacei. Labrum et mandibulae rufo-piceae. Caput thoracis latitudine, subtilissime alutaceum, fronte punctis grossiusculis parcius impressa, apice leviter emarginata, angulis anterioribus acutiusculis, haud prominentibus. Thorax latitudine vix brevior, basin versus ab apice ultra medium parum, dein fortius angustatus, angulis posterioribus obtusis, transversim convexus, parcius punctatus, punctis oblongis, in dorso biseriatis. Elytra thorace haud longiora, punctis maiusculis oblongis parioribus vage impressa, brunneo-picea. Abdomen subtilissime alutaceum, sat crebre punctatum. Pedes rufo-testacei.

Obs. Specimen a Dom. Zimmermann ex Carolina missum duplo maius, ceterum haud distinctum.

#### O b s e r v a t i o.

Speciem huius generis in resina Animae inclusam Dom. Hope Transact. Ent. Soc. Lond. II. p. 52. t. 7. f. 1. hoc modo descripsit:

*O. brunnicornis*: Niger, nitidus, tibiis anterioribus dilatatis, antennis pedibusque brunneis. Corpus cylindricum, nitidum, nigrum, long.  $4\frac{3}{4}$  lin. Caput elongatum, mandibulis intus dentatis. Antennae brunneae, pubescentes, capite longiores, articulo primo basi contracto, apice dilatato, secundo minore, tertio praecedente longiore, sequentibus 7 aequalibus, extimo ovato, antice subconico. Thorax fere quadratus, capite brevior, laevis. Scutellum triangulare, magnum. Elytra thorace longiora, parum latiora, nigra, nitida, marginibus posticis brunneis. Pedes rufo-brunnei, tibiis anterioribus dilatatis, compressis, spinulosis.

### Holotrochus Er.

#### *Tibiae muticae.*

Corpus elongatum, subcylindricum, alatum. Caput thorace minus et angustius, basi haud constrictum, oculis mediocribus, rotundatis, haud prominulis. Labrum transversum, apice subtruncatum. Mandibulae breves, validiusculae, muticae. Maxillae (t. V. f. 14. b.) malis corneis, exteriore apice barbata, interiore lineari, interne ciliata. Palpi maxillares crassiusculi, articulo primo minuto, secundo obconico, tertio hoc brevior, quarto iterum longiore, apice acuminato. Labium (t. V. f. 14. a.) mento latitudine sua haud brevior, lateribus et apice rotundato, ligula cornea, integra, paraglos-



sis minutis, linearibus, membranaceis. Palpi labiales articulo primo crassiusculo, secundo hoc brevior, tertio elongato, apice acuminato. Antennae breviusculae, apicem versus sensim leviter incrassatae, articulo primo magis minusve elongato, sub capitis margine recondito, secundo crassiusculo, obconico, penultimis pluribus transversis, ultimo globoso. Thorax coleopterorum latitudine, iisque basi arcte applicatus. Scutellum rotundato-subtriangulare. Elytra truncata. Abdomen elongatum, immarginatum cylindricum, segmento sexto conico, apice truncato. Pedes breviusculi, tibiis simplicibus, margine exteriori spinulis brevissimis ciliato, tarsis 5-articulatis, articulis 4 primis minutis, inter se aequalibus, quinto his coniunctis longitudine aequali.

## OBSERVATIONES.

1. Corpus glabrum. Thorax basi utrinque obsolete impressus. Elytra stria suturali inculpta.

2. Genus ab Osorio tibiis muticis facile distinctum, habitu Lispino non nimis dissimile, a quo autem praeter coxas anticlas exsertas abdominis segmento septimo conspicuo, statura magis cylindrica quam depressa et fronte immarginata discedit.

3. Sexus differentia latet.

\* 1. *H. volvulus*: *Nigro-piceus, subnitidus, antennis pedibusque rufis, laevis, thorace basi utrinque foveolato*. — Long.  $2\frac{1}{2}$  lin.

T. II. f. 4.

Habitat in ins. Puerto-Rico, Dom. Moritz.

Corpus elongatum, subcylindricum, omnium subtilissime punctato-coriaceum, minus nitidum, piceum. Antennae capite paulo longiores, rufae. Palpi rufi. Caput thorace angustius, fronte antice emarginata. Thorax coleopterorum latitudine, transversim subquadratus, lateribus subrectis, parallelis, angulis omnibus rectis, anterioribus deflexis, modice transversim convexus, basi apud angulos posteriores utrinque foveola minuta impressus. Elytra thorace paulo longiora, stria suturali impressa. Abdomen lateribus obsolete punctatum, segmento septimo profunde emarginato. Pedes rufi.

2. *H. cylindrus*: *Cylindricus, niger, nitidus, laevis, antennis pedibusque rufis*. — Long. 2 lin.

Habitat in ins. Puerto-Rico, Dom. Moritz.

Praecedente paulo minor et angustior, nitidus, niger. Antennae capite vix longiores, rufae. Os rufo-piceum. Caput thorace angustius, parce omnium subtilissime punctatum, fronte antice truncata. Thorax coleopterorum fere latitudine, latitudine vix brevior, lateribus rectis, parallelis, angulis omnibus rectis, anterioribus deflexis, transversim convexus, parce subtilissime punctatus. Elytra thorace parum longiora, omnium subtilissime coriacea, stria suturali impressa. Abdomen parce obsoletius punctatum, segmento septimo apice emarginato. Pedes rufi.

\* 3. *H. crassicollis*: *Brevis, cylindricus, niger, nitidus, antennis pedibusque piceis, capite, thorace elytrisque subtiliter punctatis*. — Long.  $1\frac{3}{4}$  lin.

Habitat in Madagascar, Dom. Goudot.

Praecedentibus brevior, crassior, cylindricus, niger, nitidus. Antennae

capitis longitudine, piceae. Caput thorace angustius, leviter convexum, parcius subtiliter punctatum, fronte antice leviter emarginata. Thorax coleopteris paulo latior, latitudine dimidio brevior, lateribus subrotundatis, basin versus subangustatus, fortiter transversim convexus, parcius subtiliusque licet sat distincte punctatus, postice utrinque obsolete impressus. Elytra thorace parum longiora, parce subtiliter obsoletius punctata, stria suturali sat profunda impressa. Abdomen omnium subtilissime confertissimeque punctato-coriaceum, punctis subtilibus obsoletisque sparsum, segmento septimo integro. Pedes picei.

## Subtribus tertia.

### Genuini.

*Antennae 11-articulatae.*

*Ligula membranacea.*

*Tarsi 3-articulati.*

Labrum transversum, corneum, apice limbo coriaceo instructum Bledio; utrinque lacinia membranacea auctum Platystheto, Oxytelo, Phloeonaeo, Trogophloeoo, Apocello.

Mandibulae falcatae, dentatae.

Maxillae malis membranaceis, interiore trapezoidea, interne ciliata, exteriore parallelopipeda, apice barbata.

Palpi maxillares articulo primo minuto, tertio leviter incrassato, quarto tenui, subulato.

Mentum transversum, corneum.

Ligula membranacea.

Paraglossae ligulae connatae eamque haud superantes plerisque; nullae Bledio.

Palpi labiales breves, articulo tertio subulato.

Antennae 11-articulatae, magis minusve distincte refractae.

Oculi mediocres.

Thorax ab elytris remotus, basi plerumque rotundatus.

Scutellum plerumque obsoletum.

Abdomen marginatum.

Pedes coxis intermediis

distantibus Platystheto, Oxytelo;

approximatis Bledio, Phloeonaeo, Trogophloeoo, Apocello;

tibiis anticis extus muricato-spinulosis Bledio;

extus seriatim spinulosis, Platystheto, Oxytelo, Phloeonaeo;

intermediis etiam spinulosis Bledio, Platystheto;

omnibus muticis Trogophloeoo, Apocello;

tarsis 3-articulatis, articulis 2 primis brevibus.

**Bledius Leach.***Mannerh. Brachelytr. p. 44.**Bledius Steph. Illustr. V. p. 307. — Hesperophilus ibid. p. 309.**Paraglossae nullae.**Pedes intermedii basi approximati.**Tibiae anteriores externe biseriatim spinulosae.*

Corpus elongatum, lineare, subcylindricum, alatum. Caput thorace minus et angustius, porrectum, basi vix constrictum, oculis semiglobosis, modice prominulis. Labrum (t. V. f. 15. c.) transversum, medio subsinuatum, margine antico membrana prominula limbato, laciniis lateralibus parvis, angulatum prominulis, spatio intermedio setis crassiusculis dense ciliato. Mandibulae validiores, magis minusve prominentes, vel ante apicem vel medio dentatae. Maxillae (t. V. f. 15. b.) mala interiore margine exteriori coriaceo, interiore membraneo, apicem versus spinulis breviusculis ciliato, mala exteriori longiore, coriacea, apice dense barbata. Palpi maxillares mala exteriori parum longiores, articulo primo minuto, secundo elongato-obconico, tertio hoc paulo longiore, leviter incrassato, quarto minuto, subulato. Labium (t. V. f. 15. a.) mento latitudine vix brevior, apice truncato, ligula membranea, biloba, lobis divergentibus, paraglossis nullis. Palpi labiales articulis duobus primis aequalibus, tertio his paulo longiore et tenuiore. Antennae fractae, breviusculae, articulo primo elongato, secundo tertioque obconicis, sequentibus longioribus, his sensim paululum maioribus, ultimis tribus nonnunquam distincte maioribus, ultimo ovato. Thorax coleopteris latitudine subaequalis, lateribus pone medium ad angulos posteriores subito fortius convergentibus, convexus. Elytra truncata. Abdomen lineare, marginatum, infra convexum. Pedes breviusculi, coxis intermediis approximatis, tibiis anticis leviter incrassatis, externe biseriatim fortius spinulosis, inter spinulas dense setulosis, intermediis spinularum serie armatis, posticis muticis, omnibus seriatim pilosis, tarsis triarticulatis, articulo tertio primis duobus coniunctis multo longiore.

**O b s e r v a t i o n e s.**

1. Genus a reliquis valde distinctum labri ligulaeque structura, et tibiis anticis incrassatis, fortius spinulosis. Corpus subtiliter tenuiterque pubescens.

2. *Hesperophilus* ab *Anglis* auctoribus nonnisi capite thoraceque in utroque sexu muticis distinctus.

3. Sexus differentia varia: in nonnullis caput et thorax masculorum cornuta, in aliis abdominis segmentum inferius sextum apice emarginatum, sinu autem membrana clauso, in aliis septimum in altero sexu aliter constructum, in aliis differentia quaevis me fugit.

4. Victus in riparum argilla arenave. Cuniculos fodiunt, quos per paria incolunt. Species nonnullae ripis littoribusve salsis propriae. Larvae eodem modo viciant. Species quaedam maris accolae alternatim undis submittuntur. Sectantur in quovis aetatis statu a *Dyschiriis* eorumque larvis. (Conf. *Rudd. Ent. Mag.* II. p. 160.). *Vesperis* copiosi volitant. Olent.

\* 1. *B. taurus*: *Fronte bicorni, niger, antennis pedibusque rufo-piceis, thorace subtiliter coriaceo, parcius punctato, subtiliter canaliculato, maris spinoso. — Long. 3 — 3½ lin.*

*Mannerh. Brachelytr. 44. 2. — Runde Brachelytr. Hal. 18. 2.*

*Oxyt. taurus Germ. Faun. Ins. Europ. XII. 2.*



*Oxyt. furcatus* Oliv. *Encycl. meth.* VIII. 616. 12. (*forte*)

*Var. Elytris testaceis, circa scutellum triangulariter fuscis.*

*Bledius Skrimshiri* Curtis *Britt. Ent.* III. t. 143.

*Bledius Ruddii* Steph. *Illustr. Britt. Ent.* V. t. 127. f. 3.

Habitat in Illyriae, in ins. Jonicarum, Sardiniae, Angliae littoribus marinis, haud infrequens etiam in ripis lacuum salsorum Germaniae interioris.

B. tricorni magnitudine aequalis, niger, parcius pubescens. Antennae capite sesqui longiores, apice leviter incrassatae, piceo-rufae. Os rufo-piceum. Mandibulae ante apicem fortiter unidentatae. Caput maris laevigatum vel omnium subtilissime coriaceum, nitidum, vertice transversim leviter impresso, fronte planata, utrinque ad antennarum basin cornu valido compresso, erecto, subrecto, apice pubescente armata, feminae subtiliter coriaceo, opaco, vertice fronteque medio transversim leviter impressis, hac utrinque ad antennae basin cornu brevi, lato, compresso, truncato armata. Thorax coleopterorum prope latitudine, latitudine paulo brevior, lateribus anterieus subrectis, infra medium subito fortiter conniventibus, angulis posterioribus obtusis, apice subbisinuatus, parum convexus, confertissime subtiliter coriaceus, minus fortiter parcius punctatus, subtiliter canaliculatus, maris ab apice ultra medium sensim angustatus, angulis anterioribus acutis, antrosum prominentibus, apice medio in spinam longam porrectam, apice subcompressam, barbatam, producto, canalicula dorsali ante spinae basin evanescente, feminae muticus, lateribus ultra medium parallelis, angulis anterioribus subrectis, canalicula dorsali integra. Elytra thorace paulo longiora, crebrius minus fortiter punctata, iam tota nigra, iam testacea, spatio triangulari circa scutellum fusco. Abdomen subtilissime punctato-coriaceum, subtile crebrius punctatum, nitidum, ano piceo. Pedes piceo-rufi, coxis concoloribus.

Abdominis segmentum inferius septimum maris apice subtruncatum, feminae bisinuatum.

Variat cornubus capitis thoracisque longioribus brevioribusve. Cornua capitis breviora nonnunquam posterius medio dilatata.

Obs. 1. Olivier thoracis spinam apice bifurcatam refert, forte pilis in spinae apice humore conglutinis deceptus.

2. B. Ruddii Steph. statura minore et cornubus debilioribus a genuinis haud rite distinctus.

\* 2. *B. vitulus*: Capite bicorni, nigro, testaceo, thorace subtiliter coriaceo, parcius punctato, canaliculato. — Long.  $2\frac{1}{2}$  —  $2\frac{2}{3}$  lin.

Habitat in Arabia deserta, Dom. Prof. Ehrenberg.

Statura omnino B. bicornis, testaceus, subnitidus, parcius pubescens. Antennae capite sesqui longiores, articulo tertio secundo subaequali, apicem versus sensim leviter incrassatae, rufo-piceae, medio fuscae. Os rufo-piceum. Mandibulae ante apicem fortiter unidentatae. Caput thorace paulo angustius, omnium subtilissime coriaceum, supra nigrum, subnitidum, fronte late leviter excavata, vertice transversim leviter impresso, margine laterali utrinque in cornu erectum breviusculum, latum, compressum, apice vel truncatum vel emarginatum elevata. Thorax coleopterorum latitudine, latitudine

tertia parte brevior, lateribus anterioribus rectis, postice subito subsinuato-conviventibus, angulis anterioribus acutiusculis, leviter prominulis, leviter convexus, confertissime subtiliter coriaceus, parce fortius punctatus, profundius canaliculatus. Elytra thorace paululum longiora, minus crebre sat profunde punctata. Abdomen subtilissime punctato-coriaceum, subtus vage obsoleteque punctatum. Alae albo-hyalinae.

Abdominis segmentum inferius septimum in utroque sexu apice leviter bisinuatum, lobo intermedio maris truncato, feminae acuminato.

Obs. Maribus maioribus capitis cornua maiora magis elevata et apice magis emarginata; feminae capitis structura omnino cum maribus minoribus congrunt.

\* 3. *B. bicornis*: *Fronte bicorni, niger, ore, antennis, pedibus elytrisque rufis, his sutura late nigris, thorace piceo, coriaceo, parcius punctato, canaliculato.* — Long.  $2\frac{1}{3}$  —  $2\frac{2}{3}$  lin.

*Oxyt. bicornis* Ahrens Faun. Ins. Europ. VI. 15.

*Bledius bicornis* Runde Brachelytr. Hal. 18. 4.

Habitat in ripa lacus salsi Mannsfeldiensis, frequens; in Grusinia, Dom. Marschall v. Biberstein.

*B. tricorni* dimidio minor, habitu potius *B. tauri*, niger, tenuiter pubescens. Antennae capite sesqui longiores, articulo tertio secundo paulo brevior, apicem versus leviter incrassatae, articulis ultimis tribus distincte paulo maioribus. Mandibulae modice prominentes, fortiter falcatae, ante apicem fortiter unidentatae. Os rufo-piceum. Caput thorace dimidio angustius, subtiliter confertissime coriaceum, opacum, vertice fronteque transversim leviter impressis, hac utrinque ad antennarum basin cornu brevi, lato, compresso, maris subacuminato, feminae truncato, erecto armata. Thorax coleopteris vix angustior, latitudine paulo brevior, lateribus anterioribus ultra medium subrectis, parallelis, postice subito fortiter convergentibus, angulis posterioribus rotundatis, apice utrinque subsinuatus, parum convexus, confertissime subtilius coriaceus, minus confertim sat profunde punctatus, canaliculatus, iam niger, iam fusco-piceus, opacus. Elytra thorace paulo longiora, sat crebre profundeque punctata, vel nigro-picea, lateribus rufo-testacea, vel rufo-testacea, sutura latius angustiusve picea. Abdomen subtiliter confertissime punctato-coriaceum, subtus crebrius, supra parce obsoletius punctatum, parum nitidum, apice piceo-testaceum. Pedes rufo-testacei.

Abdominis segmentum inferius septimum maris leviter emarginatum, feminae bisinuatum.

\* 4. *B. hianulus*: *Fronte bituberculata, niger, antennis, pedibus elytrisque testaceis, his macula communi picea, thorace rufo-piceo, subtiliter coriaceo, parcius punctato, canaliculato.* — Long.  $2\frac{1}{3}$  lin.

Habitat in Russia meridionali, Dom. Godet, in Grusinia, Dom. Marschall v. Biberstein.

Statura et summa affinitate *B. bicornis*, et minoribus eius speciminibus aequalis, at paulo gracilior, piceo-niger, tenuiter pubescens. Antennae capite sesqui longiores, articulo tertio secundo brevior, ultimis tribus distincte paulo maioribus, rufo-testaceae. Os rufum. Mandibulae ante apicem

sat fortiter unidentatae. Caput thorace angustius, nigro-piceum, opacum, subtilissime coriaceum, fronte utrinque longitudinaliter subimpressa, inter antennarum insertionem linea transversali profundiore exarata, medio puncto maiusculo impresso notata, maris postice late leviterque transversim impressa, margine laterali in utroque sexu supra antennarum basin modice elevato, tuberculiformi, rufo-piceo. Thorax coleopteris parum angustior, latitudine parum brevior, lateribus anterie ultra medium subrectis, postice subito fortiter convergentibus, angulis posterioribus rotundatis, parum convexus, confertissime subtiliter coriaceus, minus crebre sat fortiter punctatus, canaliculatus, rufo-piceus, minus nitidus. Coleoptera thorace paulo longiora, minus confertim punctata, dilute testacea, macula oblonga communi in media sutura nigra piceave. Abdomen subtilissime coriaceum, subtus obsolete punctatum, nigro-piceum, nitidulum, ano piceo-testaceo. Pedes testacei, coxis concoloribus.

Abdominis segmentum inferius septimum in utroque sexu apice bisinuaturn, lobo intermedio maris emarginato, feminae acuminato.

Obs. A praecedente praecipue statura graciliore, frontis margine laterali in utroque sexu haud in cornu formam elevato, elytris minus dense punctatis etc. distinctus.

- \* 5. *B. iuvenicus*: *Fronte subcornuta, niger, antennarum basi pedibusque rufis, thorace subtiliter coriaceo, parcius punctato, canaliculato.* — Long. 2 lin.

Habitat in Rossia meridionali, Dom. Godet.

*B. bicorni* valde affinis, at paulo gracilior et dimidio minor, niger, longius pubescens. Antennae capite plus sesqui longiores, articulo tertio secundo paulo brevior, apicem versus leviter incrassatae, articulis ultimis tribus distincte paulo maioribus, fuscae, articulis duobus primis rufis. Mandibulae haud prominentes. Oris partes rufae. Caput thorace angustius, subtilissime confertissimeque punctato-coriaceum, opacum, vertice late profundeque impresso, fronte late leviterque longitudinaliter excavata, inter antennarum basin linea tenui transversali exarata, margine laterali supra antennarum basin in cornu perbreve, latum, compressum, truncatum elevato. Thorax coleopterorum fere latitudine, latitudine vix brevior, lateribus anterie ultra medium rectis, parallelis, postice subito fortiterque convergentibus, basi truncatus, angulis posterioribus rotundatis, leviter convexus, subtiliter coriaceus, parum nitidus, minus crebre sat profunde punctatus, canaliculatus. Elytra thorace parum longiora, parcius et minus fortiter punctata, nigra, parum nitida, margine apicali piceo. Abdomen subtilissime punctato-coriaceum, subtus obsolete punctatum, nitidum, apice summo piceo-testaceo. Pedes testaceo-rufi, coxis anterioribus concoloribus.

Sexus differentia latet.

- \* 6. *B. tricornis*: *Niger, antennis pedibusque piceis, elytris rufis, thorace subtiliter coriaceo, dense punctato, canaliculato, maris spinoso.* — Long.  $2\frac{2}{3}$  —  $2\frac{1}{3}$  lin.

*Mannerh. Brachelytr.* 44. 1. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 456. 1. — *Runde Brachelytr. Hal.* 17. 1. — *Er. Col. March.* I. 578. 1.



- Staph. tricornis* Herbst Archiv. 149. 9. t. 30. f. 8. — Payk. Mon. Staph. 51. 37. Faun. Suec. III. 396. 38. — Oliv. Ent. III. 42. 30. 41. t. 6. f. 56.
- Oxyt. tricornis* Grav. Micr. 109. 11. Mon. 196. 11. — Latr. Hist. nat. Crust. et Ins. IX. 365. 11. — Oliv. Encycl. meth. VIII. 616. 13. — Gyll. Ins. Suec. II. 444. 1. — Guér. Iconogr. Ins. t. 9. f. 10. a. b.
- Staph. armatus* Panz. Faun. Germ. 66. 17. (male)

Habitat in Europa.

Corpus nigrum, subnitidum, tenuiter griseo-pubescent. Antennae capite sesqui longiores, articulo secundo tertioque aequalibus, ab articulo septimo inde leviter incrassatae, iam nigro-, iam rufo-piceae. Mandibulae prominentes, ante apicem fortiter dentatae. Caput maris impunctatum, laevigatum vel omnium subtilissime alutaceum, vertice impresso, fronte depressa, lateribus supra antennarum basin acute angulato-elevatis, feminae subtiliter confertissime alutaceum, fronte leviter convexa, sparsim punctata, lateribus supra antennarum basin obtuse angulato-elevatis, inter antennarum basin linea tenui transversali impressa. Thorax coleopteris parum angustior, latitudine apicis paululum brevior, basin versus sensim leviter angustatus, lateribus subrectis, paulo pone medium subito fortius convergentibus, basi truncatus, angulis posterioribus obtusis, leviter convexus, subtiliter confertissime coriaceus, magis minusve crebre punctatus, canaliculatus, feminae muticus, maris apice medio in spinam longam, porrectam, canaliculatam productus, plerumque caput superantem, individuis maioribus longiorem, minoribus brevior. Elytra thorace paulo longiora, dense profunde punctata, rufa, macula scutellari maiore minoreve nigricante. Abdomen supra utrinque parce obsoleteque, subtus crebre punctatum, ano piceo-rufo. Pedes rufo-, rarius nigro-picei.

- \* 7. *B. unicornis*: Niger, antennis pedibusque rufo-piceis, thorace subtiliter coriaceo, crebre punctato, canaliculato, maris spinoso. — Long. 2 lin.

Mannerh. Brachelytr. 45. 3. — Runde Brachelytr. Hal. 18. 3.

*Oxyt. unicornis* Germ. Faun. Ins. Europ. XII. 3.

Habitat ad Germaniae interioris lacus salsos, et in Europae meridionalis littoribus.

Statura fere *B. fracticornis*, niger, subnitidus, tenuiter pubescens. Antennae capite paulo longiores, articulo tertio secundo subaequali, apicem versus sensim incrassatae, rufo-piceae, basi magis rufo-testaceae. Mandibulae leviter prominentes, arcuatae, ante apicem fortiter unidentatae, piceo-rufae. Oris partes omnes rufae. Caput maris laeve, nitidum, vertice excavato, fronte ante oculos linea obsoleteque transversali impressa, lateribus supra antennarum basin obtusius elevato-angulata, feminae subtilissime coriaceum, fronte media foveola minuta punctiformi notata, anterieus utrinque ad antennarum basin leviter longitudinaliter impressa, margine laterali supra antennarum insertionem leviter elevato. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus anterieus ultra medium rectis, parallelis, postice subito fortiter convergentibus, basi truncatus, angulis posterioribus obtusis, leviter convexus, subtilissime coriaceus, crebrius profunde punctatus, subtiliter canaliculatus, feminae muticus, maris antice medio in spinam

longam porrectam, haud canaliculatam, apice rufo-piceam productus. Elytra thorace paulo longiora, sat crebre profundeque punctata, callo humerali margineque apicali saepius picescentibus, nonnunquam tota obscure picea. Abdomen supra utrinque parce obsoleteque, subtus sat crebre minus profunde punctatum, ano piceo. Pedes rufi, coxis anterioribus concoloribus, femoribus posticis nigris.

8. *B. mandibularis*: Testaceus, fronte nigricante, thorace obsolete punctato, canaliculato, rufo-testaceo. — Long.  $3\frac{3}{4}$  lin.

Habitat in America septentrionali (?), coll. Dom. Chevrolat.

Statura *B. tricornis*, testaceus, parum nitidus, subtiliter griseo-pubescentis. Antennae capite parum longiores, tenuiusculae, articulo tertio secundo paulo longiore, ultimis haud distincte crassioribus. Mandibulae elongatae, porrectae, parum incurvae, interne medio fortiter acuteque unidentatae, ante apicem leviter dilatatae, rufae, apice piceae. Palpi pallide testacei. Caput maiusculum, confertissime subtiliter alutaceum, parce obsoleteque punctatum, fronte inter oculos utrinque oblique impressa, posterius canalicula brevi profundiore notata, rufo-testaceum, fronte verticeque nigris. Thorax coleopterorum prope latitudine, latitudine paulo brevior, lateribus anterioribus ultra medium subrectis, parallelis, postice basin versus subito fortiterque convergentibus, basi truncatus, leviter convexus, subtilissime confertissimeque alutaceus, parce obsoleteque punctatus, canaliculatus, canalicula medio minus profunda, rufo-testaceus. Elytra thorace sesqui longiora, omnium subtilissime coriaceo-rugulosa, minus dense subtilius obsoleteque punctata. Abdomen omnium subtilissime confertim alutaceo-strigosum, infra parce obsoleteque punctatum.

Sexus differentia latet.

- \* 9. *B. rubiginosus*: Rufo-testaceus, fronte nigra, thorace crebrius fortiusque punctato, canaliculato, rufo, nitido. — Long.  $2\frac{1}{2}$  lin.

Habitat in Carolina meridionali, Dom. Zimmermann.

Habitu omnino *B. fracticornis*, at paulo maior, praecipue robustior, saturate rufo-testaceus, subnitidus, parce subtiliterque griseo-pubescentis. Antennae capite sesqui longiores, articulo tertio secundo longitudine aequali, ab articulo septimo inde leviter incrassatae, testaceae, apice rufo-piceae. Mandibulae breviores, haud prominentes, leviter falcatae, ante apicem unidentatae, piceo-rufae. Palpi testacei. Caput thorace paulo angustius, omnium confertissime subtilissimeque punctato-alutaceum, parce obsoleteque punctatum, plica longitudinali prope oculum utrumque sat elevata, fronte utrinque leviter impressa, apice angulo utroque denticulo minuto obsoleteque terminato, minus nitidum, supra nigro-piceum, ore et infra rufum. Thorax coleopteris parum angustior, latitudine haud brevior, lateribus modice rotundatis, pone medium basin versus convergentibus, angulis posterioribus rotundatis, convexus, canaliculatus, canalicula profundiore, crebrius fortiusque punctatus, interstitiis laevibus. Elytra thorace paulo longiora, dense profundeque punctata, dilutius testacea. Abdomen supra laeve, infra sparsim punctatum. Femora flavescens-testacea.

Sexus differentia latet.

10. *B. lateralis*: *Rufus, nitidus, capite nigro, elytris sutura late nigricante, thorace profunde punctato canaliculatoque, disco piceo.* — Long. 2 lin.

Habitat prope Puerto Cabello in Columbia, Dom. Moritz.

Statura omnino *B. fracticornis*, longius pubescens. Antennae capite sesqui longiores, articulo tertio secundo subaequali, apicem versus sensim incrassatae, rufae. Mandibulae falcatae, ante apicem fortius unidentatae, rufae. Palpi testacei. Caput thorace paulo angustius, nigrum, opacum, confertissime subtilissimeque coriaceum, fronte posteriorius foveola minuta punctiformi notata, antice utrinque obsolete longitudinaliter impressa, angulis anterioribus acutiusculis, prominulis. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus leviter rotundatis, a medio inde basin versus modice convergentibus, convexus, crebre fortiter profundeque punctatus, interstitiis laevibus, canalicula longitudinali, medio sat forti, basi apiceque subtiliore exaratus, rufo-piceus, nitidus, disco obscuriore. Elytra thorace paulo longiora, creberrime sat profunde punctata, rufa, subnitida, secundum suturam late picea. Abdomen supra omnium subtilissime confertissimeque, subtus crebrius fortiusque punctatum, rufum. Corpus subtus totum rufum. Pedes testacei.

- \* 11. *B. politus*: *Nitidus, niger, antennis, pedibus elytrisque testaceis, his sutura nigra, thorace piceo-rufo, parce fortius punctato, canaliculato.* — Long.  $2\frac{1}{3}$  lin.

Habitat in Carolina meridionali, Dom. Zimmermann.

Statura omnino *B. fracticornis*, at paululum maior. Corpus totum glabrum. Antennae capite sesqui longiores, articulo tertio secundo paulo brevior, ab articulo septimo inde leviter incrassatae, testaceae. Mandibulae breviores, parum curvatae, infra medium unidentatae, rufo-piceae. Palpi rufo-testacei. Caput thorace dimidio angustius, omnium subtilissime confertissimeque alutaceum, parce obsoleteque punctatum, plica longitudinali prope oculum utrumque sat elevata, fronte utrinque leviter impressa, lineaque transversali inter antennarum basin exarata, posterius inter oculos in tuberculum latiusculum leviter elevata, apice utrinque denticulo subtili terminata, nigro-piceum, minus nitidum. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus levissime rotundatis, pone medium basin versus oblique convergentibus, angulis posterioribus obtusissimis, convexus, canaliculatus, parce disperse minus subtiliter punctatus, interstitiis politis, nitidus, piceo-rufus. Elytra thorace tertia parte longiora, parce fortius punctata, nitida, testacea, sutura apicem versus attenuata nigro-picea. Abdomen omnium confertissime subtilissimeque punctato-alutaceum, infra parce obsoletius punctatum, piceo-nigrum, nitidum, apice summo rufescente. Pectus nigrum, nitidum. Pedes cum coxis testacei.

Sexus differentia latet.

- \* 12. *B. aquarius*: *Niger, antennis basi testaceis, elytris, tibiis tarsisque rufis, thorace subtilius alutaceo, opaco, crebre fortius punctato, postice subtiliter canaliculato.* — Long.  $2\frac{2}{3}$  lin.

Habitat Genevae, Dom. Chevrier. Dedit Dom. Prof. Germar.



Habitu omnino *B. tricornis* et minoribus eius individuis aequalis, niger, parum nitidus, tenuiter griseo-pubescens. Antennae capite sesqui longiores, graciliores, articulo tertio secundo paulo brevior, apicem versus sensim leviter incrassatae, nigrae, basi testaceae. Mandibulae perparum prominentes, rufae. Palpi rufi. Caput thorace dimidio angustius, confertissime subtilissimeque punctato-alutaceum, fronte utrinque longitudinaliter leviter impressa et parcius obsoletiusque punctata, inter oculos posterius linea transversali leviter exarata, inter antennarum insertionem leviter transversim impressa. Thorax coleopterorum latitudine, latitudine summa vix brevior, lateribus rotundatus, postice sinuato-coarctatus, basi truncatus, angulis posterioribus acutiusculis, prommulis, sat convexus, confertissime subtiliter alutaceus, crebre fortius punctatus, linea lata media longitudinali laevi (impunctata) posterius subtiliter canaliculata. Elytra thorace vix longiora, dense profundeque punctata, rufa, summa basi nigricantia. Abdomen confertissime subtilissimeque punctato-alutaceum, nitidulum, supra parce, infra crebrius punctatum. Pedes nigro-picei, tibiis tarsisque rufis.

Sexus differentia latet.

\* 13. *B. fracticornis*: *Niger, antennarum basi pedibusque rufis, coxis anterioribus piceis, thorace crebre fortiter punctato, canaliculato.* — Long. 2 lin.

*Mannerh. Brachelytr.* 45. 5. — *Runde Brachelytr. Hal.* 18. 5. — *Er. Col. March.* I. 579. 2.

*Staph. fracticornis* Payk. *Mon. Car. App.* 135. 17—18. *Faun. Suec.* III. 382. 19.

*Oxyt. fracticornis* Gyll. *Ins. Suec.* II. 446. 2. — *Sahlb. Ins. Fenn.* I. 408. 1

*Oxyt. pallipes Gallicus* Grav. *Mon. p.* 198.

*Bledius pallipes* Boisd. et Lacord. *Faun. Ent. Paris.* I. 458. 4.

*Var. Elytris rufis.*

*Oxyt. tricornis* var. *minor* Grav. *Mon. p.* 196.

Habitat in Europa.

*Niger*, subnitidus, tenuiter subtiliterque fusco-pubescens. Antennae capite sesqui longiores, articulis ultimis distincte crassioribus, octavo sequentibus paulo minore, tertio secundo longitudine aequali, nigrae, basi rufae. Mandibulae breviusculae, fortiter falcatae, ante apicem dentatae, rufo-piceae, basi nigrae. Caput thorace paulo angustius, opacum, parce obsoleteque subtilius punctatum, utrinque ad antennarum basin leviter impressum. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus, angulis posterioribus et basi rotundatus, convexus, evidenter canaliculatus, dense fortiterque punctatus, interstitiis laevibus. Elytra thorace tertia parte longiora, dense profunde punctata, iam nigra, iam saturatius dilutiusve rufa, basi interiore suturaque magis minusve nigris. Abdomen omnium confertissime subtilissimeque punctatum, nitidum, ano rufo-piceo. Pedes rufo-testacei, coxis etiam anticis nigro-piceis.

Abdominis segmentum inferius sextum maris apice sinuatum, sinu membrana repleto, feminae integrum.

- \* 14. *B. femoralis*: *Niger, subnitidus, ore pedibusque rufis, coxis piceis, femoribus picescentibus, thorace crebre profundeque punctato, canaliculato.* — *Long.*  $1\frac{3}{4}$  lin.

*Mannerh. Brachelytr.* 45. 9.

*Oxytelus femoralis* *Gylk. Ins. Suec. IV.* 497. 3 — 4.

Habitat in Suecia, Dom. Schüppel.

Habitus et summa affinitas praecedentis. Niger, subnitidus, subtiliter griseo-pubescent. Antennae capite sesqui longiores, articulo tertio secundo dimidio brevior, apicem versus ab articulo septimo inde modice incrassatae, articulo octavo nono minore, totae nigro-piceae. Mandibulae falcatae, paulo ante apicem acute dentatae, rufae. Palpi rufo-testacei. Caput thorace paulo angustius, opacum, parce subtiliterque obsoletius punctatum, fronte utrinque leviter impressa. Thorax coleopteris paulo angustior, latitudine summa paulo brevior, lateribus parum rotundatis, basi subito oblique angustatus, angulis posterioribus subrotundatis, convexus, evidenter canaliculatus, crebre fortius punctatus, interstitiis subtiliter alutaceis. Elytra thorace tertia parte longiora, dense profunde punctata, piceo-nigra. Abdomen omnium subtilissime alutaceum, nitidulum, subtus parcius punctatum. Pedes rufi, coxis omnibus piceis, femoribus a basi inde magis minusve alutaceum.

Sexus differentia latet.

Obs. Differt a praecedente antennis nigro-piceis, thorace minus rotundato, evidenter alutaceo, minus dense punctato, etc.

- \* 15. *B. longulus*: *Niger, nitidus, antennis elytris pedibusque rufis, thorace profunde punctato, subtiliter canaliculato.* — *Long.*  $1\frac{2}{3}$  lin.

*Er. Col. March. I.* 579. 3.

Habitat in Germania.

*B. fracticorni* dimidio fere minor, praecipue brevior, niger, nitidus, tenuiter griseo-pubescent. Antennae capite sesqui fere longiores, articulo tertio secundo dimidio brevior, apicem versus ab articulo quarto inde leniter incrassatae, articulo octavo sequente vix minore, rufae, medio subfuscescentes. Os rufum. Mandibulae minus elongatae, leviter falcatae, medio bidenticulatae. Caput thorace angustius, subtiliter obsoleteque parce punctatum, fronte utrinque ad antennarum basin leviter impressa. Thorax coleopterorum fere latitudine, latitudine summa paulo brevior, lateribus fortiter rotundatus, basi truncatus, angulis posterioribus obtusissimis, convexus, distincte canaliculatus, sat crebre profundeque punctatus, interstitiis laevibus vel oculo fortiter armato omnium subtilissime coriacea. Elytra thorace paululum longiora, densius profunde punctata, laete rufa. Abdomen omnium subtilissime confertissimeque punctatum, nitidum, ano concolore. Pedes rufi, coxis anterioribus concoloribus.

Sexus differentia latet. Abdominis segmentum septimum inferius in omnibus apice profunde bisinuatum.

- \* 16. *B. procerulus*: *Nigro-piceus, antennis basi pedibusque testaceis, elytris castaneis, thoracis longitudine, hoc dense fortiter punctato, canaliculato.* — *Long.*  $1\frac{2}{3}$  lin.

Habitat in Austria, Dom. Schüppel.

**B. fracticorni** dimidio minor et angustior, nigro-piceus, minus nitidus, parce griseo-pubescens. Antennae capite sesqui longiores, articulo tertio secundo paulo brevior, ab articulo septimo inde distinctius incrassatae, testaceae, apicem versus picescentes. Oris partes testaceae. Mandibulae haud prominentes, falcatae, ante apicem dentatae. Caput thorace parum angustius, omnium subtilissime alutaceum, sparsim subtilius obsoletiusque punctatum, fronte utrinque late leviter impressa. Thorax coleopteris parum angustior, latitudine haud brevior, lateribus subrectis, postice introrsum modice rotundatis, angulis posterioribus obtusis, basi truncatus, modice convexus, dense fortius profunde punctatus, interstitiis subtilissime alutaceis, medio profunde canaliculatus. Elytra thorace vix longiora, dense fortius profunde punctata, piceo-castanea, subnitida. Abdomen elongatum, omnium subtilissime alutaceum, nitidulum, apice rufo-piceum. Pedes testacei, coxis omnibus concoloribus.

Sexus differentia latet.

Obs. Hanc speciem, ab affinibus coleopteris thorace neque latioribus, neque longioribus distinctam, sub *Oxyteli* proceri nomine misit Dahl.

\* 17. *B. nigricans*: *Niger, subnitidus, ore, antennis, pedibus anoque rufo-testaceis, elytris margine inflexo rufo, thorace crebre profundeque punctato.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Rossia meridionali, Dom. Godet.

Habitu omnino *B. fracticornis*, at paulo minor, niger, nitidulus, densius cinereo-pubescens. Antennae capite sesqui longiores, articulo tertio secundo longitudine subaequali, ab articulo septimo inde incrassatae, basi rufo-testaceae, apice rufae. Ores partes rufo-testaceae. Mandibulae haud prominentes. Caput thorace paululum angustius, omnium subtilissime alutaceum, opacum, subtiliter obsoleteque vage punctatum, fronte utrinque ad antennarum basin leviter longitudinaliter et inter antennis linea transversali impressa. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus antè parum, posterius angulos posteriores versus fortius rotundatis, his obtusis, convexus, sat crebre fortiter profundeque punctatus, interstitiis laevibus, linea media longitudinali laevi, haud vero canaliculatus, supra niger, subtus rufo-piceus. Elytra thorace sesqui longiora, confertim sat profunde punctata, supra nigra, margine inflexo, interdum etiam callo humerali obscure rufis. Abdomen omnium confertissime subtiliter punctulatum, ano rufo-testaceo. Pedes rufo-testacei, coxis anterioribus concoloribus. Alae albae.

Abdominis segmentum sextum inferius maris apice emarginatum, sinu membrana clauso, feminae integrum.

Obs. *B. fracticorni* dimidio fere minor, antennis totis coxisque anterioribus testaceis, thorace haud canaliculato etc. facile distinctus.

\* 18. *B. dissimilis*: *Niger, nitidus, antennis, pedibus anoque rufis, elytris thorace longioribus, rufo-castaneis, sutura late nigricante, thorace crebre profundeque punctato.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Germania, Austria, Hungaria, Istria et Gallia.

Statura omnino *B. fracticornis*, niger, nitidus, tenuiter griseo-pubescens. Antennae capite sesqui fere longiores, articulo tertio secundo dimidio brevior, ab articulo septimo inde leviter incrassatae, rufo-testaceae. Oris partes rufo-



testaceae. Mandibulae falcatae, ante apicem fortiter dentatae. Caput thorace paululum angustius, omnium subtilissime alutaceum, minus nitidum, fronte utrinque subtilissime obsolete punctata, ad antennarum insertionem leviter impressa. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus anterioribus parum, postice angulos posteriores versus oblique subrotundatis, his obtusissimis, convexus, sat fortiter profundeque dense punctatus, interstitiis laevibus, linea media longitudinali laevi subelevata. Elytra thorace tertia parte longiora, profundius dense punctata, saturate rufa, nitida, secundum suturam late nigricantia. Abdomen subtilissime confertissimeque punctatum, nitidum, ano rufo. Pedes testacei, coxis anterioribus rufis.

Abdominis segmentum sextum inferius maris apice late emarginatum, sinu membrana clauso, lateribus utrinque spina longa, tenui, leviter sursum arcuata, acuta armatum, feminae integrum.

Obs. Proximus sequentibus, a *B. rufipenni* thorace multo crebrius punctato, elytris dorso nigricantibus, statura paulo minore etc., a *B. crassicolli* coleopteris thorace latioribus et longioribus distinctus. Mas praeterea abdomine infra apice spinis duabus longioribus, superne etiam conspicuis armato insignis.

\* 19. *B. rufipennis*: *Niger, nitidus, antennis, elytris pedibusque rufis, thorace parcius profunde punctato, elytris thorace longioribus.* — Long. 2 lin.

Habitat in Austria.

Statura omnino *B. fracticornis*, niger, nitidus, parcius griseo-pubescent. Antennae capite sesqui longiores, tenuiores, articulo tertio latitudine secundo subaequali, ab articulo septimo inde leviter incrassatae, totae rufo-testaceae. Mandibulae haud prominentes, falcatae, ante apicem dentatae. Caput thorace paulo angustius, omnium subtilissime alutaceum, minus nitidum, fronte utrinque leviter longitudinaliter impressa et parce subtiliterque punctata, angulis apicalibus denticulo minutissimo terminata. Thorax coleopteris paulo angustior, latitudine haud brevior, lateribus ultra medium subrectis, postice oblique subrotundatis, basi truncatus, angulis posterioribus obtusissimis, convexus, parcius at sat fortiter profundeque punctatus, interstitiis laevigatis, linea lata media longitudinali laevi, haud canaliculatus. Elytra thorace paulo longiora, minus confertim profundius punctata, laete rufa, circa scutellum parvo spatio nigricantia. Abdomen omnium subtilissime alutaceum, parce obsoleteque punctatum, ano concolore. Pedes rufi, coxis concoloribus.

Abdominis segmentum inferius sextum maris late leviter sinuatum, sinu membrana repleto, utrinque denticulo seu spina brevior, subtili, acuta, leviter sursum curvata terminato, feminae integrum.

Obs. Olim pro sequentis varietate mihi habitus, iam vero, speciminibus utriusque sexus pluribus examinatis, differentias constantes reperio. Distinctus videtur a sequente statura paulo maiore, thorace parce punctato, coleopteris thorace latioribus et evidentior longioribus, ano semper concolore etc. Abdominis structura mas medium tenet inter praecedentem et sequentem.

\* 20. *B. crassicollis*: *Niger, nitidus, antennis, elytris, ano pedibusque rufis, thorace crebre profunde punctato, elytris thorace vix longioribus.* — Long. 1½ lin.

*Bledius crassicollis* Boisd. et Lacord. Faun. Ent. Paris. I. 456. 2. — Er. Col. March. I. 580. 4.

## Habitat in Germania.

Statura fere *B. fracticornis*, at paulo minor et gracilior, niger, nitidus, tenuiter griseo-pubescent. Antennae capite duplo longiores, tenuiores, articulo tertio secundo paulo brevior, ab articulo septimo inde leviter incrassatae, rufae. Mandibulae haud prominentes, falcatae, ante apicem fortiter unidentatae, piceae. Oris partes inferiores rufae. Caput thorace parum angustius, opacum, fronte utrinque parce obsoleteque subtiliter punctata et ad antennarum insertionem fortius impressa, angulis apicalibus subdenticulatis. Thorax coleopteris haud angustior, latitudine summa parum brevior, lateribus anterieus levissime, postice cum angulis posterioribus fortiter rotundatis, basi ipsa truncata, convexus, sat crebre fortiterque punctatus, interstitiis laevibus, haud canaliculatus, linea media longitudinali laevi. Elytra thorace perparum longiora, sat crebre profundeque punctata, saturate rufa, circa scutellum nigricantia. Abdomen omnium subtilissime confertissimeque punctatum, nitidum, ano rufo. Pedes rufi, coxis concoloribus.

Abdominis segmentum inferius sextum maris apice sinuatum, sinu membrana repleto, denticulo obtusiusculo utrinque terminato, feminae integrum.

Variat ano concolore.

\* 21. *B. opacus*: *Niger, antennarum basi pedibusque testaceis, elytris rufis, sutura nigricante, thorace subtiliter coriaceo, opaco, subtiliter canaliculato, parcius obsoleteque punctato, angulis posterioribus subrotundatis.* — Long.  $1\frac{3}{4}$  lin.

*Er. Col. March. I. 581. 5.*

*Staph. opacus* Block *Ins. Vall. Plauens. 117. 7. t. 7. f. 7.*

*Bledius castaneipennis* Mannerh. *Brachelytr. 45. 6.* — *Boisd. et Lacord. Faun. Ent. Paris. I. 457. 3.*

## Habitat in Germania.

Niger, parce tenuiterque griseo-pubescent. Antennae capite sesqui fere longiores, articulo tertio secundo dimidio brevior, ab articulo septimo inde leviter incrassatae, nigrae, basi testaceae, vel testaceae, apice fuscescentes. Mandibulae parum prominentes, leviter falcatae, interne medio obsoleteque bidenticulatae, rufo-piceae. Os inferius testaceum. Caput confertissime subtiliter coriaceum, subopacum, fronte utrinque longitudinaliter et inter antennarum insertionem transversim impressa, utrinque secundum oculi marginem interiori longitudinally elevata, angulis apicalibus denticulo minuto terminatis. Thorax medio coleopteris parum angustior, latitudine tertia parte brevior, antrorsum subangustatus, lateribus praecipue ad angulos posteriores rotundatis, his obtusissimis, subrotundatis, modice convexus, confertissime subtiliter coriaceus, opacus, subtiliter canaliculatus, parcius parum profunde punctatus. Elytra thorace sesqui prope longiora, dense profunde punctata, testaceo-rufa, nitida, secundum suturam magis minusve late nigricantia. Abdomen omnium subtilissime confertissimeque punctatum, nitidum, ano piceo. Pedes rufo-testacei, coxis anterioribus concoloribus.

Sexus differentia latet.

- X \* 22. *B. erraticus*: *Niger, subnitidus, antennis, pedibus, elytris anoque rufis, thorace subtiliter coriaceo, parce obsoleteque punctato.* — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 582. 6.*

Habitat in Germania.

*B. opaco proximus, at paulo minor, niger, subnitidus, tenuiter cinereo-pubescent. Antennae capite sesqui longiores, articulo tertio secundo dimidio brevior, ab articulo septimo inde modice incrassatae, rufae, articulis duobus primis testaceis. Oris partes rufo-testaceae. Caput thorace paulo angustius, omnium subtilissime confertissimeque coriaceum, subnitidum, fronte utrinque parce subtiliter obsoleteque punctata, et ad antennarum basin leviter impressa. Thorax coleopteris parum angustior, latitudine paulo brevior, lateribus subrotundatis, a medio versus basin subsinuato-angustatis, angulis posterioribus obtusis, non vero obtusissimis, convexus, confertissime subtilissimeque coriaceus, opacus, parcius subtiliusque obsolete punctatus, linea media longitudinali laevi (impunctata), haud canaliculatus, supra nigro-vel fusco-piceus, subtus rufus. Elytra thorace paulo longiora, minus confertim punctata, testaceo-rufa, nitida. Abdomen supra laeve, nitidum, nigro-piceum, ano rufo, subtus obsoleteque punctatum, piceo-rufum. Pedes rufo-testacei.*

Sexus differentia latet.

Variat elytris prope suturam nigricantibus vel totis nigro-piceis.

Obs. A *B. opaco* thorace haud canaliculato, fronte haud transversim impressa et elytris brevioribus distinctus.

- \* 23. *B. pallipes*: *Niger, antennarum basi pedibusque testaceis, thorace subtiliter coriaceo, opaco, subtiliter canaliculato, parcius obsoleteque punctato, angulis posterioribus obtusis.* — Long.  $1\frac{3}{4}$  lin.

*Mannerh. Brachelytr. 45. 8. — Runde Brachelytr. Hal. 19. 7.*

*Oxyt. pallipes Grav. Mon. 197. 12. — Oliv. Encycl. meth. VIII. 614. 14. — Gyll. Ins. Suec. II. 447. 3. IV. 497. 3. — Sahlb. Ins. Fenn. I. 408. 2.*

Habitat in Austria, Dom. Ullrich, in Gallia, Dom. Aubé.

Statura et summa affinitas *B. opaci*. *Niger, tenuiter griseo-pubescent. Antennae capite sesqui longiores, articulo tertio secundo duplo brevior, ab articulo septimo inde sensim leviter incrassatae, piceae, basi testaceae. Mandibulae parum prominentes, leviter falcatae, medio leviter bidentatae, rufo-testaceae. Palpi testacei, articulo tertio fuscescente. Os inferius testaceum. Caput thorace perparum angustius, subtilissime coriaceum, opacum, utrinque parce obsolete punctatum, fronte utrinque ad antennarum basin impressa. Thorax coleopteris paulo angustior, latitudine tertia parte brevior, lateribus modice rotundatus, basin versus a medio inde sensim angustatus, angulis posterioribus, licet obtusis, subprominulis, leviter convexus, confertissimeque coriaceus, opacus, subtiliter canaliculatus, parcius obsolete punctatus. Elytra thorace plus sesqui longiora, dense punctata, nigra, nitida. Abdomen omnium subtilissime punctatum, nitidum, ano piceo. Pedes rufo-testacei, coxis anterioribus concoloribus.*

Sexus differentia latet.

Obs. A *B. opaco* differt thoracis angulis posterioribus prominulis, elytris longioribus, densius punctatis, nigris, fronte haud transversim impressa.



- \* 24. *B. atricapillus*: *Piceus*, capite nigro, antennarum basi, elytris pedibusque pallide testaceis, thorace subtilissime alutaceo, parce obsoleteque punctato, subtiliter canaliculato. — Long.  $1\frac{1}{2}$  lin.

Mannerh. *Brachelytr.* 45. 8. — Runde *Brachelytr.* Hal. 19. 6.

*Oxyt. atricapillus* Germ. Faun. Ins. Europ. XI. 4. — Nicolai *Col. Halens.* 40. 3.

Habitat in Germania.

Praecedente paulo minor et gracilior, piceus, parum nitidus, subtiliter griseo-pubescens. Antennae capite sesqui longiores, articulo tertio secundo multo brevior, ab articulo septimo inde modice incrassatae, testaceae, basi pallidiores, apice fuscescentes. Mandibulae leviter prominentes, minus falcatae, acutae, medio fortiter unidentatae, rufo-testaceae. Oris partes testaceae. Caput thorace paulo angustius, omnium subtilissime alutaceum, fronte utrinque ad antennarum insertionem leviter impressa, nigrum, opacum. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus anterie subrectis, fere parallelis, pone medium subitu fortiter convergentibus, angulis posterioribus cum basi leviter rotundatis, convexus, confertissime subtilissimeque alutaceus, parce subtiliter obsoleteque punctatus, subtiliter canaliculatus, rufo-piceus, parum nitidus. Elytra thorace plus sesqui longiora, densius subtiliusque punctata, pallide testacea, prope suturam leviter infusca. Abdomen parce subliterque punctatum, dorso plerumque nigrum, nitidum. Pedes cum coxis omnibus pallide testacei.

Sexus differentia latet.

- \* 25. *B. nanus*: *Niger*, nitidulus, antennarum basi, pedibus elytrisque testaceis, his sutura late nigra, thorace parce subtiliter obsoleteque punctato, subtiliter canaliculato. — Long.  $1\frac{1}{4}$  lin.

Habitat in Austria, Dom. Ullrich.

Habitu omnino *B. fracticornis*, at minutus, niger, nitidulus, tenuiter griseo-pubescens. Antennae capite sesqui longiores, articulo tertio secundo multo minore, ultimis tribus abrupte maioribus, nigrae, articulis duobus primis testaceis. Mandibulae parum prominentes, falcatae, inter medium et apicem unidentatae. Oris partes rufo-testaceae. Caput thorace parum angustius, omnium subtilissime coriaceum, fronte utrinque ad antennarum basin longitudinaliter impressa. Thorax apice coleopteris paulo angustior, basin versus angustatus, lateribus ante medium leviter, a medio inde fortiter convergentibus, angulis posterioribus obtusis, convexus, omnium subtilissime coriaceus, parce subtiliter obsoleteque punctatus, subtiliter canaliculatus, subtus utrinque piceo-rufus. Elytra thorace paulo longiora, crebrius sat profunde punctata, testacea, secundum suturam late nigricantia. Abdomen omnium subtilissime confertissimeque punctatum, nitidum. Pedes testacei, coxis anterioribus concoloribus.

Sexus differentia latet.

- \* 26. *B. pusillus*: *Niger*, antennarum basi, pedibus elytrisque rufis, thorace coriaceo, crebrius obsolete punctato, obsolete canaliculato. — Long. 1 lin.

*Er. Col. March.* I. 583. 7.

Habitat in Germania, Gallia.

Habitu prope *B. crassicollis*, at minutus, niger, tenuiter griseo-pubescent. Antennae capite sesqui longiores, articulo tertio secundo multo minore, ultimis tribus distincte licet minus abrupte maioribus, fusco-ferrugineae, articulis duobus primis testaceis. Mandibulae perparum prominentes, ante apicem obsolete unidentatae. Os rufum. Caput thorace parum angustius, omnium subtilissime alutaceum, nitidum, fronte utrinque obsolete punctata, inter antennarum insertionem sulco transversali, utrinque foveola minuta terminato, impressa. Thorax apice coleopteris parum angustior, latitudine paulo brevior, lateribus anterieus subrectis, posterius a medio inde fortiter convergentibus, angulis posterioribus obtusis, leviter coriaceus, parum nitidus, crebrius minus subtiliter obsolete canaliculatus, subtus rufus. Coleoptera thorace paulo longiora et latiora, minus confertim obsolete punctata, rufa, nitida. Abdomen omnium subtilissime punctulatum, nigrum, nitidum, ano ventrique rufo-piceis. Pedes rufo-testacei, coxis anterioribus concoloribus.

Obs. *B. nano* paulo brevior, fronte linea transversali impressa, thorace minus subtiliter coriaceo punctatoque, obsolete canaliculato, elytris secundum suturam haud nigris etc. facile distinguendus.

\* 27. *B. pygmaeus*: *Niger, antennarum basi, pedibus elytrisque testaceis, thorace crebrius obsolete punctato. — Long. 1 lin.*

*Er. Col. March. I. 583. 8.*

Habitat in Germania, Gallia.

Statura graciliore, minutus, *B. pusillo* longitudine aequalis, at angustior, niger, nitidus, tenuiter griseo-pubescent. Antennae capite sesqui longiores, articulo tertio secundo multo minore, ultimis tribus distincte maioribus, fuscae, articulis duobus primis testaceis. Mandibulae perparum prominentes, ante apicem obsolete unidentatae. Oris partes rufo-testaceae. Caput thorace paulo angustius, omnium subtilissime coriaceum, nitidum, fronte utrinque ad antennarum basin leviter impressa. Thorax coleopterorum latitudine, latitudine antica vix brevior, basin versus angustatus, lateribus anterieus parum, a medio inde fortius conniventibus, angulis posterioribus obtusis, convexus, subtiliter coriaceus, crebrius obsolete punctatus, linea media longitudinali laevi (impunctata), haud canaliculatus, supra nigro-piceus, subtus lateribus piceo-rufus. Elytra thoracis longitudine, minus crebre obsolete punctata, testaceo-rufa. Abdomen confertim subtilissime punctatum, nigrum, nitidum, apice summo piceo-rufo, subtus ab apice inde rufescens. Pedes testacei, coxis concoloribus.

Obs. *B. pusillo* proximus, statura autem graciliore, coleopteris thorace neque longioribus, neque latioribus, thorace haud canaliculato etc. specie distinguendus.

\* 28. *B. troglodytes*: *Niger, antennis, elytris pedibusque pallide testaceis, thorace subtiliter canaliculato, parcius punctato, interstitiis laevigatis, fronte apice bituberculata. — Long. 1 lin.*

Habitat in Carolina meridionali, Dom. Zimmermann.

Praecedente paulo minor, niger, nitidus, parce subtiliterque cinereo-pubescent. Antennae capite sesqui longiores, tenuiores, articulo tertio secundo multo minore, apice sensim leviter incrassatae, pallide

testaceae, apice nonnunquam picescentes. Mandibulae vix prominentes. Oris partes omnes testaceae. Caput thorace paulo angustius, laevigatum, fronte ad antennarum insertionem utrinque leviter impressa, apice utrinque tuberculo minuto terminata, inter tubercula emarginata. Thorax coleopteris paulo angustior, latitudine antica haud brevior, basin versus angustatus, lateribus leviter rotundatis, pone medium fortius conniventibus, angulis posterioribus obtusis, convexus, subtiliter canaliculatus, minus confertim subtilius punctatus, interstitiis laevigatis, nigro-piceus, nitidus. Elytra thorace sesqui longiora, dense subtiliter punctata, pallide rufo-testacea, secundum suturam obsolete infuscata. Abdomen supra laevigatum, infra parce subtiliterque punctatum, nigrum, nitidum. Pedes pallide flavo-testacei.

Sexus differentia latet.

29. *B. parvulus*: *Rufus, nitidus, capite nigro, thorace abdomineque supra piceis, elytris macula magna communi fusca, thorace canaliculato, parcius profunde punctato.* — Long.  $1\frac{1}{2}$  lin.

Habitat Novae Valentiae in Columbia, Dom. Moritz.

Habitu fere *B. nani*. Antennae capite sesqui longiores, articulo tertio secundo dimidio brevior, apicem versus sensim subincrassatae, testaceae, apice nigricantes. Os rufo-testaceum. Mandibulae fortiter falcatae, acutae. Caput thoracis fere latitudine, omnium subtilissime coriaceum, fronte subconvexa, ad antennarum basin vix impressa, nigrum, parum nitidum. Thorax coleopteris paulo angustior, latitudine parum brevior, lateribus anterioribus subrectis, pone medium subito convergentibus, angulis posterioribus obtusis, convexus, subtilius canaliculatus, minus crebre fortius profunde punctatus, interstitiis laevibus, piceus, nitidus. Coleoptera thorace paulo longiora et latiora, sat crebre profundeque punctata, pallide rufo-testacea, nitida, macula magna dorsali communi fusca. Abdomen laeve, nitidum, supra segmentis anterioribus margine postico, sexto toto piceo. Corpus subius totum rufum. Pedes testacei.

Sexus differentia latet.

30. *B. pumilio*: *Testaceus, capite nigro, pedibus elytrisque pallidis, his macula obsoleta fusca notalis, thorace parcius obsolete punctato.* — Long. vix 1 lin.

Habitat in Columbia, Dom. Moritz.

Statura parva *B. pygmaei*, at hoc etiam paulo minor, ideoque minimus in hoc genere. Antennae capite paulo longiores, articulo tertio secundo multo minore, apicem versus sensim crassiores, testaceae. Os testaceum. Mandibulae leviter falcatae, acutae, ante apicem subtiliter unidentatae. Caput nigrum, subnitidum, fronte parce subtilissime punctulata, utrinque obsolete impressa. Thorax coleopteris paulo angustior, latitudine paulo brevior, lateribus anterioribus parum, postice cum angulis posterioribus fortiter rotundatis, convexus, parcius obsoletiusque punctatus, linea media longitudinali laevi, canalicula media utrinque abbreviata subtilissima exaratus, rufo-testaceus, nitidus. Elytra thorace paulo longiora, crebre punctata, subtiliter pubescentia, pallide testacea, nitida, secundum suturam macula magna ob-



longa fusca obsoletiore notata. Abdomen saturate testaceum, nitidum. Pedes et pedes pallidè testacea.

Sexus differentia latet.

\* 31. *B. verres*: *Niger, parum nitidus, pedibus elytrisque pallide testaceis, his basi suturaque nigris, thorace subtilissime alutaceo, subtiliter punctato, canaliculato, fronte apice bidenticulata, antennarum basi maris elevata. — Long. 2 lin.*

Habitat in Sardinia, Dom. Prof. Gené.

*Niger*, subtiliter griseo-pubescent. Antennae capite duplo prope longiores, articulo tertio secundo duplo brevior, ultimis 4 distincte crassioribus, densius pubescentes, nigrae, articulo primo piceo, secundo testaceo, articulis tribus primis maris elongatis. Mandibulae porrectae, minus falcatae, acutae, maris longiores, medio dente valido, acuto, fere erecto armatae, feminae muticae. Caput thorace angustius, omnium subtilissime alutaceum, obsoletissime punctulatum, apice denticulis duobus approximatis, porrectis, maris longioribus, spinuliformibus armata, tuberculo ante oculos, cui antennae insertae sunt, in femina haud magis quam in reliquis prominulo, in mare magno, fere cylindrico et porrecto, antennarum insertionem longe ultra frontem elevata. Thorax coleopteris paulo angustior, latitudine antica paulo brevior, lateribus anterie subrectis, parallelis, pone medium subito fortiter convergentibus, angulis posterioribus rotundatis, modice convexus, subtilissime alutaceus, subnitidus, crebrius subtiliter obsoleteque punctatus, canaliculatus. Elytra thorace sesqui longiora, densius subtiliusque punctata, pallide rufo-testacea, sutura late basique nigra. Abdomen parce punctatum, parcius griseo-pilosellum, nitidum. Pedes pallide testacei, coxis omnibus et femoribus anticis nigro-piceis, femoribus posterioribus albidis.

Abdominis segmentum inferius septimum maris apice leviter bisinuatatum, feminae truncatum.

\* 32. *B. Niloticus*: *Nigricans, antennarum basi pedibusque pallide flavis, thorace transverso, subtilissime coriaceo, crebrius punctato, subtiliter canaliculato, fronte apice bituberculata. — Long. 1½ lin.*

Habitat in Aegypto, Dom. Prof. Ehrenberg.

*Nigricans*, parum nitidus, tenuiter griseo-pubescent. Antennae capite paulo longiores, articulo tertio secundo paulo brevior, apice sensim incrassatae, nigrae, articulis duobus primis testaceis. Os testaceum. Mandibulae elongatae, prominentes, leviter arcuatae, acutae, ante medium acute unidentatae. Caput thorace paulo angustius, subtilissime confertissimeque coriaceum, fronte utrinque super antennarum basin tuberculo rotundato, sat prominulo, testaceo munita, apice subretusa, bidenticulata. Thorax coleopteris parum angustior, latitudine paulo brevior, lateribus anterie rectis, parallelis, pone medium subito fortiter convergentibus, basi angulisque posterioribus subrotundatis, leviter convexus, omnium subtilissime coriaceus, subnitidus, subtiliter crebrius punctatus, subtiliter canaliculatus. Elytra thorace sesqui longiora, dense profundeque punctata. Abdomen supra laeve, subtus obsoletius punctatum. Pedes pallide testacei.

Mas mandibulis paulo longioribus, fronte subimpressa, tuberculis su-

per antennarum basin denticulisque apicalibus magis prominulis, acutis a femina distinguendus.

Variat immaturus testaceus, capite nigricante.

\* 33. *B. talpa*: *Niger, elytris nigro-cyaneis, nitidis, thorace opaco, subtiliter obsoleteque punctato, subtiliter canaliculato.* — Long. 2 lin.

*Mannerh. Brachelytr.* 45. 10. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 458. 5.

*Oxyt. talpa* Gyll. *Ins. Suec.* II. 448. 4. — *Sahlb. Ins. Fenn.* I 409. 3. — *Germ. Faun. Ins. Europ.* XI. 3. — *Zetterst. Ins. Lappon.* 73. 7.

Habitat in Suecia, Germania, Gallia.

Statura omnino *B. fracticornis*, niger, nitidus, longius griseo-pubescent. Antennae capite sesqui longiores, articulo tertio secundo multo minore, apicem versus leviter incrassatae, nigrae, articulo secundo piceo-testaceo. Mandibulae modice prominentes, leviter arcuatae, acutae, medio unidentatae, nigrae, apice piceo-rufae. Palpi nigri. Caput thorace angustius, omnium subtilissime coriaceum, opacum, vage subtiliter obsoleteque punctatum, fronte utrinque leviter longitudinaliter impressa. Thorax coleopteris dimidio angustior, latitudine parum brevior, lateribus medio fortiter rotundatus, posterius fortiter sinuato-angustatus, angulis posterioribus subprominulis, convexus, omnium subtilissime coriaceus, opacus, minus confertim subtiliter obsoleteque punctatus, subtiliter canaliculatus. Elytra thorace sesqui longiora, sat crebre fortiterque punctata, obscure coerulea, nitida. Abdomen omnium subtilissime confertissimeque punctatum, supra obsolete sparsim parceque, subtus crebrius fortiusque punctatum, nitidum. Pedes nigri, tarsis trochanteribusque piceo-testaceis.

Sexus differentia latet.

\* 34. *B. subterraneus*: *Niger, pedibus piceo-testaceis, femoribus nigricantibus, thorace opaco, subtiliter canaliculato, subtiliter obsoleteque punctato.* — Long. 1 $\frac{3}{4}$  lin.

*Er. Col. March.* I. 584. 9.

Habitat in Germania.

Niger, tenuiter griseo-pubescent. Antennae capite paulo longiores, articulo tertio secundo dimidio brevior, apicem versus sensim leviterque incrassatae, nigrae, apice saepe fusciscentes, articulis duobus primis nonnunquam testaceis. Mandibulae prominulae, tenues, parum arcuatae, acutae, medio obsolete dentatae, rufae. Palpi piceo-testacei, articulo tertio nigropiceo. Caput thorace angustius, omnium subtilissime coriaceum, opacum, subtiliter obsoleteque vage punctatum, fronte utrinque ad antennarum basin longitudinaliter impressa. Thorax coleopteris dimidio fere angustior, latitudine parum brevior, lateribus ante medium rotundatus, pone medium sinuato-angustatus, angulis posterioribus obtusis, subprominulis, convexus, omnium subtilissime coriaceus, opacus, minus confertim subtiliter obsoleteque punctatus, subtiliter canaliculatus. Elytra thorace duplo prope longiora, subtilius creberrime punctata, subnitida, pube brevissima conspersa. Abdomen omnium subtilissime confertissimeque punctatum, nitidum. Pedes coxis femoribusque nigro-piceis, trochanteribus, tibiis tarsisque piceo-testaceis.

Sexus differentia latet.

- \* 35. *B. arenarius*: *Niger, nitidus, antennis apicem versus, tibiis tarsisque fusco-testaceis, elytris testaceis, basi suturaque nigris, thorace brevi, subtiliter canaliculato, parce subtiliterque punctato.* — Long.  $1\frac{1}{2}$  lin.

*Mannerh. Brachelytr.* 46. 11. — *Er. Col. March. I.* 585. 10.

*Staph. arenarius* Payk. *Faun. Suec. III.* 382. 20.

*Oxyt. arenarius* Gyll. *Ins. Suec. II.* 448. 5. — *Sahlb. Ins. Fenn. I.* 409. 4.

*Rudd Ent. Mag. II.* p. 180.

Habitat in arena in lacum ripis, in Suecia, Germania.

*B. opaco* duplo fere minor, niger, nitidus, tenuiter griseo-pubescens. Antennae capite paulo longiores, graciliores, articulo tertio secundo multo minore et duplo brevior, ultimis tribus abrupte distincteque crassioribus, piceo-testaceae, articulis duobus primis nigris. Mandibulae modice prominentes, tenues, acutae, leviter arcuatae, medio obsolete bidentatae, rufae. Palpi nigri. Caput thorace dimidio angustius, omnium subtilissime coriaceum, parce obsoleteque punctulatum, subnitidum, fronte utrinque ad antennarum basin foveola minuta impressa, angulis apicalibus denticulo minimo prominulo acuto terminatis. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus anterioribus subrectis, pone medium angulos posteriores versus subito sinuato-angustatis, his subprominulis, convexus, omnium subtilissime coriaceus, nitidulus, subtiliter canaliculatus, minus crebre subtiliter obsoleteque punctatus. Elytra thorace sesqui longiora, confertim subtiliter punctata, pallide rufo-testacea, basi suturaque nigris, vel nigra, angulo apicali exteriori magis minusve late testaceo. Abdomen supra laeve, subtus crebre obsolete punctatum, nitidum. Pedes nigri, trochanteribus, tibiis tarsisque obscure testaceis.

Sexus differentia latet.

- \* 36. *B. debilis*: *Niger, opacus, antennis, pedibus elytrisque pallide testaceis, his macula scutellari triangulari nigra, thorace subtilissime coriaceo, subtiliter punctato canaliculatoque.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Rossia meridionali, Dom. Godet.

*B. arenario* paulo longior, tenuiter pubescens, niger, capite thoraceque opacis, abdomine subnitido. Antennae capite sesqui longiores, articulo tertio secundo paulo brevior, apicem versus sensim incrassatae, testaceae. Mandibulae porrectae, parum falcatae, tenues, edentatae, acutae, rufae. Oris partes reliquae testaceae. Caput omnium confertissime subtilissimeque coriaceum, nigrum, opacum, fronte utrinque ad antennarum insertionem vix impressa. Thorax coleopteris paulo angustior, latitudine parum longior, lateribus et basi leviter, angulis posterioribus fortiter rotundatis, apice submarginatus, leviter convexus, omnium confertissime subtilissimeque coriaceus, vage subtiliter obsoleteque punctatus, subtiliter canaliculatus, nigro-piceus, opacus. Elytra thorace paulo longiora, sat crebre subtilius punctata, pallide testacea, nitida, circa scutellum spatio parvo triangulari nigricante. Abdomen supra subtilissime confertissimeque punctatum, nigrum, nitidum, subtus crebrius punctatum. Corpus totum subtus rufo-piceum. Pedes cum coxis omnibus pallide testacei.

Sexus differentia latet.



Species mihi invisae.

37. *B. elongatus*: *Elongatus, niger, subnitidus, profunde punctatus, elytrorum macula oblonga rufa, pedibus pallidis, thorace orbiculato, obsolete canaliculato.*

*Mannerh. Brachelytr.* 45. 4.

Habitat Petropoli in terra argillacea.

Medius quasi inter *B. tricornem* et *fracticornem*, illo parum brevior, sed fere duplo angustior, thorace convexiore mutico et colore praeterea facile dignoscitur. (*Mannerh.*)

38. *B. brunnipennis*: *Niger, obscurus, elytris pedibusque brunneis.*

*Staph. brunnipennis* *Fab. Syst. El.* II. 596. 31.

Habitat in India orientali.

Medius. Antennae nigrae, basi ferrugineae. Caput et thorax scabra, nigra, obscura. Elytra subpunctata, immaculata. Abdomen atrum, ano sive ultimo segmento brunneo. Pedes brunnei. (*Fab.*)

39. *B. tuberculatus*: *Niger, elytris testaceis, ad suturam nigris, capite tuberculato.*

*Staph. tuberculatus* *Fabr. Ent. Syst. Suppl.* 181. 49—50. *Syst. El.* II. 601. 62.

Habitat in India orientali.

Parvus. Caput magnum, nigrum, mandibulis exsertis, elongatis tuberculisque duobus elevatis, obtusis. Antennae testaceae. Thorax niger. Elytra testacea, ad suturam nigra. Abdomen nigrum, ano testaceo. Pedes testacei. (*Fab.*)

40. *B. pallipennis*: *Testaceus, capite nigro, clypeo apice elevato, bidentato, thorace transverso, canaliculato. — Long. ultra 3½ lin.*

*Oxyt. pallipennis* *Say Act. Acad. Nat. Scienc. Philadelph. III. I.* 155. 1.

Habitat in Missuria.

Testaceus, punctatus, pubescens. Caput nigrum, antice sparsim punctatum, clypeo apice medio elevato, prominulo, bidentato. Antennae et carinae ad earum basin pallide rufae. Mandibulae porrectae, piceae, medio bifidae, dente superiore brevior. Palpi pallidi. Thorax latitudine brevior, rufo-piceus, linea dorsali exaratus. Elytra apice et sutura fuscescentibus. Pedes albi. (*Say.*)

41. *B. armatus*: *Dilute rufo-piceus, capite nigro, carina laterali apice picea. — Long. 2 — 2½ lin.*

*Oxyt. armatus* *Say Act. Acad. Nat. Scienc. Philadelph. III. I.* 152. 2.

Habitat in America septentrionali.

Dilute rufo-piceus, punctatus, leviter pubescens. Caput nigrum, obsolete punctatum, carina verticali super anteriorem oculorum partem abbreviata, ad antennarum basin abrupte terminata, apice picea, clypei angulis anterioribus reflexis. Antennae et palpi pallide rufi. Mandibulae piceae. Thorax latitudine haud brevior, linea longitudinali exaratus, disperse punctatus, mar-

gine nigricante. Elytra crebre distincte punctata, sutura nigricante. Femora testacea.

Mas dilutior, fronte inter oculos tuberculo munita, thorax linea longitudinali dorsali impunctata, abdomine dorso apice obscuriore. (Say).

42. *B. cordatus*: *Niger, elytris albidis, macula communi fusca.* — Long.  $1\frac{1}{2}$  lin.

*Oxyt. cordatus* Say *Transact. Amer. Philoph. Soc. Philadelph. Nov. Ser. IV.* 461. 3.

Habitat in Virginia.

Niger, subtiliter punctatus. Mandibulae piceae. Antennae pallidae, apice fuscescentes. Thorax angulis subrotundatis. Elytra flavescenti-alba, disco communi fuliginoso. Pedes flavescenti-albi, tibiis tarsisque obscurioribus.

Frequentissimus occurrit m. Octobre in cuniculis sub arenae littoralis superficie, aestu alterno opertus. Variis pusillis Crustaceis, fluctu recedente sub arenam refugientibus pasci videtur. (Say).

43. *B. fasciatus*: *Nigricans, elytris pallide flavis, abdomine testaceo, obsolete fusco-fasciato.* — Long.  $1\frac{1}{2}$  lin.

*Oxyt. fasciatus* Say *Act. Acad. Nat. Scienc. Philadelph. III.* 1. 156. 4.

Habitat in Missuria.

Nigricans, punctatus, pubescens. Caput nigrum, impunctatum, subtilissime granulatum. Os et antennae testaceae. Mandibulae piceae. Thorax nigro-piceus, grossius distincte punctatus, basi cum angulis posterioribus rotundatus, ab elytris remotus. Elytra pallide flava, basi interne et sutura fuscis, crebre grossius distincte punctata, apice obtuse rotundata. Pectus rufo-piceum. Pedes pallidiores. Abdomen supra rufescens, segmentis singulis linea transversali media obsoleta nigricante. (Say).

44. *B. melanocephalus*: *Pallide testaceus, capite pectoreque nigris, sutura fusca.* — Long. 1 lin.

*Oxyt. melanocephalus* Say *Act. Acad. Nat. Scienc. Philadelph. III.* 156. 3.

Habitat in Missouri fluminis ripis.

Pallide testaceus vel albidus. Caput nigerrimum, ore antennisque pallide testaceis. Mandibulae muticae. Elytra sutura nigricante. Pectus nigrum. Variat abdomine rufo-piceo. (Say).

45. *B. emarginatus*: *Niger, elytris albidis, sutura fuliginosa.* — Long. infra 1 lin.

*Oxyt. emarginatus* Say *Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV.* 461. 4.

Habitat in Indiana Americae septentrionalis.

Niger, punctatus. Caput crebre punctatum, at haud rugulosum, fronte inter antennis longitudinaliter bi-impressa. Antennae, mandibulae et palpi testacei. Thorax picescens, basi rotundatus, disperse distincte punctatus, linea dorsali evidenter canaliculatus. Elytra distincte punctata, vitta communi latiore fuliginosa, angulo apicali suturali obtuse rotundato. Pedes testacei, coxis femoribusque albidis.

Obs. Elytra apice ob angulum interiorem obtuse rotundatum profunde emarginata apparent. (Say).

## **Platysthetus Mannerh.**

*Mannerh. Brachelytr. p. 46.*

*Oxytelus Fam. II. Grav. Mon. p. 194.*

*Paraglossae apice liberae.*

*Pedes intermedii basi distantes.*

*Tibiae anteriores externe uni-seriatim spinulosae.*

Corpus oblongum, subdepressum, alatum. Caput sat magnum, porrectum, basi leviter constrictum, thoraci immissum, oculis rotundatis, modice prominulis. Labrum transversim subquadratum, laciniis membraneis triangularibus, acuminatis. Mandibulae modice prominentes, interne ante apicem bidentatae, basi membrana fimbriatae. Maxillae (t. V. f. 16. b.) malis membranaceis, interiore interne membranea, ciliata, exteriori longiore, apice dense barbata. Palpi maxillares mala exteriori paulo longiores, articulo primo minuto, secundo tertioque subaequalibus, apice leviter incrassatis, quarto minore, subulato. Labium (t. V. f. 16. a.) mento transversum, ligula membranea, lata, apice bisinuata, lobis lateralibus magis productis, acutis, paraglossis liberis, linearibus, falcatis, acutis, interne dense ciliatis. Palpi labiales articulo primo crassiusculo, cylindrico, secundo hoc paulo brevior, tertio secundo iterum longiore. Antennae fractae, apicem versus leniter incrassatae. Thorax transversus, caput excipiens, lateribus et basi rotundatus, ab elytris remotus. Elytra brevia, truncata, apice dehiscentia. Abdomen sursum reflectendum, supra planum, infra convexum. Pedes breviusculi, intermedii distantes, tibiis anterioribus externe uniseriatim spinulosis, anticis apice emarginatis, tarsis triarticulatis, articulo tertio reliquis duobus multo longiore.

### O b s e r v a t i o n e s.

1. Corpus glabrum. Thorax canaliculatus. Abdomen laevigatum.
2. Labri structura et pedibus intermediis basi distantibus cum Oxytelo convenit, differt autem habitu peculiari, ligula apice minus fortiter bisinuata, paraglossis basi ligulae connatis, palpis labialibus articulo ultimo haud subuliformi, pedibus intermediis fortius distantibus, tibiis intermediis externe spinulosis.
3. Mares capite maiore, plerique etiam fronte apice bispinosa, et abdominis segmentorum inferiorum structura distinguuntur.
4. Victus in fimo.
5. Metamorphosin *Pl. morsitantis* Dom. Bouché observavit et hoc modo descripsit: „Larva linearis depressa. antice angustata, parce setulosa, flava. Caput dilute piceum, lenticulare. Antennae 5-articulatae, articulis 3 primis cylindricis, tertio apice processu minuto elliptico aucto, reliquis ambobus mucronato-subulatis, penultimo apice setis duabus fortibus instructo. Mandibulae lineares, arcuatae, apice obtusae, bidentatae, dentibus piceis, apice nigricantibus. Labrum quadratum. Maxillae oblongae, intus processu angusto, apice introrsum curvato, interne setis ciliato instructae. Palpi maxillares 5-articulatae, articulo secundo longiore, introrsum arcuato. Mentum rotundatum. Ligula oblongo-quadrangularis. Palpi labiales 2-articulati, ligulae medio insertae. Prothorax capitis latitudine, mesothorace angustior, posterius utrinque papilla prominula instructus. Meso- et metathorax abdominis latitudine. Abdomen segmentis singulis utrinque parce rugosis, ultimo fortiter attenuato, caudis duabus, setosis terminato. Long. 2 lin. Victitat hieme in stercore bovino. Nympha oblongo-ovata, depressiuscula, parce setosa. Oculi nigri. Vertex setis duabus longis validis divergentibus. Mandibulae nigrae, apice tridentatae ut in imagine. Abdomen marginatum, segmento anali caudis duabus conicis terminato. Long.  $1\frac{1}{2}$  lin. Hunc statum 14 diebus absolvit. (Hist. Ins. I. 182. I. t. VIII. f. 14 — 21.)



- \* 1. *P. morsitans*: Niger, nitidus, ore, tibiis tarsisque testaceis, fronte depressa, canaliculata, thorace fortius canaliculato elytrisque crebre punctatis, longitudinaliter strigosis, his saepe testaceis. — Long.  $1\frac{1}{2}$  — 2 lin.

Mannerh. *Brachelytr.* 46. 2. — Runde *Brachelytr.* Hal. 19. 3. — Er. Col. March. I. 585. 1.

*Staph. morsitans* Payk. Mon. Curc. App. 145. 23 — 24. Faun. Suec. III. 383. 21. — Marsh. Ent. Brit. 508. 28.

*Oxyt. morsitans* Grav. Micr. 108. 9. Mon. 195. 9. — Gyll. Ins. Suec. II. 451. 7. — Sahlb. Ins. Fenn. I. 411. 6.

*Staph. trilobus* Oliv. Ent. III. 42. 20. 22. t. 5. f. 48. — Marsh. Ent. Brit. 507. 26.

*Oxyt. trilobus* Latr. Hist. nat. Crust. et Ins. IX. 364. 9. — Oliv. Encycl. meth. VIII. 615. 9.

*Staph. pallidipennis* Panz. Faun. Germ. 27. 16. Ent. Germ. 358. 46.

*Staph. arenarius* Fourcr. Ent. Paris. I. 172. 30.

*Platyth. striolatus* Boisd. et Lacord. Faun. Ent. Paris. I. 460. 2.

*Oxyt. mordax* Sahlb. Ins. Fenn. I. 411. 7.

*Platysth. mordax* Runde *Brachelytr.* Hal. 19. 2.

Habitat in Europa.

Niger, nitidus. Antennae capite sesqui longiores, articulo secundo longiore, 6 — 10 transversis, sensim crassioribus, ultimo ovato, piceae. Palpi picei. Caput maris magnum, thorace latius, feminae thorace minus et angustius, subdepressum, fronte linea postica transversali impressa sulculoque medio longitudinali sat profundis, clypei spatio impresso, sparsim, pone oculos fortius crebriusque punctatum. Thorax coleopterorum latitudine, latitudine plus dimidio brevior, lateribus et basi rotundatus, angulis posterioribus obtuse rotundatis, apice bisinuatus, angulis anterioribus rotundatis, parum convexus, sat profunde canaliculatus, crebre punctatus, praecipue versus latera, longitudinaliter strigosus. Elytra thoracis fere longitudine, minus confertim punctata, subtiliter strigosa, picea vel piceo-testacea. Abdomen nigro-pilosellum. Pedes testacei, femoribus saepe piceis.

Abdominis segmentum inferius septimum maris profunde triangulariter excisum, medio spinulis duabus subtilibus acutis armatum, feminae apice medio leviter productum.

Obs. *Oxyt. mordax* Sahlb. a minoribus huius speciei individuis haud satis distinctus mihi videtur.

- \* 2. *P. cornutus*: Niger, nitidus, fronte crebre punctata, maris apice bispinosa, thorace fortius canaliculato elytrisque obsolete punctatis, his saepius testaceis. — Long.  $1\frac{1}{3}$  —  $1\frac{2}{3}$  lin.

Mannerh. *Brachelytr.* 46. 1. — Boisd. et Lacord. Faun. Ent. Paris. I. 459. 1. — Runde *Brachelytr.* Hal. 19. 1. — Er. Col. March. I. 586. 2.

*Oxyt. cornutus* Grav. Micr. 109. 10. Mon. 195. 10. — Latr. Hist. nat. Crust. et Ins. IX. 365. 10. — Oliv. Encycl. meth. VIII. 615. 10. — Gyll. Ins. Suec. II. 450. 6. — Zetterst. Faun. Lappon. I. 94. 5. Ins. Lappon. 73. 5. — Sahlb. Ins. Fenn. I. 410. 5.

*Platysth. scybalarius* Runde *Brachelytr.* Hal. 19. 4.

Habitat in Europa; in Syria Dom. Prof. Ehrenberg.

*P. morsitanti* magnitudine aequalis, at corpus paulo magis cylindricum,

nigrum, subnitidum, omnium subtilissime alutaceum. Antennae capite plus sesqui longiores, articulo tertio secundo brevior, 6 — 10 sensim paulo crassioribus, non autem transversis, ultimo ovato, nigro-piceae. Caput maris thorace parum latius, feminae parum angustius, fronte subdepressa, sat crebre punctata, sulculo abbreviato medio sulcoque laterali longitudinalibus lineaque tenui transversali verticali impressa, margine antico in mare spinis duabus porrectis acutis validisque armato. Thorax apice coleopterorum latitudine, latitudine summa dimidio fere brevior, lateribus et basi eodem arcu rotundatis, apice emarginatus, angulis anterioribus acutis, leviter convexus, subtiliter obsoleteque sparsim punctatus, evidenter canaliculatus. Elytra vix thoracis longitudine, subtiliter obsoleteque sparsim punctata, iam nigra, iam piceo-testacea, disco dilutiore. Pedes picei, tarsis, interdum etiam tibiis testaceis.

Abdominis segmentum inferius sextum maris apice leviter emarginatum, septimum medio late impressum, apice truncatum, feminae apice leviter triangulariter productum.

Obs. Individua Europae meridionalis et Asiae occidentalis saepius elytris coloribus, Europae septentrionalis plerumque elytris disco luteis instructa sunt. Pl. scybalarius a Dom. Prof. Germar benevole communicatus cum huius speciei exemplis minutis omnino convenit.

\* 3. *P. nodifrons*: *Niger, nitidus, tibiis tarsisque testaceis, capite, thorace elytrisque sparsim punctatis, thorace profundius canaliculato, latitudine duplo brevior.* — Long.  $1\frac{1}{4}$  lin.

Mannerh. *Brachelytr.* 47. 3. — Er. Col. *March.* I. 587. 3.

*Oxyt. nodifrons* Sahlb. *Ins. Fenn.* I. 412. 8.

*Oxyt. nitens* Sahlb. *Ins. Fenn.* I. 413. 9.

*Oxyt. morsitans* var. c Gyll. *Ins. Suec.* II. 451. 7.

Habitat in Europa.

*P. morsitans* longe minor, niger, nitidus. Antennae capite duplo prope longiores, articulo tertio secundo brevior, 4 — 10 sensim paulo crassioribus, penultimis leviter transversis, ultimo oblongo-ovato, nigrae. Os piceum. Mandibulae sat elongatae. Caput maris thorace paulo maius, feminae eodem vix minus, sparsim punctatum, super oculos sulculis nonnullis impressum, linea transversali verticali magis minusve obsoleta, margine laterali super antennarum basin fortius elevato, ceterum aequale. Thorax coleopterorum latitudine vel iisdem paulo latior, latitudine antica duplo brevior, lateribus et basi eodem arcu rotundatis, apice truncatus, angulis anterioribus subrotundatis, leviter convexus, sat profunde canaliculatus, sparsim subtilius fortiusve punctatus, interstitiis laevissimis. Elytra thorace vix longiora, prope suturam subtiliter unistriata, sparsim subtilius fortiusve punctata, interstitiis laevissimis. Abdomen subtiliter griseo-pilosellum. Pedes testacei, femoribus saepius picei.

Abdominis segmentum inferius septimum maris longitudinaliter bica rinatum, apice truncatum, feminae apice leviter productum.

Obs. Haec species statura parum instabilis insigniter variat punctura, iam crebriore fortioreque (*O. nodifrons* Sahlb.), iam parciore et subtiliore (*O. nitens* Sahlb.): quae varietates speciminibus multis intermediis obviis haud rite specie distinguendae videntur.

- \* 4. *P. spinosus*: *Niger, nitidus, elytris pedibusque piceis, tarsis flavis, fronte maris apice bispinosa, capite, thorace elytrisque crebre punctatis, thorace profundius canaliculato, latitudine vix dimidio brevior.* — *Long.*  $1\frac{1}{4}$  lin.

Habitat in Lusitania, Com. Hoffmannsegg, Lutetiae, Dom. Dr. Aubé.

Subdepressus, niger, nitidus. Antennae capite duplo prope longiores, articulo tertio secundo paulo longiore, 4 — 10 sensim paulo crassioribus, penultimis haud transversis, ultimo oblongo-ovato, nigrae. Palpi nigropicei. Caput maris thoracis latitudine, feminae eodem paulo angustius, inter oculos crebre subtiliter punctatum, interstitiis laevissimis, fronte ad antennarum basin utrinque leviter impressa, vertice linea transversali tenui, super oculos sulculis duobus canaliculaque media brevissima obsoletaque longitudinalibus exarata, maris apice spinis duabus longis incumbentibus armata. Thorax apice coleopteris haud angustior, basin versus sensim modice angustatus, lateribus vix rotundatus, apice leviter bisinuatus, angulis anterioribus acutiusculis, posterioribus obtusis, basi leviter rotundatus, supra parum convexus, fortiter canaliculatus, crebrius subtiliter punctatus, interstitiis laevissimis. Elytra thorace haud longiora, parce subtiliter punctata, interstitiis laevissimis, nitida, nonnunquam fusco-picea. Abdomen subtiliter griseopilosellum. Pedes fusco-picei, tarsis flavis.

Abdominis segmentum inferius septimum maris apice truncatum, ad apicem denticulis duobus distantibus munitum, feminae apice leviter triangulariter productum.

5. *P. spiculus*: *Niger, nitidus, elytris fuscis, pedibus testaceis, capite, thorace elytrisque impunctatis, thorace subtiliter canaliculato, fronte maris apice bispinosa.* — *Long.*  $\frac{4}{5}$  lin.

Habitat prope Caracas in Columbia, Dom. Moritz.

Minutus, subdepressus, niger, nitidus. Antennae capite paulo longiores, articulo tertio secundo paulo brevior et duplo tenuiore, 4 — 10 sensim paulo crassioribus, ultimo ovato, nigrae. Palpi rufo-picei. Caput thorace paulo maius, impunctatum, fronte posterius linea transversali tenuissima et utrinque lineis tribus longitudinalibus sat profundis exarata, clypei spatio modice impresso, margine antico spinis duabus acutis incumbentibus armato. Thorax apice coleopteris latior, latitudine plus dimidio brevior, basin versus angustior, lateribus, basi angulisque posterioribus rotundatis, apice leviter bisinuatus, angulis anterioribus acutiusculis, parum convexus, subtiliter canaliculatus, ceterum laevissimus. Elytra thorace haud longiora, laevissima, dilute fusco-picea, nitida. Pedes testacei.

Abdominis segmentum inferius septimum maris apice subbisinuatum. Femina latet.

- \* 6. *P. Americanus*: *Niger, nitidus, elytris fusco-piceis, tibiis tarsisque testaceis, capite crebre punctato, fronte maris bispinosa, thorace obsolete canaliculato elytrisque omnium subtilissime punctulatis.* — *Long.*  $1\frac{1}{2}$  lin.

Habitat in Pennsylvania et Virginia, Dom. Zimmermann.

Niger, nitidus. Antennae capite plus sesqui longiores, articulo primo maris incrassato, tertio secundo paulo brevior, 6 — 10 sensim crassioribus,



8—10 transversis, ultimo subovato, nigrae. Mandibulae minus elongatae, palpique picei. Caput maris thoracis magnitudine, feminae eodem paulo minus, sat crebre punctatum, lateribus (in mare magis quam in femina) longitudinaliter strigosum, linea postica transversali subarcuata fortius impressa, clypei spatio utrinque foveolato, in mare depresso et apice spinulis duabus incumbentibus, haud ultra oris partes porrectis, minus conspicuis armato. Thorax coleopterorum prope latitudine, latitudine antica duplo brevior, lateribus, basi angulisque posterioribus eodem arcu rotundatis, leviter convexus, parce subtilissime punctulatus, canalicula media obsoletissima, vix observanda. Elytra thorace vix longiora, subtilissime punctulata, obscure fusco-picea, nitida. Pedes picei, tibiis tarsisque testaceis.

Abdominis segmentum inferius septimum maris apice truncatum, feminae leviter triangulariter productum.

### **Oxytelus Grav.**

*Mannerh. Brachelytr. p. 47.*

#### *Paraglossae liberae.*

*Pedes intermedii basi distantes.*

*Tibiae anticae externe subtiliter uniseriatim spinulosae.*

Corpus elongatum, lineare, depressum, alatum. Caput porrectum, basi constrictum, collo quasi brevi crassiusculo thoraci immissum, oculis ovalibus, leviter prominulis. Labrum (t. V. f. 17. c.) parvum, transversum, apice leviter sinuatum, laciniis membraneis triangularibus, acuminatis. Mandibulae breviusculae, plerumque parum prominentes, apice obtuse bidentatae, interne membrana apice barbata fimbriatae. Maxillae (t. V. f. 17. b.) malis coriaceis, interiore interne membranea, apicem versus barbata, exteriori longiore, apice dense barbata. Palpi maxillares parum elongati, malam exteriori vix superantes, articulo primo brevi, secundo tertio paulo longiore tertioque teretibus, quarto minuto, subulato. Labium (t. V. f. 17. a.) mento transversum, ligula membranea, lata, apice leviter bisinuata, paraglossis ligulam superantibus, liberis, acuminatis, introrsum falcatis, interne dense ciliatis. Palpi labiales articulo primo cylindrico, secundo hoc paulo longiore et tenuiore, tereti, tertio breviori, acuminato. Antennae subrefractae, apicem versus leniter incrassatae. Thorax transversus, basin versus leviter angustatus, basi subrotundatus, apice truncatus. Scutellum triangulare. Elytra brevia, truncata. Abdomen lineare, marginatum. Pedes breviusculi, coxis intermediis distantibus, tibiis externe spinulis uniseriatis ciliatae, anticis prope apicem excisis, tarsis triarticulatis, articulo primo leviter elongato, tertio reliquis duobus longitudine aequalibus.

#### **Observationes.**

1 Thorax in omnibus fere speciebus dorso longitudinaliter trisulcatus, sulco intermedio recto, lateralibus huic approximatis leviter arcuatis. Caput, thorax et elytra saepius longitudinaliter rugulosa, glabra. Abdomen parce subtiliter pubescens.

2. Sexus differentia in abdominis apice inferiore sita apud species singulas illustranda. Mares saepius capite maiore insignes; nonnunquam caput speciminibus masculis maioribus maius, minoribus minus et femineo subaequale.

3. Victus in stercore.

## \* Thorax margine laterali crenulato.

X

- \* 1. *O. rugosus*: Niger, parum nitidus, pedibus rufo-piceis, thorace trisulcato, margine crenato, fronte punctato-rugosa, antice impressa laevique. — Long.  $2\frac{1}{4}$  lin.

Er. Col. March. I. 588. 1.

*Staph. rugosus* Fabr. Syst. Ent. 267. 19. Spec. Ins. I. 338. 24. Mant. Ins. I. 222. 34. Ent. Syst. I. II. 530. 54. Syst. El. II. 601. 66. — Goeze Ent. Beitr. I. 724. 10. — Panz. Ent. Germ. 358. 43. — Marsh. Ent. Britt. 506. 24.

*Staph. carinatus* Panz. Faun. Germ. 57. 24.

*Oxyt. carinatus* Grav. Micr. 106. 6. Mon. 187. 5. a. — Latr. Hist. nat. Crust. et Ins. IX. 363. 6. Gen. Crust. et Ins. I. 295. 1. — Oliv. Encycl. meth. VIII. 613. 5. — Gyll. Ins. Suec. II. 452. 8. — Sahlb. Ins. Fenn. I. 413. 10. — Zetterst. Faun. Lappon. I. 92. 1. Ins. Lappon. 72. 1. — Mannerh. Brachelytr. 47. 1. — Boisd. et Lacord. Faun. Ent. Paris. I. 463. 3. — Runde Brachelytr. Hal. 20. 1.

*Staph. piceus* Oliv. Ent. III. 42. 20. 23. t. 3. f. 30.

*Staph. piceus* var.  $\beta$  Payk. Mon. Staph. 20. 12. Faun. Suec. III. 385. 22.

*Staph. piceus* var. 1 Walk. Faun. Paris. I. 270. 9.

*Staph. striatus* Ström Act. Nidros. IV. 35. — Müll. Prod. 99. 1122. — Goeze Ent. Beitr. I. 728. 34.

*Staph. sulcatus* Fourcr. Ent. Par. I. 168. 16.

Le *Staph. noir à corcelet noir et silloné et bordé* Geoffr. Ins. I. 367. 16.

Var. Thorace piceo, antennis elytrisque rufis.

*Oxyt. pulcher* Grav. Micr. 107. 7. — Latr. Hist. nat. Crust. et Ins. IX. 364. 7.

*Oxyt. terrestris* Boisd. et Lacord. Faun. Ent. Paris. I. 462. 2.

*Staph. laqueatus* Marsh. Ent. Britt. 513. 45.

Habitat in Europa.

Niger vel nigro-piceus, nitidus. Antennae capite sesqui longiores, articulo tertio secundo subaequali, 5 — 10 sensim paulo crassioribus, nigrae. Palpi nigro-picei. Mandibulae piceae. Caput maris thoracis magnitudine, orbiculatum, feminae duplo fere minus, oblongum, creberrime punctatum, fronte subdepressa, posterius utrinque longitudinaliter obsolete impressa, utrinque dense strigosa, clypei spatio laevi, opaco. Thorax coleopteris paulo angustior, latitudine plus dimidio brevior, basin versus modice angustatus, lateribus, basi angulisque posterioribus rotundatis, subdepressus, dense sat profunde rugoso-punctatus, dorso trisulcatus, sulcis exterioribus antice, intermedio postice angustatis, lateribus depressus, margine laterali subtiliter crenato. Elytra thorace parum longiora, dense profundeque punctata, longitudinaliter rugulosa. Abdomen subtilissime punctulatum. Pedes rufo-picei.

Abdominis segnentum inferius quintum maris apice medio tuberculo minuto valde prominulo, acutiusculo armatum; sextum apice medio subtridenticulato, septimum profunde sinuatum, trilobum, lobo intermedio brevior; feminae segmentum septimum apice profunde sinuatum, lobo intermedio triangulari, lateralibus paulo longiora.

Variat minus maturus colore dilutiore, nonnunquam elytris rufo-piceis, saepius totus magis minusve piceo-testaceus. — Varietatem punctis et rugis multo minus confertis, corpore nitido, insignem communicavit Dom. Märkel.

- \* 2. *O. insectatus*: Niger, nitidus, elytris pedibusque rufis, thorace trisulcato, fronte parcius, antice subtilissime punctata, thoracis margine crenato. — Long. 2 lin.

Grav. Mon. 189. 5. e. — Er. Col. March. I. 589. 2.

*Oxyt. carinatus* var. b Gyll. Ins. Suec. II. 453. 8.

Habitat in Europa.

Praecedente minor, praesertim angustior, niger, nitidus. Antennae capite thoraceque breviores, articulo secundo tertioque subaequalibus, 4 — 10 sensim crassioribus, penultimis vix transversis, nigrae, articulo primo nigropiceo, 2 — 4 rufo-piceis. Os cum palpis piceo-rufum. Caput maris thorace maius, feminae dimidio minus, sparsim profundius punctatum, fronte utrinque inter oculos subtilissime strigosa, clypei spatio haud depresso, parce subtiliter punctato, nitido, utrinque impresso. Thorax coleopteris latitudine aequalis, latitudine dimidio brevior, lateribus leviter rotundatus, basin versus leviter angustatus, basi truncatus, angulis posterioribus obtusis, subdepressus, sat fortiter denseque punctatus, subrugulosus, dorso trisulcatus, sulco intermedio postice, exterioribus antice obsolescentibus, margine laterali dense crenulato. Elytra thorace haud longiora, sat crebre profundeque punctata, rufa, circa scutellum infuscata. Abdomen obsolete punctulatum, apice rufo-piceum. Pedes rufi.

Abdominis segmentum inferius sextum maris tuberculo oblongo minimo obsoleteque medio instructum, septimo apice profunde bisinuato, lobo intermedio lateralibus parum brevior, feminae apice medio leviter triangulariter producto.

- \* 3. *O. fulvipes*: Niger, nitidus, antennarum basi pedibusque testaceis, thorace trisulcato, fronte parce punctata, antice laevi, thorace brevi, margine obsolete crenato. — Long. 2 lin.

Er. Col. March. I. 590. 3.

Habitat in Germania.

*O. rugoso* paulo minor, niger, nitidus. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo brevior, 4 — 11 sensim paululum crassioribus, nigrae, articulis 4 primis rufis. Os rufum. Caput maris thoracis fere magnitudine, feminae paulo minus, fronte parcius sat profunde punctata, clypei spatio laevi, nitido, utrinque impresso. Thorax coleopteris paulo angustior, latitudine duplo brevior, basin versus angustatus, lateribus anterieus fortiter rotundatus, basi truncatus, angulis posticis obtusis, subdepressus, sat profunde parce punctatus, dorso trisulcatus, sulcis integris, exterioribus subarcuatis, utrinque late impressus, impressione crebre punctata, margine laterali obsolete crenulato. Elytra thorace paulo longiora, sat profunde minus dense punctata, hinc inde longitudinaliter rugulosa. Abdomen vix punctulatum, apice piceum. Pedes rufo-testacei.

Abdominis segmentum inferius septimum maris apice bisinuatum, lobo intermedio longiore, apice emarginato, feminae apice rotundatum.



\*\*\* Thorax margine laterali integerrimo.

- \* 4. *O. piceus*: Niger, nitidus, antennarum basi, elytris pedibusque testaceis, thorace trisulcato, fronte punctata, posterius canaliculata, antice impressa laevique. — Long.  $2\frac{1}{4}$  lin.

Grav. Micr. 105. 5. Mon. 190. 6. — Latr. Hist. nat. Crust. et Ins. IX. 363. 5. t. 80. f. 4. — Oliv. Encycl. meth. VIII. 614. 6. — Gyll. Ins. Suec. II. 454. 9. — Sahlb. Ins. Fenn. I. 414. 11. — Zetterst. Faun. Lappon. I. 93. 2. Ins. Lapp. 73. 2. — Mannerh. Brachelytr. 47. 2. — Boisd. et Lacord. Faun. Ent. Paris. I. 461. 1. — Runde Brachelytr. Hal. 20. 2. — Er. Col. March. I. 590. 4.

*Staph. piceus* Linn. Syst. Nat. I. II. 686. 25. — Fab. Syst. Ent. 267. 20. Spec. Ins. I. 338. 25. Mant. Ins. I. 222. 35. Ent. Syst. I. II. 530. 55. Syst. El. II. 601. 67. — Goeze Ent. Beitr. I. 722. 25. — De Vill. Ent. I. 418. 25. — Payk. Mon. Staph. 20. 12. Faun. Suec. III. 384. 22. — Panz. Ent. Germ. 358. 44. Faun. Germ. 27. 14. — Marsh. Ent. Brit. 500. 23. — Walk. Faun. Paris. I. 270. 9.

*Staph. sulcatus* Müll. Prodr. 97. 1094.

*Staph. rugosus* Schrank Faun. Boic. I. 648. 886.

Habitat in Europa; in Syria, Dom. Prof. Ehrenberg.

Statura omnino *O. rugosi*, niger, nitidus. Antennae capite thoraceque breviores, articulo tertio secundo paulo brevior, ab articulo quinto inde abrupte incrassatae, articulis 6—10 fere transversis, 4 primis testaceis, reliquis fuscis, ultimo apice testaceo. Palpi testacei. Caput maris thorace sesqui maius, suborbiculatum, feminae dimidio minus, oblongum, crebrius subtiliusque punctatum, fronte inter oculos leviter canaliculata, clypei spatio subimpresso, laevi, nitido, margine exteriore elevato. Thorax coleopteris parum angustior, latitudine in mare duplo, in femina plus dimidio brevior, basin versus subangustatus, lateribus leviter rotundatus, angulis posterioribus obtusis, subdepressus, minus fortiter inaequaliter punctatus, dorso trisulcatus, sulcis exterioribus subarcuatis, antice obsolescentibus, intermedio integro, utrinque obsolete impressus, margine laterali integerrimo. Elytra thorace sesqui longiora, dense punctata, subtiliter longitudinaliter rugulosa, testacea. Abdomen laeve. Pedes flavi.

Abdominis segmentum inferius sextus maris apice bisinuat, lobo intermedio longiore, apice truncato, septimo bis profunde inciso, lobis lateribus acuminatis, intermedio paulo brevior, apice tricuspi, feminae segmento septimo apice bisinuato, lobo intermedio lato, leviter triangulariter producto.

Variat minus maturus corpore magis minusve pallide testaceo, capite saepius obscuriore.

- \* 5. *O. sculptus*: Niger, nitidus, antennarum basi pedibusque testaceis, thoracis disco elytris piceis, thorace trisulcato, fronte crebre punctata, canaliculata, antice impressa laevique. — Long. 2 lin.

Grav. Mon. p. 191. — Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 460. 1. — Er. Col. March. I. 591. 5.

*Oxyt. longicornis* Mannerh. Brachelytr. 48. 3. — Sahlb. Ins. Fenn. I. 415. 12.

Habitat in Europa; in Aegypto Dom. Prof. Ehrenberg, in America septentrionali, Knoch.

Praecedente parum minor, niger, nitidus. Antennae capitis thoracisque fere longitudine, articulo tertio secundo paulo longiore, sequentibus sensim paulo crassioribus, haud transversis, nigrae, articulis tribus primis testaceis,

ultimo apice ferrugineo. Os testaceum. Caput in utroque sexu thorace minus et angustius, oblongum, fronte parcius subtiliusque punctata, medio canaliculata, clypei spatio depresso, laevi, opaco, margine laterali elevato, rufo-testaceo. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus anterieus fortius rotundatis, angulis posterioribus obtusis, subdepressus, minus fortiter punctatus, dorso trisulcatus, sulcis exterioribus subarcuatis, intermedio integro, lateribus impressus subtiliterque longitudinaliter rugulosus, disco plerumque rufo-piceo. Elytra thorace tertia parte longiora, sat crebre punctata, subtiliter confertim longitudinaliter rugulosa, fusco-picea. Abdomen supra laeve, subtus punctulatum, apice piceum. Pedes flavo-testacei.

Abdominis segmentum inferius septimum maris bis profunde incisum, lobo intermedio lateralibus aequali, apice emarginato, feminae apice leviter bisinuato.

Variat minus maturus corpore testaceo, capite obscuriore, antennis praeter basin nigris.

6. *O. Bengalensis*: *Testaceus, parum nitidus, fronte nigro-picea, leviter trisulcata, thorace rufo-testaceo, dorso trisulcato.* — Long.  $2\frac{1}{4}$  lin.

Habitat in Bengala, Mus. Dom. Hope.

Praecedente paulo maior. Antennae capite thoraceque parum breviores, articulo tertio secundo sesqui longiore, 6—10 leviter transversis, fuscae, articulis 5 primis testaceis. Palpi testacei. Caput thoracis prope latitudine, fronte depressa, impunctata, postice sulco transverso profundiore distincta, inter oculos longitudinaliter obsolete trisulcata, clypei spatio leviter circulariter impresso, margine anteriore elevato, antice truncato, nigricans, margine anteriore testaceo. Thorax coleopteris vix angustior, latitudine plus dimidio brevior, lateribus rotundatus, basin versus angustatus, angulis posterioribus obtusis, planiusculus, obsolete longitudinaliter strigosus, dorso trisulcatus, sulcis exterioribus leviter flexuosis, intermedio obsolete, rufo-testaceus, opacus, margine summo laterali et basali nigricante. Elytra thorace parum longiora, plana, subtiliter longitudinaliter strigosa, subtiliter interpunctata, lateribus subtiliter marginata, sutura leviter elevata, testacea, opaca, circa suturam et prope angulum apicalem anteriorem leviter infuscata. Abdomen impunctatum, rufo-testaceum, opacum. Pectus rufo-testaceum. Pedes testacei.

- \* 7. *O. collaris*: *Niger, pedibus testaceis, femoribus tibiisque medio piceis, thorace obsolete trisulcato, inaequali, rufo, fronte apice elongata emarginataque.* — Long.  $2\frac{1}{3}$  lin.

Habitat in Terra Van-Diemenii, Dom. Schayer.

Corpus nigrum, nitidulum, thorace rufo. Antennae capite sesqui longiores, articulo tertio secundo paulo longiore, ab articulo quinto inde incrassatae, articulis 6—10 transversis, nigrae, articulis 3 primis basi rufis. Palpi nigri. Caput thoracis prope latitudine, suborbiculatum, fronte depressa, confertim punctata, intricato-rugulosa, inter oculos obsolete longitudinaliter bi-impressa, apice leviter producto, angustato, apice emarginato. Thorax coleopteris vix angustior, latitudine duplo brevior, lateribus rotundatus, ba-

sin versus angustatus, angulis posterioribus obtusis, dense inaequaliter punctatus, dorso obsolete trisulcatus, utrinque fortius late impressus. Elytra thorace paulo longiora, fortius inaequaliter punctata, longitudinaliter subrugulosa, impressionibus nonnullis parum profundis leviter inaequalia. Abdomen sparsim subtilius punctatum. Pedes testacei, femoribus dilutioribus tibiisque medio fuscis.

Abdominis segmentum inferius septimum maris apice bisinuatum, lobo intermedio parvo, leviter prominulo, apice emarginato, testaceo. Femina latet.

- \* 8. *O. Caffer: Niger, parum nitidus, tibiis tarsisque piceis, thorace bisulcato, fronte strigosa, antice subimpressa laevique, apice bisinuata. — Long. 2 lin.*

Habitat ad Promont. Bonae Spei, Dom. Prof. Lichtenstein.

Statura omnino et summa affinitas *O. sculpturati*. Corpus nigrum, minus nitidum. Antennae capite vix longiores, articulo tertio secundo prope longiore, apicem versus sensim incrassatae, articulis 6 — 10 fortiter transversis, totae nigrae. Mandibulae et palpi picei. Caput thoracis latitudine, suborbiculatum, fronte posteriori subtiliter strigosa, utrinque obsolete longitudinaliter impressa, medio posteriori leviter canaliculata, clypei spatio subdepresso, laevi, lateribus subtiliter acute marginata, apice profundius bisinuata. Thorax coleopterorum latitudine, latitudine duplo brevior, lateribus leviter rotundatus, basin versus subangustatus, angulis posterioribus fere rectis, subdepressus, dense longitudinaliter rugulosus, dorso trisulcatus, interstitiis laevioribus, sulco intermedio posteriori angustato, exterioribus subrectis, utrinque late profundius impressus. Elytra thorace paulo longiora, subtiliter punctata, dense strigosa, rugulosa. Abdomen parce subtiliter punctatum. Pedes picei, coxis femoribusque nigricantibus.

Abdominis segmentum inferius septimum maris apice late parum profunde sinuatum. Femina latet.

Obs. Ab *O. sculpturato* imprimis fronte apice bisinuata, lateribus marginata et abdomine maris haud tuberculato distinctus.

- \* 9. *O. sculpturatus: Niger, parum nitidus, pedibus testaceis, thorace trisulcato, fronte confertim strigosa, antice impressa laevioreque, apice rotundata. — Long. 2 lin.*

*Grav. Mon.* 187. 5. 6. — *Gyll. Ins. Suec.* II. 456. 10. — *Sahlb. Ins. Fenn.* I. 416. 13. — *Mannerh. Brachelytr.* 48. 4. — *Runde Brachelytr. Hal.* 20. 3. — *Er. Col. Marck.* I. 592. 6.

*Oxyt. flavipes* Boisd. et Lacord. *Faun. Ent. Paris.* I. 464. 4.

Habitat in Europa; ad Promont. Bonae Spei, Dom. Prof. Lichtenstein.

Statura fere *O. picei*, et parum minor, niger, parum nitidus. Antennae capite paulo longiores, articulo tertio secundo longitudine subaequali, apicem versus fortius incrassatae, articulis 6 — 10 transversis, nigrae. Palpi picei. Caput maris iam thorace paulo maius, suborbiculatum, iam ut in femina paulo minus, punctulatum, confertissime subtiliter longitudinaliter strigosum, fronte medio foveolata, clypei spatio subdepresso, utrinque sulcato, laeviore, maris nitidiore, feminae opaco, apice rotundato, subtiliter marginato. Thorax coleopteris parum angustior, latitudine plus dimidio brevior, lateribus



leviter rotundatus, basin versus parum angustatus, angulis posterioribus subrectis, subdepressus, punctulatus, confertim subtiliter longitudinaliter rugulosus vel strigosus, dorso trisulcatus, sulco intermedio apicem versus attenuato, exterioribus parum arcuatis, utrinque late leviter impressis. Elytra thorace paulo longiora, confertissime subtiliter strigosa, punctulata, nigropicea, opaca. Abdomen parce subtiliter punctulatum. Pedes testacei, coxis femoribusque saepe piceis.

Abdominis segmentum inferius sextum maris apice tuberculis duobus minutis testaceis munitum, septimum late profundeque sinuatum, feminae apice medio parum triangulariter producto, utrinque leviter sinuato.

\* 10. *O. inustus*: *Niger, nitidus, mandibulis rufis, pedibus testaceis, thorace obsolete trisulcato, capite parce subtiliterque punctato.* — Long. 2 lin.

*Grav. Mon.* 188. 5. c. — *Er. Col. March.* 1. 593. 7.

Habitat in Europa.

Habitu omnino *O. picei*, ad dimidio minor, niger, nitidus. Antennae capite sesqui longiores, articulo secundo tertio longitudine subaequali, ab articulo quinto inde modice incrassatae, articulis 6 — 10 transversis, totae nigrae. Mandibulae rufae. Palpi picei. Caput maris thorace paulo maius, feminae paulo minus, parce subtiliter punctatum, clypei spatio laevi, aequè nitido, utrinque obsolete impressum, antice rotundatum. Thorax coleopteris parum angustior, latitudine plus dimidio brevior, basin versus subangustatus, lateribus anterioribus leviter rotundatus, angulis posterioribus obtusis, subdepressus, minus fortiter punctatus, lateribus parcius longitudinaliter subtiliter strigosus, dorso trisulcatus, sulco intermedio distinctiore, basin versus attenuato, exterioribus obsoletioribus, parum arcuatis, utrinque leviter longitudinaliter impressus, margine laterali quasi elevato-plicato. Elytra thorace vix longiora, crebre punctata, longitudinaliter rugulosa, picea. Abdomen parce subtiliterque punctatum, subtiliter pubescens. Pedes testacei, femoribus plerumque medio piceis.

Abdominis segmentum inferius sextum maris apice tuberculis duobus minutis concoloribus munitum, septimum late profundeque sinuatum; feminae apice medio parum triangulariter productum, utrinque leviter sinuatum.

\* 11. *O. incolumis*: *Niger, nitidus, elytris pedibusque luteis, thorace aequali.* — Long. 2 lin.

Habitat in Carolina meridionali, Dom. Zimmermann.

Niger, nitidus. Antennae capite paulo longiores, articulo tertio secundo vix longiore, ab articulo quinto inde incrassatae, articulis 6 — 10 transversis, ultimo oblongo, piceae, basi rufae. Palpi rufi. Os rufo-piceum. Caput maris iam thoracis latitudine, iam ut in femina eodem dimidio angustius, fronte parcius subtiliter punctata, clypei spatio late excavato, apice obsolete bisinuato. Thorax coleopteris haud angustior, latitudine duplo brevior, lateribus parum rotundatus, basin versus vix angustatus, angulis posterioribus obtusis, leviter convexus, crebrius subtiliter punctatus, aequalis. Elytra thoracis longitudine, subtilius punctata, subtiliter longitudinaliter rugulosa, dilute lutea. Abdomen parce subtiliterque punctatum, parce pube-

scens, segmentorum singulorum margine anoque piceo-testaceis. Pedes luteo-testacei.

Abdominis segmentum inferius sextum maris apice leviter bisinuatum, septimum profunde bisinuatum, lobo intermedio longiore, apice truncato; feminae medio leviter triangulariter productum.

- \* 12. *O. luteipennis*: *Niger, nitidus, antennarum basi, elytris pedibusque testaceis, thorace trisulcato, fronte punctata, bisulcata, antice impressa laevique.* — Long. 2 lin.

*Er. Col. March* I. 593. 8.

Habitat in Germania et in montibus Uralensibus.

Habitu omnino *O. picei*, at paulo minor, niger, nitidus. Antennae capite sesqui longiores, articulo secundo tertioque subaequalibus, ab articulo quinto inde crassiusculae, 6—10 transversis, apicem versus fortius incrassatae, nigrae, articulis 5 primis testaceis. Mandibulae rufo-piceae. Palpi picei. Caput maris thoracis latitudine vel individuis minoribus eodem angustius, feminae thorace dimidio angustius, suborbiculatum, crebrius punctatum, fronte inter oculos bisulcata, clypei spatio impresso, laevi, aequè nitido, margine exteriori elevato, antice utrinque denticulo terminato. Thorax coleopteris parum angustior, latitudine duplo fere brevior, lateribus modice rotundatus, basin versus leviter angustatus, angulis posterioribus obtusis, subdepressus, sat crebre punctatus, dorso trisulcatus, sulcis omnibus sat profundis, integris, exterioribus leviter flexuosis, utrinque longitudinaliter leviter impressus, margine laterali integerrimo. Elytra thorace paulo longiora, sat crebre punctata, subtiliter longitudinaliter rugulosa, luteo-testacea. Abdomen laeve. Pedes flavi,

Abdominis segmentum inferius sextum maris apice tuberculis duobus minutis rotundatis piceo-testaceis instructum, septimum profunde bisinuatum, lobo intermedio in mucronem sat elongatum, apice impressum elongato; feminae segmentum septimum apice bisinuatum, lobo intermedio rotundato, laterales vix superante.

Obs *O. piceo* simillimus, differt autem praecipue fronte parcius punctata, inter oculos bisulcata, clypei spatio fortius impresso, apice bidenticulato, et mas abdominis segmentis inferioribus ultimis aliter formatis.

- \* 13. *O. Pensylvanicus*: *Piceus, antennarum basi, pedibus elytrisque testaceis, thorace trisulcato, fronte crebre subtilius punctata, antice impressa laevique.* — Long. 2 lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Statura omnino praecedentis. Antennae capite sesqui longiores, articulo tertio secundo brevior, ab articulo quinto inde incrassatae, articulis 6—10 transversis, ultimo oblongo, nigrae, articulis 4 primis testaceis. Os testaceum. Caput maris thoracis latitudine, orbiculatum, feminae thorace dimidio angustius, oblongum, fronte crebrius minus fortiter punctata, passim obsolete longitudinaliter rugulosa, clypei spatio impresso, laevi, nitido, margine exteriori elevato, antico maris leviter bisinuato, feminae rotundato, nigrum, nitidum. Thorax coleopterorum prope latitudine, latitudine plus dimidio brevior, basin versus parum angustatus, lateribus leviter rotundatus, angulis

posterioribus obtusis, subdepressus, crebrius punctatus, dorso trisulcatus, sulco intermedio integro, exterioribus leviter arcuatis, obsoletioribus, utrinque evanescentibus, lateribus late impressus, piceus, nitidus. Elytra thorace paulo longiora, longitudinaliter rugulosa, parcius subtiliusque punctata, lutea. Abdomen laeve, supra nigro-piceum, infra testaceum, segmentis singulis basi piceis. Pectus nigro-piceum. Pedes flavi.

Abdominis segmentum inferius septimum maris medio tuberculatum, apice profunde bisinuatum, lobo intermedio laterales vix superante, apice emarginato, feminae apice leviter triangulariter productum.

\*14. *O. Madagascariensis*: Testaceus, elytris pedibusque luteis, capite thoraceque supra nigro-piceis, hoc trisulcato, fronte subtiliter punctata, longitudinaliter strigosa, antice laevi, maris apice producta emarginataque. — Long. 2 lin.

Habitat in Madagascar, Dom. Goudot.

Statura omnino *O. insigniti*, at maximis eius individuis paulo maior. Antennae capite sesqui longiores, articulo tertio secundo longitudine subaequali, ab articulo quinto inde incrassatae, articulis 6 — 10 transversis, totae testaceae. Os testaceum. Caput maris thoracis latitudine, suborbiculatum, feminae thorace dimidio angustius, fronte inter oculos punctata, densius longitudinaliter rugulosa, clypei spatio depresso, laevi, minus nitido, margine laterali elevato, antico maris leviter producto, apice emarginato, feminae truncato, infra piceo-testaceum, supra piceo-nigricans, nitidulum. Thorax coleopteris parum angustior, latitudine plus dimidio brevior, basin versus modice angustatus, lateribus leviter rotundatis, angulis posterioribus obtusis, subdepressus, minus dense subtilius punctatus, lateribus late confertim fortius longitudinaliter rugulosus, dorso trisulcatus, sulco intermedio integro, exterioribus subarcuatis, subintegris, lateribus late per totam fere longitudinem impressus, supra nigro-piceus, nitidulus, infra testaceus. Elytra thorace paulo longiora, densius punctata, dense longitudinaliter rugulosa, lutea. Abdomen laeve, testaceum. Pectus piceo-testaceum. Pedes lutei.

Abdominis segmentum inferius septimum maris apice late emarginatum, feminae leviter productum.

\*15. *O. insignitus*: Niger, nitidus, pedibus flavis, ore, antennarum basi elytrisque testaceis, thorace piceo-rufo, trisulcato, capite laevi, fronte posterius lateribus strigosa, antice utrinque foveolata, maris apice acuminata. — Long.  $1\frac{1}{3}$  —  $1\frac{2}{3}$  lin.

Grav. Mon. 188. 5. d.

Oxyt. Americanus Mannerh. Brachelytr. 48. 5.

Habitat in America: in America septentrionali, Knoch, Com. Dejean, Dom. Zimmermann, in ins. St. Thomae, in Columbia, Dom. Moritz, in Brasilia, Dom. Thorey.

Praecedentibus minor. Antennae capite paulo longiores, articulo tertio secundo paulo longiore, ab articulo quinto inde incrassatae, articulis penultimis tribus fortius transversis, fuscae, articulo primo testaceo. Palpi testacei. Mandibulae rufo-piceae; maris elongatae, acutae, decussatae. Caput maris plerumque thoracis latitudine, feminae angustius, nigrum, nitidum,



impunctatum, fronte utrinque leviter impressa, lateribus dense longitudinaliter strigosa, clypei spatio haud distincto, utrinque foveola profundiore impresso, margine antico in mare bisinuato, medio producto, acuminato, in femina truncato. Thorax coleopteris parum angustior, latitudine duplo fere brevior, basin versus in mare fortius quam in femina angustatus, lateribus, angulis posterioribus basi que rotundatis, apice late leviter emarginatus, subdepressus, rufo-piceus, subnitidus, crebre punctatus, dorso trisulcatus, sulcis sat profundis, integris, utrinque fortius longitudinaliter impressus. Scutellum piceum. Elytra thorace paulo longiora, crebre punctata, obsolete longitudinaliter rugulosa, dilute testacea, parum nitida. Abdomen laeve, nigrum, parum nitidum, ano piceo-testaceo. Pectus nigrum. Pedes flavi.

Abdominis segmentum inferius septimum maris apice late et parum profunde sinuatum, feminae rotundatum.

Obs. Mas mandibulis elongatis capitisque margine antico medio acuminato prae reliquis huius generis speciebus insignis.

- \* 16. *O. politus*: *Niger, nitidus, pedibus testaceis, elytris fuscis, thorace trisulcato, sulcis lateralibus obsoletis, capiteque parce subtilissime punctatis.* — Long.  $1\frac{1}{4}$  lin.

Habitat in Austria, Dom. Schüppel.

Statura omnino *O. nitiduli*, at paulo minus depressus, niger, pernitidus. Antennae articulis tribus nigro-piceis, tertio secundo longitudine subaequali (reliqui desunt). Palpi testacei. Caput thorace paulo angustius, parce subtilissime punctatum, interstitiis politis, anterius utrinque profundius longitudinaliter impressum, fronte antice et lateribus subtilissime marginata. Thorax coleopteris paulo angustior, latitudine plus sesqui brevior, basin versus leviter angustatus, lateribus modice, basi levissime rotundatis, angulis posterioribus obtusis, apice utrinque subsinuatus, angulis anterioribus acutiusculis, subdepressus, parce subtiliterque punctatus, interstitiis medio laevissimis, versus latera subtilissime longitudinaliter strigosus, dorso trisulcatus, sulco intermedio profundiore, integro, exterioribus obsoletioribus, leviter arcuatis, utrinque profundius longitudinaliter impressus. Elytra thorace plus sesqui longiora, parcius subtiliter punctata, parce subtiliterque longitudinaliter strigosa, fusco-picea, nitida. Abdomen laevissimum. Pedes testacei.

Abdominis segmentum inferius septimum feminae apice medio triangulariter productum. Mas latet.

- \* 17. *O. intricatus*: *Niger, supra parum nitidus, elytrorum disco pedibusque piceo-testaceis, capite thoraceque confertissime punctato-rugosis, hoc trisulcato, fronte antice subdepressa laevioreque.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Austria, Dom. Schüppel, Smyrnae, Dr. Helfer.

Statura fere *O. nitiduli*, niger, subtus nitidus, supra subopacus. Antennae capite paulo longiores, articulo secundo tertio tenuiore, at vix brevior, apicem versus sensim leviter incrassatae, articulis 7 — 10 transversis, nigrae. Mandibulae et palpi picei. Caput maris thoracis magnitudine, feminae paulo minus, confertissime rugoso-punctatum, fronte medio obsolete canaliculata, clypei spatio impresso, utrinque sulcato, laevi, subnitido. Thorax coleopterorum prope latitudine, latitudine duplo prope brevior, basin versus suban-

gustatus, lateribus et basi leviter rotundatus, antice late leviter emarginatus, angulis anterioribus acutis, posterioribus obtusis, subrotundatis, subdepressus, confertim rugoso-punctatus, dorso trisulcatus, sulcis omnibus obsoletioribus, utrinque late leviter impressus. Elytra thorace paulo longiora, depressa, confertim punctata, dense longitudinaliter rugulosa, fusca, disco fusco-testacea. Abdomen supra densius subtiliter punctatum. Pedes piceo-testacei.

Abdominis segmentum inferius septimum maris late emarginatum, feminae apice medio leviter productum.

Obs. *O. nitidulo* magnitudine aequalis, at paulo latior, punctura multo crebriore, thoracis sulcis omnibus obsoletis, corpore supra fere opaco distinctus.

\* 18. *O. nitidulus*: *Niger, nitidulus, elytris piceis, pedibus testaceis, capite thoraceque crebrius punctatis, subtiliter rugulosis, hoc trisulcato, fronte antice subimpressa, laevi.* — Long. 1 — 1 $\frac{1}{3}$  lin.

Grav. Micr. 107. 8. Mon. 186. 4. — Latr. Hist. nat. Crust. et Ins. IX. 364. 8. — Oliv. Encycl. meth. VIII. 613. 4. — Panz. Faun. Germ. 104. 13. — Gyll. Ins. Suec. II. 458. 12. — Sahlb. Ins. Fenn. I. 416. 15. — Zetterst. Faun. Lappon. I. 93. 3. Ins. Lappon. 73. 3. — Mannerh. Brachelytr. 48. 7. — Boisd. et Lacord. Faun. Ent. Paris. I. 464. 5. — Runde Brachelytr. Hal. 20. 5. — Er. Col. March. I. 594. 9.

*Staph. piceus* Schrank Enum. Ins. Austr. 236. 449. Faun. Boic. I. 647. 885.

*Staph. piceus, var. pygmaea* Payk. Faun. Suec. III. p. 384.

*Oxyt. rugulosus* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 460. 2. — Harris Transact. Nat. Hist. Soc. Hartford I. 69. 6.

Habitat in Europa frequens, in Aegypto, Dom. Prof. Ehrenberg, in America septentrionali, Dom. Zimmermann, in Mexico sec. Say.

Minutus, niger, nitidulus. Antennae capite sesqui longiores, articulo tertio secundo paulo minore, apicem versus sensim subincrassatae, articulis ultimis tribus maioribus, nigrae. Mandibulae et palpi picei. Caput maris thoracis magnitudine, feminae eodem paulo minus, minus confertim punctatum, longitudinaliter strigosum, fronte inter oculos transversim impressa, clypei spatio impresso, laevi, in utroque latere profundius longitudinaliter impresso vel late sulcato. Thorax coleopterorum prope latitudine, latitudine plus dimidio brevior, basin versus leviter angustatus, lateribus et basi leviter rotundatis, angulis posterioribus obtusis, subdepressus, sat crebre fortiterque punctatus, longitudinaliter subrugulosus, dorso trisulcatus, sulco intermedio sat profundo, integro, exterioribus obsoletioribus, subarcuatis, utrinque late impressus, margine laterali integerrimo. Elytra thorace sesqui longiora, creberrime punctata, longitudinaliter substrigosa, fusco-picea. Abdomen supra parce subtilissime punctatum. Pedes testacei.

Abdominis segmentum sextum inferius maris apice leviter emarginatum.

Obs. *Oxyt. pedicellus* Grav. Micr. 102. 2. Mon. 185. 2. — Latr. Hist. nat. Crust. et Ins. IX. 362. 2. — Oliv. Encycl. meth. VIII. 613. 2, mihi omnino dubius, forte ab hac specie haud distinctus, at typo perduto nunquam extricandus.

\* 19. *O. complanatus*: *Niger, opacus, pedibus testaceis, elytris fuscis, his, capite thoraceque subtilissime longitudinaliter strigosis, thorace trisulcato, angulis anterioribus acutiusculis.* — Long. 1 $\frac{2}{3}$  lin.

Er. Col. March. I. 595. 10.

*Oxyt. depressus* Gyll. Ins. Suec. II. 457. 11. — Sahlb. Ins. Fenn. I. 416. 14. — Mannerh. Brachelytr. 48. 5. — Runde Brachelytr. Hal. 20. 4.

Habitat in Europa.

*O. piceo* duplo fere minor, brevisculus, depressus, niger, subnitidus, capite, thorace elytrisque opacis. Antennae capite paulo longiores, articulo tertio secundo subaequali, apicem versus sensim leniter incrassatae, articulis 6 — 10 transversis, nigrae. Mandibulae rufo-piceae. Palpi picei. Caput maris thoracis magnitudine, feminae eodem paulo minus, totum subtilissime confertissimeque longitudinaliter strigosum, fronte lateribus subtiliter marginata, posterius subtiliter interpunctata, utrinque longitudinaliter obsolete impressa, clypei spatio parum impresso, margine anteriore rotundato, parum elevato. Thorax coleopteris parum angustior, latitudine duplo prope brevior, basin versus leviter angustatus, lateribus ante medium rotundatis, apice late leviter emarginatus, angulis anterioribus acutis, basi leviter rotundatus, angulis posterioribus obtusis, subdepressus, subtilissime confertissime longitudinaliter strigosus, parce subtiliterque interpunctatus, dorso trisulcatus, sulco intermedio basin versus angustiore, exterioribus subarcuatis, obsoletioribus, utrinque leviter impressus. Elytra thorace paulo longiora, depressa, confertissime subtilissimeque longitudinaliter strigosa, obsoletius subtiliter interpunctata, nigra vel plerumque fusco-nigra, nonnunquam dilutius fusca. Abdomen supra sat dense subtiliter punctatum. Pedes testacei, femoribus interdum medio picescentibus.

Abdominis segmentum inferius sextum maris apice tuberculis duobus approximatis munitum, septimum late emarginatum, feminae apice medio triangulariter productum.

\* 20. *O. depressus*: Niger, opacus, pedibus testaceis, capite, thorace elytrisque omnium subtilissime confertissimeque longitudinaliter strigosis, thorace trisulcato, angulis anterioribus subrotundatis, abdomine confertim punctato. — Long.  $\frac{3}{4}$  — 1 lin.

Grav. Micr. 103. 3. Mon. 185. 3. — Latr. Hist. nat. Crust. et Ins. IX. 362. 3. — Oliv. Encycl. meth. VIII. 613. 3. — Panz. Faun. Germ. 104. 11. — Boisd. et Lacord. Faun. Ent. Paris. I. 465. 6. — Er. Col. March. I. 595. 11.

*Oxyt. pusillus* Mannerh. Brachelytr. 48. 8. — Sahlb. Ins. Fenn. I. 417. 16. — Runde Brachelytr. Hal. 20. 6.

*Staph. tetracarinatus* Block Ins. vall. Plauens. 116. 5. f. 5.

Var. *Elytris fuscis*.

Habitat in Europa.

Minutus, depressus, niger, subnitidus, capite, thorace elytrisque opacis. Antennae capite paulo longiores, articulo tertio secundo paulo tenuiore at vix brevior, apicem versus sensim crassiores, totae nigrae. Mandibulae et palpi picei. Caput maris thoracis magnitudine, feminae eodem paulo minus, confertissime subtilissime longitudinaliter strigosum, fronte utrinque longitudinaliter obsolete impressa, clypei spatio vix depresso, margine anteriore rotundato, leviter elevato. Thorax coleopteris paulo angustior, latitudine dimidio fere brevior, basin versus modice angustatus, lateribus parum rotundatus, apice truncatus, angulis anterioribus subrotundatis, basi leviter rotundatus, angulis posterioribus obtusis, subdepressus, confertissime subtilissimeque longitudinaliter strigosus, dorso trisulcatus, sulcis exterioribus



obsoletioribus, utrinque leviter impressus. Elytra thorace paulo longiora, depressa, confertissime subtilissimeque longitudinaliter strigosa, parce subtiliter obsoleteque interpunctata, nigra, interdum fusca. Abdomen sat dense subtiliter punctatum. Pedes testacei.

Abdominis segmentum inferius septimum maris apice late emarginatum, feminae medio leviter triangulariter productum.

\* 21. *O. pumilus*: *Niger, opacus, pedibus piceis, geniculis tarsisque testaceis, capite, thorace elytrisque subtilissime confertissimeque longitudinaliter strigosis, thorace trisulcato, angulis anterioribus subrotundatis, abdomine crebrius subtiliter punctato. — Long. 1 — 1½ lin.*

*Er. Col. March. I. 596. 12.*

Habitat Berolini, Dom. Schüppel.

Statura et summa affinitate *O. depressi*, at nonnunquam paulo maior, niger, nitidulus, capite, thorace elytrisque opacis. Antennae capite paulo longiores, articulo tertio secundo paululum minore, apicem versus leviter incrassatae, articulis 6 — 10 transversis, ultimis tribus abrupte paulo maioribus. Os piceum. Caput in utroque sexu thorace paulo minus et angustius, subtilissime confertissime longitudinaliter strigosum, fronte depressiuscula, posteriori medio foveola minuta oblonga impressa, clypei spatio subimpresso, antice rotundata et subtiliter marginata. Thorax coleopteris paulo angustior, latitudine plus dimidio brevior, lateribus modice, basi leviter rotundatus, basin versus vix angustatus, angulis posterioribus obtusis, apice truncatus, angulis anterioribus subrotundatis, subdepressus, subtilissime confertissimeque longitudinaliter strigosus, dorso trisulcatus, sulco intermedio profundiore, exterioribus latis, obsolescentibus, subarcuatis, utrinque late leviterque impressus. Elytra thorace sesqui longiora, depressa, subtiliter confertissime longitudinaliter strigosa, subtiliter parce obsoletius interpunctata. Abdomen crebrius punctulatum, nitidum. Pedes nigro-picei, geniculis tarsisque testaceis.

Abdominis segmentum inferius sextum maris late emarginatum, medio spinulis tribus, oblique retrorsum vergentibus, distantibus, anteriore paulo minore, septimum apice medio spinis duabus approximatis, pariter retrorsum directis armatum; feminae segmentum septimum inferius apice medio triangulariter productum.

Obs. Mas abdomine apice infra 5-spinoso sat insignis. Femina ab *O. depresso* praesertim pedibus obscurioribus, thorace paulo latiore, sulcis lateralibus obsoletioribus differt.

\* 22. *O. nanus*: *Niger, opacus, elytris fuscis, pedibus testaceis, femoribus picescentibus, capite thorace elytrisque omnium subtilissime confertissimeque longitudinaliter strigosis, thorace trisulcato, abdomine parce subtilissime punctato, nitido. — Long. ¾ lin.*

Habitat in America septentrionali, Dom. Zimmermann.

Minutus, *O. depresso* aequalis et proximus, niger, nitidus, capite, thorace elytrisque opacis. Antennae capite paulo longiores, articulo tertio secundo multo minore, apicem versus incrassatae, articulis 6 — 10 transversis, ultimis tribus abrupte maioribus, nigrae. Os nigro-piceum. Caput

in utroque sexu thorace paulo angustius et minus, omnium confertissime subtilissimeque longitudinaliter strigosum, fronte utrinque leviter longitudinaliter impressa, posterius medio saepius obsolete sulcata, clypei spatio vix depresso, apice immarginata. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus et basi leviter rotundatus, basin versus vix angustatus, angulis posterioribus obtusis, apice truncatus, angulis anterioribus obtusiusculis, subdepressus, omnium subtilissime confertissimeque longitudinaliter strigosus, dorso distincte trisulcatus, sulcis integris, intermedio profundiore, exterioribus leviter arcuatis, utrinque late levissime impressus. Elytra thorace paulo longiora, depressa, confertissime subtilissimeque longitudinaliter strigosa, parce subtiliter obsoleteque interpunctata. Abdomen parce subtilissime punctatum. Pedes testacei, coxis femoribusque saepius nigro-piceis.

Abdominis segmentum inferius septimum maris apice late parum profunde sinuatum, feminae medio leviter triangulariter productum.

Obs. Ab *O. depresso* imprimis abdomine parce subtilissimeque punctato, ab *O. exiguo* thorace fortius bisulcato, antennis totis nigris, mandibulis palpisque piceis distinctus.

\* 23. *O. exiguus*: *Niger, opacus, ore, antennarum basi pedibusque testaceis, elytris fuscis, capite thoraceque subtilissime longitudinaliter strigosis, thorace dorso obsolete trisulcato.* — Long.  $\frac{3}{4}$  lin.

Habitat in America septentrionali, Dom. Zimmermann.

*O. depresso* proximus et minimis eius individuis aequalis, niger, nitidus, capite, thorace elytrisque opacis. Antennae capite paulo longiores, articulo tertio secundo minore, apicem versus sensim incrassatae, articulis 6 — 10 transversis, ultimis tribus multo maioribus, nigrae, basi testaceae. Os testaceum. Caput thorace paulo angustius, subtilissime confertissimeque longitudinaliter strigosum, fronte antierius utrinque longitudinaliter impressa et inter antennas obsolete transversim sulcata, margine anteriore rotundato, haud elevato. Thorax coleopteris parum angustior, latitudine dimidio brevior, basin versus modice angustatus, lateribus ante medium fortius, basi leviter rotundatus, angulis posterioribus obtusis, apice truncatus, angulis anterioribus subrotundatis, depressus, confertissime subtilissimeque longitudinaliter strigosus, dorso obsolete trisulcatus, sulcis exterioribus subarcuatis, lateribus utrinque leviter impressus. Elytra thorace paulo longiora, depressa, confertissime subtilissimeque longitudinaliter strigosa, parce subtiliter obsoleteque interpunctata, fusca. Abdomen parce subtiliterque punctatum, subtilissime pubescens. Pedes flavo-testacei.

Abdominis segmentum inferius septimum maris apice bisinuatum, feminae late triangulariter productum.

\* 24. *O. minimus*: *Niger, opacus, ore, antennarum basi pedibusque testaceis, capite, thorace elytrisque confertissime subtilissimeque strigosis, thorace dorso fortius trisulcato.* — Long.  $\frac{3}{5}$  lin.

Habitat in Columbia, Dom. Moritz.

*O. depresso* simillimus et minimis eius individuis fere minor, depressus, niger, subnitidus, capite, thorace elytrisque opacis. Antennae capite

sesqui longiores, articulo tertio secundo minore, apicem versus sensim incrassatae, nigrae, basi testaceae. Os testaceum. Caput maris thorace paulo maius, feminae eiusdem magnitudine, confertissime omniumque subtilissime longitudinaliter strigosum, utrinque longitudinaliter impressum, medio inter oculos canalicula brevi impressum, margine anteriore rotundato, haud elevato. Thorax coleopteris paulo angustior, latitudine plus dimidio brevior, basin versus subangustatus, lateribus et basi leviter rotundatus, angulis posterioribus obtusis, apice truncatus, angulis anterioribus subrectis, subdepressus, confertissime omniumque subtilissime longitudinaliter strigosus, dorso profundius trisulcatus, sulcis exterioribus arcuatis, utrinque sat profunde impressus. Elytra thorace paulo longiora, depressa, sparsim subtiliter punctata, confertissime subtilissimeque longitudinaliter strigosa. Abdomen supra parce subtiliter obsoleteque punctulatum. Pedes testacei.

Abdominis segmentum inferius septimum maris apice late emarginatum, feminae medio leviter triangulariter productum.

### **Phloconaeus Er.**

*Er. Col. March. I. p. 597.*

*Aploderus Steph. Illustr. Ent. Britt. V. p. 321.*

*Paraglossae ligulae connatae.*

*Pedes intermediū basi approximati.*

*Tibiae anticae externe subtiliter uniseriatim spinulosae.*

Corpus elongatum, lineare, depressum, alatum. Caput porrectum, basi modice constrictum, collo quasi brevi crasso thoraci immissum, oculis rotundatis, parum prominulis. Labrum minutum, transversum, apice truncatum, laciniis membranaceis triangularibus, acuminatis. Mandibulae leviter prominentes, falcatae, interne ante apicem subtiliter bidentatae, basi membrana apice ciliata fimbriatae. Maxillae (t. V. f. 18. b.) malis membranaceis, interiore interne membranea, apice ciliata, exteriorē paulo longiore, apice dense barbata. Palpi maxillares mala exteriorē paulo longiores, articulo primo brevi, minuto, secundo apicem versus leviter incrassato, tertio huic longitudine subaequali, crassiusculo, quarto minuto, subulato. Labium (t. V. f. 18. a.) mento transversum, ligula membranea, lata, apice leviter bisinuata, lobis lateralibus productis, acutis, paraglossis linearibus, falcatis, acuminatis, liberis, ligulam superantibus, membranaceis, pubescentibus. Palpi labiales articulis tribus longitudine subaequalibus, tertio paululum tenuiore. Antennae subrefractae, longiores, apicem versus vix incrassatae. Thorax transversus, basin versus fortiter angustatus, basi apiceque truncatus. Scutellum breve, subrotundatum. Elytra truncata. Abdomen lineare, marginatum. Pedes breviusculi, coxis intermediis approximatis, tibiis posterioribus externe ciliatis, anticis externe aut setulis ciliatis, ante apicem emarginatis (Ph. caelatus), aut pilis subtilibus ciliatis, apice subangustatis (Ph. caesus), tarsis brevibus, triarticulatis.

### **Observationes.**

1. Habitus omnino generis praecedentis, at thorax sulco medio nullo, ligula uni-sinuata, paraglossae ei connatae, antennae magis elongatae, vix refractae, coxae intermediae haud distantes.



2. Nomen a Dom. Stephensio propositum reiecti, cum perperam compositum, et insuper ineptum sit: thorax nullo modo simplex.

3. Sexus differentia ut in genere praecedente.

4. Victus sub arborum cortice.

\* 1. *Ph. caelatus*: Niger, pedibus testaceis, elytris fuscis, thorace bi-impresso, tibiis anticis apice emarginatis. — Long. 2 lin.

*Er. Col. March. I. 597. 1.*

*Oxyt. caelatus* Grav. *Micr. 103. 4. Mon. 190. 7.* — *Latr. Hist. nat. Crust. et Ins. IX. 363. 4.* — *Oliv. Encycl. meth. VIII. 614. 7.* — *Panz. Faun. Germ. 104. 12.* — *Gyll. Ins. Suec. II. 459. 13.* — *Sahlb. Ins. Fenn. I. 417. 17.* — *Zetterst. Faun. Lappon. I. 93. 4. Ins. Lappon. 73. 4.* — *Mannerh. Brachelytr. 49. 9.* — *Boisd. et Lacord. Faun. Ent. Paris. I. 465. 7.* — *Runde Brachelytr. Hal. 21. 8.*

*Staph. brachypterus* Marsh. *Ent. Brit. 510. 35.*

*Staph. spinipes* Block *Ins. vull. Plauens. 116. 6. f. 6.*

Habitat in Europa.

Niger, subnitidus, pubescens. Antennae in utroque sexu vix capitis thoracisque longitudine, nigrae, basi piceae. Oris partes rufo-piceae. Caput maris thoracis magnitudine vel paulo maius, orbiculatum, feminae thorace paulo minus, oblongum, parcius obsolete punctatum, fronte posterius foveola minuta oblonga obsolete notata, anterieus utrinque ad antennarum basin fortius longitudinaliter impressa. Thorax coleopteris angustior, latitudine plus dimidio brevior, antice emarginatus, angulis anterioribus acutis, lateribus rotundatus, basin versus angustatus, angulis posterioribus cum basi rotundatis, leviter convexus, parcius obsolete punctatus, dorso foveolis duabus arcuatis impressus. Elytra thorace paulo longiora, dense fortius profunde punctata, fusca. Abdomen subtilissime coriaceum, parce subtiliterque punctulatum, apice piceum. Pedes testacei, tibiis anticis margine exteriori spinulis subtilibus ciliato, apice emarginato.

Abdominis segmentum sextum inferius maris apice leviter bisinuatum, septimum profunde emarginatum, feminae apice medio leviter triangulariter productum.

\* 2. *Ph. caesus*: Niger, pedibus rufo-testaceis, elytris fuscis, thorace bi-impresso, tibiis anticis integris. — Long. 2 lin.

*Er. Col. March. I. 598. 2.*

Habitat in Germania.

Statura et summa similitudo praecedentis. Niger, subnitidus, pubescens. Antennae feminae capitis thoracisque longitudine, maris paulo longiores, nigrae. Oris partes rufo-piceae. Caput maris thoracis latitudine, suborbiculatum, feminae paulo angustius, oblongum, crebre profundeque punctatum, fronte posterius profunde canaliculata, anterieus utrinque ad antennarum basin impressa. Thorax coleopteris paulo angustior, latitudine plus dimidio brevior, apice truncatus, lateribus basin versus angustatus, subrotundatus, basi subrotundatus, angulis anterioribus subrectis, posterioribus obtusis, leviter convexus, crebre profunde punctatus, dorso foveolis duabus arcuatis impressus, interstitio subelevato, laevi. Elytra thorace paulo longiora, dense fortiter profundeque punctata, fusca, nitidula. Abdomen subtilissime coria-

ceum, parce obsoleteque punctatum, apice fusco-piceum. Pedes rufo-testacei, tibiis anticis angustis, integris, spinulis subtilibus ciliatis.

Abdominis segmentum inferius sextum maris integrum, septimum leviter emarginatum, feminae apice medio leviter triangulariter productum.

### **Trogphloeus Mannerh.**

*Er. Col. March. 1. p. 599.*

*Trogphloeus Mannerh. Brachelytr. p. 49. Steph. Illustr. V. p. 322. — Taenosoma Mannerh. Brachelytr. p. 50. Steph. Illustr. V. p. 323. — Carpalimus Steph. ibid. p. 324.*

*Paraglossae ligulae connatae.*

*Pedes intermedii basi approximati.*

*Tibiae omnes muticae.*

*Tarsi breves.*

Corpus vel oblongum, vel plerumque elongatum, lineare, aliis subdepressum, aliis magis subcylindricum, alatum. Caput porrectum, basi plerumque leviter constrictum, oculis semiglobosis, modice prominulis. Labrum transversum, medio leviter sinuatum, laciniis membraneis parvis, triangularibus, spatio interiecto setulis rigidis dense ciliato. Mandibulae haud prominentes, apice bicuspes, interne medio bidentatae, basi membrana fimbriatae. Maxillae (t. V. f. 19. b.) mala interiore margine exteriori coriaceo, interiore membraneo, apicem versus barbato, mala exteriori paulo longiore, coriacea, apice membranea, dense barbata. Palpi maxillares parum elongati, articulo primo minuto, secundo subobconico, tertio hoc longiore crassioreque, ovato, quarto minuto, subulato. Labium (t. V. f. 19. a.) mento transverso, apice late emarginato, ligula membranea, lata, apice medio leviter sinuata, paraglossis ligulae omnino connatis eamque haud excedentibus, nonnisi spinularum serie suturae loco distinguendae. Palpi labiales ligula parum longiores, articulo primo secundo paulo brevior, tertio praecedentibus tenuiore et paulo brevior. Antennae articulo primo parum elongato, secundo tertioque plerumque reliquis paulo longioribus, ultimis tribus saepius paulo maioribus, quinto et septimo etiam nonnunquam contiguis paulo crassioribus, ultimo ovato. Thorax plerumque subcordatus. Elytra truncata. Abdomen plerumque lineare, marginatum. Pedes breviusculi, tenuiusculi, coxis intermediis approximatis, tibiis omnibus muticis, pubescentibus, tarsis brevibus, triarticulatis, articulis duobus primis brevissimis.

### **Observationes.**

1. Corpus pube subtili sericea obductum. Punctura saepius subtilissima. Thorax saepius dorso longitudinaliter bi-impressus. Tarsis brevibus, tibia dimidia brevioribus hoc genus insigne videtur.

2. Tibiis inermibus optime Com. Mannerheim a praecedentibus hoc genus distinxit. Taenosoma (T. pusillus) ill. auctor inter Omalides collocavit, tarsorum articulorum numero haud respecto, praeterea antennarum articulis tribus ultimis paulo maioribus distinctum. Temere a Dom. Stephens eiusdem insecti tarsi 5-articulati descripti sunt. Carpalimum (T. bilineatus — nomen a Leachio olim toto huic generi impositum) Dom. Stephens a Trogphloeoo antennarum articulis ultimis 3 maioribus, a Taenosomate tarsis 3-articulatis distinguit. Antennarum articuli 3 ultimi in aliis speciebus minus, in aliis magis a praecedentibus magnitudine discedunt, quod in Bledio etiam pari modo se habet.

3. Sexus differentia latet.

4. Victus aliis sub arborum cortice, aliis in floribus, aliis in pratis et silvis uliginosis.

## \* Scutellati.

- \* 1. *T. dilatatus*: *Niger, cinereo-pubescens, thorace subcordato, foveola arcuata impresso, coleopteris thorace duplo latioribus.* — Long.  $1\frac{1}{2}$  lin.

Er. Col. March. I. 599. 1.

Habitat in Germania, Gallia; in Mesopotamia etiam a Dr. Helfer lectus.

Corpus latiusculum, subdepressum, nigrum, nitidulum, cinereo-pubescens. Antennae capitis thoracisque fere longitudine, graciles, apice sensim leniter incrassatae, nigro-piceae. Mandibulae rufae. Palpi nigri. Caput thorace paulo angustius, ad antennarum basin utrinque leviter impressum, subtiliter cinereo-pubescens. Thorax coleopteris duplo fere angustior, latitudine summa dimidio brevior, lateribus anterieus valde rotundatis, postice coarctatus, basi coleopteris triplo angustior, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusis, subdepressus, sulco arcuato, ferri equini figuram aequante, thoracis discum basi et lateribus ambiente, basi fortius impresso notatus, subtiliter cinereo-pubescens. Coleoptera apicem versus sensim leviter latiora, thorace plus sesqui longiora, subdepressa, confertim subtilissime punctata, densius cinereo-pubescentia. Abdomen sat latum, apice angustatum, confertim subtilissime punctulatum, cinereo-pubescens. Pedes nigri, tibiis fuscis, tarsis fusco-testaceis.

- \* 2. *T. segnis*: *Niger, nitidulus, griseo-pubescens, pedibus rufo-piceis, thorace subcordato, postice foveola lunata impresso, coleopteris thorace plus sesqui latioribus.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Madagascar, Dom. Goudot.

Statura et summa affinitas praecedentis. Corpus nigrum, nitidulum, pube brevi subtili grisea densius vestitum. Antennae capitis thoracisque longitudine, tenuiores, apicem versus sensim levissime incrassatae, nigrae, articulo primo piceo. Palpi picei. Caput thorace angustius, dense subtilissime punctatum, fronte anterieus utrinque longitudinaliter impressa. Thorax coleopteris plus dimidio angustior, latitudine brevior, lateribus anterieus fortiter rotundatis, basin versus angustatus, basi coleopterorum basi duplo angustior, basi apiceque truncatus, angulis posterioribus obtusis, anterioribus rotundatis, parum convexus, dense subtiliter punctatus, basi impressione transversali lunata fortius notatus. Coleoptera thorace plus sesqui longiora, apicem versus sensim paululum latiora, confertissime subtiliter punctata. Abdomen crebre minus subtiliter punctatum, longius et minus subtiliter pubescens. Pedes rufo-picei.

Obs. Distinctus a praecedente punctura minus subtili, thorace latiore, postice minus constricto, sulco impresso lateribus haud adscendente etc.

- \* 3. *T. Omalinus*: *Linearis, subdepressus, niger, pedibus testaceis, elytris fuscis, thorace plus duplo longioribus, thorace obsolete carinato, basi utrinque oblique impresso.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Saxonia, Dom. Märkel, in Anglia, Dom. Rudd.

Habitu fere et statura Omalii deplanati, linearis, subdepressus, niger, nitidus, pube brevi depressa minus subtili grisea sericante densius vestitus. Antennae capitis thoracisque prope longitudine, apicem versus sensim leviter incrassatae, nigrae vel nigro-piceae, basi rufo-piceae. Mandibulae et palpi



rufo-picei. Caput thorace parum angustius, supra utrinque parcius subtiliter punctatum, ad antennarum basin utrinque sulculo profundo impressum. Thorax coleopteris dimidio angustior, latitudine paulo brevior, lateribus leviter rotundatus, basin versus leviter angustatus, basi apiceque truncatus, angulis posterioribus obtusis, anterioribus rotundatis, parum convexus, parce subtilissime punctatus, dorso sulcis duobus longitudinalibus obsoletis approximatis, interstitio subcarinato, postice utrinque oblique impressus. Elytra thorace triplo fere longiora, depressa, dense minus subtiliter punctata, fusca. Abdomen pectore vix longius, parce subtilissime punctulatum. Pedes testacei.

- \* 4. *T. angustatus*: *Linearis, subdepressus, fuscus, antennis, pedibus elytrisque testaceis, his thorace plus sesqui longioribus, thorace longitudinaliter bi-impresso.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Bavaria superiore, Dom. Dr. Walth.

Elongatus, linearis, subdepressus, pube brevi, minus subtili, flavescenti-grisea vestitus. Antennae capite thoraceque paulo longiores, fortiusculae, apicem versus sensim paululum crassiores, testaceae. Os testaceum. Caput thoracis latitudine, utrinque parce subtilissime punctulatum, inter antennas utrinque longitudinaliter impressum, nigrum, nitidulum. Thorax coleopteris dimidio angustior, latitudine haud brevior, basin versus sensim leviter angustatus, lateribus parum rotundatus, basi apiceque truncatus, angulis omnibus obtusiusculis, subrotundatis, parum convexus, subtilissime punctulatus, impressionibus duobus longitudinalibus, linea media longitudinali subcarinata disiunctis, parum profundis, subrectis notatus, fuscus, parum nitidus. Scutellum fuscum. Elytra thorace plus sesqui longiora, depressa, confertim subtiliter punctata, testacea. Abdomen parcius subtiliter punctatum, longius pubescens, nigrum, subnitidum, ano piceo-testaceo. Pectus nigrum. Pedes pallide testacei.

### \*\* E x s c u t e l l a t i.

- \* 5. *T. signatus*: *Elongatus, niger, subtiliter cinereo-pubescens, antennarum basi, thoracis angulis anterioribus, elytrorum macula pedibusque rufis, thorace 5-foveolato.* — Long.  $2\frac{1}{4}$  lin.

*Erichs. Act. Acad. Caes. Leopold. Carol. Nat. Cur. Vol. XVI. Suppl. 229. 13. (Meyen It. III. 353. 13.).*

Habitat in Chile, Dom. Prof. Meyen.

Elongatus, niger, nitidus, pube brevi cinerea vestitus. Antennae capite thorace paulo breviores, apicem versus leniter incrassatae, piceae, articulis duobus primis rufis. Os rufo-piceum. Caput thoracis latitudine, subtilissime punctatum, fronte ad antennarum basin utrinque longitudinaliter impressa, callo elevato supra antennas rufo, subtus totum obscure sanguineum. Thorax coleopteris dimidio angustior, latitudine summa paulo brevior, lateribus antice rotundatis, basin versus sensim aequaliter angustatus, basi apiceque truncatus, angulis omnibus subrotundatis, parum convexus, crebre subtiliter punctatus, 5-foveolatus, foveolis duobus posterioribus approximatis, reliquis tribus in medio transversaliter dispositis, lateralibus rotundatis, intermedia lunata, supra nigro-piceus, angulis anterioribus obscure rufis, subtus saturate rufus. Coleoptera thoracis basi duplo latiora, eodemque sesqui

longiora, confertim punctata, prope apicem versus suturam macula minuta rubra notata. Abdomen lineare, subtilissime punctulatum, segmento penultimo apice piceo. Pectus postice utrinque macula rubra signatum. Pedes omnes cum coxis rufo-testacei.

- \* 6. *T. ferrugineus*: *Rubro-testaceus, subtiliter cinereo-pubescens, thorace transverso, dorso obsolete longitudinaliter bi-impresso.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Guyana Venezuelensi, Dom. Moritz.

Maioribus *T. riparii* individuis aequalis, rubro-testaceus vel dilute rufo-ferrugineus, parum nitidus, pube brevi subtili cinerea densius vestitus. Antennae capite thoraceque paulo longiores, fortiusculae, apicem versus sensim leniter incrassatae. Caput thorace angustius, fronte utrinque ad antennarum insertionem leviter impressa, fuscescente, ore testaceo. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus leviter rotundatus, basin versus leviter angustatus, basi apiceque truncatus, angulis posterioribus obtusis, anterioribus subrotundatis, vix convexus, dorso foveolis duabus approximatis, longitudinalibus, obsoletioribus impressus. Elytra thorace sesqui longiora, confertim subtiliter punctata. Abdomen omnium confertissime subtilissimeque punctatum, longius pubescens. Pedes testacei.

- \* 7. *T. angulatus*: *Elongatus, niger, subtiliter cinereo-pubescens, elytris apice, antennis basi pedibusque rufo-testaceis, thorace basi bifoveolato, angulis anterioribus acutis, lateribus acute angulatis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Brasilia, Dom. Melly.

Linearis, niger, parum nitidus, pube subtili depressa griseo-sericea densius vestitus. Antennae capite thoraceque paulo breviores, apicem versus parum crassiores, nigrae, articulo primo testaceo, sequentibus 4 piceis. Palpi rufo-testacei, articulo tertio piceo. Caput thorace parum angustius, ad antennarum basin utrinque longitudinaliter impressum. Thorax latitudine summa paulo brevior, lateribus ante medium fortiter dilatatis, angulatis, dein basin versus emarginato-angustatus, apice emarginatus, angulis anterioribus acuminatis, basi truncatus, angulis posterioribus obtusis, parum convexus, subtilissime confertissimeque alutaceus, foveolis approximatis binis prope basin, binis obsoletioribus in medio impressus. Coleoptera thoracis basi duplo fere latiora eodemque paulo longiora, confertim subtiliter punctata, secundum suturam longitudinaliter leviter impressa, fusco-picea, margine apicali testaceo. Abdomen confertissime subtilissimeque punctatum, longius pubescens, ano picescente. Pedes rufo-testacei.

- \* 8. *T. fulvipes*: *Elongatus, niger, subtiliter cinereo-pubescens, antennarum basi pedibusque rufo-testaceis, thorace postice longitudinaliter bi-impresso, angulis anterioribus acutis, lateribus obtuse angulatis.* — Long. 1 lin.

Habitat in ins. Puerto-Rico, Dom. Moritz.

Habitu omnino *T. bilineati*, linearis, niger, parum nitidus, pube subtili depressa densa cinereo-sericea obductus. Antennae capite thoraceque paululum breviores, apicem versus vix crassiores, fuscae, basi testaceae. Palpi testacei, articulo tertio fuscescente. Caput thorace parum angustius, ad an-

tennarum basin utrinque impressum. Thorax coleopteris paulo angustior, latitudine summa paulo brevior, lateribus ante medium fortiter dilatatis, obtuse angulatis, basin versus angustatus, apice emarginatus, angulis anterioribus acutis, posterioribus obtusis, basi truncatus, parum convexus, subtilissime confertissimeque punctulatus, dorso impressionibus duabus longitudinalibus leviter arcuatis, antrorsum divergentibus notatus. Coleoptera thorace sesqui longiora eiusque basi duplo latiora, confertissime subtiliter punctata, picea, margine apicali saepe dilutiore. Abdomen subtilissime punctulatum, apice piceum, subtus segmentorum marginibus rufo-piceis. Pedes rufo-testacei, coxis omnibus concoloribus.

- \* 9. *T. morio*: *Elongatus, niger, griseo-pubescent, tarsis testaceis, thorace posterius longitudinaliter bi-impresso, lateribus subangulato, elytris fortiter punctatis.* — Long.  $1\frac{3}{4}$  lin.

Habitat in Pensylvania, Dom. Zimmermann.

Niger, subnitidus, cinereo-pubescent. Antennae capite thoraceque paulo breviores, graciles, apice levissime incrassatae, piceo-nigrae. Palpi nigropicei. Mandibulae rufo-piceae. Caput thorace angustius, ad antennarum basin utrinque impressum, subtilissime pubescens. Thorax ante medium coleopteris dimidio angustior, latitudine summa plus dimidio brevior, lateribus ante medium fortiter dilatatis, subangulatis, basin versus angustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusis, parum convexus, confertim subtiliter punctatus, subtiliter pubescens, dorso posterius impressionibus duabus longitudinalibus, medio subinterruptis notatus. Coleoptera thorace paulo longiora eiusque basi plus sesqui latiora, secundum suturam leviter impressa, creberrime sat fortiter punctata, longius pubescentia. Abdomen supra segmentis singulis basi subtilissime punctatis, parcius pubescens, subtus longius densiusque pubescens. Pedes nigri, geniculis tibiatarumque apice rufo-piceis, tarsis rufo-testaceis.

- \* 10. *T. scrobiculatus*: *Niger, nitidus, griseo-pubescent, antennarum articulo primo pedibusque rufo-piceis, thorace cordato, 7-foveolato, foveola posteriore lunata.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Saxonia montana, Dom. Märkel.

Statura fere et habitu *T. dilatati*, niger, nitidus, pube subtili fusco-grisea vestitus. Antennae capite thoraceque paulo breviores, tenuiores, apice sensim leniter incrassatae, nigrae, articulo primo piceo-rufo. Palpi picei. Caput thorace paulo angustius, dense subtilissime punctatum, fronte anteriori utrinque fortius longitudinaliter impressa. Thorax ante medium coleopteris dimidio angustior, latitudine summa dimidio brevior, lateribus ante medium fortiter rotundatis, basin versus fortius angustatus, basi coleopteris plus duplo angustior, basi apiceque truncatus, angulis posterioribus obtusis, anterioribus rotundatis, parum convexus, dense subtiliter punctatus, medio foveolis duabus rotundatis approximatis, versus angulos anteriores utrinque foveola pariter rotundata, at obsoletiore, in ipso apice foveolis duabus minoribus obsoletis, modice distantibus, inter medium et basin sulco transversali, utrinque abbreviato, arcuato vel lunato, profundiore impressus. Elytra thorace plus sesqui longiora, dense minus subtiliter punctata. Abdomen confertim subtilissime punctulatum, longius pubescens. Pedes rufo-picei



- \* 11. *T. bilineatus*: *Elongatus, niger, subtiliter cinereo-pubescens, antennarum articulo primo pedibusque rufis, thorace transverso, subcordato, subdepresso, dorso 4-foveolato. — Long. 1 — 1 $\frac{1}{4}$  lin.*

*Er. Col. March. I. 600. 3.*

*Carpalimus bilineatus Steph. Illustr. Britt. Ent. V. 467. 1.*

*Oxyt. corticinus Gyll. Ins. Suec. II. 645. 14. — Sahlb. Ins. Fenn. I. 418. 19.*

*Trogophl. corticinus Mannerh. Brachelytr. 49. 1.*

*Var. pedibus piceis.*

*Trogophl. inquilinus Er. Col. March. I. 603. 6.*

Habitat in Europa.

Niger, subnitidus, subtiliter cinereo-pubescens. Antennae capitis thoracisque longitudine, apice perparum sensim incrassatae, piceo-nigrae, articulo primo rufo. Palpi rufi, articulo tertio nigro-piceo. Mandibulae basi nigrae. Caput thorace paulo angustius, inter oculos utrinque sat profunde impressum, omnium subtilissime confertissimeque punctatum. Thorax coleopteris dimidio angustior, latitudine summa dimidio brevior, lateribus anteriori fortiter rotundatus, basin versus angustatus, apice truncatus, basi levissime rotundatus, angulis posterioribus obtusis, anterioribus acutiusculis, leviter convexus, subtilissime confertissimeque punctatus, dorso foveolis duabus linearibus, longitudinalibus, basi apiceque profundioribus, medio subinterruptis, sat fortibus notatus. Coleoptera thorace sesqui longiora eiusque basi duplo prope latiora, antice secundum suturam impressa, minus subtiliter dense punctata. Abdomen subtilissime confertim punctatum. Pedes rufi, femoribus saepius medio piceis.

Variat pedibus et nonnunquam etiam antennarum articulo primo piceis.

- \* 12. *T. memnonius*: *Elongatus, niger, cinereo-pubescens, antennarum basi pedibusque rufo-testaceis, thorace obsolete 4-foveolato. — Long. 1 $\frac{1}{4}$  lin.*

Habitat in Aegypto, Dom. Prof. Ehrenberg.

Niger, nitidulus, subtiliter cinereo-pubescens. Antennae capite thoraceque paulo breviores, tenues, apice perparum incrassatae, nigro-piceae, articulo primo rufo, sequentibus duobus rufo-piceis. Mandibulae rufae. Palpi rufi, articulo tertio nigro. Caput thorace angustius, utrinque ad antennarum basin sat fortiter longitudinaliter impressum. Thorax coleopteris dimidio angustior, latitudine summa dimidio brevior, lateribus anteriori fortiter rotundatus, basin versus angustatus, basi apiceque truncatus, angulis anterioribus posterioribusque obtusis, perparum convexus, confertissime subtilissimeque punctatus, dorso postice foveolis duabus arcuatis, obsolete, subinterruptis impressus. Coleoptera thorace sesqui fere longiora eiusque basi duplo latiora, secundum suturam anteriori leviter impressa, dense subtiliter punctata, pube brevissima minus subtili vestita. Abdomen supra omnium subtilissime punctulatum, infra densius pubescens, apice summo piceo. Pedes rufo-testacei.

Obs. *C. bilineato proximus*, differt autem antennis brevioribus gracilioribusque, thorace latiore, anteriori fortius dilatato, obsolete foveolato.

- \* 13. *T. riparius*: *Elongatus, niger, subtiliter cinereo-pubescens, antennarum articulis duobus primis pedibusque rufis, thorace transverso, subdepresso, dorso longitudinaliter bi-impresso. — Long. 1½ lin.*

*Boisd. et Lacord. Faun. Ent. Paris. I. 467. 1. — Er. Col. March. I. 600. 2.*

Habitat in Europa.

Niger, nitidulus, cinereo-pubescens. Antennae capitis thoracisque prope longitudine, crassiusculae, apicem versus sensim paulo crassiores, pubescentes, piceae, articulis duobus primis rufis. Os cum palpis rufum, his articulo tertio concolore. Caput thorace paulo angustius, utrinque ad antennarum basin sat profunde impressum, confertissime subtilissimeque punctatum, subtiliter pubescens. Thorax coleopteris paulo angustior, latitudine summa dimidio fere brevior, lateribus anterie fortiter rotundatus, basin versus angustatus, basi apiceque truncatus, angulis anterioribus acutiusculis, posterioribus obtusis, parum convexus, subtilissime confertissimeque punctatus, subtiliter pubescens; dorso posterius foveis duabus linearibus, longitudinalibus, leviter arcuatis, antice divergentibus impressus. Coleoptera thorace sesqui longiora eiusque basi sesqui latiora, secundum suturam antice leviter impressa, dense fortius punctata, dense longius pubescentia. Abdomen supra omnium subtilissime alutaceum, subtilissime punctulatum, parcius pubescens, apice summo piceo, infra densius pubescens. Pedes rufi, coxis omnibus concoloribus.

Variat elytris piceis.

Obs. C. bilineato affinis, at statura paulo maiore, magis lineari, antennis crassioribus, articulis duobus primis rufis, coleopteris minus amplis, fortius punctatis, etc. distinctus.

- \* 14. *T. elongatulus*: *Linearis, subdepressus, niger, densius cinereo-pubescens, antennarum basi pedibusque rufo-piceis, thorace subcordato, dorso longitudinaliter bi-impresso. — Long. 1 lin.*

*Er. Col. March. I. 601. 4.*

Habitat in Germania.

Elongatus, linearis, niger, parum nitidus, subtiliter cinereo-pubescens. Antennae capite thoraceque paulo longiores crassiusculae, apicem versus sensim subincrassatae, nigrae vel fuscae, basi rufae. Os rufo-testaceum, palpis articulo tertio piceo. Caput thorace parum angustius, confertissime subtilissimeque punctatum, utrinque ad antennarum basin sat profunde impressum. Thorax coleopteris parum angustior, latitudine vix dimidio brevior, lateribus leviter rotundatus, basin versus angustatus, basi apiceque truncatus, angulis anterioribus obtusis, subrotundatis, posterioribus fortiter rotundatis, parum convexus, subtilissime confertissimeque punctatus, dorso posterius foveolis duabus linearibus longitudinalibus, subarcuatis, parum profundis impressus. Coleoptera thorace paulo longiora eiusque basi sesqui latiora, secundum suturam antice subimpressa, confertissime subtiliter punctata. Abdomen elongatum, lineare, subtilissime confertim punctulatum, minus subtiliter pubescens. Pedes rufi, femoribus saepe picescentibus.

Obs. A praecedentibus statura minore, elongata et elytris subtilius punctatis distinctus.

- \* 15. *T. fuliginosus*: *Elongatus, niger, dense cinereo-pubescens, antennarum basi pedibusque rufis, thorace transverso, lateribus rotundato-ampliato, leviter convexo, basi obsolete bifoveolato. — Long. 1—1½ lin.*

*Er. Col. March. I. 602. 5.*

*Oxyt. fuliginosus Grav. Micr. 102. 1. Mon. 185. 1. — Latr. Hist. nat. Crust. et Ins. IX. 362. 1. — Oliv. Encycl. meth. VIII. 613. 1.*

Habitat in Germania.

Niger, nitidus, pube subtili cinereo-sericea densius vestitus. Antennae capite thoraceque paulo breviores, apicem versus sensim subincrassatae, rufae, apice fuscescentes. Oris partes rufae. Caput thorace multo angustius, omnium subtilissime confertissimeque punctatum, fronte utrinque ad antennarum basin profunde impressa. Thorax coleopteris parum angustior, latitudine summa dimidio fere brevior, lateribus ante medium dilatatis, rotundatis, basin versus angustatus, basi apiceque truncatus, angulis anterioribus subrectis, posterioribus obtusis, leviter convexus, medio per totam longitudinem subcarinatus, omnium subtilissime punctatus, basi obsolete bifoveolatus, impressionum dorsalium vix rudimento ullo. Coleoptera thorace paulo longiora eiusque basi sesqui latiora, secundum suturam per totam longitudinem leviter impressa, confertissime subtilissimeque punctata, nonnunquam picea. Abdomen confertissime subtilissimeque punctatum, apice summo piceo. Pedes rufi.

- \* 16. *T. flavipes*: *Elongatus, niger, cinereo-pubescens, elytris apice, antennae basi pedibusque testaceis, thorace dorso longitudinaliter bi-impresso. — Long. ¾ — 1 lin.*

Habitat in Americae meridionalis insulis St. Johannis et St. Thomae, Dom. Moritz.

Niger, nitidus, subtiliter cinereo-pubescens. Antennae capitis thoracisque longitudine, apicem versus leniter incrassatae, fuscae, articulis duobus primis testaceis. Oris partes testaceae. Caput thorace parum angustius, subtilissime obsolete punctulatum, fronte utrinque ad antennarum basin profunde foveolata. Thorax coleopteris paulo angustior, latitudine summa dimidio brevior, lateribus fortiter rotundatus, basin versus angustatus, apice truncatus, angulis anterioribus obtusis, basi cum angulis posterioribus rotundatus, leviter convexus, confertissime subtilissimeque punctatus, dorso foveolis duabus linearibus, longitudinalibus, subarcuatis, profundioribus impressis. Coleoptera thorace paulo longiora eiusque basi sesqui latiora, confertim subtiliter punctata, fusco-picea, margine apicali testaceo. Abdomen elongatum, subtilissime obsolete punctulatum. Pedes testacei.

- \* 17. *T. Niloticus*: *Elongatus, fuscus, cinereo-pubescens, capite abdomineque nigris, antennae pedibusque testaceis, thorace obsolete 4-foveolato. — Long. ¾ lin.*

Habitat in Aegypto, a Dom. Prof. Ehrenberg copiose lectus.

Minutus, fuscus, capite abdomineque nigris, cinereo-pubescens. Antennae capitis thoracisque longitudine, apicem versus leviter incrassatae, articulis ultimis tribus abrupte paulo maioribus, testaceae, apice fuscescentes. Palpi flavi. Caput thorace parum angustius, omnium subtilissime punctula-



tum, fronte utrinque impressa. Thorax coleopteris dimidio angustior, latitudine paulo brevior, basin versus angustatus, lateribus ante medium rotundatis, apice truncatus, angulis anterioribus obtusis, basi cum angulis posterioribus rotundatus, leviter convexus, omnium subtilissime punctulatus, dorso obsolete longitudinaliter bi-impressus, impressionibus medio subinterruptis. Coleoptera thorace sesqui longiora eiusque basi duplo latiora, confertim subtiliter punctata. Abdomen confertim subtilissime punctulatum, fortius pubescens. Pedes pallide flavi.

Variat minus maturus corpore testaceo, capite abdomineque fuscis.

- \* 18. *T. corticinus*: *Elongatus, niger, subtilissime cinereo-pubescens, antennis concoloribus, pedum geniculis tarsisque piceis, thorace transverso, subcordato, dorso fortius longitudinaliter bi-impresso.* — Long.  $\frac{3}{4}$  — 1 lin.

*Er. Col. March. I.* 603. 7.

*Oxyt. corticinus* Grav. *Mon.* 192. 8. — *Oliv. Encycl. meth. VIII.* 416. 8.

*Oxyt. minimus* Runde *Brachelytr. Hal.* 20. 7.

Habitat in Germania, frequens.

Niger, subnitidus, tenuiter subtiliterque cinereo-pubescens. Antennae capite thoraceque paulo breviores, minus tenues, apice leniter incrassatae, nigrae, articulis ultimis tribus distinctius abrupte paulo maioribus, articulo primo piceo-nigro. Palpi piceo-nigri. Caput thoracis prope magnitudine, confertim subtilissime punctatum, fronte ad antennarum basin utrinque foveolata. Thorax coleopteris paulo angustior, latitudine summa tertia parte brevior, lateribus rotundatus, basin versus angustatus, apice truncatus, basi cum angulis posterioribus leviter rotundatus, angulis anterioribus acutiusculis, parum convexus, confertissime subtiliter punctatus, dorso foveis duabus linearibus longitudinalibus, leviter arcuatis impressus. Coleoptera thorace plus sesqui longiora eiusque basi sesqui latiora, antice secundum suturam longitudinaliter leviter impressa, confertim subtiliter punctata. Abdomen subtiliter obsoleteque punctulatum. Pedes nigri, trochanteribus, geniculis tibiaramque apice piceis, tarsis testaceis.

Variat pedibus rufis, antennarum articulis tribus primis piceis.

- \* 19. *T. exiguus*: *Subcylindricus, niger, subtilissime cinereo-pubescens, antennis palpisque concoloribus, pedum geniculis tarsisque piceis, thorace oblongo, basin versus angustato, convexo, dorso obsolete foveolato.* — Long.  $\frac{2}{3}$  lin.

*Er. Col. March. I.* 604. 8.

*Trogophl. impressus* Boisd. et Lacord. *Faun. Ent. Paris. I.* 467. 2. (*forte*)

Habitat in Germania, Gallia.

Statura subcylindrica fere Bledii cuiusdam, minutus, niger, parum nitidus, subtilissime cinereo-pubescens. Antennae capite thoraceque breviores, apicem versus leniter sensimque incrassatae, nigrae. Palpi nigri. Mandibulae rufae. Caput thoracis magnitudine, basi vix constrictum, fronte utrinque ad antennarum basin leviter impressa. Thorax coleopteris parum angustior, latitudine summa vix brevior, lateribus ante medium modice rotundatus, basin versus angustatus, basi subrotundatus, apice truncatus, angulis antero-

ribus obtusiusculis, posterioribus obtusis, subrotundatis, satis convexus, subtilissime confertim punctatus, impressionibus duabus oblongis dorsalibus obsoletissimis. Coleoptera thorace paulo longiora eiusque basi sesqui latiora, confertim subtiliter punctata. Abdomen subtiliter punctulatum. Pedes picei, trochanteribus, geniculis, tibiæ apice tarsisque rufescentibus, interdum tibiis etiam totis rufis.

- \* 20. *T. punctatellus*: *Niger, nitidulus, parce cinereo-pubescent, pedibus piceis, capite thoraceque crebre, elytris dense fortiterque punctatis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Carinthia, Dom. Schmidt.

Niger, nitidulus, pube subtili cinerea parcius vestitus, haud sericans. Antennae capite thoraceque paulo breviores, apicem versus leviter incrassatae, articulo secundo tertio multo maiore, crasso, quinto contiguis paulo crassiore, ultimis tribus paulo maioribus, nigrae, articulo primo rufo-piceo. Oris partes testaceae, palpis nigro-piceis. Caput thoracis prope latitudine, crebre profundius et minus subtiliter punctatum, linea media longitudinali laevi, fronte ad antennarum basin utrinque foveolata. Thorax coleopteris paulo angustior, latitudine summa dimidio fere brevior, lateribus fortius rotundatus, basin versus angustatus, basi apiceque truncatus, angulis anterioribus subrotundatis, posterioribus obtusis, modice convexus, aequalis, crebre minus subtiliter profunde punctatus, linea media longitudinali laevi subcarinata. Coleoptera thorace paulo latiora et eiusdem basi sesqui latiora, dense fortiter profundeque punctata. Abdomen parce subtilissime punctulatum. Pedes picei, trochanteribus, geniculis tarsisque testaceis.

- \* 21. *T. foveolatus*: *Niger, subtiliter cinereo-pubescent, pedibus piceis, thorace obsolete 4-foveolato, elytris dense fortius punctatis.* — Long.  $\frac{2}{3}$  lin.

*Oxyt. foveolatus* Sahlb. Ins. Fenn. I. 419. 20.

Habitat in Germania, Dom. Schüppel, in Sardinia, Dom. Prof. Gené.

T. corticino paulo minor, praecipue brevior, niger, minus nitidus, subtiliter cinereo-pubescent. Antennae capite thoraceque multo breviores, minus graciles, articulo secundo tertio multo maiore, crassiusculo, apice leniter incrassatae, nigrae, basi piceae. Palpi nigro-picei. Mandibulae testaceae. Caput thorace paulo angustius, opacum, fronte ad antennarum basin utrinque profunde impressa. Thorax coleopteris parum angustior, latitudine summa dimidio brevior, lateribus fortius rotundatus, basin versus angustatus, basi apiceque truncatus, angulis posterioribus rotundatis, anterioribus obtusis, supra leviter convexus, opacus, dorso foveolis 4 obliquis, binis prope basin, binis in medio sitis, omnibus obsoletioribus impressus. Coleoptera thorace sesqui longiora eiusque basi sesqui latiora, dense fortius profunde punctata. Abdomen subtilissime punctulatum. Pedes picei, geniculis tarsisque testaceis.

- \* 22. *T. troglodytes*: *Niger, subtilissime cinereo-pubescent, antennarum basi, pedibus elytrisque piceis, his apice testaceo-marginatis, thorace obsolete 4-foveolato.* — Long.  $\frac{2}{3}$  lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura omnino praecedentis, niger, minus nitidus, capite thoraceque opacis. Antennae capitis thoracisque longitudine, minus graciles, articulo secundo tertio crassiore, apicem versus sensim leviter incrassatae, nigricantes, basi piceae. Palpi picei. Caput thoracis fere latitudine, omnium subtilissime alutaceum, fronte depressiuscula, ad antennarum basin utrinque sulculo profundiore impressa. Thorax coleopteris paulo angustior, latitudine summa tertia parte brevior, lateribus, basi et angulis posterioribus rotundatis, basin versus angustatus, apice truncatus, angulis anterioribus rectis, supra parum convexus, omnium subtilissime alutaceus, subtilissime obsoleteque punctulatus, dorso foveolis quatuor obsoletioribus, binis prope basin, binis in medio sitis impressus. Coleoptera thorace paulo longiora eiusque basi sesqui latiora, minus dense subtiliter punctata, nitida, nigro-picea, margine apicali testaceo. Abdomen parcius subtilissime punctulatum. Pedes picei, trochanteribus, geniculis, tibiis apice tarsisque testaceis.

Obs. Distinctus a praecedente thorace minus transverso, elytris paulo brevioribus, minus dense et minus fortiter punctatis, neque concoloribus.

\* 23. *T. pusillus*: *Elongatus, subdepressus, niger, subtiliter cinereo-pubescent, antennarum basi pedibusque luteis, elytris fusco-testaceis, thorace transversim subcordato, dorso obsoletius 4-foveolato.* — Long.  $\frac{3}{4}$  lin.

*Er. Col. March. I. 605. 9.*

*Aleochara pusilla* Grav. *Micr. 78. 17. Mon. 153. 16.* — *Latr. Hist. nat. Crust. et Ins. IX. 383. 17.* — *Gyll. Ins. Suec. II. 409. 31.*

*Taenodoma pusillum* Mannerh. *Brachelytr. 51. 2.*

*Oxyt. fuliginosus* Gyll. *Ins. Suec. II. 460. 14.* — *Sahlb. Ins. Fenn. I. 418. 18.* — *Zetterst. Ins. Lappon. 73. 6.*

*Trogophl. corticinus* Boisd. et Lacord. *Faun. Ent. Paris. I. 468. 3.*

Habitat in Europa.

Minutus, linearis, subdepressus, niger, parum nitidus, dense subtiliterque cinereo-pubescent. Antennae capite paulo longiores, fusco-testaceae, articulo primo testaceo, ultimis tribus distincte crassioribus, fuscis, tertio secundo multo minore, quinto contiguus paulo crassiore. Palpi testacei, articulo tertio fusco. Os testaceum. Caput thorace vix angustius, fronte inter oculos utrinque sulco sat profundo impressa. Thorax coleopteris paulo angustior, latitudine summa paulo brevior, lateribus leviter rotundatus, basin versus modice angustatus, basi subrotundatus, apice truncatus, angulis omnibus obtusis, subrotundatis, parum convexus, vix punctulatus, dorso foveis duabus oblongis, sublinearibus, medio interruptis, basi apiceque profundioribus, leviter impressus. Coleoptera thorace sesqui longiora eiusque basi sesqui latiora, confertissime subtiliter punctata, basi secundum suturam leviter impressa, fusco-testacea. Abdomen confertissime subtilissimeque punctatum, apice piceum. Pedes flavo-testacei.

\* 24. *T. tenellus*: *Linearis, niger, subtilissime cinereo-pubescent, antennarum basi pedibusque testaceis, elytris fuscis, thorace subquadrato, dorso obsolete impresso.* — Long.  $\frac{3}{4}$  lin.

*Er. Col. March. I. 605. 10.*

*Taenodoma gracile* Mannerh. *Brachelytr. 51. 1. (forte)*

Habitat in Germania.



Minutus, valde elongatus, linearis, subdepressus, niger, parum nitidus, subtilissime pubescens. Antennae capite sesqui longiores, crassae, apicem versus fortius incrassatae, articulo tertio secundo multo minore, quinto septimoque contiguis paulo crassioribus, ultimis tribus distincte maioribus, fuscae, articulo primo testaceo. Os testaceum. Caput thoracis magnitudine, omnium subtilissime confertissime punctatum, fronte inter antennis utrinque foveolata. Thorax coleopteris paulo angustior, latitudine vix brevior, lateribus parum rotundatis, basin versus subangustatus, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrectis, parum convexus, confertissime subtilissimeque punctatus, dorso obsolete bi-impressus, impressionibus basi profundioribus. Elytra thorace duplo prope longiora, confertissime subtiliter punctata, iam nigro-fusca, iam fusco-testacea. Abdomen confertim subtilissime punctatum, apice summo piceo. Pedes testacei.

Variat thorace elytrisque piceis.

\* 25. *T. subtilis*: *Linearis, niger, subtilissime cinereo-pubescens, antennis pedibusque testaceis, elytris fuscis, thorace oblongo, basin versus leviter angustato, dorso obsolete impresso.* — Long.  $\frac{3}{4}$  lin.

*Er. Col. March. I. 606. 11.*

Habitat Berolini.

Statura minuta, tenui praecedentis, eique proximus, niger, parum nitidus, subtiliter cinereo-pubescens. Antennae capite sesqui longiores, crassiusculae, articulo tertio secundo multo minore, ultimis tribus distincte crassioribus, totae testaceae. Os testaceum. Caput thoracis latitudine vel eodem paulo latius, basi leviter constrictum, subtilissime confertissimeque punctulatum, inter antennis utrinque fortiter foveolatum. Thorax coleopteris paulo angustior, latitudine summa haud brevior, lateribus anterieus rotundatus, basin versus angustatus, basi cum angulis posterioribus leviter rotundatus, angulis anterioribus obtusiusculis, parum convexus, omnium subtilissime confertissimeque punctatus, basi obsolete bi-impressus. Coleoptera thorace sesqui longiora eiusque basi plus sesqui latiora, secundum suturam longitudinaliter obsolete impressa, fusca. Abdomen confertissime subtilissimeque punctatum, apice piceum. Pedes breves, testacei.

Obs. A praecedente praecipue thorace paulo longiore, basin versus distincte angustato, antennis totis testaceis distinguitur.

### **Apocellus Er.**

*Paraglossae ligulae connatae.*

*Pedes intermedii basi approximati.*

*Tibiae omnes muticae.*

*Tarsi modice elongati.*

Corpus minutum, elongatum, lineare, apterum. Caput maiusculum, exsertum, porrectum, rotundatum, collo quasi tenui thoraci immissum, oculis mediocribus, leviter prominulis. Labrum latitudine paulo brevius, medio leviter sinuatum, laciniis membranaceis linearibus. Mandibulae breves, haud exsertae, apice fortiter tridentatae, interne basi membrana margine barbata fimbriatae. Maxillae (t. V. f. 20. b.) malis membranaceis, apice barbatis, interiore interne ciliata, modice elongatis, exteriori longiore. Palpi maxil-

lares mala exteriore vix longiores, articulo primo parvo, secundo apicem versus leviter incrassato, tertio hoc paulo longiore, crassiusculo, quarto parvo, subulato. Labium (t. V. f. 20. a.) mento transverso, ligula lata, membranacea, apice leviter emarginata, paraglossis ligulae connatis eandemque haud superantibus. Palpi labiales ligula paulo longiores, articulis longitudine crassitieque decrescentibus. Antennae refractae, modice elongatae, apicem versus sensim incrassatae. Thorax coleopteris angustior, subglobosus. Scutellum breviusculum, triangulare. Elytra breviuscula, truncata. Abdomen lineare, marginatum, supra planum, subtus convexum. Pedes elongati, graciles, coxis anticis conicis, liberis, intermediis subapproximatis, tibiis omnibus teretibus, pubescentibus, tarsis triarticulatis, articulis duobus primis brevibus, tertio elongato.

#### O b s e r v a t i o n e s.

1. Habitu Stilici vel Falagriae reliquis Oxytelinis omnino alienus videtur. Corpus laevigatum, parce pilosellum.

2. Characteribus plerisque, oris imprimis structura, cum Trogophloeo convenit, distinctus autem habitu, corpore haud pubescente, pedibus magis elongatis, tarsis minus brevibus, tibia dimidia longioribus.

3. Sexus differentia in abdominis apice inferiore sita in speciebus describendis illustranda.

4. De victu nihil constat.

\* 1. *A. sphaericollis*: *Piceus, nitidus, abdomine nigro, antennarum basi pedibusque piceo-testaceis.* — Long.  $1\frac{1}{2}$  lin.

T. II. f. 5.

*Lathrobium sphaericolle* Say Transact. Amer. Philoph. Soc. Philadelph. Nov. Ser. IV. 455. 6.

Habitat in Carolina, Dom. Zimmermann.

Antennae capitis thoracisque fere longitudine, apicem versus sensim incrassatae, fusco-piceae, basi piceo-testaceae. Os piceo-testaceum. Caput thoracis fere latitudine, suborbiculatum, piceum, laevissimum, fronte utrinque longitudinaliter impressa, impressione subtilissime strigosa, margine laterali super antennarum basin plicato-elevato. Thorax coleopteris paulo angustior, subglobosus, laevissimus, piceus, nitidus. Elytra thorace haud longiora, laevissima, basi summa subtilissime strigosa, picea, disco nigra, nitida. Abdomen laeve, nigrum. Pedes piceo-testacei. Corpus totum pilis raris erectis instructum.

Abdominis segmentum inferius sextum maris margine apicali medio dilatato, reflexo, leviter emarginato, subtiliter ciliato, flavido, septimum profunde excisum, lateribus lobatum, lobis conniventibus, feminae apice medio leviter triangulariter productum.

\* 2. *A. ustulatus*: *Testaceus, elytris abdomineque apice nigris.* — Long.  $1\frac{1}{3}$  lin.

Habitat in Ins. St. Johannis Americae, Dom. Moritz.

Flavo-testaceus, nitidus, pilis parvis erectis vestitus. Antennae capitis thoracisque longitudine, apicem versus sensim incrassatae, testaceae, basi dilutiores. Caput thorace latius, suborbiculatum, laevissimum, fronte utrinque leviter longitudinaliter impressa, margine laterali super antennarum basin

plicato-elevato. Thorax parvus, coleopteris duplo fere angustior, subquadratus, leviter convexus, laevissimus. Elytra thorace paulo longiora, laevissima, piceo-nigra, nitida, basi piceo-testacea. Abdomen laeve, segmentis 4 primis testaceis, reliquis piceo-nigris. Pedes pallide testacei.

Abdominis segmentum inferius sextum maris apice medio leviter emarginatum, septimum profunde sinuatum, feminae apice rotundatum.

\* 3. *A. albipes*: *Piceus, nitidus, antennis basi testaceis, abdominis segmento primo pedibusque albidis, femoribus apice fuscis.* — Long. 1 lin.

Habitat in Columbia, Dom. Moritz.

Saturate fusco-piceus, nitidus, parce pilosus. Antennae capitis thoracisque longitudine, apicem versus sensim incrassatae, fuscae, basi testaceae, articulo ultimo ferrugineo. Palpi testacei. Caput thorace latius, suborbiculatum, laevissimum, rufo-piceum, fronte utrinque ad antennarum basin subimpressa, margine laterali super antennarum insertionem plicato-elevato. Thorax coleopteris multo angustior, latitudine haud brevior, lateribus anterioribus leviter rotundatus, basin versus subangustatus, convexus, laevissimus. Elytra thorace paulo longiora, laevissima. Abdomen segmento primo albido. Pedes albi, femoribus apice ad medium usque fuscis.

Variat capite rufo-piceo.

### Subtribus quarta.

## Coprophilini.

*Antennae 11-articulatae.*

*Ligula membranea.*

*Tarsi 5-articulati.*

Labrum transversum, corneum,  
apice utrinque lacinia membranea instructum Coprophilo, Acrognatho, Deleasteri;

apice toto membrana limbatum Syntomio.

Mandibulae muticae vel medio subdentatae,  
haud prominentes Coprophilo, Deleasteri, Micralymmati, Syntomio;

prominentes, acutae Acrognatho.

Maxillae malis coriaceis membranaceisve, exteriore apice, interiore interne barbata.

Palpi maxillares filiformes, articulo primo brevi.

Ligula membranea, biloba.

Paraglossae ligulae aequales.

Palpi labiales breviusculi, filiformes.

Antennae 11-articulatae,

rectae Coprophilo, Deleasteri, Micralymmati, Syntomio;

subrefractae Acrognatho;

articulis ultimis distincte crassioribus subclavatae Acrognatho, Syntomio.



Oculi mediocres.

Thorax basi truncatus, ab elytris parum remotus.

Scutellum distinctum.

Abdomen marginatum.

Pedes coxis intermediis approximatis, tibiis anterioribus spinulosis Coprophilo, Acrognathos; omnibus muticis Deleasteri, Micralymmati, Syntomio; tarsis 5-articulatis.

#### O b s e r v a t i o.

Discedunt ab Oxytelinis genuinis praecipue palpis filiformibus et tarsis 5-articulatis. Deleaster Anthophago, reliqua huius subtribus genera Omalio ab auctoribus adscripta sunt, a quibus, licet habitu iis similes, praecipue ocellis nullis, elytris pectore haud longioribus, abdominis segmento septimo exserto distinguuntur.

### Coprophilus Latr.

Latr. Regn. Animal. IV, p. 439. Nov. Annal. Mus. Hist. nat. I, p. 89.

*Paraglossae ligulae connatae.*

*Antennae rectae, apicem versus sensim subincrassatae.*

*Tibiae anteriores extus spinulosae.*

Corpus elongatum, lineare, subdepressum, alatum. Caput porrectum, basi haud coarctatum, thoraci immissum eodemque angustius, oculis subovalibus, mediocribus, parum prominulis. Labrum (t. V. f. 21. c.) transversum, apice subsinuatum, laciniis membraneis minutis, triangularibus, interne ciliatis. Mandibulae haud prominentes, validae, medio subdentatae. Maxillae (t. V. f. 21. b.) mala interiore externe coriacea, interne membranea margine interiore ciliato, mala exteriori coriacea, apice dense barbata. Palpi maxillares articulo primo minuto, secundo elongato-obconico, tertio hoc paulo brevior, quarto secundo prope longior, elongato-ovato, apice subacuminato. Labium (t. V. f. 21. a.) mento transversum, apicem versus leviter angustatum, apice truncatum, ligula membranea, lata, apice late emarginata, paraglossis linearibus, ligulam haud excedentibus, membraneis, interne spinulis ciliatis. Palpi labiales articulis tribus longitudine aequalibus, ultimo ovato, subacuminato. Antennae rectae, apicem versus sensim subincrassatae, articulo primo crassiusculo, secundo tertio paulo brevior, 4 — 10 sensim paulo crassioribus et magis obconicis, ultimo ovato. Thorax coleopteris parum angustior, basin versus subangustatus. Scutellum triangulare. Elytra truncata, angulo exteriori obtuso. Abdomen lineare, late marginatum. Pedes breviusculi, validiusculi, coxis intermediis carinula tenuissima disiunctis, tibiis integris, omnibus seriatim setulosis, anterioribus extus uni-seriatim spinulosis, tarsis 5-articulatis, articulis 4 primis brevibus, quinto his coniunctis subaequali.

#### O b s e r v a t i o n e s.

1. Habitus aliquatenus Oxyteli. Corpus glabrum, capite thoraceque minus subtiliter punctatis, elytris seriatim punctatis, abdomine subtiliter pubescente.

2. Mas abdominis segmento ventrali septimo apice minus rotundato a femina distinguitur.

3. Victus sub lapidibus.

- \* 1. *C. striatulus*: *Niger, nitidus, ore pedibusque piceis, thorace bifoveolato, elytris fortiter punctato-striatis.* — Long. 3 lin.

*Er. Col. March. I.* 609. 1.

*Staph. striatulus* *Fab. Ent. Syst. I. II.* 529. 27. *Syst. El. II.* 596. 35. — *Panz. Ent. Germ.* 354. 19.

*Staph. rugosus* *Oliv. Ent. III.* 42. 30. 42. t. 4. f. 43.

*Omalium rugosum* *Grav. Micr.* 115. 7. *Mon.* 203. 11. — *Latr. Hist. nat. Crust. et Ins. IX.* 372. 7. — *Oliv. Encycl. meth. VIII.* 476. 11. — *Gyll. Ins. Succ. II.* 233. 30.

*Anthobium rugosum* *Boisd. et Lacord. Faun. Ent. Paris. I.* 469. 1.

*Coprophilus rugosus* *Guér. Iconogr. Regn. An. Ins. t. 10. f. 2.*

*Oxyt. laesus* *Germ. Faun. Ins. Europ. XV.* 2.

Habitat in Europa.

Corpus nigrum piceumve, nitidum. Antennae fere capitis thoracisque longitudine, apicem versus paululum incrassatae, nigro-piceae, apice rufescentes. Caput sat crebre inaequaliter punctatum, fronte super oculos striga transversali, utrinque ad antennarum insertionem fovea longitudinali impressa. Thorax coleopteris paulo angustior, latitudine parum brevior, basin versus leviter angustatus, lateribus antrorsum modice rotundatus, apice subemarginatus, angulis anterioribus leviter prominulis, subdepressus, crebrius punctatus, basi foveolis duabus approximatis, medio fovea elongata, lineari, canaliculae instar impressus, lateribus late marginatus, margine obsolete crenulato. Scutellum punctis nonnullis impressum. Elytra thorace sesqui longiora, sulcata, sulcis fortiter punctatis, interstitiis laevibus, apice confer-tim ruguloso-punctata. Abdomen subtiliter punctulatum. Pedes rufo-picei.

### **Acrognathus Er.**

*Er. Col. March. I.* p. 607.

*Paraglossae apice liberae.*

*Antennae subrefractae, articulis ultimis crassioribus.*

*Tibiae anteriores spinulosae.*

Corpus elongatum, lineare, alatum. Caput porrectum, basi leviter contractum. Labrum (t. V. f. 22. c.) breve, medio sinuatum, utrinque lacinia sat elongata, acuminata, membranea, interne pilis longioribus ciliata auctum. Mandibulae elongatae, prominentes, falcatae, acutae. Maxillae (t. V. f. 22. b.) mala interiore membranea, margine exteriori corneo, interiore ciliato, apice barbata; mala exteriori cornea, apice barbata. Palpi maxillares modice elongati, articulo primo minuto, secundo tertioque inter se longitudine subaequalibus, quarto his singulis sesqui fere longiore, apice acuminato. Labium (t. V. f. 22. a.) mento brevi, apice truncato, ligula membranea, sinu profundo biloba, lobis sublinearibus, divergentibus, paraglossis his connatis, eosque haud superantibus, apice summo liberis. Palpi labiales subelongati, articulo primo secundoque fere aequalibus, tertio his tenuiore et parum longiore. Antennae subrefractae, articulo primo longiusculo crassiusculoque, secundo tertioque leviter elongatis, obconicis, hoc secundo paulo longiore, ultimis 3 vel 5 distincte crassioribus. Thorax coleopteris paulo angustior, basin versus subangustatus. Scutellum triangulare. Elytra truncata, angulo exteriori rotundato. Abdomen lineare, late marginatum. Pedes parum elongati, intermedii approximati, tibiis omnibus setulosis, posterioribus apice

oblique truncatis, anterioribus a medio apicem versus sensim attenuatis, extus uniseriatim spinulosis, tarsis 5-articulatis, articulis 4 primis minutis, inter se aequalibus, quinto his coniunctis subaequali.

#### O b s e r v a t i o.

Habitus fere Trogophloeï, at tibiae spinulosae, tarsi 5-articulati, et palporum articulus ultimus elongatus. Mandibulae prominentes affinitatem quandam cum Bledio probant. Antennae subrefractae, sub tuberculo frontis marginali insertae, articulis ultimis tribus saepe distinctius maioribus, ut in genere laudato utroque.

- \* 1. *A. mandibularis*: *Rubro-testaceus, capite piceo, elytris subtiliter subseriatim punctatis.* — Long. 3 lin.

Er. Col. March. I. 608. 1.

*Omalium mandibulare* Gyll. Ins. Suec. IV. 468. 29 — 30.

Habitat in Germania, Gallia, Suecia, rarior.

Corpus rubro-testaceum, fere lateritium, opacum. Antennae capitis thoracisque fere longitudine, articulis ultimis tribus abrupte paulo maioribus, ultimo ovato, concolores. Caput fronte antice subretusa, parce punctata, nigricante, tuberculo utrinque marginali super antennarum insertionem elevato dilutius. Oculi fortiter prominuli. Thorax coleopteris dimidio angustior, latitudine parum longior, basin versus leviter angustatus, lateribus et basi leviter rotundatus, apice truncatus, angulis anterioribus subrotundatis, posterioribus subrectis, leviter convexus, subtilissime alutaceus, parcius punctatus, linea media subelevata laevi, lateribus tenuiter marginatus, saturate rubro-testaceus, subtiliter pubescens. Elytra thorace sesqui longiora, subtilissime alutacea, subtiliter subseriatim punctata, lineis tribus subelevatis laevibus, dilutius rubro-testacea, subtiliter griseo-pubescencia. Abdomen subtilissime alutaceum, subtiliter pilosellum. Pedes testacei.

Abdominis segmentum inferius secundum maris tuberculo minuto apicali munitum, feminae muticum.

- \* 2. *A. cephalotes*: *Piceus, capite maiusculo, oblongo, thorace elongato, dorso bisulcato, rufo, elytris dense seriatim punctatis, piceo-testaceis.* — Long. 3½ lin.

Habitat in Corcyra, Dom. Schüppel.

Latitudine praecedentis, at magis elongatus linearisque. Antennae capite thoraceque multo breviores, articulis 5 ultimis abrupte distincte maioribus, ultimo subovato, rufae. Os rufum, mandibulis piceis. Caput thorace paulo latius, oblongum, basi perparum constrictum, fronte lateribus crebre, medio biseriatim profundius punctata, nigro-piceum, basi rufescens. Oculi haud prominuli. Thorax coleopteris paulo angustior, latitudine paulo longior, basin versus leviter angustatus, lateribus subrectis, basi leviter rotundatus, apice late leviter emarginatus, angulis anterioribus acutiusculis, posterioribus obtusis, modice convexus, dorso longitudinaliter bisulcatus, sulcis et in utroque latere plaga longitudinali leviter flexuosa crebre punctatis, piceo-rufus, nitidus, uti caput pube erecta parca subtili vestitus. Elytra thorace sesqui prope longiora, dorso striatim, apice disperse crebreque fortius punctata, subtiliter griseo-pubescencia, piceo-testacea, dorso nigricantia. Abdomen crebre punctulatum, subtiliter griseo-pubescens, piceum, nitidum. Pectus nigro-piceum. Pedes breviusculi, testacei.



\* 3. *A. palpalis*: Testaceus, abdomine ante apicem, pectore fronteque piceis, elytris fortius seriatim punctatis. — Long. 1 lin.

*Er. Col. March. I.* 608. 2.

Habitat Berolini, Dom. Schüppel, in Saxonia, Dom. Märkel, in ins. Coreyra, Dom. Schüppel.

Minutus, subdepressus, testaceus, vix nitidus. Antennae capitis thoracisque longitudine, articulis septimo et octavo praecedentibus crassioribus, ultimis tribus his iterum maioribus et crassioribus, ultimo subovato, apice abrupte subacuminato, flavae, articulis ultimis 5 fuscis. Caput thoracis latitudine, basi minus distincte constrictum, fronte planiuscula, subtilissime alutacea, parcius punctata, picescente, ore flavo. Palpi eiusdem structurae atque in praecedentibus, at paulo crassiores. Oculi leviter prominuli. Thorax coleopteris dimidio angustior, latitudine paulo longior, basin versus leviter angustatus, lateribus subrectis, basi leviter rotundatus, apice leviter emarginatus, angulis anterioribus acutiusculis, posterioribus obtusis, leviter convexus, subtilissime alutaceus, vage punctatus, linea media sat lata subelevata laevi. Elytra thorace sesqui fere longiora, subdepressa, dorso sat distincte regulariterque seriatim punctata, flavo-testacea, subnitida, subtilissime pubescentia. Abdomen ante apicem piceum. Pectus piceum. Pedes flavi.

Abdominis segmentum inferius septimum maris apice subrotundatum, feminae acuminatum.

### **Deleaster Er.**

*Er. Col. March. I.* p. 610.

*Paraglossae liberae.*

*Palpi maxillares articulo ultimo penultimo sesqui longiore.*

*Antennae rectae, filiformes.*

*Tibiae omnes muticae.*

Corpus modice elongatum, subdepressum, alatum. Caput exsertum, porrectum, collo quasi cylindrico thoraci immissum, subovatum, oculis rotundatis, mediocribus, sat prominulis. Labrum (t. V. f. 24. c.) transversum, apice leviter sinuatum, laciniis membraneis triangularibus, interne ciliatis. Mandibulae parum prominentes, muticae. Maxillae (t. V. f. 24. b.) malis membranaceis, interiore interne, exteriore apice barbatis. Palpi maxillares parum elongati, articulo primo minimo, secundo elongato-obconico, tertio hoc brevior, quarto secundo paulo longiore, subacuminato. Labium (t. V. f. 24. a.) mento transversum, ligula membranea, lata, apice late profundeque sinuata, paraglossis liberis, linearibus, acutis, ligulam haud superantibus. Palpi labiales ligulam parum excedentes, articulis duobus primis cylindricis, secundo paulo brevior, tertio primo aequali, subovato. Antennae elongatae, validiusculae, rectae, articulo primo leviter elongato, secundo hoc triplo minore, subobconico, tertio iterum elongato, reliquis longitudine aequalibus, et sensim paululum crassioribus, ultimo ovato. Thorax coleopteris multo angustior, subcordatus. Scutellum apice rotundatum. Elytra sat ampla, singula apice leviter rotundata. Abdomen latum, late marginatum. Pedes elongati, graciles, coxis intermediis approximatis, tibiis omnibus subtiliter

pubescentibus, tarsis 5-articulatis, articulis 4 primis obcordatis, subaequalibus, quinto his coniunctis paulo brevior.

### O b s e r v a t i o n e s .

1. Habitus omnino Anthophagi, at huius tribus, et accuratius inspectus sat distinctus.

2. Dom. Curtis hoc insectum pro generis *Lesteuae* typo proposuit, at oris partes Anthophagi exhibuit.

3. Mas differt a femina tarsis anterioribus articulis 4 primis leviter dilatatis, subtus subtiliter spongiosis, abdominis segmento inferiore septimo apice truncato neque rotundato.

4. Victus riparius.

\* 1. *D. dichrous*: Testaceus, capite nigro, abdomine piceo. — Long. 3 lin.

*Er. Col. March. I.* 611. 1.

*Anthophagus dichrous* Grav. *Micr.* 188. 1. *Mon.* 221. 6.

*Lesteua dichroa* Latr. *Hist. nat. Crust. et Ins.* IX. 369. 7. — Guér. *Iconogr.*

*Regn. An. Ins. t.* 10. *f.* 3. — Mannerh. *Brachelytr.* 55. 1. — Boisd. et Lacord.

*Faun. Ent. Paris. I.* 479. 1. — Runde *Brachelytr. Hal.* 23. 1.

*Lesteua Leachii* Curt. *Britt. Ent.* VII. t. 303.

*Staph. Brassicae* Schrank *Faun. Boica. I.* 644. 873. (*forte*)

Habitat in Germania, Gallia, Anglia.

Antennae capite thoraceque longiores, rufo-ferrugineae. Caput thorace sesqui latius, impunctatum, fronte medio convexa, lateribus super oculos et inter antennis longitudinaliter impressa, sub antennis linea sat profunda transversaliter divisa, nigrum, subnitidum, ore late rufo. Thorax coleopteris duplo fere angustior, latitudine summa paulo brevior, ante medium fortiter rotundatus, basin versus angustatus, basi apiceque truncatus, leviter convexus, parce punctatus, minus distincte canaliculatus, basi super scutellum et lateribus paulo infra medium foveola sat magna impressus, rufus, subnitidus. Scutellum parce subtilissime punctulatum. Elytra thorace duplo latiora et duplo longiora, subdepressa, dense subtiliter punctata, subtiliter flavescenti-pubescentia, rufo-testacea, subnitida. Abdomen dorso depressum, subtiliter punctulatum, subtilissime tenuiter flavescenti-pubescenti, subnitidum, iam nigrum, iam piceum, iam piceo-testaceum, apicem versus magis nigricans. Pectus rufo-testaceum. Pedes testacei.

### **Micralymma Westw.**

*Westwood Magaz. of Zool. and Bot. II. p.* 129. *t.* 4.

*Palpi maxillares articulo ultimo penultimo triplo longiore.*

*Antennae rectae, filiformes.*

*Tibiae omnes muticae.*

Corpus elongatum, sublineare, depressum, apterum. Caput oblongum, thorace paulo angustius, porrectum, basi vix constrictum, oculis mediocribus, rotundatis, parum prominulis. Labrum transversum, apice bisinuatum, margine ciliato (W.). Mandibulae haud prominentes, muticae. Maxillae mala interiore gracili, apice acuto intus setoso, exteriori maiore, sub apice articulata (Westw.). Palpi maxillares articulo primo minuto, secundo obco-

nico, tertio hoc brevior, quarto iterum longior, apice subacuminato. Mentum transversum. Ligula biloba. Palpi labiales breves, 3-articulati, articulis sensim decrescentibus (Westw.). Antennae filiformes, apicem versus sensim paulo crassiores, articulo primo crassiusculo, secundo hoc paulo brevior, tenuioreque, subobconico, tertio hoc longitudine aequali, at iterum paulo tenuiore, quarto sequentibusque subglobosis, sensim paulo latoribus, penultimis sensim transversis, ultimo breviter ovato, apice subacuminato. Thorax coleopteris paululum angustior, basin versus leviter angustatus. Scutellum breve, apice leviter rotundatum. Elytra breviora, apice leviter rotundata. Abdomen elongatum, ab apice basin versus sensim subangustatum, late marginatum. Pedes breviusculi, intermediis basi approximati, tibiis omnibus muticis, tarsis 5-articulatis, articulis 4 primis perbrevis, inter se aequalibus, subtus pubescentibus, quinto his coniunctis paulo brevior.

#### O b s e r v a t i o n e s .

1. Corpus parcius subtiliter pubescens. Caput, thorax et elytra parce, abdomen paulo crebrius punctata.

2. Habitus obiter observatus, Aleocharam quandam, e. g. Homalotam simulat, Oxytelinis alienus, at characteres huius tribus et subtribus.

3. Sexus differentia latet.

4. Victus in littore maris, intra aestus surgentis et collabentis limites, quo fit, ut aestu alterno animalculum retegatur et altera vice per quatuor horas obtegatur.

5. Larva et nympha, a Dom. Johnston cum imagine lectae hoc modo a Dom. Westwood describuntur: „Larvae corpus elongatum, angustum. Caput oblongum, depressum. Mandibulae falcatae, intus medio fortiter dentatae. Maxillae stipite elongato, mala 2-articulata, articulis longitudine subaequalibus, secundo tenuiore. Palpi maxillares 4-articulati, articulis 2 primis crassis, primo brevior, tertio quartoque tenuioribus, subaequalibus. Labium quadrangulum, stipitibus palporum labialium cylindricis, liberis. Palpi labiales tenues, 2-articulati. Antennae 4-articulae, articulo primo, secundo quartoque longitudine subaequalibus, tertio duplo longior, processu laterali aucto. Prothorax segmentis sequentibus maior, his subaequalibus, excepto ultimo minore, ano tubuloso, utrinque cauda setosa 2-articulata instructo. Pedes breves, praeter coxam 4-articulati, unguiculo singulo instructae. Nympha thoracis margine antico setis duabus perlongis et nonnullis brevioribus instructo, abdomine lateribus setoso, apice lobulis duabus angustis terminato.

1. *M. brevipenne*: Nigrum, subnitidum, parce subtiliter punctatum, antennis ante basin geniculisque piceis. — Long.  $1\frac{2}{3}$  lin.

*Omalium brevipenne* Gyll. Ins. Suec. II. 234. 31. — Zetterst. Faun. Lappon. I. 56. 18. Ins. Lappon. 54. 26.

*Micralymma Johnstonis* Westw. Mag. of Zool. et Bot. II. p. 130. t. 4. f. 1.

Habitat in littore maris Sueciae et Angliae. Specimen a Dom. Westwood missum e coll. Dom. Aubé descripsi.

Nigrum, subnitidum, parce subtiliter pubescens. Antennae capitis thoracisque prope longitudine, nigrae, articulo secundo tertioque piceis. Caput thorace paulo angustius, depressum, parce subtiliterque punctatum, fronte inter oculos et inter antennarum insertionem utrinque obsolete impressa. Thorax latitudine anteriore paululum brevior, basin versus angustatus, lateribus ante medium fortiter rotundatis, pone medium subsinuatis, basi apiceque truncatus, angulis omnibus rotundatis, parum convexus, parce subtiliterque punctatus, dorso longitudinaliter depressus, linea media longitudinali



subelevata laevi. Elytra thorace haud longiora, apice singula leviter rotundata, basin versus sensim leviter angustata. Abdomen crebrius subtiliusque punctulatum. Pedes nigri, geniculis piceis.

### **Syntomium Curt.**

*Curt. Britt. Ent. V. t. 228.*

*Paraglossae ligulae connatae.*

*Palpi maxillares articulo quarto tertio minore, acuminato.*

*Antennae rectae, articulis ultimis tribus abrupte crassioribus.*

*Tibiae omnes mulicae.*

Corpus breviusculum, crassiusculum, alatum. Caput thorace paulo minus et angustius, porrectum, basi haud constrictum, oculis mediocribus, subglobosis, leviter prominulis. Labrum transversum, medio sinuatum, margine antico membrana prominula limbato, laciniis lateralibus parvis, angulatim prominentibus, sinu interiecto setis crassiusculis dense ciliato. Mandibulae haud prominentes, breviusculae, crassiusculae, muticae, intus basi membrana ciliata fimbriatae. Maxillae (t. V. f. 23. b.) malis breviusculis, interiore membranacea, interne ciliata, exteriore coriacea, apice barbata. Palpi maxillares breviores, articulo primo minimo, secundo obconico, tertio paulo maiore, crassiusculo, quarto hoc minore, acuminato. Labium (t. V. f. 23. a.) mento transversim subquadrato, ligula lata, medio sinuata, membranea, apice medio spinulis duabus brevibus, acutis, corneis instructa, paraglossis ligulae aequalibus, eidemque connatis, sutura nonnisi setularum serie indicata. Palpi labiales ligulam vix superantes, articulis duobus primis aequalibus, tertio paulo minore. Antennae rectae, articulo primo secundoque crassiusculis, tertio sequentibus longiore, obconico, 4—8 parvis, subglobosis, ultimis tribus distincte crassioribus, clavam oblongam formantibus. Thorax transversus, coleopteris angustior. Elytra apice singula rotundata, interne coniunctim sinuata. Abdomen coleopterorum latitudine, supra planum, lateribus late marginatum, subtus convexum. Pedes breviusculi, coxis intermediis approximatis, tibiis omnibus simplicibus, tarsis 5-articulatis, articulis 4 primis brevibus, aequalibus, quinto his coniunctis aequali.

### **O b s e r v a t i o n e s.**

1. Corpus fortiter punctatum, tenuiter pubescens, colore metallico insigne.
2. Sexus differentia latet.
3. Victus in silvis umbrosis. Incessus lentus tardusque.
4. Larva glomeriformis, ovata, convexa, laevis, nigro-aenea, nitida, cum imagine occurrit. Specimina sicca e Carinthia a Dom. Schmidt cum imagine missa magis cum Silpharum, quam cum reliquorum Staphylinorum larvis congruere videntur, mihi haud rite examinata.

\* 1. *S. aeneum*: *Viridi-aeneum, nitidum, antennis apice pedibusque rufopiceis, capite, thorace elytrisque profunde punctatis.* — Long. paulum ultra 1 lin.

*Omalium aeneum* Müller Germ. Mag. IV. 216. 17. — Gyll. Ins. Suec. IV. 466. 22 — 23.

*Syntomium nigro-aeneum* Curtis Britt. Ent. V. t. 228.

Habitat in Germania, Suecia, Anglia, locis umbrosis humidis lente tardeque per humum repens (Müll.).

Corpus obesulum, virescenti-aeneum, nitidum, subtilissime pubescens. Antennae capitis thoracisque prope longitudine, crassiusculae, pilosellae, nigrae, apicem versus rufo-piceae. Caput thorace duplo minus, crebre fortiter punctatum, inter antennis bifoveolatum. Thorax coleopteris paulo angustior, latitudine duplo fere brevior, lateribus fortiter rotundatis, basin versus sinuato-angustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusiusculis, fortius convexus, creberrime fortiter profundeque punctatus, linea media longitudinali laevi subcarinata, basi bifoveolatus, margine laterali subtiliter crenulato. Elytra thorace paulo longiora, leviter convexa, basi media subelevata, confertissime fortiter profundeque punctato-subrugosa. Abdomen laeve, lateribus rotundatum, margine laterali elevato. Corpus subtus nigrum, pectore punctato, abdomine laevi. Pedes rufo picei.

## Tribus octava.

### P i e s t i n i.

*Stigmata prothoracica occulta.*

*Coxae posticae transversae, anticae globosae, haud prominentes.*

*Trochanteres postici simplices.*

Labrum breve, transversum, apice medio sinuatum, corneum, utrinque lacinia interne ciliata auctum, lacinia

triangulari, membranea, labro contigua plerisque;

rhomboidea, membranea, labro continua Leptochiro.

Mandibulae validae, dentatae, prominentes, saepius porrectae Leptochiro, Piesto,

muticae, interne lacinia membranea ciliata fimbriatae, parum prominentes reliquis;

maris cornutae Prognathae.

Maxillae stipite extus sub palporum basi leviter dilatata, mala inferiore angustiore, brevior, interne barbata, exteriore segmento inferiore minore corneo, apicali membraneo,

dilatato, semicirculari plerisque;

parallelo Hypotelo.

Palpi maxillares filiformes, articulo primo minuto.

Mentum corneum, labii partes membraneas praeter ligulam obtegens.

Ligula membranea saepius biloba.

Paraglossae nullae plerisque;

ligulae connatae Hypotelo.

Palpi labiales breves, filiformes.

Antennae sub frontis margine laterali insertae, 11-articulatae, rectae, filiformes, nonnunquam corpore longiores.

Caput porrectum, thoraci immissum, basi leviter constrictum.

Thorax totus corneus, spatio etiam pone coxas anticae clauso, coleopterorum saepius latitudine,

basi ab elytris remotus Leptochiro, Piesto, Isomalo;

elytrorum basi applicatus Lispino, Prognathae, Hypotelo.



Scutellum distinctum.

Elytra pectoris longitudine.

Abdomen segmentis sex compositum, septimo haud exserendo, immarginatum, cylindricum *Leptochiro*;

immarginatum, subdepressum *Lispino*;

tenuissime marginatum *Isomalo*;

late marginatum *Piesto*, *Prognathae*, *Hypotelo*.

Pedes breviusculi, validiusculi, coxis anticis parvis, globosis, ex acetabulis haud prominentibus, intermediis approximatis, trochanteribus posticis simplicibus, tibiis, anticis,

marginē exteriore crenato *Leptochiro*, *Piesto*;

integris, anterioribus spinulosis *Prognathae*, *Hypotelo*;

muticis *Lispino*, *Isomalo*;

spinis terminalibus altera validiore, tibiarum anticarum valida, tarsis 5-articulatis, simplicibus.

#### Observationes.

1. Proxima tribui praecedenti, quacum a Latreillio confusa est, at coxis anticis globosis haud exsertis, abdominis segmento septimo nullo modo conspicuo etc. satis distincta. Labri structura in utraque eadem.

2. Mas nonnunquam fronte cornuta spinosae, interdum etiam antennis elongatis insignis. Notae sexuales in abdominis apice haud observantur.

3. Victus sub arborum putrescentium cortice.

### **Leptochirus Germ.**

*Germ. Spec. Ins. nov. p. 35.*

*Mandibulae exsertae, dentatae.*

*Tibiae anticae serratae.*

*Abdomen immarginatum.*

Corpus mediocre, elongatum, depressum, abdomine angusto, cylindrico, alatum. Caput exsertum, porrectum, subquadratum, fronte interdum antrorsum prominente, oculis parvis, rotundatis, prominulis. Labrum (t. V. f. 7. c.) emarginatum, laciniis membraneis longioribus, parallelipipedis, intus ciliatis. Mandibulae validae, porrectae, fortiter dentatae. Maxillae (t. V. f. 7. b.) malis membraneis, interiore oblonga, intus barbata, exteriore suborbiculata, apice pubescente. Palpi maxillares articulo primo minuto, secundo elongato, subclavato, tertio hoc plus duplo breviorē, obconico, quarto tertio paulo longiore, cylindrico, apice subacuminato. Labium (t. V. f. 7. a.) mento antrorsum angustato, lateribus sinuato, apice medio obsolete undentato, ligula bifida, laciniis linearibus, divergentibus, membranaceis, intus ciliatis. Palpi labiales ligulae subaequales, articulis duobus primis cylindricis, tertio paulo minore, acuminato. Antennae capite thoraceque haud longiores, articulo primo magno, clavato, secundo minuto, globoso, reliquis moniliformibus. Thorax coleopterorum latitudine, quadratus, basi medio paulum productus, ab elytris remotus. Scutellum parvum triangulare. Coleoptera quadrata, angulis posterioribus rotundatis. Abdomen coleopteris duplo angustius, subcylindricum, immarginatum, segmento penultimo reliquis paulo longiore, ultimo conico. Pedes breves, femoribus incrassatis, tibiis angustis, anticis margine exteriore dense serrato, posterioribus muticis, tarsis brevissimis, debilibus, articulis quatuor primis minutis, aequalibus.

## O b s e r v a t i o n e s.

1. Corpus parce pilosellum, capite, thorace elytrisque supra glaberrimis. Thorax linea impressa canaliculatus. Elytra laevigata, stria suturali impressa.

2. A Latreillio aliisque nonnullis auctoribus cum Zirophoro commutatus, quocum fronte saepius cornuta, mandibulis exsertis dentatis et tibiis anticis serratis convenit, a quo autem praeter oris structuram abdomine immarginato cylindrico facile distinguitur.

3. Sexus differentia latet.

4. Victus gregarius sub arborum putrescentium cortice.

\* 1. *L. maxillosus*: *Niger, nitidus, fronte mutica, canaliculata, antice utrinque foveolata, apice producta, biloba.* — Long.  $4\frac{1}{3}$  —  $6\frac{1}{3}$  lin.

*Var. a: Thorace pectore, elytrisque rufo-castaneis.*

*Leptochirus brunneo-niger* Perty *Delect. An. Art. p. 32. t. 7. f. 7.*

*Leptochirus Cayennensis* Laporte *Etud. Ent. I. 125. 2. t. 4. f. 5.*

*Var. b: Thorace, pectore elytrisque rufo-castaneis, his apice nigris.*

*Cucuius maxillosus* Fab. *Syst. El. II. 93. 8.*

Habitat in America meridionali: Brasilia, Cayenna, Columbia, Mexico.

Niger, nitidus. Antennae capite thoraceque paulo breviores, pilosellae. Caput thorace paulo angustius, fronte plana, canaliculata, apice utrinque foveolata, medio producta, biloba. Thorax subquadratus vel latitudine parum brevior, lateribus rectis, apice late leviter emarginatus, planus, canaliculatus. Coleoptera thoracis longitudine latitudineque, plana, stria suturali impressa, margine inflexo punctulato. Abdomen parce pubescens, segmentis singulis basi utrinque parum-punctatis. Pedes nigri, tarsis rufo-piceis, femoribus posticis fortius incrassatis.

Variat thorace, pectore elytrisque aut nigerrimis aut rufo-castaneis, his interdum apice nigris.

Frons nonnunquam, forte secundum sexum (an in femina?) ante apicem transversim subimpressa.

\* 2. *L. scoriaceus*: *Niger, nitidus, fronte mutica, canaliculata, apice bi-impressa, thorace transversim subquadrato.* — Long. 9 lin.

*Germ. Spec. Ins. nov. 35. 58. t. 1. f. 1.*

*Zirophorus coriaceus* Griff. *Animal Kingd. XIV. p. 306. t. 32. f. 4.*

*Var. Plus duplo minor, long. 5 lin.*

Habitat in Brasilia; in Columbia interiore, Dom. Buquet; in Mexico, collect. Dom. Chevrolat.

Corpus totum nigerrimum, tarsis solis rufo-piceis. Antennae capiti thoracique longitudine subaequales, pilosellae. Caput thorace paulo angustius, fronte depressa, profunde canaliculata, antice haud protuberante, foveolis duabus transversis terminata. Thorax latitudine paulo brevior, lateribus rectis, parallelis, apice late leviter emarginatus, planus, canaliculatus, laevis. Elytra thorace paululum longiora, plana, laevissima, stria suturali impressa, lateribus inflexis obsolete parceque punctatis. Abdomen utrinque parce punctatum.

- \* 3. *L. Mexicanus*: *Niger, nitidus, fronte mutica, canaliculata, apice bi-impressa, thorace quadrato. — Long. 7 lin.*

Habitat in Mexico.

Corpus totum nigerrimum, nitidum, tarsis solis rufo-piceis. Antennae capitis thoracisque longitudine, pilosellae. Caput thorace paulo angustius, fronte profunde canaliculata, apice haud producta, foveolis duabus transversim terminata, punctis quatuor notata, binis ad apicem approximatis, binis in medio, magis distantibus. Thorax latitudine haud brevior, lateribus rectis, parallelis, apice truncatus, subdepressus, laevis, profunde canaliculatus. Elytra thorace paululum longiora, subdepressa, laevissima, stria suturali impressa, lateribus inflexis parce obsoleteque punctatis. Abdomen lateribus parce obsoleteque punctatum.

Obs. Differt a praecedente statura graciliore, minus plana, thorace latitudine haud brevior, fronte 4-punctata.

- \* 4. *L. laticeps*: *Niger, nitidus, fronte mutica, plana, antice bifoveolata. — Long. 5½ — 6 lin.*

Habitat in Java, Dom. De Haen.

Niger, nitidus, tarsis solis rufo-piceis. Antennae capite thoraceque breviores, pilosellae. Caput thoracis latitudine, fronte plana, laevissima, haud canaliculata, apice foveolis duabus minutis profundis distantibus impressa. Thorax latitudine brevior, lateribus rectis, basin versus subangustatus, apice late leviter emarginatus, planus, laevis, subtilius canaliculatus. Coleoptera thorace subangustiora et vix longiora, depressa, laevissima, stria suturali impressa, lateribus inflexis crebrius punctatis. Abdomen lateribus crebrius punctatum.

- \* 5. *L. unicolor*: *Niger, nitidus, fronte postice canaliculata, antice subimpressa, margine antico tridentato. — Long. 6 lin.*

Laporte Etud. Ent. I. 125. 1.

Habitat in Java, Dom. De Haen.

Statura praecedentium, picco-niger, nitidus. Antennae capite thoraceque paulo breviores, dense flavescenti-pilosae. Palpi rufo-picei. Caput thorace angustius, fronte profunde canaliculata, clypei spatio subdepresso, linea semicirculari sat profunde impressa circumdato, apice tridentato. Thorax latitudine brevior, lateribus levissime rotundatis, planus, laevis, profunde canaliculatus. Elytra thorace paulo longiora, plana, laevia, stria suturali impressa, lateribus inflexis parce punctatis. Abdomen lateribus parce punctatum. Pedes dilute rufo-picei.

- \* 6. *L. ebeninus*: *Niger, nitidus, fronte postice profunde canaliculata, antice subimpressa, margine antico bidentato. — Long. 3½ lin.*

Habitat in Madagascar, Dom. Goudot.

Praecedente plus duplo minor, nigerrimus, nitidus. Antennae vix capitis thoracisque longitudine, crassiusculae, parce pilosellae. Palpi rufo-picei. Caput thorace angustius, fronte profunde canaliculata, clypei spatio subimpresso, linea semicirculari impressa circumscripto, margine laterali elevato, apice denticulo prominulo terminato. Thorax latitudine brevior, la-



teribus subrectis, antrorsum subangustatis, subdepressus, profunde canaliculatus, lateribus parce sat distincte punctatus. Elytra thorace longiora, depressa, laevia, stria suturali impressa, margine laterali punctorum serie notato. Abdomen parce obsolete punctulatum. Pedes picei, tarsis dilute rufo-piceis.

OBS. *Leptochirus convexus* Laporte (Etud. Ent. I. p. 132.) absque dubio idem, at descriptio manca non satis quadrat, quam ob rem hoc insectum pro novo proponendum putavi.

7. *L. brevicornis*: *Niger, nitidus, thorace piceo, fronte excavata, postice canaliculata, antice utrinque elevato-producta.* — Long. 5 lin.

Habitat in Java, collect. Dom. Chevrolat.

Habitu omnino *L. unicoloris*, niger, nitidus, thorace piceo, nitido. Antennae capite thoraceque paulo breviores, dense flavescenti-pilosae. Palpi rufo-picei. Caput thorace angustius, fronte tota excavata, lateribus elevatis, antice elevato-productis, postice canaliculata, clypei spatio linea impressa semicirculari terminato, margine antico mutico. Thorax coleopteris paulo lator, postice latitudine duplo fere brevior, antrorsum angustatus, leviter convexus, dorso canaliculatus, praeterea laevissimus, lateribus parce punctatis. Elytra thorace longiora, fere plana, laevissima, stria suturali impressa. Abdomen laeve, parce flavescenti-pilosellum, segmentis singulis margine postico piceo. Pedes flavescenti-piloselli, nigro-picei, tarsis rufescensibus.

- \* 8. *L. bispinus*: *Niger, nitidus, fronte bispinosa.* — Long. 3 $\frac{3}{4}$  lin.

Habitat in Java, Dom. De Haen.

Corpus totum nigerrimum, nitidum. Antennae capitis thoracisque prope longitudine, pilosellae, piceae, articulo primo nigro. Mandibulae elongatae. Caput thorace paulo angustius, fronte plana, antice in spinas duas porrectas, sat validas, acutas producta, inter spinas profunde impressa. Thorax latitudine brevior, lateribus parallelis, vix rotundatis, apice leviter late emarginatus, planus, laevissimus, subtilius canaliculatus. Elytra thorace vix longiora, plana, laevia, stria suturali impressa, lateribus punctis paucis impressa. Abdomen laeve. Pedes trochanteribus tibiisque piceis, his apicem versus sensim dilutioribus, tarsis dilute rufo-piceis.

- \* 9. *L. Javanicus*: *Niger, nitidus, thorace sanguineo, fronte bispinosa.* — Long. 7 lin.

Laporte Etud. Ent. I. 126 3. t. 4. f. 6.

Habitat in Java, Dom. De Haen.

Corpus nigerrimum, nitidum, thorace sanguineo. Antennae capite thoraceque paulo breviores, dense pilosellae, piceae, articulo ultimo ferrugineo. Mandibulae valde elongatae. Caput thorace angustius, fronte plana, antice in spinas duas sat validas longasque, acutas, porrectas producta, inter spinas excavata. Thorax latitudine brevior, lateribus perparum rotundatus, apicem versus subangustatus, planus, obsolete canaliculatus, lateribus punctis parvis impressus. Elytra thorace paulo longiora, plana, laevissima, stria suturali impressa, margine laterali punctorum serie notato. Abdomen segmentis singulis basi utrinque punctorum piliferorum serie transversali impressis. Pedes nigri, tarsis dilute rufo-piceis, femoribus anticis sanguineis.

**Lispinus Er.**

*Mandibulae muticae, haud prominentes.*

*Tibiae anticae integerrimae.*

*Abdomen immarginatum.*

Corpus lineare, subdepressum, alatum. Caput porrectum, fronte antice rotundata, subtiliter marginata, oculis rotundatis, leviter prominulis. Labrum minutum, utrinque lacinia triangulari membranea auctum. Mandibulae simplices, falcatae, porrectae, at minutae et haud prominentes. Maxillae (t. V. f. 8. b.) mala exteriore segmento primo sive basali corneo, secundo subsemicirculari, membraneo, costulis radialibus plicato, basi laevi, mala interiore extus cornea, apice denticulo introrsum curvato armata, margine interiore membraneo, apicem versus spinulis ciliato. Palpi maxillares breves, articulo primo minuto, secundo longiore, obconico, tertio brevi, crassiusculo, quarto iterum longiore, cylindrico. Labium (t. V. f. 8. a.) mento antrorsum angustato, utrinque bisinuato, apice rotundato, ligula sat ampla, triloba, lobis lateralibus latis, interne pubescentibus intermedio multo brevior, emarginato, Palpi ligula paulo longiores, articulis longitudine subaequalibus. Antennae capite thoraceque haud longiores, crassiusculae, articulo primo tertioque reliquis transversis paulo longiores. Thorax quadrangulus, basi coleopteris applicatus. Scutellum minutum, triangulare. Elytra truncata. Abdomen lineare, subdepressum, immarginatum. Pedes breves, femoribus crassiusculis, compressis, tibiis tenuibus, omnibus muticis, tarsis minutis.

## O b s e r v a t i o n e s.

1. Corpus glabrum, saepius punctatum, fronte semper mutica, elytris stria suturali impressa. A praecedente genere mandibulis tibiisque muticis, thorace basi truncato et minus ab elytris remoto, abdomine depressiusculo, a reliquis abdomine immarginato praeter oris structuram distinguitur.

2. Sexus differentia latet.

- \* 1. *L. attenuatus*: *Piceus, nitidus, pedibus rufis, crebre punctatus, fronte bifoveolata, thorace basi utrinque longitudinaliter impresso, apicem versus subangustato.* — Long.  $2\frac{1}{3}$  lin.

Habitat in ins. Puerto-Rico, Dom. Moritz.

Corpus elongatum, sublineare, subdepressum, piceum, nitidum. Antennae capite thoraceque paulo breviores, crassiusculae, apicem versus leniter incrassatae, articulo tertio elongato, piceae. Os rufo-piceum. Caput thorace paulo angustius, crebrius subtiliter punctatum, fronte leviter convexa, inter oculos bi-impressa. Thorax basi coleopterorum latitudine, latitudine paulo brevior, apicem versus sensim leviter angustatus, lateribus subrectis, apice subemarginatus, parum convexus, creberrime punctatus, punctis oblongis, medio linea subtilissima utrinque abbreviata canaliculatus, basi utrinque longitudinaliter leviter impressus. Elytra thorace paulo longiora, sat crebre parum profunde punctata, stria suturali impressa. Abdomen obsoletissime punctulatum, apice rufo-piceum. Pedes rufi.

- \* 2. *L. laticollis*: *Rufo-piceus, nitidus, parce subtiliterque punctatus, fronte aequali, thorace latiore, basi utrinque longitudinaliter impresso, apicem versus subangustato.* — Long. 2 lin.

Habitat in Cuba, Dom. Riehl.

Statura et summa affinitate praecedentis, rufo-piceus, nitidus. Anten-

nae capite thoraceque vix breviores, apicem versus leniter incrassatae, articulo tertio elongato. Caput thorace angustius, parce subtilissime punctulatum, fronte leviter convexa, aequali. Thorax coleopteris paululum latior, latitudine dimidio brevior, apicem versus leviter angustatus, lateribus leviter rotundatis, apice late emarginatus, parum convexus, parce subtiliter punctatus, basi ad angulum utrumque posteriorem longitudinaliter impressus. Elytra thorace paululum longiora, parce subtiliter punctata, stria suturali impressa. Abdomen apicem versus leniter attenuatum, obsolete subtiliter punctatum. Pedes rufi, tarsis testaceis.

Obs. Distinctus a praecedente corpore minus cylindrico, punctura multo parciore subtilioreque et fronte haud foveolata.

- \* 3. *L. striola*: *Piceus, nitidus, pedibus rufis, capite thoraceque crebre punctatis, hoc basi utrinque longitudinaliter impresso, lateribus parallelis, elytris parce subtiliter punctatis, disco striola longitudinali notatis.* — Long. 2 lin.

Habitat in Columbia et in Americae meridionalis insulis.

Elongatus, linearis, subdepressus, piceus, nitidus. Antennae capite thoraceque breviores, apicem versus parum incrassatae, articulo secundo tertioque subaequalibus, piceae, articulo ultimo ferrugineo. Palpi rufo-picei. Caput thorace paulo angustius, crebre subtilius punctatum, fronte parum convexa. Thorax coleopterorum prope latitudine, latitudine parum brevior, lateribus subrectis, parallelis, apice leviter emarginatus, perparum convexus, crebre minus subtiliter punctatus, basi utrinque sulculo lato longitudinali impressus. Elytra thorace parum longiora, parce subtiliterque punctata, stria suturali striolaque basi apiceque abbreviata disci impressa. Abdomen parce subtiliter punctatum. Pedes rufi.

- \* 4. *L. angustatus*: *Piceus, nitidus, pedibus rufis, capite thoraceque crebre, elytris abdomineque parcius punctatis, fronte bifoveolata, thorace postice utrinque impresso, basin versus angustato.* — Long. 2 lin.

#### T. II. f. 6.

Habitat in Madagascar, Dom. Goudot.

Elongatus, linearis, subdepressus, piceus, nitidus. Antennae capitibus thoracisque prope longitudine, crassae, pone medium subincrassatae, piceae, articulo secundo tertioque aequalibus. Caput thorace paulo angustius, crebrius punctatum, fronte antice utrinque impressa. Thorax ante medium coleopterorum latitudine, latitudine paulo brevior, lateribus leviter rotundatis, pone medium angustatus, basi truncatus, angulis posterioribus rectis, apice sat profunde emarginatus, subdepressus, crebrius punctatus, medio linea subtilissima canaliculatus, basi ad angulum posteriorem utrinque foveola oblonga impressus. Elytra thorace longiora, parcius punctata, stria suturali impressa. Abdomen minus crebre punctatum, lateribus punctorum maiorum serie notatum. Pedes rufi.

5. *L. linearis*: *Piceus, subnitidus, pedibus testaceis, parce punctatus, fronte aequali, thorace basi utrinque profunde longitudinaliter impresso, basin versus subangustato.* — Long. 1 $\frac{3}{4}$  lin.

Habitat in montibus prope Puerto Cabello in Columbia, Dom. Moritz.



Elongatus, linearis, subdepressus, piceus, minus nitidus. Antennae capite paulo longiores, apicem versus subincrassatae, articulo secundo tertioque aequalibus, rufo-piceae. Caput thoracis fere latitudine, subtiliter parcius punctatum, fronte aequali hand impressa. Thorax coleopterorum prope latitudine, latitudine parum brevior, lateribus subrectis, basin versus sensim subangustatus, apice leviter emarginatus, subdepressus, sat crebre punctatus, basi ad angulum posteriorem utrinque sulco antrorsum sensim evanescente longitudinali impressus. Elytra thorace paulo longiora, crebrius punctata, stria profunda suturali impressa. Abdomen parcius obsoletiusque punctatum, apice rufo-piceum. Pedes rufo-testacei.

\* 6. *L. tenellus*: Piceus, nitidus, pedibus rufis, capite, thorace elytrisque parce punctatis, fronte bifoveolata, thorace basi utrinque longitudinaliter impresso, leviter coarctato. — Long.  $1\frac{1}{2}$  lin.

Habitat in Columbia et in Americae meridionalis ins. St. Johannis et Puerto-Rico, Dom. Moritz.

Habitu omnino praecedentium, at multo minor, elongatus, sublinearis, subdepressus, piceus, nitidus. Antennae capitis thoracisque prope latitudine, apicem versus leviter incrassatae, articulo secundo tertioque subaequalibus, dilute rufo-piceae. Caput parce subtilissime punctatum, fronte utrinque impressa. Thorax coleopterorum prope latitudine, latitudine paulo brevior, lateribus modice rotundatus, basin leviter angustatus, angulis posterioribus rectis, antice subemarginatus, subdepressus, subtiliter sparsim punctatus, basi ad angulum posteriorem utrinque sulco longitudinali impressus. Elytra thorace sesqui fere longiora, parce subtilissime punctata, stria suturali impressa. Abdomen subtilissime punctato-coriaceum, segmentorum marginibus apiceque dilutioribus. Pedes rufo-testacei.

7. *L. exiguus*: Rufo-testaceus, nitidus, laevis, thorace basi utrinque impresso, basin versus angustato. — Long. 1 lin.

Habitat in ins. Puerto-Rico, Dom. Moritz.

Habitu omnino *L. tenelli*, at dimidio fere minor, rufo-testaceus, nitidus. Antennae capite paulo longiores, apice modice incrassatae, articulo secundo tertioque subaequalibus. Caput thorace parum angustius, laeve, fronte utrinque longitudinaliter impressa. Thorax antice coleopterorum prope latitudine, latitudine vix brevior, lateribus parum rotundatus, basin versus leviter angustatus, angulis posterioribus rectis, apice truncatus, subdepressus, laevis, basi ad angulum posteriorem utrinque foveola sat profunda impressus. Elytra thorace parum longiora, laevia, stria suturali obsoleta. Abdomen omnium subtilissime punctato-coriaceum. Pedes pallide testacei.

### Piestus Grav.

Grav. Mon. p. 223. — Laporte Etud. Ent. I. p. 126.  
Zirophorus Dalm Act. Holm. 1821. p. 371. Anal. Ent. p. 23.  
Trichocoryne Gray Animal Kingd. XIV. p. 306.

*Mandibulae dentatae.*

*Palpi maxillares articulo ultimo penultimo longiore.*

*Tibiae anticae externe crenatae.*

*Abdomen marginatum.*

Corpus oblongum, depressum, alatum. Caput thorace minus et angu-

stius, fronte inaequali, interdum cornubus duobus antrorsum directis horizontalibus munita, oculis parvis, parum prominulis. Labrum (t. V. f. 9. c.) transversum, apice truncatum, angulis anterioribus rotundatis, laciniis membranis angustis, linearibus. Mandibulae validae, porrectae, dentatae. Maxillae (t. V. f. 9. b.) malis membranaceis, interiore intus, exteriori apice barbatis, hac externe rotundato-dilatata. Palpi maxillares articulo primo minuto, secundo elongato, subarcuato, tertio hoc duplo brevior, quarto secundo paulo longiore, cylindrico, apice obtuse acuminato. Labium (t. V. f. 9. a.) mento antice rotundato, ligula membranacea, medio incisa, apice truncata, barbata. Palpi labiales breves, ligulam vix superantes, articulis longitudine subaequalibus. Antennae elongatae, interdum corpore longiores, articulo primo longiusculo, secundo parvo, obconico, tertio iterum elongato, clavato, apice saepe penicillato, sequentibus cylindricis, pubescentibus. Thorax coleopterorum latitudine, transversus, quadrangulus, basi medio productus, ab elytris remotus. Scutellum parvum, triangulare. Elytra truncata. Abdomen coleopteris vix angustius, minus elongatum, segmento penultimo reliquis paulo longiore, ultimo obtuso. Pedes breves, femoribus crassiusculis, tibiis angustis, anticis infra medium compressis, extus denticulatis, spina apicalis validiore instructis, intermediis infra medium extus spinulis setiformibus ciliatis, posticis muticis, tarsis minutis, articulo ultimo reliquos, inter se aequales, coniunctos longitudine prope superans.

#### Observationes.

1. Corpus tenuiter pubescens, capite, thorace elytrisque glaberrimis. Thorax linea media longitudinali integra exaratus, ad angulos posteriores utrinque sulculo magis minusve abbreviato impressus. Elytra striis vel sulcis 5 vel 6 laevibus cre-natisve exarata.

2. Huius generis speciem unam Fabricius inter Cucuios, alteram Olivier inter Oxytelos proposuerunt; tertiam generis Piesti typum descripsit Gravenhorst. Postea Dalman genus Zirophorus condidit, quod cum Leptochiro Germ. confusum Latreille recepit (Regn. Anim. V. p. 438.). Nomen Gravenhorstianum, Leptochiro probe distincto, Dom. Laporte restituit, forte Lucii Buquetii collectione edoctus, cui notiones vel de genere, vel de specie Fabriciana communicavimus.

3. Mas in nonnullis speciebus antennis longioribus et articulo earum tertio fortius penicillato distinctus; in plerisque sexus differentia latet.

4. Victus sub arborum putrescentium corticibus. Hoc genus Americae calidiori proprium videtur.

5. Larvam forte speciei cuiusdam Cayennensis Ziroph. longicornis sibi vocatam Dom. Lacordaire in silvis Conanensibus cum imagine observavit, long. 4 linearum, corpore curto, utrinque leviter attenuato, griseo, albedo-venoso, capite nigro, mandibulis validis, crassis, bifidis, dente superiore elevato ut in imagine; reliquis oribus partibus vix oculo armato distinguendis, exceptis palpis duobus, articulo aciculari acutissimo terminatis; corporis segmentis singulis papillis binis transversim positissimis instructis; incessu admodum lento; odore foetido. Specimina nonnulla diebus 8 praeterlapsis in folliculum oblongum sese condiderunt, at imago non prodiiit. (Nov. Annal. Mus. Hist. nat. II. p. 65.)

\* 1. *P. bicornis*: Fronte bispinosa, niger, nitidus, abdomine apice rufo, thorace laevi, subtiliter canaliculato, elytris striatis. — Long.  $4\frac{1}{3}$  lin.

Laporte Etud. Ent. I. 128. 1.

*Oxytelus bicornis* Oliv. Encycl. meth. Ins. VIII. 615. 11.

*Zirophorus fronticornis* Dalm. Act. Holm. 1821. 373. 1. Anal. Ent. 24. 1. t. 4. f. 1.

*Zirophorus striatus* Guér. Iconogr. Ins. t. 9. f. 12.

*Var. Thorace, pectore, elytris pedibusque rufo-castaneis.*

*P. Oxytelinus* Laporte Etud. Ent. I. 128. 2.

Habitat in Columbia, Dom. Gory, Surinami, Dom. Schüppel.

Niger, nitidus. Antennae corporis dimidiam longitudinem superantes, nigrae, ab articulo quinto inde pube griseo-flavescente dense vestitae, interne longius dense pilosae, quasi barbatae. Mandibulae porrectae, apice acuto, leviter introrsum verso, extus ante apicem denticulo minuto munitae. Caput laeve, fronte aequali, antice fovea rhombea, parum profunda, fundo medio elevato impressa, apice spinis duabus porrectis, rectis, approximatis, acutis, sat longis validisque armata. Thorax latitudine dimidio brevior, lateribus rectis, ad angulos posteriores oblique emarginatis, angulis anterioribus rotundatis, planatus, leviter canaliculatus, basi angulo utroque posteriore leviter impresso. Elytra thorace parum longiora, striis 5 simplicibus exarata. Abdomen segmentis singulis basi subtiliter punctatis, penultimo apice ultimoque toto rufis, rufo-pilosis. Pedes nigri, tarsis piceis.

\* 2. *P. Lacordairei*: *Fronte bispinosa, niger, nitidus, ano pedibusque piceis, thorace laevi, subtiliter canaliculato, elytris striatis.* — Long.  $3\frac{1}{2}$  lin.

*Laporte Etud. Ent. I. 129. 4.*

Habitat in Cayenna, a Dom. Com. Dejean nomine Zirophori gregarii communicatus.

*P. spinoso* longitudine fere aequalis, at multo angustior et minus depressus, niger, nitidus. Antennae corpore tertia parte breviores, subtiliter griseo-pubescentes, parce pilosellae, articulis tribus primis glabris. Mandibulae capitis longitudine, porrectae, sat validae, rectae, ante apicem leviter introrsum arcuatum dentatae. Caput fronte excavata, apice spinis duabus validis, porrectis, parallelis, rectis, acutis, mandibulas longitudine aequantibus, distantibus armata. Thorax latitudine dimidio brevior, lateribus rectis, ad angulos posteriores oblique subemarginatis, angulis anterioribus subrotundatis, planatus, leviter canaliculatus. Elytra thorace paulo longiora, striis 5 subtiliter crenatis exarata, extus medio striola brevi subtili impressa. Abdomen minus subtiliter crebre punctulatum, segmentorum summis marginibus anoque piceis. Pedes rufo-picei.

\* 3. *P. spinosus*: *Fronte bispinosa, rufo-castaneus, nitidus, capite abdomineque nigris, thorace medio subtiliter punctulato canaliculatoque, elytris punctato-striatis.* — Long.  $3\frac{1}{3}$  lin.

*Laporte Etud. Ent. I. 128. 3. t. 4. f. 3.*

*Cucuius spinosus* Fab. Syst. El. II. 93. 9. — Schönk. Syn. Ins. III. 53. 8.

*Zirophorus (?) spinosus* Dalm. Anal. Ent. 25. 3.

*Piastus oxytelinus* Ferty Delect. An. art. p. 33. t. 9. f. 9.

Habitat in Cayenna, Dom. Buquet.

Antennae corpore parum breviores, nigrae, subtiliter cinereo-pubescentes, parce pilosellae, articulis tribus primis castaneis, glabris. Mandibulae porrectae, capite paulo breviores, debiles, rectae, ante apicem dente inferiore



introrsum vergente armatae. Caput piceo-nigrum, nitidum, fronte subimpressa, canalicula media brevissima sulcoque arcuato inter antennis exarata, apice spinis duabus porrectis acutis valde distantibus armata. Thorax latitudine duplo fere brevior, lateribus rectis, ad angulos posteriores oblique emarginatis, angulis anterioribus subrotundatis, planus, medio subimpressus, leviter canaliculatus, basi foveola longitudinali impressus, medio foveolisque basalibus punctulatis, rufo-castaneus, nitidus. Elytra thorace paulo longiora, sulcis 5 sat profundis, fortiter punctatis exarata, externe striola abbreviata aliaque apicali impressa, rufo-castanea, nitida. Abdomen nigrum, nitidum, segmentis singulis basi crebre subtilius punctatis, penultimo apice ultimoque toto rufo-castaneis. Pectus rufo-castaneum. Pedes rufi.

Obs. Specimen a Fabricio descriptum in Museo regio Hafniensi asservatum, ipsum vidi, quare locum citatum ad hanc speciem spectare mihi compertum est, licet descriptio allata non omnino convenit.

4. *P. capricornis*: Fronte bidentata, rufo-piceus, nitidus, abdomine nigro-piceo, thorace subtiliter canaliculato, punctulato, elytris punctato-striatis. — Long.  $3\frac{3}{4}$  lin.

Laporte Etud. Ent. I. 129. 5. t. 4. f. 4.

Habitat in ins. Guadeloupe, collect. Dom. Chevrolat.

Statura omnino praecedentis, at paululum maior. Antennae corpore paulo longiores, parce pilosellae, nigrae, articulis 3 primis rufo-piceis, glabris. Mandibulae haud porrectae, dextra extus cornu falcato, sinistra dente obtuso armata. Caput fronte inter oculos foveolis duabus minusculis impressa, margine antico emarginato, utrinque dente prominente armato. Thorax latitudine duplo prope brevior, lateribus rectis, ad angulos posteriores oblique emarginatis, angulis anterioribus subrotundatis, planus, disco sparsim punctatus, tenuiter canaliculatus. Elytra thorace longiora, striis 5 punctatis exarata, externe striola abbreviata media aliaque apicali impressa. Abdomen crebrius punctulatum, nigro-piceum. Tibiae anticae obsolete crenatae.

Obs. A Dom. Laporte Cayenna pro huius insecti loco natali indicata, an rite?

5. *P. fulvipes*: Fronte mutica, rufo-piceus, nitidus, pedibus fulvis, thorace subtilissime punctulato, canaliculato, elytris 5-sulcatis. — Long.  $3\frac{1}{2}$  lin.

Habitat in ins. Guadeloupe, Dom. Chevrolat.

P. penicillato maior, magis elongatus, rufo-piceus, nitidus. Antennae corpore paulo longiores, piceo-rufae, dense pubescentes, articulis tribus primis glabris, primo interne acute dentato, dente pilorum fasciculo penicillato. Caput fronte antice truncata, inter oculos utrinque profunde oblique impressa, spatio interiecto triangulari, leviter elevato, laterali incrassato, tumido. Thorax latitudine vix dimidio brevior, lateribus rectis, ad angulos posteriores oblique emarginatis, angulis anterioribus leviter rotundatis, minus depressus, subtilissime punctulatus, sat fortiter canaliculatus, basi utrinque ad angulos posteriores sulculo brevi profundo impressus. Elytra thorace paulo longiora, striis 5 sat profundis, impunctatis exarata, interstitio primo medio bipunctato, extus ante medium lineola abbreviata tenui impressa. Abdomen crebre subtiliter punctulatum, ano segmentorumque singulorum margine postico dilutiore. Pedes testaceo-rufi.

- \* 6. *P. penicillatus*: *Fronte mutica, niger, nitidus, ano pedibusque piceis, thorace laevi, subtiliter canaliculato, elytris 5-striatis.* — Long.  $2\frac{1}{3}$  — 3 lin.

*Ziophorus penicillatus* Dalm. Act. Holm. 1821. 373. 2. Anal. ent. 24. 2. t. 4. f. 2.

Habitat in Americae meridionalis insulis: St. Barthelemy (Dalm.), Puerto-Rico. Dom. Moritz.

Niger, nitidus. Antennae corpori longitudine aequales vel eodem paulo longiores, articulo primo tuberculato, tuberculo pilorum fasciculo penicillato. Mandibulae mediocres, apice arcuatae bidentataeque. Caput fronte apice truncata, inter oculos utrinque sat profunde oblique impressa, spatio medio triangulari elevato. Thorax latitudine plus dimidio brevior, lateribus rectis, ad angulos posteriores oblique emarginatis, angulis anterioribus rotundatis, planatus, subtilius canaliculatus, basi utrinque ad angulum posteriorem sulculo brevi sat profundo impressus. Elytra thorace paulo longiora, striis 5 sat profundis, in fundo obsoletius punctatis excarata, interstitiis subtilissime obsolete punctulatis, extus medio lineola brevi tenuique impressa. Abdomen segmentis singulis basi utrinque subtiliter punctatis, penultimo apice ultimoque piceis. Pedes picei.

- \* 7. *P. erythropus*: *Fronte mutica, piceus, nitidus, pedibus rufis, thorace subtilissime punctulato, canaliculato, elytris 5-striatis.* — Long.  $2\frac{1}{3}$  lin.

*Trichocoryne striata* Gray An. Kingd. XIV. p. 306. t. 32. n. 5. (forte)

Habitat in Cuba, Dom. Riehl et collect. Dom. Chevrolat.

Habitu omnino *P. minuti*, at sesqui maior, saturate nigro-piceus, nitidus. Antennae longitudine corporis, nigricantes, subtiliter pubescentes, articulis tribus primis rufo-piceis, glabris, primo pilorum fasciculo introrsum vergente penicillato. Caput fronte antice truncata, inter oculos utrinque oblique impressa, spatio interiecto triangulari subelevato, plica transversali subarcuata leviter elevata a vertice distincta. Thorax latitudine dimidio brevior, lateribus subrectis, ad angulos posteriores oblique emarginatis, minus planus, parce subtilissime punctatus, sat fortiter canaliculatus, basi utrinque ad angulos posteriores sulculo brevi sat profunde impressus. Elytra thorace paulo longiora, striis 5 sat profundis, subtilius punctatis exarata, interstitiis subtiliter punctulatis, externe apice, nonnunquam etiam ante medium, striola tenui impressa. Abdomen sat crebre punctatum, segmentis 4 primis supra medio laevioribus. Pedes testaceo-rufi.

Obs. A *P. minuto* elytrorum striis punctatis facile distinguendus, a *P. penicillato* colore, corpore minus plano, elytrorum striis fortius punctulatis etc. distinctus.

- \* 8. *P. minutus*: *Fronte mutica, castaneus, nitidus, thorace laevi, subtilius canaliculato, elytris 5-striatis.* — Long. 2 lin.

*Piestus sulcatus* Laporte Etud. Ent. I. 129. 6. (forte)

Habitat in Columbia, Dom. Moritz.

Habitu omnino *P. penicillati*, at dimidio fere minor, castaneus, nitidus, capite thoraceque obscurioribus, piceis. Antennae maris corporis prope longitudine, articulo primo pilorum fasciculo penicillato, feminae corpore

breviores, articulo primo simplice. Caput thorace angustius, fronte utrinque leviter impressa et profunde oblique arcuatim sulcata, vertice plicula subtili elevata transversa instructo. Thorax latitudine plus dimidio brevior, lateribus subrectis, ad angulos posteriores oblique emarginatis, fere planus, laevis, canalicula tenui exaratus, basi ad angulum posteriorem utrinque sulculo brevi sat profundo impressus. Elytra thorace parum longiora, striis 5 simplicibus sat profundis exarata, externe ad apicem striae sextae rudimento notata. Abdomen supra segmentis singulis utrinque subtiliter punctatis, subtus totum crebre distinctius punctatum. Pedes rufi.

- \* 9. *P. pygmaeus*: *Fronte mutica, niger, nitidus, ore, antennis pedibus, elytrisque rufo-castaneis, thorace subtiliter punctato, canaliculato, elytris 5-striatis.* — Long.  $1\frac{1}{4}$  — 2 lin.

*Laporte Etud. Ent. I. 130. 8.*

Habitat in Brasilia; in Columbia Dom. Moritz.

Praecedenti affinis, at paulo minor, praecipue angustior. Antennae maris corpore longiores, feminae paulo breviores, fuscae, pubescentes, articulis tribus primis rufis, glabris, primo pilorum penicillo in mare crasso, in femina tenui instructo. Caput nigrum, ore castaneo, fronte utrinque subimpressa, inter antennis linea transversali, medio fortiter sursum arcuata, exarata. Thorax latitudine dimidio brevior, lateribus subrectis, postice ad angulos posteriores oblique emarginatis, angulis posterioribus ipsis prominulis, acutiusculis, anterioribus subrotundatis, planus, subtiliter punctulatus, canalicula media sat profunda et utrinque sulco profundo longiore longitudinali impressus, niger, nitidus, lateribus nonnunquam castaneis. Scutellum nigro-piceum, nitidum. Elytra thorace paulo longiora, striis 5 simplicibus sat profundis exarata, externe ad apicem striae sextae rudimento instructa, interstitiis subtilissime punctulatis, rufo-castanea, nitida. Abdomen supra subtiliter, subtus utrinque evidentius punctatum, nigro-piceum, nitidum, segmentorum marginibus anoque rufo-castaneis. Pectus piceum. Pedes rufi.

- \* 10. *P. sulcatus*: *Fronte mutica, castaneus, nitidus, thorace subtilissime punctulato, canaliculato, elytris 6-sulcatis.* — Long.  $2\frac{1}{3}$  lin.

*Grav. Mon. 224. 1. — Lepell. et Serv. Encycl. meth. Ins. X. p. 123.*

Habitat in Parà Brasiliae, Sieber.

Statura omnino *P. minuti*, castaneus, nitidus. Antennae maris corporis longitudine, feminae paulo breviores, articulo primo in mare pilorum fasciculo instructo, feminae nudo. Caput thorace angustius, subtiliter obsolete punctulatum, fronte inter antennis utrinque profunde arcuatim impressa. Thorax latitudine plus dimidio brevior, lateribus leviter rotundatis, ad angulos posteriores fortiter oblique emarginatis, angulis anterioribus subrotundatis, planatus, subtilissime punctulatus, linea longitudinali media sat profunda exaratus, basi utrinque sulco longiore sat forti longitudinali impressus. Elytra thorace paulo longiora, sulcis sex integris, sat profundis, subtiliter crenatis exarata, interstitio marginali subcarinato. Abdomen sat crebre, minus subtiliter punctatum, ano dilutiore. Pedes rufi.



11. *P. Mexicanus*: *Fronte mutica, niger, parum nitidus, pedibus rufo-piceis, capite thoraceque punctatis, hoc canaliculato, utrinque sulcato, elytris punctato-sulcatis. — Long. 2½ lin.*

*Laporte Etud. Ent. I. 130. 7.*

Habitat in Mexico, collect. Dom. Chevrolat.

Subdepressus, niger, parum nitidus. Antennae corpore breviores, nigrae, subtiliter pubescentes, articulis 3 primis rufo-piceis, glabris. Palpi piceo-rufi. Caput crebre minus subtiliter punctatum, fronte inter oculos utrinque leviter oblique impressa, antice rotundata. Thorax latitudine plus dimidio brevior, antrorsum angustatus, lateribus rotundatis, pone medium profunde oblique emarginatis, sinu utrinque denticulo prominulo terminato, sat crebre subinaequaliter minus subtiliter punctatus, canalicula sat profunda media et utrinque sulco sat lato laterali, ultra medium adscendente impressus. Elytra thorace sesqui longiora, dorso sulcis 6 punctatis exarata, interstitiis angustis, elevatis, laevibus. Abdomen crebre punctatum, ano, segmentorumque singulorum margine postico rufo-piceo. Pedes rufo-picei.

#### O b s e r v a t i o.

Eleusis Laporte (Etud. Ent. I. p. 131.) Piesto affinis videtur, mihi omnino incognita et dubia, hoc loco forte inserenda. Genus intermedium descriptum est inter Leptochirum et Piestum, antennis brevioribus ut in illo, at apicem versus incrassatis, palpis articulo brevi, tenui acuminatoque terminatis, corpore depresso, capite maximo, quadrato, mandibulis porrectis, arcuatis, thorace brevissimo, capite minore, antice truncato, postice rotundato, coleopteris quadratis. Species singula, *E. tibialis*, nigra, capite fortiter punctato, thorace 4-foveolato, elytris abdomineque laevibus, tibiis et oris partibus rufescentibus, long. 3 lin. a Goudotio ex ins. Madagascar allata.

### **Prognatha Latr.**

*Latr. Regn. Animal. IV. p. 439. Nov. Annal. Mus. Hist. nat. I. p. 89.*

*Siagonium Kirby et Spence Introd. 1. t. 1. f. 3. — Curt. Britt. Ent. 1. t. 23.*

*Mandibulae edentatae, maris cornutae.*

*Palpi maxillares articulo ultimo penultimo duplo longiore.*

*Tibiae anteriores spinulosae.*

*Abdomen marginatum.*

Corpus elongatum, depressum, alatum. Caput, praesertim maris, sat magnum, fronte antice rotundata, medio immarginata, in mare prope oculum utrumque cornu horizontali, antrorsum porrecto, acuto armatum, oculis parvis, rotundatis, leviter prominulis. Labrum transversum, medio leviter sinuatum, laciniis membraneis parvis, angustis, linearibus. Mandibulae falcatae, acutae, parum prominentes, feminae simplices, maris extrorsum dilatatae, cornu magno, (mandibula ipsa maiore), horizontali, falcato, mandibulam simulante armatae. Maxillae (t. V. f. 10. b.) mala interiore angusta, exteriori semicirculari, utraque membranea, apice ciliata. Palpi maxillares articulo primo minuto, secundo modice elongato, tertio hoc multo brevior, quarto iterum elongato, secundo paulo longiore, apice subacuminato. Labium (t. V. f. 10. a.) mento transversum, lateribus leviter angustato, apice truncato, ligula membranea, biloba. Palpi labiales breves, ligulam vix superantes, articulis tribus subaequalibus, ultimo subacuminato. Antennae maris fere corporis longitudine, feminae capite thoraceque paulo longiores, articulo primo reliquis

crassiore, secundo tertioque elongato-obconicis, reliquis cylindricis. Thorax quadrangulus, basi coleopteris arcte applicatus. Scutellum parvum, triangulare. Elytra truncata. Abdomen coleopterorum latitudine, lineare, marginatum. Pedes breviusculi, femoribus praecipue posticis leviter incrassatis, tibiis angustis, anticis margine exteriori subtiliter serratis, spina apicali maiore terminatis, tarsis minutis, articulis quatuor primis perbrevibus, subtus setulosus.

#### O b s e r v a t i o n e s.

1. Corpus parce tenuiterque pubescens, capite thoraceque glabris, thorace aequali, haud canaliculato, elytris subseriatim punctatis.

2. Mas capite maiore, fronte excavata, utrinque cornuta, mandibulis pariter cornutis, antennis longioribus distinctus.

3. Victus sub arborum cortice.

X \* 1. *P. quadricornis*: *Nigro-picea, nitida, elytris parce seriatim punctatis, castaneis.* — Long. 2 lin.

*Boisd. et Lacord. Faun. Ent. Paris. I. 454.*

*Siagonium 4-corne Kirby et Spence Introd. I. t. 1. f. 3. — Curt. Britt. Ent. I. t. 23.*

*Prognatha rufipennis Blondel Annal. Scienc. Nat. X. p. 414. t. 18. f. 14 — 15. — Guér. Iconogr. Regn. An. Ins. t. 10. f. 1.*

Habitat in Anglia, Dom. Hope, Fontibus Bellaqueis Galliae, Dom. Aubé, Halae Saxonum, Dom. Alexander Klug.

*Picea, nitida.* Antennae maris corpore tertia parte breviores, feminae eodem dimidio vix longiores, pubescentes, rufae. Caput maris thoracis magnitudine, parce subtiliter punctatum, fronte late leviter excavata, antice laeviore, cornubus divergentibus, frontis apicem paulo superantibus, feminae thorace minus et angustius, crebrius minus subtiliter punctatum, fronte antice rotundata, lateribus marginata, inter antennis utrinque impressa. Thorax coleopteris paulo angustior, latitudine summa paulo brevior, lateribus leviter rotundatis, basin versus angustatus, basi coarctatus, apice leviter bisinuatus, basi truncatus, angulis posterioribus subrectis, subdepressus, minus confertim subtilius licet profunde punctatus, aequalis. Elytra thorace sesqui longiora, subdepressa, praeter striam suturalem integram striis 4 obliquis postice abbreviatis, punctatis notata, lateribus et apice passim disperse punctata, rufo-castanea, nitida. Abdomen crebrius punctatum, apice rufo. Pedes rufi.

\* 2. *P. humeralis*: *Nigro-picea, elytris crebre striatim punctatis, macula humerali rufo.* — Long. 2 lin.

*Siagonium humerale Germ. Faun. Ins. Europ. XVIII. 6.*

Habitat in Hungaria montana, Dom. Dr. Frivaldsky.

Statura et summa similitudo praecedentis. Corpus nigrum, nitidum. Antennae maris corpore paulo longiores, feminae paulo breviores, obscure rufae. Caput magnum, in utroque sexu thorace paulo latius, subtiliter punctatum, fronte maris parcius subtiliusque punctata, anterieus leviter declivi, sublaevi, postice transversim impressa, vertice fere plicato-elevato, cornubus parallelis, frontis apicem paulo superantibus, feminae crebrius punctata, anterieus laeviore, utrinque fortius impressa, antice rotundata. Thorax coleo-

pteris parum angustior, latitudine paulo brevior, lateribus rotundatus, basi truncatus, angulis posterioribus acutiusculis, depressus, parcius disperse minus fortiter at sat profunde punctatus, aequalis. Elytra thorace plus sesqui longiora, depressa, praeter striam suturalem integram striis senis, apice paululum abbreviatis, punctatis notata, lateribus crebrius subtiliter punctata, subtilissime longitudinaliter strigosa, nigro-picea, plaga magna humerali rufa. Abdomen laevigatum, ano piceo. Pedes rufo-picei.

Obs. Antennis in utroque sexu longioribus, thorace basi fortius coarctato, elytris longioribus, crebrius punctato-striatis, abdomine laevigato, etc. a praecedente satis distincta.

### **Isomalus Er.**

*Mandibulae muticae.*

*Palpi maxillares articulo ultimo penultimo minore.*

*Tibiae muticae.*

*Abdomen subtiliter marginatum.*

Corpus elongatum, lineare, valde depressum, planum, forte apterum. Caput porrectum, magnum, quadratum, in utroque sexu muticum, fronte antice rotundata, immarginata, oculis parvis, rotundatis, parum prominulis. Labrum minutum, transversum, angulo apicali utroque lacinia angusta membranea aucto. Mandibulae in utroque sexu minutae, muticae, vix prominentes, basi latiusculae, sensim acuminatae, introrsum falcatae. Maxillae (t. V. f. 11. b.) mala interiore angusta exterioreque semicirculari membranais, hac basi cornea. Palpi maxillares breviusculi, articulo primo minimo, secundo tertio dimidio brevior, quarto secundo parum longiore, subacuminato. Labium (t. V. f. 11. a.) mento brevi, antrorsum angustato, utrinque bisinuato, apice rotundato, ligula lata membranea, apice trisinuata, ciliata, lobis lateralibus divergentibus, sinu intermedio setis duabus, lateralibus pilorum fasciculo instructis. Palpi labiales ligula vix longiores, articulis decrescens, ultimo subacuminato. Antennae capite thoraceque haud longiores, filiformes, parce pilosellae, articulo primo paulo crassiore, secundo tertioque reliquis paulo tenuioribus, 4 — 10 sensim paulo crassioribus, ultimo ovato. Thorax basin versus fortiter angustatus, lateribus subangulatus, ab elytris remotus. Scutellum semicirculare. Elytra truncata. Abdomen coleopterorum latitudine, lineare, subtiliter marginatum. Pedes breves, femoribus subcompressis, tibiis angustis, omnibus simplicibus, extus versus apicem setulis dense ciliatis, tarsis minutis.

### **Observationes.**

1. Corpus glabrum. Caput posterius secundum marginem lateralem sulculo longitudinali exaratum. Thorax haud canaliculatus, lateribus saepius denticulatus. Elytra laevia, stria suturali nulla. A genere praecedente praeter oris structuram mandibulis fronteque in utroque sexu muticis, thorace ab elytris remoto, abdomine subtiliter marginato, tibiis muticis etc. discedit.

2. Genus Chasolium Laporte (Etud Ent. I. p. 132.) vix ab Isomalo distinctum, at tibiis anticis fortiter angulatis, externe subtilissime denticulatis, mandibulis prominentibus, thorace medio excavato, lateribus fortiter elevato descriptum, quam ob rem nomen illud recipere dubitavi. Species descripta Madagascarinensis, Ch. Ernestini, cum Isomalo complanato convenire videtur.

3. Mas capite paulo maiore a femina differe videtur. Reliquae differentiae latent.



- \* 1. *I. complanatus*: Niger, nitidus, capite thoraceque denticulato subtilissime canaliculatis, elytris testaceis, pedibus rufis. — Long.  $1\frac{3}{4}$  lin.

T. II. f. 7.

Habitat in Madagascar, Dom. Goudot.

Corpus nigrum, nitidum. Antennae piceae, apice nigricantes. Caput magnum, planum, subtilissime punctatum, medio subtiliter obsoleteque canaliculatum, utrinque stria profunda marginali exaratum. Thorax antice coleopterorum latitudine, apice late submarginatus, lateribus ante medium subrectis, a medio inde fortiter angustatus, spatio aequali inter medium et basin dente prominulo angulatus, basi truncatus, planus, subtilissime obsoleteque punctulatus, subtiliter obsolete canaliculatus, angulo anteriore utroque striola brevi sat lata notato. Elytra thorace paulo longiora, parce obsolete punctata, plana, testacea, nitida, margine omni suturaque infuscat. Abdomen planum, laeve, subnitidum, segmentis singulis tenuiter piceo-marginatis. Pedes rufo-picei.

- \* 2. *I. humilis*: Niger, nitidus, antennis pedibusque piceis, elytris discoluteis, thorace obsolete denticulato. — Long.  $1\frac{1}{2}$  lin.

Habitat in Brasilia, Dom. Melly.

Corpus valde depressum, nigrum, nitidum. Antennae capite thoraceque paulo breviores, piceae. Os rufescens. Palpi testacei. Caput thorace paululum latius, laeve, pone oculos stria marginali longitudinali profunda impressum. Thorax antice coleopteris paulo angustior, basin versus angustatus, lateribus rotundatis, pone medium obsolete denticulatis, laevis, dorso late subimpresso, medio obsolete subtiliter canaliculato, apice utrinque foveola minuta impressus. Elytra thorace paulo longiora, laevia, nitida, luteo-testacea, undique fusco-marginata. Abdomen laeve, segmentis singulis piceo-marginatis. Pedes picei, tibiis tarsisque testaceis.

- \* 3. *I. interruptus*: Supra niger, nitidus, subtus piceus, capite thoraceque obsolete denticulato laevibus. — Long.  $1\frac{3}{4}$  lin.

Habitat Carthagenae in Columbia, Dom. Gory.

Corpus planum, supra nigrum, nitidulum, subtus rufo-piceum. Antennae capitis thoracisque prope longitudine, piceae, articulis primis rufis. Oris partes rufae. Caput thoracis latitudine, ad oculorum marginem anteriorem sulculo brevi longitudinali utrinque exaratum, ceterum laevissimum. Thorax antice fere coleopterorum latitudine, latitudine summa haud brevior, lateribus rotundatus, basin versus fortiter angustatus, pone medium obsolete denticulatus, laevissimus, apice utrinque prope angulos anteriores leviter impressus. Elytra thorace sesqui longiora, laevia. Abdomen laeve, segmentis singulis piceo-marginatis. Pedes rufi.

4. *I. bicolor*: Nitidus, testaceo-rufus, capite, thorace, elytris, ano femoribusque anterioribus nigris, thorace mutico. — Long.  $2\frac{1}{2}$  lin.

Habitat in Mexico, Mus. Dom. Chevrolat.

Reliquis maior. Antennae capite sesqui longiores, pilosellae, piceae. Os piceum. Caput thoracis latitudine, subquadratum, parce subtiliter pun-

ctatum, punctis in media fronte crebrioribus et fortioribus, utrinque pone oculos sulco longitudinali marginali lato profundoque impressum, nigrum, nitidum. Thorax antice coleopterorum latitudine, lateribus ultra medium subrectis parallelisque, dein leviter rotundatis, basin versus fortiter angustatus, angulis posterioribus paululum prominulis, subrectis, planus, parce subtilissime punctatus, apice utrinque prope angulos anteriores foveola maiore, basi ad angulos posteriores foveola minore minus profunde impressus, niger, nitidus. Scutellum nigrum. Elytra thorace tertia parte longiora, laevia, nigra, nitida. Abdomen laeve, segmentis 4 primis testaceo-rufis, ultimis duobus nigris. Pectus cum pedibus posticis testaceo-rufum. Pedes antici rufo-picei, femoribus piceo-nigris.

\* 5. *I. testaceus*: *Testaceus, nitidus, elytris apice picescentibus, thorace mutico*. — Long.  $1\frac{1}{3}$  lin.

Habitat in Madagascar, Dom. Goudot.

I. complanato duplo minor. Antennae capite sesqui longiores, rufo-testaceae. Caput thoracis latitudine, oblongum, planum, parce subtilissime punctatum, punctis in media fronte crebrioribus et distinctioribus, lateribus posteriori sulculo longitudinali marginali longiore exaratum, rufo-testaceum, nitidum. Thorax antice coleopteris parum angustior, latitudine summa haud brevior, lateribus rotundatis, a medio inde basin versus fortiter angustatus, parce omnium subtilissime punctulatus, disco late impresso, rufescenti-testaceus, nitidus. Scutellum testaceum. Elytra thorace tertia parte longiora, parce subtilissime punctata, nitida, testacea, apice picescentia. Abdomen laeve, piceo-testaceum, nitidum, segmentis singulis pallido-marginatis. Pedes dilute testacei.

## **Hypotelus Er.**

*Mandibulae muticae.*

*Palpi maxillares articulo ultimo penultimo duplo longiore.*

*Tibiae anteriores spinulosae.*

*Abdomen marginatum.*

Corpus minutum, lineare, subdepressum. Caput thorace paulo minus, porrectum, fronte antice rotundata, immarginata, oculis rotundatis, prominulis. Labrum breve, transversum, medio sinuatum ciliatumque setulis rigidis, laciniis membraneis parvis, subtiliter ciliatis. Mandibulae parvae, haud prominentes, muticae, interne ciliatae. Maxillae (t. V. f. 12. b.) malis oblongis, interiore membranea, apice ciliata, exteriore apice extrorsum oblique truncata spisseque barbata. Palpi maxillares articulo primo minuto, secundo maiusculo, obconico, tertio hoc paulo brevior, quarto his coniunctis aequali, cylindrico, apice obtuse acuminato. Labium (t. V. f. 12. a.) mento subsemicirculari, ligula lata, transversa, membranea, medio leviter incisa. Palpi labiales breves, crassiusculi, articulo primo maiore, secundo tertioque minutis. Antennae capitis thoracisque fere longitudine, crassiusculae, articulo primo fusiformi, medio introrsum subdentato, reliquis obconicis, 5—10 subaequalibus, ultimo ovato. Thorax quadrangulus, basi coleopteris applicatus. Scutellum triangulare. Elytra truncata. Abdomen coleopterorum latitudine, lineare, marginatum. Pedes breviusculi, femoribus leviter incrassatis, tibiis

anticis apicem versus sensim subdilatatis intermediisque angustis margine exteriori spinulis subtilibus ciliatis, posticis prope apicem nonnullis tantummodo spinulis munitis, tarsis minutis, articulo ultimo reliquis coniunctis vix aequali.

### Observationes.

1. Corpus parce pubescens, capite, thorace elytrisque glabris. Thorax haud canaliculatus. Elytra stria suturali impressa. Habitus fere Prognathae, a qua discedit palporum maxillarium articulo tertio brevi, maxillarum mala exteriori haud dilatata, mento apice rotundato, ligula minus fortiter biloba. Mandibularum mala exteriori parallela et paraglossis ciliarum serie distinctis ab omnibus reliquis huius tribus generibus discrepat et ad Oxytelinos accedit.

2. Sexus differentia latet.

\* 1. *H. pusillus*: *Niger, nitidus, elytris pedibusque testaceis*. — Long. 1 lin.

T. II. f. 8.

Habitat in Brasilia.

*Niger, nitidus*. Antennae capitis thoracisque prope longitudine, fuscae, articulis tribus primis rufis. Os rufum. Caput thorace paulo angustius, basi leviter constrictum, subtiliter punctatum, fronte parum convexa, inter antennis utrinque obsolete impressa. Thorax medio coleopterorum fere latitudine, latitudine parum brevior, postice leviter angustatus, lateribus rotundatus, basi apiceque truncatus, angulis anterioribus subrotundatis, posterioribus obtusis, lateribus subtiliter marginatis, parum convexus, subtiliter licet sat distincte punctatus, lineola subtilissima canaliculatus. Elytra thorace parum longiora, sat distincte minus crebre punctata, stria suturali impressa, testacea, nitida, margine anteriore, exteriori suturaque tenui spatio infuscatis. Abdomen subtiliter obsoleteque parce punctatum, parce pilosellum, apice rufo-piceum. Pedes rufo-testacei.

2. *H. praecox*: *Testaceus, nitidus, capite nigro, thorace rufo*. — Long. 1 lin.

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Praecedente haud longior at paulo latior. Antennae capite thoraceque paulo longiores, validiusculae, pilosellae, fuscae, articulis tribus primis testaceis. Palpi testacei. Caput thorace paulo angustius, fronte levissime convexa, parce subtiliter punctata, inter antennis utrinque leviter foveolata, nigrum, nitidum. Thorax coleopteris haud angustior, latitudine summa paulo brevior, lateribus fortiter rotundatis, a medio inde basin versus fortiter angustatus, basi apiceque truncatus, angulis anterioribus obtusiusculis, posterioribus obtusis, subdepressus, parcius subtiliter punctatus, linea tenui longitudinali subtilissime canaliculatus, testaceo-rufus, nitidus. Elytra thorace sesqui longiora, parcius subtiliter punctulata, stria suturali impressa, nitida, testacea, apice picescentia. Abdomen parce omnium subtilissime punctatum, piceo-testaceum, nitidum. Pectus nigro-piceum. Pedes rufo-testacei.



## Tribus nona.

### **Phloeocharini.**

*Stigmata prothoracica occulta.*

*Coxae posticae transversae, anticae conicae, exsertae.*

*Trochanteres postici fulcrantes.*

*Ocelli nulli.*

---

Labrum transversum, corneum, simplex vel apice limbo membraneo prominulo instructum.

Mandibulae breviusculae, falcatae, parum exsertae, intus lacinia membranea ciliata fimbriatae.

Maxillae mala interiore rhombea interne, exteriori oblonga apice barbata.

Palpi maxillares articulo primo parvo, filiformes Olisthaero;

articulo tertio incrassato, quarto parvo, subulato Phloeocharidi.

Mentum corneum, perbreve, transversum, apice truncatum.

Ligula membranea, biloba.

Paraglossae ligulae aequales, connatae Phloeocharidi; ligulae longe breviores Olisthaero.

Palpi labiales filiformes.

Antennae sub frontis margine laterali insertae, 11-articulatae, rectae, filiformes.

Caput porrectum, thoraci immissum, basi haud aut vix constrictum.

Thorax coleopterorum latitudine iisque basi applicatus, subtus spatio pone coxas abque integmento corneo, membrana obducto, coxis obtecto.

Scutellum distinctum.

Elytra pectoris longitudine.

Abdomen elongatum, marginatum, segmentis sex distinctis, septimo retractili, rarius conspicuo.

Pedes breviusculi, coxis anticis conicis, exsertis, intermediis approximatis, posticis transversis, tibiis spinulis terminalibus subtilioribus, tarsis 5-articulatis,

anticis dilatatis Phloeocharidi;

simplicibus Olisthaero.

## O b s e r v a t i o n e s.

1. Duo tantummodo genera neque ultra tres species continens haec tribus ab omnibus satis distincta. Proxima videtur Omalinis, at ocelli nulli, maxillarum mala interior mutica, elytra pectore haud longiora et abdomen late marginatum. Affinis praecedentibus etiam duabus, at coxae posticae fulcrantes et labrum laciniis destitutum, ab Oxytelinis praeterea abdominis segmento septimo retractili, a Piestinis hoc segmento exserendo et coxis anticis conicis distincta. Quodammodo ad Tachyporinos etiam (imprimis Boletobium) se vergit, at stigmata antica occulta.

2. Sexus differentia vix observanda.

3. Victus sub arborum cortice.

**Olisthaerus Dej.***Palpi maxillares filiformes.*

Corpus elongatum, lineare, depressum, alatum. Caput exsertum, basi constrictum, ore attenuato, oculis parvis, rotundatis, vix prominulis. Labrum transversum, apice truncatum, simplex. Mandibulae leviter prominentes, muticae, apice falcatae. Maxillae (t. V. f. 4. b.) stipite modice elongato, mala exteriore cornea, apice membranacea barbataque, interiore membranacea, interne barbata. Palpi maxillares articulo primo minuto, secundo modice elongato, subarcuato, apicem versus sensim incrassato, tertio hoc parum brevior, leviter obconico, quarto hoc paulo minore, acuminato. Labium (t. V. f. 4. a.) mento brevi, transversum, partes labii membraneas parum obtegens, ligula rotundata, apice bifida, paraglossis ligula multo brevioribus, apice liberis. Palpi labiales articulis duobus primis cylindricis, inter se aequalibus, tertio oblongo-ovato, illis singulis minore. Antennae apicem versus sensim leviter incrassatae, articulis tribus primis obconicis, longitudine subaequalibus, primo paulo crassiore, quarto quintoque subglobosis, sequentibus sensim paulo magis transversis, ultimo breviter ovato. Thorax coleopterorum latitudine, basin versus subangustatus, basi apiceque truncatus. Scutellum sat magnum, triangulare. Elytra apice interne recte, externe oblique truncata. Abdomen elongatum, lineare, lateribus marginatum. Pedes breviusculi, intermedii basi approximati, tibiis parcius spinulosi, tarsi parum elongati, articulis 4 primis sensim leviter aequaliterque decrescentibus, quinto his coniunctis paulo brevior.

## O b s e r v a t i o n e s.

1. Caput et thorax laevisima. Elytra substriata, stria suturali profundiore impressa. Abdomen pubescens parceque pilosellum.

2. Victus sub Pinuum putrescentium cortice.

\* 1. *O. megacephalus*: *Rufo-castaneus, nitidus, capite thoracis latitudine, piceo.* — *Long. prope 3 lin.*

*Omalius megacephalum* Zetterst. *Faun. Lappon. I. 56. 17. Ins. Lappon. 54. 25.*

Habitat in Lapponia, Dalecarlia.

Corpus laete castaneum, glabrum, pernitidum, abdomine obscuriore, pubescente. Antennae capite thoraceque parum breviores, rufo-ferrugineae. Caput thoraci latitudine vix inferius, inter antennas linea levi arcuata impressum, ceterum laevisimum, fronte picea. Thorax coleopterorum latitudine, latitudine dimidio brevior, lateribus leviter rotundatis, pone medium leviter subsinuato-angustatus, apice late submarginatus, basi truncatus, an-

gulis posterioribus rectis, subdepressus, laevisissimus. Scutellum laevisissimum. Elytra thorace tertia parte longiora, crebrius longitudinaliter substriata. Abdomen supra parcius et obsoletius, subtus crebrius sat distincte punctatum, parcius lutescenti-pubescent, segmentis superioribus 5 primis basi nigris. Pedes magis rufi.

Obs. Utrumque sexum novi, at differentiae sexuales me fugiunt.

\* 2. *O. substriatus*: *Rufus, nitidus, capite thorace angustiore elytris piceis.* — Long.  $2\frac{2}{3}$  lin.

*Omalium substriatum* Gyll. Ins. Suec. II. 232. 29. — Sahlb. Ins. Fenn I. 288. 27. — Zetterst. Faun. Lappon. I. 55. 16. Ins. Lappon. 53. 24.

Habitat in Lapponia, Dalecarlia.

Antennae capitis thoracisque prope longitudine, rufae. Caput thorace dimidio angustius, depressum, laevisissimum, inter antennas transversim impressum, nigro-piceum, pernitidum, ore rufo. Thorax coleopterorum latitudine, latitudine dimidio brevior, lateribus leviter rotundatis, ad summam basin subsinuatis, basi apiceque truncatus, angulis posterioribus rectis, subdepressus, laevisissimus, basi prope angulum posteriorem utrinque obsolete foveolatus, rufus, pernitidus. Scutellum rufum. Elytra thorace sesqui longiora, substriata, nigro-picea, pernitida, margine summo apicali rufescente. Abdomen supra infraque sat crebre punctatum, parcius pubescens, rufum, nitidum. Pectus rufum. Pedes rufi.

Mas abdominis segmento anali minore et magis acuminato, septimo prominulo distinguendus.

Obs. A praecedente statura paulo minore, capite minus grandi, elytris longioribus, abdomine dorso evidentius crebriusque punctato, colore corporis magis rufo, elytrorum nigro-piceo, constanter differre videtur.

## Phloeocharis Mannerh.

*Mannerh. Brachelytr. p. 50.*

*Palpi maxillares subulati.*

Corpus pusillum, elongatum, subdepressum, alatum. Caput thorace minus et angustius, basi leviter constrictum, porrectum, oculis semiglobosis, prominulis. Labrum transversum, medio leviter emarginatum, apice membraneo-limbato. Mandibulae haud porrectae, ante medium unidentatae, margine interiore ad dentem usque limbo membraneo ciliato fimbriato. Maxillae (t. V. f. 5. b.) malis coriaceis, interiore apice introrsum oblique truncato barbata, exteriore hanc paululum superante apice barbata. Palpi maxillares breviusculi, articulo primo minuto, secundo leviter elongato, tertio incrassato, quarto tenui, subulato. Labium (t. V. f. 5. a.) mento brevi, transverso, antorsum angustato, apice truncato, ligula membranea, brevi, biloba, paraglossis liberis, ligulam vix superantibus, membraneis, interne ciliatis. Palpi labiales articulo primo cylindrico, secundo hoc multo brevior, tertio primo longitudine aequali, at tenuiore, apice subacuminato. Antennae capite thoraceque breviores, articulo primo cylindrico secundoque ovato crassiusculis, tertio tenuiore, obconico, 4 — 10 subglobosis, sensim paulo maioribus et crassioribus, ultimo ovato. Thorax sat am-



plus, transversus. Scutellum triangulare. Elytra apice truncata, ad angulum anteriorem emarginata. Abdomen marginatum, apice leviter angustatum. Pedes breviusculi, intermedii basi approximati, tibiis pubescentibus, tarsis breviusculis, articulis 4 primis brevibus, quinto his coniunctis longitudine aequali, anticis leviter dilatatis.

#### O b s e r v a t i o n e s.

1. Corpus totum punctulatum, minus subtiliter pubescens.
2. Sexus differentia latet.
3. Victus sub truncorum cortice sicco.

\* 1. *Ph. subtilissima*: Griseo-pubescens, fusco-nigra, elytris abdominisque segmentorum marginibus rufo-brunneis, ore, antennis pedibusque testaceis. — Long.  $\frac{3}{4}$  lin.

Mannerh. *Brachelytr.* 50. 1. — *Er. Col. March.* 1. 612. 1.

Habitat in Europa sub cortice Pinus.

Fusco-nigra, parum nitida, pube minus subtili depressa fulvescenti-grisea vestita. Antennae testaceae. Caput minutum, subtilissime punctulatum, ore testaceo. Thorax coleopterorum latitudine, latitudine plus dimidio brevior, lateribus leviter rotundatis, antrorsum subangustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, levissime convexus, subtilissime punctulatus, dorso obsolete longitudinaliter bi-impressus. Elytra thorace sesqui longiora, densius punctata, rufo-brunnea. Abdomen subtiliter punctulatum, segmentorum singulorum margine rufo-piceo. Pedes rufo-testacei.

---

## T r i b u s d e c i m a .

### O m a l i n i .

*Stigmata prothoracica occulta.*

*Coxae posticae transversae, anticae conicae, exsertae.*

*Trochanteres postici fulcrantes.*

*Ocelli duo.*

---

Labrum transversum, corneum,  
apice membrana medio sinuata vel biloba auctum Lestueae, Olophro, Deliphro, Omalio;  
utrinque lacinia parva acuminata membranacea ciliata instructum Lathrimaeo.

Mandibulae breves, falcatae,  
dentatae Anthophago, Lestueae;  
muticae reliquis.

Maxillae malis membraneis, inferiore brevior, apice unco corneo armata, superiore apice barbata,  
lineari Anthophago, Acidotae, Omalio, Anthobio;  
externe rotundato-dilatata Arpedio, Lathrimaeo, Deliphro;  
utraque lineari, elongata, cornea Lestueae, Olophro.

Palpi maxillares filiformes, articulo primo minuto.

Mentum corneum, transversum, antrorsum angustatum, apice truncatum.

Ligula membranacea, lata, biloba, paraglossis nullis.

Palpi labiales breviores, filiformes.

Antennae sub frontis margine laterali insertae, 11-articulatae, rectae, filiformes.

Caput porrectum, thoraci immissum, fronte ocellis duobus instructa.

Thorax lateribus marginatus, basi coleopterorum basi approximatus.

Scutellum distinctum.

Elytra pectore magis minusve longiora, nonnunquam abdominis maximam partem obtegentia, lateribus immarginata, angulo apicali posteriore fortiter rotundato.

Abdomen depressum, subtiliter marginatum, apice rotundatum, segmentis 6 distinctis.

Pedes coxis anticis conicis, exsertis, intermediis approximatis vel nonnunquam tenui spatio distantibus, posticis transversis, trochanteribus posticis fulcrantibus, tibiis aut inermibus aut spinulosis, spinis terminalibus subtilibus, tarsis 5-articulatis, posterioribus

articulis 4 primis sensim decrescentibus, primo magis minusve elongato Anthophago, Lestuae, Arpedio, Acidotae;

articulis 2 primis aequalibus, sequentibus duobus paulo longioribus Olophro, Lathrimaeo, Deliphro;

articulis 4 primis aequalibus, brevibus, coniunctis articulo ultimo longitudine aequalibus Omalio, Anthobio;

unguiculis simplicibus plerisque;

lacinia membranacea instructis Anthophago \*.

#### Observationes.

1. Tribus valde distincta, ocellis duobus frontalibus, maxillarum mala inferiore apice unco corneo terminata, elytris pectore longioribus, apice exteriori rotundatis insignis, e generibus Anthophago et Omalio Grav. composita, exclusis speciebus nonnullis, quae aut inter Oxytelinos Coprophilinos aut inter Protei-ninos genera peculiaria constituunt.

2. Abdominis segmentum sextum in mare truncatum, vel inferne emarginatum, septimum prominulum; femina stylo bifido ex ano saepius prominulo instructa.

### Anthophagus Grav.

Grav. Micr. p. 120. — Er. Col. March. I. p. 614.

*Mandibulae ante apicem bidentatae.*

*Maxillae malis membranaceis.*

*Palpi maxillares articulo ultimo penultimo aequali.*

*Tibiae muticae.*

*Tarsi postici articulo primo elongato.*

Corpus oblongum, subdepressum, alatum. Caput exsertum, thoracis latitudine vel eodem latius, rhombeum, basi constrictum, collo quasi cylindrico thoraci immissum, oculis medioeribus, rotundatis, prominulis. Ocelli pone oculorum marginem superiorem siti. Labium transversum, apice subsinuatum, Mandibulae apice acuto introrsum falcato, ante apicem acute bidentatae. Maxillae (t. V. f. 25. b.) malis membranaceis, exteriori apice, interiori intus barbatis, hac unco corneo terminata. Palpi maxillares modice elongati, articulo primo minuto, reliquis tribus inter se longitudine aequalibus, tertio leviter obconico, quarto apice subacuminato. Labium (t. V. f. 25. a.) mento brevi, transversum, apice vix sinuato, ligula membranacea, biloba, lobis interne sinuatis. Palpi labiales articulo primo breviusculo, secundo longiore, crassiusculo, tertio iterum minore subacuminato. Antennae filiformes, satis elongatae gracilesque. Thorax coleopteris multo angustior, subcordatus seu basin versus angustatus. Scutellum triangulare. Elytra pectus perparum excedentia. Abdomen late marginatum. Pedes elongati, graciles, tarsis articulo primo leviter elongato, quarto subtus pilorum fasciculo instructo, quinto plerumque laciniis duabus membraneis, unguiculo ipso longitudine aequalibus instructo.

#### Observationes.

1. Corpus iam glabrum iam subtilissime subtiliterve pubescens. Caput, thorax et elytra punctulata. Frons ante ocellos lineolis duabus obliquis insculpta.



2. Habitu, thorace subcordato pedibusque elongatis, gracilibus ad Caraborum nonnullorum genera v. c. ad Dromium accedunt Anthophagus et Lestea.

3. Mas differt abdominis segmento septimo infra conspicuo. Specierum nonnullarum (*A. armiger*, *alpinus*) mares capite dilatato, antice bispinoso mandibulisque elongatis crassatisque excellunt.

4. Victus in herbis, fruticibus et floribus.

\* Unguiculi basi membrana lineari, libera, unguiculo ipso fere aequali, instructi.

X

\* 1. *A. armiger*: Rufo-testaceus, pectore abdomineque nigro-piceis, elytris testaceis, thorace transversim quadrato, crebrius punctato, basi foveolato, fronte maris apice bispinosa. — Long.  $2\frac{1}{2}$ —3 lin.

Grav. Micr. 122. 3. Mon. 221. 5.

*Staph. bicornis* Block Ins. vall. Plauens. 118. 10. f. 10.

Habitat in Germaniae montanis. Suecicum etiam specimen dedit Dom. Schüppel.

*A.* testaceo paulo maior et latior. Antennae capite thoraceque multo longiores, testaceae. Caput parce subtiliterque punctulatum, inter oculos striolis duabus obliquis impressum, rufo-testaceum, basi nonnunquam piceum, feminae vix thoracis latitudine, maris thorace latius, medio late at parum profunde impressum, apice ad antennarum insertionem spina valida porrecta utrinque armatum, mandibulis prominentibus, incrassatis, incurvis, decussatis. Thorax coleopteris duplo fere angustior, latitudine dimidio brevior, lateribus anterieus leviter rotundatis, basin versus parum angustatus, angulis posterioribus rectis, parum convexus, margines versus distinctius crebriusque, medio parcius obsoletiusque punctatus, basi supra scutellum foveola transversa impressus, rufo-testaceus, nitidus, medio interdum piceus. Scutellum laeve, rufo-testaceum, rarius piceum. Elytra thorace plus duplo longiora, sat crebre punctata, parce subtiliter pubescentia, luteo-testacea, nitida. Abdomen parce subtiliter punctulatum, tenuissime griseo-pubescent, nigrum, nitidum, margine saepius irregulariter obscure testaceo. Pectus nigro-piceum. Pedes rufo-testacei.

Variat elytris brevioribus, thorace vix duplo longioribus.

\* 2. *A. alpinus*: Nigro-piceus, antennarum basi, elytris pedibusque testaceis, thorace subquadrato, vage punctato. — Long.  $1\frac{1}{2}$ —2 lin.

Mas capite magno, apice bispinoso, mandibulis magnis, incurvis.

*Anthophag. mandibularis* Gyll. Ins. Suec. IV. 460. 2—3. — Sahlb. Ins. Fenn. I. 275. 4.

*Anthophag. maxillosus* Zetterst. Faun. Lappon. I. 45. 3. Ins. Lappon. 48. 4.

Fem. capite thorace angustiore, mutico.

*Staph. alpinus* Fab. Ent. Syst. I. II. 526. 33. Syst. El. II. 598. 43. — Payk. Faun. Suec. III. 387. 27. — Oliv. Ent. III. 42. 32. 45. t. 6. f. 55.

*Lestea alpina* Latr. Hist. nat. Crust. et Ins. IX. 368. 1. Gen. Crust. et Ins. I. 297. 2. — Lepell. et Serv. Encycl. meth. Ins. X. 216. 1. — Runde Brachelytr. Hal. 24. 5.

*Anthophag. alpinus* Grav. Micr. 188. 2. Mon. 220. 1. — Zetterst. Faun. Lapp. I. 46. 4. Ins. Lappon. 48. 5.

*Anthophag. alpinus* var. *a* Gyll. Ins. Suec. II. 194. 2.

- Anthophag. Lapponticus* Sahlb. *Ins. Fenn. I.* 275. 3. — *Zetterst. Ins. Lappon.* 47. 2.  
*Lesteua Lapponica* Mannerh. *Brachelytr.* 56. 5.  
*Lesteua flavipennis* Boisd. et Lacord. *Faun. Ent. Paris. I.* 480. 2.

Habitat in Suecia boreali; in Alpibus Tyrolensibus, Dom. Prof. Ahrens, Helveticis Dom. Heer, in montibus Silesiae, Dom. Märkel.

A. Caraboide dimidio minor. Antennae capite thoraceque longiores, nigrae, articulis 2, 3 vel 4 primis testaceis. Caput parce subtiliter obsoleteque punctatum, inter oculos lineolis duabus obliquis impressum, nigropiceum, subnitidum, ore testaceo, feminae thoracis latitudine, maris thorace multo latius, inaequale, apice ante antennarum insertionem spina porrecta utrinque armatum, mandibulis valde prominentibus, incrassatis, incurvis. Thorax coleopteris dimidio angustior, latitudine vix brevior, lateribus anterieus modice rotundatis, basin versus leviter angustatus, angulis posterioribus rectis, leviter convexus, vage sat distincte punctatus, dorso longitudinaliter obsolete bi-impresso, spatio interiecto subelevato, laevi, nigro- vel fusco-piceus, subnitidus, margines versus rufescens. Scutellum laeve, piceum. Elytra thorace duplo longiora crebre fortiterque punctata, pallide testacea. Pectus et abdomen nigra, nitida. Pedes rufo-testacei.

Obs. Femina mare multo frequentior: Dom. Märkel singulum inter quadragenas vel quinquagenas feminas marem observavit.

- \* 3. *A. Omalinus*: *Nigro-piceus, nitidus, capite thoraceque rufo-piceis, antennis, pedibus elytrisque testaceis, thorace transversim subquadrato, parce obsoleteque punctato.* — *Long.*  $1\frac{1}{2}$  lin.

*Zetterst. Faun. Lappon. I.* 46. 5. *Ins. Lappon.* 48. 6.

*Anthophag. alpinus* var. *b* Gyll. *Ins. Suec. IV.* 194. 2.

*Anthophag. alpinus* Sahlb. *Ins. Fenn. I.* 274. 2.

*Lesteua alpina* Mannerh. *Brachelytr.* 56. 6.

Habitat in Lapponia, Dom. Zetterstedt, in Dalecarlia, Dom. Schönherr, — „in Pini sylvestris frondibus” Gyll. „in foliis Betulae albae” Zetterst.

A. alpino paulo minor et paulo magis dilatatus. Antennae capite thoraceque paulo longiores crassiusculae, totae laete testaceae. Caput in utroque sexu thorace angustius, vix punctatum, striolis duabus obliquis inter oculos sat profundis impressum, piceo- vel rufo-testaceum, subnitidum. Thorax coleopterorum basi dimidio angustior, latitudine paulo brevior, lateribus ante medium leviter rotundatis, basin versus vix angustatus, angulis posterioribus rectis, leviter convexus, aequalis, punctis obsolete, raris, ad marginem basalem crebrioribus impressus, rufo-testaceus, subnitidus. Scutellum laeve, piceo-testaceum. Elytra thorace plus duplo longiora, apicem versus sensim dilatata, crebrius punctata, dilute testacea. Pectus et abdomen nigra, nitida. Pedes testacei.

Mas nonnisi antennis longioribus abdominisque segmento sexto infero emarginato distinguitur.

- \* 4. *A. Austriacus*: *Niger, ore, antennis, pedibus elytrisque testaceis, thorace transversim quadrato, subtiliter vage punctato.* — *Long.*  $2\frac{1}{2}$  lin.

Habitat in Austria, Dom. Schüppel, in Carinthia, Dom. Schmidt.

Statura fere A. testacei, niger, capite thoraceque parum nitidis, sub-

aeneis. Antennae capite thoraceque longiores, crassiusculae, obscurae testaceae, articulis 4 — 6 vel 4 — 8 fusciscentibus. Os testaceum. Caput in utroque sexu thorace angustius, vage punctatum, inter oculos striolis duabus obliquis impressum, antice ad antennarum basin utrinque foveolatum. Thorax coleopteris paulo angustior, latitudine tertia parte brevior, lateribus antemedium leviter rotundatis, basin versus parum angustatus, angulis posterioribus obtusiusculis, parum convexus, vage subtiliter punctatus, dorso longitudinaliter obsolete bi-impressus, spatio interiecto impunctato, marginibus obscurae testaceis. Scutellum laeve, nigro-piceum. Elytra thorace duplo longiora, sat crebre fortiterque punctata, testacea. Abdomen confertim subtiliter subrugoso-punctatum, subtiliter griseo-pubescent, nigrum, nitidum. Pectus nigrum. Pedes testacei, femoribus omnibus maris incrassatis.

\* 5. *A. Caraboides*: *Rufo-testaceus, abdomine ante apicem capiteque piceis, thorace subcordato, subtiliter punctato.* — Long. 2 lin.

Grav. Micr. 120. 1. Mon. 221. 2. — Zetterst. Faun. Lappon. I. 45. 2. Ins. Lappon. 47. 3. — Er. Col. March. I. 614. 1.

*Staph. Caraboides* Linn. Faun. Suec. n. 854 Syst. Nat. I. II. 685. 20. — Fabr. Syst. Ent. 267. 18. Spec. Ins. I. 337. 23. Mant. Ins. I. 222. 33. Ent. Syst. I. II. 530. 53. Syst. El. II. 601. 65. — Schrank Enum. Ins. Austr. 234. 443. — Goeze Ent. Beitr. I. 720. 20. — De Vill. Ent. I. 417. 20. — Cederh. Faun. Ingr. 119. 366. — Walk. Faun. Paris. I. 271. 10. — Payk. Mon. Staph. 19. 11. Faun. Suec. III. 386. 25. — Oliv. Ent. III. 42. 22. 26. t. 2. f. 17. — Panz. Ent. Germ. 358. 42. — Marsh. Ent. Britt. 521. 67.

*Lestea Caraboides* Mannerh. Brachelytr. 55. 3. — Runde Brachelytr. Hal. 23. 3. Anthophag. Caraboides var. a. b Gyll. Ins. Suec. II. 192. 1. — Sahlb. Ins. Fenn. I. 274. 1.

*Staph. fulvus* Degeer Ins. IV. 25. 12.

*Lestea testacea* Boisd. et Lacord. Faun. Ent. Paris. I. 482. 5.

Habitat in Europa.

Antennae capite thoraceque multo longiores, testaceae. Caput in utroque sexu thoracis latitudine, parcius punctulatum, inter oculos striolis duabus obliquis sat profundis impressum, rufo-testaceum, plerumque magis minusve picescens, ore semper rufo-testaceo. Thorax coleopteris duplo angustior, latitudine haud brevior, lateribus anterieus rotundatus, basin versus angustatus, angulis posterioribus subrectis, parum convexus, parcius subtiliter punctatus, dorso nonnunquam obsolete longitudinaliter bi-impressus, rufo-testaceus, parum nitidus. Scutellum laeve, rufo-testaceum, nitidum. Elytra thorace duplo longiora, sat crebre fortiterque punctata, dilutius testacea. Abdomen rufo-testaceum, postice macula magna picea, apice summo iterum testaceo. Pectus testaceum. Pedes flavi.

\* 6. *A. abbreviatus*: *Testaceus, abdomine apice, capite pectoreque nigropiceis, elytris fascia fusciscente, thorace subcordato.* — Long.  $1\frac{2}{3}$  — 2 lin.

*Carabus abbreviatus* Fab. It. Norvag. p. 263. Spec. Ins. 313. 85. Mant. Ins. I. 205. 120. Ent. Syst. I. 1. 167. 188. Syst. El. I. 209. 215. — Panz. Ent. Germ. 367. 114 — 115. Faun. Germ. 36. 2. — Illig. Käf. Pr. 205. 90.

*Lestea angusticollis* Mannerh. Brachelytr. 56. 4.

Habitat in Europa.

Praecedente paulo gracilior. Antennae capite thoraceque multo lon-



giores, testaceae. Palpi pallidi. Caput in utroque sexu thoracis latitudine, parcius punctatum, inter oculos striolis duabus obliquis sat profundis impressum, nigro-piceum, nitidum, ore testaceo. Thorax coleopteris duplo angustior, latitudine summa haud brevior, lateribus anterieus rotundatus, basin versus angustatus, angulis posterioribus subrectis, parum convexus, parcius subtiliter punctatus, nonnunquam longitudinaliter obsolete bi-impressus, rufo-testaceus, subnitidus, disco nonnunquam fusciscente. Scutellum laevissimum, piceo-testaceum, nitidum. Elytra thorace plus duplo longiora, crebre profundius punctata, parce subtilissime griseo-pubescentia, nitida, lutea, fascia media sinuata dilute fusca, nonnunquam magis minusve obsolescente, sutura nonnunquam infuscata. Abdomen parcius subtiliter punctatum, nitidum, iam nigrum basi testaceum, iam testaceum apice late nigro-piceum. Pectus nigro-piceum. Pedes pallide flavi.

Obs. Proximus praecedenti et forte mera eius varietas, statura autem graciliore, thorace basin versus paulo magis angustato, elytris fascia fusciscente, pectore piceo etc. distinguendus videtur.

\* 7. *A. scutellaris*: Testaceus, nitidus, abdomine ante apicem, fronte, scutello pectoreque nigro-piceis, thorace crebre punctato, elytris luteis, circa scutellum infuscatis. — Long.  $2\frac{1}{4}$  lin.

Habitat Genevae. Specimen a Dom. Chevrier sibi communicatum dedit Dom. Prof. Germar.

Praecedentibus paulo maior. Antennae capite thoraceque longiores, validiores, testaceae. Palpi dilute testacei. Caput thoracis latitudine, fronte posterius crebre fortius punctata, striolis duabus obliquis impressa, anterieus parce punctata, leviter impressa et utrinque foveolata, nitidum, rufo-testaceum, supra posterius nigro piceum. Thorax coleopteris dimidio angustior, latitudine vix brevior, lateribus ante medium rotundatus, basin versus angustatus, angulis posterioribus rectis, parum convexus, sat crebre profundius punctatus, linea media longitudinali laevi, rufescenti-testaceus, nitidus, parce subtiliter griseo-pubescent. Scutellum laevissimum, nigro-piceum, nitidum. Elytra thorace duplo longiora, densius profunde punctata, subtiliter griseo-pubescentia, macula communi triangulari circa scutellum nigricante. Abdomen parcius subtilissime punctulatum, subtiliter pubescens, testaceum, nitidum, macula maiore rotundata ante apicem nigro-picea. Pectus nigro-piceum. Pedes flavi.

Obs. A praecedentibus duobus thorace pubescente, a sequente colore, elytris brevioribus, thorace lateribus minus rotundato distinctus.

\* 8. *A. testaceus*: Rufo-testaceus, nitidus, capite thoraceque rufis, hoc subquadrato, crebre punctato. — Long.  $2\frac{1}{4}$  lin.

Grav. Micr. 121. 2. Mon. 221. 3. — Er. Col. March. I. 615. 2.

*Lestena testacea* Mannerh. Brachelytr. 55. 2. — Runde Brachelytr. Hal. 23. 2.

*Lestena Caraboides* Boisd. et Lacord. Faun. Ent. Paris. I. 480. 3.

*Anthophag. Caraboides* var. c Gyll. Ins. Suec. II. 192. 1. — Sahlb. Ins. Fenn. I. 274. 1.

Habitat in Europa, in fruticibus.

A. Caraboide sesqui fere maior, rufo-testaceus, nitidus, capite thoraceque saturatoribus, testaceo-rufis. Antennae capite thoraceque multo lon-

giores. Caput in utroque sexu thorace paulo angustius, crebrius sat distincte punctatum, antice laevius, fronte planiuscula, inter oculos striolis duabus, vix obliquis, profundis impressum. Thorax coleopteris plus dimidio angustior, latitudine haud brevior, lateribus anterieus fortius rotundatus, basin versus leviter angustatus, angulis posterioribus rectis, leviter convexus, sat crebre fortius punctatus, postice linea media longitudinali abbreviata laevi, subelevata, subtiliter griseo-pubescentia. Scutellum laeve. Elytra plus duplo longiora, crebre fortiterque punctata, tenuiter griseo-pubescentia. Abdomen parcius subtiliter punctulatum, subtiliter griseo-pubescentia, ante apicem saepe fuscescens. Pedes dilutius testacei.

Variat interdum capite thoraceque fuscescentibus: forte *A. Caraboidis* var. d et e Gyll. Sahlb. (l. supra c.) huc referendae.

X

\* 9. *A. praeustus*: *Rufo-testaceus, nitidus, abdomine ante apicem elytrorumque macula magna apicali nigris, thorace transversim subquadrate, crebre punctato.* — Long.  $2\frac{1}{4}$  lin.

Müller Germ. Mag. IV. 228. 25.

*Lesteua praeusta* Runde Brachelytr. Hal. 23. 2.

*Lesteua bimaculata* Boisd. et Lacord. Faun. Ent. Paris. I. 481. 4.

Habitat in Germania, Gallia, Hungaria.

Statura et summa affinitas *A. testacei*, rufo-testaceus, nitidus. Antennae capite thoraceque multo longiores, crassiusculae. Caput in utroque sexu thorace angustius, fronte depressa, crebrius fortiusque punctata, striolis duabus obliquis profundis impressa, antice laeviore. Thorax coleopteris dimidio angustior, latitudine paulo brevior, lateribus anterieus fortius rotundatus, basin versus leviter angustatus, angulis posterioribus rectis, parum convexus, crebre fortius punctatus, linea media longitudinali subelevata, antice abbreviata, laevi, subtiliter griseo-pubescentia. Scutellum laeve. Elytra thorace plus duplo longiora, sat crebre fortiter punctata, longius griseo-pubescentia, macula magna rotundata apicali nigra, marginem haud attingente. Abdomen parce subtilissime punctulatum, parce pubescens, nitidum, supra infraque macula magna nigra ante apicem.

### \*\* Unguiculi simplices.

\* 10. *A. plagiatus*: *Niger, pubescens, crebre punctatus, fronte impressa, thorace cordato, basi foveolato, antennis pedibusque fusco-testaceis.* — Long. 2 —  $2\frac{1}{2}$  lin.

Var. a: *Elytris thorace plus duplo longioribus, plaga discoidali rufescente.*

Grav. Micr. 123. 5. 189. 3. Mon. 222. 8. — Gyll. Ins. Suec. II. 195. 3. — Zetterst. Faun. Lappon. I. 44. 1. Ins. Lappon. 47. 1. — Sahlb. Ins. Fenn. I. 276. 5. — Germ. Faun. Ins. Europ. XIII. 5.

*Staph. plagiatus* Fab. Ent. Syst. Suppl. 180. 32. Syst. El. II. 597. 42. — Payk. Faun. Suec. III. 387. 26.

*Lesteua plagiata* Mannerh. Brachelytr. 56. 7. — Boisd. et Lacord. Faun. Ent. Paris. I. 482. 6. — Runde Brachelytr. Hal. 24. 6.

Var. b: *Elytris thorace plus duplo longioribus, sutura sanguinea.*

*Lesteua suturalis* Boisd. et Lacord. Faun. Ent. Paris. I. 283. 7.

*Var. c: Elytris thorace plus duplo longioribus, concoloribus.*

*Staph. plagiatus var. Payk. Faun. Suec. III. 387. 26.*

*Anthophag. plagiatus var. b Gyll. Ins. Suec. II. 195. 3. — Sahlb. Ins. Fenn. I. 276. 5.*

*Anthophag. nigrita Müll. Germ. Mag. IV. 226. 24.*

*Var. d: Elytris thorace vix duplo longioribus, fusco-piceis, long. 2 lin.*

*Lestea longipes Mannerh. Brachelytr. 56. 9. (forte)*

*Var. e: Thorace canaliculato, elytris thorace vix duplo longioribus, nigris, concoloribus.*

*Lestea globulicollis Mannerh. Brachelytr. 56. 8.*

*Anthophag. globulicollis Zetterst. Ins. Lappon. 49. 8.*

*Var. f: Thorace canaliculato, elytris thorace sesqui longioribus, disco rufescentibus.*

Habitat in Europa.

A. testaceo plerumque maior et latior, niger, subnitidus, tenuiter griseo-pubescens. Antennae capite thoraceque multo longiores, piceo-rufae, basi interdum nigricantes, articulo ultimo elongato. Os piceo-rufum. Caput in utroque sexu thorace angustius, crebrius punctatum, inaequale, fronte inter oculos impressa, striolisque duabus obliquis sulcata, medio obsoletius distinctiusve canaliculata. Thorax coleopteris dimidio angustior, latitudine paulo brevior, lateribus fortiter rotundatus, basi coarctatus, angulis posterioribus rectis, subprominulis, leviter convexus, crebre fortius punctatus, basi supra scutellum impressione transversa, et utrinque in ipso angulo baseos foveola minuta rotundata notatus, dorso interdum medio longitudinaliter obsolete impressus, in aliis distinctius obsoletiusve canaliculatus, in aliis linea media longitudinali laeviore, posterius subelevata. Scutellum laeve. Elytra sat crebre fortiterque punctata, nigra, iam plaga media longitudinali, iam sutura sanguineis, iam immaculata. Abdomen subtiliter subtus densius punctulatum. Pedes piceo-rufi.

Obs. 1. Maiore individuorum copia examinata varietates citatae specie mihi non satis distinctae videntur. Varietates a et c elytris thorace plus duplo longioribus inter se conveniunt et praeter elytrorum colorem vix differunt. Antennarum basis, palpi et pedes in utraque iam nigro-, iam rufo-picei, iam rufi. Var. b, genuinis iam paulo maior, iam dimidio minor, elytrorum longitudine cum iis congruens, at thorace lateribus paulo minus rotundato, forte specie distincta, at characteres haud sufficere videntur. Hanc ex Austria misit Dahl nom. Anthophagi variegati et variantis. — Var. d, nom. Anthophagi Kunzei a Dom. Heer ex Helvetiae montibus missa, elytris brevioribus, genuinis paulo minor, thorace parum convexo, aequali a var. e distincta, quae colore et statura cum var. e congruit, et thorace paululum convexiore, distinctius canaliculato et elytris paulo brevioribus discedit. Specimina eius e Dalecarlia a Dom. Schönherr sibi communicata dedit Dom. Schüppel. — Var. f tandem, cuius specimen Tyrolense dedit Dom. Walzl, colore cum var. a, thoracis structura cum var. e convenit, ab omnibus autem coleopteris multo brevioribus et thorace vix latioribus longius discedit.

2. *Staph. nigrophthalmus* Block Ins. Vall. Plauens. 117. 9. f. 9. huius speciei specimen immaturum colore testaceo exhibere videtur.

\* 11. *A. caesus: Testaceus, fronte, thorace elytrisq. brunneis, abdomine ante apicem brunneo, thorace oblongo, profunde canaliculato. — Long. 3½ lin.*

Habitat in America septentrionali, Knoch.



Antennae elongatae, dimidio corpore longiores, articulo ultimo paululum incrassato, testaceae. Caput thorace plus dimidio angustius, vage punctatum, inaequale, inter oculos impressum, saturate rufo-testaceum, supra posterius nigro-brunneum. Thorax coleopteris paulo angustior, latitudine paulo longior, lateribus fortiter rotundatus, basin apicemque versus aequaliter angustatus, angulis posterioribus minutis, acutiusculis, prominulis, convexus, canalicula profunda, basi foveola terminata, exaratus, sat crebre fortiterque punctatus, tenuiter griseo-pubescent, subtile saturate testaceus, supra nigro-brunneus, nitidus, margine antico posticoque rufescente. Scutellum piceum, impressum. Elytra thorace paulo longiora, crebre sat fortiter punctata, brunnea, apicem versus sensim nigro-picea. Abdomen testaceum, ante apicem nigricans. Pectus testaceum. Pedes elongati, testacei.

Species mihi invisae.

12. *A. brunneus*: *Rufo-brunneus, pedibus abdomineque pallidioribus, thorace linea foveolaque dorsali impresso. — Long. infra 2½ lin.*

*Say Act. Acad. Nat. Scienc. Philadelph. III. p. 158.*

Habitat in fluminis Missouri ripis.

Rufo-brunneus, punctatus, breviter pubescens. Caput inaequale, inter oculos et inter antennis impressum. Antennae, palpi et pedes testacei. Mandibulae apice piceae. Thorax dense punctatus, subrotundatus, angulis posterioribus rectis, linea dorsali basi foveola terminata canaliculatus. Elytra dense punctata, angulo posteriore laterali rotundato, suturali acuto. Abdomen pallide rufo-brunneum, dorso macula fusca ante apicem, segmentis singulis undique fusco-marginatis. (Say.)

13. *A. verticalis*: *Rufo-brunneus, elytris segmentorumque abdominis marginibus testaceis. — Long. infra 2 lin.*

*Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. p. 463.*

Habitat in Pensylvania.

Castaneus, punctatus, leviter pubescens. Caput thorace obscurius, sparsim punctatum, verticis disco impresso, inter antennis inaequale, antennis et ore pallidioribus. Thorax convexus, utrinque rotundatus, basi contractus, linea impressa basi puncto dilatato terminata evidenter canaliculatus. Elytra rufo-testacea, punctata, apice obscura. Abdomen linea dorsali impressa, segmentis singulis in utroque latere macula apicali triangulari. Pedes femoribus testaceis, tibiis tarsisque dilutioribus.

Obs. Affinis *A. brunneo*, at abdominis segmenta haud fusco-marginata, singula autem macula triangulari, ad marginem posteriorem fortius dilatata et marginem lateralem prope attingente signata. (Say.)

## Lesteua Latr.

Latr. Précis d. caract. d. gen. d. Ins. p. 75. — Er. Col. March. I. p. 616.

*Mandibulae medio unidentatae.*

*Maxillae malis elongatis, corneis.*

*Palpi maxillares articulo ultimo penultimo triplo longiore.*

*Tibiae muticae.*

*Tarsi postici articulo primo elongato.*

Corpus oblongum, subdepressum, alatum apterumve. Caput exsertum, thorace vix angustius, rhombeum, basi constrictum, oculis mediocribus, rotundatis, sat prominulis. Ocelli super oculorum marginem superiorem siti. Labrum (t. V. f. 26. c.) breve, apice leviter sinuatum, membrana medio subsinuata auctum. Mandibulae leviter falcatae, acutae, medio denticulo armatae. Maxillae (t. V. f. 26. b.) malis corneis, elongatis, linearibus, exteriore barbatula, interiore unco terminata, interne barbata. Palpi maxillares articulo primo tertioque brevi, secundo quartoque elongatis, quarto tertio triplo longiore. Labium (t. V. f. 26. a.) mento transversum, apicem versus angustatum, apice truncato, ligula membranacea, biloba, lobis rotundatis. Palpi labiales articulo primo tertioque elongatis, secundo brevi. Antennae filiformes, graciles, articulo primo cylindrico, sequentibus subobconicis, ultimo ovato, apice acuminato. Thorax coleopteris angustior, lateribus rotundatus, basin versus angustatus. Scutellum triangulare. Elytra pectus parum superantia, angulo apicali exteriore rotundato. Abdomen late marginatum, apice subacuminatum. Pedes elongati, graciles, tibiis pubescentibus, posticis articulo primo elongato.

### O b s e r v a t i o n e s.

1. Corpus subtiliter pubescens. Caput, thorax et elytra dense punctata. Frons utrinque longitudinaliter leviter impressa, impressione inter oculos foveolata, antice clypei spatio impressione semicirculari distincto.

2. Mas abdominis segmento sexto subtruncato, segmento igitur septimo conspicuo a femina distinguitur.

3. Habitus omnino generis praecedentis, quocum confunditur. Oris structura omnino distinctiva. Palporum maxillarium differentia facillime diagnosi potest. Praeterea frons in *Lesteua* utrinque longitudinaliter impressa, in *Anthophago* medio lineolis duabus obliquis insculpta differentiam quandam habitualement suppediat.

4. Species nonnisi Europaeae cognitae. Victus ad rivulos in herbis indigenas praecipue in muscis rivulorum affluvie humidis observavi.

5. Latreillianum servavi nomen huic generi, quod speciei eius vulgari conditum fuit. Nomen Gravenhorstianum praecedenti generi proprium maneat.

\* 1. *L. bicolor*: Alata, nigra, pubescens, crebre punctata, antennis pedibusque fusco-testaceis, thoracis angulis posterioribus rectis, elytris thorace plus duplo longioribus, fuscis. — Long. 2 lin.

Er. Col. March. I. 616. 1.

*Staph. bicolor* Fab. Ent. Syst. I. II. 529. 47. Fab. Syst. El. II. 600. 60. — Walk. Faun. Paris. I. 270. 8.

*Lesteua punctulata* Latr. Hist. nat. Crust. et Ins. IX. 369. 2. Gen. Crust. et Ins. IX. 297. 1. t. 9. f. 1.

*Staph. obscurus* Payk. Faun. Suec. III. 388. 28.

*Anthophag. obscurus* Grav. Micr. 122. 4. Mon. 221. 7. — Gyll. Ins. Suec. II. 195. 4. — Zetterst. Faun. Lappon. I. 47. 6. Ins. Lappon. 49. 7. — Sahlb. Ins. Fenn. I. 176. 6.

- Lestena obscura* Mannerh. *Brachelytr.* 57. 10. — *Boisd. et Lacord. Faun. Ent. Paris.* 1. 483. 8. — *Runde Brachelytr. Hal.* 24. 7.  
*Anthophag. intermedius* Grav. *Mon.* 221. 4.  
*Carabus dimidiatus* Panz. *Faun. Germ.* 36. 3. *Ent. Germ.* 367. 114 — 115.  
*Carabus Staphylinoides* Marsh. *Ent. Britt.* 464. 87.  
*Staph. multipunctatus* Block *Ins. vall. Plauens.* 117. 8. f. 8. (*forte*)  
*Staph. macroelytron* Fourcr. *Ent. Par.* 1. 164. 4.  
*Le Staph. noir à long étuis* Geoffr. *Hist. d. Ins.* 1. 362. 4.

Habitat in Europa.

Nigra, nitida, tenuiter griseo-pubescent. Antennae capite thorace paululum longiores, piceo-testaceae, articulis duobus primis saepe fuscescentibus. Palpi testacei. Caput thorace paulo angustius, sat crebre punctatum, fronte utrinque leviter longitudinaliter impressa, impressione utraque inter oculos foveolata. Thorax coleopteris dimidio angustior, latitudine summa paulo brevior, lateribus anterie fortiter rotundatus, postice coarctatus, angulis posterioribus rectis, leviter convexus, confertim minus subtiliter punctatus, basi supra scutellum foveola transversa impressus, interdum etiam versus apicem medio longitudinaliter obsolete impressus. Scutellum laeve. Elytra thorace plus duplo longiora, dense fortius punctata, fusca, iam nigrofusca, interdum subaenea, iam fusco-testacea. Abdomen confertim subtiliter punctatum. Pedes piceo-testacei.

- \* 2. *L. pubescens*: *Alata, nigra, densius pubescens, subtiliter punctata, ore, antennis pedibusque rufis, thoracis angulis posterioribus acutiusculis, elytris thorace duplo longioribus, fuscis.* — *Long.* 2 lin.

*Mannerh. Brachelytr.* 57. 12. — *Er. Col. March.* 1. 617. 2.

Habitat in Germania.

Statura omnino et summa similitudo *L. bicoloris*. Nigra vel fusco-nigra, nitidula, pube tenui cinerea densius vestita. Antennae capite thorace multo longiores, graciles, rufae, basi dilutiores. Palpi rufi. Caput thorace angustius, sat crebre subtilius punctatum, fronte utrinque profundius longitudinaliter impressa, ore piceo-rufo. Thorax coleopteris dimidio angustior, latitudine vix brevior, lateribus ante medium fortiter rotundatus, basin versus angustatus, angulis posterioribus acutiusculis, leviter prominulis, parum convexus, sat crebre subtiliter punctatus, basi supra scutellum obsolete punctatus. Elytra thorace duplo longiora, sat crebre subtilius punctulata, fusca vel fusco-nigra. Abdomen confertim subtilissime punctulatum. Pedes rufo-testacei.

Obs. 1. Thorax nonnunquam medio dorso utrinque foveolatus: specimen tale descripsit Com. Mannerheim.

2. Differt a *L. bicolor* pubescentia densiore, punctura subtiliore, antennis longis, gracilibus, fronte utrinque profundius longitudinaliter impressa, thoracis angulis posterioribus acutiusculis etc.

- \* 3. *L. maura*: *Alata, nigra, pubescens, antennis pedibusque piceis, thoracis angulis posterioribus obtusis, elytris thorace plus duplo longioribus.* — *Long.* 1 $\frac{3}{4}$  lin.

Habitat in Sardinia, Dom. Prof. Gené.

*L. bicolor* paulo minor, nigra, nitidula, densius griseo-pubescent.



Antennae capite thoraceque vix longiores, piceae, articulis duobus primis nigricantibus. Palpi picei. Caput thorace paulo angustius, fronte utrinque longitudinaliter, antice transversim impressa. Thorax coleopteris dimidio angustior, latitudine summa haud brevior, lateribus anterieus fortiter rotundatus, basin versus angustatus, angulis posterioribus obtusis, leviter convexus, confertim subtilius punctatus, dorso medio nonnunquam obsolete bifoveolatus. Scutellum laeve. Elytra thorace plus duplo longiora, nigro-fusca. Abdomen crebrius subtilissime punctulatum. Pedes iam nigro-, iam rufo-picei.

Obs. Proxima *L. bicolori*, at statura paululum minore, antennis paulo brevioribus, frontis impressionibus longitudinalibus profundioribus, inter oculos haud foveolatis et thoracis angulis posterioribus obtusioribus distincta.

- \* 4. *L. Sicala: Picea, subnitida, tenuiter griseo-pubescent, thorace brevior, lateribus foveolato, dorso obsolete bi-impresso, elytris thorace plus duplo longioribus. — Long. 2 lin.*

Habitat in Sicilia, Dom. Grohmann.

*L. bicolor* paulo brevior et latior, saturate rufo-picea, nitidula, tenuiter griseo-pubescent, ore, antennis pedibusque rufis. Caput thorace dimidio angustius, dense profunde punctatum, fronte planiuscula, inter oculos utrinque obsolete impressa. Thorax coleopteris dimidio angustior, latitudine summa tertia parte brevior, lateribus anterieus fortiter rotundatus, basi angustatus, angulis posterioribus obtusis, perparum convexus, crebre profunde punctatus, dorso obsolete longitudinaliter bi-impressus, lateribus utrinque medio foveola minuta rotundata notatus. Scutellum parce punctatum. Elytra thorace plus duplo longiora, subdepressa, fortius profunde dense punctata. Abdomen crebrius subtiliter punctulatum, dorso nigricans.

- \* 5. *L. punctata: Aptera, picea, pubescens, crebre fortius punctata, antennis pedibusque rufis, elytris thorace vix longioribus, fortiter punctatis. — Long. vix 2 lin.*

*Er. Col. March. I. 618. 3.*

Habitat in Germania.

*L. bicolor* paulo minor, nigro-, rarius brunneo- vel rufo-picea, subnitida, tenuiter pubescens. Antennae capite thoraceque longiores, crassiusculae, rufae. Caput thorace paulo angustius, apicem versus attenuatum, sat crebre profunde punctatum, fronte inter oculos utrinque oblique impressa, antice inter antennas utrinque foveolata, ore rufo. Thorax coleopteris angustior, latitudine summa vix brevior, lateribus ante medium fortiter rotundatis, angulis posterioribus subrectis, postice fortiter angustatus, leviter convexus, sat crebre fortius punctatus, lateribus medio foveolatus. Scutellum punctulatum. Elytra thorace vix duplo longiora, sat crebre fortiter punctata. Abdomen subtilissime punctulatum. Pedes rufi.

Species mihi invisae, forte huius generis.

6. *L. longula: Nigro-fusca, densius pubescens, subtilius creberrime punctulata, antennis, elytris pedibusque testaceis, thorace latiore, cordato, elytris longioribus, fronte bifoveolata.*

*Mannerh. Brachelytr. 57. 11.*

Habitat in Fennia meridionali.

*L. bicolori* similis, subtilius et crebrius punctulata, thorace latiore, elytrisque thorace triplo longioribus, dum in *L. bicolori* elytra thorace vix duplo longiora (?), quare ut species diversa consideranda. (Mannerh.)

### **Arpedium Er.**

*Er. Col. March. I. p. 618.*

*Mandibulae muticae.*

*Maxillae malis membranaceis.*

*Palpi maxillares articulo ultimo penultimo longitudine aequali.*

*Tibiae muticae.*

*Tarsi postici articulo primo elongato.*

Corpus oblongum, subdepressum, iam alatum, iam apterum. Caput exsertum, thorace angustius, subtriangulare, basi colli ad iustar sat distincte coarctatum, fronte immarginata, oculis parvis, rotundatis, haud prominulis. Ocelli super oculorum marginem superiorem siti. Labrum transversum, medio subsinuatum. Mandibulae breviusculae, validiusculae, acutae, muticae. Maxillae (t. V. f. 27. b.) mala interiore membranacea, extus cornea, oblique truncata, margine ciliato, mala exteriori membranacea, apice subsemicirculariter dilatata, margine barbato. Palpi maxillares articulo primo minuto, secundo, tertio quartoque inter se longitudine subaequalibus, hoc subacuminato. Labium (t. V. f. 27. a.) mento apicem versus sensim angustato, apice truncato, ligula membranacea, subsinuata, lobis exterioribus paulo magis prominulis, ciliatis. Palpi labiales articulis tribus longitudine subaequalibus. Antennae filiformes, graciles, articulis 2 — 10 elongato-subobconicis, ultimo ovato, subacuminato. Thorax coleopteris paulo angustior, transversim subquadratus. Scutellum triangulare. Elytra pectus paulo superantia, angulo exteriori fortiter rotundato. Abdomen depressum, tenuiter marginatum. Pedes modice elongati, tibiis pubescentibus, tarsis anticis leviter dilatatis, subtus villosis, articulis 4 primis transversis, cordatis, quinto his coniunctis longitudine aequali, posterioribus tenuibus, articulis 4 primis simplicibus, subtus longius pubescentibus, articulo quinto his coniunctis brevior, postici articulo primo elongato.

#### **Observationes.**

1. Caput et thorax parcius, elytra densius punctata. Frons ante ocellos lineola brevi longitudinali, ad antennarum insertionem foveola insculpta. Thorax margine laterali medio foveola notatus.

2. Habitus fere *Omalii* floralis, at thoracis forma magis *Acidotam* refert. Maxillarum mala exteriori dilatata *Lathrimaeo* propinquum, tarsorum autem structura et palporum maxillarium articulis ultimis duobus longitudine aequalibus distinctum.

3. Mas abdominis segmento sexto superiore et inferiore truncato, septimo prominulo distinctus.

4. Victus sub lapidibus et sub truncorum cortice.

\* 1. *A. quadrum*: Oblongum, subdepressum, nigrum, antennis pedibusque testaceis, thorace subquadrato parce, elytrisque crebre fortiter punctatis piceis, limbo testaceo. — Long. 2 lin.

*Er. Col. March. I. 619. 1.*

*Omalium quadrum* Grav. Mon. 215. 18. — Oliv. *Encycl. meth. VIII.* 478. 18. —

*Gyll. Ins. Suec. IV. 461. 3—4. — Sahlb. Ins. Fenn. I. 278. 3. — Zetterst. Ins. Lappon. 50. 4. — Mannerh. Brachelytr. 52. 9. — Boisd. et Lacord. Faun. Ent. Paris. I. 487. 4.*

*Anthobium castaneum* Runde *Brachelytr. Hal. 23. 3.*

Habitat in Germania, Suecia.

Corpus oblongum, subdepressum, nigrum, nitidum, glabrum. Antennae capite thoraceque paulo longiores, apicem versus haud incrassatae, rufae, articulo primo testaceo. Palpi testacei. Caput thorace plus duplo minus, parce punctatum, fronte inter oculos et inter antennis foveolis binis sat profundis impressa. Thorax coleopteris vix angustior, latitudine parum brevior, subquadratus, lateribus parum rotundatus, basi apiceque truncatus, angulis omnibus rotundatis, subdepressus, parcius inaequaliter sat fortiter punctatus, ad marginis lateralis medium utrinque foveolatus, margine omni rufo-testaceo. Scutellum parce punctatum. Elytra thorace duplo prope longiora, nonnisi abdominis segmentum primum tegentia, subdepressa, crebre sat fortiter profundeque interius subseriatim punctata, fusco-picea, nitida, limbo exteriori et apicali fusco-testaceo. Abdomen parce subtilissime obsoletissimeque punctulatum, parce subtiliter pubescens, minus nitidum. Pedes rufo-testacei.

Mas tibiis anticis medio dentatis distinctus.

\* 2. *A. brachypterum*: *Piceum, parum nitidum, parce pubescens, capite thoraceque subquadrato parce subtiliterque punctatis, hoc utrinque profunde foveolatum, elytris crebre fortius punctatis.* — Long.  $1\frac{2}{3}$  lin.

*Omalius brachypterum* Grav. *Micr. 114. 5. Mon. 208. 9. — Latr. Hist. nat. Crust. et Ins. IX. 372. 5. — Oliv. Encycl. meth. VIII. 476. 9.*

*Omalius impressum* Zetterst. *Faun. Lappon. I. 52. 9. Ins. Lappon. 51. 14.*

*Omalius Gyllenhalii* Sahlb. *Ins. Fenn. I. 286. 23. — Zetterst. Ins. Lapp. 52. 16.*

*Anthobium Gyllenhalii* Mannerh. *Brachelytr. 54. 10.*

Habitat in Germania, Lapponia, Fennia.

Corpus oblongum, subdepressum, piceum, parum nitidum, minus subtiliter at parce pubescens. Antennae rufo-testaceae, pilosellae. Caput thorace duplo minus, parce subtiliter obsoleteque punctatum, fronte inter oculos striolis duabus minutis, profundis, apice inter antennis foveolis duabus minutis impressa, margine apicali tumidulo, saturatius piceum. Thorax coleopteris vix lator, latitudine paulo brevior, lateribus leviter rotundatus, basi apiceque truncatus, angulis omnibus rotundatis, supra parum convexus, parce subtilissime punctulatus, ad marginem lateralem utrinque paulo ante medium foveola minuta rotundata sat profunda impressus. Scutellum parce subtiliter obsoleteque punctatum. Elytra thorace sesqui longiora, depressa, creberrime minus subtiliter punctata. Abdomen crebrius subtiliter punctulatum. Pedes testacei.

Variat 1, Colore iam magis nigro-, iam magis rufo-piceo, elytris in illis saepius dilutioribus;

2, Elytris thorace iam vix iam plus sesqui-longioribus. Specimina illa aptera, haec alata esse videntur.

Obs. Dom. Prof. Zetterstedt Om. Gyllenhalii ab O. pubescente statura paulo maiore, pubescentia tenuiore, pedibus paulo dilutioribus differe observat, quae



autem differentiae forte non satis constantes me fugiunt; specimina nostra Laponica Fennicaque tam inter se quam cum typo Gravenhorstiano conveniunt, quare nomini Gravenhorstiano illa subiunxi.

- \* 3. *A. humile*: *Lineare, depressum, pubescens, nigrum, capite, thorace elytrisque piceo-testaceis, dense aequaliter punctatis. — Long. vix ultra 1 lin.*

Habitat in montibus Uralensibus, Prof. Eversmann.

Habitu omnino Omalii plani, et minoribus eius individuis aequale, lineare, depressum, pube subtili brevi depressa fulvescenti-grisea densius vestitum. Antennae capitis thoracisque prope longitudine, apicem versus sensim leviterque incrassatae, articulis 4 primis laete testaceis, reliquis fusciscentibus. Caput thoracis prope latitudine, breve, planum, aequale, subtilius at sat profunde dense punctatum, piceo-testaceum, nitidum. Thorax coleopterorum prope latitudine, latitudine dimidio brevior, lateribus leviter rotundatus, basi apiceque truncatus, angulis posterioribus obtusis, subrotundatis, depressus, dense minus subtiliter profunde punctatus, aequalis, piceo-testaceus, nitidus. Elytra thorace duplo longiora, subdepressa, dense fortius profunde punctata, piceo-testacea, nitida. Abdomen parce subtilissime punctatum, nigrum, nitidum. Pectus nigrum, nitidum. Pedes flavo-testacei.

Obs. Exemplum singulum palporum tarsorumque structura huius generis esse videtur.

*Species mihi invisae, forte huius generis.*

4. *A. Sibiricum*: *Oblongum, glabrum, nigrum, antennis, thoracis limbo, elytris pedibusque testaceis.*

*Omalium Sibiricum* Mannerh. *Brachelytr.* 52. 8.

Habitat prope Barnaul Sibiriae.

A. quadro simillimum, parcius punctatum, capite thoraceque respecta magnitudine multo minoribus et abdomine longiore fere unice diversum. (Mannerh.)

### **Acidota Leach.**

*Mannerh. Brachelytr. p. 55. — Steph. Illustr. V. p. 358. — Boisd. et Lacord. Faun. Ent. Paris. I. p. 476. — Er. Col. Murch. I. p. 620.*

*Mandibulae muticae.*

*Maxillae malis membranaceis.*

*Palpi maxillares articulo ultimo penultimo vix longiore.*

*Tibiae spinulosae.*

*Tarsi postici articulo primo elongato.*

Corpus elongatum, subdepressum, alatum. Caput exsertum, thorace angustius et multo minus, oblongum, basi vix constrictum, fronte apice marginata, oculis parvis, prominulis. Ocelli super oculorum marginem superiorem positi. Labrum breve, transversum, integrum. Mandibulae validiusculae, at breves, muticae. Maxillae (t. V. f. 28. b.) malis membranaceis, exteriori interiorem paulo superante, hac interne ciliata, illa apice barbatula. Palpi maxillares modice elongati, articulo primo minuto, secundo longiore tertioque hoc paulo brevior leviter obconicis, quarto secundo longitudine

aequali, tertio paululum longiore, oblongo-ovato, subacuminato. Labium (t. V. f. 28. a.) mento transverso, apice truncato, ligula membranea, profunde biloba, lobis interne sinuatis. Palpi labiales articulo primo longiore, secundo hoc duplo brevior, tertio primo aequali, subacuminato. Antennae filiformes, articulo primo tumidulo, ovato, secundo tertio paululum brevior, tertioque subobconicis, sequentibus longitudine sensim licet levissime decrescentibus, penultimis crassitie vix longioribus, ultimo oblongo-ovato, acuminato. Thorax sat amplus, apicem versus subangustatus, lateribus late marginatis. Scutellum triangulare. Elytra abdominis primum segmentum obtegentia, angulo exteriori rotundato. Abdomen late marginatum, parallelum, apice rotundato-acuminatum. Pedes breviores, tibiis subtiliter spinulosis, tarsis omnibus simplicibus: antici articulis 4 primis inter se aequalibus, brevibus, quinto his coniunctis longitudine aequali, intermedii articulo primo elongato, sequentibus tribus inter se aequalibus, brevibus, quinto reliquis quatuor coniunctis paulo brevior, postici articulo primo elongato, secundo subelongato, tertio quartoque inter se aequalibus, brevibus, quinto reliquis coniunctis dimidio brevior.

## O b s e r v a t i o n e s.

1. Corpus glabrum. Caput et thorax fortius, abdomen subtilius, elytra subseriatim punctata. Frons antice ad antennarum insertionem utrinque foveolata impressave.

2. A Deliphro tarsorum posteriorum articulo primo magis quam secundo elongato, ligula profunde biloba etc., a reliquis Omalinis pedibus validiusculis, tibiis distincte spinulosis differt. Ligulae structura cum Anthophago et Anthobio congruit.

3. Mas segmento utroque sexto truncato, septimo conspicuo distinctus.

4. Victus in sylvis sub muscis.

\* 1. *A. crenata*: *Picea*, antennis pedibusque rufis, thorace leviter convexo, aequali, elytris fortiter striato-punctatis. — Long. 3 lin.

*Mannerh. Brachelytr.* 55. 3. — *Steph. Illustr. V. t.* 127. f. 6. — *Boisd. et Lacord. Faun. Ent. Paris.* I. 476. 1. — *Er. Col. March.* I. 620. 1.

*Staph. crenatus* *Fab. Ent. Syst.* I. II. 525. 26. *Syst. El.* II. 596. 34. — *Payk. Mon. Curc. App.* 146. 45. *Faun. Suec.* III. 403. 47. — *Panz. Ent. Germ.* 354. 18. — *Walk. Faun. Paris.* I. 272. 15.

*Omal. crenatum* *Oliv. Encycl. meth.* VIII. 476. 6. — *Gyll. Ins. Suec.* II. 230. 27. — *Sahlb. Ins. Fenn.* I. 288. 26. — *Zetterst. Faun. Lappon.* I. 55. 15. *Ins. Lappon.* 53. 23.

*Omal. castaneum* *Grav. Mon.* 207. 8. — *Oliv. Encycl. meth.* VIII. 476. 8.

*Var. Rufo-testacea.*

*Omal. rufum* *Grav. Micr.* 115. 6. *Mon.* 207. 7. — *Latr. Hist. nat. Crust. et Ins.* IX. 372. 6. — *Oliv. Encycl. meth.* VIII. 476. 7.

*Acidota rufa* *Mannerh. Brachelytr.* 55. 1. (*forte*)

Habitat in Europa.

Corpus elongatum, lineare, glabrum, iam obscure castaneo-piceum, abdominis segmentorum marginibus rufis, iam totum rufum vel rufo-testaceum. Antennae capite thoraceque paulo longiores, apice vix crassiores, rufo-ferrugineae. Palpi testacei. Caput parvum, thorace quadruplo minus, crebre profundeque punctatum, fronte plana, antice inter antennis bifoveolata, margine antico elevato. Thorax basi coleopterorum prope latitudine,

latitudine haud brevior, lateribus leviter rotundatus, antrorsum leviter angustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusis, supra leviter convexus, crebre profundeque punctatus; linea media utrinque abbreviata laevi, margine laterali explanato, relevato, medio subfoveolato. Scutellum parum-punctatum. Elytra thorace plus sesqui longiora, fortiter profundeque interne sat regulariter striato-punctata. Abdomen minus fortiter crebre punctatum. Pedes rufi.

- \* 2. *A. cruentata*: Rufo-testacea, thorace subdepresso, longitudinaliter bi-impresso, elytris thorace duplo longioribus, interne seriatim punctatis. — Long.  $2\frac{1}{3}$  lin.

Mannerh. Brachelytr. 55. 2. — Er. Col. March. 1. 621. 2.

Omal. rufum Gyll. Ins. Suec. II. 228. 26. — Sahlb. Ins. Fenn. I. 288. 25.

Habitat in Germania, Suecia.

Praecedente minor, brevior et magis depressa, rufo-testacea, nitida, glabra. Antennae capite thoraceque longiores, crassiusculae, apicem versus haud incrassatae, articulis 4 — 10 cylindricis, sensim paulo brevioribus, prioribus eorum crassitie longioribus, longius pubescentes. Caput thorace triplo fere minus, crebre fortiter punctatum, fronte inter oculos bifoveolata, anterius longitudinaliter utrinque impressa, margine antico elevato. Thorax coleopteris parum angustior, latitudine dimidio fere brevior, lateribus leviter rotundatus, antrorsum subangustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, parum convexus, crebre sat profunde punctatus, dorso depressiusculus, obsolete longitudinaliter bi-impressus, ad marginis lateralis medium utrinque foveola rotundata sat profunda notatus. Scutellum basi punctatum. Elytra thorace duplo fere longiora, profunde punctata, punctis ad suturam in series sat regulares dispositis. Abdomen minus fortiter crebrius punctatum, dorso rufo-piceum, segmentorum marginibus rufo-testaceis. Pectus piceum. Pedes rufo-testacei.

- \* 3. *A. ferruginea*: Rufo-testacea, nitida, thorace depresso, obsolete longitudinaliter bi-impresso, elytris thorace paulo longioribus, interne subseriatim punctatis. — Long. 2 lin.

Boisd. et Lacord. Faun. Ent. Paris. I. 477. 2. (forte)

Omalium quadrum Zetterst. Faun. Lappon. I. 54. 14.

Omalium quadratum Zetterst. Ins. Lappon. 53. 22.

Habitat in Suecia. Exemplum a Gyllenhal nomine Omal. rufi missum dedit Dom. Schüppel.

Praecedente plus dimidio minor, rufo-testacea, nitida, glabra. Antennae capitis thoracisque longitudine, crassiusculae, apicem versus haud incrassatae, articulis 4 — 10 cylindricis, primis eorum crassitie vix longioribus, ultimis subbrevioribus, pilosellae. Caput thorace triplo fere minus, crebrius profundiusque punctatum, punctis in media fronte rarioribus, hac inter oculos impressionibus duabus obliquis, postice conniventibus, antice foveola profundiore terminatis, inter antennas foveolis duabus latioribus notata, margine anteriore praecipue ad latera elevato. Thorax coleopterorum latitudine, latitudine dimidio brevior, antrorsum vix angustatus, lateribus perparum rotundatus, basi apiceque truncatus, angulis posterioribus subrectis, antero-



ribus rotundatis, parum convexus, crebre fortius profunde punctatus, dorso depressus, obsolete longitudinaliter bi-impressus, utrinque prope marginis lateralis medium foveola rotundata profunde impressus. Scutellum basi punctatum. Elytra thorace tertia parte longiora, subdepressa, crebre profundeque punctata, punctis interne in series nonnullas minus regulares, geminatas digestis. Abdomen crebre fortius punctatum. Pedes dilutius testacei.

Obs. Distincta videtur a praecedente statura minore, antennis elytrisque brevioribus, his minus regulariter striato-punctatis, abdomine minus subtiliter punctato, etc.

- \* 4. *A. subcarinata*: *Pubescens, rufo-picea, capite, thoracis abdominisque dorso nigricantibus, thorace subcarinato, elytris crebre profunde punctatis.* — Long.  $2\frac{1}{4}$  lin.

Habitat Bostoniae in America septentrionali, Dom. Zimmermann.

Statura omnino praecedentium, subdepressa, pube minus subtili fulvescenti vestita. Antennae capite thoraceque vix longiores, articulis 4—10 subcylindricis, sensim paulo crassioribus at vix brevioribus, rufae. Palpi rufi. Caput thorace quadruplo prope minus, crebrius minus fortiter at profunde punctatum, fronte subaequali, inter oculos et inter antennis foveolis binis minutis levius impressa, margine antico leviter elevato, nitidum, nigrum, antice rufescens. Thorax coleopteris parum angustior, latitudine paululum brevior, antrorsum subangustatus, lateribus modice rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusis, leviter convexus, crebrius fortiusque profunde punctatus, dorso depressiusculus, longitudinaliter late obsolete bi-impressus, interstitio subcarinato, laevi, utrinque prope marginis lateralis medium foveola rotunda impressus, margine laterali explanato, relevato, obsolete crenulato, piceus, nitidus, limbo omni rufescente. Scutellum basi punctatum. Elytra thorace plus sesqui longiora, margine laterali planato, apicali depresso, tota aequaliter crebre fortiter profundeque punctata, rufo-picea, nitida. Abdomen crebre minus subtiliter punctulatum, subtus piceo-rufum, supra nigro-piceum, limbo laterali rufescente, nitidum. Pectus piceo-rufum. Pedes rufi.

Obs. Nomine *Omalii subcarinati* Say i. litt. communicata.

## **Olophrum Er.**

*Er. Col. March. I. p. 622.*

*Mandibulae muticae.*

*Maxillae malis elongatis, corneis.*

*Palpi maxillares articulo penultimo ultimo duplo brevior.*

*Tibiae muticae.*

*Tarsi postici articulis duobus primis subelongatis, inter se aequalibus.*

Corpus crassiusculum, convexiusculum, ovatum vel oblongum, iam alatum, iam apterum. Caput thorace multo minus et angustius, subdeflexum, basi haud constrictum, fronte immarginata, oculis parvis, subprominulis. Ocelli super oculorum marginem superiorem siti. Labrum (t. V. f. 29. c.) transversum, integrum, apice membrana prominula biloba auctum. Mandibulae validiusculae, subrectae, muticae, apice acuto introrsum curvato. Ma-

xillae (t. V. f. 29. b.) malis elongatis, corneis, interiore interne tenuiter ciliata, exteriore apice barbatula. Palpi maxillares articulo primo minuto, secundo sat elongato, tertio brevior, subobconico, quarto iterum longior, praecedente duplo fere longior, subacuminato. Labium (t. V. f. 29. a) mento latitudine baseos brevior, antrorsum valde angustato, apice subemarginato, ligula membranea, apice truncata vel subsinuata. Palpi labiales articulo primo elongato, secundo tertioque minutis, illo paulo brevior. Antennae filiformes, articulo primo parum crassior, secundo tertio vix brevior, sequentibus subcylindricis, longitudine sensim levissime decrescentibus, penultimo crassitie sua vix longior, ultimo ovato, acuminato. Thorax sat amplus, coleopterorum fere longitudine. Elytra sat ampla, abdominis maiorem minoremve partem obtegentia. Abdomen tenuiter marginatum, apice subacuminatum. Pedes minus elongati, tibiis inermibus, pubescentibus, tarsis anticis articulis 4 primis subdilatatis, transversim cordatis, quinto his coniunctis paulo brevior, intermediis articulo primo subelongato, sequentibus tribus brevibus, inter se aequalibus, quinto his coniunctis (primo excepto) longitudine aequali, posticis articulis primis duobus subelongatis, tertio quartoque brevioribus, quinto praecedentibus tribus coniunctis longitudine fere aequali. Tarsorum omnium articuli 4 primi subtus longius densiusque pubescentes.

#### O b s e r v a t i o n e s .

1. Corpus glabrum. Caput, thorax et elytra punctata. Abdomen glabrum. Frons inter oculos lineolis foveolisve nullis insculpta.
2. Maxillarum structura *Lesteuae* proximum.
3. Mas abdominis segmento sexto truncatulo, septimo prominulo distinctus.
4. Victus in sylvis sub musco, foliis deciduis, lapidibus.

\* 1. *O. piceum*: *Convexum, piceum, nitidum, antennis pedibusque rufis, elytris fortiter vage punctatis.* — Long:  $2\frac{1}{4}$  lin.

*Er. Col. March. I. 623. 1.*

*Omalium piceum* Gyll. *Ins. Suec. II. 200. 3.* — *Zetterst. Faun. Lappon. I. 49. 3. Ins. Lappon. 50. 5.* — *Mannerh. Brachelytr. 51. 4.* — *Boisd. et Lacord. Faun. Ent. Paris. I. 485. 1.* — *Runde Brachelytr. Hal. 21. 1.*

*Silpha Blattoides Ahrens Act. nov. Halens. II. 11. 20. 12. t. 2. f. 3.*

Habitat in Germania, Hungaria, Anglia, Suecia.

Corpus breve, convexum, brunneo-piceum, saepe etiam rufo-piceum vel testaceum. Antennae capitis thoracisque fere longitudine, rufae vel rufo-testaceae. Caput maiusculum, crebrius fortiusque punctatum, inter antennis foveolis duabus minutis impressum, ore testaceo. Thorax coleopterorum latitudine, latitudine duplo fere brevior, basi levissime rotundatus, apice leviter emarginatus, lateribus rotundatus, angulis anterioribus posterioribusque aequaliter rotundatis, apice basi latitudine aequali, margine laterali late explanato, disco convexo, sat crebre profunde fortiterque punctatus. Scutellum laeve. Elytra thorace duplo longiora, basi thoracis medio angustiora, apicem versus sensim leviter dilatata, convexa, fortiter profundeque sat crebre aequaliter punctata. Abdomen laeve, apice piceo-testaceum. Pedes rufi vel testacei.

- \* 2. *O. obtectum*: Nigro-piceum, antennis pedibusque rufis, elytris dense vage punctatis. — Long. 2 lin.

Habitat in America septentrionali, Dom. Zimmermann.

Statura omnino et summa affinitas praecedentis. Nigro-piceum, minus nitidum. Antennae capitis thoracisque longitudine, rufo-testaceae. Caput crebrius punctatum, inter oculos et inter antennis obsolete bifoveolatum, ore testaceo. Thorax coleopteris paulo angustior, latitudine duplo brevior, basi levissime rotundatus, apice leviter emarginatus, lateribus rotundatus, angulis anterioribus rotundatis, posterioribus obtusis, subrotundatis, leviter transversim convexus, lateribus marginatus, margine autem haud explanato, dense punctatus, marginibus rufo-piceis. Scutellum parce punctatum. Elytra thorace plus duplo longiora, apicem versus sensim leniter dilatata, dense sat profunde punctata, margine apicali rufo-piceo. Abdomen laeve, apice testaceum. Pedes rufi, femoribus nonnunquam picescentibus.

Obs. Differt ab *O. piceo* corpore minus convexo, punctura subtiliore, capite minore, thorace angustiore, elytris longioribus.

- \* 3. *O. fuscum*: Leviter convexum, piceum, antennarum basi pedibusque rufis, elytris fortiter punctatis, punctis interne subseriatis. — Long. 2 lin.

Er. Col. March. I. 623. 2.

*Omalius fuscum* Grav. Mon. 211. 16. — Oliv. Encycl. meth. VIII. 478. 17.

Habitat in Germania.

Habitu et summa affinitate *O. picei*, at dimidio minus, angustius, minus convexum, brunneo-piceum, nitidulum. Antennae capite thoraceque paulo longiores, piceae, apice rufescentes, articulo primo rufo, interdum totae rufae. Caput crassiusculum, vage crebrius sat fortiter punctatum, fronte inter oculos foveolis duabus obsoletis, inter antennis duabus profundioribus impressa, palpis mandibulisque rufo-piceis. Thorax coleopterorum basi latitudine aequalis, latitudine dimidio brevior, lateribus leviter rotundatus, apice basi angustior, angulis anterioribus rotundatis, posterioribus obtusis, subrotundatis, margine laterali late explanato, disco modice convexo, crebrius vage fortius punctatus, versus margines rufescens. Scutellum punctis paucis profundis impressum. Elytra thorace duplo prope longiora, apicem versus sensim leniter dilatata, parum convexa, crebre sat fortiter punctata, punctis anterioribus versus suturam subseriatis. Abdomen laeve. Pedes rufi.

Obs. Variat minus maturum corpore rufo-testaceo: tale specimen *O. assimili* simile, distinctum autem statura paulo maiore, elytris longioribus, densius punctatis, frontis foveolis anticis maioribus.

- \* 4. *O. assimile*: Testaceum, abdomine piceo, elytris fortiter punctatis. — Long.  $1\frac{2}{3}$  lin.

Er. Col. March. I. 624. 3.

*Staph. assimilis* Payk. Faun. Suec. III. 409. 53.

*Omalius assimile* Gyll. Ins. Suec. II. 202. 4. — Sahlb. Ins. Fenn. I. 278. 4. —

Mannerh. Brachelytr. 51. 5.

Habitat in Germania, Suecia, in silvis, sub foliis deciduis putrescentibus.

Habitu omnino *O. picei*, at plus duplo minus. Antennae capite thoraceque paulo longiores, apicem versus paulo crassiores, pilosellae, testaceae.



Caput crassiusculum, crebre profunde punctatum, apice inter antennas foveolis duabus minutis notatum, rufo-testaceum, nitidum. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus leviter rotundatus, angulis anterioribus rotundatis, posterioribus obtusiusculis, basi apiceque latitudine aequalibus, margine laterali explanato, disco convexus, fortius profunde sat crebre punctatus, ad medium marginis lateralis utrinque foveolatus, rufo-testaceus, nitidus. Scutellum parce punctatum, rufo-testaceum. Elytra thorace sesqui longiora, parum convexa, crebrius fortiter profundeque punctata, testacea. Abdomen testaceo-piceum. Pectus testaceum. Pedes flavo-testacei.

\* 5. *O. boreale*: Oblongum, nigro-piceum, nitidum, antennarum basi, thoracis elytrorumque limbo pedibusque testaceis, thorace parcius profunde punctato, lateribus postice subsinuato. — Long. 2 lin.

*Omalium boreale* Gyll. Ins. Suec. II. 198. 1. — Sahlb. Ins. Fenn. I. 277. 1. — Zetterst. Faun. Lappon. I. 48. 1. Ins. Lappon. 49. 1. — Mannerh. Brachelytr. 51. 1. *Staph. borealis* Payk. Mon. Curc. App. 146. 47 — 48. — Faun. Suec. III. 411. 57. (mas)

Habitat in Lapponia, Dalecarlia, Dom. Schüppel.

Corpus oblongum, subdepressum, nigro-piceum, nitidum, glabrum. Antennae capite thoraceque paulo longiores, tenuiores, apicem versus vix incrassatae, articulo ultimo oblongo, apice abrupte subacuminato, piceae, basi dilutiores, rufae. Palpi picei. Caput crebre fortius punctatum, inter antennas obsoletius bifoveolatum, apice laevigatum. Thorax coleopteris paulo angustior, latitudine plus dimidio brevior, lateribus fortiter rotundatis, basin versus leviter angustatis, postice subsinuatis, basi levissime rotundatus, apice truncatus, angulis posterioribus obtusis, at nullo modo rotundatis, anterioribus rotundatis, leviter convexus, parcius profunde punctatus, dorso ante basin et margine laterali leviter explanato ante medium foveolatis, limbo omni dilutiore, rufo-piceo. Scutellum laeve. Elytra thorace plus duplo longiora, crebre, fortiter profundeque punctata, punctis suturam versus subsinuatis, margine apicali testaceo. Pectus minus fortiter punctatum. Abdomen supra laevigatum, subtus parcius punctulatum, apice piceo-testaceum. Pedes dilute rufo-picei.

Obs. *O. fusco* affine, distinctum autem statura angustiore, thorace lateribus postice subsinuato, angulis posterioribus nullo modo rotundatis.

\* 6. *O. rotundicolle*: Oblongum, convexum, piceum, thorace dense profundius punctato, angulis posterioribus fortiter rotundatis. — Long.  $2\frac{2}{3}$  lin.

*Omal. rotundicolle* Sahlb. Ins. Fenn. I. 281. 11. — Zetterst. Ins. Lappon. 49. 3. — Mannerh. Brachelytr. 51. 3.

Habitat in Fennia, sec. Sahlberg, in Dalecarlia, Dom. Schüppel.

Habitu omnino *O. consimilis*, at paulo maius, piceum, nitidum, glabrum. Antennae capitis thoracisque prope longitudine, tenuiores, apicem versus haud incrassatae. Palpi testacei. Caput dense fortius profunde punctatum, inter antennas leviter bifoveolatum, fronte planiuscula, basi incrassatum. Thorax coleopteris vix angustior, latitudine parum brevior, apicem versus subangustatus, lateribus rectis, basi leviter rotundatus, apice leviter bisinua-

tus, angulis anterioribus obtusis, posterioribus late fortiterque rotundatis, fortiter convexus, dense fortius profunde punctatus, punctis in medio dorso fortioribus et minus confertis, anterieus medio longitudinaliter impressus. Scutellum profunde punctatum. Elytra thorace duplo prope longiora, modice convexa, dense fortiter profundeque punctata, punctis vix seriatis, nonnisi abdominis segmentum primum obtegentia. Abdomen supra laeve, subtus parce obsoletius punctatum, parum nitidum. Pectus profunde punctatum. Pedes piceo-rufi.

- \* 7. *O. alpestre*: Oblongum, convexum, piceum, elytris limbo apicali testaceo, thorace dense profundeque punctato, lateribus postice subsinuato. — Long.  $2\frac{1}{2}$  lin.

Habitat in Helvetiae alpidibus, Dom. Prof. Heer.

O. fusco vix latius, at sesqui fere longius, convexum, piceum, nitidum, glabrum. Antennae capitis thoracisque fere longitudine, apicem versus vix incrassatae, rufo-piceae, basi dilutiores. Caput fortius profunde punctatum, fronte antice inter antennis transversim impressa utrinque subfoveolata, margine antico incrassato, laevigato, ore piceo-rufo. Thorax coleopteris parum angustior, latitudine dimidio fere brevior, basi apiceque truncatus, lateribus rotundatis, postice subsinuatis, basin versus subangustatus, angulis anterioribus rotundatis, posterioribus obtusiusculis, leviter prominulis, valde convexus, dense profunde punctatus, ante basin nonnunquam obsolete impressus, limbo omni rufo-piceo. Scutellum crebre profundeque punctatum. Elytra thorace duplo longiora, abdominis partem minorem tegentia, convexa, fortiter profundeque crebre punctata, punctis suturam versus subseriatis, rufo-picea, limbo apicali testaceo. Abdomen laeve, apice summo piceo-testaceo. Pedes rufi vel picei.

- \* 8. *O. consimile*: Oblongum, nigrum, subnitidum, antennis, pedibus elytrisque piceis, his margine apicali testaceo, thorace convexo, confertim subtilius punctato, lateribus postice subsinuato. — Long.  $2\frac{1}{3}$  lin.

*Omal. consimile* Gyll. Ins. Suec. II. 199. 2. — Sahlb. Ins. Fenn. I. 277. 2. — Zetterst. Faun. Lappon. I. 48. 2. Ins. Lappon. 49. 2. — Mannerh. Brachelytr. 51. 2. *Staph. borealis* fem. Payk. Faun. Suec. III. 411. 57.

Habitat in Lapponia, Dalecarlia, Dom. Schüppel, in Rossia, Redowsky.

O. boreali paulo magis oblongum, nigrum, subnitidum, glabrum. Antennae capitis thoracisque prope longitudine, tenuiores, apicem versus haud incrassatae, piceae. Palpi picei. Caput minus fortiter densius punctatum, fronte inter antennis bifoveolata. Thorax coleopteris paulo angustior, lateribus rotundatis, postice subsinuatis, basi leviter rotundatus, apice subsinuatus, angulis anterioribus rotundatis, posterioribus obtusis, leviter prominulis, minus fortiter licet profunde confertim punctatus, ante basin medio impressus, limbo laterali anguste explanato, parcius profundiusque punctato. Scutellum parce profundeque punctatum. Elytra thorace plus duplo longiora, aequaliter densius fortiter profundeque punctata, picea, limbo apicali testaceo, abdominis tertiam fere partem obtegentia. Abdomen supra laeve, subtus parcius obsoletiusque punctatum. Pectus fortiter punctatum. Pedes picei.

## Species mihi invisae.

9. *O. emarginatum*: Nigro-piceum, thorace postice rotundato, antice emarginato. — Long. infra 2 lin.

*Omalium rotundicolle* Say Transact. Amer. Philoph. Soc. Philadelph. Nov. Ser. IV. 462. 2.

Habitat in Pennsylvania.

Grosse profundeque crebre punctatum. Antennae fusco-piceae, articulis tribus primis testaceis. Thorax transversim ovalis, antice profunde emarginatus, aequaliter punctatus, angulis posterioribus rotundatis, marginibus piceis. Elytra aequaliter punctata, longa, margine apicali piceo. Abdomen supra piceum, segmentorum marginibus apicalibus dilutioribus, infra pice-scens. Pedes saturate testacei. (Say.)

**Lathrimaeum Er.**

Er. Col. March. I. p. 624.

*Mandibulae muticae.*

*Maxillae malis membranaceis, exteriore dilatata.*

*Palpi maxillares articulo ultimo penultimo longiore.*

*Tibiae muticae.*

*Tarsi postici articulis duobus primis subelongatis, inter se aequalibus.*

Corpus latum, subdepressum, alatum apterumve. Caput thorace multo minus et angustius, basi sat distincte constrictum, fronte marginata, oculis mediocribus, rotundatis, prominulis. Ocelli super oculorum marginem superiorem siti. Labrum (t. V. f. 30. c.) transversum, truncatum, apice utrinque lacinia lineari membranacea auctum. Mandibulae sat breves, muticae. Maxillae (t. V. f. 30. b.) mala interiore membranacea, extus cornea, denticulo corneo terminata, apice oblique truncata, ciliata, exteriore membranacea, apice subsemicirculariter dilatata, barbata. Palpi maxillares articulo primo minuto, secundo sat elongato, tertio brevior, quarto hoc plerumque duplo longior, apice subacuminato. Labium (t. V. f. 30. a.) mento brevi, antrorsum angustato, apice truncato, ligula membranacea, transversa, apice truncata, medio leviter incisa. Palpi labiales breviusculi, articulo primo tertioque subaequalibus, secundo his brevior. Antennae filiformes, articulo secundo tertio paululum brevior, sequentibus longitudine sensim levissime decrescentibus, penultimo crassitie sua haud brevior, ultimo ovato. Thorax amplus, coleopteris latitudine subaequalis, lateribus late marginatus. Scutellum triangulare. Elytra sat ampla, abdominis maximam partem tegentia, angulo apicis exteriore rotundato. Abdomen breviusculum, apice subacuminatum. Pedes minus elongati, tibiis simplicibus, pubescentibus, tarsis anticis simplicibus, articulis 4 primis brevibus, inter se aequalibus, articulo quinto his coniunctis longitudine aequali, intermediis articulis 4 primis longitudine decrescentibus, primo subelongato, quinto his coniunctis brevior, posticis articulis duobus primis subelongatis, inter se aequalibus, tertio brevior, quarto brevi, quinto his coniunctis brevior.

## O b s e r v a t i o n e s.

1. Caput, thorax et elytra punctata, plerumque glabra; caput distincte foveolatum; thorax magis minusve distincte canaliculatus, dorso postico saepius foveis duabus obliquis postice conniventibus obsolete impressus; elytra ampla.



2. Habitus fere generis praecedentis, at maxillarum, malae neque elongatae, neque corneae, exterior externe rotundato-ampliata, ut in Arpedio; ab Olophro palporum maxillarum articulo ultimo minus elongato et fronte profundius foveolata distinguendum.

3. Sexus differentia ut in praecedentibus.

4. Victus in silvis sub musco et foliis putrescentibus.

\* 1. *L. melanocephalum*: Testaceum, capile nigro thoraceque transverso, canaliculato, late marginato, apice profunde emarginato profunde punctatis, elytris amplis, creberrime fortiter subseriatim punctatis. — Long. 2 lin.

*Silpha melanocephala* Illig. *Schneid. Mag. V.* 596. 4. *Kaef. Preuss.* 356. 2. — *Panz. Ent. Germ. I.* 122. 19. *Faun. Germ.* 25 5.

Habitat in Germania.

Corpus oblongo-subquadratum, parum convexum, testaceum, nitidum, glabrum. Antennae capitis thoracisque fere longitudine, articulis 4 — 10 sensim paulo brevioribus et crassioribus, ultimo subovato, apice acuminato, nigrae, articulis 4 primis testaceis. Palpi testacei. Caput thorace triplo fere minus, creberrime punctatum, apice laeve, supra utrinque longitudinaliter impressum, inter antennis utrinque foveolatum, margine anteriore utrinque elevato, nigro-subaeneum vel piceum, nitidum. Thorax coleopteris parum angustior, latitudine media duplo fere brevior, lateribus rotundatis, basi truncatus, apice profunde emarginatus, angulis anterioribus acutiusculis, prominentibus, posterioribus subrectis, leviter prominulis, parum convexus, distinctius canaliculatus, basi supra scutellum leviter impressus, lateribus late explanatis, medio impressis. Scutellum parce punctatum. Elytra thorace plus duplo longiora, abdomen fere totum tegentia, apice coniunctum rotundata, parum convexa, creberrime sat fortiter punctata, punctis plerisque in series sat regulares dispositis. Abdomen laeve. Pectus profunde punctatum, interdum piceo-testaceum, Pedes testacei.

\* 2. *L. luteum*: Testaceum, pectore nigro, thorace transverso, apice profundius emarginato, dorso utrinque obsolete oblique impresso, profunde punctato, elytris amplis, fortiter creberrime subseriatim punctatis. — Long. 1 $\frac{2}{3}$  lin.

Habitat in Lusitania.

*L. melanocephalo* affine, testaceum, subnitidum, glabrum. Antennae capitis thoracisque longitudine, apicem versus sensim leniterque incrassatae, articulo ultimo ovato, apice fusciscentes. Caput thorace duplo minus, posterius fortius sat crebre, antierius subtiliter parce punctatum, fronte inter oculos et inter antennis utrinque impressa, margine laterali supra antennarum insertionem elevato. Thorax coleopteris parum angustior, latitudine plus dimidio brevior, lateribus rotundatus, basi truncatus, apice emarginatus, angulis anterioribus rotundatis, posterioribus obtusiusculis, subdentatis, leviter convexus, crebre sat fortiter punctatus, dorso pone medium foveolis duabus oblongis, obliquis, postice conniventibus, obsolete impressus, margine laterali minus late explanato, medio foveolato. Scutellum parce punctatum. Elytra ampla, abdomen fere totum tegentia, thorace plus duplo longiora, modice convexa, sat crebre fortiterque subseriatim punctata. Abdomen supra

laeve, subtus lateribus obsoletius punctulatum. Pectus fortiter punctatum, nigrum. Pedes testacei.

Obs. Intermedium quasi inter *L. melanocephalum* et *atrocephalum*, ab utroque autem thorace haud canaliculato distinctum.

\* 3. *L. atrocephalum*: Testaceum, capite nigro-piceo thoraceque transverso, obsolete canaliculato, apice leviter emarginato profunde punctatis, elytris crebre fortiterque subseriatim punctatis. — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 625. 1.*

*Omali. atrocephalum* Gyll. *Ins. Suec. IV. 463. 4—5.* — *Boisd. et Lacord. Faun. Ent. Paris. I. 486. 3.*

*Omali. melanocephalum* Runde *Brachelytr. Hal. 22. 3.*

*Silpha melanocephala* Marsh. *Ent. Britt. 127. 39.*

Habitat in Germania, Suecia.

Subovatum, leviter convexum, testaceum, nitidum, glabrum. Antennae capitis thoracisque longitudine, apice sensim leviterque incrassatae, testaceae, apice fuscae. Caput thorace duplo fere minus, posterius sat crebre fortiterque, anterieus parce subtiliterque punctatum, fronte inter oculos oblique, inter antennis profundius utrinque impressa, margine laterali supra antennarum basin leviter elevato, piceum, nitidum. Thorax coleopteris paulo angustior, latitudine duplo fere brevior, lateribus rotundatus, basi truncatus, apice leviter late emarginatus, angulis anterioribus rotundatis, posterioribus obtusiusculis, subdentatis, modice convexus, sat fortiter profundeque dense punctatus, obsoletius canaliculatus, basi supra scutellum obsolete impressum, margine laterali minus late explanato, modice foveolato. Scutellum parce punctatum. Elytra ampla, abdomen fere totum tegentia, thorace plus duplo longiora, leviter convexa, sat fortiter crebreque subseriatim punctata. Abdomen laeve. Pectus dense profundeque punctatum.

Variat rarissime corpore nigro, antennis concoloribus, pedibus solis piceis. Specimen singulum Dom. Schüppel Berolini legit.

\* 4. *L. fuscum*: Fuscum, nitidum, capite nigro, thorace elytrisque confertissime punctatis, thorace transverso, obsolete canaliculato, dorso utrinque oblique impresso, apice leviter emarginato. — Long.  $1\frac{1}{4}$  lin.

*Er. Col. March. I. 626. 2.*

Habitat in Germania.

*L. atrocephalo* minus, praesertim angustius et magis depressum, obscure fuscum, subnitidum, glabrum. Antennae capitis thoracisque longitudine, apice sensim leviter incrassatae, fusco-testaceae, apice nigrae. Caput thorace duplo fere minus, creberrime punctatum, apice laevi, fronte inter oculos utrinque impressa, inter antennis utrinque profunde foveolata, margine laterali supra antennarum basin elevato, nigrum, ore rufo-piceo. Thorax coleopterorum basi parum angustior, latitudine duplo fere brevior, lateribus rotundatus, basi truncatus, apice leviter emarginatus, angulis anterioribus subrotundatis, posterioribus obtusis, parum convexus, dense punctatus, obsolete canaliculatus, postice foveis duabus oblongis, obliquis postice conniventibus impressus, margine laterali minus late explanato, medio foveolato, nigro-fuscus, marginibus dilutioribus. Scutellum parce punctatum. Elytra thorace duplo longiora, abdomen dimidium tegentia, apicem versus

sensim leviter dilatata, subdepressa, confertissime punctata, fusca vel fusco-testacea. Abdomen subtilissime punctulatum, nigro-piceum. Pectus fortiter punctatum. Pedes fusco-testacei.

- \* 5. *L. sordidum*: *Dilute testaceum, nitidum, pectore abdomineque nigro-piceis, capite, thorace elytrisque fortiter profundeque punctatis, thorace transverso, apice truncato.* — Long.  $1\frac{1}{2}$  lin.

Habitat in America septentrionali, e collectione Knochiana.

*L. atrocephalo* angustius, magis depressum, flavo-testaceum, nitidum, pectore abdomineque piceis. Antennae capitis thoracisque longitudine, apicem versus sensim leviterque incrassatae. Caput thorace duplo fere minus, parcius punctatum, fronte inter oculos et inter antennis foveolis binis oblongis minutis impressa. Thorax coleopterorum basi parum angustior, latitudine dimidio brevior, lateribus rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusis, parum convexus, sat fortiter profundeque vage punctatus, dorso foveola media duabusque obliquis, omnibus valde obsoletis, lateribus utrinque foveola profundiore impressus. Scutellum laeve. Elytra thorace duplo longiora, abdominis tertiam fere partem obtegentia, apicem versus sensim leviterque dilatata, subdepressa, secundum suturam impressa, crebre fortiter profundeque punctata. Abdomen parce obsoletissime punctulatum. Pedes pallide testacei.

- \* 6. *L. canaliculatum*: *Nigrum, pubescens, ore, antennarum basi, thorace subquadrato pedibusque testaceis, elytris fortiter punctatis, flavis.* — Long. 2 —  $2\frac{1}{3}$  lin.

Habitat in Germaniae, Austriae, Hungariae, Syrmiae montibus.

Antennae capite thoraceque paulo longiores, apicem versus sensim subincrassatae, fuscae, articulis 4 primis testaceis. Caput thorace dimidio fere angustius, oblongum, parcius obsoletiusque punctatum, fronte depressa, inter oculos utrinque sulco minus profundo impressa, nigrum, ore late testaceo. Thorax coleopterorum basi parum angustior, maris latitudine perparum, feminae dimidio brevior, lateribus leviter rotundatis, basi truncatus, apice subemarginatus, angulis anterioribus rotundatis, posterioribus obtusiusculis, leviter convexus, minus confertim punctatus, subtiliter canaliculatus, margine laterali posterius depresso, testaceum, subnitidum. Scutellum iam testaceum, iam nigrum. Elytra thorace duplo prope longiora, apicem versus sensim paulo latiora, subdepressa, densius sat fortiter punctata, flavo-testacea, abdominis nonnisi basin tegentia. Abdomen parce obsoleteque punctulatum, maris nigrum, nitidum, apice piceum, ano acuminato, feminae rufo-testaceum. Pectus parce punctatum, nigrum. Pedes testacei.

Mas femoribus anterioribus leviter incrassatis, anticis medio subtus denticulo armatis, tibiis intermediis incurvis, medio interne emarginatis, sinu superius denticulo terminato, femina pedibus omnibus simplicibus. Praeterea femina thorace brevior et abdomine testaceo distincta.

Obs. Spurium in hoc genere videtur, a reliquis speciebus capite elongato, inter antennarum basin haud foveolato, thorace convexo, margine laterali haud dilatato, tarsis elongatis, corpore pubescente et differentia sexuali discedens, oris autem structura partiumque proportionem non tam mihi differre videtur, ut generi peculiari constituendo characteres satis sint.



**Deliphrum Er.***Er. Col. March. I. p. 627.**Mandibulae muticae.**Maxillae malis membranaceis, exteriore dilatata.**Palpi maxillares articulo ultimo penultimo longiore.**Tibiae spinulosae.**Tarsi postici articulis duobus primis subelongatis, inter se aequalibus.*

Corpus latum, subdepressum, alatum. Caput thorace minus et angustius, obtusiusculum, basi modice constrictum, fronte immarginata, oculis mediocribus, rotundatis, modice prominulis. Ocelli super oculorum marginem superiorem siti. Labrum breve, truncatum, apice membrana brevi biloba auctum. Mandibulae breves, validiusculae, acutae, simplices. Maxillae mala interiore membranacea, extus cornea, denticulo corneo terminata, apice oblique truncata, ciliata, exteriore membranea, apice dilatata, rotundata, barbata. Palpi maxillares articulo primo minuto, secundo sat elongato, tertio brevior, quarto tertio multo longior. Labium mento transversum, antrosum angustatum, apice truncatum, ligula membranea, transversa, medio submarginata. Palpi labiales breviusculi, articulo primo tertioque longioribus, secundo brevior. Antennae filiformes, articulo tertio secundo parum longior, sequentibus longitudine sensim levissime decrescentibus, penultimo crassitie haud brevior, ultimo ovato. Thorax subquadrangulus, angulis rotundatis. Scutellum triangulare. Elytra longiora, angulo exteriore rotundato. Abdomen marginatum, antice parallelum, postice rotundatum, apice subacuminatum. Pedes minus elongati, tibiis subtiliter at distincte spinulosis, tarsis anticis articulis 4 primis brevibus, subcordatis, haud dilatatis, inter se aequalibus, quinto his coniunctis brevior, intermediis articulis duobus primis subelongatis, tertio quartoque brevibus, utrisque inter se aequalibus, quinto his coniunctis brevior, posticis articulis duobus primis leviter elongatis, inter se aequalibus, tertio paulo brevior, quarto brevi, quinto reliquis coniunctis multo brevior.

**Observationes.**

1. Affine generi praecedenti, imprimis oris etiam structura, distinctum praecipue tibiis spinulosis. Habitus quoque corpore laevior, capite thoraceque fere laevibus, statura etiam saepius magis elongata divergens. Habitus propius ad *Omalium* accedit, distinguitur autem tarsorum articulis primis leviter elongatis, antennis gracilioribus et maxillarum mala exteriore haud dilatata.

2. Sexus differentia ut in praecedentibus.

\* 1. *D. tectum*: Nigrum, nitidum, subtiliter punctulatum, antennarum basi, thoracis transversi, obsolete canaliculati margine, elytris pedibusque testaceis. — Long.  $1\frac{1}{2}$  lin.

*Er. Col. March. I. 627. 1.*

*Staph. tectus* Payk. Mon. *Staph.* 68. 48. *Faun. Suec. III.* 411. 56. — *Oliv. Ent. III.* 42. 36. 52. t. 3. f. 21.

*Omal. tectum* Grav. Mon. 213. 19. — *Oliv. Encycl. meth. VIII.* 478. 19. — *Gyll. Ins. Suec. II.* 202. 5. — *Sahlb. Ins. Fenn. I.* 278. 5. — *Zetterst. Faun. Lappon. I.* 49. 4. *Ins. Lappon.* 50. 6. — *Mannerh. Brachelytr.* 52. 11. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 487. 5.

*Omal. laeve* Grav. Mon. 211. 15. — *Oliv. Encycl. meth. Ins. VIII.* 477. 15.

Habitat in Germania, Suecia.

Breviusculum, latum, parum convexum, nigrum, nitidum, capite thoraceque subaeneis. Antennae capite thoraceque longiores, apicem versus sensim leviter incrassatae, nigrae, articulis 3 primis testaceis. Palpi picei. Caput breviusculum, obsolete subtilissime punctulatum, fronte ad oculos utrinque oblique sulcata. Thorax coleopterorum basi paulo angustior, latitudine duplo prope brevior, lateribus rotundatus, antrorsum subangustatus, basi apiceque truncatus, angulis omnibus obtusis, subrotundatis, leviter convexus, crebre subtiliter punctatus, subtiliter obsoleteque canaliculatus, margine laterali et basali testaceo. Elytra thorace plus duplo longiora, ampla, abdominis dimidiam partem tegentia, apicem versus sensim paulo latiora, creberrime minus fortiter punctata, lutea. Abdomen omnium subtilissime punctulatum, segmento ultimo testaceo. Pedes testacei, coxis omnibus concoloribus.

- \* 2. *D. crenatum*: Brunneum, nitidum, antennarum basi pedibusque testaceis, capite thoraceque subtilius punctatis, hoc dorso medio bipunctato, elytris punctato-striatis. — Long.  $2\frac{1}{4}$  lin.

*Omal. crenatum* Grav. Micr. 114. 4. Mon. 207. 6.

Habitat in Germania.

Habitu omnino Omalii brunnei, at sesqui maius, praecipue latius, subdepressum, rufo-brunneum, nitidum, glabrum. Antennae capitis thoracisque fere longitudine, apicem versus vix incrassatae, articulo quarto reliquis duplo minore, 5 — 10 subobconicis; ultimo ovato, fusco-nigrae, articulis 4 primis testaceis. Caput subtiliter vage punctatum, fronte inter oculos striolis duabus obliquis obsoletioribus, et ad antennarum insertionem utrinque foveola minus profunda impressa. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus leviter rotundatis, angulis posterioribus subrectis, depressis, leviter convexus, crebrius subtiliter punctatus, dorso medio punctis duobus maioribus impressus. Scutellum parce subtiliter punctatum. Elytra thorace vix duplo longiora, crebre punctato-striata. Abdomen parce subtilissime punctulatum. Pedes testacei.

- \* 3. *D. arcticum*: Piceo-testaceum, nitidum, thorace transverso, aequali elytrisque crebrius vage punctatis. — Long.  $1\frac{1}{2}$  lin.

Habitat in Lapponia, Dom. Schüppel.

Habitu omnino praecedentis, at plus duplo minus, subdepressum, piceo-testaceum, nitidum, glabrum. Antennae capite thoraceque paulo longiores, apicem versus sensim leniter incrassatae, pilosellae, testaceae. Caput parce subtiliter punctatum, antice laevigatum, inter oculos foveolis binis obsoletioribus, inter antennas binis distinctioribus impressum. Thorax coleopterorum prope latitudine, latitudine duplo prope brevior, lateribus leviter rotundatus, apice leviter emarginatus, basi truncatus, angulis posterioribus subrectis, anterioribus obtusiusculis, leviter convexus, angulis posterioribus depressis, ad marginem lateralem utrinque pone medium leviter impressus, ceterum aequalis, aequaliter crebrius subtiliusque punctatus. Scutellum laeve. Elytra thorace duplo prope longiora, parcius minus subtiliter profunde punctata, punctis nullo modo seriatis, abdominis vix quartam partem obtegantia. Abdomen dilatatum, laeve. Pectus punctatum. Pedes testacei.

- \* 4. *D. algidum*: Oblongum, nigrum, nitidum, antennarum articulo primo pedibusque rufis, thorace transverso, confertim subtiliter punctato, elytris confertissime punctatis. — Long. 2 — 2½ lin.

Habitat in montibus Hercyniae, tempore vernali in nive lectum a Dom. Saxesen.

Statura oblonga, subdepressa Arpedii quadri, nigrum, nitidulum, glabrum. Antennae capite thoraceque paulo longiores, apicem versus haud incrassatae, nigrae, articulo primo obscure rufo. Os inferne piceo-rufum, palpis maxillaribus nigro-piceis. Caput thorace parce subtiliterque punctatum, fronte utrinque prope oculos oblique impressa, antice inter antennis utrinque leviter bifoveolata, margine apicali tumidulo. Thorax coleopteris parum angustior, latitudine dimidio brevior, antrorsum subangustatus, lateribus leviter rotundatis, basi apiceque truncatus, angulis omnibus subrotundatis, subdepressus, confertim subtiliter punctatus, margine laterali ante medium foveolato. Scutellum parce punctulatum. Elytra thorace triplo prope longiora, apicem versus sensim subdilatata, subdepressa, confertissime punctata, punctorum paulo maiorum sat distantium seriebus tribus notata, maris apice recte truncata, abdominis basin vix superantia, feminae apice oblique truncata, paulo longiora, abdominis dimidiam partem tegentia. Abdomen laeve. Pedes rufi.

- \* 5. *D. angustatum*: Oblongum, nigrum, nitidum, antennarum basi pedibusque testaceis, capite thoraceque angustiore quadrato laevibus, elytris confertim subseriatim punctatis, fuscis, limbo apicali testaceo. — Long. 1½ lin.

Habitat in Thuringia, Austria.

Corpus oblongum, subdepressum, nigrum, nitidum. Antennae capite thoraceque paulo longiores, apicem versus sensim subincrassatae, nigrae, basi testaceae. Palpi nigri, basi testacei. Caput oblongum, parcius subtilissime punctatum, fronte inter oculos foveolis duabus minutis profundis notata, inter antennis transversim impressa. Thorax coleopteris plus dimidio angustior, latitudine parum brevior, lateribus vix rotundatis, basin versus subangustatus, basi apiceque truncatus, angulis omnibus obtusis, subrotundatis, leviter convexus, parce omnium subtilissime punctulatus, margine summo laterali piceo. Scutellum impressum. Elytra thorace plus duplo longiora, subdepressa, confertim punctata, punctis hinc inde in series dispositis, fusca, nitida, limbo apicali testaceo. Abdomen laeve, ano acuminato. Pedes testacei, coxis anterioribus concoloribus.

### **Omalium Grav.**

*Er. Col. March. I. p. 628.*

*Anthobium Mannerh. Brachelytr. p. 53.*

*Mandibulae muticae.*

*Maxillae malis membranaceis.*

*Palpi maxillares articulo ultimo penultimo longiore.*

*Tibiae subtilissime spinulosae.*

*Tarsi articulis 4 primis brevissimis, aequalibus.*

Corpus oblongum vel elongatum, subdepressum, alatum. Caput thorace



minus et angustius, basi sat profunde distincteque constrictum, fronte lateribus marginata, oculis mediocribus, rotundatis, prominulis. Ocelli super oculorum marginem superiorem siti. Labrum transversum, apice sinuatum, membrana brevissima ciliata prominula auctum. Mandibulae breves, muticae. Maxillae (t. V. f. 31. b.) malis membranaceis, exteriore oblonga, apice barbatula, interiore extus cornea, apice denticulo terminata, oblique truncata ciliataque. Palpi maxillares articulo primo minuto, secundo modice elongato, tertio brevi, quarto iterum longiore, apice subacuminato. Labium (t. V. f. 31. a.) mento transverso, apicem versus angustato, apice truncato, ligula membranea, biloba, lobis rotundatis. Palpi labiales breviusculi, articulo primo tertioque longioribus, secundo brevior. Antennae apicem versus leviter incrassatae, articulo primo maiore, crassiusculo, secundo tertioque subaequalibus, penultimis transversis, ultimo subovato. Thorax coleopteris saepe paulo angustior, lateribus marginatus. Scutellum triangulare. Elytra abdominis basin obtegentia, angulo apicis exteriore rotundato. Abdomen magis minusve elongatum, lateribus marginatum, apice rotundatum. Pedes breviusculi, tibiis subtilissime et vix perspicue spinulosis, tarsis brevibus, articulis 4 primis brevissimis, inter se aequalibus, haud dilatatis, quinto his coniunctis aequali vel longiore.

## O b s e r v a t i o n e s.

1. Corpus aliis glaberrimum, aliis subtilissime pubescens, plerisque oblongum, nonnunquam lineare, subdepressum. Antennarum articuli penultimi plerisque transversi. Ceterum a praecedentibus hoc genus tarsorum structura sat distinctum.

2. Sexus differentia ut in praecedentibus.

3. Victus aliis in herbis, fruticeis, aliis sub arborum cortice.

4. Nomen Leachianam, quod Com. Mannerheim huic tribuit, sequenti generi restitui.

\* 1. *O. rivulare*: Nigrum, nitidulum, antennarum basi pedibusque testaceis, elytris fusco-testaceis, capite thoraceque crebre punctatis, hoc foveolis duabus oblongis, subarcuatis, profundis. — Long. 2 lin.

Grav. Micr. 116. 8. Mon. 209. 12. — Latr. Hist. nat. Crust. et Ins. IX. 373. 8. t. 80. f. 6. Gen. Crust. et Ins. 1. 298. 1. — Oliv. Encycl. meth. VIII. 477. 12. — Gyll. Ins. Suec. II. 214. 14. — Sahlb. Ins. Fenn. I. 282. 13. — Zetterst. Faun. Lappon. I. 51. 7. Ins. Lappon. 51. 12. — Er. Col. March. I. 629. 1.

Staph. rivularis Payk. Mon. Staph. 65. 46. Faun. Suec. III. 407. 50. — Oliv. Ent. III. 42. 35. 49. t. 3. f. 27. — Panz. Faun. Germ. 27. 15. Ent. Germ. 357. 37. — Marsh. Ent. Britt. 513. 13.

Anthobium rivulare Mannerh. Brachelytr. 53. 1. — Boisd. et Lacord. Faun. Ent. Paris. I. 470. 2. — Runde Brachelytr. Hal. 22. 1.

Omal. incisum Grav. Mon. 209. 12. b.

Staph. cursor Müll. Prodr. 97. 1095. — Goeze Ent. Beitr. I. 727. 22.

Var. Elytris fuscis.

Omal. Oxyacanthae Boisd. et Lacord. I. 471. 4.

Habitat in Europa.

Nigrum, nitidulum. Antennae capite thoraceque parum breviores, apicem versus sensim modice incrassatae, articulis 7 — 10 transversis, nigrae, basi rufae. Caput sat crebre punctatum, apice laevissimum, fronte postice foveolis duabus minoribus, rotundatis, profundis, antice utrinque foveola oblonga, latiore, minus profunda impressa, oris partibus testaceis. Thorax

coleopteris angustior, latitudine plus dimidio brevior, lateribus anterior rotundatus, basin versus leviter angustatus, angulis posterioribus rectis, anterioribus rotundatis, parum convexus, sat crebre profundeque punctatus, dorso foveolis duabus oblongis, subarcuatis, basi latioribus profundioribusque, et ad apicem plerumque foveola tertia intermedia minuta obsoletiore impressus, limbo basali et laterali saepe testaceo. Scutellum laeve, piceum, vel piceo-testaceum. Elytra thorace duplo longiora, dense sat fortiter profundeque punctata, secundum suturam subimpressa, fusco-testacea, vel fusco-picea, nitidula. Abdomen subtilissime punctulatum, nigrum, margine piceo, apice summo testaceo. Pedes testacei.

- \* 2. *O. monilicorne*: Nigrum, nitidulum, antennarum basi pedibusque rufis, capite thoraceque parce subtiliter punctatis, hoc foveolis duabus oblongis obsoletioribus impresso. — Long.  $1\frac{1}{2}$  lin.

Gyll. Ins. Suec. II. 219. 18. — Sahlb. Ins. Fenn. I. 283. 5.

Habitat in Suecia, Dom. Schüppel, in montibus Hercyniae, Dom. Saxesen.

*O. rivulari* paulo minus, magis elongatum, nigrum, nitidum. Antennae capite thoraceque paulo breviores, ab articulo sexto inde abrupte crassiores, articulis 7—10 transversis, nigrae, articulis 5 primis rufis. Caput parvum, punctis raris subtilibus sparsum, fronte postice foveolis duabus minutis, rotundatis, profundis, antice inter antennis utrinque foveola maiore, profundiore impressa, oris partibus rufis. Thorax coleopteris angustior, latitudine dimidio brevior, lateribus antice cum angulis anterioribus rotundatis, basin versus leviter angustatus, angulis posterioribus subrectis, parum convexus, margine laterali posterius depresso, dorso foveolis duabus oblongis parum profundis impressus, sparsim subtiliter punctatus. Scutellum laeve. Elytra thorace duplo longiora, subtilius licet sat profunde confertim punctata, nigropicea, nitida. Abdomen omnium subtilissime parce punctulatum, apice summo rufo-piceo. Pedes rufo-testacei.

- \* 3. *O. exiguum*: Nigrum, nitidulum, pedibus testaceis, elytris fusco-piceis, capite thoraceque dense punctatis, hoc foveolis duabus oblongo-ovatis, profundis impresso. — Long. 1 lin.

Gyll. Ins. Suec. II. 218. 17. — Zetterst. Faun. Lappon. I. 51. 7. Ins. Lapp. 51. 13. *Anthobium exiguum* Mannerh. Brachelytr. 53. 3.

Habitat in Germania.

Habitu fere *O. rivularis*, at quadruplo minus, nigrum, subnitidum. Antennae capitis thoracisque prope longitudine, crassiusculae, ab articulo septimo inde apicem versus sensim leniter incrassatae, articulis penultimis duobus leviter transversis, fusco-nigrae. Caput confertim punctatum, antice laevigatum, fronte utrinque longitudinaliter impressa, impressione utraque postice foveola minuta rotundata sat profunda, antice foveola oblonga profundiore notata, medio longitudinaliter leviter convexa, ore piceo. Thorax coleopteris angustior, latitudine dimidio brevior, lateribus anterioribus cum angulis anterioribus rotundatis, basin versus leviter angustatus, angulis posterioribus subrectis, parum convexus, margine laterali posterius depresso, dorso pone medium foveolis duabus oblongo-ovatis profundis impressus, confertim sat profunde subtilius punctatus. Scutellum laeve. Elytra thorace duplo

longiora, fortius profunde denseque punctata, subrugulosa, secundum suturam ultra medium leviter impressa, nigro-fusca, subnitida. Abdomen omnium subtilissime punctulatum. Pedes testacei.

- \* 4. *O. Oxyacanthae*: Nigrum, nitidulum, pedibus testaceis, elytris fusco-piceis, capite thoraceque crebre punctatis, hoc sulcis duobus subarcuatis impresso. — Long. parum ultra 1 lin.

Grav. Mon. 210. 12. c.

*Anthob. caesum* Boisd. et Lacord. Faun. Ent. Paris. I. 470. 3.

Habitat in Germania, Gallia.

Habitu fere *O. rivularis* at plus triplo minus, *O. exiguo* prope aequale, nigrum, nitidulum. Antennae capitis thoracisque fere longitudine, ab articulo septimo inde apicem versus sensim leniterque incrassatae, nigro-fuscae. Caput crebre punctatum, apice laevigatum, fronte posterius inter oculos et antice inter antennas foveolis binis oblongis profundioribus impressa, ore rufo. Thorax coleopteris paulo angustior, latitudine duplo fere brevior, basin versus parum magis quam apicem versus angustatus, angulis anterioribus obtusis, posterioribus subrectis, parum convexus, margine laterali posterius et angulis posterioribus depressis, dorso impressionibus duabus sublinearibus, extrorsum leviter arcuatis, et antice foveola tertia minore oblonga intermedia notatus, sat crebre subtilius at sat profunde punctatus. Elytra thorace plus duplo longiora, subdepressa, sat fortiter profundeque dense punctata, secundum suturam ultra medium subimpressa, fusca. Abdomen omnium subtilissime punctulatum. Pedes testacei.

- \* 5. *O. fossulatum*: Nigrum, subnitidum, pedibus testaceis, elytris fuscis, capite thoraceque crebre subtiliter punctatis, hoc dorso foveis duabus ovatis profundis impresso, apicem versus angustato. — Long. 1 $\frac{2}{3}$  lin.

Er. Col. March. I. 630. 2.

*Omal. caesum* Gyll. Ins. Suec. II. 215. 15. — Sahlb. Ins. Fenn. I. 283. 11.

*Anthob. caesum* Mannerh. Brachelytr. 53. 2.

Habitat in Germania.

*O. rivulari* paulo minus, praecipue brevius, nigrum. Antennae capitis thoracisque longitudine, apicem versus ab articulo septimo inde sensim leviterque incrassatae, fusco-testaceae, basi nigricantes. Caput subtilius confertissime punctatum, vix nitidum, fronte posterius sulculis longitudinalibus anticeque foveolis oblongis binis profundis impressa. Thorax coleopteris dimidio angustior, latitudine dimidio brevior, lateribus leviter rotundatus, antrorsum, non vero basin versus, angustatus, angulis anterioribus subrotundatis, posterioribus acutiusculis, parum convexus, margine laterali late explanato, pone medium impresso, dorso pone medium foveolis duabus ovatis, antrorsum attenuatis, profundis, approximatis et plerumque ante apicem impressione oblonga obsolete notatus, subtiliter crebre punctatus, subnitidus, marginibus saepe piceis. Scutellum laeve. Elytra thorace duplo longiora, subdepressa, lateribus latius marginata, secundum suturam et infra basin subimpressa, confertim fortius punctata, nigro-picea vel fusco-testacea. Abdomen laeve, nitidum. Pedes testacei.



- \* 6. *O. caesum*: *Nigrum, opacum, pedibus testaceis, elytris fuscis, capite thoraceque confertissime punctatis, hoc dorso foveolis duabus oblongis profundis impresso, basin versus angustato.* — Long.  $1\frac{1}{2}$  lin.

Grav. Mon. 209. 12. a. — Er. Col. March. I. 630. 3.

Omal. *Oxyacanthae* Gyll. Ins. Suec. II. 217. 16.

Anthob. *Oxyacanthae* Mannerh. Brachelytr. 53. 3. — Runde Brachel. Hal. 23. 2.

Habitat in Germania.

Statura et summa affinitas praecedentis. Nigrum. Antennae capitis thoracisque longitudine, apice leviter incrassatae, nigrae, apicem versus fusco-testaceae. Caput confertissime punctatum, subrugulosum, opacum, fronte planiuscula, posterius lineolis duabus longitudinalibus profundis, antice utrinque foveola obsoletiore impressa. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus cum angulis anterioribus rotundatus, basin versus leviter angustatus, angulis posterioribus subrectis, parum convexus, margine laterali explanato, pone medium fortius impresso, dorso foveolis duabus oblongis, sat profundis, tertiaque ante apicem minore, obsoletiore, oblonga, intermedia impressus, dense punctatus, nigro-piceus, parum nitidus, marginibus omnibus piceis vel fusco-testaceis. Scutellum laeve. Elytra thorace duplo longiora, subdepressa, lateribus angustius marginata, secundum suturam et pone basin subimpressa, dense fortius profunde punctata, fusca, parum nitida. Abdomen subtilissime punctulatum, parum nitidum. Pedes testacei.

- \* 7. *O. repandum*: *Nigrum, opacum, antennis pedibusque rufis, thorace elytrisque rufo-piceis, disco nigricantibus, capite thoraceque creberrime punctatis, hoc dorso foveolis duabus oblongis impresso.* — Long.  $1\frac{1}{2}$  lin.

Habitat in America septentrionali, Dom. Zimmermann.

Statura et summa affinitas praecedentium. Antennae capitis thoracisque fere longitudine, apice sensim leviter incrassatae, rufae. Caput subtiliter confertissime punctatum, fronte plana, posterius sulculis duobus longitudinalibus profundis, antice inter antennas utrinque foveola rotundata, minus profunda impressa, supra nigrum, parum nitidum, subtus cum ore rufo-piceum. Thorax coleopteris angustior, latitudine dimidio fere brevior, lateribus leviter rotundatus, margine explanato, pone medium impresso, repando, angulis posterioribus subrectis, apice leviter emarginatus, angulis posterioribus acutiusculis, dorso parum convexus, foveolis duabus minus amplis profundisque, oblongis, antice sulculo obsoleto desinentibus impressus, confertim subtiliter punctatus, parum nitidus, piceo-rufus, disco fuscescens. Scutellum laeve, nigrum. Elytra thorace plus sesqui longiora, subdepressa, lateribus angustius marginata, secundum suturam leviter impressa, confertissime fortius sat profunde punctata, parum nitida, piceo-rufa, dorso nigricantia. Abdomen subtilissime punctulatum, nigrum, vix nitidum, ano rufo-piceo. Pectus nigrum. Pedes rufi.

- \* 8. *O. minimum*: *Depressum, nigrum, opacum, ore pedibusque testaceis, elytris fuscis, punctatis, thorace 4-foveolato, lateribus angulato.* — Long.  $\frac{2}{3}$  lin.

Er. Col. March. I. 631. 5.

Habitat Berolini, Dom. Schüppel.

O. pusillo affine, at paulo minus, lineare, subdepressum, nigrum, opacum. Antennae capite thoraceque multo breviores, articulis duobus primis crassiusculis, secundo subgloboso, dein ab articulo sexto inde iterum crassiores, piceae. Os testaceum. Caput triangulare, supra impunctatum, utrinque longitudinaliter impressum, fronte lateribus et apice subtiliter marginata. Thorax coleopteris paulo angustior, latitudine plus dimidio brevior, lateribus rotundatis, medio angulatis, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusis, foveis quatuor oblongis magnis profundisque, exterioribus ad marginem lateralem sitis, omnibus a basi ultra medium adscendentibus, impressus. Elytra thorace duplo prope longiora, depressa, confertim distincte punctata, fusca. Abdomen subtilissime punctulatum. Pedes testacei.

\* 9. *O. pusillum*: *Depressum nigrum, opacum, laeve, ore, antennis pedibusque testaceis, thorace dorso bifoveolato.* — Long. 1 lin.

Grav. Mon. 205. 5. — Oliv. Encycl. meth. VIII. 475. 3. — Gyll. Ins. Suec. II. 220. 19. — Sahlb. Ins. Fenn. I. 284. 16. — Er. Col. March. I. 631. 4.

Anthob. pusillum Mannerh. Brachelytr. 53. 5.

Habitat in Europa sub cortice Pinus.

Lineare, depressum, impunctatum, opacum, subsericeo-micans, nigrum, thorace elytrisque saepe piceis vel piceo-testaceis. Antennae capite thoraceque paulo breviores, apice incrassatae, articulis 6 — 10 transversis, rufo-testaceae, articulis penultimis 4 fuscis. Os rufo-testaceum. Caput fronte planiuscula, postice utrinque ante ocellum puncto minuto, apice foveis duabus maioribus impressa. Thorax coleopteris angustior, latitudine duplo fere brevior, lateribus anterioribus modice rotundatus, basin versus leviter angustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus rectis, subdepressus, dorso pone medium foveis duabus oblongis sat profundis, in medio marginis lateralis foveola rotundata impressus. Elytra depressa, thorace duplo fere longiora. Abdomen elongatum, lineare, segmentis margine postico testaceo. Pedes testacei.

\* 10. *O. florale*: *Nigrum, nitidum, pedibus rufis, thorace obsolete bifoveolato, elytris subseriatim punctatis.* — Long. 2 lin.

Er. Col. March. I. 633. 7.

Staph. floralis Payk. Mon. Staph. 67. 47. Faun. Suec. III. 406. 49. — Fab. Ent. Syst. I. II. 530. 52. — Walk. Faun. Paris. I. 272. 16.

Stenus floralis Fab. Syst. El. II. 604. 6.

Omal. Viburni Grav. Micr. 117. 9. Mon. 210. 13. — Latr. Hist. nat. Crust. et Ins. IX. 373. 9. — Oliv. Encycl. meth. VIII. 477. 13. — Gyll. Ins. Suec. II. 224. 23. — Sahlb. Ins. Fenn. I. 285. 21.

Anthob. Viburni Mannerh. Brachelytr. 54. 7. — Boisd. et Lacord. Faun. Ent. Paris. I. 472. 5. — Runde Brachelytr. Hal. 23. 4.

Staph. rufipes Fourcr. Ent. Paris. I. 168. 15.

Le Staph. noir à corcelet lisse et bordé Geoffr. Hist. Ins. I. 367. 15.

Habitat in Europa.

Statura et magnitudo omnino O. rivularis, nigrum, nitidum, subtilissime

pubescens. Antennae capite thoraceque breviores, crassiusculae, apicem versus ab articulo quinto inde incrassatae, nigrae, apice magis minusve ferrugineo-testaceae. Palpi picei. Caput posterius parcius punctatum, anterius laevigatum, fronte posterius lineolis duabus minutis, antice utrinque foveola rotundata impressa. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus modice rotundatus, tenuiter marginatus, basin apicemque versus aequaliter angustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, leviter convexus, crebre punctatus, dorso foveolis duabus oblongis, valde obsoletis impressus. Scutellum laeve. Elytra thorace vix duplo longiora, subdepressa, confertim subruguloso-punctata, punctis praecipue versus suturam subseriatis. Abdomen subtilissime punctulatum. Pedes rufi, femoribus interdum picescentibus.

Variat thoracis margine elytrisque picei, antennisque totis rufis.

\* 11. *O. nigrum*: *Nigrum, nitidum, antennarum basi pedibusque rufis, thorace margine piceo, dorso obsoletius bifoveolato, elytris subseriatim punctatis.* — Long.  $1\frac{1}{2}$  lin.

Grav. Mon. 212. 17. — Oliv. Encycl. meth. VIII. 478. 17.

*Omalium Salicinum* Gyll. Ins. Suec. IV. 467. 24 — 25.

*Anthob. Salicinum* Mannerh. Brachelytr. 54. 12.

Habitat in Suecia, Dom. Schüppel, in Bavaria, Dom. Dr. Walzl.

Praecedenti affine, at paulo minus, nigrum, nitidum, subtilissime pubescens. Antennae capitis thoracisque fere longitudine, ab articulo quinto inde leniter incrassatae, nigrae, articulis 5 primis rufis. Palpi rufi. Caput parcius punctatum, fronte inter oculos striolis duabus minutis, distantibus, inter antennis utrinque foveola maiore, sat profunda, rotundata impressa. Thorax coleopteris paulo angustior, latitudine plus dimidio brevior, lateribus modice rotundatis, latius marginatis, apicem versus paulo magis quam basin versus angustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, dorso modice convexus, sat crebre punctatus, foveolis duabus oblongis obsoletioribus impressus, margine praecipue laterali piceo-rufo. Scutellum parce subtiliter punctatum. Elytra thorace duplo longiora, subdepressa, confertim subruguloso-punctata, punctis passim subseriatis, nigro-picea. Abdomen subtiliter punctulatum. Pedes rufo-testacei.

\* 12. *O. Salicis*: *Nigrum, nitidum, antennis pedibusque rufis, thorace fortius punctato, dorso bifoveolato, elytris fortius subseriato-punctatis.* — Long. 2 lin.

Gyll. Ins. Suec. II. 226. 24. — Sahlb. Ins. Fenn. I. 286. 22.

*Anthob. Salicis* Mannerh. Brachelytr. 54. 11.

Habitat in Suecia, Dom. Schüppel.

Statura omnino *O. brunnei*, lineare, subdepressum, nigrum, nitidum, glabrum. Antennae capite thoraceque paulo breviores, ab articulo septimo inde leviter incrassatae, rufae. Palpi rufi. Caput parce punctatum, apice laevigatum, fronte inter oculos foveolis duabus minutis, punctiformibus, inter antennis foveis duabus maioribus rotundatis impressa, ore rufo-piceo. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus leviter rotundatus, basin apicemque versus aequaliter angustatus, basi apiceque trun-



catus, angulis anterioribus obtusis, posterioribus subrectis, leviter convexus, crebre fortius profundiusque punctatus, dorso foveis duabus oblongis approximatis leviter impressis, margine laterali praecipue posterius explanato, rufo-piceo. Scutellum parce subtiliter punctatum. Elytra thorace duplo prope longiora, subdepressa, fortius profundiusque dense punctata, punctis versus saturam confertioribus subseriatis, saepe confluentibus. Abdomen parcius subtiliter punctatum. Pedes rufo-testacei.

- \* 13. *O. brunneum*: *Rufo-testaceum, nitidum, capite, ano elytrorumque macula apicali nigris, capite thoraceque crebre punctatis, elytris crebre punctato-striatis.* — Long. 2 lin.

*Oliv. Encycl. meth. VIII.* 475. 5. — *Gyll. Ins. Suec. II.* 227. 25. — *Sahlb. Ins. Fenn. I.* 287. 24. — *Er. Col. March. I.* 633. 8.

*Staph. brunneus Payk. Mon. Staph.* 63. 45. *Faun. Suec. III.* 404. 47.

*Omal. brunneum var. 2 Grav. Micr.* 114. 2.

*Anthob. brunneum Mannerh. Brachelytr.* 54. 13. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 475. 7.

*Staph. melanocephalus Fab. Mant. Ins. I.* 222. 32. *Ent. Syst. I. II.* 529. 29. *Syst. El. II.* 601. 63. — *Panz. Ent. Germ.* 357. 38.

*Staph. melanocephalus Cederh. Faun. Ingr.* 119. 365. (*forte*)

Habitat in Europa, sub arborum cortice.

Rufo-testaceum, nitidum, glabrum. Antennae capitis thoracisque longitudine, apicem versus leniter incrassatae, articulis ultimis 5 crassioribus, ultimo ovato, rufo-ferrugineae. Caput crebrius vage punctatum, fronte posterius ante ocellos punctis duobus profundis, minutis, antice inter antennis utrinque fovea latiore impressa, nigro-piceum, ore rufo. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, modice convexus, crebre sat profunde punctatus, dorso obsolete bi-impressus, margine laterali depresso, angusto, posterius latiore. Scutellum parce punctatum. Elytra thorace vix duplo longiora, crebre striato-punctata, macula apicali rotundata nigra. Abdomen supra parcius subtiliter punctulatum, infra dense fortius punctatum, segmentis ultimis duobus nigro-piceis. Pedes rufo-testacei.

- \* 14. *O. lucidum*: *Rufum, nitidum, capite, pectore, elytrorum sutura apiceque abdominisque apice nigris, capite thoraceque crebre punctatis, elytris punctato-striatis.* — Long.  $1\frac{1}{4}$  lin.

*Er. Col. March. I.* 634. 9.

*Staph. brunneus var. β Payk. Mon. Staph.* 64. 45. *Faun. Suec. III.* 405. 47.

*Omal. brunneum var. b Gyll. Ins. Suec. II.* 227. 25. — *Sahlb. Ins. Fenn. I.* 287. 24.

*Omal. brunneum var. 1 Grav. Mon.* 206. 5. — *Latr. Hist. nat. Crust. et Ins. IX.* 371. 3.

Habitat in Germania, Gallia, Suecia, sub cortice Pinus.

Habitu *O. brunnei*, at plus triplo minus, subtilissime pubescens. Antennae capite thoraceque paulo breviores, apice crassiores, articulis 6—10 transversis, ultimo globoso, rufo-ferrugineae. Caput creberrime fortiter punctatum, fronte convexiuscula, antice bifoveolata, nigro-piceum, nitidum, ore rufo. Thorax coleopteris paulo angustior, latitudine dimidio fere bre-

vior, lateribus modice rotundatus, basi apiceque truncatus, angulis anterioribus obtusis, subdeflexis, posterioribus obtusiusculis, subdepressis, modice convexus, sat fortiter creberrime punctatus, dorso pone medium obsolete bifoveolatus, rufus, nitidus. Scutellum parce punctatum, piceum, nitidum. Elytra thorace vix duplo longiora, sat fortiter crebreque punctato-striata, rufo-testacea, sutura apiceque late nigris. Abdomen crebre subtilius punctulatum, rufo-testaceum, dorso fere toto ab apice inde nigro-piceo. Pectus nigro-piceum. Pedes testacei.

\* 15. *O. vile: Elongatum, nigricans, subtiliter pubescens, antennarum basi pedibusque testaceis, thorace leviter bifoveolato, elytris crebre profundius punctatis.* — Long. vix ultra 1 lin.

Habitat in Saxonia montana, Dom. Märkel.

Statura praecedentis, at paululum minus, elongatum, subdepressum, piceo-nigrum, nitidum, pube brevissima subtili aequaliter adpersum. Antennae capite thoraceque breviores, articulis duobus primis crassiusculis, ab articulo sexto iterum incrassatae, articulis 6 — 10 fortiter transversis, nigricantes, articulis 6 primis rufo-testaceis. Caput crebrius punctatum, anterius laevius, fronte antice utrinque impressa. Thorax coleopteris paulo angustior, latitudine parum brevior, lateribus leviter rotundatus, basin versus paulo magis quam anterius angustatus, basi apiceque truncatus, angulis posterioribus subrectis, parum convexus, dense punctatus, dorso obsolete longitudinaliter bi-impressus. Elytra thorace duplo prope longiora, subdepressa, dense minus subtiliter punctata, punctis nonnunquam seriatis. Abdomen parce omnium subtilissime obsolete punctulatum. Pedes testacei.

Obs. Thoracis impressiones nonnunquam obsoletae.

\* 16. *O. striatum: Nigrum, pedibus testaceis, thorace obsolete canaliculato bifoveolatoque, elytris punctato-striatis.* — Long. 1 lin.

Grav. Micr. 119. 12. Mon. 216. 23. — Latr. Hist. nat. Crust. et Ins. IX. 374. 12. — Oliv. Encycl. meth. VIII. 479. 23. — Gyll. Ins. Suec. II. 231. 28. — Er. Col. March. I. 632. 6.

Anthob. striatum Mannerh. Brachelytr. 55. 15. — Boisd. et Lacord. Faun. Ent. Paris. I. 475. 11.

Staph. minutus Oliv. Ent. III. 42. 38. 56. t. 6. f. 53.

Habitat in Europa.

Minutum, nigrum, subnitidum. Antennae capitis thoracisque fere longitudine, ab articulo septimo inde leniter incrassatae, nigrae, articulo primo plerumque testaceo, sequentibus saepe piceis. Caput parcius subtiliter punctatum, fronte convexiuscula, posterius utrinque obsolete impressa, antice profundius bifoveolata. Thorax coleopteris angustior, latitudine duplo fere brevior, antrorsum subangustatus, lateribus parum rotundatus, apice leviter emarginatus, basi truncatus, angulis posterioribus subrectis, parum convexus, utrinque ad antennarum marginem lateralem obsolete impressus, dorso longitudinaliter obsolete bi-impressus, obsolete canaliculatus, crebre punctatus. Scutellum laeve. Elytra ampla, thorace plus duplo longiora, apicem versus sensim dilatata, confertim subtilius punctato-striata, iam nigra, iam picea. Abdomen laeve, apice piceum, segmento apicali nonnunquam testaceo. Pedes testacei.

- \* 17. *O. pygmaeum*: Oblongo-ovatum, leviter convexum, rufo-testaceum, pubescens, capite, thorace elytrisque crebre subtiliter punctatis. — Long.  $1\frac{1}{4}$  lin.

Gyll. Ins. Suec. II. 223. 22. — Mannerh. Brachelytr. 52. 7. — Boisd. et Lacord. Faun. Ent. Paris. I. 488. 6.

Staph. pygmaeus Payk. Faun. Suec. III. 410. 55.

Habitat in Germania, Knoch, in Suecia, Dom. Schüppel.

Oblongo-ovatum, leviter convexum, rufo-testaceum, nitidum, minus subtiliter griseo-pubescens. Antennae capitis thoracisque fere longitudine, pilosellae, articulis ultimis 5 sensim crassioribus, 7 — 10 transversis, ultimo subovato. Caput sat crebre punctatum, fronte antice inter antennas utrinque longitudinaliter impressa, margine laterali leviter elevato. Thorax coleopteris paulo angustior, latitudine duplo fere brevior, lateribus leviter rotundatis, medio subangulatis, apicem versus paulo magis quam basin versus angustatus, basi apiceque truncatus, angulis anterioribus obtusis, deflexis, posterioribus obtusis, modice convexus, dense punctatus, aequalis, angulis posterioribus late depressis. Elytra thorace duplo fere longiora, leviter convexa, dense punctata. Abdomen omnium subtilissime parce punctulatum.

Mas trochanteribus posticis dilatatis, securiformibus et abdominis segmento quinto inferiore apice medio reflexo, acute bidentato insignis.

- \* 18. *O. inflatum*: Ovatum, convexum, piceo-rufum, nitidum, subglabrum, capite, thorace elytrisque crebre profundeque punctatis. — Long. 1 —  $1\frac{1}{4}$  lin.

Gyll. Ins. Suec. III. 700. 22 — 23. — Sahlb. Ins. Fenn. I. 285. 20. — Zetterst. Faun. Lappon. I. 53. 12. Ins. Lappon. 52. 19. — Mannerh. Brachelytr. 51. 6.

Habitat in Suecia, Dom. Schüppel, in Fennia, Dom. Sahlberg, in Germania Dom. Ratzeburg.

*O. pygmaeo* fere aequale, at brevius et magis convexum, piceo-testaceum, nitidum, vix pubescens. Antennae capitis thoracisque longitudine, longius pilosellae, articulis 7 — 10 abrupte maioribus, subperfoliatis, ultimo ovato. Caput sat crebre punctatum, apice laeve, fronte antice inter antenas obsolete bifoveolata. Thorax coleopteris paulo angustior, latitudine vix duplo brevior, lateribus leviter rotundatus, margine reflexo, obsolete crenulato, basi truncatus, angulis posterioribus rectis, antrorsum subangustatus, apice leviter emarginatus, angulis anterioribus acutiusculis, convexus, creberrime sat profunde punctatus. Scutellum laeve. Elytra thorace duplo longiora, lateribus leviter rotundata, convexa, sat fortiter profundeque punctata. Abdomen omnium subtilissime parce punctulatum, apice testaceum. Pedes testacei.

- \* 19. *O. rufulum*: Oblongum, rufo-testaceum, parce pubescens, elytris fortiter subseriatim, capite thoraceque subtiliter punctatis, hoc basin versus angustato. — Long.  $1\frac{1}{4}$  lin.

Habitat in Austria, Dom. Ullrich.

*O. pygmaeo* fere aequale, at minus ovatum, subdepressum, rufo-testaceum, nitidum, minus subtiliter parce pubescens. Antennae capite thoraceque paulo longiores, crassae, articulo tertio quartoque tenuiores, apicem ver-



sus sensim leviterque subincrassatae, pilosellae. Caput parce subtilissime punctulatum, fronte antice obsolete bifoveolata. Thorax coleopteris dimidio angustior, latitudine dimidio brevior, lateribus anterieus modice rotundatus, basin versus leviter angustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus rectis, leviter convexus, aequalis, crebrius subtiliter punctatus, angulis posterioribus depressis. Scutellum laeve. Elytra thorace vix duplo longiora, minus confertim at sat profunde subseriato-punctata. Abdomen laeve, apicem versus piceum, ano testaceo. Pedes testacei.

20. *O. brevicorne: Elongatum, subdepressum, subtiliter pubescens, nigrum, nitidulum, antennarum basi pedibusque rufo-testaceis, thorace leviter bi-impresso, elytris thorace duplo longioribus, confertim punctatis, subrugulosis. — Long. 1 lin.*

Habitat in Sardinia, Mus. Reg. Taurin.

Habitu omnino *O. monilicornis*, at plus duplo minus, nigrum, nitidulum, pube brevissima subtilissimaque grisea tenuiter vestitum. Antennae capite paulo longiores, dense pubescentes, articulis 5 primis rufis, sequentibus crassioribus, 6 — 10 transversis, ultimo subgloboso, apice obtuse acuminato, nigris. Palpi rufo-testacei. Caput thorace paulo angustius, subtiliter punctulatum, fronte inter oculos utrinque obsolete sulcata, antice inter antennas distinctius bi-impressa. Thorax coleopteris paulo angustior, latitudine paulo brevior, lateribus ante medium modice rotundatus, basin versus paulo magis quam antrorsum angustatus, basi apiceque truncatus, angulis anterioribus obtusis, posterioribus subrectis, leviter convexus, dense minus subtiliter profunde punctatus, dorso pone medium foveolis duabus oblongis longitudinalibus approximatis leviter impressus, angulis posterioribus depressis, subtus piceus. Elytra thorace duplo longiora, apice truncata, minus subtiliter confertim punctata, longitudinaliter obsolete strigosa. Abdomen segmento primo solo obtecto, lineare, parce subtilissime punctatum, glabrum. Pedes rufo-testacei.

- \* 21. *O. lineare: Elongatum, subdepressum, glabrum, nitidum, rufum, capite thoraceque crebre punctatis, hoc dorso foveolis duabus oblongis impresso, elytris crebre subseriatim punctatis. — Long. 1½ lin.*

Zetterst. Faun. Lapon. I. 53. 13. Ins. Lapp. 53. 21. — Er. Col. March. I. 636. 13.

Habitat in Lapponia (Zetterst.), in Germania, Dom. Schüppel.

Statura fere *O. deplanati*, at angustius et magis depressum, lineare, glabrum, nitidum, rufum. Antennae capite thoraceque multo breviores, crassiusculae, apicem versus incrassatae, articulis 6 — 10 fortiter transversis, perfoliatis, pilosellae. Caput subtilius sat profunde creberrime punctatum, fronte antice inter antennas foveolis duabus rotundatis profundius impressa. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus rotundatus, basin versus leviter angustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posticis subrectis, dorso subdepressus, leviter longitudinaliter bi-impressus, creberrime punctatus. Scutellum punctis nonnullis subtilibus impressum. Elytra thorace duplo fere longiora, subdepressa, sat profunde crebreque punctata, punctis subseriatis, praecipue suturam versus. Abdomen subtiliter punctatum, apicem versus rufo-piceum. Pedes testacei.

- \*22. *O. testaceum*: *Elongatum, subdepressum, subtiliter pubescens, rufo-testaceum, nitidum, capite, thorace elytrisque crebre punctatis, thorace aequali.* — Long.  $1\frac{1}{2}$  lin.

*Omal. pygmaeum* Grav. Mon. 206. 4.

*Omal. brunneum* var. 3 Grav. Micr. 114. 3.

*Omal. deplanatum* var. b Er. Col. March. I. 635. 10.

Habitat in Germania.

*Elongatum, subdepressum, rufo-testaceum, nitidum, parcius subtiliter pubescens.* Antennae capitis thoracisque prope longitudine, apicem versus leviter incrassatae, ab articulo sexto inde distincte paulo crassiores, articulis penultimis autem vix transversis. Caput thorace angustius, subtriangulare, confertim minus subtiliter punctatum, antice bifoveolatum. Thorax coleopteris parum angustior, latitudine plus dimidio brevior, lateribus rotundatus. basin versus leviter angustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, leviter convexus, aequalis, dense minus subtiliter punctatus. Scutellum punctatum. Elytra thorace plus sesqui longiora, dense minus subtiliter punctata. Abdomen parce omnium subtilissime punctulatum, apicem versus subdilatatum. Pedes dilutius testacei.

Obs. *O. deplanato* proximum, at parcius pubescens, fortius punctatum, antennis longioribus, abdomine sublaevi etc. satis distinctum. Ab *O. pygmaeo* corpore lineari, subdepresso, thorace basin versus angustato etc. differt.

- \*23. *O. deplanatum*: *Elongatum, subdepressum, subtiliter pubescens, piceo-nigrum, nitidum, antennis, pedibus elytrisque rufis, capite, thorace elytrisque confertissime subtiliter punctatis.* — Long.  $1\frac{1}{2}$  lin.

Gyll. Ins. Suec. II. 222. 21. — Sahlb. Ins. Fenn. I. 284. 18. — Er. Col. March. I. 635. 10.

*Anthob. deplanatum* Mannerh. Brachelytr. 54. 14.

*Omal. depressum* Grav. Micr. 113. 2. Mon. 205. 2. — Latr. Hist. nat. Crust. et Ins. IX. 371. 2. — Oliv. Encycl. meth. VII. 475. 2.

*Anthob. depressum* Boisd. et Lacord. Faun. Ent. Paris. I. 474. 9.

*Var. Thorace dorso obsolete bifoveolato.*

*Anthob. oblongum* Boisd. et Lacord. Faun. Ent. Paris. I. 473. 8.

Habitat in Europa.

Corpus lineare, subdepressum, densius subtiliter pubescens, nigro-piceum, subnitidum. Antennae capite thoraceque breviores, crassiusculae, apicem versus vix incrassatae, articulis 6 — 10 transversis, fere perfoliatis, totae rufae, vel rufo-testaceae. Caput thorace angustius, depressum, creberrime punctatum, fronte antice inter antennas foveolis duabus minutis impressa, ore rufo-testaceo. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus anterieus rotundatus, basin versus modice angustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, leviter convexus, confertim subtiliter punctatus, dorso foveolis duabus oblongis valde obsolete impresso, piceo-niger, margine laterali basali-que rufo-testaceo. Scutellum subtiliter punctulatum, nigro-piceum. Elytra thorace plus sesqui longiora, subdepressa, confertim subtiliter punctata, rufo-testacea. Abdomen subtilissime punctulatum, piceum, nonnunquam rufo-piceum vel rufum, segmento quinto nigro. Pedes rufi.

- \* 24. *O. concinnum*: *Elongatum, subdepressum, parce omnium subtilissime pubescens, nigrum, nitidum, antennis pedibusque rufis, capite thoraceque subtilius crebre punctatis, elytris punctato-subrugulosis, rufis piceisve.* — *Long.*  $1\frac{1}{2}$  *lin.*

*Var. thorace dorso obsolete bifoveolato.*

*Staph. concinnum Marsh. Ent. Britt. 510. 34.*

Habitat in Germania.

Statura et summa affinitas praecedentis. Lineare, subdepressum, nigrum, nitidum, capite, thorace elytrisque parce omnium subtilissime pubescentibus vel subglabris, abdomine pube subtili depressa distinctiore vestito. Antennae capite paulo longiores, crassiusculae, apicem versus haud incrassatae, articulis 4 — 10 transversis, piceo-rufae. Os piceum. Caput thoracis prope latitudine, suborbiculatum, depressiusculum, crebre subtilius punctatum, antice obsolete bifoveolatum. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus leviter rotundatus, basin versus subangustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, parum convexus, crebre subtilius punctatus, dorso iam omnino aequali, iam foveis duabus oblongis magis minusve obsolete impresso, limbo laterali plerumque rufo-piceo. Elytra thorace duplo prope longiora, crebre punctata, longitudinaliter substrigosa, iam nigro- iam rufo-picea. Abdomen subtiliter punctatum. Pedes rufo-testacei.

Obs. Praecedenti proximum et simillimum, punctura autem capitis thoracisque minus subtili et minus densa, elytris longitudinaliter substrigosis, nec non capite, thorace elytrisque subglabris distinctum videtur.

- \* 25. *O. planum*: *Elongatum, subdepressum, nigrum, nitidum, antennarum basi pedibusque testaceis, elytris fusco-testaceis, thorace obsolete canaliculato.* — *Long.* 1 —  $1\frac{1}{3}$  *lin.*

*Grav. Micr. 112. 1. Mon. 204. 1. — Latr. Hist. nat. Crust. et Ins. IX. 370. 1. — Oliv. Encycl. meth. VIII. 475. 1. — Gyll. Ins. Suec. II. 221. 20. — Sahlb. Ins. Fenn. I. 284. 17. — Zetterst. Faun. Lappon. I. 53. 11. Ins. Lappon. 52. 18. — Er. Col. March. I. 636. 12.*

*Staph. planum Payk. Mon. Curc. App. 145. 11 — 12. Faun. Suec. III. 405. 48.*

*Anthob. planum Mannerh. Brachelytr. 54. 6. — Boisd. et Lacord. Faun. Ent. Paris. I. 475. 10.*

*Staph. flavipes Fab. Ent. Syst. I. II. 527. 39. Fab. Syst. El. II. 599. 48. — Panz. Ent. Germ. 356. 29.*

*Staph. flavipes Linn. Faun. Suec. n. 856. Syst. Nat. I. II. 685. 22. — Müll. Faun. Frid. 23. 226. Prodr. 97. 1099. — Goeze Ent. Beitr. I. 721. 22. — De Vill. Ent. 418. 22. (forte)*

Habitat in Europa.

Elongatum, lineare, subdepressum, nigrum, nitidum, glabrum. Antennae capite thoraceque paulo breviores, rufo-testaceae, articulis 6 — 11 abrupte crassioribus, fuscis. Os rufo-testaceum. Caput thorace paulo angustius, parce subtiliterque punctatum, fronte plana, posterius inter oculos et antice inter antennas foveolis binis impressa, oculis sat prominulis. Thorax coleopteris vix angustior, latitudine plus dimidio brevior, lateribus sat fortiter rotundatus, basin versus coarctatus, basi apiceque truncatus, angulis anterioribus subrotundatis, posterioribus acutiusculis, prominulis, subdepressus, sat



crebre subtiliter punctatus, obsolete canaliculatus, dorso nunquam utrinque obsolete impressus, apice utrinque foveola rotundata, in margine ipso notatus. Scutellum laeve. Elytra thorace duplo longiora, depressa, sat crebre subtiliter punctata, piceo-testacea vel fusco-picea. Abdomen subtilissime punctulatum. Pedes rufo-testacei.

- \* 26. *O. philanthus*: *Subdepressum, nigrum, ore, antennis pedibusque testaceis, elytris flavis, limbo apicali nigro.* — Long.  $1\frac{1}{2}$  lin.

*Staph. philanthus* Fab. *Ent. Syst. Suppl.* 181. 46. *Syst. El.* II. 600. 58.

Habitat ad promont. Bonae Spei, Dom. Bergius.

Oblongum, depressum, nigrum, nitidum. Antennae capite thoraceque breviores, articulis ultimis 6 abrupte crassioribus, 6 — 10 transversis, ultimo subovato, laete testaceae. Os testaceum. Caput subtilissime parcius punctatum, fronte utrinque sulco longitudinali flexuoso, parum profundo impressa. Thorax coleopteris parum angustior, latitudine dimidio brevior, lateribus leviter rotundatis, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusiusculis, subdepressus, sat crebre subtiliter punctatus, medio obsolete canaliculatus. Scutellum laeve. Elytra thorace plus sesqui longiora, apice coniunctim leviter rotundata, depressa, confertissime subtiliter punctata, flava, margine laterali fusciscente, limbo apicali lato nigro. Abdomen subtiliter punctulatum. Pedes testacei.

- \* 27. *O. pedicularium*: *Oblongum, subdepressum, testaceum, pectore, scutello, elytris apice, abdomine ante apicem picescentibus, thorace foveolis duabus oblongis obsoletis impresso.* — Long. 1 lin.

Habitat in ins. Puerto-Rico, Dom. Moritz.

*O. pusillo* haud longius, at sesqui latius, depressum, testaceum, subnitidum. Antennae capite thoraceque paulo breviores, apice fortius incrassatae, articulis 7 — 10 transversis, ultimo subgloboso, testaceae, apicem versus fusciscentes. Caput subtilissime punctulatum, fronte posterioribus binis impressis, rotundatis sat profundis, antice inter antennis maioribus binis impressis, rufo-testaceum. Thorax coleopteris parum angustior, latitudine duplo brevior, lateribus leviter rotundatis, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusiusculis, subdepressus, subtilissime punctulatus, dorso foveolis duabus oblongis, distantibus, obsoletioribus impressus, testaceus. Scutellum piceum. Elytra thorace duplo longiora, subtiliter minus confertim punctata, flavo-testacea, apice subpicescentia. Abdomen parce subtilissime punctulatum, testaceum, ante apicem subpicescens. Pectus piceum. Pedes flavo-testacei.

Species mihi invisae, forte huius generis.

28. *O. laeviusculum*: *Depressum, nigrum, nitidum, antennarum basi, thoracis marginibus pedibusque testaceis, thorace transverso, foveolato, vix punctulato, elytris longioribus fuscis, subtiliter punctulatis.*

*Gyll. Ins. Suec. IV.* 464. 13—14. — *Sahlb. Ins. Fenn.* I. 282. 12. — *Zetterst. Ins. Lappon.* 51. 11.

*Var. b*, thorace toto rufescente, corpore subtus rufo-piceo.

*Var. c*, ut *var. b*, sed abdomen etiam supra rufo-testaceum.

Habitat in Suecia.

*O. rivulari* simile et affine, sed paulo maius, latius, thorax vix et elytra subtilius punctulata etc. Caput maiusculum, porrectum, subtriangulare, depressum, nigrum, nitidum, obsolete punctulatum, foveis duabus verticis profundis, brevibus, aliisque duabus anterioribus oblongis impressum; os testaceum, palpis apice fuscis; oculi magni, laterales, globosi, nigri. Antennae capite cum thorace longiores, pubescentes et pilosae, articulo primo magno, clavato, testaceo, secundo breviori obconico, eiusdem coloris, tertio aequae longo, obconico, apice fusco, 4, 5 brevioribus, nodosis, 6 — 10 sensim multo maioribus, cyathiformibus, nigris, ultimo adhuc maiore, ovato, acuminato, concolore. Thorax transversus, basi apiceque truncatus, lateribus ante medium parum ampliatus, posterius nonnihil angustior, angulis rectis, supra parum convexus, inaequalis, foveis nempe duabus mediis oblongis, at obsolete impressus et intra margines, posterius latius, depressus, obsolete punctulatus, aut niger, nitidus, lateribus posterius testaceis, ut in a, aut totus rufo-testaceus, ut in b et c. Scutellum breve, triangulare piceum, subtilissime punctulatum. Elytra antice thoracis basi latiora, humeris rotundatis, thorace duplo longiora, apice truncata, angulo exteriori rotundato, supra depressa, confertim subtiliter vage punctulata, aut saturatius picea, aut dilutius fusco-testacea, basi apiceque semper dilutiora. Abdomen pone elytra vix illorum longitudine, subdepressum, laeve, nitidum, marginibus elevatis, apice rotundatum, in a, b nigrum, in var. c rufo-testaceum. Corpus subtus in a et b nigrum, nitidum, laeve, in var. c rufo-piceum. Pedes medioeres, toti rufo-testacei. (Gyll.)

29. *O. Fahraei*: Depressum, glabriusculum, nitidum, nigrum, elytris piceis, pedibus testaceis, thorace subquadrato, lateribus subrotundato, posterius paulo angustato elytrisque elongatis, crebre punctatis. — Long. 1—1½ lin.

Zetterst. Ins. Lappon. 52. 17.

Habitat in Lapponia, Norvagia.

Affine *O. pubescenti*, sed glabritie corporis, antennarum forma etc. distinctum. Ab *Arped. brachyptero* (*O. Gyllenhali*), cui etiam subsimile videtur, antennarum colore etc. mox differt. Antennae fere thorace breviores, extrorsum distincte crassiores, totae nigricantes. Os et palpi nigra. Thorax subconvexus, punctulatus, sed vix foveolis impressus, structura ut in diagnosi. Elytra thorace duplo longiora, picea, crebre et certo situ subseriatim punctulata. Pedes flavo-testacei. (Zett.)

30. *O. fenestrale*: Elongatum, subtiliter crebre punctatum, pubescens, piceo-nigrum, annis elongatis, fuscis, pedibus fusco-testaceis, thorace rotundato, obsolete canaliculato. — Long. 1¼ lin.

Zetterst. Faun. Lappon. I. 50. 6. Ins. Lappon. 50. 10.

Habitat in Lapponia Tornensi.

Magnitudo *O. rivularis*. Caput triangulare, subtiliter parce punctulatum, fronte inter antennas bifeveolata, nitidum, nigrum, ore piceo. Antennae thorace longiores, filiformes, pubescentes, nigro-fuscae. Thorax subtransversus, rotundatus, undique marginatus, supra subconvexus, in medio longitudinaliter leviter canaliculatus, totus pubescens, nitidus, niger. Scutellum

magnum, triangulare, punctatum. Elytra thorace plus quam duplo longiora, creberrime et subtiliter, attamen profundius, quam ille, punctata, pubescentia, nigro-picea. Alae amplae, albo-hyalinae. Corpus nitidum, nigrum, ano piceo. Pedes obscure testacei. (Zetterst.)

31. *O. pubescens*: *Brevius, subtiliter crebre punctatum, pubescens, piceo-nigrum, antennis mediocribus fuscis, thorace rotundato, non canaliculato.* — *Long. circiter 1 lin.*

*Zetterst. Faun. Lappon. I. 52. 10. Ins. Lappon. 52. 15.*

Habitat in Lapponia.

*O. fenestrali* valde affine et simile, sed paulo minus, antennis elytris, ut videtur, brevioribus, thorace non canaliculato, et pube paulo longiore supra vestitum, distinctum. Antennae capite thoraceque parum longiores, filiformes. Thorax subtiliter punctatus, non canaliculatus, niger. Elytra thorace duplo longiora, crebre subrugoso-punctata, pubescentia, fere pilosa, picea. Pedes fusco-lutei.

32. *O. alpinum*: *Ovatum, convexum, tenuissime pubescens, profunde punctatum, totum testaceum, pedibus pallidioribus, antennis extrorsum sub-fuscescentibus.* — *Long.  $\frac{3}{4}$  lin.*

*Zetterst. Ins. Lappon. 52. 20.*

Habitat in Lapponia meridionali.

*O. pygmaeo* affine et sat simile, sed paulo brevius, et latius, elytris paulo longioribus profundiusque punctatis, corpore pallidior etc. abunde distinctum. Totum nitidum, omnium subtilissime rigide pubescens, flavescenti-testaceum, oculis nigris, et vertice macula fusca. Antennae thorace vix longiores, pubescentes, testaceae, extrorsum crassiores, fuscuscentes. Thorax convexus, transversus, lateribus subrotundatus, totus punctatus, sed foveolis nullis impressus. Elytra thorace fere dimidio longiora, distincte punctata, punctis certo situ seriatim dispositis. Abdomen pone elytra eorum longitudine, laeve, Pedes flavescentes. (Zett.)

33. *O. luridum*: *Oblongo-ovatum, rufo-piceum, antennis extrorsum gradatim crassioribus pedibusque testaceis, thorace postice angustiore, elytris confertissime punctatis.*

*Gyll. Ins. Suec. III. 701. 22 — 23. — Sahlb. Ins. Fenn. I. 285. 19.*

Habitat in Smolandia, Fennia.

Magnitudo et fere statura *O. pygmaei*, sed distinctum, thorace postice angustiore, etc.; ab *O. inflato* differt: corpore longiore, minus convexo, antennarum structura etc. Caput subtriangulare, brunneum, punctatum, inter antennas vix impressum, ore testaceo; oculi laterales, subglobosi, nigri. Antennae capite thoraceque longiores, pallide testaceae, tenue pubescentes, articulo primo valde elongato, apice incrassato, obconico, secundo etiam crasso, sed brevi; tertio tenui, longiusculo, obconico; quarto parvo, globoso, 5—10 gradatim crassioribus, truncatis, ultimo adhuc crassiore, ovato. Thorax longitudine latior, basi apiceque truncatus, lateribus tenue marginatus, ante medium rotundato-dilatatus, posterius angustior, angulis subrectis, supra



modice convexus, aequalis, rufo-piceus, parum nitidus, crebre punctatus. Scutellum parvum, triangulare, piceum, laeve. Elytra thorace duplo longiora, antice thoracis basi latiora, posterius nonnihil dilatata, apice truncata, angulo exteriori rotundato, supra in dorso subdepressa, saturatius picea, confertissime et sat profunde punctata, pilis minutissimis griseis parce adspersa. Abdomen pone elytra illorum fere longitudine, latum, marginatum, piceum, laeve, ano subacuminato, testaceo. Corpus subtus pallidius rufo-testaceum, nitidum, laeve. Pedes tenues, pallide testacei. (Gyll.)

34. *O. marginatum*: Piceum, punctatum, antennis pedibusque obscure rufis. — Long. 1 lin.

Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 463. 1.

Habitat in Missuria superiore.

Corpus oblongo-ovale, piceum, punctatum. Caput disperse, apice obsolete punctatum, inter antennis utrinque foveola, et basi utrinque sulculo impressum. Antennae cinereo-pilosellae, obscure rufo-piceae, basi rufae. Palpi rufi. Thorax transversim subquadratus, aequaliter punctatus, margine posteriore obscure piceo. Elytra oblongo-quadrata, ultra abdominis medium elongata, apice angulis exterioribus rotundatis, interioribus acuminatis, substriata, margine exteriori humerum versus piceo. Abdomen supra margine laterali et segmentorum singulorum margine posteriore obscure rufo. Pedes picei, apice dilutiores. (Say.)

35. *O. cursor*: Nitidulum, nigrum, elytris pedibusque picescentibus. — Long.  $1\frac{1}{3}$  lin.

Grav. Mon. 208. 10.

Habitat in Borussia.

Caput orbiculatum, thorace paulo minus, laeve, nitidum, nigrum, antennis, capite thoraceque paulo longioribus, fuscis. Thorax convexiusculus, suborbiculatus, coleopteris paulo minor, nitidus, laevis, niger. Coleoptera subtransversa, laevia, fusco-picea. Pedes ochracei. (Grav.)

Obs. Species dubia, typo in collectione Hellwigiana perditio aegre unquam recognoscenda.

### **Anthobium Leach.**

Steph. Illustr. V p. 335. — Er. Col. March. 1. p. 637.

*Mandibulae muticae.*

*Maxillae malis membranceis.*

*Palpi maxillares articulo ultimo penultimo longiore.*

*Tibiae muticae.*

*Tarsi articulis 4 primis aequalibus, brevibus, dilatatis.*

Corpus latiusculum, breviusculum, subdepressum, alatum. Caput thorace multo minus et angustius, breviusculum, basi distincte constrictum, fronte plana, immarginata, oculis sat magnis, rotundatis, prominulis. Ocelli inter oculos siti. Labrum transversum, apice subsinuatum. Mandibulae breves, muticae. Maxillae (t. V. f. 32. b.) malis membranaceis, exteriori oblonga, apice barbatula, interiore extus cornea, apice denticulo terminata,

oblique subemarginata ciliataque. Palpi maxillares articulo primo minuto, secundo modicè elongato, tertio brevior, quarto hoc sesqui longior, apice subacuminato. Labium (t. V. f. 32. a.) mento antrorsum angustato, apice truncato, ligula biloba, lobis coniunctis profunde emarginatis. Palpi labiales breviusculi, articulo primo tertioque longioribus, secundo duplo brevior. Antennae filiformes vel apicem versus perparum incrassatae, articulo tertio secundo paulo longior. Thorax coleopteris parum angustior, lateribus tenuiter marginatus. Scutellum triangulare. Elytra sat ampla, plerumque abdominis maximam partem obtegentia, angulo apicali exteriore rotundato. Abdomen lateribus rotundatum, late marginatum, apice subacuminatum. Pedes breves, femoribus sat validis, tibiis subcurvatis, pubescentibus, tarsis brevibus, articulis 4 primis brevibus, inter se longitudine aequalibus, magis minusve dilatatis, subtus dense pubescentibus, quarto cordato, quinto illis coniunctis longitudine aequali.

#### Observationes.

1. Corpus plerisque latiusculum, breviusculum, elytris maximam abdominis partem obtegentibus, aut glabrum aut subtiliter pubescens.

2. Tarsorum structura generi praecedenti proximum, at articuli primi magis dilatati, ceterum antennis gracilioribus, oculis maioribus et magis prominentibus, ocellis non super sed inter oculorum marginem superiorem sitis discedens. Ligula eadem forma ac in *Anthophago* et *Acidota*.

3. Mas abdominis segmento inferiore sexto profundius sinuato quam in reliquis *Omalinis* distinctus. Nonnullarum specierum feminae elytris angulo apicali inferiore producto, acuminato insignes.

4. Victus in floribus.

\* 1. *A. florale*: *Nigrum, nitidulum, antennis, ore pedibusque testaceis, elytris thorace plus duplo longioribus, apice interiore in utroque sexu truncatis.* — Long.  $1\frac{1}{2}$  lin.

*Mannerh. Brachelytr.* 54. 8.

*Omal. florale* Grav. *Micr.* 118. 10. *Mon.* 210. 14. — *Latr. Hist. nat. Crust. et Ins.* IX. 375. 10. — *Oliv. Encycl. meth.* VIII. 477. 14.

*Staph. floralis* Panz. *Faun. Germ.* 11. 12. *Ent. Germ.* 358. 41.

Habitat in Germania, Gallia.

A. minuto duplo maius, praecipue latius, nigrum, nitidulum, capite thoraceque glabris, elytris abdomineque parce omnium subtilissime pubescentibus. Antennae maris capitis thoracisque longitudine, feminae capite thoraceque paulo breviores, apicem versus leviter incrassatae, lacte testaceae. Os testaceum. Caput parce subtiliter obsoleteque punctatum, fronte inter oculos et inter antennis foveolis binis impressa, utrinque ad oculum subtiliter longitudinaliter strigosa. Thorax coleopteris angustior, latitudine duplo fere brevior, lateribus leviter rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusiusculis, leviter transversim convexus, parcius subtiliter punctatus, ad marginem lateralem pone medium utrinque impressus. Scutellum laeve. Elytra lata, thorace plus duplo longiora, apice interiore truncata, angulo suturali acutiusculo, confertissime punctata, iam nigro-picea, iam testaceo-fusca, nitida. Abdomen nigrum, apice summo testaceo. Pedes testacei.

- \* 2. *A. triviale*: *Nigrum, ore, antennis pedibusque testaceis, thorace piceo, elytris fusco-testaceis, thorace duplo longioribus, apice interiore in utroque sexu truncato. — Long.  $1\frac{1}{2}$  lin.*

*Er. Col. March. I. 639. 2.*

*Anthob. florale Boisd. et Lacord. Faun. Ent. Paris. I. 472. 6. (forte)*

Habitat in Germania, Gallia, Anglia.

Praecedenti affine, at angustius et magis elongatum, nigrum, subnitidum, glabrum, abdomine subtilissime pubescente. Antennae capite thoraceque longiores, graciliores, apicem versus leniter incrassatae, testaceae. Os testaceum. Caput fronte inter oculos et inter antennis foveolis binis impressa, ad oculum utrinque subtiliter longitudinaliter strigosa. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus leviter rotundatus, basin versus subangustatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusiusculis, parum convexus, sparsim minus subtiliter punctatus, nigro-piceus, marginibus rufescentibus. Scutellum laeve. Elytra thorace plus duplo longiora, dense sat fortiter punctata, fusco-testacea, limbo apicali dilutiore, apice interiore truncato. Abdomen nigrum vel piceum, apice summo testaceo. Pedes testacei.

- \* 3. *A. nigrum*: *Nigrum, nitidum, antennarum basi pedibusque testaceis, elytris thorace plus duplo longioribus, profunde punctatis, fuscis, apice interiore truncato. — Long.  $1\frac{1}{3}$  lin.*

Habitat in Helvetia.

Statura fere et summa affinitate *A. floralis*, nigrum, nitidum, parce subtiliter pubescens. Antennae maris capitis thoracisque fere longitudine, feminae iisdem paulo breviores, apicem versus sensim leviter incrassatae, nigro-fuscae, articulis 5 primis testaceis. Os testaceum. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus leviter rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusis, leviter convexus, lateribus pone medium depressis, aequaliter sat crebre minus subtiliter profunde punctatus, linea media longitudinali laeviore. Scutellum laeve. Elytra ampla, thorace plus duplo longiora, angulo apicali interiore in utroque sexu truncato, dense fortiter profundeque punctata, nigro-fusca, nitida. Abdomen parce subtilissime punctatum. Pedes testacei.

Obs. Antennis bicoloribus et punctura fortiore facile ab *A. floralis* distinctum.

- \* 4. *A. obsoletum*: *Nigrum, parum nitidum, ore, antennarum basi pedibusque testaceis, elytris thorace duplo longioribus, fuscis, apice interiore truncato. — Long.  $1\frac{1}{2}$  lin.*

Habitat in Lusitania.

Statura fere praecedentium at paulo minus, subdepressum, nigrum, parum nitidum, capite thoraceque glabris, elytris abdomineque pube subtilissima brevissimaque adspersis. Antennae capite thoraceque paulo breviores, apicem versus leniter incrassatae, testaceae, articulis ultimis 5 fuscis. Caput vage subtiliter punctatum, inter oculos obsolete bifoveolatum, inter antennis utrinque obsolete impressum, ad oculum utrinque subtilissime strigosum, ore testaceo. Thorax coleopteris angustior, latitudine dimidio brevior, late-



ribus modice rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, depressus, vage subtiliter punctatus. Scutellum laeve. Elytra thorace duplo longiora, dense subtilius punctata, fusca, apice interiore truncato. Abdomen subtilissime punctulatum, apice summo testaceo. Pedes testacei.

Mas latet.

- \* 5. *A. excavatum*: *Elongatum, nigrum, nitidum, antennarum basi, tibiis tarsisque testaceis, fronte late excavata, antice utrinque carinulata, thorace longitudinaliter impresso, dorso utrinque tuberculato, elytris crebre punctatis, apice coniunctim rotundatis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Helvetia, Dom. Schüppel.

Elongatum, lineare, subdepressum, nigrum, glabrum. Antennae capite thoraceque parum breviores, apicem versus sensim incrassatae, nigrae, articulis 5 primis testaceis. Palpi testacei. Os piceum. Caput thorace paulo angustius, impunctatum, fronte late at minus profunde excavata, antice in utroque latere carinula elevata sat prominula acutaque longitudinali instructa, minus nitidum. Thorax coleopteris paulo angustior, latitudine duplo prope brevior, lateribus rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusis, supra parum convexus, angulis posterioribus depressis, dorso late longitudinaliter impressus, impressionis margine utrinque ante medium leviter elevato et tuberculo minuto prominulo instructo, basi et lateribus crebrius subtiliter punctulatus, minus nitidus. Scutellum laevisimum. Elytra thorace triplo prope longiora, apice coniunctim rotundata, dense minus subtiliter profundius punctata, fusco-picea, nitida. Abdomen laeve, nigrum, nitidum, apice testaceum. Pedes nigro-picei, geniculis, tibiis tarsisque testaceis.

Femina latet.

- \* 6. *A. abdominale*: *Rufo-testaceum, elytris testaceis, thorace plus duplo longioribus, apice interiore in utroque sexu truncato, thorace angulis posterioribus obtusis, dorso foveolis duabus longitudinalibus magis minusve obsolete impresso, abdomine maris nigro.* — Long.  $1\frac{1}{2}$  —  $1\frac{3}{4}$  lin.

*Er. Col. March I.* 639. 8.

*Omalius abdominale Grav. Mon.* 219. 27. — *Oliv. Encycl. meth. VIII.* 480. 27.

— *Runde Brachelytr. Hal.* 22. 7.

*Dermestes semicoleoptratus Panz. Faun. Germ.* 24. 6.

Habitat in Germania.

Corpus oblongum, subdepressum, glabrum, abdomine subtilissime parce pubescente. Antennae capite thoraceque paulo longiores, apicem versus leviter incrassatae, testaceae. Caput vage punctatum, fronte inter oculos lineolis duabus longitudinalibus, inter antennis foveolis duabus obsolete impressa, rufo-testaceum, subnitidum, basi nigricante. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus rotundatus, basi apiceque truncatus, angulis omnibus obtusis, anterioribus subrotundatis, subdepressus, minus confertim subtiliter punctatus, linea leviter impressa, medio obsoleta canaliculatus, dorso foveolis duabus longitudinalibus, approximatis, interdum sat distinctis, saepius obsolete impressus, rufo-testaceus, parum nitidus,

medio in mare saepe longitudinaliter nigricans. Scutellum laeve, iam nigrum, iam piceo-testaceum. Elytra thorace plus duplo longiora, dense fortius punctata, testacea, circa scutellum nonnunquam fusciscentia. Abdomen maris nigrum, feminae testaceum, iam immaculatum. Pectus maris iam nigrum, iam testaceum, feminae testaceum. Pedes testacei.

- \* 7. *A. limbatum*: *Rufo-testaceum, elytris testaceis, apice interiore in utroque sexu truncato, thorace angulis posterioribus rectis, obsolete canaliculato. — Long.  $1\frac{1}{2}$  lin.*

Habitat in Hercyniae floribus Salicis, Dom. Saxesen, in Austria, Dom. Ullrich.

A. abdominali affine, at duplo fere minus, angustius, glabrum, abdomine apice subtiliter pilosello. Antennae apicem versus leniter incrassatae, capite thoraceque paulo longiores, testaceae, articulis ultimis 5 nigro-fuscis. Caput parce punctatum, fronte inter oculos lineolis duabus abbreviatis longitudinalibus, inter antennas foveolis duabus minutis impressa, rufo-testaceum, nitidum, basi nigricans. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, parum convexus, parce subtiliter punctatus, dorso obsolete canaliculatus, rufo-testaceus, subnitidus, longitudinaliter infuscatus. Scutellum laeve, iam nigrum, iam piceo-testaceum. Elytra thorace duplo longiora, sat dense fortius punctata, apice in utroque sexu truncata, luteo-testacea, circa scutellum nonnunquam infuscata. Abdomen laeve, nigrum. Pectus nigrum. Pedes testacei.

Variat elytris fuscis.

Obs. Differt ab A. abdominali praecipue statura angustiore et thoracis angulis posterioribus subrectis, femina mari haud dissimili.

- \* 8. *A. ophthalmicum*: *Rufo-testaceum, confertissime subtiliter punctatum, elytris flavo-testaceis, thorace duplo longioribus, apice interiore in utroque sexu truncato. — Long. 1 lin.*

*Staph. ophthalmicus* Payk. Faun. Suec. III. 419. 54.

*Omal. ophthalmicum* Gyll. Ins. Suec. II. 215. 7. — Germ. Faun. Ins. Europ. XIII.

6. — Mannerh. Brachelytr. 53. 14. — Boisd. et Lacord. Faun. Ent. Paris. I. 490.

9. — Runde Brachelytr. Hal. 22. 5.

*Omal. pallidum* Grav. Mon. 217. 25. — Oliv. Encycl. meth. Ins. VIII. 480. 25.

Habitat in Suecia, Germania.

A. minuto paulo latius, rufo-testaceum, parum nitidum, glabrum, abdomine parcius subtilissime pubescente. Antennae capitis thoracisque longitudine, apicem versus leviter incrassatae, testaceae, apice fusciscentes. Caput confertim subtiliter punctatum, fronte planiuscula, inter antennas foveolis duabus minutis impressa. Thorax coleopteris paulo angustior, latitudine plus dimidio latior, lateribus leviter rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, subdepressus, confertissime subtiliter punctatus, obsolete canaliculatus. Scutellum laeve. Elytra thorace duplo longiora, subdepressa, confertim fortius profundeque punctata, dilutius testacea. Abdomen subtilissime punctulatum. Pedes testacei.

- \* 9. *A. Sorbi*: Testaceum, parce subtiliterque punctatum, elytris flavo-testaceis, thorace duplo longioribus. — Long.  $\frac{3}{4}$  lin.

*Er. Col. March. I.* 640. 5.

*Omal. Sorbi* Gyll. *Ins. Succ. II.* 206. 8. — *Mannerh. Brachelytr.* 53. 15. — *Runde Brachelytr. Hal.* 22. 6.

*Omal. ophthalmicum* Grav. *Mon.* 216. 24. — *Oliv. Encycl. meth. VIII.* 479. 24. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 490. 9.

*Silpha lutea* Marsh. *Ent. Britt.* 128. 41.

*Var. a: Abdomine nigro.*

*Omal. testaceum* Grav. *Mon.* 218. 26. — *Oliv. Encycl. meth. VIII.* 480. 26.

*Var. b: Pectore abdominisque apice nigris.*

*Var. c: Pectore abdomineque nigris.*

*Silpha torquata* Marsh. *Ent. Britt.* 127. 40.

Habitat in Europa.

A. minuto duplo minus, luteo-testaceum, subnitidum, elytris pedibusque dilutioribus. Antennae capitis thoracisque fere longitudine, apicem versus sensim incrassatae, pilosellae, pallide testaceae, apicem versus fusciscentes. Caput parce obsoleteque punctulatum, fronte inter oculos et inter antennis foveolis binis minutis impressa. Thorax coleopteris angustior, maris latitudine dimidio brevior, feminae duplo fere brevior, lateribus leviter rotundatis, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, depressis, parum convexus, vage subtiliter punctulatus. Elytra thorace duplo vel plus duplo longiora, apice coniuncta subrotundata, dense fortius punctata. Abdomen iam testaceum, versus apicem saepe infuscatum, iam nigrum. Pectus iam testaceum, iam nigrum. Pedes pallide testacei.

- \* 10. *A. tempestivum*: Nigrum, antennarum basi thoraceque rufo-testaceis, ore, pedibus elytrisque luteis, his circa scutellum infuscatis. — Long. 1 lin.

Habitat in Sardinia, Dom. Prof. Gené.

Statura omnino A. minuti, glabrum, abdomine parce subtiliterque pubescentibus. Antennae capitis thoracisque fere longitudine, apicem versus leniter incrassatae, testaceae, articulis ultimis 5 nigricantibus. Caput thorace vix angustius, fronte depressa vel subimpressa, crebre punctata, inter antennis utrinque leviter impressa, nigrum, nitidum, ore testaceo, fronte media macula piceo-testacea. Thorax coleopteris dimidio prope angustior, latitudine dimidio brevior, lateribus leviter rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusis, leviter convexus, crebre subtilius punctatus, obsolete canaliculatus, rufo-testaceus, nitidus. Scutellum laevissimum, nigrum. Elytra thorace plus duplo longiora, crebre fortius profundeque punctata, apice coniunctim leviter rotundata, lutea, nitida, circa scutellum nigricantia. Pectus et abdomen nigra, nitidula. Pedes testacei.

Femina latet.

- \* 11. *A. scutellare*: Rufo-testaceum, scutello, pectore abdomineque nigris, elytris testaceis, thorace duplo longioribus, apice interiore maris truncato, feminae acuminato. — Long. vix 1 lin.

*Omal. testaceum* Boisd. et Lacord. *Faun. Ent. Paris. I.* 489. 8.



Habitat in Austria, Gallia.

Statura omnino *A. minuti*, rufo-testaceum, parum nitidum, totum subtiliter tenuiterque pubescens. Antennae capitis thoracisque longitudine, apicem versus leniter incrassatae, testaceae, apicem versus fuscescentes. Caput subtilissime punctulatum, fronte inter oculos punctis duobus profundis, inter antennis foveolis duabus minutis obsoletis impressa, basi summa nigricans. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, leviter convexus, subtiliter punctulatus. Scutellum laeve, nigro-piceum. Elytra thorace plus duplo longiora, subdepressa, confertim fortius punctata, testacea, apice interiore in mare truncato, in femina producto, acuminato. Abdomen et pectus nigra. Pedes testacei.

Obs. *Omal. testaceum* Grav., secundum exemplum typicum ab auctore benevole communicatum, ad *A. Sorbi* retuli.

\* 12. *A. minutum*: Nigrum, antennarum basi pedibusque testaceis, elytris fuscis, maris apice interiore truncato, feminae acuminato. — Long. 1 lin.

*Er. Col. March. I.* 638. 1.

*Silpha minuta* Fab. *Ent. Syst. I.* 254. 26. *Syst. El. I.* 342. 25. — *Panz. Ent. Germ.* 122. 18. — *Herbst. Col. V.* 207. 32. — *Schönh. Syn. II.* 132. 27.

*Omal. Ranunculi* Grav. *Micr.* 110. 11. *Mon.* 215. 20. — *Latr. Hist. nat. Crust. et Ins. IX.* 374. 11. — *Oliv. Encycl. meth. VIII.* 478. 20. — *Gyll. Ins. Suec. II.* 204. 6. — *Sahlb. Ins. Fenn. I.* 279. 6. — *Zetterst. Faun. Lappon. I.* 50. 5. *Ins. Lappon.* 50. 7. — *Mannerh. Brachelytr.* 52. 12. — *Boisd. et Lacord. Faun. Ent. Paris. I.* 489. 7. — *Runde Brachelytr. Hal.* 22. 4.

Habitat in Europa, in floribus pratensis.

Parvum, depressiusculum, nigrum, subnitidum, glabrum, abdomine parce subtilissime pubescente. Antennae capite thoraceque vix breviores, crassiusculae, testaceae, apice fuscae. Caput parce subtiliter punctulatum, fronte longitudinaliter bi-impressa, ore testaceo. Thorax coleopteris angustior, latitudine dimidio prope brevior, transversim quadratus, lateribus leviter rotundatis, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus rectis, parum convexus, margine laterali depresso, parcius subtiliter punctulatus, utrinque ad marginis lateralis medium obsolete foveolatus. Scutellum laeve. Elytra abdominis maximam partem obtegentia, subdepressa, dense fortius punctata, punctis hinc inde subseriatis, fusca vel testaceo-fusca, subnitida, apice interiore in mare truncato, in femina producto, acuminato, dehiscente. Abdomen laeve, apice summo piceo. Pedes validiusculi, testacei.

\* 13. *A. flavipenne*: Nigrum, parum nitidum, antennarum basi, pedibusque testaceis, thorace brevi, canaliculato, elytris thorace plus duplo longioribus, pallide testaceis. — Long. 1 lin.

Habitat in Lapponia, a Dom. Prof. Zetterstedt pro *Om. Ranunculi* Gyll. communicatum.

Statura et summa affinitate *A. minuti*, nigrum, parum nitidum, glabrum, abdomine subtilissime pubescente. Antennae capite thoraceque paulo breviores, apicem versus haud incrassatae, testaceae, articulis ultimis 5 nigris. Palpi testacei. Mandibulae rufae. Caput subtilissime parce punctulatum,

fronte antice utrinque leviter oblique impressa, inter antennas profundius foveolata. Thorax coleopteris paulo angustior, latitudine duplo brevior, lateribus leviter rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusis, parum convexus, margine laterali posterior subdepresso, medio foveolato, parce subtiliter punctulatus, medio leviter canaliculatus. Scutellum laeve. Elytra thorace plus duplo longiora, subdepressa, sat dense fortius punctata, flavo-testacea, angulo interiore maris truncato. Abdomen subtilissime punctulatum, nitidulum. Pedes testacei.

Femina latet.

\*Obs. Ab A. minuto praeter colorem antennis brevioribus, apicem versus haud incrassatis, thorace latiore, brevior, canaliculato, elytris brevioribus sat distinctum.

- \* 14. *A. montanum*: *Subdepressum, nigrum, parum nitidum, pedibus antennisque testaceis, his apice nigris, elytris thorace plus triplo longioribus, fuscis, maris apice coniunctim rotundatis, feminae acuminatis.* — *Long. vix 1 lin.*

Habitat in Austriae Helvetiaeque alpibus.

A. minuto paulo minus, subdepressum, nigrum, minus nitidum, glabrum. Antennae capite thoraceque paulo longiores, testaceae, articulis ultimis tribus paulo maioribus, nigricantibus. Os piceum. Caput thorace paulo angustius, vix punctulatum, fronte postice utrinque obsolete oblique bisulcata, antice leviter bifoveolata. Thorax coleopteris paulo angustior, latitudine dimidio brevior, lateribus leviter rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus rectis, subdepressus, parce subtiliter obsoleteque punctatus, subtilissime canaliculatus. Scutellum laeve. Elytra thorace plus triplo longiora, in utroque sexu abdomen fere totum tegentia, maris apice coniunctum fortiter rotundata, feminae angulo apicali interiore acuminato, subdepressa, crebre punctata, fusca, basi nigricantia, apice indeterminate testacea. Pedes testacei.

Obs. Ab A. minuto statura minore, magis depressa, colore capitis thoracis-que opaco, hoc parce obsolete punctato, elytris multo longioribus, maris apice haud truncatis, antennis longioribus, articulis ultimis tribus solis maioribus distinctum.

- \* 15. *A. luteicorne*: *Subdepressum, nigrum, parum nitidum, ore, antennis pedibusque luteis, elytris fuscis, thorace plus duplo longioribus, feminae apice acuminatis.* — *Long. vix 1 lin.*

Habitat in Sicilia, Dom. Grohmann.

Statura et summa affinitas praecedentis. Nigrum, parum nitidum, glabrum. Antennae luteo-testaceae (articuli ultimi tres desunt). Os testaceum. Caput thorace paulo angustius, vix punctulatum, fronte inter oculos bifoveolata, inter antennas transversim impressa. Thorax coleopteris paulo angustior, latitudine vix dimidio brevior, lateribus leviter rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus subrectis, subdepressus, parce subtiliter obsoleteque punctatus, obsolete subtiliter canaliculatus. Scutellum laeve. Elytra thorace plus duplo longiora, subdepressa, dense hinc inde subseriatim punctata, fusca, apice testaceo-marginata, angulo apicali interiore acuminato. Pedes testacei.

## Mas latet.

Obs. Praecedenti proximum, et imprimis elytris multo brevioribus, vix ultra abdominis dimidiam partem tegentibus distinctum.

- \* 16. *A. anale*: Nigrum, nitidum, ano, pedibus antennisque testaceis, his apice fuscis, elytris fuscis, fortius punctatis, maris thorace duplo longioribus, apice rotundatis, feminae thorace plus triplo longioribus, acuminatis. — Long.  $\frac{2}{3}$  — 1 lin.

Habitat in montibus Saxoniae, Austriae, Croatiae.

Statura omnino *A. minuti*, nigrum, nitidum, glabrum. Antennae vix capitis thoracisque longitudine, apice sensim leniterque incrassatae, testaceae, plerumque apicem versus fuscесcentes. Os piceum. Caput thorace angustius, fronte inter oculos lineola obliqua, inter antennis foveola minuta levique utrinque impressa, ceterum laeve. Thorax coleopteris paulo angustior, latitudine paulo brevior, lateribus fortius rotundatus, basi apiceque truncatus, angulis omnibus obtusis, modice convexus, parce obsoleteque punctatus. Scutellum laeve. Elytra maris thorace duplo longiora, apice coniuncta rotundata, feminae thorace plus triplo longiora, angulo apicali interiore acuminato, dense fortius profundiusque punctata, fusca, apice saepius testaceo-limbata. Abdomen ano maris laete rufo-testaceo, feminae piceo. Pedes testacei.

Obs. Elytris fortius punctatis imprimis ab affinibus distinctum.

- \* 17. *A. luteipenne*: Nigrum, ore, antennis, pedibus coleopterisque luteis, his thorace triplo longioribus, maris apice rotundatis, feminae acuminatis. — Long. 1 lin.

Habitat in Saxonia montana, Dom. Märkel.

Statura omnino *A. longipennis*, subdepressum, nigrum, opacum, glabrum. Antennae capitis thoracisque longitudine, apicem versus vix incrassatae, testaceae. Os testaceum. Caput thorace angustius, impunctatum, fronte utrinque leviter longitudinaliter impressa. Thorax coleopteris paulo angustior, latitudine plus dimidio lator, lateribus fortius rotundatus, basi apiceque truncatus, angulis anterioribus rotundatis, posterioribus obtusis, parum convexus, parce obsoleteque punctatus, subtiliter obsoleteque canaliculatus. Scutellum laeve, testaceum. Elytra thorace triplo longiora, maris apice coniuncta rotundata, feminae angulo interiore acuminato, crebre minus fortiter punctata, interstitiis subtilissime rugulosis, luteo-testacea, vix nitida. Abdomen ano maris testaceo, feminae piceo. Pedes testacei.

- \* 18. *A. longipenne*: Testaceum, coleopteris thorace plus triplo longioribus, maris apice truncatis, feminae coniunctim rotundatis. — Long. 1 lin.

Er. Col. March. I. 640. 4.

Habitat in Germania, Suecia.

Statura *A. minuti*, licet colore corporis testaceo et elytris multo longioribus valde distinctum. Corpus glabrum. Antennae capitis thoracisque prope longitudine, apicem versus leniter incrassatae. Caput subtilissime parce punctulatum, foveolis duabus minutis inter oculos, duabusque inter antennis impressum, rufo-testaceum, subnitidum, summa basi nigricante. Thorax



coleopteris paulo angustior, latitudine plus dimidio latior, lateribus leviter rotundatus, basi apiceque truncatus; angulis anterioribus rotundatis, posterioribus obtusiusculis, margine laterali antice anguste, postice latius depresso, disco leviter convexus, parce obsoleteque punctulatus, utrinque ad marginis lateralis medium subfoveolatus, rufo-testaceus, parum nitidus. Scutellum rufo-testaceum, laeve. Elytra thorace plus triplo longiora, abdomen fere totum tegentia, praesertim in femina, dense fortius punctata, testacea, subnitida, apice interno in mare truncato, in femina magis producto, angulo suturali acutiusculo, cum elytro altero coniunctim rotundato. Abdomen maris nigrum, apice testaceum, feminae totum testaceum. Pectus et pedes testacea.

Obs. Specimina duo feminea elytris apice interiore longius producto acuminatis, cum maribus et feminis genuinis e Croatia a Dom. Parreyss communicata, specie haud differre videntur.

Species mihi invisae, forte huius generis.

19. *A. fimetarium*: Rufo-piceum, nitidum, crebre punctatum, antennarum basi, thoracis elytrorumque limbo pedibusque pallidioribus, thorace transverso, canaliculato, elytris substriatis, abdomen fere totum obtegentibus, maris truncatis, feminae apice interiore sinuato-acuminatis.

*Omal. fimetarium* Mannerh. *Brachelytr.* 52. 10.

Habitat in stercore in ins. Sitcha Americae borealis.

Delipthro tecto maius, fortius punctatum, thorace brevior, canaliculato, elytrisque longioribus, substriatis, in femina sinuatis diversum. (Mannerh.)

20. *A. Lapponicum*: Nigrum, obscurum, subtilissime punctatum, antennarum basi, thoracis lateribus, elytris pedibusque pallidis, thorace convexo.

*Omal. Lapponicum* Mannerh. *Brachelytr.* 52. 13. — Zetterst. *Ins. Lappon.* 50. 9.

Habitat in Lapponia.

A. minuto (Om. Ranunculi) duplo minus, subtilius punctulatum, thorace convexo, posterius minus angustato et elytris brevioribus facile distinguendum. (Mannerh.)

#### O b s e r v a t i o n e s.

1. Huic tribui forte adscribendum est genus mihi invisum:

### **Boreaphilus Sahlberg.**

Palpi anteriores 4-articulati, articulo tertio clavato, ultimo parvo, subulato, posteriores minuti, filiformes. Mandibula elongata, tenuis, falcata, integra. Antennae subfiliformes, extrorsum paulo crassiores. Corpus postice dilatatum. Caput orbiculatum, subexsertum. Thorax oblongus.

1. *B. Henningianus*: Piceus, punctulatus, parce pubescens, ore, antennis pedibusque testaceis, segmentorum ventralium marginibus abdominis neque supra piceo-nigris, abdomine ovato, lateribus elevato-marginatis. — Long.  $1\frac{1}{4}$  lin.

Habitat in Lapponia boreali rarissime. In littore lacus Mandujarvi sub quisquiliis tria specimina die 9. Augusti 1830 legit F. Sahlberg.

Statura corporis Anthophagis quodammodo similis. Caput thorace latius, piceum, subnitidum, crebre punctatum, inaequale, fronte late et sat profunde excavata, versus thoracem parum impresso-emarginatum. Os et palpi piceo-testacea. Antennae longe ante oculos in angulo frontis insertae, longitudine thoracis, extrorsum paulo crassiores, testaceae, articulo primo elongato, incrassato, secundo et tertio paulo brevioribus et angustioribus, tribus insequentibus brevibus, 7 — 10 parum incrassatis, ultimo longiore, ovato, acuto. Oculi nigri, prominuli. Thorax latitudine longior, antice paulo angustior, pro emarginatura capitis parum rotundato-productus, lateribus anguste marginatus, paulo ante medium angulato-dilatatus, postice truncatus, parce pubescens, fusco-piceus, foveis tribus, una utrinque laterali et tertia profundiore paulo ante basin impressis. Scutellum minutum, subtriangulare, fusco-piceum. Elytra thorace paulo latiora, eodemque et latitudine sua fere dimidio longiora, posterius parum dilatata, apice subtruncata, angulis exterioribus rotundatis, supra ferrugineo-picea, crebre punctata, parce pubescentia. Abdomen ovatum, depressum, lateribus late elevato-marginatis, nigro-piceum, vix punctulatum, subpubescens, marginibus segmentorum piceo-nigris. Pedes elongati, rufo-testacei.

*Sahlb. Ins. Fenn. I. 433. 1. — Zetterst. Ins. Lappon. 72. 1.*

2. Alterum genus, quod huius tribus esse videtur, est *Coryphium* Kirby, a Stephensio Ent. Britt V. p. 344. descriptum, mihi invisum et incognitum. Corpus ei depressum, subpubescens. Caput triangulare, inaequale, oculis prominulis. Antennae filiformes, apicem versus sensim incrassatae, articulo basali valido, basi angustato, sequentibus duobus clavatis, reliquis turbinatis, ultimo ovato, acuminato. Palpi breves, maxillares articulo penultimo rotundato, ultimo fortius incrassato, truncato. Thorax capite angustior, lateribus subtiliter marginatus. Elytra oblonga, dimidio abdomine longiora. Abdomen dilatatum, marginatum, apice acuminato. Pedes graciles, simplices, tarsis 5-articulatis, articulo ultimo praecedentibus duobus coniunctis longiore.

A *Boreaphilo* haud distinctum esse hoc genus supponerem, nisi palporum descriptiones impedirent, quo minus alterum alteri subjungam. Non autem omnino deest quod dubitem, *Boreaphili* palpos maxillares rite observatos esse. Ceterum capite latiore et thorace angustiore utrumque genus inter *Omalinos* sat insigne.

---

**Tribus undecima.**  
**Proteinini.**

*Stigmata prothoracica occulta.*

*Coxae posticae transversae, anticae cylindricae, haud prominentes.*

*Trochanteres postici fulcrantes.*

---

Labrum transversum, corneum,  
apice membrana biloba auctum Proteino.

Mandibulae breves, haud prominentes, muticae.

Maxillae mala exteriore coriacea, apice barbata, interiore cornea, apice  
unco duplici terminata Proteino,  
unco simplici terminata Megarthro, Phloeobio.

Palpi maxillares articulo primo minuto, brevi, secundo incrassato,  
tertio hoc minore, quarto angustiore, subacuminato.

Mentum corneum.

Ligula membranea,  
biloba Proteino, Megarthro, Phloeobio;  
integra Micropeplo.

Paraglossae distinctae Proteino, Megarthro, Phloeobio;  
ligulae aequales Proteino, Phloeobio;  
ligulam longe superantes Megarthro.

Palpi labiales perbreves.

Antennae sub clypei margine laterali insertae,

11-articulatae, saepius clavatae vel apice leviter incrassatae, liberae plerisque;

9-articulatae, capitulatae, receptae Micropeplo.

Caput thoraci immissum, porrectum,

ocellis nullis plerisque;

ocello singulo Phloeobio.

Thorax marginatus, infra spatio pone coxas anteriores corneo.

Scutellum distinctum.

Elytra pectore longiora,

abdomen fere totum tegentia Proteino;

abdominis summam basin tegentia Megarthro, Phloeobio, Glypto-  
mati, Micropeplo.



Abdomen segmentis 6 distinctis, septimo nonnunquam eminulo, marginatum Proteino, Megarthro, Phloeobio, Micropeplo; immarginatum Glyptomati.

Pedes breves, coxis anticis cylindricis, oblique positis, haud exsertis, intermediis

approximatis Proteino, Phloeobio;

tenui spatio distantibus Megarthro, Glyptomati;

late distantibus Micropeplo;

posticis transversis, trochanteribus posticis fulcrantibus, tibiis, inermibus, tarsis

5-articulatis Proteino, Megarthro, Phloeobio;

3-articulatis Glyptomati, Micropeplo.

#### O b s e r v a t i o.

Proxima haec tribus praecedenti, at coxis anticis haud exsertis et paraglossis plerumque distinctis discedit. Coxarum anticarum forma cum Nitidulis convenit.

### **Proteinus Latr.**

*Latr. Précis d. caract. gén. des Ins. p. 9.*

*Antennae 11-articulatae, liberae, articulis ultimis tribus maioribus.*

*Tarsi 5-articulati.*

Corpus parvum, ovatum, leviter convexum. Caput thorace multo minus, subtriangulare, clypeo rotundato, oculis sat magnis, prominulis, ocellis omnino nullis. Labrum transversum, medio subsinuatum, apice membrana brevi biloba auctum. Mandibulae minutae, muticae. Maxillae (t. V. f. 33. b.) mala interiore cornea, apice acute bidentata, interne membrana tenui aucta, exteriore coriacea, apice barbata. Palpi maxillares breviusculi, articulo primo brevi, tenui, secundo magno, incrassato, obconico, tertio brevi, transverso, quarto tenuiore, longiore, tereti. Labium (t. V. f. 33. a.) mento transversum, antrorsum angustatum, apice truncato, ligula membranea, biloba, lobis truncatis, paraglossis ligulae connatis eamque haud excedentibus. Palpi labiales breves, ligulam haud superantes, articulo primo longiore, secundo brevissimo, tertio minuto, secundo autem longiore. Antennae articulis duobus primis maioribus crassioribusque, ultimis tribus sensim maioribus, ultimo subgloboso. Thorax transversus. Scutellum triangulare. Elytra sat ampla, abdominis segmenta duo anteriora obtegentia, apice truncata, angulo exteriore rotundata. Abdomen latiusculum, lateribus marginatum, apice acuminatum. Pedes breviusculi, coxis intermediis mesosterni carina tenui longitudinali discretis, tibiis omnibus simplicibus, tarsis 5-articulatis, articulis 4 primis subaequalibus, quinto his coniunctis multo brevioribus.

#### O b s e r v a t i o n e s.

1. Corpus subtiliter pubescens, punctulatum.
2. Habitus fere Nitidularum quarundam, imprimis Cercorum, quibus etiam ab auctoribus multis associatur.
3. Mas abdominis segmento inferiore sexto emarginato cognoscitur.
4. Victus in floribus et in fungis.

- \* 1. *P. brachypterus*: *Niger, nitidulus, antennarum articulo primo pedibusque testaceis, elytris crebre punctulatis*. — Long.  $\frac{3}{4}$  lin.

*Latr. Hist. nat. Crust. et Ins. X. 46. 1. Gen. Crust. et Ins. I. 298. 1. — Lepell. Enc. meth. X. 215. 1. — Mannerh. Brachelytr. 57. 1. — Boisd. et Lacord. Faun. Ent. Paris. I. 491. 1. — Runde Brachelytr. Hal. 24. 1. — Er. Col. March. I. 642. 1.*

*Dermestes brachypterus* *Fab. Ent. Syst. I. I. 235. 46. Syst. El. I. 320. 45. — Payk. Faun. Suec. I. 288. 14*

*Cateretes brachypterus* *Herbst Col. V. 13. 2. t. 45. f. 2. — Gyll. Ins. Suec. I. 251. 6. — Sahlb. Ins. Fenn. I. 84. 5. — Schönk. Syn. II. 150. 6.*

*Omal. brachypterus* *Gyll. Ins. Suec. II. 207. 9. — Sahlb. Ins. Fenn. I. 279. 7.*

*Omal. ovatum* *Grav. Mon. 215. 22. — Oliv. Encycl. meth. Ins. VIII. 479. 22.*

Habitat in Europa, in floribus et in fungis.

Breviter ovatus, leviter convexus, niger, nitidus, subtiliter griseo-pubescent. Antennae capitis thoracisque longitudine, nigrae, articulo primo testaceo. Oris partes testaceae, labro palpisque maxillaribus piceis. Caput laeve, fronte ad oculos utrinque oblique obsolete impressa. Thorax coleopteris angustior, latitudine duplo brevior, lateribus leviter rotundatus, a medio inde antrorsum angustatus, apice leviter emarginatus, basi subinnatus, angulis anterioribus obtusis, posterioribus subrectis, leviter transversim convexus, laevis, margine basali et apicali nonnunquam rufo-piceo. Scutellum laeve. Elytra thorace plus duplo longiora, ampla, lateribus leviter rotundata, leviter convexa, subtilius crebre punctata. Abdomen subtilissime punctulatum. Pedes rufo-testacei, coxis anterioribus concoloribus.

- \* 2. *P. brevicollis*: *Niger, subnitidus, antennis piceis, pedibus testaceis, elytris creberrime punctatis, fusco-piceis*. — Long. 1 lin.

Habitat in Italia, Dom. Schüppel, Lutetiae, Dom. Aubé.

Ovatus, subdepressus, parce minus subtiliter griseo-pubescent, niger, nitidus, capite thoraceque minus nitidis. Antennae capitis thoracisque longitudine, piceae, clava nigra, articulo primo rarius rufescente, maris fortius clavatae, feminae graciliores, apice lenius incrassatae. Palpi picei. Caput laeve, fronte utrinque ad oculi marginem interiorem oblique impressa. Thorax coleopterorum basi parum angustior, latitudine plus duplo brevior, lateribus rotundatus, a medio inde apicem versus leviter angustatus, apice levissime emarginatus, basi subbisinnatus, angulis posterioribus acutiusculis, anterioribus rotundatis, parum convexus, laevis, angulis posterioribus depressis. Scutellum laevissimum. Elytra thorace plus duplo longiora, creberrime minus subtiliter punctata, obscure fusco-picea, nitida, humeris dilutioribus. Abdomen subtilissime punctulatum, pube brevissima subtilissime adpersum. Pedes testacei, coxis anticis concoloribus.

Obs. Differt a praecedente statura paulo maiore, subdepressa, antennarum basi haud testacea, elytris distinctius punctatis, etc.

- \* 3. *P. macropterus*: *Niger, nitidulus, antennarum articulis duobus primis pedibusque testaceis, elytris piceis, crebre punctatis*. — Long.  $\frac{2}{3}$  lin.

*Er. Col. March. I. 643. 2.*

*Omal. macropterus* *Gyll. Ins. Suec. II. 209. 10. — Sahlb. Ins. Fenn. I. 280. 8.*

Habitat in Germania.

*P. brachyptero paulo minor, breviter ovalis, convexus, niger, nitidulus, parce subtiliterque griseo-pubescens. Antennae capite thoraceque paulo breviores, piceae, articulis duobus primis late rufo-testaceis. Caput laeve, fronte inter oculos utrinque oblique impressa, apice leviter bifoveolata. Thorax coleopterorum basi parum angustior, latitudine duplo prope brevior, lateribus leviter rotundatus, a basi apicem versus aequaliter subangustatus, basi subsinuatus, apice late emarginatus, angulis anterioribus rotundatis, posterioribus acutiusculis, leviter convexus, laevis. Scutellum laevissimum. Elytra thorace plus duplo longiora, lateribus leviter rotundata, leviter convexa, crebre subtiliter punctata. Abdomen subtilissime punctulatum. Pedes testacei, coxis anticis concoloribus.*

Obs. Differt a *P. brachyptero* antennarum articulis duobus primis testaceis, thorace non a medio sed a basi inde antrorsum angustato, lateribus minus, angulis anterioribus magis rotundatis, posterioribus magis acutis. — Gyllenhal hanc speciem thorace punctulato a *P. brachyptero* distinguit, thorax autem revera laevis est, et puncturae speciem nonnisi pilis subtilissimis offert.

\* 4. *P. atomarius: Fusco-piceus, antennis pedibusque pallidis, elytris crebre punctulatis. — Long.  $\frac{2}{5}$  lin.*

Habitat in Germania.

Habitu omnino *P. brachypteri*, at triplo vel quadruplo minor, breviter subovalis, parum convexus, fusco-piceus, nitidulus, subtiliter pubescens. Antennae capitis thoracisque longitudine, pallide flavescentes, apice fuscescentes. Oris partes testaceae. Caput laeve, inter oculos utrinque foveola rotundata obsolete impressum, apice obsolete bifoveolatum. Thorax basi coleopteris paulo angustior, latitudine duplo brevior, lateribus leviter rotundatus, a basi inde antrorsum subangustatus, apice leviter emarginatus, basi subbisinuatus, angulis anterioribus rotundatis, posterioribus acutiusculis, leviter convexus, laevis. Elytra thorace triplo longiora, ampla, lateribus leviter rotundata, leviter convexa, crebre subtiliter punctata. Abdomen apice dilutius. Pedes pallide testacei.

Obs. Proximus praecedenti, at statura duplo minore et thorace paulo breviori distinctus videtur.

### **Megarthus Kirby.**

*Steph. Illustr. Britt. Ent. V. p. 330.*

*Megarthus* \* *Er. Col. March. I. p. 644.*

*Antennae 11-articulatae, articulo ultimo maiore.*

*Frons oculo nullo.*

*Tarsi 5-articulati.*

Corpus parvum, latiusculum, subdepressum, alatum. Caput thorace multo minus et angustius, subtriangulare, basi constrictum, collo quasi crassiusculo thoraci immissum, porrectum, fronte marginata, oculis subglobosis, sat prominulis, ocellis nullis. Labrum frontis margine reconditum, transversum, breve, apice membrana integra prominula fimbriata. Mandibulae minutae, muticae. Maxillae (t. V. f. 34. b.) mala interiore cornea, elongata, apice acuminata, interne setulis paucis ciliata, exteriori coriacea, apice membrana barbataque, interiorem parum superante. Palpi maxillares brevius-



sculi, articulo primo minimo, secundo sat magno, obconico, tertio brevi, quarto secundo paulo brevior, tenuiusculo, acuminato. Labium (t. V. f. 34. a.) mento latitudine parum brevior, antrorsum sensim angustato, apice truncato, ligula membranea, brevi, medio subsinuata, paraglossis ligulam longe superantibus, liberis, acuminatis. Palpi labiales breviusculi, articulis duobus primis subaequalibus, tertio paulo minore. Antennae breviusculae, apicem versus sensim subincrassatae, articulo ultimo crassiusculo, subglobozo, basalibus duobus reliquis multo maioribus crassioribusque. Thorax transversus, coleopterorum prope latitudine, lateribus late marginatus. Elytra abdominis nonnisi summam basin tegentia, apice truncata, angulo exteriore obtuso. Abdomen breviusculum, apicem versus angustatum, late marginatum. Pedes breviusculi, coxis intermediis subapproximatis, mesosterni carinula discretis, tibiis omnibus muticis, tarsis 5-articulatis, articulis 4 primis brevibus, aequalibus, quinto his coniunctis brevior.

## O b s e r v a t i o n e s.

1. Corpus punctatum, pube brevi adpersum, thorace profunde canaliculato, angulis posterioribus excisis insigne.

2. Mas abdominis segmento inferiore sexto apice emarginato, nec non pedum posteriorum structura distinctus.

3. Victus sub arborum cortice.

\* 1. *M. depressus*: *Niger, opacus, pedibus ferrugineis, thorace lateribus rotundato.* — Long. 1 lin.

*Er. Col. March. I. 644. 1.*

*Staph. depressus* Payk. *Mon. Staph. 70. 49. Faun. Suec. III. 412. 58.* — *Oliv. Ent. III. 42. 36. 51. t. 3. f. 26.*

*Omal. depressum* Gyll. *Ins. Suec. II. 210. 11.* — *Sahlb. Ins. Fenn. I. 280. 9.* — *Zetterst. Ins. Lappon. 50. 8.* — *Mannerh. Brachelytr. 53. 16.*

*Phloeobium depressum* Boisd. et Lacord. *Faun. Ent. Paris. I. 494. 4.*

*Omal. macropterum* Grav. *Mon. 215. 21.* — *Oliv. Encycl. meth. VIII. 479. 21.*

Habitat in Europa.

Corpus parvum, latiusculum, subdepressum, nigrum, opacum, omnium subtilissime pubescens. Antennae piceae, articulis duobus primis crassioribus nigris. Palpi nigro-picei. Caput omnium subtilissime rugulosum, fronte utrinque late impressa, antice rotundata, margine subincrassato. Thorax coleopteris parum angustior, latitudine duplo brevior, basi apiceque truncatus, lateribus rotundatis, antrorsum modice angustatus, angulis anterioribus obtusis, posterioribus emarginatis, denticulis duobus obtusiusculis, leviter convexus, confertim subruguloso-punctatus, tenuiter at profunde canaliculatus, margine laterali parum depresso. Scutellum punctulatum, obsoletius canaliculatum. Elytra thorace duplo longiora, dense punctata, nigro-picea. Abdomen subtiliter punctulatum, ano piceo. Pedes rufo-ferruginei, femoribus saepe nigro-piceis.

Pedes posteriores maris femoribus subincrassatis, tibiis leviter incurvis.

\* 2. *M. sinuato-collis*: *Niger, opacus, pedibus rufis, thorace canaliculato, lateribus angulato.* — Long. 1 lin.

*Phloeobium sinuato-colle* Boisd. et Lacord. *Faun. Ent. Paris. I. 493. 3.*

Habitat in Helvetia, Gallia.

Statura omnino praecedentis, nigrum, opacum, subtiliter pubescens. Antennae fusco-nigrae. Palpi picei. Caput subtiliter rugulosum, utrinque late longitudinaliter impressum, margine anteriore lateribus parum elevato. Thorax coleopteris vix angustior, latitudine duplo brevior, antrorsum subangustatus, basi leviter trisinuatus, apice emarginatus, lateribus subrotundatus, medio angulatus, angulis anterioribus obtusis, posterioribus emarginatis, denticulis obtusiusculis, disco leviter convexus, creberrime rugoso-punctatus, tenuiter at sat profunde canaliculatus, lateribus explanatus. Scutellum confertim punctatum. Elytra thorace vix duplo longiora, confertissime sat fortiter punctata, nigro-picea. Abdomen obsolete punctulatum, subtilissime rugulosum, ano piceo-testaceo. Pedes rufi.

\* 3. *M. denticollis*: Nigro-piceus, opacus, antennarum articulo primo pedibusque rufis, thoracis margine rufo. — Long. 1 lin.

*Omalium denticolle* Beck Beitr. 26. 40. t. 7. f. 40.

*Megarthus marginicollis* Er. Col. March. I. 644. 2.

*Phloeobium marginicolle* Boisd. et Lacord. Faun. Ent. Paris. I. 492. 1.

*Silpha hemiptera* var. *α* Illig. Käf. Pr. 355. 1.

Habitat in Europa.

Statura praecedentium, niger, opacus. Antennae piceae, articulo primo rufo. Caput ruguloso-punctatum, utrinque late longitudinaliter impressum, interstitio subcarinato, margine anteriore acute elevato. Thorax basi coleopteris haud angustior, latitudine vix duplo brevior, apice truncatus, basi subtrisinuatus, lateribus parum rotundatis, angulis anterioribus rotundatis, posterioribus acute emarginatis, bidenticulatis, disco leviter convexus, rugoso-punctatus, tenuiter at profunde canaliculatus, lateribus explanatis rufis. Scutellum crebre punctatum, canaliculatum. Elytra subdepressa, sat fortiter creberrime punctata, margine apicali rufo-piceo. Abdomen punctulatum, nigrum, apice rufo-ferrugineum. Pedes rufi.

Pedes maris intermediis femoribus incrassatis, tibiis leviter incurvis, postici femoribus incrassatis, trochanteribus fortiter dilatatis, tibiis antecapicem emarginatis, apice uncinatis.

\* 4. *M. hemipterus*: Ferrugineus, opacus, capite nigro. — Long. 1 lin.

Er. Col. March. I. 645. 3.

*Silpha hemiptera* Illig. Schneid. Mag. V. 597. 5. — Panz. Faun. Germ. 25. 6.

*Silpha hemiptera* var. *β* Illig. Käf. Pr. 355. 1.

*Omal. depressum* var. *c* Gyll. Ins. Suec. II. 699 11.

*Staph. melanocephalus* Oliv. Ent. III. 42. 38. 55. t. 4. f. 52.

*Phloeobium nitiduloides* Boisd. et Lacord. Faun. Ent. Paris. I. 493. 2.

*Megarthus rufescens* Steph. Illustr. V. t. 127. f. 5. (mas)

Habitat in Europa.

Praecedentibus paululum maior, ferrugineus, opacus, subtiliter griseo-pubescens. Antennae testaceae, apice fuscescentes. Caput parvum, ruguloso-punctatum, utrinque late impressum, margine anteriore acute elevato, nigrum, ore ferrugineo. Thorax basi coleopterorum latitudine, latitudine plus duplo brevior, antrorsum angustatus, basi subtrisinuatus, apice

leviter emarginatus, lateribus subrotundatis, angulis anterioribus obtusis, subrotundatis; posterioribus leviter emarginatis, denticulo anteriore obtuso, posteriore minimo, acutiusculo, disco leviter convexus, subtilius punctatus, sat profunde canaliculatus, margine laterali late explanato, medio late impresso. Scutellum crebre punctatum. Elytra thorace duplo fere longiora, sat fortiter dense rugoso-punctata, secundum marginem lateralem impressa. Abdomen punctulatum.

Pedes maris femoribus posterioribus leviter incrassatis, tibiis intermediis leviter incurvis, simplicibus, posticis interne sinuatis, medio dentatis.

### **Phloeobium Dej.**

*Megarthrus* \*\* *Er. Col. March. I. p. 646.*

*Antennae 11-articulatae, articulo ultimo maiore.*

*Frons ocello singulo.*

*Tarsi 5-articulati.*

Corpus parvum, latiusculum, subdepressum, alatum. Caput thorace parum angustius, margine laterali et antico dilatato, hoc truncato, oculis globosis, prominulis, ocello singulo in media fronte. Labrum deflexum. Mandibulae minutae, muticae. Maxillae mala interiore cornea, apice acuminata, exteriori coriacea, apice membranosa barbataque, interiorem haud superante. Palpi maxillares articulo primo minimo, secundo sat magno, incrassato, tertio hoc paulo brevior et tenuior, quarto iterum tenuior, at secundo longitudine aequali, acuminato. Labium mento latitudine paulo brevior, antrosum sensim angustato, apice subtruncato, ligula membranosa, biloba, paraglossis latis, ligulam haud superantibus. Palpi labiales breviusculi, articulo primo brevi, secundo oblongo, cylindrico, tertio hoc paulo brevior, tenui, acuminato. Antennae sub frontis margine dilatato insertae, tenues, articulo primo clavato secundoque oblongo-obconico paulo crassioribus, penultimis tribus leviter incrassatis, ultimo oblongo-ovato, crassiusculo. Thorax transversus, coleopterorum latitudine, lateribus late marginatus. Elytra abdominis nonnisi summam basin obtegunt, apice truncata, angulo exteriori subrotundato. Abdomen breviusculum, latiusculum, apice rotundato-attenuatum, late marginatum. Pedes breviusculi, coxis intermediis approximatis, tibiis muticis, tarsis 5-articulatis, articulis 4 primis brevibus, aequalibus, quinto his coniunctis longitudine subaequali.

### **O b s e r v a t i o n e s.**

1. Genus in hac familia ocello unico singulare, praecedenti generi propinquum, at corpore minus depresso capitis structura, thorace minus distincte canaliculato et angulis posterioribus minus evidenter excisis, antennis longioribus, ligula minus brevi distinctum. Nomen huic Dejeanium servavi.

2. Sexus differentia ut in praecedente.

3. Victus sub arborum cortice.

\* 1. *Ph. clypeatum*: Testaceum, opacum, capitis margine anteriore dilatato, angulato, thorace basi utrinque denticulato. — Long. 1 lin.

*Silpha clypeata* Müll. *Germ. Mag. IV. 204. 12.* — *Germ. Faun. Ins. Europ. V. 5.*

*Megarthrus clypeatus* *Er. Col. March. I. 646. 4.*

*Phloeobium corticale* *Boisd. et Lacord. Faun. Ent. Paris. I. 494. 5.*



Habitat in Germania, Gallia.

Megarthro hemiptero paulo minor, praesertim angustior, totus testaceus, opacus, subtiliter pubescens. Antennae nigrae. articulo ultimo testaceo. Caput sat magnum, profunde punctatum, fronte leviter convexa, antice leviter excavata, margine dilatato, antice truncato, lateribus bis inciso. Thorax coleopteris haud angustior, basi utrinque subsinuatus, medio magis prominente, ante angulum anteriorem denticulo minuto instructus, apice subsinuatus, lateribus parum rotundatis, angulis omnibus subrectis, parcius punctatus, minus profunde canaliculatus, medio convexus, lateribus late explanatis, pone medium impressis, margine obsolete crenulato. Scutellum parce punctatum. Elytra thorace plus sesqui longiora, leviter convexa, sat profunde minus crebre punctata, secundum marginem lateralem subtilissime crenulatum medio impressa. Abdomen obsolete punctulatum.

Pedes maris femoribus posterioribus vix incrassatis, tibiis intermediis incurvis.

### Glyptoma.

*Thoracophorus Motschoulsky Bull. d. l. Soc. Imp. de Moscou. 1837. V. 98.*

*Antennae 11-articulatae, liberae.*

*Tarsi 3-articulati.*

*Abdomen immarginatum.*

Corpus minutum, oblongum, lineare, alatum. Caput porrectum, thoracis latitudine, suborbiculatum, basi modice constrictum, oculis parvis, subrotundatis, haud prominulis. Mandibulae haud prominentes. Palpi filiformes. Mentum trapezoidum, antrorsum angustatum. Antennae 11-articulatae, ant totae aut apice incrassatae. Thorax coleopterorum latitudine, marginatus, basi coarctatus. Scutellum triangulare. Elytra truncata. Abdomen immarginatum. Pedes breves, intermediis basi tenui spatio distantes, tibiis omnibus simplicibus, tarsis triarticulatis, articulis duobus primis brevissimis.

### Observationes.

1. Corpus opacum, plerumque glabrum. Thorax impressionibus et elevationibus inaequalis: elevationes longitudinales costas saepe referunt. Elytra costis longitudinalibus sculpta. Abdomen depresso-cylindricum, aequale.

2. Tarsis triarticulatis et corporis sculptura Micropeplo affinis, a quo autem corpore lineari, elytris metathoracem vix excedentibus, abdomine immarginato, antennis distincte 11-articulatis, liberis, etc. satis discrepat. Ab Oxytelinis coxis anticis haud exsertis et a Piestinis, quibus in multis propinquum videtur, tarsis triarticulatis differt. Habitus Colydiorum genera quaedam in animum revocat, at abdominis structura prohibet, quo minus pro Colydio quodam, Bitomae affini, elytris abbreviatis, habeatur.

3. Nomen generis, a Dom. Motschoulsky propositum mutandum putavi, minus ob cacophoniam — Thoracophorus enim vocandum esset — quam quia Thorax terminus est partis, qua Insecta omnia instructa sunt. Cf. Linn. Phil. Bot. §. 250. Fabr. Phil. Ent. §. 33.

4. Sexus differentia latet.

5. Victus sub arborum cortice.

\* 1. *G. crassicorne: Castaneum, rude punctatum, elytris unicastatis, antennis crassis, cylindricis.* — Long.  $1\frac{1}{2}$  lin.

T. II. f. 9.

Habitat Novae Valentiae in Columbia, Dom. Moritz.

Corpus elongatum, castaneum, opacum, tenuiter pubescens. Antennae capitis thoracisque longitudine, crassae, articulis 2 — 10 aequalibus, ultimo apice obtuse acuminato, praecedentibus haud crassiore. Caput thoracis latitudine, suborbiculatum, fronte convexa, immarginata, crebre sat fortiter punctata. Thorax subcordatus, lateribus anterioribus rotundatis, postice coarctatus, leviter convexus, punctis creberrimis grossiusculis cicatricosis, postice utrinque foveolatus. Scutellum minutum, triangulare. Elytra thorace sesqui longiora, confertim punctata, punctis grossis exterioribus profundis, interioribus obsoletioribus, costa singula longitudinali media elevata integra. Abdomen dense, grosse, minus profunde punctatum. Pedes rufi.

Obs. Antennarum structura a reliquis speciebus satis differt, at reliquae notae omnino conveniunt, quare rite sub eodem genere militare videtur.

- \* 2. *G. corticinum*: Castaneum, thorace lateribus crenulato elytrisque costatis, antennis articulis ultimis 5 crassioribus. — Long. parum ultra 1 lin.

*Thoraxophorus corticinus* Motschoulsky Bull. d. l. Soc. Imp. des Nat. de Moscou. 1837. V. p. 98. t. 7. f. A.

Habitat in Italia, Dom. Schüppel, Lutetiae collect. Dom. Aubé, qui specimen unicum in trunco fagi inter pusillas Formicas legit. Varsoviae etiam ad radices arborum sub cortice a D. D. Waga et Motschoulsky lectum.

Saturate rufo-ferrugineum. Antennae capite thoraceque paululum breviores, crassiusculae, articulis ultimis 5 crassioribus, 7 — 10 transversis, apicali apice obtuse acuminato. Caput thorace paulo angustius, fronte lateribus acute marginata, inter oculos longitudinaliter 4-carinata, carinulis interioribus postice duplicatis. Thorax coleopteris paulo angustior, latitudine paulo brevior, marginatus, postice subito angustatus, angulis anterioribus dentiformibus, prominulis, subdepressus, supra sexcostatus, costis externis submarginalibus, interstitiis vage rugulosis, margine laterali subtiliter crenulato. Scutellum maiusculum, medio impressum. Elytra thorace paulo longiora, 5-costata, costis alternis elevatioribus, extima marginali, interstitiis vage elevato-rugulosis. Abdomen subtiliter rugulosum, segmento sexto laevi, testaceo. Pedes rufi.

- \* 3. *G. costale*: Castaneum, thorace lateribus integro elytrisque costatis, antennis brevibus, apice subclavatis. — Long. vix ultra 1 lin.

Habitat in America septentrionali.

Statura praecedentis, at paulo minus. Castaneum, opacum. Antennae capite paulo longiores, articulis tribus ultimis modice incrassatis, clavam oblongam minus distinctam formantibus. Caput thoracis prope latitudine, marginatum, lineis pluribus elevatis longitudinalibus sculptum, clypei spatio laevi. Thorax coleopterorum fere latitudine, subquadratum, angulis anterioribus acutiusculis, prominulis, posterioribus oblique truncatis vel submarginatis, truncaturae angulo posteriore denticuli instar prominulo, subdepressus, costis longitudinalibus 6 sat elevatis sculptus, costis externis submarginalibus, reliquis 4 discoidalibus, interstitiis singulis linea singula longitudinali elevata subtili notatis. Elytra costata, costis alternis obsoletio-

ribus, alternis (3 dorsalibus 1-que laterali) elevatioribus, interstitiis punctulatis. Abdomen segmentis 5 prioribus apice granulosis, sexto laevi.

- \* 4. *G. denticolle*: *Castaneum, thorace inaequali, lateribus quadridentato, elytris costatis, antennis minutis, clavatis*. — *Long. parum ultra 1 lin.*

Habitat in ins. Puerto-Rico, Dom. Moritz.

Corpus testaceo-ferrugineum, capite, thorace abdomineque supra nigricantibus. Antennae capite perparum longiores, articulo primo longiore cylindrico secundoque crassioribus, 3 — 8 tenuibus, arcte contiguis, inter se subaequalibus, ultimis tribus incrassatis, clavam oblongam formantibus. Caput thoracis longitudine, fronte lateribus et antice marginata, medio bituberculata, clypei spatio linea arcuata impressa terminato. Thorax coleopteris paululum angustior, latitudine brevior, subdepressus, costis, tuberculis foveolisque inaequalis, margine laterali dilatato, 4-dentato, postice subito angustato. Elytra regulariter costata, costis tribus dorsalibus, quarta infra marginem lateralem, interstitiis subtiliter punctulatis. Abdomen subtilissime punctatoprugulosum.

5. *G. exile*: *Rufo-ferrugineum, thorace inaequali, lateribus integro, elytris tricostatis, antennis brevibus, clavatis*. — *Long.  $\frac{3}{4}$  lin.*

Habitat in Americae merid. ins. St. Thomae, Dom. Moritz.

Praecedentibus plus duplo minus, testaceo-ferrugineum, opacum. Antennae capite sesqui longiores, articulis ultimis tribus distincte maioribus, clavam oblongam formantibus. Caput thoracis fere latitudine, lateribus acute marginatum, fronte plagis duabus longitudinalibus elevatis. Thorax coleopterorum prope latitudine, angulis anterioribus rectis, haud prominulis, lateribus anterieus rectis, dein pone medium subito sinuato-angustatis, subdepressus, modice inaequalis, lineis 5 sat subtilibus elevatis longitudinalibus sculptus, externis submarginalibus. Elytra thorace sesqui longiora, costata, costis tribus dorsalibus, quarta laterali, interstitiis laevibus. Abdomen laeve.

6. *G. sculptile*: *Fuscum, capite nigro, thorace rufo, quadrisulcato, lateribus integro, elytris multicostatis, antennis subclavatis*. — *Long.  $\frac{3}{4}$  lin.*

Habitat in valle Araguensi Columbiae, Dom. Moritz.

Minutum, praecedenti aequale, at paulo gracilius. Antennae capite sesqui longiores, tenuiusculae, articulis ultimis tribus distincte maioribus, testaceae. Caput thoracis latitudine, subtiliter marginatum, fronte subdepressa, lineis duabus elevatis obsoletioribus longitudinalibus notata, nigrum, opacum. Thorax antice coleopteris parum angustior, angulis anterioribus rectis, haud prominulis, lateribus anterieus leviter rotundatis, pone medium constrictis, depressus, sulcis quatuor longitudinalibus sat latis, interioribus rectis, antice abbreviatis, exterioribus profundioribus, integris, repandis inaequalis, fuscorufus, opacus. Elytra thorace sesqui longiora, striata, interstitiis elevatis, alternis elevatioribus, fusca, opaca. Abdomen laeviusculum, fuscum, opacum, segmentis singulis margine apicali testaceo. Pedes rufo-testacei.



**Micropeplus Latr.***Latr. Gen. Crust. et Ins. IV. p. 377.*

*Antennae receptae, 9-articulatae, capitulatae.*  
*Tarsi 3-articulati.*

Corpus minutum, subovatum, crassiusculum, alatum. Caput a thorace receptum eodemque minus et angustius, subtriangulare, oculis mediocribus, semiglobosis, modice prominulis. Labrum transversum, integrum, corneum. Mandibulae parvae, muticae, haud prominentes. Maxillae (t. V. f. 35. b.) mala interiore angusta, cornea, exteriore trapezoidea, apice dilatata, rotundata, barbata. Palpi maxillares articulo primo tenui, minuto, secundo sat magno, fortiter incrassato, tertio brevi transverso, quarto conico. Labium (t. V. f. 35. a.) mento sat magno, ante apicem constricto, ligula brevi, angustiore, rotundata, membranea, palpis brevissimis, articulis decrescentibus, ultimo fere obsoleto. Antennae 9-articulatae, articulis duobus primis crassiusculis, primo maiusculo, ovato, secundo hoc minore, apice attenuato, sequentibus tribus tenuibus, elongatis, penultimis tribus haud crassioribus at brevibus, ultimo maximo, globoso, capitulum antennae globosum formante. Thorax transversus, basi coleopterorum latitudine, iisque arcte applicatus, antrorsum angustatus, lateribus late explanato-marginatis, subtus excipiendis antennis secundum marginem lateralem profunde scrobiculatis. Scutellum sat magnum, triangulare. Elytra ampla, at nonnisi abdominis summam basin obtegentia. Abdomen acuminatum. Pedes breves, tenues, simplices, intermedii basi distantes, tarsis brevibus, triarticulatis.

## O b s e r v a t i o n e s.

1. Corporis habitus singularis sculptura singulari, praeterea antennae capitulatis, receptis et pedibus intermediis late distantibus insigne hoc genus, a multis auctoribus perperam *Nitidulis* adscriptum.

2. Antennarum capitulum non solidissimum sed articulis tribus connatum videtur, unde forte antennae revera 11-articulati haberi possint.

3. Tarsi a Dom. Curtis loco apud M. tesserulam citato 4-articulati descripti, mihi autem non ultra tres articuli observati sunt.

4. Mas abdominis segmento inferiore sexto apice emarginato distinctus.

5. Victus in quisquiliis.

\* 1. *M. porcatus*: *Niger, antennarum basi pedibusque rufis, elytris basi leviter convexis, 4-costatis, interstitiis crebre punctatis.* — Long. 1 lin.

*Charp. Hor. Ent. p. 202. t. 8. f. 9. — Er. Col. March. I. 646. 1.*

*Staph. porcatus Ent. Syst. I. II. 530. 56. Syst. El. II. 602. 68. — Payk. Mon.*

*Staph. 71. 50. Faun. Suec. III. 413. 59. — Oliv. Ent. III. 42. 35. 50. t. 4. f. 33.*  
*— Panz. Ent. Germ. 368. 45.*

*Nitidula porcata Marsh. Ent. Britt. 137. 26.*

*Omalium porcatum Gyll. Ins. Suec. II. 211. 12.*

*Nitidula sulcata Herbst Col. V. 247. 22. t. 54. f. 6. — Panz. Ent. Germ. I. 129. 31. — Schönk. Syn. II. 148. 64.*

Habitat in Europa.

*Niger*, opacus. Antennae rufae, capitulo piceo. Caput confertim subtiliter rugulosum, fronte depressa, antice posticeque linea transversa, medio linea longitudinali impressa, margine tenuiter elevato, prope oculos angulato. Thorax basi coleopterorum fere latitudine, longitudine duplo fere latior, an-

trorsum valde angustatus, lateribus medio obtuse angulatis, angulis posterioribus acutiusculis, disco modice convexo, lineis elevatis impresso-areolato, areolis subtilissime confertissimeque rugulosis, lateribus late explanatis, medio puncto diaphano piceo, margine elevato. Scutellum laeve. Elytra thorace sesqui longiora, leviter convexa, apice depressa, sutura costisque quatuor elevatis, costis exterioribus duabus apice paulum abbreviatis, interstitiis lateribusque crebre profunde punctatis, his costis duabus elevatis, inferiore marginali, arcuata. Abdomen supra segmentis 2—5 4-foveatis, foveis magnis, profundis, carinis valde elevatis angustis divis, sexto confertim subtiliter punctulato. Pedes rufi.

\* 2. *M. caelatus*: *Niger, antennarum basi pedibusque rufis, elytris basi convexis, 4-costatis, interstitiis crebre punctatis.* — Long.  $\frac{4}{5}$  lin.

*Er. Col. March* I. 647. 2.

Habitat in Germania.

M. porcato minor, praecipue brevior, niger, opacus. Antennae rufae, capitulo piceo. Caput omnium confertissime subtiliter punctato-rugulosum, fronte haud depressa, linea longitudinali levi tenuique transversa antica impressum. Thorax basi coleopterorum latitudine, longitudine plus duplo lator, antrorsum valde angustatus, lateribus medio subangulatis, disco convexo, confertissime subtiliter punctato-rugoso, rugis elevatis parum distinctis areolato, lateribus late explanatis, margine elevato. Scutellum laeve. Elytra thorace sesqui latiora, elevato-convexa, apice depressa, dorso costis 4, lateribus 2, suturaque elevatis, interstitiis crebre fortiterque punctatis, ut in M. porcato. Abdomen segmentis 2—4 4-foveatis, foveis magnis, profundis, carinis angustis divis, quinto sextoque confertissime rugoso-punctatis, illo basi 4-foveolato. Pedes rufi.

Obs. M. porcato valde affinis, differt autem statura brevior, elytris magis elevatis, abdominis segmento quinto basi tantum foveolato, thorace obsolete cancellato etc.

3. *M. fulvus*: *Testaceus, elytris subelevatis, tricostatis, interstitiis punctatis.* — Long. 1 lin.

Habitat Lutetiae, Mus. Dom. Chevrolat.

Testaceus, immaculatus. Caput inter antennarum basin transversim impressum, fronte posteriorius lineis nonnullis elevatis subtilibus longitudinalibus sculpta, apice in altero sexu acuminata, in altero obtusa. Thorax latitudine coleoptera parum excedens, latitudine summa duplo fere brevior, antrorsum angustatus, angulis posterioribus rectis, disco leviter convexo, lineis elevatis cancellato-areolato, lateribus late explanatis, margine elevato. Elytra subelevata, apice depressa, sutura costisque tribus dorsalibus, extima marginali, elevatis, interstitiis crebre sat profunde punctatis, lateribus sat crebre profundeque punctatis, costis duabus elevatis, superiore recta, costae marginali aequali, inferiore arcuata, marginali, itidem sat fortiter elevata. Abdomen segmentis 2—4 basi 4-foveolatis, foveolis transversis, sat profundis, interstitiis angustissimis discretis, quinto basi 4-foveolato, interstitio medio elevato, tuberculiformi.

Obs. M. Staphilinoidi perquam affinis, at differt thorace minus dilatato, abdominis segmenti quinti tuberculo longe minus prominulo, elytris minus depressis, margine laterali bicarinato, carina laterali superiore haud, ut in illo, obsoleta.

- \* 4. *M. Staphylinoides*: *Piceus, thoracis lateribus, antennis pedibusque rufis, elytris 3-costatis, interstitiis punctatis.* — Long. 1 lin.

*Nitidula Staphylinoides* Marsh. Ent. Britt. 137. 25.

*Micropepl. Maillei* Guér. Iconogr. Ins. t. 10. f. 4.

Habitat in Anglia, Dom. Spence, in Gallia Dom. Aubé, in Italia, Dom. Schüppel.

*M. porcato* paulo minor et angustior, piceus, nitidus. Antennae rufae. Caput posterius subtiliter rugulosum, antice laeve, fronte depressa, linea longitudinali anticaque transversali impressa, margine elevato, apice in aliis individuis acuminato, in aliis obtuso. Thorax coleopteris fere lator, latitudine plus duplo brevior, antrorsum valde angustatus, lateribus medio subangulatis, angulis posterioribus acutis, disco leviter convexo, lineis elevatis cancellato-areolato, piceo, lateribus late explanatis, rufis, margine leviter elevato. Scutellum obsolete impressum. Elytra thorace paulo longiora, subdepressa, costis tribus dorsalibus suturaque elevatis, interstitiis crebre sat profunde punctatis, lateribus sat crebre profunde punctatis, costis duabus elevatis, superiore recta, subtili, obsoleta, inferiore arcuata, marginali, sat fortiter elevata. Abdomen segmentis 2 — 4 4-foveatis, foveis magnis, profundis, carinis tribus divis, carina intermedia magis elevata, quinto sextoque laevibus, illo basi 4-foveolato, carina valde elongata a basi ad medium percurrente, apice acuminata cristato. Pedes rufi.

- \* 5. *M. tesserula*: *Niger, antennarum basi pedibusque rufis, elytris 3-costatis, interstitiis laevibus.* — Long.  $\frac{2}{3}$  lin.

*Curtis Ent. Britt. V. t. 204.*

*Omal. Staphylinoides* Gyll. Ins. Suec. II. 213. 13. — *Sahlb. Ins. Fenn. I. 280. 10.*

Habitat in Suecia; in Carinthia, Dom. Schmidt.

*M. porcato* duplo minor, niger, nitidulus. Antennae nigrae, basi rufae. Caput fronte depressa, longitudinaliter trisulcata, sulco intermedio profundiore, antice linea transversali impressa, margine laterali tenuiter elevato. Thorax coleopterorum latitudine, latitudine duplo brevior, antrorsum valde angustatus, angulis posterioribus subrectis, disco parum convexo, lineis elevatis cancellato-areolato, lateribus late explanatis, piceis, puncto medio pelucido, margine subelevato. Scutellum laeve. Elytra thorace paulo longiora, leviter convexa, apice depressa, costis tribus dorsalibus suturaque elevatis, interstitiis lateribusque laevibus, his unicostatis, costa arcuata, marginali, sat elevata. Abdomen segmentis 2 — 4 minus fortiter 4-foveatis, ultimis duobus laevibus. Pedes rufi.

### Species mihi invisae.

6. *M. obtusus*: *Testuceus, capite antice rotundato.*

*Newman Ent. Magaz. II. 201.*

Habitat prope Halifax.

Totus pallide brunneus. *M. Staphylinoides* minus elongatus, at paulo lator, elytrorum costis magis elevatis et capite antice non acuminato sed rotundato distinctus.

Obs. *M. fulvo* affinis, at distinctus videtur.



## O b s e r v a t i o.

Huic forte tribui inseratur genus, ab auctore hoc modo descriptum :

**Pseudopsis Newman.**

*Newman Ent. Mag. II. p. 313.*

Caput elongatum, angustum, antice rotundatum, epicranio valde depresso, partibus lateralibus supra oculos iterumque parte media longitudinali elevatis, clypeo prono, rotundato; antennae filiformes, 11-articulatae, extus incrassatae, articulo apicali conico; palpi maxillares articulo apicali elongato, acuto, tenuissimo, proximo quadruplo maiore; alia instrumenta cibaria haud examinata. Prothorax depressus, fere circularis, sed antice et postice paulo truncatus, lateribus lineisque 4 disci longitudinalibus perspicue elevatis. Elytra valde depressa, marginibus, sutura lineisque 2 singuli disci longitudinalibus perspicue elevatis. Abdominis segmenta 7 nuda, haud elytris tecta, media depressa, lateribus elevata, apicem versus magnitudine pedetentim decrescentia, apice elongato, angusto.

*Ps. sulcatus: Niger, ore, antennis pedibusque fuscis.*

Caput nigrum, ore fusco, fronte depressa, linea utrinque laterali supra oculos tertiaeque inter oculos longitudinalibus elevatis, hac obsoletiore et minus regulari. Antennae fuscae, moniliformes, 11-articulatae, articulo apicali subconico, penultimis transversis, basin versus sensim minoribus. Palpi maxillares articulo apicali longo, tenui, acuminato, praecedente incrassato, apicali quadruplo longiore. Thorax capite latior, depressus, margine laterali lineisque quatuor longitudinalibus omnino rectis disci evidenter elevatis, interiectis sulcis 5. Coleoptera thorace latiora, depressa, margine, sutura lineisque duabus evidenter elevatis, lineis non omnino rectis, sed apicem versus sensim suturae magis approximatis. Abdomen segmentis septem omnino liberis, medio depressis, lateribus elevatis, sensim decrescentibus, ultimo elongato, angusto. Thorax, elytra et abdomen obscure nigra. Pedes fusci. — Long.  $1\frac{1}{2}$  lin. — Captus in insula Vectis a Dom. Walker.

## ADDENDA ET EMENDANA.

### Myrmedonia.

p. 41.

#### 10. *M. obscura*:

Secundum specimen a Dom. Westermann amice communicatum accuratius describenda:

*Rufo-testacea. crebrius punctata, capite abdomineque nigris, hoc segmento superiore primo apice trispinoso. — Long. 3 lin.*

Antennae capite vix duplo longiores, crassae, articulo secundo tertio duplo brevior, 4 — 10 transversis, ultimo conico, ferrugineae, articulis 3 primis rufo-testaceis. Caput thoracis latitudine, fronte planiuscula, alutacea, obsoletius punctata, nigrum, minus nitidum, ore testaceo. Thorax coleopteris paulo angustior, latitudine vix brevior, basin versus subangustatus, lateribus leviter rotundatus, subdepressus, sat crebre fortius punctatus, subtiliter fulvo-pubescent, rufo-testaceus, nitidus. Elytra thoracis longitudine, crebre fortius punctata, subtiliter fulvo-pubescentia, rufo-testacea, nitida. Abdomen supra parce, infra crebre punctatum, supra segmento primo apice producto, spinis tribus sat elongatis, acutis, lateralibus paulo longioribus, terminato, quinto medio obsolete tuberculato, sexto apice obsolete tridentato, nigrum, nitidum, segmentis singulis inferioribus piceo-marginatis, ano rufo-piceo. Pectus rufo-testaceum. Pedes testacei.

Obs. Proxima certe *M. spinigerae*, sed distincta punctura fortiore, thorace longiore et abdominis structura, neque sexu solo differens. Femina utriusque speciei latet.

### Falagria.

p. 52.

#### 10. *F. thoracica*: adde

*Aleocho. ruficollis* Waltl Isis. 1838. IV. 268. 4.

**Tachyusa.**

p. 73 adde

- \* 7 — 8. *T. uvida*: *Nigra, subnitida, antennis pedibusque testaceis, elytris fuscis, abdomine lineari, thorace subquadrato fronteque late canaliculatis.* — Long.  $1\frac{1}{2}$  lin.

Habitat in insula Vectis frequens ad littora maris sub fucis putrescentibus, Dom. Rudd.

Statura omnino et summa affinitate Tachysae laesae, nigra, subnitida, densius subtiliusque griseo-pubescent. Antennae capite thoraceque longiores, apicem versus perparum incrassatae, totae testaceae. Caput thoracis latitudine, suborbiculatum, dense evidenter licet subtiliter punctatum, depressiusculum, fronte late, feminae leviter, maris profunde canaliculata. Thorax coleopteris dimidio angustior, latitudine haud brevior, basin versus leviter angustatus, lateribus rectis, basi leviter rotundatus, apice utrinque oblique truncatus, angulis anterioribus fortiter rotundatis, subdepressus, dense subtiliter punctatus, secundum totam longitudinem in mare late profundeque in femina levius canaliculatus. Elytra thoracis latitudine, subdepressa, confertissime subtiliter punctata. Abdomen lineare, nitidum, minus subtiliter punctatum, parcius pubescens, apice summo fulvo-tomentoso, segmentis aequalibus. Pedes testacei.

Obs. Distincta a *T. laesa* punctura et pube minus subtilibus, fronte in utroque sexu canaliculata, thorace magis canaliculato quam excavato, antennis totis testaceis etc.

p. 74 adde

- \* 9 — 10. *T. chalybea*: *Nigro-cyanea, nitidula, subtilissime griseo-pubescent, thorace subquadrato, leviter canaliculato, tarsis testaceis.* — Long.  $1\frac{3}{4}$  lin.

Habitat in ins. Vectis. Nomine Ischnopodae chalybeae a Dom. Rudd communicata.

Habitu fere *T. umbraticae*, at paulo maior, corpore cyanescenti-nigro, capite, thorace elytrisque cyaneis, nitidula, omnium subtilissime punctulata, pube subtilissima fusco-sericea obducta. Antennae capite thoraceque paulo longiores, tenues, apicem versus leviter incrassatae, articulo tertio secundo paulo longiore, nigrae. Caput thoracis prope latitudine, suborbiculatum, basi fortiter constrictum, fronte media leviter impressa, ore nigro. Thorax coleopteris dimidio angustior, latitudine haud brevior, neque basin neque apicem versus angustatus, lateribus anterioribus leviter, angulis anterioribus fortiter rotundatis, apice truncatus, basi leviter rotundatus, subdepressus, canalicula parum profunda, anterioribus obsolescente impressus. Coleoptera thorace paulo longiora. Abdomen coleopteris paulo angustius, parallelum, segmentis subaequalibus, anterioribus basi profundius transversim impressis. Pedes nigri, tarsis testaceis.

Obs. Forte a *T. coerulea* (p. 76. n. 14.) haud distincta, at huic frons impressa, et thorax canaliculatus.

- \* 10 — 11. *T. immunita*: *Nigra, subtilissime griseo-pubescent, elytris fuscis, pedibus testaceis, thorace subquadrato, maris late leviter impresso, feminae basi obsolete foveolato, abdomine parallelo.* — Long.  $1\frac{1}{2}$  lin.

Habitat in Anglia, Dom. Rudd.

*Hemiteles yagor*



Statura et summa similitudine Homalotae elongatae, subdepressa, minus gracilis, nigra, minus nitidula, dense subtilissime griseo-pubescent. Antennae capitis thoracisque longitudine, minus tenues, apicem versus sensim subincrassatae, articulo secundo tertioque subaequalibus, fuscae, articulis basalibus nigricantibus. Palpi fuscii. Caput thoracis latitudine, orbiculatum, fronte leviter convexa, omnium subtilissime punctulatum. Thorax coleopteris paulo angustior, latitudine haud brevior, neque basin neque apicem versus angustatus, lateribus angulos anteriores versus parum rotundatis, subdepressus, subtilissime dense punctatus, iam dorso late leviter, iam basi obsolete impressus. Elytra thorace paulo longiora, dense subtilissime punctata. Abdomen parallelum segmentis subaequalibus, superioribus 4 anterioribus subtilissime punctulatis, 2 posterioribus sublaevibus. Pedes testacei.

Obs. Anceps in hoc genere, Homalotam elongatulam statura, colore et punctura simulat, at tarsorum posticorum articulo primo satis elongato instructa. Thorace vel late impresso vel subaequali forte sexu variat.

### Homalota.

p. 93.

23. *H. plana*: Eandem ex America septentrionali misit Dom. Zimmermann.

p. 130. adde

- \* 108 — 109. *H. paedida*: Nigra, opaca, dense subtiliter punctata, fusco-pubescent, antennis longis, crassis, nigris, pedibus elytrisque luteis, his circa scutellum et lateribus infuscatis. — Long.  $1\frac{1}{4}$  lin.

Habitat in Madagascar, Dom. Goudot.

Statura et summa affinitate *H. lividipennis*, at plus duplo minor et magis depressa, nigra, dense subtiliter punctata, pube subtili depressa fusca obducta. Antennae capite thoraceque paulo longiores, validae, articulo secundo tertioque subaequalibus, obconicis, 4—10 cylindricis, crassitie haud brevioribus, ultimo praecedentibus ambobus aequali, subacuminato, nigrae. Caput thorace angustius, fronte leviter convexa. Thorax basi coleopterorum latitudine, antrosum leviter angustatus, latitudine dimidio brevior, lateribus modice rotundatus, apice leviter emarginatus, angulis posterioribus obtusis, anterioribus rotundatis, deflexis, subdepressus, aequalis. Coleoptera thorace paululum longiora, depressa, sordide lutea, circa scutellum et lateribus fuliginosa. Abdomen apicem versus angustatum, apice pilosellum, infra segmentis sinuatis apice late testaceo-marginatis. Pedes flavi.

### Oxypoda.

p. 151. adde

- \* 21 — 22. *O. contractula*: Brevis, nigra, subnitida, subtiliter pubescens, dense punctata, antennis pedibusque ferrugineis, thorace transverso, basi bisinuato, apice rotundato. — Long. 1 lin.

Habitat in Madagascar, Dom. Goudot.

*O. curtulae* affinis et eadem haud longior, at duplo prope latior, Aleochara morione latior et convexior, habitu fere Tachypori, nigra, nitidula, densius pube subtili fusca sericante vestita, dense minus subtiliter punctata.

Antennae capite thoraceque paulo longiores, articulo secundo tertio maiore, praecipue crassiore, apicem versus vix incrassatae, articulo ultimo maiusculo, ovato, rufo ferrugineae, articulo ultimo piceo. Caput inflexum, ore leviter producto, testaceo. Thorax coleopteris latitudine aequalis, latitudine baseos plus dimidio brevior, basi profundius bisinuatus, angulis posterioribus acutis, lateribus et apice eodem arcu rotundatus, convexus, obsolete piceo-marginatus. Coleoptera thorace vix longiora, apice late emarginata, dorso depressiuscula, minus subtiliter punctata. Abdomen breviusculum, apicem versus fortius angustatum, acuminatum, segmentis piceo-marginatis, lateribus nigro-setulosum. Pedes rufo-ferruginei.

Obs. Ore producto in hoc genere aliena, et Gymnusae vel Myllaenae propior videtur, at palpi labiales distincte articulatae. Habitus etiam potius Aleocharae quam Oxypodae prae se fert, at palpis labialibus triarticulatis, articulo tertio penultimo duplo longiore ab illo genere distat. Thorace basi sinuato, punctura et imprimis antennarum structura *O. curtulae* proxima.

### Aleochara.

p. 160. adde

- \* 3 — 4. *A. loricata*: *Rufa, nitida, abdomine nigro, ano pedibusque testaceis, capite, thorace elytrisque parce profundeque, abdomine crebrius punctatis.* — Long. 2 lin.

Habitat in Guinea; specimen singulum in Formicae cuiusdam nido repertum amicissime dedit Dom. Westermann.

Antennae breves, capite sesqui longiores, articulo tertio secundo sesqui prope longiore, apicem versus leniter incrassatae, subcompressae, articulis 6 — 10 transversis, ultimo breviter ovato, apice acuminato, totae testaceae. Caput thorace plus duplo angustius, fronte punctis paucis maioribus setuliferis notata, ceterum laevissimum, rufo-piceum, ore testaceo. Thorax amplius, coleopteris paulo latior, latitudine duplo prope brevior, antrorsum leviter angustatus, lateribus et angulis omnibus rotundatis, apice emarginatus, basi subbisinuatus, fortius transversim convexus, punctis maiusculis setuliferis sparsis, latera versus crebrioribus, interstitiis laevissimis, rufus, nitidus. Scutellum piceum. Coleoptera thoracis longitudine, apice late emarginata, angulo apicali exteriore rotundato, punctis raris grossiusculis setuliferis sparsis notata, interstitiis laevissimis, rufo-testacea, limbo laterali et apicali picescente. Abdomen crassiusculum, apicem versus subangustatum, segmentis quatuor anterioribus crebrius punctatis, setulis depressis nigris vestitis, nigris, nitidis, ultimis duobus laevibus, testaceis. Pectus rufum. Pedes testacei.

### Placusa.

p. 196.

4. *P. infima*: *Forte huc referatur:*

*Aleochar. tachyporoides* Walz Isis 1838. IV. 268. 5.

### Dinarda.

p. 201.

1. *D. dentata*: *Specimina maiora minoraque commixta in Formicae rufae nidis lecta Dom. Märkel benevole communicavit.*

**Myllaena.**

p. 210.

2. *M. intermedia*: adde*Centroglossa attenuata* Matthews Ent. Mag. V. 196. 2.

p. 211.

3. *M. minuta*: adde*Centroglossa minuta* Matth. Ent. Mag. V. 196. 3.

Obs. Species reliquae tres a Dom. Matthews huic generi adscriptae thorace coleopteris latitudine aequali a praecedentibus distinctae, mihi haud extricatae, forte Oxypodae adscribendae.

**Gymnusa.**

p. 212.

\* 2. *G. laticollis*: adde*Deinopsis fuscatus* Matthews Ent. Mag. V. p. 193.

Dom. Matthews labium palpis et ligulae laciniis perditis mutilatum delineavit et palpos maxillares tarsosque perperam 3-articulatos l. c. exhibuit. Genus *Deinopsis* igitur delendum, nomen autem specificum in *G. fuscata* mutandum.

**Hypocypus.**

p. 216.

2. *H. discoideus*: adde*Cypha biguttata* Matthews Ent. Mag. V. p. 198. (forte)**Conurus.**

p. 222.

5. *C. crassus*: adde*Tachyp. moestus* Say Transact. Am. Philosoph. Soc. Philad. Nov. Ser. IV. 466. 2.

p. 226.

11. *C. cinctulus*: adde*Tachypor. opicus* Say Transact. Am. Philosoph. Soc. Philad. Nov. Ser. IV. 467. 5.Nomen in *C. opicus* mutandum.**Tachyporus.**

p. 233.

3. *T. rufus*: adde*Tachyp. formosus* Matthews Ent. Mag. V. p. 197.Nomen in *T. formosum* mutandum.

p. 237.

11. *T. arduus*: adde*Tachyp. iocosus* Say Transact. Am. Philosoph. Soc. Philad. Nov. Ser. IV. 466. 3.Nomen in *T. iocosum* mutandum.



p. 241.

18. *T. brunneus*: adde

*Tachyp. faber* Say Transact. Amer. Philosoph. Soc. Philad. Nov. Ser. IV. 468. 7.

p. 242. adde

22. *T. acandus*: *Abbreviatus, testaceus, thorace lateribus pallidiore.* — Long. infra 1 lin.

Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 467. 6.

Habitat in Indiana Americae septentrionalis.

Testaceus, politus. Caput nigro-piceum. Antennae fuscae, articulis 4 primis obscure flavescentibus. Palpi flavescentes, articulo ultimo aciculari. Thorax testaceus, disco et basi fusciscentibus, lateribus pallidioribus et fere pellucidibus. Elytra rufo-testacea, leviter pubescentia, regione scutellari fusciscente, angulo apicali suturali rotundato, unde coniuncta acute emarginata apparent. Abdomen elytris duplo brevius, nigro-aeneum, pubescens. Pedes pallide flavi. Tibiae spinulosae. — Similis *T. iocoso*, at latior, licet abdomine fortiter contracto. (Say.)

23. *T. acuductus*: *Nigro-piceus, thoracis angulis posterioribus, antennis pedibusque testaceis, elytris minutissime acuductis.* — Long. 1 lin.

Kirby Faun. Boreali-Amer. 90. 1.

Habitat in America boreali, prope Aedes Cumberlandianas, sub lat. 54°.

Obscure piceus, glaber, laevis, nitidus. Antennae et os testacea. Thorax laevis, coleopteris latior, angulis posterioribus testaceis. Elytra capite thoraceque coniunctis longiora, picea, angulo apicali exteriori testaceo, oculo probe armato inspecta subtilissime aciculata, praecipue prope suturam, immixtis punctis minutissimis. Abdomen perbreve, conicum, vix marginatum, piceum, segmentis apice dilutioribus. (Kirby.)

24. *T. affinis*: *Nigro-piceus, nitidus, antennis pedibusque testaceis, elytris castaneis, punctulatissimis.* — Long. 1 lin.

Kirby Faun. Boreali-Amer. 91. 2.

Habitat in America boreali. Lectus cum praecedente.

*T. acuducto* similimus: thoracis angulis posterioribus haud testaceis, elytris pallide castaneis, dense subtiliter punctatis, nullo modo aciculatis distinctus. (Kirby.)

## Tachinus.

p. 252.

17. *T. gibbulus*: *Nomen in T. ventriculum mutandum et addendum:*

*Tachyp. ventriculus* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 466. 3.

p. 256. adde

\* 20—21. *T. luridus*: *Nigro-piceus, nitidus, thoracis limbo laterali et abdominis segmentorum marginibus rufo-testaceis, elytris luteo-testaceis.* — Long. 2½ lin.

Habitat in Americae septentrionalis montibus Catskill, Dom. Zimmermann.

Statura *T. flavipedis*, nigro-piceus, nitidus. Antennae capitis thoracis-que longitudine, articulo tertio secundo paulo longiore, 5 — 10 crassioribus, aequalibus, subobconicis, ultimo ovato, nigrae, articulis 4 primis piceis. Palpi picei. Caput subtilissime punctulatum. Thorax coleopterorum latitudine, latitudine baseos dimidio brevior, antrorsum leviter angustatus, apice levissime emarginatus, angulis anterioribus perparum prominentibus posterioribusque obtusis, subrotundatis, leviter convexus, dense subtilissime alutaceo-strigosus, parce subtilissime interpunctatus, margine apicali basalique tenui et laterali latiore rufescenti-testaceo. Elytra thorace sesqui longiora, parcius subtiliter punctata, luteo-testacea, nitida, postice leviter infuscata. Abdomen distinctius punctatum, pube brevi adpersum, segmentis 5 anterioribus margine apicali, sexto toto rufescenti-testaceis. Pedes piceo-testacei.

Mas latet.

Femina abdominis segmento superiore sexto trifido, laciniis lateralibus convergentibus, intermedia longiore, acuminata; inferiore integro, apice triangulariter producto.

p. 258. adde

23 — 24. *T. fulvipes*: *Niger, nitidulus, antennarum basi apiceque pedibusque testaceis, elytris thorace paulo longioribus, hoc angulis posterioribus acuminatis. — Long. 2 $\frac{2}{3}$  lin.*

Habitat in Java, Mus. Dom. Westermann.

*T. rufipede* paulo brevior et latior, depressiusculus, niger, nitidulus, capite, thorace elytris-que omnium subtilissime punctulatis. Antennae capite thoraceque breviores, articulo tertio secundo paulo brevioribus, articulis 5 — 10 crassioribus, sensim paulo brevioribus, piceis, ultimo ovato, luteo, 4 primis rufo-testaceis. Caput fronte leviter convexa, ore concolore, palpis testaceis. Thorax basi coleopterorum latitudine, latitudine baseos duplo prope brevior, angustatus, lateribus rotundatus, basi late emarginatus, angulis posterioribus retrorsum spectantibus, acutis. Elytra thorace paulo longiora, obscure brunnea, basi et sutura late nigricantibus, limbo laterali nigerrimo. Abdomen crebre minus subtiliter punctatum. Pedes rufo-testacei.

Mas latet.

Femina abdominis segmento superiore sexto quadrifido, laciniis subaequalibus, acutis.

p. 261.

29. *T. axillaris*: adde

*Tachyp. fumipennis* Say *Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV.* 464. 3.

Nomen in *T. fumipennem* mutandum.

## Trichophyus.

p. 268.

\* 1. *Tr. pilicornis*: *Specimen huius insecti rarissimi integrum Dom. Mürkel benevole dedit, ex quo antennarum descriptio hoc modo emendetur:*

Antennae capite thoraceque paulo longiores, articulo primo secundo-

que crassitie solita, primo subcylindrico, secundo obconico, reliquis omnibus aequaliter tenuibus, longitudine subaequalibus, ultimo reliquis paululum longiore, undique pilosis.

## Boletobius.

p. 280.

### 21. *B. pygmaeus*: adde

*Megacronus elegans* Matthews Ent. Mag. V. p. 197. (forte)

*Tachin. trimaculatus* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 464. 3.

Specimen ex America septentrionali, ab hac specie nullo modo distinctum Dom. Zimmermann misit.

p. 281. adde

### 23. *B. obsoletus*: Niger, thorace, elytrorum arcu, abdominis segmentorum marginibus pedibusque flavescentibus. — Long. ultra $2\frac{1}{2}$ lin.

*Tachin. obsoletus* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 464. 2.

Habitat in Virginia, mense Octobre.

Niger. Antennae fuscae, articulis 4 primis flavescentibus, ultimo apice fulvo. Os testaceum. Thorax testaceus. Elytra certo situ obsolete striata, stria suturali impressa, basi, sutura et margine apicali testaceis. Abdomen segmentis singulis apice piceis. Pedes pallide flavescentes. (Say.)

### 24. *B. cincticollis*: Niger, thoracis margine posteriore, elytrorum stria pedibusque flavescentibus. — Long. infra 2 lin.

*Tachin. cincticollis* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 465. 4.

Habitat in Indiana Americae septentrionalis.

Niger. Os flavescens, palpis pallide flavis. Antennae fuscae, articulis 3 primis flavescentibus, quarto piceo. Thorax margine basali et margine laterali basin versus dilatato testaceis. Scutellum piceum. Elytra apice substriata, stria suturali impressa et punctorum remotorum seriebus tribus, prima prope suturam, secunda in medio dorsi, tertia in margine laterali, pallide flavescentia, macula ovali, dilatata, ab apice ultra medium extensa, coeruleo-nigra, suturam haud attingente, cum linea marginali nigricante, ante humerum abbreviata confluenta. Abdomen supra segmentis singulis apice piceis. Pedes pallide flavi. — A reliquis thoracis margine posteriore discolore distinctus. (Say.)

## Mycetoporus.

p. 284.

### 4. *M. lepidus*: Hanc quoque speciem Dom. Zimmermann ex America septentrionali misit, cui forte subiungatur:

*Tachinus humidus* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 465. 5.



p. 285.

6. *M. pronus*: *Forte huc allegatur*:*Mycetop. brevicornis* Matthews Ent. Mag. V. p. 197.**Platyprosopus.**

p. 294. adde

3. *P. tamulus*: *Niger, griseo-pubescent, elytris, ano, antennis pedibusque rufis.* — Long.  $6\frac{1}{2}$  lin.

Habitat in Bengala, Mus. Dom. Westermann.

*P. elongato* laud longior, at crassior, niger, subnitidus, totus griseo-pubescent, pube capitis densiore, thoracis parciore. Antennae capite sesqui longiores, articulo tertio secundo sesqui longiore, et palpi dilute rufi. Labrum castaneum. Caput thoracis prope latitudine, oblongum, lateribus rectis, totum dense subtilius punctatum. Thorax coleopteris vix angustior, latitudine paulo longior, basin versus subangustatus, apice truncatus, basi leviter rotundatus, dense subtilius punctatus, linea media longitudinali laevi. Elytra thorace tertia parte longiora, dense subtiliter punctata, rubra. Abdomen dense subtiliter punctatum, apice attenuatum, segmentis ultimis duobus rufis. Pedes cum coxis anticis rufi.

4. *P. fuliginosus*: *Niger, fusco-pubescent, elytris fuscis, pedibus rufis.* — Long. 6 lin.

Habitat in Bengala, Mus. Dom. Westermann.

*P. elongato* crassior, niger, opacus, pube brevi depressa fusca densius aequaliter vestitus. Antennae capite thoraceque paulo breviores, articulo tertio secundo duplo prope longiore, rufae. Palpi rufo-testaceae. Caput thoracis prope latitudine, oblongo-ovatum, creberrime subtiliter punctatum. Thorax coleopterorum latitudine, latitudine paulo longior, basin versus subangustatus, apice truncatus, basi leviter rotundatus, confertissime subtiliter punctatus, obsolete subtiliter carinatus. Elytra thorace paulo longiora, dense subtiliter punctata, fusca, opaca. Abdomen apicem versus angustatum, dense subtiliter punctatum, supra nigrum, segmentis singulis apice castaneo-marginatis, penultimo apice, ultimo toto rufis, infra rufo-piceum, segmento penultimo apice, ultimo toto rufo. Pedes dilute rufi.

Obs. Proximus praecedenti, at paulo robustior, opacus, pube fusca, punctura capitis thoracisque subtiliore et conferta, elytris brevioribus et colore distinctus.

**Xantholinus.**

p. 313. adde

\* 14 — 15. *X. aulicus*: *Laete cyaneus, nitidus, antennis pedibusque nigris, elytris rufis, capite subtriangulari, laevi.* — Long.  $5\frac{1}{2}$  lin.

Habitat in Brasilia, Dom. Westermann.

Corpus laete cyaneum, splendidum. Antennae capite paulo longiores, articulo tertio secundo paulo longiore, nigrae, articulo secundo piceo, ultimo apice fulvo. Caput thorace paulo latius, antrorsum angustatum, basi emarginatum, angulis posterioribus rotundatis, laeve vel parce omnium subtilissime punctatum, sulcis intermediis frontis anticae conniventibus, punctis

prope oculos utrinque tribus, et serie punctorum in capitis basi transversali impressum. Thorax latitudine paulo longior, basin versus vix angustatus, lateribus subrectis, apice truncatus, angulis anterioribus subrectis, transversim convexus, laevissimus, intra angulum utrumque anteriorem puncto singulo maiusculo notatus. Scutellum obsolete punctatum. Elytra thorace paulo longiora, dorso parcius obsoletiusque subtiliter, lateribus crebrius distinctiusque punctata, rufa, purpureo-nitentia. Abdomen punctatum, parcius subtiliter griseo-pubescentius. Pedes nigri, tarsis testaceis.

p. 319.

29. *X. glabratus*: adde

*Xanth. occidentalis* Walzl It. Andalus. II. p. 58.

*Staph. ferrugineus* Rossi Faun. Etr. I. 248. 610. Ed. Hellw. 307. 610.

p. 326.

42. *X. consentaneus*: adde

*Xanth. cephalus* Say Transact. Am. Philosoph. Soc. Philad. Nov. Ser. IV. 452. 2.

*Gyrophyn. assimilis* Kirby Faun. Boreali-Amer. 88. 1.

Nomen in *X. cephalum* mutandum.

p. 329.

47. *X. Hesperius*: adde

*Xanth. limbatus* Walzl It. Andalus. II. p. 57.

p. 330.

48. *X. obscurus*: Forte huic adscribatur:

*Xanth. hamatus* Say Transact. Am. Philosoph. Soc. Philad. Nov. Ser. IV. 453. 3.

## Leptacinus.

p. 335.

4. *L. batychrus*: adde

*Xanth. tener* Walzl It. Andalus. II. p. 59.

## Palaestrinus.

p. 344.

1 — 2. *P. mutillarius*: Niger, thoracis apice, abdomine infra et fasciis duabus dorsalibus cinereo-villosis, elytris rufis, fulvo-pubescentibus. — Long. 5 lin.

Habitat in Bengala, Mus. Dom. Westermann.

Niger. Antennae capite paulo longiores, articulo tertio secundo sesqui longiore, articulis 4 — 10 sensim paulo brevioribus, ultimo apice truncato et inferne subtiliter acuminato. Palpi rufo-picei, filiformes, apice truncati. Caput thorace paulo latius, orbiculatum, fortiter convexus, sat fortiter dense punctatum, punctis confluentibus longitudinaliter rugulosum, pube longiore cinerea parcius vestitum. Thorax coleopteris duplo angustior, basin versus angustatus, latitudine antica haud longior, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus subrotundatis, fortiter con-

vexus vel elevatus, dense fortiter punctatus, disco subtiliter nigro-pubescente, limbo cinereo-villoso. Scutellum nigrum, fulvo-pubescent. Elytra thorace longiora, depressa, rufa, villis depressis fulvis vestita. Abdomen coleopteris angustius, supra nigro-pubescent, fasciis duabus cinereo-villosis, posteriore segmentum quartum occupante, anteriore basali, lateribus descendente et maculam permagnam triangularem includente; infra segmentis 4 anterioribus aequaliter cinereo-villosis, quinto nigro-, sexto cinereo-pubescente. Pectus cinereo-pubescent. Pedes picei, cinereo-pubescentes, coxis anticis castaneis, tibiis anticis incrassatis, externe laevigatis glabratiss.

p. 345. Inter Palaestrinum et Staphylinum novum genus inducendum est:

### **Caranistes Er.**

*Antennae rectae.*

*Palpi articulo ultimo apice truncato.*

*Pedes intermedii basi leviter distantes.*

*Tarsi posteriores deplanati.*

Corpus elongatum, alatum. Caput thorace maius, grossum, basi fortiter constrictum, oculis parvis, vix prominulis, intra marginem lateralem in capitis facie anteriore sitis, ovalibus, obliquis. Labrum transversum, bilobum, corneum, fortiter setosum, limbo laterali angusto, bilobo. Mandibulae validae. Maxillae mala exteriori breviori, apice barbata. Palpi maxillares articulo primo minuto, secundo tertioque leviter obconicis, subaequalibus, quarto paulo longiore, cylindrico, apice subcompresso, truncato. Palpi labiales articulo primo minuto, secundo obconico, tertio paulo longiore, cylindrico, apice compresso truncatoque. Antennae filiformes, apice compressae. Thorax coleopteris angustior. Scutellum triangulare. Elytra apice rotundata. Abdomen apicem versus subangustatum, marginatum. Pedes coxis intermediis parum distantibus, tibiis omnibus spinulosis, tarsis anticis dilatatis, tibia paulo latioribus, posterioribus, depressis, articulis 4 primis sensim decrescentibus.

### **O b s e r v a t i o.**

Tarsis posterioribus depressis cum Haematode, Cordylaspide et Scaphaeo conveniens, at maxillarum mala exterior haud elongata et tarsi posteriores ipsi angusti. A Staphylino praeter tarsorum structuram palpis maxillaribus apice truncatis discedit. Habitu proximus videtur Palaestrino, at pedes intermedii parum distantes, tibiae anticae haud incrassatae, tarsi antici tibia latiores, et tarsi posteriores depressi.

1. *C. Westermanni*: Niger, capite, thorace elytrisque obscure coeruleis, abdomine macula magna dorsali atro-tomentosa, apice ferrugineo, pedibus castaneis. — Long.  $7\frac{1}{2}$  lin.

Habitat in Bengala, Mus. Dom. Westermann.

Antennae capite paulo longiores, filiformes, apice compressae, articulo tertio secundo duplo longiore, 4—10 sensim paululum brevioribus crassioribusque, penultimis latitudine paulo brevioribus, ultimo truncatulo, medio abrupte subacuminato, fuscae, subtiliter densius pubescentes, articulo primo castaneo. Palpi rufo-picei. Caput thorace multo maius, coleopterorum latitudine, crassum, lateribus subrectis, basi emarginatum, angulis posterioribus rotundatis, infra nigrum, supra convexum, dense punctato-rugosum,



nigro-coeruleum, opacum, subtiliter cinereo-pubescentia. Thorax coleopteris angustior, basin versus angustatus, latitudine antica vix longior, basi cum angulis posterioribus rotundatus, apice utrinque oblique truncatus, angulis anterioribus acutiusculis, prominulis, convexus, confertim punctato-rugulosus, subtiliter cinereo-pubescentia, lateribus nigro-pilosus. Scutellum atro-tomentosum. Elytra thorace paulo longiora, apice rotundata, subtiliter punctato-rugulosa, nigro-coerulea, opaca, subtiliter cinereo-pubescentia, apice flavescenti-ciliata, ad suturam densius nigro-pubescentia. Abdomen nigrum, segmento penultimo apice, ultimo toto ferrugineo, tenuiter griseo-pubescentia, parcius nigro-pilosum, segmentis secundo tertioque dorsalibus macula magna e tomento aterrimo. Pedes rufo-castanei, coxis nigricantibus.

### Staphylinus.

p. 349.

4. *St. villosus*: adde

*Creophilus villosus* Kirby Faun. Boreali-Amer. 95. 1.

p. 360.

21. *St. cingulatus*:

*St. chrysurus* Kirby Faun. Boreali-Amer. 94. 1,

e Nova Scotia, licet minor, long.  $5\frac{1}{2}$  lin., nullo modo distinctus videtur.

p. 361. adde

22 — 23. *St. marmoratus*: *Niger, tomento fusco-nebuloso vestitus, antennarum basi, palpis pedibusque testaceis, femoribus medio nigris. — Long. 5 lin.*

Habitat in Java, Mus. Dom. Westermann.

Statura omnino *St. murini*, niger, vix nitidus. Antennae articulo tertio secundo sesqui longiore, quarto crassitie longiore, sequentibus transversis, nigrae, articulis 4 primis testaceis. Palpi testacei. Labrum testaceum. Caput, thorax et elytra tomento denso depresso fulvo-fuscoque variegato obducta. Caput thorace paulo latius, subdepressum. Thorax coleopteris angustior, basin versus subangustatus, latitudine paulo longior, lateribus rectis, basi cum angulis posterioribus rotundatis, angulis anterioribus acutiusculis, subdepressus. Scutellum macula magna nigro-tomentosa. Elytra thorace vix longiora. Abdomen segmentis duobus anterioribus dorsalibus fulvo-tomentosis, nigro-variegatis, reliquis subtiliter nigro-tomentosis, basi e pube cinerea maculatim variegatis, infra omnibus basi cinereo-pubescentibus. Pectus cinereo-pubescentia. Pedes testacei, femoribus supra et annulo medio nigris.

p. 376. adde

\* 50 — 51. *St. mysticus*: *Niger, fusco-pubescentia elytris brunneis abdomineque nigro-maculosis, tibiis tarsisque rufo-piceis. — Long. 7 lin.*

Habitat in America septentrionali, Dom. Riehl.

*St. maculoso* affinis, at minor et magis depressus, *St. fusco-maculato* aequalis, niger, opacus. Antennae capite paulo longiores, articulo tertio secundo sesqui longiore, sequentibus crassitie haud brevioribus, ultimo apice

abrupte subacuminato, piceae, articulo primo infra testaceo. Palpi rufi. Caput thorace paulo minus et angustius, antrorsum leviter angustatum, subdepressum, supra dense punctatum, fusco-pubescens. Thorax coleopteris vix angustior, latitudine sublongior, lateribus subrectis, parallelis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus rectis, leviter convexus, dense subtiliter punctatus, subtilissime carinatus, subtiliter fusco-pubescens. Scutellum macula magna rotundata atro-holasericea. Elytra thorace vix longiora, dense subtiliter punctata, brunnea, nigro-maculosa, e pube fuscescente nigricanteque subtili obsolete variegata. Abdomen dense subtiliter punctulatum, subtiliter fulvescenti-pubescens, supra segmentis singulis lituris duabus nigro-tomentosis signatis, ultimo rufo-piceo. Pedes tibiis tarsisque rufo-piceis.

Obs. A *St. maculoso* distinctus colore obscuriore, statura minore et magis depressa, thorace subtiliter punctato.

p. 389.

75. *St. fuscicornis*: adde

*Ocypus scabrosus* Curtis Linn. Transact. XVIII. 196. 28.

### Ocypus.

p. 409.

8. *O. masculus*: adde

*Staph. aethiops* Waltl It. Andal. II. p. 56.

Nomen in *St. aethiopem* mutandum. — Specimina illaesa densius nigro-tomentosa, abdominis segmentis dorsalibus basi macula minuta media e pube lutea signatis.

### Belonuchus.

p. 426. adde

Species mihi invisae, forte huius generis:

14. *B. ephippiatus*: Niger, elytris rufis. — Long. ultra 6 lin.

*Staph. ephippiatus* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser. IV. 448. 1.

Habitat in Mexico.

Niger, politus. Caput linea longitudinali inter oculos, orbitis superioribus punctis 3 vel 4, super hos serie e punctis 4 vel 5 et singulis 2 vel 3, basi utrinque serie e punctis 4 impressum. Palpi picei. Antennae articulo ultimo oblique truncato. Thorax semiovalis, punctis utrinque paucis dispersis impressus, seriebus dorsalibus punctis 4 vel 5 compositis. Scutellum dense punctatum, nigrum. Elytra cinnamomeo-rufa, subtiliter pubescentia punctataque. Abdomen supra tenuiter pubescens, nigro-coerulescens, versicolor. (Say.)

15. *B. connexus*: Niger, elytris pedibusque rufis. — Long. ultra 3 lin.

*Staph. connexus* Say Transact. Am. Philosoph. Soc. Philad. Nov. Ser. IV. 448. 2.

Habitat in Mexico.

Niger, politus. Caput orbitis punctatis et utrinque disperse punctatum, fronte inter antennis fortiter impressa, et supra hanc impressionem utrinque puncto singulo notata. Palpi nigro-picei. Thorax seriebus dorsalibus punctis ultra 5 fortibus compositis, lateribus utrinque punctis parvis dispersis impressus. Elytra cinnamomeo-rufa, brevissime pubescentia, subtiliter punctata. Pedes elytris concolores. — Similis praecedenti, at pedibus rufis, abdomine supra nigerrimo, haud versicolore, clypeo medio leviter producto. (Say.)

## Philonthus.

p. 437.

16. *Ph. aeneus*: *Locis citatis adde*

*Phil. politus* Kirby Faun. Boreali-Amer. p. 91. 1.

*Phil. mandibularis* Kirby Faun. Boreali-Amer. p. 92. 2.

Obs. *Ph. politum* Kirby ad marem, *Ph. mandibularem* ad feminam referas.

p. 457. adde

\* 47 — 48. *Ph. inquietus*: *Niger, nitidus, antennarum basi, elytrorum limbo pedibusque testaceis, elytris dense profundeque punctatis, capite ovato.* — Long.  $2\frac{1}{2}$  lin.

Habitat in America septentrionali, Dom. Zimmermann.

Statura omnino *Ph. sordidi* et minoribus eius individuis aequalis, niger, nitidus, capite thoraceque nigro-subaeneis, politis. Antennae capite thoraceque paulo breviores, articulo tertio secundo paulo longiore, 4 — 10 subcylindricis, longitudine et crassitie subaequalibus, ultimo ovato, apice truncato et inferne subacuminato, nigrae, articulo primo infra, secundo toto testaceis. Palpi piceo-testacei. Caput thoracis prope latitudine, ovatum, leviter convexum, fronte supra punctis paucis sparsis, inter oculos utrinque punctis duobus oblique positis impressus. Thorax coleopteris dimidio angustior, latitudine paulo longior, lateribus subrectis, antrorsum vix angustatus, basi rotundatus, apice truncatus, angulis anterioribus obtusiusculis, deflexis, posterioribus obtusis, leviter convexus, seriebus dorsalibus punctis 4 sat remotis, postico paulo remotiore, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis omnibus minus subtilibus. Scutellum crebre punctatum, tenuiter griseo-pubescent. Elytra thorace paulo longiora, densius fortiusque punctata, parce griseo-pubescentia, limbo laterali, apicali suturalique tenui testaceo. Abdomen crebre punctulatum, griseo-pubescent, segmentis omnibus apice late testaceo-limbatis. Pedes testacei, coxis anterioribus concoloribus, coxis anticis femoribusque dilutioribus, tarsis anticis in utroque sexu simplicibus.

p. 461. adde

54 — 55. *Ph. iopterus*: *Niger, nitidus, elytris crebre punctatis, violaceis, splendidis.* — Long. 4 lin.

Habitat Bogotae in Columbia interiore, Dom. De Haen.

*Ph. candenti* proximus, nigerrimus, nitidus, capite thoraceque nitidissimis, elytris splendide violaceis. Antennae capite paulo longiores, articulo tertio secundo paulo longiore, 4 — 10 brevibus, leviter transversis, ultimo



subgloboso, apice leviter emarginato et inferne acuminato. Mandibulae longiores, falcatae, acutae. Caput thorace latius, breviter obovatum, fronte subdepressa, supra punctis sparsis, inter oculos utrinque duobus transversim positis, fortioribus notata, antice medio foveolata. Thorax coleopteris duplo angustior, latitudine longior, basin versus vix angustatus, lateribus subrectis, basi leviter rotundatus, angulis anterioribus deflexis, fortiter rotundatis, posterioribus obtusis, transversim convexus, seriebus dorsalibus punctis 5, aequaliter distantibus compositis, lateribus utrinque punctis 6 impressis, punctis omnibus minus subtilibus. Scutellum subtilius crebre punctatum. Elytra thorace paulo longiora, crebre minus fortiter punctata, pube subtili depressa nigra tenuiter vestita. Abdomen apicem versus leviter angustatum, subtiliter parcius punctulatum, parcius nigro-pubescent. Pedes elongati, graciles, coxis intermediis distantibus, tibiis omnibus subtiliter spinulosi, tarsis anticis simplicibus. (Mas.)

Obs. Praeter elytrorum colorem a Ph. candente imprimis antennis multo brevioribus, thorace longiore, angulis anterioribus fortius rotundatis distinctus.

56. *Ph. ebeninus*: Huic adscribendus videtur:

*Staph. quisquiliarius var. longicornis* Waltl. It. Andalus. II. p. 57.

p. 470. adde

69 — 70. *Ph. promptus*: Niger, nitidus, coxis anticis testaceis, elytris brevioribus, fortius punctatis, capite ovato. — Long.  $3\frac{1}{2}$  lin.

Habitat in Pennsylvania, Dom. Zimmermann.

Statura Ph. bipustulati, niger, nitidus, capite thoraceque pernitidis. Antennae capitis thoracisque prope longitudine, articulo tertio secundo paulo longiore, 4 — 10 subaequalibus, crassitie paulo longioribus, ultimo ovato, apice truncato et inferne acuminato, totae nigrae. Palpi concolores. Caput thorace plus dimidio angustius, ovatum, punctis parvis utrinque super oculos, et utrinque duobus transversim positis inter oculos impressum. Thorax coleopteris paulo angustior, latitudine baseos vix longior, antrorsum leviter angustatus, lateribus subrectis, basi cum angulis posterioribus rotundatus, apice truncatus, angulis anterioribus deflexis, subrotundatis, convexus, seriebus dorsalibus punctis 5, aequaliter distantibus, compositis, lateribus punctis praeter marginalia utrinque 5 impressus, punctis minus subtilibus. Scutellum subtiliter punctulatum. Elytra thoracis longitudine, sat crebre minus subtiliter punctata, glabra. Abdomen dense subtiliter punctatum, subtiliter cinereo-pubescent, segmentis ventralibus margine apicali rufo-piceo. Pedes picei, coxis anticis totis laete testaceis, intermediis approximatis, tibiis omnibus spinulosi.

Tarsi antici feminae leviter dilatati. Mas latet.

Obs. Elytris brevioribus, fortius punctatis ab affinis distinctus.

71. *Ph. flavolimbatus*: Eundem Dom. Zimmermann e Carolina meridionali misit, nonnisi elytrorum limbo laterali obsoleto distinctum.

p. 473.

74. *Ph. ventralis*: Forte huc pertinet:

*Staph. dimidiatus* Say Transact. Amer. Philosoph. Soc. Philadelph. Nov. Ser.

IV. 450. 6.

p. 476.

80. *Ph. rufimanus*:

Habitat etiam in Helvetia, Sardinia.

Varietatem pedibus anterioribus nigris, coxis solis laete rufo-testaceis, cum genuino in Sardinia lectam nuperrime Prof. Gené misit.

p. 481. adde

88 — 89. *Ph. tachiniformis*: *Niger, thorace sanguineo, pedibus piceis.*  
— *Long.*  $2\frac{1}{3}$  lin.

*Staph. tachiniformis* Say *Transact. Amer. Philosoph. Soc. Nov. Ser. IV.* 450. 7.

Habitat in Mexico.

*Niger.* Caput parvum, ovale, politum, punctis utrinque duobus inter oculos, 8 vel 10 supra oculos impressum. Antennae fuscae, basi obscure piceae. Os piceum. Thorax sanguineus, seriebus dorsalibus punctis 5, a basi et ab apice remotis, compositis, lateribus punctis utrinque duobus, margine tribus, triangulariter positis impressus. Elytra subaenea, crebre subtiliter punctata, pube depressa vestita. Pedes testacei, posticis, tibiis et tarsis obscurioribus. (Say.)

p. 485.

96. *Ph. fulvipes*: *Locis citatis adde*

*Phil. fulvipes* Kirby *Faun. Boreali-Amer.* 93. 4.

p. 486.

97. *Ph. brunneus*: *Forte huc referatur:*

*Phil. picatus* Kirby *Faun. Boreali-Amer.* 93. 3.

p. 493.

111. *Ph. (?) stenoides*: *adde*

*Staph. stenoides* Grav. *Mon.* 90. 86.

p. 494. adde

\* 112 — 113. *Ph. pretiosus*: *Viridi-aeneus, capite thoraceque cupreo-auratis, fulgidis.* — *Long.* 8 lin.

Habitat Bogotae in Columbia interiore, Dr. De Haen.

Insectum splendidissimum, corpore laete viridi-aureo, nitido, capite thoraceque supra purpurascenti-aureis, splendidis. Antennae capite thoraceque paulo breviores, apicem versus haud incrassatae, articulo tertio secundo sesqui longiore, 4 — 10 sensim paulo brevioribus, ultimo ovato, apice oblique subemarginato, inferne acuminato, nigrae, articulis tribus primis viridibus. Caput thorace paulo latius suborbiculatum, sparsim punctatum, fronte media laevi. Thorax coleopteris plus dimidio angustior, latitudine longior, basin versus leviter angustatus, lateribus anterieus rotundatis, posterius subsinuatis, basi rotundatus, apice truncatus, angulis anterioribus deflexis, subrectis, posterioribus obtusis, convexus, seriebus dorsalibus irregularibus, 7 — 9-punctatis, lateribus punctis parvis impressus. Scutellum parce punctatum. Elytra thorace paulo longiora, parcius et minus fortiter punctata, parcius nigro-pubescentia. Abdomen sensim angustatum, parce sub-

tiliterque punctatum, tenuiter nigro-pubescent. Pedes virides, coxis inter-  
mediis approximatis, tibiis posterioribus spinulosis, tarsis nigris.

Tarsi antici maris modice dilatati. Femina latet.

p. 499.

117. *Ph. orphanulus* pro *Ph. orphano* substituendus.

p. 501.

124 — 125. *Ph. ferrugatus*: *Elongatus, depressus, ferrugineus, fronte nigra, abdomine piceo, capite subquadrato*. — Long.  $4\frac{1}{2}$  lin.

Habitat in Java, Mus. Dom. Westermann.

Statura elongata, depressa omnino Belonuchi haemorrhoidalis. Antennae capite sesqui longiores, articulo primo leviter elongato, tertio secundo paulo longiore, sequentibus sensim paulo brevioribus et crassioribus, 5 — 10 transversis, ultimo ovato, apice oblique submarginato et inferne acuminato, ferrugineae, medio fusciscentes, summo apice flavescente. Mandibulae longiores, falcatae, acutae. Caput thorace latius, subquadratum, fronte depressa, fortius parce punctata, punctis medio subseriatis, nitidum, supra nigro-infra rufo-piceum. Thorax coleopteris angustior, basin versus angustatus, latitudine antica vix longior, lateribus subrectis, basi rotundatus, apice utrinque oblique truncatus, angulis anterioribus acutiusculis, prominulis, posterioribus obtusis, depressus, seriebus dorsalibus 8 — 10-punctatis, lateribus parce punctatus, saturate ferrugineus, nitidus, oblique inspectus subsericans. Scutellum rufo-piceum, crebre punctatum. Elytra thorace paulo longiora, depressa, densius punctata, punctorum maiorum distantium seriebus duabus notata, saturate ferruginea, nitida, griseo-pubescentia. Abdomen apicem versus sensim attenuatum, minus dense punctatum, griseo-pubescent pilosumque, piceum, segmentis 4 primis apice obscure ferrugineo-marginatis, ultimis duobus apice testaceis. Pectus piceum. Pedes ferruginei, coxis intermediis modice distantibus, tibiis omnibus spinulosis.

Tarsi antici feminae leviter dilatati. Mas latet.

p. 504.

130. *Ph. haematurus*: adde

*Staph. apicalis* Say Transact. Am. Philosoph. Soc. Philad. Nov. Ser. IV. 451. 8.

Nomen in *Ph. apicalem* mutandum.

p. 515. adde speciem mihi invisam, forte huius generis:

151. *Ph. apicicornis*: *Nigro-coeruleus, capite transverso, laevigato, punctato, ore, mandibulis basi ferrugineis, antennis basi rufis, medio nigris, apice flavis, thorace oblongo, postice rotundato, angulis anticis acuminatis, supra laevigato, vage punctato, elytris brunneo-aeneis, micantibus, subrugosis, villosis, abdomine marginato, subrugoso, villosio, pedibus ferrugineis*. — Long. 5 — 6 lin.

*Staph. apicicornis* Guér. Voyag. d. l. Coquille, Zool. II. II. p. 62. Ins. t. 1. f. 21.

Habitat in Nova Guinea.

Antennae coloribus insignes: articulis 3 primis rufis, sequentibus 5 nigris, ultimis 3 ochraceis. Corpus totum nigro-coerulescens, oris partibus, mandibularum basi pedibusque rufo-ferrugineis, elytris rubro-cupreis, violaceo-micantibus. (Guér.)



**Lathrobium.**

p. 603. adde

- \* 26 — 27. *L. Stilicinum: Elongatum, nigrum, elytris piceis, antennis pedibusque ferrugineis, capite suborbiculato thoraceque subovato confertissime punctatis, subrugulosis.* — Long.  $2\frac{2}{3}$  lin.

Habitat in Sardinia, Dom. Prof. Gené.

Corpus elongatum, subdepressum, nigrum, minus nitidum, totum subtilissime griseo-pubescens. Antennae capite thoraceque paulo breviores, articulo secundo tertioque subaequalibus, sequentibus sensim paulo brevioribus, rufo-ferrugineae. Mandibulae rufo-piceae. Palpi rufo-testacei. Caput thorace paulo latius, suborbiculatum, supra leviter convexum, confertissime punctatum, subrugulosum. Thorax coleopteris paulo angustior, latitudine paulo longior, basin versus subangustatus, angulis posterioribus rotundatis, antice apicem versus rotundato-attenuatus, subdepressus, confertissime punctatus, subrugulosus. Elytra thorace paulo longiora, crebre fortius punctata, picea, nitidula. Abdomen dense subtiliter punctatum. Pedes ferruginei.

Mas abdominis segmento inferiore quinto obsolete longitudinaliter impresso, apice leviter emarginato, sexto basi bicarinato, apice triangulariter exciso, femina eodem rotundato-producto.

Variat elytris piceo-testaceis.

Obs. Capitis thoracisque forma et punctura non minus ac *L. scabricolle* Stilici habitum prae se ferens, at huius generis, *L. scabricolli* affine, at alatum, corpore minore, subdepresso, thorace magis elongato, elytris longioribus etc. perquam distinctum.

**Sunius.**

p. 638.

1. *S. filiformis: adde*

*Paederus filum* Walth. Isis. 1838. IV. 267. 1.

**Paederus.**

p. 650.

1. *P. littoralis:*

*P. confinis* Zetterst. Ins. Lappon. 69. 2,

ab auctore benevole communicatus *P. littoralis* genuino paulo minor et alatus, ceterum autem nullo modo distinctus, ut non possim quin huius varietatem eum habeam. Deleatur igitur apud *P. riparium*. (p. 654.)

p. 660. adde

- \* 20 — 21. *P. melanurus: Alatus, testaceus, abdomine apice nigro.* — Long. 3 lin.

Habitat Ticini, inundante Ticino fluvio a Dom. Gené olim lectus, nuper benevole communicatus.

Statura omnino *P. riparii*, colore corporis cum capite, pectore elytrisque laete testaceo, abdominis ultimis duobus segmentis solis nigerrimis eximius. Antennae capitis thoracisque longitudine, articulo tertio secundo duplo longiore, 4 — 10 sensim paulo brevioribus et crassioribus, ultimo oblongo-ovato, apice oblique acuminato, flavescenti-testaceae, apicem versus fusciscentes. Palpi immaculati. Caput thoracis latitudine, orbiculatum, supra leviter convexum, parce subtiliterque punctulatum, utrinque ad antel-

narum basin subimpressum, parce nigro-pilosellum. Thorax coleopterorum latitudine, latitudine paulo longior, basin versus leviter angustatus, lateribus anterieus rotundatis, subovatus, leviter convexus, parce obsoletissime punctatus, parce pilosus. Elytra thorace paulo longiora, crebre profundeque punctata, parce nigro-pilosella. Abdomen parcius punctulatum, nigro-pilosellum. Pedes dilute testacei, geniculis subfusciscentibus.

Obs. Descriptus a Dom. Prof. Gené in opusculo mihi invisio: De quibusdam Coleopteris Italiae novis aut rarioribus. Tentam. inaug., Ticin. Reg. 1830.

p. 666.

37. *P. australis*: adde

Guér. *Voyag. d. l. Coquille, Zool. II. II. p. 63.*

P. ripario statura et coloribus proximus, at pedibus totus nigris. (Guér.)

Obs. Quo differat a *P. melampode* non patet.

p. 667. adde

39. *P. cyanipennis*: *Ater, nitidus, elytris rugosis, elongatis, cyaneis, antennis brunneis, basi ferrugineis, pedibus atris.* — Long. 3 lin.

Guér. *Voyag. d. l. Coquille, Zool. II. II. p. 62. Ins. t. I. f. 22.*

Habitat Limae.

Niger, nitidus. Caput orbiculatum, punctatum, pilosellum. Antennae brunneae, pubescentes, articulis tribus primis rufescentibus. Thorax globosus, nitidus, parce disperse punctatus, pilosellus. Elytra oblonga, rugosa, laete violacea, splendida. Abdomen nigrum, rugosum, pilosellum, marginatum. Pedes graciles, nigri. (Guér.)

### Bledius.

p. 760.

1. *B. taurus*: adde

*Bled. Stephensii* Westwood *Zool. Journ. III. p. 61, 301, 509. t. 11. f. 4.*

### Platystethus.

Obs. Falso plerisque locis *Platysthetus* scripsi.

p. 783.

3. *Pl. nodifrons*: adde

*Oxyt. asphaltinus* Waltl *Isis. 1838. IV. 267. 2.*

### Phloeonaeus.

p. 800.

1. *Ph. caelatus*: adde

*Oxyt. biarcuatus* Matthews *Ent. Mag. V. p. 198.*

### Prognatha.

p. 837.

1. *P. quadricornis*: adde

*Siagon. 4-corne* Westw. *Zool. Journ. III. p. 61.*

### Lesteua.

p. 857.

5. *L. punctata*: *Var. pallida, immatura huius speciei esse videtur:*

*Anthophag. villosus* Waltl. *Isis. 1838. IV. 268. 3.*

# EXPLICATIO TABULARUM.

---

## Tab. I.

- Fig. 1. *Peliusa labiata*.  
„ 2. *Euryusa sinuata*.  
„ 3. *Pronomaea rostrata*.  
„ 4. *Tanygnathus terminalis*.  
„ 5. *Holius analis*.  
„ 6. *Diochus nanus*.  
„ 7. *Scytalinus serpentinus*.  
„ 8. *Palaminus variabilis*.  
„ 9. *Oedichirus Paederinus*.

## Tab. II.

- Fig. 1. *Ophites versatilis*.  
„ 2. *Echiaster longicollis*; a *labium*, b *maxilla*, c *labrum* E. *melanuri*.  
„ 3. *Megalops cephalotes*; a *labium*, b *maxilla*.  
„ 4. *Holotrochus volvulus*.  
„ 5. *Apocellus sphaericollis*.  
„ 6. *Lispinus angustatus*.  
„ 7. *Isomalus complanatus*.  
„ 8. *Hypotelus pusillus*.  
„ 9. *Glyptoma crassicorne*.

## Tab. III.

- Fig. 1. *Myrmedoniae (humeralis) labium*.  
„ 2. *Autaliae (impressae) labium*.  
„ 3. *Falagriae (sulcatae) labium*.  
„ 4. *Bolitocharae (lunulatae) labium*.  
„ 5. *Ocaleae (castaneae) labium*.  
„ 6. *Caloderae (nigritae) labium*.  
„ 7. *Tachyusae (coarctatae) labium*.  
„ 8. *Homalotae (elongatulae) labium*.  
„ 9. *Aleocharae (fuscipedis) labium*.  
„ 10. *Oligotae (pusillimae) labium*.  
„ 11. *Gyrophaenae (nanae) labium*.  
„ 12. *Peliusae (labiatae) labium*.  
„ 13. *Placusae (infimae) labium*.  
„ 14. *Euryusae (sinuatae) labium*.  
„ 15. *Dinardae (dentatae) labium*.  
„ 16. *Lomechusae (emarginatae) labium*.  
„ 17. *Silusae (rubiginosae) labium*.  
„ 18. *Pronomaeae (rostratae) labium*.  
„ 19. *Gymnusae (brevicollis) labium*.



Fig. 20. *Myllaenae* (dubiae) *labium*.

- „ 21. *Myrmedoniae* (humeralis) *maxilla*.
- „ 22. *Homalotae* (elongatulae) *maxilla*.
- „ 23. *Euryusae* (sinuatae) *maxilla*.
- „ 24. *Dinardae* (rubiginosae) *maxilla*.
- „ 25. *Lomechusae* (emarginatae) *maxilla*.
- „ 26. *Silusae* (rubiginosae) *maxilla*.
- „ 27. *Pronomaeae* (rostratae) *maxilla*.
- „ 28. *Gymnusae* (brevicollis) *maxilla*.
- „ 29. *Myllaenae* (dubiae) *maxilla*.

#### Tab. IV.

Fig. 1. *Hypocypiti* (granuli): a *labium*, b *maxilla*.

- „ 2. *Conuri* (pubescentis): *labium*.
- „ 3. *Tachypori* (obtusi): a *labium*, b *maxilla*.
- „ 4. *Habroceri* (capillaricornis): a *labium*, b *maxilla*, c *antenna*.
- „ 5. *Tachini* (rufipedis): a *labium*, b *maxilla*.
- „ 6. *Boletobii* (atricapilli): a *labium*, b *maxilla*.
- „ 7. *Mycetopori* (lepidi): a *labium*, b *maxilla*.
- „ 9. *Acylophori* (glabricollis): a *labium*, b *maxilla*.
- „ 10. *Eurypori* (picipedis): a *labium*, b *maxilla*.
- „ 11. *Astrapaei* (Ulmi): a *labium*, b *maxilla*.
- „ 12. *Philonthi* (splendendis): a *labium*, b *maxilla*.
- „ 13. *Heterothopis* (dissimilis): a *labium*, b *maxilla*.
- „ 14. *Staphylini* (maxilloso) *labium*.
- „ 15. *Ocypodis* (morionis) *labium*.
- „ 16. *Haematodis* (bicoloris): a *labium*, b *maxilla*.
- „ 17. *Cordylaspidis* (pilosus): a *labium*, b *maxilla*.
- „ 18. *Scariphaei* (luridipennis): a *labium*, b *maxilla*, c *tarsus posticus*,  
superne et a latere inspectus.
- „ 19. *Sterculiae* (formicarum): a *labium*, b *maxilla*.
- „ 20. *Scytalini* (serpentini): a *labium*, b *maxilla*.
- „ 21. *Xantholini* (glabrati): a *labium*, b *maxilla*.
- „ 22. *Leptacini* (batychoi): a *labium*, b *maxilla*.
- „ 23. *Holisi* (humilis): a *labium*, b *maxilla*.
- „ 24. *Othii* (fulvipennis): a *labium*, b *maxilla*.
- „ 25. *Platyprosopi* (Beduini): a *labium*, b *maxilla*, c *labrum*.
- „ 26. *Lathrobii* (brunnipedis): a *labium*, b *maxilla*.
- „ 27. *Doliceonis* (lathrobiodis): a *labium*, b *maxilla*.
- „ 28. *Cryptobii* (fracticornis): a *labium*, b *maxilla*.
- „ 29. *Lithocharidis* (ochraceae): a *labium*, b *maxilla*.
- „ 30. *Paederi* (riparii): a *labium*, b *maxilla*.
- „ 31. *Stilici* (rufipedis): a *labium*, b *maxilla*.
- „ 32. *Ophitis* (versatilis): a *labium*, b *maxilla*.
- „ 33. *Sunii* (angustati): a *labium*, b *maxilla*.
- „ 34. *Pinophili* (latipedis): a *labium*, b *maxilla*.

- Fig. 35. *Taenodematis* (cyanescentis): a *labium*, b *maxilla*.  
 „ 36. *Palamini* (variabilis): a *labium*, b *maxilla*.  
 „ 37. *Oedichiri* (Paederini): a *labium*, b *maxilla*.  
 „ 38. *Procirri* (Lefebvrei): a *labium*, b *maxilla*.

### Tab. V.

- Fig. 1. *Dianoi* (coerulescentis): a *labium*, b *maxilla*.  
 „ 2. *Steni* (speculatoris): a *labium*, b *maxilla*.  
 „ 3. *Euaestheti* (scabri): a *labium*, b *maxilla*.  
 „ 4. *Olisthaeri* (megacephali): a *labium*, b *maxilla*.  
 „ 5. *Phloeocharidis* (subtilissimae): a *labium*, b *maxilla*.  
 „ 6. *Oxyperi* (rufi): a *labium*, b *maxilla*.  
 „ 7. *Leptochiri* (scoriacei): a *labium*, b *maxilla*, c *labrum*.  
 „ 8. *Lispini* (angustati): a *labium*, b *maxilla*.  
 „ 9. *Piesti* (penicillati): a *labium*, b *maxilla*, c *labrum*.  
 „ 10. *Prognathae* (humeralis): a *labium*, b *maxilla*.  
 „ 11. *Isomali* (complanati): a *labium*, b *maxilla*.  
 „ 12. *Hypoteli* (pusilli): a *labium*, b *maxilla*.  
 „ 13. *Osorii* (atri): a *labium*, b *maxilla*.  
 „ 14. *Holotrochi* (volvuli): a *labium*, b *maxilla*.  
 „ 15. *Bledii* (tricornis): a *labium*, b *maxilla*, c *labrum*.  
 „ 16. *Platystethi* (morsitantis): a *labium*, b *maxilla*.  
 „ 17. *Oxyteli* (picei): a *labium*, b *maxilla*, c *labrum*.  
 „ 18. *Phloeonaei* (caelati): a *labium*, b *maxilla*.  
 „ 19. *Trogophloeii* (elongatuli): a *labium*, b *maxilla*.  
 „ 20. *Apocelli* (sphaericollis): a *labium*, b *maxilla*.  
 „ 21. *Coprophili* (striatuli): a *labium*, b *maxilla*, c *labrum*.  
 „ 22. *Acrognathi* (mandibularis): a *labium*, b *maxilla*, c *labrum*.  
 „ 23. *Syntomii* (aenei): a *labium*, b *maxilla*.  
 „ 24. *Deleasteris* (dichroi): a *labium*, b *maxilla*, c *labrum*.  
 „ 25. *Anthophagi* (alpini): a *labium*, b *maxilla*.  
 „ 26. *Lesteuae* (punctatae): a *labium*, b *maxilla*, c *labrum*.  
 „ 27. *Arpedii* (quadri): a *labium*, b *maxilla*.  
 „ 28. *Acidotae* (crenatae): a *labium*, b *maxilla*.  
 „ 29. *Olophri* (picei): a *labium*, b *maxilla*, c *labrum*.  
 „ 30. *Lathrimaei* (atrocephali): a *labium*, b *maxilla*, c *labrum*.  
 „ 31. *Omali* (rivularis): a *labium*, b *maxilla*.  
 „ 32. *Anthobii* (floralis): a *labium*, b *maxilla*.  
 „ 33. *Proteini* (brachypteri): a *labium*, b *maxilla*.  
 „ 34. *Megarhtri* (hemipteri): a *labium*, b *maxilla*.  
 „ 35. *Micropepli* (porcati): a *labium*, b *maxilla*.

Obs. Labia omnia absque palpo labiali sinistro delineavi, ut ligulae paraglossarumque figura distinctius conspiciatur.

# INDEX NOMINUM.

Synonyma litteris obliquis distincta sunt.

	pag.		pag.		pag.
<i>Achenium</i>	581	<i>abdominalis</i>	146	<i>collaris</i>	43
2. <i>aequatum</i>	583	<i>aenescens</i>	111	<i>complana</i>	90
<i>anale</i>	580	<i>aethiops</i>	67. 68	<i>coracina</i>	75
5. <i>basale</i>	585	<i>affinis</i>	184	<i>corticalis</i>	77
<i>Caucasicum</i>	580	<i>agaricola</i>	132	40. <i>crassicornis</i>	177
<i>cordatum</i>	582	<i>alternans</i>	145	<i>crassicornis</i>	134. 163
6. <i>debile</i>	585	26. <i>amplicollis</i>	171	<i>crassiuscula</i>	162
1. <i>depressum</i>	582	<i>analis</i>	114. 115. 154	<i>curta</i>	166
<i>depressum</i>	586	35. <i>angulata</i>	175	<i>deplanata</i>	60. 110
4. <i>ephippium</i>	584	<i>angustata</i>	136	<i>depressa</i>	99
8. <i>humile</i>	586	<i>angustula</i>	91	<i>depressiuscula</i>	90
9. <i>ieiunum</i>	587	<i>annularis</i>	135	29. <i>deserta</i>	173
3. <i>planum</i>	583	<i>aterrima</i>	104. 133	<i>dichroa</i>	107
7. <i>striatum</i>	585	<i>atra</i>	73	3. <i>dilatata</i>	160
10. <i>tenellum</i>	587	<i>atramentaria</i>	111	<i>dimidiata</i>	80
<i>Acidota</i>	860	<i>atrata</i>	198	<i>dubia</i>	210
1. <i>crenata</i>	861	<i>bifoveolata</i>	115	18. <i>duplicata</i>	167
2. <i>cruentata</i>	862	<i>bilineata</i>	168	<i>elegantula</i>	158
3. <i>ferruginea</i>	862	17. <i>bimaculata</i>	167	<i>elongatula</i>	89. 90. 101
<i>rufa</i>	861	9. <i>bipunctata</i>	163	25. <i>erythroptera</i>	171
4. <i>subcarinata</i>	863	16. <i>bisignata</i>	166	<i>evanescens</i>	114
<i>Acrognathus</i>	816	<i>Boleti</i>	103. 191	<i>excavata</i>	59
2. <i>cephalotes</i>	817	<i>borealis</i>	138	<i>excusa</i>	212
1. <i>mandibularis</i>	817	<i>brevicollis</i>	212	<i>exigua</i>	175
3. <i>palpalis</i>	818	<i>brevicornis</i>	39	<i>exilis</i>	90. 115
<i>Acylphorus</i>	519	10. <i>brevipennis</i>	163	<i>flavipes</i>	124
<i>Ahrensii</i>	519	6. <i>breviuscula</i>	161	<i>flavitaris</i>	74
3. <i>femoralis</i>	520	4. <i>Cayennensis</i>	161	<i>fracticornis</i>	54
1. <i>glabricollis</i>	519	<i>canaliculata</i>	36	<i>fulgida</i>	40
2. <i>longicornis</i>	520	<i>carboraria</i>	75	15. <i>fumata</i>	166
4. <i>picipes</i>	521	<i>carnivora</i>	163. 212	<i>fumata</i>	170
5. <i>pronus</i>	521	<i>castanoptera</i>	103	<i>funesta</i>	38
<i>Adelobium</i>	576	2. <i>chrysorrhoea</i>	160	<i>fungi</i>	131
<i>brachypterum</i>	576	<i>cincta</i>	58	<i>fusca</i>	137
<i>Agrodes</i>	301	<i>cinnamomea</i>	127	<i>fuscicornis</i>	39
<i>elegans</i>	303	<i>circellaris</i>	98	1. <i>fuscipes</i>	156
<i>Aleochara</i>	159	<i>coerulea</i>	76	<i>fuscula</i>	133



	pag.		pag.		pag.
<i>graminicola</i>	81	<i>pulchella</i>	157	7. <i>limbatum</i>	894
<i>grandicollis</i>	39	<i>pulchra</i>	58	18. <i>longipenne</i>	898
<i>haemorrhoea</i>	135	30. <i>pulla</i>	173	15. <i>luteicorne</i>	897
38. <i>haemorrhoidalis</i>	176	<i>pulla</i>	144	17. <i>luteipenne</i>	898
<i>haemorrhoidalis</i>	170	<i>pumilio</i>	195	12. <i>minutum</i>	896
<i>humeralis</i>	38	<i>pusilla</i>	811	14. <i>montanum</i>	897
<i>impressa</i>	47	<i>pusillima</i>	179	3. <i>nigrum</i>	892
<i>impressifrons</i>	141	<i>pygmaea</i>	134. 137	<i>oblongum</i>	885
<i>inguinalis</i>	98	<i>quisquiliarum</i>	84	4. <i>obsoletum</i>	892
<i>inguinula</i>	116	<i>reptans</i>	77	8. <i>ophthalmicum</i>	891
<i>intricata</i>	163	33. <i>restricta</i>	174	<i>planum</i>	886
32. <i>laevigata</i>	174	<i>ricularis</i>	47	<i>pusillum</i>	879
<i>laevigata</i>	162	<i>ruficollis</i>	915	<i>rivulare</i>	875
<i>laevis</i>	37	24. <i>ruficornis</i>	170	<i>rugosum</i>	816
21. <i>lanuginosa</i>	168	<i>ruficornis</i>	142. 143	<i>Salicinum</i>	880
<i>lata</i>	159	<i>rufilabris</i>	63	<i>Salicis</i>	880
5. <i>lateralis</i>	161	7. <i>rufipennis</i>	162	11. <i>scutellare</i>	895
<i>lateralis</i>	152	<i>scapularis</i>	137	9. <i>Sorbi</i>	895
<i>laticollis</i>	39	<i>sericans</i>	103	<i>striatum</i>	882
<i>limbata</i>	37	<i>socialis</i>	102	10. <i>tempestivum</i>	895
<i>linearis</i>	81. 91	81. <i>spissicornis</i>	173	2. <i>triviale</i>	892
<i>lividipennis</i>	129	<i>strumosa</i>	203	<i>Viburni</i>	879
<i>longicornis</i>	81. 103. 129	<i>sulcata</i>	49. 50	<i>Anthophagus</i>	847
<i>longiuscula</i>	81. 145	<i>sulcatula</i>	50	6. <i>abbreviatus</i>	850
3-4. <i>loricata</i>	918	<i>suturalis</i>	157	2. <i>alpinus</i>	848
<i>lucida</i>	58	<i>tachyporoides</i>	918	1. <i>armiger</i>	848
<i>lugens</i>	39	14. <i>taeniata</i>	165	4. <i>Austriacus</i>	849
<i>lunulata</i>	58	<i>tenella</i>	136	12. <i>brunneus</i>	851
<i>macilenta</i>	137	<i>tenera</i>	140	11. <i>caesus</i>	853
<i>Mannerheimii</i>	139	<i>tenuis</i>	77	5. <i>caraboides</i>	850
<i>marginalis</i>	138	<i>teres</i>	77. 89	<i>dichrous</i>	819
<i>melanaria</i>	151	<i>terminalis</i>	89	<i>globulicollis</i>	853
<i>minuta</i>	211	<i>testacea</i>	68	<i>intermedius</i>	856
22. <i>moerens</i>	169	8. <i>tristis</i>	162	<i>Kunzei</i>	853
23. <i>moesta</i>	170	<i>umbrata</i>	134. 144	<i>Lapponicus</i>	849
<i>moesta</i>	81. 171	28. <i>vaga</i>	172	<i>mandibularis</i>	848
34. <i>morion</i>	175	<i>validicornis</i>	129	<i>maxillosus</i>	848
<i>nana</i>	184	11. <i>verberans</i>	164	<i>nigrita</i>	853
<i>nigra</i>	54	39. <i>villosa</i>	177	<i>obscurus</i>	855
<i>nigricollis</i>	64	12. <i>vulnerata</i>	164	3. <i>Omalinus</i>	849
<i>nigritula</i>	103	1. <b>ALEOCHARINI</b>	33	10. <i>plagiatus</i>	852
20. <i>nitida</i>	168	<i>Anodus</i>	416	9. <i>praeustus</i>	852
<i>nitidula</i>	183	<i>falcifer</i>	418	7. <i>scutellaris</i>	851
19. <i>notula</i>	167	<i>messor</i>	418	8. <i>testaceus</i>	851
<i>obfuscata</i>	134	<i>morio</i>	417	<i>varians</i>	853
<i>obscura</i>	54	<b>Anthobium</b>	890	<i>variegatus</i>	853
37. <i>obscurella</i>	176	6. <i>abdominale</i>	893	13. <i>verticalis</i>	854
<i>opaca</i>	144	16. <i>anale</i>	898	<i>villosus</i>	933
<i>pallidula</i>	140	<i>brunneum</i>	881	<i>Aploderus</i>	799
<i>parva</i>	139	<i>caesum</i>	877	<i>Apocellus</i>	812
<i>pellucida</i>	152	<i>castaneum</i>	859	3. <i>albipes</i>	814
27. <i>peltata</i>	172	<i>deplanatum</i>	885	1. <i>sphaericollis</i>	813
<i>picea</i>	54	<i>depressum</i>	885	2. <i>ustulatus</i>	813
<i>pilicornis</i>	268	5. <i>excavatum</i>	893	<i>Araeocerus</i>	669
<i>plana</i>	93	<i>exiguum</i>	876	<i>niger</i>	672
<i>polita</i>	190	19. <i>finetarium</i>	899	<i>Araeocnemis</i>	301
36. <i>praetexta</i>	176	13. <i>flavipenne</i>	896	<i>flagellicornis</i>	304
<i>procerula</i>	145	1. <i>florale</i>	891	<i>fulgens</i>	303
<i>prolixa</i>	63. 155	<i>Gyllenhalii</i>	859	<i>pubescens</i>	304
13. <i>puberula</i>	165	20. <i>Lapponicum</i>	899	<i>Arpedium</i>	858

	pag.		pag.		pag.
2. brachypterum	859	21. opacus	771	<i>carbonaria</i>	75
3. humile	860	40. pallipennis	779	<i>castanoptera</i>	103
1. quadrum	858	23. pallipes	772	<i>cincta</i>	59
4. Sibiricum	860	<i>pallipes</i>	767	<i>cinnamomea</i>	127
<i>Astenus</i>	637	29. parvulus	775	<i>circellaris</i>	98
- <i>angustatus</i>	640	11. politus	766	<i>collaris</i>	43
<i>procerus</i>	638	16. procerulus	768	<i>complana</i>	90
<i>Astilbus</i>	36	30. pumilio	775	<i>compressa</i>	136
<i>Astrapaeus</i>	552	26. pusillus	773	<i>crassicornis</i>	135
<i>rufipes</i>	415	27. pygmaeus	774	<i>depressa</i>	99
1. Ulmi	553	9. rubiginosus	765	<i>depressiuscula</i>	90
<i>Ulmicus</i>	553	<i>Ruddii</i>	761	<i>evanescens</i>	111
<i>Atemeles</i>	202	19. rufipennis	770	<i>excavata</i>	110
<i>Autalia</i>	46	<i>Skrimshiri</i>	761	<i>exilis</i>	90
1. impressa	47	35. subterraneus	777	<i>flavipes</i>	124
2. rivularis	47	33. talpa	777	<i>forticornis</i>	65
<i>Belonuchus</i>	419	1. taurus	760	<i>fracticornis</i>	81
8. agilis	423	6. tricornis	763	<i>funesta</i>	38
9. avidus	423	28. troglodytes	774	<i>fungi</i>	131. 132
15. connexus	927	39. tuberculatus	779	<i>fuscula</i>	133
2. dichrous	420	7. unicornis	764	<i>haemorrhoea</i>	121
14. ephippiatus	927	31. verres	776	<i>humeralis</i>	38
13. formosus	426	2. vitulus	761	<i>impressifrons</i>	141
9. gagates	424	<i>Boletobius</i>	268	<i>inguinalis</i>	98
1. haemorrhoidalis	420	1. analis	269	<i>limbata</i>	37
6. latro	422	14. atricapillus	276	<i>linearis</i>	81
10. minax	424	10. axillaris	274	<i>longiuscula</i>	81. 91
5. mordens	422	9. bicolor	274	1. lucida	58
12. mundus	425	7. cernuus	272	2. lunulata	58
13. planceps	425	24. cincticollis	922	<i>luridipennis</i>	136
3. Satyrus	421	17. cinctus	278	<i>minutissima</i>	134
4. xanthopterus	421	2. cingulatus	270	<i>nigriceps</i>	100
<i>Bisnius</i>	427	13. dimidiatus	276	<i>nigricollis</i>	64
<i>Bledius</i>	760	20. exoletus	280	<i>nigritula</i>	103
12. aquarius	766	4. formosus	271	3. obliqua	59
35. arenarius	778	3. inclinans	271	<i>oblonga</i>	90
41. armatus	779	16. lunulatus	278	<i>pallidula</i>	140
24. atricapillus	773	<i>merdarius</i>	272. 281	<i>parvula</i>	139
3. bicornis	762	11. niger	275	<i>patruelis</i>	132
38. brunnipennis	779	5. obscurus	272	<i>planiuscula</i>	90
<i>castaneipennis</i>	771	23. obsoletus	922	<i>prolixa</i>	63
42. cordatus	780	22. pulchellus	281	<i>pulchra</i>	58
20. crassicollis	770	21. pygmaeus	280	<i>pumilio</i>	195
36. debilis	778	8. rufus	273	<i>quisquiliarum</i>	84
18. dissimilis	769	6. seriatius	272	<i>reptans</i>	77
37. elongatus	779	15. speciosus	277	<i>sericans</i>	103
45. emarginatus	780	12. striatus	275	<i>socialis</i>	102
22. erraticus	772	18. trimaculatus	279	<i>tenella</i>	136
43. fasciatus	780	19. trinitatus	279	<i>teres</i>	89. 114
14. femoralis	768	<i>Bolitochara</i>	57	<i>terminalis</i>	89
13. fracticornis	767	<i>agaricola</i>	132	<i>validicornis</i>	129
4. hinnulus	762	<i>analis</i>	114	4. varia	59
5. iuvenus	763	<i>angustula</i>	91	<i>Boreaphilus</i>	899
10. lateralis	766	<i>annularis</i>	135	1. Henningianus	899
11. longulus	768	<i>atra</i>	73	<i>Brachydirus</i>	353
8. mandibularis	765	<i>atramentaria</i>	111	<i>xanthocerus</i>	353
44. melanocephalus	780	<i>atrata</i>	198	<i>Bryocharis</i>	268
25. nanus	773	<i>aterrima</i>	104	<i>analis</i>	270
17. nigricans	769	<i>bifoveolata</i>	115	<i>Cafius</i>	294. 427
32. Niloticus	776	<i>Boleti</i>	103. 109	<i>apicalis</i>	426

	pag.		pag.		pag.
<i>aterrimus</i>	492	4. pubescens	221	<i>Deliphrum</i>	872
<i>fucicola</i>	454	6. tumidus	223	4. algidum	874
<i>fulminans</i>	295	12. varians	227	5. angustatum	874
<i>nanus</i>	479	13. venustus	227	3. arcticum	873
<i>nigritulus</i>	492	COPROPHILINI	814	2. crenatum	873
<i>pumilus</i>	492	<i>Coprophilus</i>	815	1. tectum	873
<i>splendidulus</i>	478	<i>rugosus</i>	816	<i>Dermestes</i>	
<i>xantholoma</i>	452	1. striatulus	816	<i>brachypterus</i>	903
<i>Callicerus</i>	85	<i>Cordylaspis</i>	311	semicoleopratus	893
<i>hybridus</i>	85	1. pilosus	312	<i>Dianous</i>	688
<i>obscurus</i>	84	<i>tuberculatus</i>	342	1. coerulescens	689
<i>Spencei</i>	84	<i>Coryphium</i>	900	<i>Diglossa</i>	208
<i>Calodera</i>	64	<i>Creophilus</i>	347	1. mersa	209
10. aethiops	68	<i>badipennis</i>	400	<i>Dinarda</i>	200
2. forticornis	65	<i>chloris</i>	401	1. dentata	201
4. humilis	65	<i>ciliaris</i>	351	<i>Diochus</i>	300
5. longitarsis	66	<i>Erithacus</i>	401	1. nanus	301
3. nigricollis	64	<i>erythrocephalus</i>	351	<i>Dolicaon</i>	576
3. nigrita	65	<i>fasciatus</i>	349	5. biguttulus	578
12. protensa	69	<i>maxillosus</i>	348	4. gracilis	578
9. riparia	68	<i>oculatus</i>	352	3. haemorrhous	577
7. rubens	67	<i>variegatus</i>	351	2. Illyricus	577
6. rubicunda	66	<i>villosus</i>	349	1. Lathrobioides	576
<i>testacea</i>	68	<i>Cryptobium</i>	561	<i>Drusilla</i>	35
8. uliginosa	66	26. agile	575	<i>canaliculata</i>	36
11. umbrosa	69	8. albipes	566	<i>exarata</i>	72
<i>Carabus</i>		10. antiquum	567	<i>Echiaster</i>	636
<i>abbreviatus</i>	850	17. assimile	570	1. longicollis	637
<i>dimidiatus</i>	856	1. badium	562	2. melanurus	637
<i>Staphylinoides</i>	856	3. bicolor	563	<i>Eleusis</i>	836
<i>Caranistes</i>	925	20. brevicorne	572	<i>tibialis</i>	836
1. Westermanni	925	2. Carolinum	563	<i>Emus</i>	346
<i>Carpalinus</i>	801	21. curticorne	572	<i>aeneicollis</i>	413
<i>bilineatus</i>	806	22. curtum	572	<i>aeneocephalus</i>	412
<i>Cateretes</i>		12. dispar	568	<i>angustatus</i>	417
<i>brachypterus</i>	903	7. fasciatum	565	<i>attenuatus</i>	543
<i>Centroglossa</i>	209	11. femorale	567	<i>boops</i>	548
<i>attenuata</i>	919	27. fracticorne	575	<i>brunnipes</i>	410
<i>conuroides</i>	210	9. fulvipes	566	<i>Carinthiacus</i>	381
<i>minuta</i>	919	15. geniculatum	569	<i>chalcocephalus</i>	412
<i>Chasolium</i>	838	14. latebricola	569	<i>chloropterus</i>	373
Ernestini	838	6. longipenne	565	<i>chrysocephalus</i>	371
<i>Cyllaeus</i>	7	4. melanocephalum	564	<i>cyaneus</i>	405
<i>Conurus</i>	218	23. obscurum	573	<i>erythropterus</i>	378
3. australis	221	18. ochropus	571	<i>floralis</i>	526
9. basalis	225	13. pallipes	578	<i>fossor</i>	377
8. binotatus	224	5. prolixum	564	<i>fuscatus</i>	417
19. bipunctatus	230	25. pulchellum	574	<i>fuscomaculatus</i>	376
7. bipustulatus	223	24. scitum	574	<i>hirtus</i>	346
(11) <i>cinctulus</i>	226	16. similipenne	570	<i>impressus</i>	530
5. crassus	222	19. sulphuripes	571	<i>inauratus</i>	363
15. fumatus	228	<i>Cucuius</i>		<i>laevigatus</i>	529
16. fuscus	229	<i>maxillosus</i>	825	<i>lateralis</i>	525
2. grossus	220	<i>spinosus</i>	832	<i>marginalis</i>	361
10. interruptus	225	<i>Cypha</i>	214	<i>maxillosus</i>	348
1. littoreus	219	<i>biguttata</i>	919	<i>molochinus</i>	535
17. lividus	229	<i>Deinopsis</i>	919	<i>morio</i>	417
14. obsoletus	228	<i>fuscatus</i>	919	<i>murinus</i>	362
11. opicus	919	<i>Deleaster</i>	818	<i>nebulosus</i>	361. 373
18. pedicularius	230	1. dichrous	819	<i>nigripes</i>	416



<i>nitidus</i>	pag. 526	10. thoracica	pag. 52	8. dispar	pag. 186
<i>occultus</i>	526	20. venustula	55	7. dissimilis	186
<i>olens</i>	405	<i>Gabrius</i>	427	11. flaveola	188
<i>picipennis</i>	413	<i>Glyptoma</i>	908	5. gentilis	185
<i>praecox</i>	542	2. corticinum	909	10. lucidula	187
<i>pubescens</i>	372	3. costale	909	15. manca	190
<i>rufipalpis</i>	418	1. crassicorne	908	19. minima	192
<i>rufipes</i>	415	4. denticolle	910	3. nana	184
<i>scintillans</i>	546	5. exile	910	<i>nana</i>	185
<i>similis</i>	409	6. sculptile	911	2. nitidula	183
<i>speciosus</i>	360	<i>Goerius</i>	404	<i>nitidula</i>	184
<i>stercorarius</i>	380	<i>Goniodes</i>	202	16. polita	190
<i>subuliformis</i>	516	<i>Gymnurus</i>	679	14. socia	189
<i>tristis</i>	537	<i>cyanescens</i>	680	12. splendidula	188
<i>vicinus</i>	552	<i>vestitus</i>	680	17. strictula	191
<i>Encephalus</i>	183	<i>Gymnusa</i>	211	6. vinula	186
<i>complicans</i>	183	1. brevicollis	212	<i>Habrocerus</i>	242
<i>Eristethus</i>	746	<i>dubia</i>	210	1. capillaricornis	243
<i>ruficapillus</i>	747	2. fuscata	919	<i>Haematodes</i>	340
<i>scaber</i>	746	(2.) laticollis	212	1. bicolor	340
<i>Euaesthetus</i>	746	<i>Gyrohyphus</i>	306	<i>Hesperophilus</i>	760
3. Americanus	746	<i>alternans</i>	297	<i>Heterothops</i>	515
4. immarginatus	748	<i>analisis</i>	314	2. binotatus	516
2. ruficapillus	747	<i>assimilis</i>	924	3. dissimilis	517
1. scaber	746	<i>aterrimus</i>	492	5. exilis	518
<i>Eulisus</i>	306	<i>batychnus</i>	335	6. nitens	518
<i>ater</i>	318	<i>consentaneus</i>	326	1. praevius	516
<i>chalybaeus</i>	309	<i>flavipennis</i>	326	4. 4-punctulus	517
<i>Mannerheimii</i>	310	<i>fulminans</i>	295	<i>Holius</i>	298
<i>Euryporus</i>	553	<i>glaber</i>	325	2. analis	299
1. picipes	554	<i>glabratus</i>	319	3. debilis	299
2. puncticollis	554	<i>lentus</i>	326	4. Guildingii	300
<i>Euryusa</i>	199	<i>linearis</i>	335	1. humilis	299
1. sinuata	199	<i>longiceps</i>	332	<i>Holotrochus</i>	757
<i>Falagria</i>	48	<i>macrocephalus</i>	297	3. crassicollis	758
21. albicincta	56	<i>melanocephalus</i>	296	2. cylindrus	758
23. albofasciata	56	<i>merdarius</i>	319	1. volvulus	758
<i>caesa</i>	49	<i>meridionalis</i>	323	<i>Homalota</i>	80
7. concinna	51	<i>nanus</i>	479	42. aemula	102
22. deliciata	56	<i>nigriceps</i>	297	20. aequata	92
4. discisa	50	<i>nigritulus</i>	492	84. alternata	119
11. discreta	53	<i>ochraceus</i>	328	118. ambigua	134
2. dissecta	49	<i>ochropterus</i>	320	70. analis	114
6. divisa	51	<i>parumpunctatus</i>	335	99. anceps	125
12. fissula	53	<i>pilicornis</i>	296	122. angustata	136
17. gratilla	54	<i>procerulus</i>	511	19. angustula	91
<i>lineolata</i>	52	<i>punctulatus</i>	328	120. annularis	135
19. munda	55	<i>pyropterus</i>	319	22. arcana	93
18. naevula	55	<i>relucens</i>	319	117. aterrima	133
16. nigra	54	<i>splendidulus</i>	479	62. atramentaria	111
15. obscura	54	<i>tenuicornis</i>	338	89. atratula	121
9. percisa	52	<i>tricolor</i>	332	117. atricilla	101
<i>picea</i>	54	<i>xantholoma</i>	452	66. autumnalis	113
<i>polita</i>	50	<i>Gyrophaena</i>	182	76. bisulcata	117
5. scissa	50	4. affinis	184	128. borealis	138
14. soluta	53	<i>amabilis</i>	185	27. brevicornis	95
8. stricta	51	18. Boleti	191	36. brunnea	98
1. sulcata	49	1. complicans	183	33. caesula	97
<i>sulcata</i>	50	9. congrua	187	7. callicera	84
3. sulcatula	50	13. corruscata	189	91. cauta	122

	pag.		pag.		pag.
92. celata	122	50. marcida	106	2. discoideus	216
88. cinctella	121	129. marginalis	138	<i>flavicornis</i>	181
104. cinnamomea	127	78. melanura	117	<i>globulus</i>	215
34. circellaris	98	21. nigella	92	<i>granulum</i>	215
68. clancula	114	38. nigrifrons	100	3. laeviusculus	216
115. clientula	133	83. nigripennis	119	1. longicornis	215
100. concolor	126	101. notha	126	5. pulicarius	217
54. conformis	108	67. oblita	113	4. seminulum	217
119. crassicornis	134	40. oblonga	101	Hypotelus	840
81. curticolis	118	80. obsoleta	118	2. praecox	841
30. cuspidata	96	4. occulta	83	1. pusillus	841
11. debilicornis	87	45. ochracea	104	<i>Ino</i>	8
24. debilis	94	114. orbata	132	<i>Ischnopoda</i>	70
82. decipula	119	116. orphana	133	<i>Ischnosoma</i>	251
60. deplanata	110	108—9. paedida	917	<i>Isomalus</i>	838
37. depressa	99	5. pagana	83	4. bicolor	839
52. dichroa	107	72. paleola	115	1. complanatus	839
75. didyma	116	132. pallidula	110	2. humilis	839
<i>dimidiata</i>	80	131. parva	139	3. interruptus	839
106. discors	128	8. pavens	85	5. testaceus	840
17. elongatula	89	49. picta	106	Lathrimaeum	869
59. excavata	110	23. plana	93	3. atrocephalum	870
71. exilis	115	<i>plana</i>	96	6. canaliculatum	871
90. fervidula	121	35. prociua	98	4. fuscum	870
63. festinans	112	53. propinqua	108	2. luteum	869
79. flavipennis	118	<i>pulicaria</i>	122	1. melanocephalum	869
97. flavipes	124	<i>quisquiliarum</i>	85	5. sordidum	871
112. fungi	131	13. ravilla	88	Lathrobium	588
126. fusca	137	93. recondita	123	13. angustatum	596
65. gemina	112	3. rigidicornis	82	7. angusticolle	593
25. gracilenta	94	110. rubicula	130	28. armatum	603
10. gracilicornis	86	127. scapularis	137	<i>badium</i>	562
1. graminicola	81	56. semilutea	104	8. bicolor	593
12. gregaria	87	85. seminigra	120	<i>bicolor</i>	563
121. haemorrhoea	135	105. semirufa	128	<i>biguttulum</i>	578
41. hepatica	102	<i>sericans</i>	103	30. brevicorne	604
102. hirta	126	87. silacea	120	1. brunnipes	588
28. illota	95	43. socialis	102	<i>castaneum</i>	596
31. immersa	96	44. sodalis	104	<i>cinctum</i>	569
103. importuna	127	95. sordidula	123	20. collare	599
134. impressifrons	141	51. spelaea	107	<i>confluentum</i>	615
58. incana	109	86. stigma	120	<i>depressum</i>	582
73. inconspicua	116	94. subida	123	25. dilatatum	602
74. inquinula	116	98. subsinuata	125	18. dimidiatum	598
29. intrusa	96	123. tenella	136	17. discolor	598
14. labilis	88	140. tenera	140	24. dividiuum	601
57. laesicollis	109	109. terminata	130	2. elongatum	589
9. languida	86	111. testudinea	131	<i>elongatum</i>	590
18. linearis	91	46. trimaculata	105	22. filiforme	601
<i>livida</i>	129	2. umbonata	82	<i>fratricorne</i>	575
108. lividipennis	129	77. ustulata	117	3. fulvipenne	590
107. longicornis	129	48. varia	106	<i>gracile</i>	578
47. lunata	105	15. velata	88	<i>Gravenhorstii</i>	568
55. lurida	108	113. vernacula	131	<i>humile</i>	586
124. luridipennis	136	64. vestigialis	112	10. labile	594
96. luteicornis	124	6. vestita	84	<i>laevigatum</i>	605
16. luteipes	89	61. viduata	111	<i>latipes</i>	674
69. luteola	114	32. vilis	97	<i>lineare</i>	592. 600. 601
26. macella	95	Hygronoma	79	16. longiusculum	597
125. macilenta	137	1. dimidiata	80	26. longulum	602
130. Mannerheimii	139	Hypocypsus	214	15. Lusitanicum	597

	pag.		pag.		pag.
<i>millepunctatum</i>	619	<i>caraboides</i>	850. 851	3. emarginata	204
<i>minimū</i>	607	<i>dichroa</i>	819	4. inflata	205
<i>minutum</i>	602	<i>flavipennis</i>	849	2. paradoxa	202
5. multipunctum	591	<i>globulicollis</i>	853	1. strumosa	208
19. nitidum	599	<i>Laponica</i>	849	<i>Medon</i>	610
<i>obsoletum</i>	623	<i>Leachii</i>	819	<i>Ruddii</i>	611
23. pallidum	601	<i>longipes</i>	853	<i>Megacronus</i>	268
<i>pallipes</i>	568	6. longula	857	<i>elegans</i>	922
14. pectorale	596	3. maura	856	<i>formosus</i>	271
<i>phoenicurum</i>	580	<i>obscura</i>	856	MEGALOPINI	750
9. picipes	594	<i>plagiata</i>	852	<i>Megalops</i>	751
<i>pilosum</i>	595	<i>praeusta</i>	852	1. caelatus	751
<i>politum</i>	596	2. pubescens	856	3. cephalotes	752
21. punctatum	600	5. punctata	857. 933	2. punctatus	752
29. puncticolle	604	<i>punctulata</i>	855	<i>Megarthus</i>	904
<i>punctulatum</i>	591	4. Sicula	857	<i>clypeatus</i>	907
11. quadratum	595	<i>suturalis</i>	852	3. denticollis	906
<i>rufescens</i>	619	<i>testacea</i>	850. 851	1. depressus	905
<i>ruficorne</i>	614	<i>Lispinus</i>	828	4. hemipterus	906
4. rufipenne	591	4. angustatus	829	<i>marginicollis</i>	906
<i>rufiventre</i>	613	1. attenuatus	828	<i>rufescens</i>	906
27. scabricolle	603	7. exiguus	830	2. sinuatocollis	905
12. scutellare	596	2. laticollis	828	<i>Micralymma</i>	819
<i>similipenne</i>	570	5. linearis	829	1. brevipenne	820
6. spadicum	592	3. striola	829	<i>Johnstonis</i>	820
<i>sphaericolle</i>	813	6. tenellus	830	<i>Microcera</i>	179
26—27. stilicinum	932	<i>Lithocharis</i>	610	<i>Micropeplus</i>	911
<i>striatum</i>	585	8. angularis	614	2. caelatus	912
<i>terminatum</i>	595	<i>bicolor</i>	614	3. fulvus	912
<i>Leistotrophus</i>	358	13. biseriata	617	<i>Maillei</i>	913
<i>Gravenhorstii</i>	359	3. brunnea	612	6. obtusus	913
<i>Leptacinus</i>	333	1. castanea	610	1. porcatus	911
4. batychnus	335	19. cinnamomea	620	4. Staphylinoides	913
1. brevicornis	334	10. confluent	615	5. tesserula	913
8. clavicornis	337	17. corticina	619	<i>Microsaurus</i>	523
6. debilis	336	15. curtula	618	<i>Molosoma</i>	752
2. filarius	334	27. debilis	625	<i>latipes</i>	757
9. nothus	338	12. depressa	616	<i>Mycetoporus</i>	281
3. parumpunctatus	335	4. diluta	612	5. Americanus	285
7. subtilis	337	11. dorsalis	616	<i>2-maculatus</i>	284
5. tenellus	336	30. exilis	627	<i>brevicornis</i>	923
10. tenuicornis	338	6. ferruginea	613	4. lepidus	284
<i>Leptochirus</i>	825	21. fumata	622	3. longulus	283
8. bispinus	827	2. fuscula	611	8. lucidus	286
7. brevicornis	827	18. hepatica	620	7. nanus	286
<i>brunneo-niger</i>	825	22. infusata	622	<i>pullidulus</i>	288
<i>Cayennensis</i>	825	20. limbata	621	6. pronus	285
<i>convexus</i>	827	9. macularis	615	2. punctus	283
6. ebeninus	826	7. melanocephala	614	1. splendens	282
9. Javanicus	829	26. nigrifluta	625	9. splendidus	287
4. laticeps	826	14. oblita	618	<i>Myllaena</i>	209
1. maxillosus	825	25. obscura	624	1. dubia	210
3. Mexicanus	826	24. obsoleta	623	2. intermedia	210
2. scoriceus	825	23. ochracea	623	3. minuta	211
5. unicolor	826	16. posticata	619	<i>Myrmedonia</i>	35
<i>Lestoea</i>	855	5. rufiventris	613	12. armata	41
<i>alpina</i>	848. 849	29. sordida	626	14. atricapilla	42
<i>angusticollis</i>	850	<i>testacea</i>	623	16. boops	42
1. bicolor	855	<i>Lomechusa</i>	202	20. camura	44
<i>bimaculata</i>	852	<i>dentata</i>	201	1. canaliculata	36



	pag.		pag.		pag.
17. collaris	43	1. Paederinus	685	21. lineare	884
19. coriacea	43	Oligota	179	14. lucidum	881
15. frontalis	42	6. apicata	182	33. luridum	889
8. fulgida	40	2. atomaria	180	<i>macropterum</i>	903. 905
5. funesta	38	5. flavicornis	181	<i>mandibulare</i>	817
4. humeralis	38	4. granaria	181	34. marginatum	890
23. inscita	45	1. pusillima	179	<i>megacephalum</i>	844
21. labida	44	3. subtilis	180	<i>melanocephalum</i>	870
3. limbata	37	Olisthaerus	843	8. minimum	878
7. lugens	39	1. megacephalus	843	2. monilicorne	876
9. marginicollis	40	2. substriatus	844	11. nigrum	880
22. munda	44	Olophrum	863	<i>ophthalmicum</i>	894. 895
10. obscura	41. 915	7. alpestre	867	<i>ovatum</i>	903
18. ochracea	43	4. assimile	865	4. Oxyacanthae	877
25. parilis	45	5. boreale	866	<i>Oxyacanthae</i>	875
2. plicata	36	8. consimile	867	<i>pallidum</i>	894
6. rigida	39	9. emarginatum	868	27. pedicularium	887
13. speciosa	42	3. fuscum	865	26. philanthus	887
11. spinigera	41	2. obiectum	865	<i>piceum</i>	864
26. tumidula	46	1. piceum	864	25. planum	886
24. turbida	45	6. rotundicolle	866	<i>porcatum</i>	911
Nitidula		X. OMALINI	846	31. pubescens	889
<i>porcata</i>	911	Omalium	874	9. pusillum	879
<i>Staphylinoides</i>	913	<i>abdominale</i>	893	17. pygmaeum	883
<i>sulcata</i>	911	<i>aeneum</i>	821	<i>pygmaeum</i>	885
Ocalea	60	32. alpinum	889	<i>quadratum</i>	862
2. badia	61	<i>assimile</i>	865	<i>quadrum</i>	858. 862
1. castanea	60	<i>atrocephalum</i>	870	<i>Ranunculi</i>	896
6. decumana	62	<i>boreale</i>	866	7. repandum	878
5. murina	62	<i>brachypterum</i>	859. 903	1. rivulare	875
4. procera	61	20. brevicorne	884	<i>rotundicolle</i>	866. 868
7. proluxa	13	<i>brevipenne</i>	820	19. rufulum	883
8. rufilabris	63	13. brunneum	881	<i>rufum</i>	861. 862
3. spadicea	61	<i>brunneum</i>	881. 885	<i>rugosum</i>	816
Ocypus	403	6. caesum	878	<i>Salicinum</i>	880
8. Aethiops	927	<i>caesum</i>	877	12. Salicis	880
6. alpestris	408	<i>castaneum</i>	861	<i>Sibiricum</i>	860
11. alpicola	411	24. concinnum	886	<i>Sorbi</i>	895
18. ater	416	<i>consimile</i>	867	<i>Staphylinoides</i>	913
10. brunnipes	410	<i>crenatum</i>	861. 873	16. striatum	882
20. cerdo	417	35. cursor	890	<i>substriatum</i>	844
21. compressus	418	<i>denticolle</i>	907	<i>tectum</i>	872
14. cupreus	412	23. deplanatum	885	22. testaceum	885
2. cyaneus	405	<i>deplanatum</i>	885	<i>testaceum</i>	895
22. falcifer	418	<i>depressum</i>	885. 905. 907	<i>Viburni</i>	879
15. fulvipennis	413	3. exiguum	876	15. vile	882
12. fuscatus	411	29. Fahraei	888	Ophites	627
3. italicus	406	30. fenestrale	888	2. Raphidioides	628
4. macrocephalus	407	<i>fimelarium</i>	899	3. velitaris	629
(8) <i>masculus</i>	409	10. florale	879	1. versatilis	628
5. megacephalus	407	<i>florale</i>	891	OSORINI	753
19. morio	417	5. fossulatum	877	Osorius	753
1. olens	405	<i>fuscum</i>	865	1. ater	754
17. pedator	415	<i>Gyllenhalii</i>	859	3. Brasiliensis	753
13. picipennis	412	<i>impressum</i>	859	<i>brunnicornis</i>	757
9. picipes	409	<i>incisum</i>	875	7. cylindricus	756
<i>scabrosus</i>	927	18. inflatum	883	5. inciscirurus	755
7. similis	408	<i>laeve</i>	872	2. intermedius	754
16. Tataricus	414	28. laeviusculum	887	8. latipes	757
Oedichirus	684	20. Lapponicum	899	4. piceus	755

6. rugifrons	pag. 756	<i>brunneus</i>	pag. 241	<i>furcatus</i>	pag. 761
Othius	294	<i>caelatus</i>	751	11. incolumis	791
1. fulvipennis	295	<i>cellaris</i>	219	2. insectatus	787
5. macrocephalus	297	<i>chrysomelinus</i>	236	15. insignitus	793
2. melanocephalus	295	<i>cinctus</i>	558	17. intricatus	794
4. pilicornis	296	<i>erythropterus</i>	234	10. inustus	791
3. punctipennis	296	5. femoralis	558	<i>laesus</i>	816
Oxyopoda	141	<i>flavipes</i>	255	<i>latipes</i>	757
10. abdominalis	146	<i>hypnorum</i>	234	<i>longicornis</i>	788
7 alternans	145	7. lateralis	559	12. luteipennis	792
28. amicta	154	<i>lunulatus</i>	278	14. Madagascariensis	793
29. analis	154	4. maior	557	<i>melanocephalus</i>	780
26. cingulata	153	3. Mannerheimii	557	24. minimus	798
21—22. contractula	917	<i>marginatus</i>	235. 259	<i>minimus</i>	809
27. corticina	153	<i>marginellus</i>	263	<i>mordax</i>	782
19. cunctans	150	2. maxillosus	556	<i>morsitans</i>	782
17. cuniculina	149	<i>melanocephalus</i>	236	22. nanus	797
21. curtula	151	<i>merdarius</i>	236. 284	<i>nitens</i>	783
37. elegantula	158	<i>minutus</i>	235	18. nitidulus	795
18. exigua	150	<i>picipes</i>	554	<i>nodifrons</i>	783
16. exoleta	149	<i>pygmaeus</i>	281	<i>pallipennis</i>	779
13. ferruginea	147	<i>rufipes</i>	254	<i>pallipes</i>	767. 772
33. fumida	156	1. rufus	556	<i>pedicellus</i>	795
30. gracilis	154	<i>Schönherri</i>	556	13. Pensylvanicus	792
12. helvola	147	8. stygicus	558	4. piceus	788
24. lateralis	152	<i>subterraneus</i>	259	16. politus	794
20. lentula	150	<i>suturalis</i>	245	<i>pulcher</i>	786
<i>lividipennis</i>	129	<i>testaceus</i>	221	21. pumilus	797
6. longiuscula	144	<i>thoracicus</i>	281	<i>pusillus</i>	796
2. luteipennis	143	3-maculatus	279	1. rugosus	786
<i>luteipennis</i>	142	6. vittatus	558	<i>rugulosus</i>	795
34. maura	156	VII. OXYTELINI	749	9. sculpturatus	790
22. melanaria	151	O. GENUINI	759	5. sculptus	788
<i>obfusca</i>	132. 134	Oxytelus	785	<i>talpa</i>	777
4. opaca	144	<i>americanus</i>	793	<i>taurus</i>	760
23. pellucida	152	<i>arenarius</i>	778	<i>terrestris</i>	786
15. praecox	148	<i>armatus</i>	779	<i>tricornis</i>	764
<i>procerula</i>	145	<i>asphaltinus</i>	933	<i>unicornis</i>	764
31. proluxa	155	<i>atricapillus</i>	773	IV. PAEDERINI	560
15. promiscua	148	6. Bengalsis	789	<i>Paederus</i>	649
3. prospera	143	<i>biarcuatus</i>	933	15. aequinoctialis	657
36. pulchella	157	<i>bicornis</i>	831	10. aestuans	655
32. ruficollis	155	<i>caesus</i>	800	<i>angustatus</i>	640
1. ruficornis	142	8. Caffer	790	37. australis	666. 933
9. sagulata	146	<i>carinatus</i>	786. 787	<i>bicolor</i>	614
25. sericata	153	7. collaris	789	<i>biguttatus</i>	690. 735
<i>sericea</i>	176	19. complanatus	795	<i>binotatus</i>	645
35. suturalis	156	<i>cordatus</i>	780	16. Brasiliensis	658
8. togata	145	<i>cornutus</i>	782	2. brevipennis	651
5. umbrata	144	<i>corticinus</i>	806. 809	<i>brunnipes</i>	589
OXYPORINI	522	20. depressus	796	6. caligatus	652
Oxyporus	555	<i>depressus</i>	796	3. Capensis	651
<i>abdominalis</i>	233	<i>emarginatus</i>	780	<i>castaneus</i>	610
<i>analis</i>	232	23. exiguus	798	30. chalybeus	664
<i>angularis</i>	556	<i>fasciatus</i>	780	<i>cinctus</i>	639
<i>atricapillus</i>	276	<i>femoralis</i>	768	36. coarctatus	666
<i>bicolor</i>	270.	<i>flavipes</i>	790	31. coerulescens	664
2-maculatus	219	<i>fratricornis</i>	767	<i>columbinus</i>	657
2-pustulatus	260	<i>fuliginosus</i>	808. 811	<i>confinis</i>	654. 932
Blumenbachii	559	3. fulvipes	787	20. conspicuus	660

	pag.		pag.		pag.
<i>corticinus</i>	619	<i>vestitus</i>	84	20. <i>coelites</i>	440
39. <i>cyanipennis</i>	933	<i>Palaestrinus</i>	343	<i>coeruleipennis</i>	433
25. <i>cianocephalus</i>	662	2. <i>aureus</i>	344	139. <i>compressicollis</i>	508
<i>dimidiatus</i>	488	1—2. <i>mutillarius</i>	924	62. <i>corruscus</i>	465
<i>discopunctatus</i>	643	1. <i>Sykesii</i>	344	57. <i>corvinus</i>	462
<i>diversicollis</i>	650	<i>Palaminus</i>	681	4. <i>cribratus</i>	431
<i>elongatus</i>	589	2. <i>piceus</i>	681	99. <i>cribricollis</i>	487
28. <i>erythroderus</i>	663	1. <i>pilosus</i>	682	9. <i>cyanipennis</i>	433
<i>extensus</i>	638	4. <i>testaceus</i>	683	73. <i>debilis</i>	472
23. <i>extraneus</i>	661	5. <i>variabilis</i>	683	24. <i>decorus</i>	442
33. <i>fastuosus</i>	665	<i>Pelecyphorus</i>	553	<i>denigrator</i>	450
19. <i>ferus</i>	659	<i>picipes</i>	554	124. <i>dimidiatipennis</i>	500
<i>filiformis</i>	595. 638	<i>Peliusa</i>	192	86. <i>dimidiatus</i>	480
<i>filum</i>	932	1. <i>labiata</i>	193	<i>dimidiatus</i>	475
<i>Finisterrae</i>	650	<i>Pella</i>	36	75. <i>discoideus</i>	474
<i>fracticornis</i>	575	<i>Philonthus</i>	426	56. <i>ebeninus</i>	461. 929
<i>fragilis</i>	630	16. <i>aeneus</i>	437. 928	132. <i>egregius</i>	505
<i>fulgidus</i>	319	68. <i>aeruginosus</i>	469	138. <i>elatus</i>	508
34. <i>fulvicornis</i>	665	<i>agilis</i>	470	145. <i>elongatulus</i>	511
<i>fulvipennis</i>	295	148. <i>agnatus</i>	513	136. <i>ephippium</i>	507
<i>fuscipes</i>	652	44. <i>albilabris</i>	455	107. <i>exiguus</i>	491
<i>glaberrimus</i>	575	34. <i>albipes</i>	449	<i>fasciatus</i>	397
9. <i>Javanus</i>	654	79. <i>alcyoneus</i>	476	69. <i>feralis</i>	469
<i>impressus</i>	36	105. <i>alumnus</i>	490	124—5. <i>ferrugatus</i>	931
21. <i>laetus</i>	660	22. <i>ambiguus</i>	441	133. <i>fervidus</i>	505
<i>laevigatus</i>	605	111. <i>analis</i>	495	23. <i>fibularius</i>	442
7. <i>limnophilus</i>	653	82. <i>angularius</i>	477	60. <i>figulus</i>	464
1. <i>littoralis</i>	650	100. <i>angusticollis</i>	487	98. <i>fimbriolatus</i>	486
<i>littoralis</i>	656	<i>anthrax</i>	473	46. <i>fimetarius</i>	456
13. <i>littorarius</i>	656	130. <i>apicalis</i>	931	21. <i>flagrans</i>	441
4. <i>longipennis</i>	651	151. <i>apicicornis</i>	931	53. <i>flavipennis</i>	460
<i>longiusculus</i>	643	14. <i>asphaltinus</i>	436	71. <i>flavolimbatus</i>	471. 929
5. <i>Madagascariensis</i>	652	<i>assimilis</i>	526	<i>flavopterus</i>	450
18. <i>mandibularis</i>	659	108. <i>astutus</i>	492	126. <i>fonticola</i>	501
22. <i>melampus</i>	660	109. <i>aterrimus</i>	492	<i>frontalis</i>	536
<i>melanocephalus</i>	614	19. <i>atramentarius</i>	440	42. <i>fucicola</i>	454
20—1. <i>melanurus</i>	932	18. <i>atratus</i>	439	96. <i>fulvipes</i>	485. 930
32. <i>memnonius</i>	664	<i>attenuatus</i>	546	91. <i>fumarius</i>	482
14. <i>Mexicanus</i>	657	129. <i>Baltimorensis</i>	503	58. <i>fumigatus</i>	463
38. <i>morio</i>	667	114. <i>bicolor</i>	495	10. <i>furvus</i>	434
<i>ochraceus</i>	623	128. <i>bilineatus</i>	503	<i>fuscicornis</i>	470
<i>orbiculatus</i>	631. 632. 634	33. <i>bimaculatus</i>	448	48. <i>fuscus</i>	457
12. <i>peregrinus</i>	656	135. <i>binotatus</i>	506	32. <i>gilvipes</i>	448
<i>procerus</i>	638	67. <i>bipustulatus</i>	468	55. <i>gratiosus</i>	461
<i>quadratus</i>	611	127. <i>bistriatus</i>	502	(130.) <i>haematurus</i>	504
8. <i>riparius</i>	653	50. <i>blandus</i>	458	112. <i>helvolus</i>	494
<i>riparius</i>	650. 651	<i>boops</i>	548	37. <i>hepaticus</i>	451
<i>rubricollis</i>	623	<i>brevicornis</i>	461	118. <i>herilis</i>	497
26. <i>ruficollis</i>	662	97. <i>brunneus</i>	486. 930	147. <i>humilis</i>	512
<i>rugepunctatus</i>	659	54. <i>cadens</i>	460	6. <i>hybridus</i>	432
35. <i>rutilicornis</i>	666	<i>capucinus</i>	531	85. <i>immundus</i>	479
11. <i>Sabaeus</i>	655	15. <i>carbonarius</i>	437	<i>impressus</i>	530
<i>semicyaneus</i>	681	63. <i>Caucasicus</i>	466	72. <i>infirmus</i>	472
<i>speculator</i>	650	30. <i>cautus</i>	446	47—48. <i>inquietus</i>	928
24. <i>tamulus</i>	661	87. <i>celer</i>	480	2. <i>intermedius</i>	429
19. <i>tempestivus</i>	658	45. <i>cephalotes</i>	455	54—5. <i>iopterus</i>	928
<i>testaceus</i>	592	<i>chrysurus</i>	367	88. <i>irregularis</i>	481
29. <i>thoracicus</i>	663	43. <i>cicatricosus</i>	454	124. <i>labilis</i>	500
27. <i>tricolor</i>	663	7. <i>cinctulus</i>	432	81. <i>laetus</i>	477
<i>tricolor</i>	331	140. <i>cinerascens</i>	509	8. <i>laevicollis</i>	433



<i>laevigatus</i>	pag. 529	17. <i>scutatus</i>	pag. 438	1. <i>spinifer</i>	pag. 178
3. <i>laminatus</i>	430	<i>scybalarius</i>	470	VIII. <b>PIESTINI</b>	823
<i>lateralis</i>	525	141. <i>sericeus</i>	509	<i>Piestus</i>	830
35. <i>lepidus</i>	450	146. <i>sobrinus</i>	512	1. <i>bicornis</i>	831
137. <i>leucotus</i>	507	47. <i>sordidus</i>	456	4. <i>capricornis</i>	833
92. <i>lomatus</i>	482	1. <i>splendens</i>	429	7. <i>erythropus</i>	834
25. <i>lucens</i>	443	84. <i>splendidulus</i>	478	5. <i>fulvipes</i>	833
149. <i>ludicrus</i>	514	111. <i>stenoides</i>	493	2. <i>Lacordairei</i>	832
103. <i>lustralis</i>	489	66. <i>stragulus</i>	468	11. <i>Mexicanus</i>	836
106. <i>luxurians</i>	491	<i>subfuscus</i>	445	8. <i>minutus</i>	834
<i>mandibularis</i>	928	61. <i>suspectus</i>	465	<i>oxytelinus</i>	832
27. <i>marginatus</i>	444	51. <i>suturalis</i>	459	6. <i>penicillatus</i>	834
<i>maurorufus</i>	541	88—9. <i>Tachiniformis</i>	930	9. <i>pygmaeus</i>	835
<i>maurus</i>	551	101. <i>tenuis</i>	488	3. <i>spinosus</i>	832
95. <i>micans</i>	484	12. <i>testaceipennis</i>	435	10. <i>sulcatus</i>	835
52. <i>microcephalus</i>	459	90. <i>thoracicus</i>	481	<i>sulcatus</i>	834
113. <i>mirabilis</i>	494	104. <i>trepidus</i>	489	V. <b>PINOPHILINI</b>	668
<i>moestus</i>	464	<i>tristis</i>	537	<i>Pinophilus</i>	669
<i>molochinus</i>	535	<i>trossulus</i>	492	13. <i>Aegyptius</i>	673
<i>multipunctatus</i>	498	64. <i>turbatus</i>	466	13. <i>albipes</i>	675
11. <i>nigerrimus</i>	434	94. <i>turbidus</i>	484	20. <i>brevicollis</i>	679
89. <i>nigrita</i>	481	28. <i>umbratilis</i>	445	18. <i>brunneus</i>	678
36. <i>nitidulus</i>	450	59. <i>umbrinus</i>	464	3. <i>caliginosus</i>	671
13. <i>nitidus</i>	435	<i>variabilis</i>	525	9. <i>Capensis</i>	673
<i>ochropus</i>	462	70. <i>varians</i>	470	5. <i>complanatus</i>	672
<i>opacus</i>	470	<i>varians</i>	461	19. <i>crassicollis</i>	678
117. <i>opulentus</i>	497	41. <i>variegatus</i>	453	11. <i>flavipes</i>	674
122. <i>orphanulus</i>	499	<i>varicolor</i>	537	14. <i>fulvipes</i>	676
39. <i>orphanus</i>	452	31. <i>varius</i>	447	7. <i>Javanus</i>	672
(122.) <i>orphanus</i>	499	74. <i>ventralis</i>	473. 929	10. <i>latipes</i>	674
<i>pachycephalus</i>	457	83. <i>vernalis</i>	478	6. <i>niger</i>	672
125. <i>pacificus</i>	501	134. <i>viduus</i>	506	8. <i>palmatus</i>	677
76. <i>palliatu</i>	474	38. <i>vilis</i>	451	4. <i>picipes</i>	671
150. <i>palmula</i>	514	93. <i>virgo</i>	483	15. <i>suffusus</i>	676
121. <i>parumpunctatus</i>	499	<i>viridanus</i>	430	13. <i>sulphuripes</i>	675
<i>pediculus</i>	531	40. <i>xantholoma</i>	452	1. <i>tenebrosus</i>	670
<i>picatus</i>	930	<i>xanthopterus</i>	421	17. <i>terminatus</i>	677
29. <i>piceatus</i>	446	116. <i>xanthopygus</i>	496	2. <i>torosus</i>	670
<i>picipes</i>	514	Phloeobium	907	<i>Pityophilus</i>	669
49. <i>placidus</i>	458	1. <i>clypeatum</i>	907	<i>Placusa</i>	194
26. <i>politus</i>	443. 928	<i>corticale</i>	907	6. <i>adscita</i>	197
112—3. <i>pretiosus</i>	930	<i>depressum</i>	905	7. <i>atrata</i>	198
144. <i>procerulus</i>	511	<i>marginicolle</i>	906	1. <i>complanata</i>	194
143. <i>prolixus</i>	510	<i>nitiduloides</i>	906	5. <i>despecta</i>	197
69—70. <i>promptus</i>	929	<i>sinuatocolle</i>	905	3. <i>humilis</i>	196
142. <i>pruinosis</i>	510	IX. <b>PHLOEOCHARINI</b>	842	4. <i>infima</i>	196
110. <i>puella</i>	493	Phloeocharis	844	2. <i>pumilio</i>	195
102. <i>pullus</i>	488	1. <i>subtilissima</i>	845	<i>pumilio</i>	196
119. <i>punctifer</i>	498	Phloeonaesus	799	<i>Platygnemus</i>	340
120. <i>punctus</i>	498	1. <i>caelatus</i>	800. 933	<i>lateritius</i>	340
<i>punctus</i>	499	2. <i>caesus</i>	800	<i>Platyprosopus</i>	292
78. <i>quisquiliarius</i>	475	Phloeopora	76	1. <i>beduinus</i>	293
<i>rigidicornis</i>	456	5. <i>columbina</i>	78	2. <i>elongatus</i>	294
77. <i>rubidus</i>	475	2. <i>corticalis</i>	77	4. <i>fuliginosus</i>	923
5. <i>ruficollis</i>	431	3. <i>latens</i>	78	3. <i>tamulus</i>	923
80. <i>rufimanus</i>	476	6. <i>religata</i>	77	<i>Platystethus</i>	781
131. <i>rufipennis</i>	504	4. <i>subtusa</i>	78	6. <i>Americanus</i>	784
<i>rufocinctus</i>	511	<i>Physetops</i>	414	2. <i>cornutus</i>	782
65. <i>sanguinolentus</i>	467	<i>tararicus</i>	414	<i>mordax</i>	782
<i>scitus</i>	528	<i>Phytosus</i>	177	1. <i>morsitans</i>	782

	pag.		pag.		pag.
3. nodifrons	783. 933	39. scintillans	549	<i>Americanus</i>	342
<i>scybalarius</i>	782	6. scitus	528	<i>Sphenoma</i>	142
<i>striolatus</i>	782	32. semiobscurus	544	<i>abdominalis</i>	146
5. spiculus	784	42. tenellus	542	III. STAPHILININI	290
4. spinosus	784	27. umbrinus	541	ST. GENUINI	339
<i>Porus</i>	36	45. vicinus	552	<i>Staphylinus</i>	345
<i>ochraceus</i>	43	36. virgulatus	547	<i>aeneicollis</i>	413
<i>Procirrus</i>	685	12. viridulus	532	<i>aeneocephalus</i>	379. 351
1. Lefebvrei	686	5. xanthopus	527		412
<i>Prognatha</i>	836	<i>Raphirus</i>	523	<i>aeneus</i>	429. 430. 437. 444
2. humeralis	837	<i>Remus</i>	427		681
1. quadricornis	837	<i>sericeus</i>	509	<i>aethiops</i>	927
<i>rufipennis</i>	837	<i>Rugilus</i>	629	77. Afer	389
<i>Pronomaea</i>	207	<i>bicolor</i>	614	<i>affinis</i>	297. 332
1. rostrata	208	<i>castaneus</i>	611	<i>agilis</i>	470. 472
XI. PROTEININI	901	<i>dentatus</i>	636	<i>albipes</i>	449
<i>Proteinus</i>	902	<i>fragilis</i>	630	<i>alpinus</i>	848
4. atomarius	904	<i>fuscus</i>	611	<i>alternans</i>	297
1. brachypterus	903	<i>laevigatus</i>	605	<i>amoenus</i>	433
2. brevicollis	903	<i>orbiculatus</i>	631. 632. 634	<i>analis</i>	269. 275. 528
3. macropterus	903	<i>rubricollis</i>	623	92. anceps	396
<i>Pseudopsis</i>	914	<i>rufipes</i>	631	<i>angularis</i>	275
1. sulcatus	914	<i>Scaphidium</i>		<i>angustatus</i>	630. 640
<i>Quedius</i>	523	<i>acuminatum</i>	215	<i>anonymus</i>	348
34. attenuatus	546	<i>longicorne</i>	215	<i>anthrax</i>	473
37. boops	548	<i>Scariphaeus</i>	342	46. antiquus	373
17. brevis	535	1. luridipennis	343	<i>apicalis</i>	931
15. bupthalmus	534	<i>Schizochilus</i>	358	<i>apicicornis</i>	931
10. capucinus	531	<i>Brasiliensis</i>	359	3. arcticus	348
11. chlorophanus	532	<i>Scimbalium</i>	579	<i>arenarius</i>	778. 782
33. collaris	545	1. anale	580	<i>armatus</i>	764
4. cruentus	527	2. planicolle	580	<i>assimilis</i>	865
16. curtus	534	3. testaceum	581	<i>ater</i>	416
1. dilatatus	524	<i>Scopaeus</i>	604	<i>aterrimus</i>	470. 492
24. fimbriatus	539	2. didymus	606	<i>atratus</i>	438. 439. 463
14. frenatus	533	6. exiguus	608	<i>atricapillus</i>	276
3. fulgidus	525	8. fasciatellus	609	<i>atricillus</i>	528
20. fuliginosus	537	5. infirmus	608	<i>atro-coerulescens</i>	406
35. hyperboreus	547	1. laevigatus	605	<i>atrox</i>	383
8. impressus	530	4. minimus	607	<i>attenuatus</i>	542. 546
29. infuscatus	543	3. minutus	606	<i>aureus</i>	344
13. labiatus	533	9. pulchellus	609	<i>auricomus</i>	369
7. laevigatus	529	7. pygmaeus	608	71. aurifluus	357
2. lateralis	525	<i>Scytalinus</i>	305	<i>azurescens</i>	405
40. lucidulus	550	1. serpentinus	305	<i>azureus</i>	383
28. maurorufus	542	<i>Siagonium</i>	836	99. badiipennis	400
44. maurus	551	<i>humerule</i>	837	<i>badius</i>	390
18. melochinus	535	<i>quadriforne</i>	837	<i>balteatus</i>	348
31. monticola	544	<i>Silpha</i>		<i>Baltimorensis</i>	503
38. oblitteratus	549	<i>blattoides</i>	864	<i>batychnus</i>	335
23. ochropterus	538	<i>clypeata</i>	907	<i>bicinctus</i>	381
9. pediculus	531	<i>hemiptera</i>	906	<i>bicolor</i>	495. 855
25. peltatus	540	<i>lutea</i>	895	<i>bicornis</i>	848
41. peregrinus	550	<i>melanocephala</i>	869. 870	<i>biguttatus</i>	691
21. picipes	537	<i>minuta</i>	896	<i>binaculatus</i>	279. 448. 470
22. planicus	538	<i>torquata</i>	895	<i>binotatus</i>	506
23. praecox	540	<i>Silusa</i>	205	40. biplagiatus	370
<i>praecox</i>	540	1. rubiginosa	206	<i>bipunctatus</i>	163. 168
30. rufipes	543	2. rubra	207	<i>bipustulatus</i>	168. 468
43. rufocinctus	551	<i>Smilar</i>	341		470. 690

	pag.		pag.		pag.
<i>blandus</i>	458	<i>concolor</i>	524	<i>flavopterus</i>	450. 485. 529
<i>Blattinus</i>	235	<i>conformis</i>	474	<i>flavopunctatus</i>	381
<i>Boleti</i>	191	<i>conicus</i>	235	<i>floralis</i>	879. 891
<i>Bombylius</i>	346	<i>connexus</i>	927	<i>fodiens</i>	377
<i>boops</i>	458	<i>contaminatus</i>	467	<i>formosus</i>	426
<i>borealis</i>	866. 867	<i>corruscus</i>	465	70. <i>fossator</i>	386
<i>brachypterus</i>	163. 800	<i>crassicollis</i>	411	52. <i>fossor</i>	377
<i>Brassicae</i>	819	<i>crenatus</i>	861	<i>fracticornis</i>	328. 767
<i>brevicollis</i>	212	<i>cruentatus</i>	861	<i>fragilis</i>	457
<i>brevicornis</i>	461	<i>cruentus</i>	527	<i>frontalis</i>	536
<i>brunneus</i>	98. 486. 881	<i>cupreus</i>	413	<i>fulgens</i>	303
<i>brunnipennis</i>	779	80. <i>cupricollis</i>	391	<i>fulgidus</i>	295. 319. 526
<i>brunnipes</i>	410	<i>cursor</i>	875		527
<i>bupthalmus</i>	706	33. <i>cyanelytrius</i>	366	<i>fuliginosus</i>	537
63. <i>Buquetii</i>	383	<i>cyaneus</i>	405	<i>fulminans</i>	295
54. <i>caesareus</i>	378	41. <i>cyanicollis</i>	370	<i>fulvipennis</i>	590
29. <i>calidus</i>	364	<i>cyanipennis</i>	433	62. <i>fulvipes</i>	381
73. <i>caliginosus</i>	388	82. <i>cyanomelas</i>	392	<i>fulvipes</i>	450. 485
103. <i>canaliculatus</i>	403	69. <i>Dauricus</i>	386	65. <i>fulvomaculatus</i>	384
<i>canaliculatus</i>	36	<i>debilis</i>	472	<i>fulvopterus</i>	295
<i>capucinus</i>	531	<i>decorus</i>	442	<i>fulvus</i>	850
<i>Caraboides</i>	850	<i>denigrator</i>	450	<i>fumarius</i>	482
<i>carbonarius</i>	437. 447	<i>dentatus</i>	589	<i>fuscatus</i>	411. 416
<i>carinatus</i>	786	<i>depressus</i>	905	75. <i>fusciornis</i>	389
56. <i>castaneus</i>	379	<i>diaphanus</i>	325	<i>fuscipennis</i>	444. 526
<i>castanopterus</i>	377	<i>dilatatus</i>	524. 537	<i>fuscipes</i>	163. 264. 410
<i>celer</i>	480	68. <i>dimidiatus</i>	385	51. <i>fuscomaculatus</i>	376
<i>cellaris</i>	219. 221	<i>dimidiatus</i>	480. 929	<i>fuscus</i>	137. 457
60. <i>chalconcephalus</i>	381	<i>discoideus</i>	470. 474	72. <i>gentilis</i>	387
<i>chalconcephalus</i>	412. 413	<i>dispar</i>	232. 235. 236	<i>geometricus</i>	162
100. <i>chloris</i>	401		241. 245	<i>glaber</i>	325. 326. 529
44. <i>chloropterus</i>	372	<i>distinctus</i>	526	<i>glabratus</i>	319
<i>chloroventris</i>	537	<i>divisus</i>	37	<i>glabricollis</i>	519
84. <i>chrysaster</i>	393	<i>dubius</i>	448	<i>globulifer</i>	409
38. <i>Chrysis</i>	369	<i>ebeninus</i>	461	90. <i>Goryi</i>	395
42. <i>chrysocephalus</i>	371	<i>edentulus</i>	471	<i>gracilis</i>	640
86. <i>chrysocomus</i>	394	<i>elegans</i>	323. 332	<i>gregarius</i>	654
<i>chrysomelinus</i>	242. 235	<i>elongatus</i>	328. 589	<i>Groenlandicus</i>	526
	236. 239	<i>emarginatus</i>	203. 204	89. <i>haemorrhoidalis</i>	394
<i>chrysopterus</i>	384. 394	<i>emmesus</i>	324	<i>haemorrhoidalis</i>	367. 420
34. <i>chrysopygus</i>	366	<i>ephippiatus</i>	927	<i>Havaniensis</i>	490
36. <i>chrysurus</i>	367	101. <i>Erithacus</i>	401	87. <i>Herculeanus</i>	394
<i>chrysurus</i>	926	8. <i>erythrocephalus</i>	351	30. <i>hilaris</i>	365
<i>cinctulus</i>	432	67. <i>erythrocnemus</i>	385	1. <i>hirtus</i>	346
<i>cinctus</i>	257. 514. 530	<i>erythropennis</i>	372	79. <i>Hottentottus</i>	390
<i>cinerascens</i>	509	3. <i>erythropterus</i>	377	<i>hybridus</i>	361
5. <i>cinerarius</i>	350	<i>erythropterus</i>	377. 378	<i>hypnorum</i>	235
21. <i>cingulatus</i>	360	<i>erythropus</i>	383	<i>immaculatus</i>	379
78. <i>cinnamopterus</i>	390	14. <i>facialis</i>	354	<i>immundus</i>	473. 479
<i>clavicornis</i>	694. 706. 732	94. <i>fasciatus</i>	397	<i>impressus</i>	47. 530
	734	<i>fasciatus</i>	184. 348	27. <i>inauratus</i>	363
<i>coenosus</i>	436	37. <i>faustus</i>	368	6. <i>incanus</i>	350
<i>coeruleipennis</i>	433	<i>Fellmanni</i>	547	<i>infuscatus</i>	359
<i>coerulescens</i>	405. 439	85. <i>femoratus</i>	393	<i>intermedius</i>	429
<i>collaris</i>	43	64. <i>ferox</i>	384	<i>interruptus</i>	370
97. <i>columbinus</i>	400	<i>ferrugineus</i>	924	<i>inversus</i>	531
<i>compressicollis</i>	509	17. <i>fetialis</i>	357	32. <i>iopterus</i>	366
<i>compressus</i>	418	<i>finetarius</i>	456. 472	<i>iracundus</i>	526
15. <i>comptus</i>	356	<i>flavescens</i>	350	<i>irregularis</i>	481
<i>concinus</i>	461. 886	<i>flavipes</i>	886	<i>Juno</i>	694



	pag.		pag.		pag.
<i>laevicollis</i>	433. 536	<i>morio</i>	411. 417	81. <i>praelongus</i>	391
<i>laevigatus</i>	263. 529	<i>morsitans</i>	419. 782	<i>procerulus</i>	511
<i>laminatus</i>	429. 430	<i>multipunctatus</i>	498. 856	<i>puberulus</i>	413
9. <i>lanio</i>	352	24. <i>murinus</i>	361	43. <i>pubescens</i>	371
<i>Lapponicus</i>	536	<i>murinus</i>	361	<i>pubescens</i>	219. 221. 371
<i>laqueatus</i>	786	50—1. <i>mysticus</i>	926	<i>pullus</i>	484
61. <i>latebricola</i>	382	98. <i>myrtilinus</i>	400	<i>punctatostratus</i>	284
<i>lateralis</i>	525	<i>nanus</i>	184. 479	<i>punctatus</i>	498. 589
<i>laticeps</i>	437	22. <i>nebulosus</i>	360	<i>punctipennis</i>	296
<i>laticollis</i>	536	<i>nebulosus</i>	348	<i>punctulatus</i>	328. 332. 411
47. <i>latro</i>	374	<i>niger</i>	409. 429	<i>punctus</i>	498. 499
<i>latus</i>	259	<i>nigriceps</i>	99	102. <i>purpurascens</i>	402
<i>lentus</i>	326	<i>nigricollis</i>	64	<i>pusillus</i>	485
<i>lepidus</i>	450	<i>nigrita</i>	481	<i>pygmaeus</i>	528. 883
25. <i>leucomus</i>	362	<i>nigritulus</i>	492	35. <i>pyraister</i>	367
<i>leucophthalmus</i>	412	<i>nigritulus</i>	437	<i>pyropterus</i>	319
<i>leucopus</i>	73	<i>nigrofulvus</i>	556	<i>quadratus</i>	595
<i>limbatus</i>	37. 310	<i>nigrophthalmus</i>	553	<i>4-punctatus</i>	546
<i>linearis</i>	332. 335	<i>nitens</i>	409. 448	<i>quisquiliarius</i>	475
<i>littoreus</i>	219. 279	<i>nitidicollis</i>	448	39. <i>regalis</i>	369
<i>longiceps</i>	332	<i>nitidulus</i>	235. 241. 450	<i>relucens</i>	320
<i>longicornis</i>	929	<i>nitidus</i>	319. 435. 465	<i>rigidicornis</i>	456
<i>lucens</i>	443	525. 530. 551		<i>riparius</i>	650. 654
<i>lucidus</i>	448. 472	66. <i>nobilis</i>	384	<i>rivularis</i>	875
<i>lucens</i>	409	<i>obscurus</i>	41. 416. 417	<i>rotundicollis</i>	439
88. <i>lugubris</i>	394		855	<i>ruficollis</i>	431. 662
<i>lugubris</i>	389	<i>obtusus</i>	232	<i>ruficornis</i>	543
<i>lunulatus</i>	58. 278	<i>ochraceus</i>	328. 332	<i>rufilabris</i>	519
91. <i>luridipes</i>	395	<i>ochropterus</i>	381	<i>rufipennis</i>	426. 504
<i>luridus</i>	484	<i>ochropus</i>	462	<i>rufipes</i>	254. 255. 263
59. <i>lutarius</i>	381	45. <i>ochropygus</i>	373		543. 879
<i>macrocephalus</i>	401	10. <i>oculatus</i>	352	<i>rufitarsis</i>	526
<i>macroelytron</i>	855	<i>olens</i>	405	<i>rufocinctus</i>	551
<i>maculatus</i>	219	<i>opacus</i>	470. 771	<i>rufus</i>	555
50. <i>maculosus</i>	375	<i>ophthalmicus</i>	405. 894	<i>rugifrons</i>	406
<i>maior</i>	405	<i>orbiculatus</i>	631. 634	<i>rugosus</i>	786. 788. 816
23. <i>marginalis</i>	361	16. <i>ornatus</i>	356	<i>rutilus</i>	318
<i>marginalis</i>	263. 444	<i>palliat</i>	474	<i>sanguinolentus</i>	467
<i>marginellus</i>	263. 530	<i>pallidipennis</i>	782	<i>saphyrinus</i>	309
22—23. <i>marmoratus</i>	926	<i>pallidus</i>	281	28. <i>sapphirinus</i>	364
<i>martialis</i>	235	<i>palmula</i>	514	18. <i>scenicus</i>	357
<i>masculus</i>	409	<i>parumpunctatus</i>	335	<i>scintillans</i>	549
<i>maurorufus</i>	541. 542. 546	<i>parvicornis</i>	470	<i>scitus</i>	528
<i>maurus</i>	551	<i>parvulus</i>	479	49. <i>scriptus</i>	375
2. <i>maxillosus</i>	348	<i>pedator</i>	415	95. <i>segmentarius</i>	397
<i>maxillosus</i>	405	<i>penetrans</i>	412	<i>semiobscurus</i>	544
<i>megacephalus</i>	407	<i>peregrinus</i>	550	74. <i>sepulchralis</i>	388
<i>melanocephalus</i>	281. 295	<i>philanthus</i>	887	<i>sericans</i>	486
	881. 906	57. <i>phoenicurus</i>	380	<i>sericeicollis</i>	413
<i>melanurus</i>	235	<i>piceus</i>	786. 788. 795	<i>sericeus</i>	412
<i>mendax</i>	485	<i>picipennis</i>	412. 413. 536	<i>serraticornis</i>	524
<i>mesdarius</i>	236. 272. 284	<i>picipes</i>	409. 537	26. <i>Sibiricus</i>	363
<i>mesomelinus</i>	256	<i>pilicornis</i>	296	<i>silphoides</i>	245
<i>metallicus</i>	438. 439	<i>pilosus</i>	342	<i>similis</i>	408. 417. 438
<i>micans</i>	484	<i>plagiatus</i>	852		733
<i>microcephalus</i>	459	<i>planatus</i>	511	<i>simplex</i>	470
<i>minutus</i>	882	<i>planus</i>	465. 886	<i>socialis</i>	106
<i>moestus</i>	464	<i>politus</i>	411. 444. 498	<i>sordidus</i>	103. 45
<i>molochinus</i>	535	<i>porcatus</i>	911	<i>spinipes</i>	86
<i>mordax</i>	406	<i>praecox</i>	540. 542	<i>splendens</i>	282. 41










	pag.		pag.		pag.
<i>splendidulus</i>	478	83. violaceus	392	15. egenus	698
58. stercorarius	380	<i>violaceus</i>	303. 406	97. elongatus	741
<i>striatulus</i>	816	55. vulpinus	379	45. excubitor	714
<i>striatus</i>	275. 786	11. xanthocerus	353	30. exiguus	706
<i>strigatus</i>	413	<i>xantholoma</i>	452	<i>femorialis</i>	708
<i>strumosus</i>	201. 203	<i>xanthopygus</i>	496	43. femoratus	713
<i>subcyaneus</i>	406	<i>zonatus</i>	390	<i>femorellus</i>	720
<i>subfuscus</i>	455	VI. STENINI	687	78. filum	731
<i>subpunctatus</i>	411	Stenus	689	82. flavicornis	733
<i>subterraneus</i>	259	<i>Aceris</i>	728	74. flavipes	729
<i>subuliformis</i>	457. 517	26. aemulus	704	<i>floralis</i>	879
<i>sulcatus</i>	49. 50. 786	71. aerosus	727	<i>foraminosus</i>	703
	788	39. agilis	711	40. fossilatus	711
96. superbus	398	95. anguinus	740	89. frontalis	737
<i>tachiniformis</i>	930	85. annularis	735	<i>fulvipes</i>	716
<i>Tataricus</i>	414	103. arculus	744	76. fuscicornis	730
<i>tectus</i>	872	46. Argus	714	49. fuscipes	716
<i>tenellus</i>	551	<i>Argus</i>	717	<i>fuscipes</i>	717
<i>tenuipennis</i>	408	99. ariolus	742	73. geniculatus	728
<i>tenuis</i>	488	9. asphaltinus	695	<i>geniculatus</i>	727. 733
31. tepidus	365	10. ater	696	93. gracilis	739
93. terminalis	396	42. aterrimus	712	3. guttula	691
<i>terminatus</i>	457	21. atratulus	701	69. hospes	726
<i>tessellatus</i>	361	91. attenuatus	738	50. humilis	716
13. testaceus	354	35. augur	709	72. impressus	728
<i>testaceus</i>	457	63. bifoveolatus	723	19. incanus	700
<i>tetracarinatus</i>	796	1. biguttatus	690	23. incrassatus	702
<i>thoracicus</i>	481. 662	<i>biguttatus</i>	689. 691. 692	8. intricatus	694
76. tomentosus	389	4. bimaculatus	692	87. iugalis	736
<i>tomentosus</i>	219. 221	59. binotatus	721	98. iunceus	741
<i>tricolor</i>	326. 331. 383	<i>binotatus</i>	722	7. Juno	694
	614	2. bipunctatus	691	<i>Juno</i>	692
<i>tricornis</i>	764. 767	<i>bipunctatus</i>	746	<i>Kirbyi</i>	691. 693
<i>trilobus</i>	782	<i>bipustulatus</i>	690	12. labilis	697
<i>trimaculatus</i>	279. 281	<i>boops</i>	694. 699. 706	67. languidus	725
<i>tristis</i>	412. 537	16. bupthalmus	699	101. latifrons	743
<i>tuberculatus</i>	779	<i>bupthalmus</i>	694. 700	94. lubricus	739
<i>Ulni</i>	553		704. 732. 734	41. lustrator	712
<i>Ulmineus</i>	553	88. callosus	737	100. paganus	742
<i>umbratilis</i>	445	55. campestris	719	77. pallipes	731
<i>umbrinus</i>	464	27. canaliculatus	704	<i>pallipes</i>	728
<i>unicolor</i>	405	<i>canaliculatus</i>	697. 699	38. palpalis	710
<i>Uralensis</i>	416	11. carbonarius	696	<i>palposus</i>	699
<i>ustulatus</i>	295	<i>carbonarius</i>	699	75. palustris	729
<i>vagepunctatus</i>	318	47. cautus	715	66. picipennis	725
<i>variabilis</i>	525. 527	58. chalcites	721	62. planus	723
<i>varians</i>	447. 461. 470. 480	84. cicindeloides	734	61. plantaris	722
7. variegatus	351	<i>cicindeloides</i>	706	<i>proboscideus</i>	728. 729
<i>variegatus</i>	589	20. cinerascens	701	44. proditor	713
<i>varius</i>	447	51. circularis	717	92. prolixus	738
12. velitaris	354	<i>circularis</i>	717	32. providus	707
<i>velox</i>	485	<i>clavicornis</i>	699. 732	53. pumilio	718
<i>velutinus</i>	385	<i>coerulescens</i>	689	102. punctatus	744
<i>ventralis</i>	473	6. colon	693	29. pusillus	705
19. venustus	357	17. colonus	699	106.4-punctatus	745
<i>vernalis</i>	232. 478	90. colubrinus	737	<i>riparius</i>	732
20. versicolor	359	104. contractus	744	105. rotundatus	745
<i>viduatus</i>	375	68. cordatus	726	13. ruralis	697
4. villosus	349	37. croceipes	710	14. rusticus	724
<i>villosus</i>	361. 362	52. declaratus	717	33. scrutator	708

	pag.		pag.		pag.
<i>similis</i>	733	1. <i>aeneum</i>	821	<i>niger</i>	275
83. <i>solutus</i>	734	<i>nigro-aeneum</i>	821	6. <i>nitidulus</i>	247
31. <i>speculator</i>	706	<i>Tachinus</i>	244	<i>obsoletus</i>	922
5. <i>stigmula</i>	693	<i>analis</i>	269. 270	<i>pullens</i>	254
14. <i>stygius</i>	698	12. <i>apicalis</i>	250	28. <i>pallipes</i>	261
70. <i>subaeneus</i>	727	<i>atricapillus</i>	276. 278	3. <i>piceus</i>	246
60. <i>subimpressus</i>	722	<i>atricaudatus</i>	278	23. <i>picipes</i>	257
34. <i>sylvester</i>	708	(29.) <i>axillaris</i>	261	2. <i>pictus</i>	246
79. <i>tarsalis</i>	732	<i>axillaris</i>	274	5. <i>pulchellus</i>	247
65. <i>tempestivus</i>	724	30. <i>basalis</i>	262	<i>pullus</i>	254
96. <i>teres</i>	740	<i>batychnus</i>	258	<i>punctus</i>	283
56. <i>unicolor</i>	720	<i>bicolor</i>	274	<i>pygmaeus</i>	281
48. <i>vafellus</i>	715	<i>biplagiatus</i>	359	4- <i>punctulus</i>	517
80. <i>vestigialis</i>	732	27. <i>bipustulatus</i>	260	36. <i>rufipennis</i>	266
<i>Sterculia</i>	301	14. <i>brevicollis</i>	251	19. <i>rufipes</i>	254
1. <i>coelestina</i>	302	10. <i>brunneus</i>	249	<i>rufipes</i>	255
3. <i>elegans</i>	303	<i>castaneus</i>	255	18. <i>rutilus</i>	253
5. <i>formicaria</i>	304	<i>cernuus</i>	272	<i>signatus</i>	254
2. <i>fulgens</i>	303	4. <i>cinctellus</i>	246	1. <i>silphoides</i>	245
4. <i>Leprieuri</i>	304	<i>cincticollis</i>	922	<i>sordidus</i>	264
<i>Stilicus</i>	629	<i>cinctus</i>	278	<i>splendidus</i>	287
6. <i>affinis</i>	633	33. <i>collaris</i>	264	<i>striatus</i>	275
9. <i>angularis</i>	634	7. <i>convexus</i>	248	25. <i>subterraneus</i>	259
11. <i>carinatus</i>	635	<i>corticinus</i>	264	<i>suturalis</i>	245
12. <i>dentatus</i>	636	15. <i>depressus</i>	251	11. <i>terminalis</i>	250
1. <i>fragilis</i>	630	35. <i>discoideus</i>	266	<i>trimaculatus</i>	279
7. <i>fuscipes</i>	633	<i>dissimilis</i>	517	<i>tristis</i>	284
5. <i>geniculatus</i>	632	<i>dubius</i>	255	17. <i>ventriculus</i>	920
10. <i>lucialis</i>	635	8. <i>elatus</i>	248	II. TACHYPORINI	213
8. <i>orbiculatus</i>	634	34. <i>elongatus</i>	265	<i>Tachyporus</i>	231
2. <i>rufipes</i>	631	25. <i>fimbriatus</i>	258	2. <i>abdominalis</i>	233
4. <i>similis</i>	631	32. <i>fimetarius</i>	264	<i>abdominalis</i>	233. 241
3. <i>subtilis</i>	631	<i>flavellus</i>	265	22. <i>acaudus</i>	920
<i>Sunius</i>	637	20. <i>flavipes</i>	255	23. <i>acuductus</i>	920
5. <i>angustatus</i>	640	<i>formosus</i>	271	24. <i>affinis</i>	920
10. <i>attenuatus</i>	643	<i>frigidus</i>	256	<i>analis</i>	232
6. <i>bimaculatus</i>	641	23—24. <i>fulvipes</i>	921	7. <i>anticus</i>	235
16. <i>binotatus</i>	645	29. <i>fumipennis</i>	921	(11.) <i>arduus</i>	237
17. <i>bipunctatus</i>	645	(17.) <i>gibbulus</i>	252. 920	<i>batychnus</i>	258
9. <i>bisignatus</i>	642	9. <i>hepaticus</i>	249	<i>bicolor</i>	259
12. <i>coarctatus</i>	644	22. <i>humeralis</i>	256	<i>bimaculatus</i>	223
15. <i>curtulus</i>	645	<i>humeralis</i>	261	<i>binotatus</i>	224
20. <i>exiguus</i>	647	<i>humidus</i>	922	<i>bipunctatus</i>	231
1. <i>filiformis</i>	638. 932	13. <i>humilis</i>	251	<i>bipustulatus</i>	223. 260
18. <i>hybridus</i>	646	<i>immaturus</i>	263	18. <i>brunneus</i>	241
4. <i>intermedius</i>	640	<i>inclinans</i>	271	<i>capillariicornis</i>	243
22. <i>labeo</i>	648	<i>intermedius</i>	263	<i>cellaris</i>	219
8. <i>lateralis</i>	642	<i>laticollis</i>	254. 263	8. <i>chrysomelinus</i>	235
7. <i>limbatus</i>	641	<i>lepidus</i>	284	<i>collaris</i>	264
3. <i>linearis</i>	639	<i>lunulatus</i>	276. 278	<i>corticinus</i>	265
11. <i>longiusculus</i>	643	20—21. <i>luridus</i>	920	<i>crassus</i>	222
21. <i>naevius</i>	647	<i>marginalis</i>	245	<i>dissimilis</i>	517
2. <i>prolixus</i>	639	<i>marginatus</i>	259	6. <i>erythropterus</i>	234
24. <i>pusio</i>	649	21. <i>marginellus</i>	263	<i>faber</i>	920
19. <i>stigma</i>	646	16. <i>melanarius</i>	252	<i>fimbriatus</i>	258
12. <i>testaceus</i>	643	<i>melanocephalus</i>	251	<i>fimetarius</i>	255. 264
14. <i>tristis</i>	644	24. <i>memnonius</i>	258	3. <i>formosus</i>	919
23. <i>troglydites</i>	648	<i>merdarius</i>	269. 271	<i>fuscus</i>	229
<i>Syntomium</i>	821	<i>nanus</i>	286	<i>gibbulus</i>	252












	pag.		pag.		pag.
<i>granulum</i>	215	6. exarata	72	9. morio	805
<i>humeralis</i>	257	4. ferialis	71	17. Niloticus	808
12. humerosus	238	11. flavitarsis	74	3. Omalinus	803
6. hypnorum	234	10. fugax	74	20. punctatellus	810
<i>immaturus</i>	263	10—11. immunita	916	23. pusillus	811
<i>interruptus</i>	224	7. laesa	73	13. riparius	807
11. iocosus	919	<i>rubicunda</i>	66	10. scrobiculatus	805
<i>lateralis</i>	238	5. scitula	72	2. segnis	802
<i>laticollis</i>	263	9. umbratica	73	5. signatus	803
<i>lepidus</i>	284	7—8. uvida	916	25. subtilis	812
<i>limbatus</i>	245	Taenodema	679	24. tenellus	811
<i>littoreus</i>	219	4. aenea	681	22. troglodytes	810
<i>marginalis</i>	245	1. cyanescens	680	<i>Tympanophorus</i>	402
<i>marginatus</i>	235	3. semicyanea	681	<i>canaliculatus</i>	403
<i>marginellus</i>	263	2. vestita	680	<i>Velleius</i>	523
<i>memnonius</i>	258	<i>Taenosoma</i>	801	<i>dilatatus</i>	524
<i>moestus</i>	919	<i>gracile</i>	811	<i>luridipennis</i>	343
16. nanus	240	<i>pusillum</i>	811	XANTHOLININI	291
21. nigriceps	242	Tanygnathus	288	Xantholinus	306
19. nigricornis	242	2. collaris	289	22. anachoreta	316
<i>nitidulus</i>	241	3. laticollis	289	17. analis	314
20. obscurellus	242	1. terminalis	288	26. ater	318
1. obtusus	232	<i>Tasgius</i>	415	49. attenuatus	330
<i>opicus</i>	919	<i>confinis</i>	416	14—15. aulicus	923
<i>pallipes</i>	261	<i>rufipes</i>	415	43. brunneipennis	327
<i>pedicularius</i>	229. 230	<i>Thoraxophorus</i>	908	<i>cadaverinus</i>	320
<i>pubescens</i>	221	<i>corticinus</i>	909	32. calidus	321
<i>pullus</i>	254	<i>Triacrus</i>	398	1. canaliculatus	307
<i>punctus</i>	283	<i>dilatatus</i>	398	33. castanopterus	321
14. pusillus	239	<i>Trichocoryne</i>	830	16. castigatus	315
<i>pusillus</i>	240	<i>striata</i>	834	42. cephalus	924
13. ruficollis	239	<i>Trichoderma</i>	410	11. chalcopterus	312
<i>ruficollis</i>	238. 241	<i>Trichophya</i>	267	6. chalybeus	309
<i>rufipes</i>	255	<i>pilicornis</i>	268. 921	10. chloropterus	311
(3.) <i>rufus</i>	233	Trichophyus	267	37. collaris	324
4. saginatus	234	1. pilicornis	268	(42.) <i>consentaneus</i>	326
<i>sagrinatus</i>	236	<i>Trichopygus</i>	515	9. cyanopterus	311
15. scitulus	240	<i>dissimilis</i>	516	38. decorus	324
<i>scutellaris</i>	241	<i>nitens</i>	518	23. dispilus	317
<i>sericeus</i>	221	<i>pumilio</i>	517	3. divisus	308
<i>signatus</i>	254	<i>Trigonophorus</i>	399	36. elegans	323
9. solutus	236	<i>myrtillinus</i>	400	39. emmesus	324
<i>splendidus</i>	287	Trogophloeus	801	<i>episcopalis</i>	335
<i>subterraneus</i>	259	7. angulatus	804	30. erythrocephalus	320
<i>suturalis</i>	245	4. angustatus	803	14. festivus	315
10. tersus	237	11. bilineatus	806	2. flavipennis	307
17. transversalis	240	18. corticinus	809	28. fulgidus	319
<i>trinotatus</i>	220	<i>corticinus</i>	806. 811	<i>fulgidus</i>	319
<i>truncatellus</i>	230	1. dilatatus	802	21. gaginginus	316
<i>ventriculus</i>	920	14. elongatulus	807	40. glaber	325
Tachyusa	69	19. exiguus	809	29. glabratus	319
8. atra	73	6. ferrugineus	804	<i>hamatus</i>	924
3. balteata	71	16. flavipes	808	47. Hesperius	329
13. carbonaria	75	21. foveolatus	810	44. humeralis	327
9—10. chalybea	916	15. fuliginosus	808	18. illucens	315
2. coarctata	71	8. fulvipes	804	13. insignis	312
14. coerulea	72	<i>impressus</i>	809	4. laticeps	308
1. constricta	70	<i>inquilinus</i>	806	41. lentus	325
12. coracina	75	12. memnonius	806	8. limbatus	310

	pag.		pag.		pag.
<i>limbatus</i>	924	<i>parumpunctatus</i>	335	51. tricolor	331
52. linearis	332	34. pectoralis	322	27. vagepunctatus	318
<i>longiceps</i>	332	16. phoenicopterus	314	20. vagus	315
31. lubricus	321	<i>pilicornis</i>	296	14. volvulus	317
15. lucidus	313	50. procerus	331	<i>Zirophorus</i>	830
5. lynceus	309	46. punctulatus	328	<i>coriaceus</i>	825
7. Mannerheimii	310	<i>pyropterus</i>	319	<i>fronticornis</i>	831
<i>meridionalis</i>	323	45. rufescens	328	<i>penicillatus</i>	834
<i>minutus</i>	335	35. rufipennis	322	<i>spinosus</i>	832
48. obscurus	330	25. rutilus	318	<i>striatus</i>	832
<i>occidentalis</i>	924	12. speciosus	312	<i>Zyras</i>	36
<i>ochraceus</i>	329. 332	<i>tener</i>	924	<i>Haworthi</i>	40















1.



2.



3.



4.



5.



6.



7.



8.



9.



10.



11.



12.



13.



14.



15.



16.



17.



18.



19.



20.



21.



22.




23.




24.



25.



26.



27.



28.



29.



